Histological divergence underlying globular body shapes in ornamental goldfish

Kinya G Ota太田欽也[1,✉], Gembu Abe阿部玄武[2], Chen-Yi Wang王貞懿[1], Ing-Jia Li李穎佳[1], and Paul Gerald Layague Sanchez[1]

[1]Laboratory of Aquatic Zoology, Marine Research Station, Institute of Cellular and Organismic Biology, Academia Sinica, Yilan, 26242, Taiwan
[2]Division of Developmental Biology, Department of Functional Morphology, School of Life Science, Faculty of Medicine, Tottori University, Nishi-cho 86, Yonago 683-8503, Japan

Body shape diversity in vertebrates reflects a complex interplay between functional demands, environmental constraints, and internal developmental mechanisms. Various environments have promoted diverse morphological adaptations not only under the natural but also domesticated conditions. One of the examples of the drastic morphological changes in body shape under the artificial condition (i.e. with significant human intervention) is the ornamental goldfish (*Carassius auratus*). Ornamental goldfish were originally derived from wild crucian carp. Over time, they diversified into numerous strains with strikingly different body shapes. In this study, we compared the body shapes of representative goldfish strains: the single-tail common goldfish (wild-type), *Ryukin*, *Oranda*, *Pearl scale*, and *Ranchu*. Our analysis revealed that the *Ryukin* and *Pearl scale* strains exhibit significantly greater body circularity in dorsal view compared to the other strains. Further anatomical and histological analyses showed that *Pearl scale* goldfish possess a thicker lateral body wall along with increased adipose tissue accumulation and reduced muscle fiber density, unlike *Ryukin* goldfish. These findings suggest that similar globular body shapes in different goldfish strains have arisen through distinct developmental pathways, exemplifying morphological convergence accompanied by histological divergence. We discussed the adipose accumulation in *Pearl scale* goldfish in relation to natural examples, aiming to investigate possible links between function, morphology, and histology.

circularity | adipose tissue | muscle fibers | *Carassius auratus*
Correspondence: *otakinya@gate.sinica.edu.tw*

BACKGROUND

The diversity of body shapes observed in aquatic vertebrates reflects adaptations to the physical constraints and ecological demands of their environments. For example, streamlined forms, such as those found in sharks, tunas, dolphins, and diving birds, have evolved independently across different vertebrate lineages, highlighting the advantage of this morphology for active swimming in open water; similarly, many other body shapes have also arisen convergently, as can be seen from early vertebrates to more derived lineages; for example, see Janvier (1996); Kardong (2006); Lampart-Kałużniacka and Heese (2000); Li et al. (2020); Liem et al. (2001); Nelson et al. (2016); Van Wassenbergh et al. (2015); Wainwright and Turingan (1997). This repeated emergence of the different types of morphologies across diverse taxonomic groups suggests that body shape of vertebrates is evolvable, representing adaptation to distinct environments.

Whereas natural selection in the wild has shaped body forms suited for survival under natural ecological constraints, domestication has introduced a distinct set of selective pressures. A particularly notable case is that of ornamental goldfish, which show a variety of body shapes evolved under aesthetic rather than wild ecological constraints (Le Verger et al. 2024; Ota 2021). Ornamental goldfish were derived from the wild crucian carp (*Carassius auratus*), which show a typical streamlined body shape. Initially, color variations of the single-tail common goldfish were isolated (Fig. 1A-C), and over time, several distinctive morphotypes were selected by breeders and fanciers (Fig.1D-G) (Chen 1954, 1956; Hervey and Hems 1948; Ota 2021; Ota and Abe 2016; Smartt 2001). These include phenotypes such as popped eyes, bifurcated caudal fins, and the absence of dorsal fins (Chen et al. 2020b; Hervey and Hems 1948; Ota 2021; Ota and Abe 2016; Smartt 2001). Notably, the globular body shape—a rounded and compact form—often emerged concurrently with these mutations in eyes and fins (Asano and Kubo 1972; Kon et al. 2020; Matsui 2006). Thus, this globular morphology is recognized as a defining characteristic of many ornamental goldfish strains (Smartt 2001).

For example, *Ryukin*, *Oranda*, *Ranchu*, and *Pearl scale* are ornamental goldfish strains known for their globular body shape (Asano and Kubo 1972; Ota 2021; Smartt 2001) (Fig. 1D–G). Because ornamental goldfish have traditionally been appreciated from a dorsal view (i.e. from ponds and/or opaque fish bowls), their external morphology has been shaped by artificial selection favoring a rounded appearance from a dorsal view. As a result, globular body shapes have become genetically fixed in these populations. Furthermore, if we add an explanation about the nature of this selection, it can be described as follows: this selection appears to have been based primarily on external form, rather than internal phenotypic features. This raised the question whether the globular body shapes seen in different ornamental strains emerged from similar types of modifications, or from distinct internal changes including developmental and tissue reorganization.

To address the question posed above, we conducted comparative analyses of representative ornamental goldfish strains (Fig. 1). Recognizing that goldfish fanciers and breeders have traditionally viewed the fish from above and selected for rounder body shapes (Chen 1954; Smartt 2001), we first focused on dorsal morphology and performed shape analysis

Fig. 1. Dorsal view of five ornamental goldfish. A–C. the single-tail common goldfish (wild-type). D. *Ryukin*. E. *Oranda*. F. *Pearl scale*. G. *Ranchu*. The individuals of single-tail, *Ryukin*, and *Oranda* are derived from our previous report (Li et al. 2019; Tsai et al. 2013). The photographs of *Pearl scale* and *Ranchu* are derived from our YouTube channel (https://www.youtube.com/@LAQZL ; see Goldfish EvoDevo series; accessed 2025-06-30). Panel labels ending in odd numbers show the original photographs. Panel labels ending in even numbers show the image with the color-coded region of interest (ROI). The colored ROIs indicate the approximately identified trunk regions. These goldfish individuals are ranging from 5 cm to 15 cm in standard length. The individuals shown in panels A1–B2 were obtained by artificial fertilization and subsequently reared at the Yilan Marine Research Station. The individuals of panels A, B, and C are denoted as Wt-Jp-2019, Wt-Jp-2015, and Wt-Tw-2015, respectively, in Fig. 2 and Table 1. The numbers on each individual correspond to the suffix of the Handle ID in Table 1 (e.g., the "-01" in Wt-Jp-2019-01). Source image available at ark:37281/k5f8z2s55, licensed under CC-BY-NC-SA 4.0, © Kinya G. Ota (2025).

from this perspective. We then carried out detailed anatomical and histological examinations to identify internal features contributing to the globular body features. We further discuss how artificial selection has historically influenced goldfish morphology and consider whether similar external forms across strains have arisen through shared or distinct developmental mechanisms—potentially representing a case of convergent and parallel evolution at the different phenotypic levels.

MATERIALS AND METHODS

Goldfish strains

All goldfish strains used in this study were maintained at the Yilan Marine Research Station (Institute of Cellular and Organismic Biology, Academia Sinica, Taiwan) (Fig. 1). Among them, the wild-type strains were derived from single-tail goldfish originating from Japanese and Taiwanese popu-lations; see also (Ota 2021; Tsai et al. 2013) for a detailed definition of the wild-type goldfish. Progenies used in this study were obtained by artificial fertilization as described previously (Li et al. 2019; Ota 2021; Ota and Abe 2016; Ota et al. 2025). In addition, the *Ryukin*, *Oranda*, *Ranchu*, and *Pearl scale* strains were purchased from local ornamental fish distributors in Taiwan. Some of these individuals were directly used for experiments, whereas others were bred to obtain progenies from their parental stocks. For progenies obtained and reared in our laboratory, their origin is specified in the corresponding figure legends.

Measurement of circularity

Goldfish individuals were photographed from the dorsal side (Fig. 1). The body circularity of goldfish specimens was measured using images captured under controlled conditions. Specimens were recorded against a uniform background (either black or white) using a digital camera or video record-

ing (Olympus; E-M1 Mark III). Images captured from a dorsal view were analyzed to determine the body trunk area and perimeter using the image analysis software Fiji (Schindelin et al. 2012). Details on how the trunk region was defined, as well as the validity of the criteria used, are described in the Results section. These parameters were manually detected. Circularity was calculated based on the detected area and perimeter, using the following formula:

$$\text{Circularity} = \frac{4\pi \times \text{Area}}{\text{Perimeter}^2}$$

Anatomical data analysis

After the data acquisition from live fish, the fish were anesthetized with MS-222 (E10521, Sigma). The anesthetized specimens were fixed overnight in 10% formalin. All experimental procedures were conducted in accordance with the animal care and use protocol approved by Academia Sinica (IACUC Protocol 23-10-2072). The fixed specimens were transversely sectioned at the anterior end of the pelvic fin using a microtome blade. The transverse sections were then photographed with a digital camera (Olympus; E-M1 Mark III). From transverse section images, we identified (1) the maximum body width (mbw), defined as the longest distance between the left and right lateral surfaces of the body in the transverse section image, and (2) the internal cavity width (icw), defined as the distance between the left and right boundaries of the body cavity wall. The lateral body wall thickness (lbw)—defined as the combined thickness of the tissue from the skin surface to the body cavity wall on both sides—was calculated by subtracting icw from mbw. In addition, the circularity of the body surface contour and of the coelomic cavity contour was measured from the transverse section images.

Histological analysis

The 10% formalin fixed specimens from the anatomical data acquisition were dehydrated through ethanol series. After dehydration, specimens were cleared in Lemosol A (5989-27-5, FUJIFILM Wako Pure Chemical Corporation), embedded in paraffin, and sectioned to 5 μm using a microtome (RM2245, Leica). The sections were placed on microscope slides (Platinum-pro, Matsunami). For the conventional histological analyses, the section was deparaffinized with Histo-Clear II (HS-202, National Diagnostics), gradually hydrated with ethanol series, and stained by alcian blue, hematoxylin, and eosin as described previously (Li et al. 2019; Ota et al. 2025). The slide was dehydrated with ethanol series, immersed in Lemosol A, and sealed by coverslip with Entellan new mounting media (107961, Sigma-Aldrich). Histological sections were observed under a microscope, and images were captured focusing on the lateral side of the body wall area. From the histological images, regions containing both muscle and adipose cells were selected, with a focus on areas where adipose cells were densely distributed. Muscle bundles or groups of muscle bundles that were surrounded or adjacent to adipose cells were visually identified and selected. Based on

the distribution patterns of muscles and adipose cells, the histological images were scored (see the Results part). The analyses and visualization of the data were conducted by RStudio and SuperPlotsOfData (Goedhart 2021), which is accessible at https://huygens.science.uva.nl/SuperPlotsOfData/.

Fig. 2. Circularity of five ornamental goldfish The plot and distribution is based on the image of Fig. 1. The distributions are calculated by SuperPlotsOfData (https://huygens.science.uva.nl/SuperPlotsOfData/). The small dots indicate the circularity of each individual. The large dots indicate the means of circularity. Original data is in Table 1. Source image available at https://n2t.net/ark:/37281/k5f8z2s55, licensed under CC-BY-NC-SA 4.0, © Kinya G. Ota (2025).

RESULTS

To examine the body shape variation among ornamental goldfish strains, we measured the circularity of their dorsal views. Given that the early breeders and fanciers selected their preferred goldfish primarily through visual inspection from dorsal view, we also outlined the body shape of the goldfish and measured the circularity of this outline based on our semi-quantitative visual inspection (Fig. 1). We defined the perimeter of the measured area to include the trunk region, extending from the posterior end of the head to the approximated cloacal area (Fig. 1). Because the cloaca is located on the ventral side and is not directly visible from the dorsal view, its position was approximated using the lateral body width and/or the position of the anal fin. The anal fin, although ventral and normally invisible from above, can be intermittently observed during handling of goldfish, which allowed us to infer the cloaca position. Based on these acquired values, we calculated the circularity of the selected area (Fig. 2 and Table 1). Our approach—though based on visually biased estimations—resembles the way early breeders perceived goldfish and can thus be considered a reasonable proxy for assessing overall body roundness.

From our calculation of the circularity in these strains, it is revealed that the globular body ornamental goldfish, including *Ryukin*, *Oranda*, *Pearl scale*, and *Ranchu* strains, showed higher circularities than their wild-type counterparts (Fig. 2). Of those, *Pearl scale* and *Ryukin* goldfish seem to be more circular than the other globular body ornamental gold-

fish (Fig. 2 and Table 1). We also then compared how the internal anatomical features are different among these ornamental goldfish strains, by observing the section plane of the wild-type, *Ryukin*, *Oranda*, *Pearl scale*, and *Ranchu* goldfish strains (Fig. 3A-E). This observation indicated that the lateral body wall thickness is significantly wider in *Pearl scale* goldfish in comparison with the other goldfish strains (the green and blue bars in Fig. 3A-E). *Pearl scale* goldfish also showed a higher ratio of the lateral body wall thickness of body width than the others (Fig. 3F). Crucially, while *Ryukin* and *Pearl scale* individuals showed quite similar circularity (Fig. 2), their anatomical and histological characteristics seem to be highly differentiated from each other. For instance, the transverse section of *Pearl scale* is clearly more circular than that of *Ryukin* (Fig. 3B,D). Moreover, although body size, age, and feeding conditions varied among individuals due to their maintenance in different aquarium tanks, the whitish appearance of the lateral body wall tissue—likely indicating lipid-rich tissue—in the *Pearl scale* goldfish (indicated by black asterisks in Fig. 3D) suggests that this strain possesses distinct morphological and histological characteristics.

To further investigate the aforementioned possibility, we obtained a set of data consisting of the circularity of the body and anatomical features of the transverse section plane from seven individuals each from wild-type, *Ryukin*, and *Pearl scale* goldfish (Fig. 4 to 6). The comparison of these three strains confirmed that the *Ryukin* and *Pearl scale* goldfish strains exhibit higher circularity, with more *Pearl scale* goldfish individuals having higher circularities than *Ryukin* individuals (Fig. 4). Exceptionally, one of the *Pearl scale* gold-

Table 1. Measured circularity values of five ornamental goldfish

Handle ID	Strain	Area*	Perimeter*	Circularity
Wt-Jp-2019-01	wild-type	99357	1292.941	0.747
Wt-Jp-2019-02	wild-type	116019	1320.087	0.837
Wt-Jp-2019-03	wild-type	102129	1345.216	0.709
Wt-Jp-2019-04	wild-type	107958	1359.783	0.734
Wt-Jp-2015-01	wild-type	122932	1453.877	0.731
Wt-Jp-2015-02	wild-type	83594	1186.909	0.746
Wt-Jp-2015-03	wild-type	55259	933.59	0.797
Wt-Jp-2015-04	wild-type	52059	948.765	0.727
Wt-Jp-2015-05	wild-type	71527	1136.819	0.696
Wt-Tw-2015-01	wild-type	210301	1988.55	0.668
Wt-Tw-2015-02	wild-type	145679	1662.477	0.662
Wt-Tw-2015-03	wild-type	185946	1790.948	0.729
Wt-Tw-2015-04	wild-type	204436	1950.582	0.675
Wt-Tw-2015-05	wild-type	227211	1981.262	0.727
RY-2019-01	Ryukin	156483	1419.801	0.975
RY-2019-02	Ryukin	144149	1373.393	0.96
RY-2019-03	Ryukin	156362	1439.078	0.949
OR-2019-01	Oranda	142041	1399.848	0.911
OR-2019-02	Oranda	119059	1257.733	0.946
OR-2019-03	Oranda	122228	1303.913	0.903
PRL-YT-01	Pearl scale	108198	1181.011	0.975
PRL-YT-02	Pearl scale	157832	1430.106	0.97
PRL-YT-03	Pearl scale	123265	1261.756	0.973
RA-YT-01	Ranchu	86237	1070.774	0.945
RA-YT-02	Ranchu	113144	1275.322	0.874
RA-YT-03	Ranchu	83144	1095.311	0.871

*Area and Perimeter are expressed in pixels as measured directly from digital images.

Fig. 3. Comparison of the transverse sections in five ornamental goldfish. A-E. Transverse section of the wild-type (Wt), *Ryukin* (RY), *Oranda* (OR), *Pearl scale* (PRL), and *Ranchu* (RA) strain. The blue and green lines indicate the internal cavity width (icw) and the lateral body wall thickness (lbw), respectively. F. Ratio between lbw and maximum body width (mbw). Abbreviations: df, dorsal fin; nt, neural tube; plv, pelvic fin. Scale bars = 1cm. Source image available at https://n2t.net/ark:/37281/k5f8z2s55, licensed under CC-BY-NC-SA 4.0, © Kinya G. Ota (2025).

fish individuals has a lower circularity than three *Ryukin* individuals (Fig. 4B2, B3, B7, and C7; these panels are indicated by the black and white asterisks). The relatively low circularity observed in this *Pearl scale* individual is likely attributable to its laterally expanded body shape (Fig. 4). These results further confirmed that there are significant differences between the *Pearl scale* and *Ryukin* goldfish strains.

Next, our results of the transverse section provided consistent and solid results in the differences of the lateral body wall thickness of these examined goldfish individuals (Fig. 5 and 6). The transverse section views in these individuals also indicated that the *Pearl scale* goldfish individuals had a clearly more circular transverse section and showed the largest lateral body wall thickness in comparison with the other strain individuals (Fig. 5). The similar width of the lateral body wall thickness of the wild-type and *Ryukin* strain indicated that although the circularity of the *Ryukin* strain is higher than the wild-type goldfish, this difference stems from the increase of the width of the coelom area in the transverse section view (Fig. 5AB and 6). On the other hand, *Pearl scale* goldfish, although with inter-individual variations, tend to increase the lateral body wall thickness to form the globular body shape (Fig. 6BC).

To clarify further how the body wall and coelom areas changed, the circularities of these structures were measured in transverse section views and compared among the three

Ota *et al.* | Anatomy and Histology in Ornamental Goldfish

strains, including wild-type, *Ryukin*, and *Ranchu* (Fig. 7A). The circularity values of the body surface and coelom in transverse section views indicated significant differences between the wild-type and the two globular body shape goldfish strains. *Ryukin* and *Pearl scale* showed quite similar values, and an increase in the circularity of the coelom occurred in both *Ryukin* and *Pearl scale*, suggesting that the *Pearl scale* goldfish possess a *Ryukin*-like coelom shape while simultaneously exhibiting a thickened lateral body wall (Fig. 6 and 7B). This tendency is also observed in the relationship between body surface circularity in transverse sections and dorsal view circularity (Fig. 7C). These results revealed that the globular body goldfish strains are diverged from the wild-type goldfish in the anatomical features. Moreover, while *Ryukin* and *Pearl scale* exhibit comparable anatomical characteristics, their tissue properties appear to be distinct.

To further examine the similarities and differences at the tissue level between *Ryukin* and *Pearl scale* goldfish, we compared histological features using seven individuals per strain, as described above. We succeeded in observing the lateral area of the trunk region in the transverse section of those individuals at the level of the pelvic fin (Fig. 8). In this observation, we clearly recognized the differences in the ratio between the muscle fibers and adipose tissue in the mid trunk region, as we observed in the transverse hand section views (Fig. 3 and 5).

The trunk muscle region of the *Ryukin* goldfish showed the conventional muscle tissue (Fig. 8A), i.e. we did not detect adipose tissue surrounding the muscle fibers from our analyzed *Ryukin* individuals (0/7 individuals; Fig. 8A). In contrast, the histological section of the *Pearl scale* goldfish individuals consistently showed muscle fibers surrounded by adipose cells (7/7 individuals; Fig. 8B). Thus, unlike other goldfish that altered anatomical proportions without accompanying histological modifications, *Pearl scale* goldfish appears to have achieved a globular body through both anatomical changes and additional histological modifications involving reorganization of muscle fibers and adipose tissue.

Table 2. Measured circularity values of wild-type, *Ryukin*, and *Pearl scale*

Handle ID	strain	mbw*	icw*	lbw	lbw / mbw	d-circ**	s-circ**	c-circ**
2025-0219-Wt-01	wild-type	43.511	32.374	11.137	0.256	0.728	0.785	0.866
2025-0219-Wt-02	wild-type	35.871	29.266	6.604	0.184	0.691	0.781	0.89
2025-0219-Wt-03	wild-type	39.108	27.453	11.655	0.298	0.72	0.771	0.868
2025-0219-Wt-04	wild-type	42.475	30.043	12.432	0.293	0.715	0.804	0.905
2025-0219-Wt-05	wild-type	37.425	27.086	10.338	0.276	0.804	0.771	0.806
2025-0219-Wt-06	wild-type	38.245	28.036	10.209	0.267	0.715	0.764	0.862
2025-0219-Wt-07	wild-type	33.669	25.878	7.791	0.231	0.698	0.735	0.841
2025-0225-RY01	Ryukin	47.309	33.799	13.511	0.286	0.891	0.838	0.909
2025-0225-RY02	Ryukin	53.029	41.698	11.331	0.214	0.961	0.887	0.97
2025-0225-RY03	Ryukin	54.583	38.849	15.734	0.288	0.954	0.912	0.953
2025-0225-RY04	Ryukin	40.597	32.504	8.094	0.199	0.938	0.824	0.951
2025-0225-RY05	Ryukin	41.957	35.029	6.928	0.165	0.891	0.853	0.959
2025-0225-RY06	Ryukin	36.216	25.014	11.201	0.309	0.913	0.839	0.917
2025-0225-RY07	Ryukin	38.158	28.122	10.036	0.263	0.961	0.837	0.925
2025-0225-PRL01	Pearl scale	105.237	64.058	41.18	0.391	0.974	0.972	0.952
2025-0225-PRL02	Pearl scale	82.446	54.173	28.273	0.343	0.969	0.92	0.956
2025-0225-PRL03	Pearl scale	84.777	45.54	39.237	0.463	0.97	0.917	0.973
2025-0225-PRL04	Pearl scale	74.417	47.612	26.806	0.36	0.967	0.911	0.931
2025-0225-PRL05	Pearl scale	71.532	49.345	22.187	0.31	0.974	0.892	0.937
2025-0225-PRL06	Pearl scale	66.755	49.662	17.094	0.256	0.982	0.881	0.936
2025-0225-PRL07	Pearl scale	64.036	40.338	23.698	0.37	0.952	0.915	0.949

*The values of mbw and icw were obtained from image data and are expressed in pixels. **d-circ, s-circ, and c-circ refer to dorsal-, surface-, and coelom-circularity, respectively.

Fig. 4. Comparison of circularity between the wild-type, *Ryukin*, and *Pearl scale*. A. Dorsal view of the wild-type. B. Dorsal view of *Ryukin*. C. Dorsal view of *Pearl scale*. The light blue, red, and magenta colored areas indicate the approximately identified trunk regions. Black asterisks indicate *Ryukin* individuals showing values approaching those of the *Pearl scale*. White asterisk indicates a *Pearl scale* individual exhibiting values close to those of the *Ryukin*. All individuals shown in this figure were obtained by artificial fertilization and reared in our laboratory. The corresponding data are summarized in Table 2. Scale bars = 1cm. Source image available at https://n2t.net/ark:/37281/k5f8z2s55, licensed under CC-BY-NC-SA 4.0, © Kinya G. Ota (2025).

DISCUSSION

Our comparative analyses revealed that ornamental goldfish can be categorized into three groups in their internal anatomical and histological features: wild-, *Ryukin*-, and *Pearl scale* type (Fig. 9). Although *Ryukin* progeny has a globular body shape, their body wall thickness is quite similar to wild-type, with almost equivalent histological features of the lateral body wall tissue (Fig. 9AB). On the other hand, investigated *Pearl scale* goldfish individuals tend to show a wider lateral body wall (Fig. 6, 8 and 9). We also further revealed

Fig. 5. Comparison of the transverse section between the wild-type, *Ryukin*, and *Pearl scale*. A-C. Transverse section of the wild-type (Wt), *Ryukin* (RY), *Pearl scale* (PRL) strains. The blue and green lines indicate the internal cavity width (icw) and the lateral body wall thickness (lbw), respectively. Panel labels correspond to the individual ID in Fig. 4. Numbers of left lower corner in each panel represent circularity and l/m, which is the ratio of the lbw and maximum body width (mbw). White asterisk indicates a *Pearl scale* individual exhibiting values close to those of the *Ryukin*. The corresponding data are summarized in Table 2. Scale bars = 1cm. Source image available at https://n2t.net/ark:/37281/k5f8z2s55, licensed under CC-BY-NC-SA 4.0, © Kinya G. Ota (2025).

Fig. 6. Comparison of circularity and the l/m between the wild-type, *Ryukin*, and *Pearl scale* ornamental goldfish. Light blue, red, and magenta indicate the wild-type, *Ryukin*, and *Pearl scale* goldfish, respectively. The data were derived from Fig. 5 and Table 2.

Fig. 7. Comparison of circularity of body surface and coelom in transverse section between the wild-type, *Ryukin*, and *Pearl scale* ornamental goldfish A. Transverse section images from Fig. 5 with outlines of the body surface and coelom traced. B. Plots of body surface circularity and coelom circularity. Light blue, red, and magenta indicate the wild-type, *Ryukin*, and *Pearl scale* goldfish, respectively. The data were derived from Fig. 5 and Table 2 Source image available at https://n2t.net/ark:/37281/k5f8z2s55, licensed under CC-BY-NC-SA 4.0, © Kinya G. Ota

Fig. 9. Schematic representation of three different trunk histologies of ornamental goldfish. A. Wild-type. B. *Ryukin*-type, C. *Pearl scale* type. Red, yellow, green, and white areas indicate the relative trunk musculoskeletal area, coelom, neural tube, and adipose tissue. Source image available at https://n2t.net/ark:/37281/k5f8z2s55, licensed under CC-BY-NC-SA 4.0, © Kinya G. Ota (2025).

Fig. 8. Histological comparison of *Ryukin* and *Pearl scale* goldfish A–B. Eosin–hematoxylin–Alcian blue-stained sections of the lateral body wall in two globular-bodied ornamental goldfish. Panel numbers correspond to individual IDs shown in Fig. 3. Tissue samples were taken from the region approximately at the level of the pelvic fins, corresponding to the transverse section indicated by the green bars in Fig. 5. Black asterisks indicate muscle fibers surrounded by adipose tissue. The regions containing integumental tissue (epidermis, dermis, scale, and related connective tissues) are labeled in panels B2, B3, B5, and B6 by bracket-shaped bars. Scale bars = 0.1 mm. Source image available at https://n2t.net/ark:/37281/k5f8z2s55, licensed under CC-BY-NC-SA 4.0, © Kinya G. Ota (2025).

that the histological features of *Pearl scale* are highly differentiated from that of the *Ryukin* strain (Fig. 8).

Pearl scale-type ornamental goldfish, along with other related strains, are documented to have been established as early as the Ming dynasty and maintained in ceramic containers traditionally used for fish keeping (Chen 1954). The use of such top-view-oriented containers suggests that morphological selection in ornamental goldfish has historically emphasized dorsal appearance (Fig. 1). From this evidence, it can be inferred that the phenotypic features which can be observed from the lateral side, for example, ratio of the epaxial and hypaxial region, lateral line position, and so on, have not been the initial subject of the selective pressure. Similarly, it is also easily assumed that the internal anatomy, for instance, the volume and circularity of the coelom as well as the lateral body wall thickness, have not been the subject of the attention by the breeders and fanciers (Fig. 3 and 6).

Thus, the currently observed phenotypes are naturally interpreted as a consequence of the selective pressure toward externally visible body shape from the dorsal view, without attention to the internal anatomy and histology. Moreover, it is reasonable to assume that none of the breeders or fanciers have applied any selective pressure specifically to *Pearl scale* goldfish to increase the lateral body wall tissue and its adipose cells. Taken together, *Pearl scale* and other globular body-shaped goldfish might have been under nearly equivalent selective pressures to establish attractive morphological features from the dorsal view. Coincidentally, two distinct histological phenotypic features emerged: an increase of adipose tissue in the lateral body wall of the *Pearl scale* strain, and no such histological modifications in the other globular goldfish like the *Ryukin* strain (Fig. 3 and 6).

To clarify the process underlying the emergence of the globular body shape, we need to further discuss how distinct internal phenotypes could arise, given that all examined goldfish belong to the same interbreeding species, *Carassius auratus*. Throughout breeding history, ornamental goldfish have been extensively interbred by breeders and fanciers with the aim of establishing new strains (Chen 1954; Smartt 2001). The *Ryukin*-type globular body shape, or a tendency toward this morphology, is shared by several modern ornamental goldfish strains, suggesting that interbreeding has occurred among ancestral strains exhibiting the *Ryukin*-type body form. In contrast, the accumulation of adipose tissue appears to be a distinctive morphological feature of the *Pearl*

scale (Fig. 3,5 and 9), which warrants further discussion as a unique histological trait.

In fact, strains derived from hybridization with the *Pearl scale* goldfish are well known among breeders and fanciers, and are commonly documented in books and on websites. For example, according to Ye and Qu (2017), several compound variation strains are introduced, such as the *Pearl scale* body type with a cranial hood (as in the *Oranda* strain; see Fig. 1E). However, Ye and Qu (2017) also mentioned that these strains were established only recently, most likely being consolidated as distinct lineages during the 20th century. Taken together, this evidence suggests that, from the perspective of genetic and evolutionary processes, the globular body shape formed through adipose tissue accumulation in the *Pearl scale* may involve a different genetic background from that of the *Ryukin*-type globular body shape.

At present, due to the lack of intensive and extensive genetic or genomic analyses specifically targeting the globular body of ornamental goldfish, we have no convincing data that directly addresses this issue. Nevertheless, the results of genome sequencing analyses of the *Pearl scale* goldfish may provide useful insights (Kon et al. 2020). According to admixture analyses of multiple goldfish genomes, while the genome sequence of the *Ryukin* and *Oranda* goldfish (including its sub-groups) consists of various populations, that of the *Pearl scale* goldfish appears to consist of a single ancestral population (Kon et al. 2020), suggesting isolated genomic features of this goldfish strain. Moreover, the fact that two independent goldfish genome sequencing projects have not investigated any candidate loci for globular body shape suggests that the responsible genetic factors in such a phenotype may not be easily detected (Chen et al. 2020a; Kon et al. 2020). This, in turn, implies that the globular body shape might result from the accumulation of multiple mutated alleles rather than the action of a single major locus. Therefore, the accumulation of multiple variant alleles is considered to have been important in producing the globular body shape, but the manner of this accumulation likely differed between the *Ryukin*-type and the *Pearl scale*-type. The former seems to have arisen through repeated crossbreeding among multiple lineages, whereas the latter appears to have established through a breeding process that emphasized intrastrain crosses to generate a pure line. Thus, it is reasonable to assume that globular body shape is formed in at least two ways in the ornamental goldfish.

To advance this discussion, it is necessary to specify which phenotypic features are clearly dissimilar between *Ryukin* and *Pearl scale*. Our morphological comparisons had limitations that precluded accurate quantitative evaluation between *Ryukin* and *Pearl scale*, as the methods relied largely on subjective visual inspection. In addition, the transverse section analyzed was limited to a single level (at the level of the pelvic fin) (Fig. 3 and 5). Given these constraints, it is difficult to rigorously quantify how dorsal-view circularity relates to that of the coelom and body surface in transverse sections of *Ryukin* and *Pearl scale*, even though differences were observed in these values (Fig. 7).

Therefore, further analyses will be necessary to determine, in a purely geometric manner independent of subjective recognition, where the differences in overall globularity lie between *Ryukin* and *Pearl scale*. In other words, breeders, fanciers, and many observers including ourselves and early researchers may perceive that *Pearl scale* appears rounder than *Ryukin* (Smartt 2001), but it remains to be clarified whether this perceived roundness of *Pearl scale* is primarily attributable to the shape of the coelom or to the influence of adipose tissue. A purely geometric assessment of such differences will thus remain a task for future studies.

Beyond the geometric differences of these two strains, it is certain that, at the histological level, *Pearl scale* possesses accumulated adipose tissue not observed in *Ryukin*. Taking this fact into account, for convenience, we describe the evolutionary process as if each strain itself responded to the selective pressures imposed by breeders and fanciers. The *Ryukin* strain specialized in expanding the coelom area and thereby achieved a highly rounded body shape without modifying subcutaneous adipose tissue. By contrast, the *Pearl scale* strain accumulated adipose tissue within the muscle layer in addition to expanding the coelom. Since *Ryukin* and *Pearl scale* exhibit visually similar coelom circularity at the transverse section plane (Fig. 7 and 9BC), *Pearl scale* goldfish may share some anatomical features with non-*Pearl scale* strains. Nevertheless, these two globular body shape types appear to rely on distinct developmental systems for adipose tissue, implying that similar external morphologies can arise through different underlying developmental and histogenesis processes, resembling cases of convergent or parallel evolution in nature.

To avoid unnecessary confusion or debate, we emphasize that our observations are not readily comparable across taxa, unlike streamlined fish morphologies. While such similarities point to a resemblance with convergent or parallel evolution, placing our findings within broader theoretical debates on convergence remains complex. For instance, several voluminous teleost species, including pufferfish, boxfish, and lumpfish, may appear superficially comparable to globular goldfish strains (Lampart-Kałużniacka and Heese 2000; Li et al. 2020; Nelson et al. 2016; Van Wassenbergh et al. 2015; Wainwright and Turingan 1997). In reality, some of these voluminous teleosts exhibit modifications in rib position and morphology compared with conventional streamlined teleosts, making them superficially comparable to the relationship between globular body shape ornamental goldfish and wild-type forms (Li et al. 2015). However, their body forms have been shaped by markedly different selective pressures. Thus, although the comparison is not straightforward, our findings invite consideration of alternative approaches that may provide a more appropriate framework for interpreting these cases.

A more fruitful alternative approach, then, is to move beyond purely functional comparisons, and instead examine how morphological convergence can arise alongside divergence at the tissue level. When the focus shifts from viewing morphology in relation to function to considering it in rela-

tion to tissue organization, there appear to be cases where more direct relationship can be drawn. For example, our comparison of *Ryukin* and *Pearl scale* goldfish suggests that, within the same species *Carassius auratus*, the same morphological outcome has been produced under similar selective pressures but through additional histological modifications in the *Pearl scale* goldfish lineage. This perspective may allow us to extend the discussion of convergent and parallel evolution. In particular, the sub-cutaneous adipose tissue of the *Pearl scale* goldfish is comparable with several examples.

One striking example is the blubber of marine mammals. Although blubber represents an adaptation to aquatic life—functioning primarily as thermal insulation—and thus differs fundamentally in function from the adipose tissue observed in *Pearl scale* goldfish, the similarity in histological organization is noteworthy. Blubber is typically stratified into three layers: an outer layer containing epidermis, dermis, and adipose cells; a middle layer composed predominantly of adipose tissue; and an inner layer consisting of a mixture of adipose and muscle (Funasaka et al. 2024; Gómez-Campos et al. 2015; Harper et al. 2008; Hashimoto et al. 2015; Montie et al. 2008; Parry 1949). Remarkably, the sub-cutaneous structure observed in *Pearl scale* goldfish closely resembles this three-layered architecture, although the extent of resemblance varies among individuals (Fig. 8B2, B3, B4, B5, B6). This resemblance implies that the adipose tissue layer may possess the greatest evolutionary plasticity among these three layers, and thus may be the most prone to modification under divergent selective pressures. Moreover, similar patterns of adipose–muscle composition are also observed in humans with obesity, in related laboratory animal models, and in meat-producing livestock such as beef cattle and pigs (Claussnitzer et al. 2015; Craig and Moon 2011; David et al. 2016; Den Broeder et al. 2015; Faillaci et al. 2018; Gandarillas and Bas 2009; Green et al. 2018; Landgraf et al. 2017; Li et al. 2023; McMenamin et al. 2013; Park Seung Ju 2018; Pond 1978, 1992; Smemo et al. 2014; Spurlock and Gabler 2008; Tan and Jiang 2024; Visscher et al. 2017; Zang et al. 2018).

Since the conceptual framework regarding the evolution of adipose tissue is still developing (Ottaviani et al. 2011; Pond 1992), further studies are required to enable comprehensive comparative analyses among the examples mentioned above. Despite the functional differences among these cases, remarkably similar histological features are observed. This is largely attributable to the simple and highly conserved layer structures of dermis, adipose tissue, and muscle fibers across vertebrates. Moreover, the recurrent emergence of similar traits in diverse lineages suggests the presence of deeply conserved developmental mechanisms, as also implied by research demonstrating shared molecular pathways of obesity between zebrafish and mammals (Oka et al. 2010). At the same time, difficulties in identifying the genetic basis of subcutaneous adipose accumulation are evident even in established model species (Claussnitzer et al. 2015; Smemo et al. 2014). Such difficulties may partly explain why no specific loci responsible for the globular body shape phenotype in goldfish have yet been reported, despite multiple large-scale genome sequencing projects (Chen et al. 2020b; Kon et al. 2020).

These histologically simple yet genetically complex characteristics may themselves reflect a key aspect of adipose tissue evolution—namely, that traits such as adipose tissue accumulation are highly plastic and strongly influenced by environmental context (Pond 1992). In other words, adipose deposition can be either advantageous or disadvantageous depending on ecological and selective conditions as has been broadly recognized in evolutionary studies of human obesity; for example, Qasim et al. (2018), Speakman (2013), Wells (2012). The case of the *Pearl scale* goldfish exemplifies this principle: a globular body shape, unlikely to confer any advantage in the natural habitat of *Carassius auratus*, became highly favored under selection for ornamental purposes. Importantly, this morphology appears to have been achieved through distinct developmental modifications, including both enlargement of the body cavity and accumulation of subcutaneous adipose tissue. Moreover, the process of genetic fixation of this morphology might be related with the environmental factors, diet and water conditions. In fact, since we observed well-established ornamental goldfish strains and/or its progenies in this study, their morphologies can be recognized as the consequence of developmental modifications, some of which are already evident even before hatching, that are now likely fixed by many rounds of selections. Still, it remains naturally assumed that the globular phenotypes might show a certain plasticity in the ancestral population of ornamental goldfish. We suggest that further developmental genetic studies of the globular body goldfish will provide valuable insights into how polygenic, environmentally responsive traits evolve under selective pressures, and into the evolutionary dynamics of highly plastic organs and tissues.

ACKNOWLEDGEMENTS

We are grateful to Wen-Hui Su (SHUEN-SHIN Breeding Farm) and You Syu Huang (formerly of the Aquaculture Breeding Institute, Hualian) for their valuable technical advice on goldfish breeding in Taiwan. We also wish to thank the current and former members of the Yilan Marine Research Station, Institute of Cellular and Organismic Biology: the late Hung-Tsai Lee, Chia-Chun Lee, Chihi-Chiang Lee, and Tsai Han Chuan for their assistance with the maintenance of aquarium systems; Shu-Hua Lee and Sky Wu for occasionally providing brine shrimp when our own supply was lacking; Shi-Chieh Liu for designing aquarium systems; and Chi-Fu Hung, Jhih-Hao Wei, and Fei Chu Chen for administrative support. We are further indebted to Konstantin Khalturin for suggesting comparative approaches with pufferfish. We also wish to thank Qichang Ye for generously donating books relevant to this study. In addition, we extend our sincere appreciation to Teng Yu-Feng (Big Wind Technology Co., Ltd.), and F. L. Lu (YuanLi Instrument Co., Ltd.) for their assistance in arranging various detailed and often demanding equipment logistics. Finally, we acknowledge the support of the Taiwan Zebrafish Core Facility at Academia Sinica.

FUNDING

The funding was provided by National Science and Technology Council (Grant No. 112-2311-B-001-033 and 113-2311-B-001 -032 -MY3), Academia Sinica through the Postdoctoral Scholar Program (Grant No. 235g), JSPS KAKENHI (Grant No. JP22K06232), and Takeda Science Foundation (Grant No. 2022036015).

AUTHOR ORCID

ⓘD Kinya G. Ota 太田欽也 https://orcid.org/0000-0002-6306-6790

ⓘD Gembu Abe 阿部玄武 https://orcid.org/0000-0002-9905-8506

ⓘD Chen-Yi Wang 王貞懿 https://orcid.org/0000-0003-4342-5140

ⓘD Ing-Jia Li 李穎佳 https://orcid.org/0000-0001-6407-5181

ⓘD Paul Gerald Layague Sanchez https://orcid.org/0000-0001-6213-8927

AUTHOR CONTRIBUTIONS

Conceptualization: Kinya G. Ota 太田欽也 (KGO). **Data curation**: KGO, Paul Gerald Layague Sanchez (PGLS). **Formal Analysis**: KGO, PGLS. **Funding Acquisition**: KGO, PGLS. **Investigation**: KGO. **Methodology**: KGO, Chen-Yi Wang 王貞懿 (CYW), Ing-Jia Li 李穎佳 (IJL), PGLS. **Project Administration**: KGO. **Resources**: CYW, IJL, KGO. **Supervision**: KGO, PGLS. **Image acquisition**: KGO. **Writing – Original Draft**: KGO. **Writing – Review & Editing**: KGO, Gembu Abe (GA), PGLS, IJL.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

AVAILABILITY OF DATA AND MATERIALS

Data underlying the figures and results presented in this manuscript are available in Depositor at https://n2t.net/ark:/37281/k5f8z2s55.

REFERENCES

Asano, H. and Kubo, Y. 1972. Variations of spinal curvature and vertebral number in goldfish. Japanese Journal of Ichthyology. **19 (4):**223–231. Publisher: The Ichthyological Society of Japan.

Chen, D., Zhang, Q., Tang, W., Huang, Z., Wang, G., Wang, Y., Shi, J., Xu, H., Lin, L., and Li, Z. 2020a. The evolutionary origin and domestication history of goldfish (Carassius auratus). Proceedings of the National Academy of Sciences. **117 (47):**29775–29785. Publisher: National Acad Sciences.

Chen, D., Zhang, Q., Tang, W., Huang, Z., Wang, G., Wang, Y., Shi, J., Xu, H., Lin, L., Li, Z., Chi, W., Huang, L., Xia, J., Zhang, X., Guo, L., Wang, Y., Ma, P., Tang, J., Zhou, G., Liu, M., Liu, F., Hua, X., Wang, B., Shen, Q., Jiang, Q., Lin, J., Chen, X., Wang, H., Dou, M., Liu, L., Pan, H., Qi, Y., Wu, B., Fang, J., Zhou, Y., Cen, W., He, W., Zhang, Q., Xue, T., Lin, G., Zhang, W., Liu, Z., Qu, L., Wang, A., Ye, Q., Chen, J., Zhang, Y., Ming, R., Van Montagu, M., Tang, H., Van de Peer, Y., Chen, Y., and Zhang, J. 2020b. The evolutionary origin and domestication history of goldfish (Carassius auratus). Proceedings of the National Academy of Sciences. **117 (47):**29775–29785. doi: 10.1073/pnas.2005545117. https://doi.org/10.1073/pnas.2005545117. Publisher: Proceedings of the National Academy of Sciences.

Chen, S. C. 1954. A historty of the domestication and the factors of the varietal formation of the common goldfish, Carassius auratus. Act Zoologica Sinica. **6 (2):**89–116.

Chen, S. C. 1956. A history of the domestication and the factors of the varietal formation of the common goldfish, Carassius auratus. Scientia Sinica. **5:**287–321.

Claussnitzer, M., Dankel, S. N., Kim, K.-H., Quon, G., Meuleman, W., Haugen, C., Glunk, V., Sousa, I. S., Beaudry, J. L., Puviindran, V., Abdennur, N. A., Liu, J., Svensson, P.-A., Hsu, Y.-H., Drucker, D. J., Mellgren, G., Hui, C.-C., Hauner, H., and Kellis, M. 2015. *FTO* Obesity Variant Circuitry and Adipocyte Browning in Humans. New England Journal of Medicine. **373 (10):**895–907. doi: 10.1056/NEJMoa1502214. https://www.nejm.org/doi/full/10.1056/NEJMoa1502214. _eprint: https://www.nejm.org/doi/pdf/10.1056/NEJMoa1502214.

Craig, P. M. and Moon, T. W. 2011. Fasted Zebrafish Mimic Genetic and Physiological Responses in Mammals: A Model for Obesity and Diabetes? Zebrafish. **8 (3):**109–117. doi: 10.1089/zeb.2011.0702. http://www.liebertpub.com/doi/10.1089/zeb.2011.0702.

David, C. J., Veena, R., and Kumaresan, G. 2016. High cholesterol diet induces obesity in zebrafish. PloS one. **8:**e66970. https://mahendrapublications.com/Document/MP787017_Manuscript_2016_09_12_0330.pdf.

Den Broeder, M. J., Kopylova, V. A., Kamminga, L. M., and Legler, J. 2015. Zebrafish as a Model to Study the Role of Peroxisome Proliferating-Activated Receptors in Adipogenesis and Obesity. PPAR Research. **2015:**1–11. doi: 10.1155/2015/358029. http://www.hindawi.com/journals/ppar/2015/358029/.

Faillaci, F., Milosa, F., Critelli, R. M., Turola, E., Schepis, F., and Villa, E. 2018. Obese zebrafish: A small fish for a major human health condition. Animal Models and Experimental Medicine. **1 (4):**255–265. doi: 10.1002/ame2.12042. https://onlinelibrary.wiley.com/doi/10.1002/ame2.12042.

Funasaka, N., Suzuki, M., Hosono, M., Shindo, H., Kawamura, K., Inamori, D., and Yoshioka, M. 2024. Blubber biopsy in common bottlenose dolphins using a novel biopsy puncher: Evaluation of the impact on living individuals and possibility of applications in cetacean research. Journal of Experimental Zoology Part A: Ecological and Integrative Physiology. **341 (5):**499–508. doi: https://doi.org/10.1002/jez.2797. https://onlinelibrary.wiley.com/doi/abs/10.1002/jez.2797. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/jez.2797.

Gandarillas, M. and Bas, F. 2009. The domestic pig (Sus scrofa domestica) as a model for evaluating nutritional and metabolic consequences of bariatric surgery practiced on morbid obese humans. Ciencia e investigación agraria. **36:**163 – 176. http://www.scielo.cl/scielo.php?script=sci_arttext&pid=S0718-16202009000200002&nrm=iso. Publisher: scielocl.

Goedhart, J. 2021. Superplotsofdata—a web app for the transparent display and quantitative comparison of continuous data from different conditions. Molecular biology of the cell. **32 (6):**470–474.

Green, A. J., Hoyo, C., Mattingly, C. J., Luo, Y., Tzeng, J.-Y., Murphy, S. K., Buchwalter, D. B., and Planchart, A. 2018. Cadmium exposure increases the risk of juvenile obesity: a human and zebrafish comparative study. International Journal of Obesity. **42 (7):**1285–1295. https://www.nature.com/articles/s41366-018-0036-y. Publisher: Nature Publishing Group UK London.

Gómez-Campos, E., Borrell, A., Correas, J., and Aguilar, A. 2015. Topographical variation in lipid content and morphological structure of the blubber in the striped dolphin. Scientia Marina.

79 (2):189–197. doi: 10.3989/scimar.04093.25A. https://scientiamarina.revistas.csic.es/index.php/scientiamarina/article/view/1585.

Harper, C., McLellan, W., Rommel, S., Gay, D., Dillaman, R., and Pabst, D. 2008. Morphology of the melon and its tendinous connections to the facial muscles in bottlenose dolphins (Tursiops truncatus). Journal of Morphology. **269 (7):**820–839. doi: https://doi.org/10.1002/jmor.10628. https://onlinelibrary.wiley.com/doi/abs/10.1002/jmor.10628. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/jmor.10628.

Hashimoto, O., Ohtsuki, H., Kakizaki, T., Amou, K., Sato, R., Doi, S., Kobayashi, S., Matsuda, A., Sugiyama, M., Funaba, M., Matsuishi, T., Terasawa, F., Shindo, J., and Endo, H. 2015. Brown Adipose Tissue in Cetacean Blubber. PLOS ONE. **10 (2):**1–14. doi: 10.1371/journal.pone.0116734. https://doi.org/10.1371/journal.pone.0116734. Publisher: Public Library of Science.

Hervey, G. F. and Hems, J. 1948. The goldfish. Batchworth Press.

Janvier, P. 1996. Early vertebrates. Number QE851 J36. Oxford University Press. Oxford.

Kardong, K. V. 2006. Vertebrates: comparative anatomy, function, evolution. Number QL805 K35 2006. McGraw-Hill. New York.

Kon, T., Omori, Y., Fukuta, K., Wada, H., Watanabe, M., Chen, Z., Iwasaki, M., Mishina, T., Matsuzaki, S.-i., Yoshihara, D., Arakawa, J., Kawakami, K., Toyoda, A., Burgess, S. M., Noguchi, H., and Furukawa, T. 2020. The Genetic Basis of Morphological Diversity in Domesticated Goldfish. Current Biology. **30 (12):**2260–2274.e6. doi: 10.1016/j.cub.2020.04.034. https://www.sciencedirect.com/science/article/pii/S0960982220305480.

Lampart-Kałużniacka, M. and Heese, T. 2000. Morphological characteristics of South Baltic lumpfish, Cyclopterus lumpus L., 1758. Acta Ichthyologica et Piscatoria. **30:**99–110. https://aiep.pensoft.net/article/25735/download/pdf/. Publisher: Pensoft Publishers.

Landgraf, K., Schuster, S., Meusel, A., Garten, A., Riemer, T., Schleinitz, D., Kiess, W., and Kőrner, A. 2017. Short-term overfeeding of zebrafish with normal or high-fat diet as a model for the development of metabolically healthy versus unhealthy obesity. BMC Physiology. **17 (1):**4. doi: 10.1186/s12899-017-0031-x. http://bmcphysiol.biomedcentral.com/articles/10.1186/s12899-017-0031-x.

Le Verger, K., Kűng, L. C., Fabre, A.-C., Schmelzle, T., Wegmann, A., and Sánchez-Villagra, M. R. 2024. Goldfish phenomics reveals commonalities and a lack of universality in the domestication process for ornamentation. Evolution Letters. **8 (6):**774–786. doi: 10.1093/evlett/qrae032. https://doi.org/10.1093/evlett/qrae032. _eprint: https://academic.oup.com/evlett/article-pdf/8/6/774/60781621/qrae032.pdf.

Li, I. J., Chang, C. J., Liu, S. C., Abe, G., and Ota, K. G. 2015. Postembryonic staging of wild-type goldfish, with brief reference to skeletal systems. Developmental Dynamics. **244 (12):**1485–1518. doi: 10.1002/dvdy.24340.

Li, I.-J., Lee, S.-H., Abe, G., and Ota, K. G. 2019. Embryonic and post-embryonic development of the ornamental twin-tail goldfish. Developmental Dynamics. **(23):**1–33. doi: 10.1002/dvdy.15.

Li, R., Xiao, Q., Liu, Y., Li, L., and Liu, H. 2020. Computational investigation on a self-propelled pufferfish driven by multiple fins. Ocean Engineering. **197:**106908. doi: 10.1016/j.oceaneng.2019.106908. https://www.sciencedirect.com/science/article/pii/S002980181930993X.

Li, X., Ge, G., Song, G., Li, Q., and Cui, Z. 2023. Effects of nutritionally induced obesity on metabolic pathways of zebrafish. International Journal of Molecular Sciences. **24 (3):**1850. https://www.mdpi.com/1422-0067/24/3/1850. Publisher: MDPI.

Liem, K. F., Bemis, W. E., Walker, W. F., and Grande, L. 2001. Functional anatomy of the vertebrates: an evolutionary perspective. Harcourt College Publishers. New York. Publisher: Harcourt College Publishers New York.

Matsui, Y. 2006. [Research on Goldfish: From Science and Hobby]. [Seizando Shoten]. [reprint edition] edition. https://ndlsearch.ndl.go.jp/books/R100000002-I000008402415. In Japanese.

McMenamin, S. K., Minchin, J. E., Gordon, T. N., Rawls, J. F., and Parichy, D. M. 2013. Dwarfism and Increased Adiposity in the gh1 Mutant Zebrafish vizzini. Endocrinology. **154 (4):**1476–1487. doi: 10.1210/en.2012-1734. https://academic.oup.com/endo/article/154/4/1476/2423447.

Montie, E. W., Garvin, S. R., Fair, P. A., Bossart, G. D., Mitchum, G. B., McFee, W. E., Speakman, T., Starczak, V. R., and Hahn, M. E. 2008. Blubber morphology in wild bottlenose dolphins (Tursiops truncatus) from the Southeastern United States: Influence of geographic location, age class, and reproductive state. Journal of Morphology. **269 (4):**496–511. doi: https://doi.org/10.1002/jmor.10602. https://onlinelibrary.wiley.com/doi/abs/10.1002/jmor.10602. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/jmor.10602.

Nelson, J. S., Grande, T. C., and Wilson, M. V. 2016. Fishes of the World. John Wiley & Sons. Hoboken, New Jersey.

Oka, T., Nishimura, Y., Zang, L., Hirano, M., Shimada, Y., Wang, Z., Umemoto, N., Kuroyanagi, J., Nishimura, N., and Tanaka, T. 2010. Diet-induced obesity in zebrafish shares common pathophysiological pathways with mammalian obesity. BMC Physiology. **10 (1):**21. doi: 10.1186/1472-6793-10-21. https://bmcphysiol.biomedcentral.com/articles/10.1186/1472-6793-10-21.

Ota, K. G. 2021. Goldfish Development and Evolution. Springer. Singapore. 1 edition. https://doi.org/10.1007/978-981-16-0850-6.

Ota, K. G. and Abe, G. 2016. Goldfish morphology as a model for evolutionary developmental biology. Wiley Interdisciplinary Reviews: Developmental Biology. **5 (3):**272–295. doi: 10.1002/wdev.224. http://doi.wiley.com/10.1002/wdev.224.

Ota, K. G., Abe, G., Wang, C.-Y., Li, I.-J., Sanchez, P. G. L., and Chi, T.-C. 2025. Evolutionary Insights into Muscle Fiber Distribution in the Twin Tails of Ornamental Goldfish. Zoological Studies. **64 (4):**1–22.

Ottaviani, E., Malagoli, D., and Franceschi, C. 2011. The evolution of the adipose tissue: A neglected enigma. General and Comparative Endocrinology. **174 (1):**1–4. doi: https://doi.org/10.1016/j.ygcen.2011.06.018. https://www.sciencedirect.com/science/article/pii/S0016648011002644.

Park Seung Ju, Beak Seok-Hyeon, J. D. J. S. K. S. Y. J. I. H. P. M. Y. K. H. J. F. D. M. N. S. W. Y. S. P. B. M. 2018. Genetic, management, and nutritional factors affecting intramuscular fat deposition in beef cattle — A review. Asian-Australas J Anim Sci. **31 (7):**1043–1061. doi: `10.5713/ajas.18.0310`. `http://www.animbiosci.org/journal/view.php?number=24006`. _eprint: http://www.animbiosci.org/journal/view.php?number=24006.

Parry, D. A. 1949. The Structure of Whale Blubber, and a Discussion of its Thermal Properties. Journal of Cell Science. **s3-90 (9):**13–25. doi: `10.1242/jcs.s3-90.9.13`. `https://doi.org/10.1242/jcs.s3-90.9.13`. _eprint: https://journals.biologists.com/jcs/article-pdf/s3-90/9/13/3311841/joces_s3-90_9_13.pdf.

Pond, C. M. 1978. Morphological Aspects and the Ecological and Mechanical Consequences of Fat Deposition in Wild Vertebrates. Annual Review of Ecology and Systematics. **9:**519–570. `http://www.jstor.org/stable/2096760`. Publisher: Annual Reviews.

Pond, C. M. 1992. An evolutionary and functional view of mammalian adipose tissue. Proceedings of the Nutrition Society. **51 (3):**367–377. doi: `10.1079/PNS19920050`.

Qasim, A., Turcotte, M., de Souza, R. J., Samaan, M. C., Champredon, D., Dushoff, J., Speakman, J. R., and Meyre, D. 2018. On the origin of obesity: identifying the biological, environmental and cultural drivers of genetic risk among human populations. Obesity Reviews. **19 (2):**121–149. doi: `https://doi.org/10.1111/obr.12625`. `https://onlinelibrary.wiley.com/doi/abs/10.1111/obr.12625`.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. 2012. Fiji: an open-source platform for biological-image analysis. Nature methods. **9 (7):**676–682.

Smartt, J. 2001. Goldfish Varieties and Genetics: Handbook for Breeders. John Wiley & Sons.

Smemo, S., Tena, J. J., Kim, K.-H., Gamazon, E. R., Sakabe, N. J., Gómez-Marín, C., Aneas, I., Credidio, F. L., Sobreira, D. R., Wasserman, N. F., Lee, J. H., Puviindran, V., Tam, D., Shen, M., Son, J. E., Vakili, N. A., Sung, H.-K., Naranjo, S., Acemel, R. D., Manzanares, M., Nagy, A., Cox, N. J., Hui, C.-C., Gomez-Skarmeta, J. L., and Nóbrega, M. A. 2014. Obesity-associated variants within FTO form long-range functional connections with IRX3. Nature. **507 (7492):**371–375. doi: `10.1038/nature13138`. `https://doi.org/10.1038/nature13138`.

Speakman, J. R. 2013. Evolutionary perspectives on the obesity epidemic: Adaptive, maladaptive, and neutral viewpoints. Annual Review of Nutrition. **33 (Volume 33, 2013):**289–317. doi: `https://doi.org/10.1146/annurev-nutr-071811-150711`. `https://www.annualreviews.org/content/journals/10.1146/annurev-nutr-071811-150711`.

Spurlock, M. E. and Gabler, N. K. 2008. The Development of Porcine Models of Obesity and the Metabolic Syndrome1,2. The Journal of Nutrition. **138 (2):**397–402. doi: `https://doi.org/10.1093/jn/138.2.397`. `https://www.sciencedirect.com/science/article/pii/S0022316622095578`.

Tan, Z. and Jiang, H. 2024. Molecular and Cellular Mechanisms of Intramuscular Fat Development and Growth in Cattle. International Journal of Molecular Sciences. **25 (5)**. doi: `10.3390/ijms25052520`. `https://www.mdpi.com/1422-0067/25/5/2520`.

Tsai, H.-Y., Chang, M., Liu, S.-C., Abe, G., and Ota, K. G. 2013. Embryonic development of goldfish (Carassius auratus): a model for the study of evolutionary change in developmental mechanisms by artificial selection. Developmental dynamics : an official publication of the American Association of Anatomists. **242 (11):**1262–83. doi: `10.1002/dvdy.24022`. `http://www.ncbi.nlm.nih.gov/pubmed/23913853`.

Van Wassenbergh, S., van Manen, K., Marcroft, T. A., Alfaro, M. E., and Stamhuis, E. J. 2015. Boxfish swimming paradox resolved: forces by the flow of water around the body promote manoeuvrability. Journal of The Royal Society Interface. **12 (103):**20141146. doi: `10.1098/rsif.2014.1146`. `https://doi.org/10.1098/rsif.2014.1146`. Publisher: Royal Society.

Visscher, P. M., Wray, N. R., Zhang, Q., Sklar, P., McCarthy, M. I., Brown, M. A., and Yang, J. 2017. 10 Years of GWAS Discovery: Biology, Function, and Translation. The American Journal of Human Genetics. **101 (1):**5–22. doi: `https://doi.org/10.1016/j.ajhg.2017.06.005`. `https://www.sciencedirect.com/science/article/pii/S0002929717302409`.

Wainwright, P. C. and Turingan, R. G. 1997. EVOLUTION OF PUFFERFISH INFLATION BEHAVIOR. Evolution. **51 (2):**506–518. doi: `10.1111/j.1558-5646.1997.tb02438.x`. `https://doi.org/10.1111/j.1558-5646.1997.tb02438.x`.

Wells, J. C. K. 2012. The evolution of human adiposity and obesity: where did it all go wrong? Disease Models & Mechanisms. **5 (5):**595–607. doi: `10.1242/dmm.009613`. `https://doi.org/10.1242/dmm.009613`. _eprint: https://journals.biologists.com/dmm/article-pdf/5/5/595/1864027/595.pdf.

Ye, Q. and Qu, L. 2017. Goldfish of China: Descriptions and Illustrations of Diversed Goldfish in China. The Straits Publishing & Distribution Group. Fujian, China.

Zang, L., Maddison, L. A., and Chen, W. 2018. Zebrafish as a model for obesity and diabetes. Frontiers in cell and developmental biology. **6:**91. `https://www.frontiersin.org/journals/cell-and-developmental-biology/articles/10.3389/fcell.2018.00091/full`. Publisher: Frontiers Media SA.