Opposing Copy Number Variation Dynamics Shape Adaptation to Glucose and Galactose in

Diploid Yeast

Prachitha Nagendra[1], Saket Choudhary[2], Supreet Saini[*,1]

[1]Department of Chemical Engineering, Indian Institute of Technology Bombay, Mumbai, India 400 076

[2]Koita Centre for Digital Health, Indian Institute of Technology Bombay, Mumbai, India 400 076

* Corresponding Author. Email: saini@che.iitb.ac.in, Phone: +91 22 2576 7216

Running Title: *CNVs Drive Yeast Adaptation*

Abstract

How do structural variants and point mutations jointly shape long-term adaptation in constant

environments? We evolved six replicate diploid *Saccharomyces cerevisiae* populations for 1,200

generations in glucose or galactose and quantified fitness every 300 generations under controlled

history manipulations. Whole-genome sequencing across four time points per population revealed

that adaptation was dominated by copy-number variations (CNVs), with single-nucleotide

polymorphisms (SNPs) rare across lines and time. CNVs were strongly environment-specific: glucose-

evolved populations showed early, widespread telomeric deletions encompassing carbon-use

modules, followed by a reduction of deletion burden and appearance of compensatory duplications;

galactose-evolved populations accumulated persistent telomeric and sub-telomeric duplications.

Many of the same genes were targeted in opposite directions across environments (deleted in

glucose, duplicated in galactose), linking structural remodeling to divergent physiological strategies.

History-dependent fitness costs were pronounced early for glucose-evolved lines after galactose

preculture but diminished by ~900 generations, whereas costs deepened in galactose-evolved lines

after glucose preculture. A linear modeling framework identified ancestral fitness as the strongest

predictor of evolved fitness, with evolutionary time contributing modestly and preconditioning/recovery

exerting subtle, time-dependent effects. Together, these results argue that predictable, environment-

specific CNV trajectories, rather than SNP accumulation, underlie the bulk of adaptation, and that

structural variation operates within constraints imposed by the starting genotype and regulatory

architecture.

Introduction

Natural environments are rarely constant[1]. Organisms frequently encounter transient changes in nutrient availability, temperature, or biotic interactions, which require populations not only to adapt to their current environment but also to maintain resilience against future fluctuations[2–4]. A central question in evolutionary biology is whether the capacity to tolerate such variability is encoded in the genetic architecture of populations, even when they have evolved in a single, stable environment[5–7]. Understanding these constraints is key to predicting evolutionary trajectories[8–11].

Experimental evolution in microbes provides a tractable system to address these questions, as populations can be propagated under controlled conditions, and phenotypic and genetic changes can be monitored over hundreds of generations[12–16]. In this study, we evolved twelve populations of diploid *Saccharomyces cerevisiae* (*S. cerevisiae*) in either glucose or galactose for 1,200 generations, environments that differ in metabolic requirements and gene regulation. In recent years, many studies have examined the physiological responses and adaptive strategies of yeast in glucose and galactose environments[17–24]. By assaying fitness in the home environment, both when sub-cultured from home environment and when brought after growth in the alternative environment, we quantify specialization, pleiotropic costs, and history-dependent effects.

While the role of single-nucleotide polymorphisms (SNPs) in driving adaptation has been often studied, the role of copy number variations (CNVs), including gene duplications and deletions, is relatively less understood[14,25–32]. Recent work has shown that although CNVs contribute to adaptive traits across taxa, their role is especially clear in microbial systems[28,31,33–37]. In *Saccharomyces cerevisiae*, for example, duplications at the *CUP1* locus confer copper resistance, while industrial strains repeatedly acquire CNVs that enhance tolerance to toxic byproducts and improve metabolic efficiency[38]. More broadly, recurrent CNVs in transporter and metabolic genes have been documented across fungi and pathogens, highlighting their role in rapid adaptation, ecological specialization, and environment-specific trade-offs[39–44]. Despite this, the contribution of CNVs to evolutionary responses remains underexplored, particularly in the context of fluctuating or contrasting environments[32,45].

Here, we systematically track CNVs and SNPs in glucose- and galactose-evolved populations across four time points, combining genome sequencing with phenotypic assays. This design allows us to ask: (1) How does adaptation in an environment affect fitness in an alternative environment across time? (2) Which genomic changes underlie specialist and generalist phenotypes? (3) Are there recurrent or predictable patterns of CNVs that correspond to fitness trajectories?

Our results show that adaptation is primarily driven by CNVs rather than SNPs. Glucose-evolved populations initially specialize, displaying early fitness costs that are later mitigated, while galactose-evolved populations accumulate persistent duplications that reinforce specialization. Many of the same genes are targeted in opposing ways in the two environments, revealing contrasting strategies

72   of genomic remodeling in similar environments. Overall, these findings highlight the central role of

73   CNVs in shaping predictable adaptive trajectories and environment-dependent trade-offs, providing

74   insight into the genetic architecture underlying specialization and generalist evolution.

Results

Adaptive dynamics and genomic architecture of glucose- and galactose-evolved populations.
We evolved six replicate diploid *Saccharomyces cerevisiae* populations in either 0.5% glucose or
0.5% galactose for 1,200 generations. Fitness was assayed every 300 generations, while whole-
genome sequencing of all six glucose- and six galactose-adapted populations, each at four time
points was performed to identify the underlying genetic basis of adaptation **(Figure 1A)**.

In both environments, populations initially experience a fitness increase in their home environment,
which later fluctuated mildly. Upon assaying fitness by preculturing in the same medium, we see that
by 1,200 generations, glucose-adapted populations grew ~20–30% faster than their ancestor in
glucose, while galactose-adapted populations improved ~15–25% in galactose **(Figure 1B,
Supplement Figure S1)**.

Fitness changes correspond to environment-specific copy number variation dynamics.
Despite the different selective environments, a consistent pattern emerged: single-nucleotide
polymorphisms (SNPs) were rare across all lines and time points **(Figure 1C, Supplement Sheet 1)**.
Instead, the dominant mode of genomic change was copy-number variation (CNV), which occurred
repeatedly and independently in multiple replicate lines **(Figure 1D-E, Supplement Sheet 2)**. Thus,
adaptation in both glucose and galactose populations was primarily driven by copy number variations
rather than point mutations.

Fitness gains were accompanied by widespread CNVs whose signatures differed sharply between the
two environments. Overall, glucose lines carried roughly equal numbers of deletions and duplications,
while galactose lines accumulated several-fold more duplications than deletions, a bias that remained
consistent across all four time points **(Supplement Sheet 3)**.

However, duplications and deletions exhibited strong contrasts, depending on the environment in
which the populations were evolved in. In glucose-evolved populations, deletions were frequent,
particularly in telomeric regions of all chromosomes, except chromosome I and II **(Figure 2,
Supplement Figure S2)**. The extent of these deletions was most pronounced in the early intervals
(first 300 generations) but declined with further evolution, suggesting that early adaptive gains
involved large number of deletions that were later refined or compensated by subsequent changes.

In contrast, galactose-evolved populations exhibited widespread telomeric and sub-telomeric
duplications **(Figure 3. Supplement Figure S3)** except in chromosome II. These duplications were
evident from as early as 300 generations and were maintained or expanded throughout 1,200
generations, indicating persistent selective benefit in the galactose environment. Strikingly, nearly 100
genes that were recurrently deleted in glucose-evolved lines were found duplicated in galactose-
evolved lines during some stage of the evolution experiment, highlighting the opposing selective

pressures acting on the same genetic loci as populations adapted in the two environments **(Figure 4,**
Supplement Figure S4).

Specialization trajectories are mirrored in genomic remodeling.
Adaptation in glucose increases fitness. However, we asked if there are particular costs associated
with this adaptation. Specifically, we investigated the cost of adaptation in the ability of the cells to
switch environments and exhibit growth. To quantify this, we compared growth of the glucose-evolved
lines and the ancestor in glucose, when the cells were first pre-cultured in galactose. Glucose-evolved
populations showed a strong penalty or growth defect at 300 generations because of prior exposure
to galactose, consistent with cost of specialization on glucose **(Figure 5)**. However, this penalty
declined sharply by generation 900, reaching nearly zero, before increasing slightly again later in the
experiment. This dynamic trajectory suggested a transient phase of specialization followed by the
acquisition of mutations conferring generalist performance.

The genomic data are consistent with this interpretation. In glucose lines, early adaptation coincided
with extensive telomeric deletions, including in regions encompassing genes involved in galactose
metabolism and alternative carbon source utilization. Pathway-level analyses revealed frequent
targeting of galactose metabolism, pyruvate metabolism, fructose and mannose metabolism, and the
glyoxylate pathway. **(Supplement Figure S5)** These deletions likely conferred rapid specialization to
glucose but incurred fitness costs when preculture history involved galactose. Later in evolution,
however, the extent of deletions decreased, suggesting the appearance of compensatory CNVs
(including duplications) that may have restored broader metabolic flexibility and enabled the observed
decline in fitness penalties.

In contrast, galactose-evolved populations showed steadily increasing penalties across the 1,200
generations, consistent with a trajectory of deepening specialization **(Figure 5)**. The underlying CNV
patterns reinforced this phenotype: galactose lines accumulated persistent duplications in telomeric
and subtelomeric regions, often in loci involved in glycolysis/gluconeogenesis, starch and sucrose
metabolism, and galactose metabolism. **(Supplement Figure S6)** Rather than being compensated,
these duplications were maintained and amplified over time, reinforcing a genetic architecture that
promoted strong specialization and increasing costs of cross-environment exposure.

Physiological history dependence and its genomic correlates.
We further tested the influence of short-term growth history on fitness. Galactose-evolved evolved
populations grown briefly in glucose (~15 generations) and then returned to galactose displayed
reduced growth rates in galactose, but this penalty was mitigated when, following growth in glucose,
cells were re-exposed to galactose for 7 or 15 generations **(Figure 6A)**. This plasticity suggests that
the specialized physiology of galactose-adapted lines is sensitive to immediate history but can be
restored with sufficient exposure to galactose. Genomic signatures provide a plausible explanation:
the duplications enriched in galactose-evolved lines were heavily biased toward genes encoding

metabolic enzymes and transporters, creating a cellular state optimized for galactose but disrupted by
prior glucose exposure.

Glucose-evolved populations behaved differently. During the first 600 generations, the duration of
glucose preculture (7 vs. 15 generations) influenced measured growth rates, consistent with
continued sensitivity to short-term history **(Figure 6B, Supplement Figure S6)**. However, beyond
generation 600, this effect disappeared: fitness became independent of preculture duration. This
history-independence parallels the genomic trend in which extensive deletions present early in
evolution diminished later, with glucose lines acquiring duplications that likely stabilized metabolism
and reduced sensitivity to environmental history.

Predictability of evolutionary outcomes
To investigate the contributors to fitness of the evolved populations in different conditions, we
analyzed our dataset using a linear model that incorporated ancestor fitness, preconditioning
environment, recovery duration, evolutionary time, and their interactions. ANOVA revealed that the
fitness of the ancestral strain was by far the strongest predictor of evolved fitness ($\eta^2 \approx 0.71$, $p <$
10^{-62}), demonstrating that the initial genotype largely sets the baseline for subsequent adaptation.
Evolutionary time also contributed significantly to fitness changes ($\eta^2 \approx 0.22$, $p < 10^{-28}$), consistent
with the gradual accumulation of adaptive changes observed in both glucose- and galactose-evolved
lines.

Other main effects, including preconditioning environment and recovery duration, were individually
non-significant. However, modestly significant interactions with evolutionary time were detected
(preconditioning × evolution: $p \approx 0.0014$; recovery × evolution: $p \approx 0.027$), indicating that
environmental context can subtly modulate the pace or trajectory of adaptation. Higher-order
interactions contributed minimally to overall variance in fitness.

Linear regression analysis further supported these conclusions. The coefficient for ancestral fitness
was large and highly significant ($\beta = 1.324$, $p < 0.001$), whereas the effect of evolutionary time per
generation was small but detectable ($\beta = 8.7 \times 10^{-5}$, $p < 0.001$). Interaction terms were largely
negligible, emphasizing that while environmental history and short-term physiological conditions can
influence fitness to a limited extent, the initial genetic background dominates subsequent evolutionary
outcomes.

These findings suggest that the evolutionary response in our experiment is highly predictable at the
level of gross fitness changes: the starting genotype establishes a strong constraint on the magnitude
and direction of adaptation, while environmental factors such as preconditioning and recovery
modulate the response only subtly. When interpreted alongside our genomic analyses, these results
imply that predictable structural changes, particularly environment-specific CNVs in key metabolic loci,

act within the bounds set by the ancestral genotype to produce incremental but repeatable fitness
gains.
196
Opposing genomic strategies underlie divergent adaptive outcomes.
Taken together, our phenotypic and genomic analyses reveal contrasting strategies of adaptation in
glucose and galactose. Glucose populations undergo an early phase of specialization marked by
widespread deletions of galactose- and carbon-metabolism loci, which is later tempered by
compensatory CNVs that restore generalist growth. In contrast, galactose populations accumulate
persistent duplications in many of the same loci, reinforcing specialization and maintaining high costs
of cross-environment preculture. The observation that nearly 100 genes are deleted in glucose but
duplicated in galactose underscores that the two environments exert opposing selective pressures on
overlapping genetic targets.
206
207 Thus, specialization appears to be a common first step in adaptation, but only in glucose do
208 subsequent genomic changes enable escape toward generalist performance. In galactose, persistent
209 duplication-driven remodeling enforces continued specialization, locking populations into a trajectory
210 of increasing costs. Interestingly, ancestor fitness was found to be a strong predictor of evolutionary
211 change in our experiments and analysis.
212

Discussion.

Across 1,200 generations in single-carbon environments, adaptation of diploid *S. cerevisiae* was dominated by copy number variations rather than point mutations. Glucose-evolved populations showed early, extensive telomeric deletions associated with history-dependent fitness costs after galactose preculture; these costs diminished as evolution proceeded. In contrast, galactose-evolved populations accumulated persistent telomeric and subtelomeric duplications that reinforced specialization. Notably, many of the same loci were targeted in opposite directions across environments (deleted in glucose, duplicated in galactose). Together, these results indicate that environment-specific CNVs, more than SNPs, underpin predictable phenotypic trajectories and trade-offs.

CNVs and SNPs differ sharply in both baseline mutation rates and context dependence, which helps explain why copy number variants dominate our experiment. In budding yeast, the single-nucleotide mutation (SNM) rate is of the order of $\sim 10^{-10}$ per site per generation (genome-wide $\sim 10^{-3}$ per genome per generation) and can vary several-fold with environment, but remains comparatively low[46,47]. By contrast, structural events occur orders of magnitude more frequently per affected locus, with rates that depend on ploidy, growth state, and local sequence architecture; for example, tandem-array changes at CUP1 can occur near 10^{-7} per division, and LOH rates commonly exceed base-substitution rates and shift with ploidy and growth conditions[48–50].

More broadly, our findings position CNVs as first-order levers of evolvability in microbes[31,33]. Parallel yet contrasting CNV signatures emerged in the two environments, i.e., mixed deletions/duplications in glucose versus duplication-biased remodeling in galactose. These changes to the genome likely map onto opposite physiological strategies with environment-specific trade-offs. The concentration of changes in telomeric regions and enrichment for transport/metabolic modules suggest that dosage tuning of peripheral genomic neighborhoods provides a rapid route to adaptation when nutrient regimes differ[51–54].

Second, the predictability and constraints of evolution are evident in our data: a linear modeling framework shows that ancestral fitness is the strongest determinant of evolved fitness, while evolutionary time contributes modestly and preconditioning/recovery exert only subtle, time-dependent effects[5,55–58]. This indicates that structural remodeling proceeds within bounds set by the starting genotype and network architecture, yielding incremental but repeatable fitness gains.

Finally, these results motivate targeted tests and applications. Mechanistically, dissecting causality and reversibility of specific CNVs (engineered gain/loss, LOH dynamics) can clarify how populations exit specialist states[33,59–63]. Ecologically and industrially, controlled fluctuation schedules could be used to steer CNV trajectories toward desired phenotypes (e.g., fermentation robustness, drug tolerance control)[64–66]. Methodologically, combining barcoded lineage tracking with long-read assemblies should resolve CNV architectures over time and link structural changes to fitness under

253   explicit history manipulations, including the observed loss of history dependence in glucose lines after

254   ~600 generations[15,67–69].

Methods

Strains used and experimental evolution.

A diploid SK1 (*a/α*) strain was generated by mating two isogenic haploids:

- *SK1 MATa ARS314::TRP1 ho::LYS2 lys2 ura3 leu2::hisG his3::hisG trp1::hisG*
- *SK1 MATα ARS314::LEU2 ho::LYS2 lys2 ura3 leu2::hisG his3::hisG trp1::hisG*

For each evolution experiment, the diploid strain was streaked from frozen stock onto YPD agar plates. A single colony was inoculated into glycerol–lactate medium (3 mL 100% glycerol, 2 mL 40% lactate, 50 mg complete amino acid mixture, 660 mg yeast nitrogen base per 100 mL) and cultured for 48 h.

From this starter culture, 1% (v/v) inoculum was transferred daily into fresh medium to establish six independent replicate populations in each of two carbon environments:

- Glucose (0.5% w/v) supplemented with 50 mg complete amino acid mixture and 0.66 mg yeast nitrogen base
- Galactose (0.5% w/v) supplemented with the same additives

All populations were propagated by serial 1:100 dilutions every 24 h and maintained at 30°C, 250rpm shaking conditions for a total of ~1200 generations.

Growth Rate Measurements as a Proxy for Adaptation

Growth rates were quantified at four evolutionary time points: 300, 600, 900, and 1200 generations. At each point, populations were revived from frozen stocks and inoculated into the same carbon source in which they had been evolved. Optical density (OD) measurements were then recorded to estimate growth rate, which served as a proxy for fitness.

We denote these assays as follows:

- ggg: populations evolved in glucose, passaged in glucose, and assayed in glucose.
- GGG: populations evolved in galactose, passaged in galactose, and assayed in galactose.
- gGg: populations evolved in glucose, passaged in galactose, and assayed in glucose.
- GgG: populations evolved in galactose, passaged in glucose, and assayed in galactose.

This design ensured that the measured growth rates reflected adaptation to the *native selection environment* of each evolving population.

Reciprocal Environment Assays

294 To assess the stability and plasticity of adaptation, evolved populations were exposed to the non-home
295 carbon source for a short duration (~15 generations) and subsequently returned to their original
296 environment. For glucose-evolved populations, the non-home environment was 0.5% galactose
297 medium, while for galactose-evolved populations, the non-home environment was 0.5% glucose
298 medium.
299
300 Fitness was assayed in the home environment at three distinct stages:
301
302 • Immediately after growth in the non-home environment
303 • Following ~7 generations of regrowth in the home environment
304 • Following ~15 generations of regrowth in the home environment
305
306 At each assay point, 100–200 μL of culture (volume adjusted according to OD) was inoculated into 10
307 mL of the relevant assay medium. Cultures were incubated at 30 °C with shaking at 250 rpm, and optical
308 density (OD) measurements were collected at regular intervals. Growth dynamics were quantified by
309 calculating a composite growth rate, incorporating both the lag phase duration and the exponential
310 growth rate derived from the OD growth curves.
311
312 Each kinetic assay was performed in triplicate for every population at every time point. In parallel,
313 ancestral strains were revived and assayed under identical conditions, providing a consistent baseline
314 for estimating relative fitness gains during evolution.
315
Whole-Genome Sequencing and Reference Genome Construction
Evolved populations at four evolutionary time points (300, 600, 900, and 1200 generations) were revived
in 5 mL of their respective evolution environments. Cells were harvested, and genomic DNA was
extracted using the phenol–chloroform method[70]. Sequencing libraries were prepared, and 150 bp
paired-end sequencing was performed on the DNBSEQ platform at ~100× coverage by Haystack
Analytics. The ancestral diploid SK1 strain was also sequenced at the same depth for comparison.
322
To construct a draft SK1 reference genome, three ancestral haploid SK1 strains from our laboratory
were independently sequenced at ~100× coverage. Each haploid genome was aligned to the S288C
reference genome using BWA-MEM, and variants were called with GATK v4.4.0.0. Common variants
across the three strains were identified and incorporated into the reference using
FastaAlternateReferenceMaker, generating a modified reference genome. This process of alignment,
variant calling, and incorporation of common polymorphisms was iterated until convergence, yielding a
high-confidence SK1 draft genome[71].
330
The diploid ancestral SK1 genome was then aligned to this draft reference, and the resulting BAM file
was used to polish the assembly with Pilon. Gene annotations were transferred from the S288C genome
to the SK1 draft genome using Liftoff, yielding a polished and annotated SK1 reference genome[72,73].

This polished SK1 genome was subsequently used as the reference for alignment of all evolved
population sequencing data.

Variant Calling and Filtering
For each evolved population, sequencing reads were aligned to the polished SK1 reference genome
using BWA-MEM. Duplicate reads were identified and marked with Picard Tools. Variant calling was
performed with GATK4, following the recommended best practices pipeline: HaplotypeCaller,
GenomicsDBImport, and GenotypeGVCFs. Variants were jointly called across all time points for each
replicate line to maintain consistency in variant representation. Variant annotation was carried out using
SnpEff[72].

From the resulting VCF files, total read depth and allele-specific read depths (reference and alternate
alleles) were extracted for each sample. Variants were filtered out based on the following conditions:

• the average coverage across all time points was <30×; and
• all populations across the four time points had <25× coverage; and
• the allele frequency remained unchanged across time.

Mutations were considered fixed when the alternate allele frequency reached thresholds of:

• 0.5–0.6 for heterozygous alternate alleles,
• 0.9–1.0 for homozygous alternate alleles, or
• 0–0.1 for homozygous reference alleles, relative to the reference genome.

. All variants were visually validated using the Integrated Genome Viewer (IGV).

Because the number of fixed SNPs was low and many were shared across replicate populations, variant
calls for one representative replicate line (across all time points in both environments) were
independently verified using an alternative approach. Variants were re-called with bcftools mpileup,
followed by filtering with the same thresholds as above. All the 5 SNPs called using GATK in glu1
population (glucose-evolved, replicate line1) were present in the variant list obtained from bcftools
mpileup. There were no unique variants to GATK but 1 unique variant to bcftools mpileup. In the case
of gal1 population (galactose-evolved, replicate line1), 3 SNPs were common to both the methods and
1 unique to GATK. This confirms the presence of few fixed variants in the population.
The   raw   sequencing   data   of   all   evolved   populations   is   available   at
http://www.ncbi.nlm.nih.gov/bioproject/1298216.

Copy Number Variation (CNV) Analysis

To ensure comparable sequencing coverage across samples, the median coverage across the genome was calculated for each evolved replicate at every time point and in both evolution environments **(Supplementary Figure S7).**

A custom coverage-based pipeline was used to identify copy number variants (CNVs) in all evolved populations. First, per-site read depth across the genome was calculated using samtools depth. To detect contiguous genomic regions exhibiting CNVs, the genome was divided into 500 bp windows, and the median coverage per window was determined. Each window's coverage was then normalized by the genome-wide median coverage for that sample. To account for standing variation present in the ancestor, the ratio of normalized coverage in evolved populations to that of the ancestral reference was computed for each window, yielding a measure of relative coverage. All the windows that had a relative coverage less than 0.25 were excluded from further analysis.

To classify CNV states, we implemented an unsupervised Hidden Markov Model (HMM) allowing discrete states of 0, 0.5, 1, 1.5, 2, 3, and 4. A CNV state of 1 indicated equal copy number in evolved and ancestral genomes, values <1 indicated deletions, and values >1 indicated duplications in the evolved populations[72].

To validate the CNV calls, one representative replicate from each environment was independently analyzed at a time point that exhibited maximum CNVs (300 generations for glucose and 1200 generations for galactose), along with the ancestor, using CNVpytor, a read depth-based tool for CNV analysis[74]. A bin size of 500bp was used, and CNV calls with p-value<0.0001, and Q0<0.5 were filtered . The copy number in each window of evolved populations was normalized with that in the ancestor to account for only the changes that occurred during the experiment. The results were compared with the coverage-based pipeline used in this study. In the glucose-evolved line, results obtained from CNVpytor had more deletions than duplications. On comparing the two methods, the coverage-based pipeline had an accuracy of 86.58%, precision of 86.94% and recall of 86.58%. In the case of galactose-evolved line, duplications were higher than deletions, and the coverage-based pipeline could predict this with accuracy of 79.58% (precision and recall of 79.61% and 79.58% respectively) when compared to CNVpytor. This confirms that the trends of deletions in glucose and duplications in galactose are consistently observed not just in our pipeline but also in another independent tool.

CNV Annotation and Pathway Enrichment Analysis

Using the CNV states obtained from coverage-based analysis, all windows with a state of 1 (no change relative to ancestor) were excluded. The remaining windows were categorized as duplications (>1) or deletions (<1). Each set of windows was annotated to identify the genes contained within the windows for every replicate at each time point. Genes of unknown function were filtered out, and annotations

414   across time points were merged within each replicate to obtain the total number of unique genes
415   affected by CNVs. To visualize genomic variation, the average CNV state per window was calculated
416   across replicates and plotted for all time points in both environments.
417

For Pathway Enrichment Analysis (PEA), only CNVs shared across all replicates were considered. A t-
test was performed for each genomic window based on the relative coverage ratio across replicates.
Windows with $p < 0.05$ and a CNV state $\neq 1$ in at least four replicates were identified as high-confidence
CNVs. These were further separated into duplications and deletions, annotated, and filtered to exclude
genes of unknown function. For each time point and environment, a nonredundant list of unique genes
showing CNVs across replicates was compiled.
424

Because the resulting gene lists were large (300–1200 genes), they were randomly partitioned into bins
of 100 genes, and KEGG pathway enrichment analysis was performed on each bin using the
clusterProfiler package in R[74]. This random segregation and enrichment analysis was repeated 3–12
times, depending on gene list size. The adjusted p-values from all runs were combined using Fisher's
method, and pathways were ranked by gene count and adjusted p-value to identify the most significantly
affected pathways. This analysis was performed independently for each time point in both evolution
environments.
432

433 **Acknowledgements**

434 This work was supported by the DBT/Wellcome Trust India Alliance grant IA/S/19/2/504632 awarded
435 to SS.

Figure 1. Experimental design, fitness gains, and a genomic overview of adaptation. (A) Schematic of the experimental evolution. Six independent diploid *S. cerevisiae* populations were propagated for 1,200 generations in either glucose or galactose. Populations were sampled at 0, 300, 600, 900, and 1,200 generations for phenotyping and whole-genome sequencing. **(B)** Home-environment fitness trajectories (mean relative growth rate ± s.e.m., *n* = 6 lines per environment) show steady improvement in both media; glucose-evolved lines (blue) rise rapidly and then plateau, while galactose-evolved lines (red) increase more gradually. **(C)** SNP accumulation is rare. Bars show the total number of fixed SNPs detected across all six replicate lines over 1,200 generations in each environment (joint calling across time points; stringent depth and allele-frequency filters; visual confirmation in IGV). **(D–E)** Copy-number variation (CNV) dominates genome change and differs by environment. Box-and-whisker plots summarize CNV across 1,200 generations. (D) deletions only and (F) duplications only. Boxes show interquartile range with median; whiskers span full range; ✕ indicates mean. Glucose-evolved lines carry abundant deletions, whereas galactose-evolved lines show persistent, higher duplication loads, exhibiting environment-specific CNV strategies.

Figure 2. CNV dynamics in glucose-evolved populations across 1,200 generations. CNV states
for the six glucose-evolved lines at 300, 600, 900, and 1,200 generations. The *x*-axis lists the 16 *S.
cerevisiae* chromosomes. Each point is a 500-bp genomic window; values are normalized to the
ancestor such that state = 1 indicates no change, <1 indicates deletion, and >1 indicates duplication.
The colored horizontal bars denote the mean CNV state for that chromosome at that time point. At
300 generations, windows with deletion states (≈0.6–0.95) are widespread and concentrated near
chromosome ends, revealing early, extensive telomeric loss. Through 600–900 generations, the
deletion burden decreases (mean states trend toward 1). By 1,200 generations, most chromosomes
sit close to state 1 with fewer and weaker deletions, consistent with refinement/compensation of early
structural changes. Overall, the trajectory indicates that early telomeric deletions dominate adaptation
in glucose, followed by later remodeling that reduces deletion extent and introduces targeted
duplications.

Figure 3. Persistent subtelomeric duplications in galactose-evolved populations. CNV states for the six galactose-evolved lines at 300, 600, 900, and 1,200 generations. The *x*-axis lists the 16 *S. cerevisiae* chromosomes. Each point represents a 500-bp window; state = 1 indicates no change, <1 deletion, >1 duplication. The colored horizontal bar marks the mean CNV state for that chromosome at that time point. Across all time points, windows with duplication states (>1.0–1.7) are abundant and cluster near chromosome ends, revealing early-onset, persistent subtelomeric duplications under galactose selection except in chromosome II. Unlike the glucose-evolved trajectories (Figure 2), deletions are rare and mean chromosome states remain consistently >1 from 300 to 1,200 generations, indicating sustained duplication loads rather than deletion-first remodeling. Together, these patterns demonstrate a duplication-biased CNV program in galactose that is stable over evolutionary time and concentrated in telomeric neighborhoods enriched for transport/metabolic modules.

Figure 4. **Divergent copy number evolution in glucose versus galactose.** Many genes deleted
during adaptation in glucose appear duplicated during adaptation in galactose, visible as rows that are
red in the left block and blue in the right block. This highlights environment specific resolution of
selection on the same loci over evolutionary time. Each row is a gene. Columns show four time points
in glucose (300, 600, 900, 1200 generations) followed by four time points in galactose (300, 600, 900,
1200 generations). Color encodes the direction of copy number change relative to the ancestor: red
indicates deletions, blue indicates duplications, white indicates no detected change (scale bar at right).

Figure 5. History-dependent fitness costs after cross-environment preculture. For each evolved
background, fitness cost was computed as the relative decrease in composite growth rate in the home
environment when cells were precultured in the non-home carbon source. Composite growth rate
integrates lag and exponential growth from OD time courses (Methods). Lines show the mean cost
across six replicate populations at each evolutionary time point (300, 600, 900, 1,200 generations);
blue, glucose-evolved populations precultured in galactose and assayed in glucose; red, galactose-
evolved populations precultured in glucose and assayed in galactose. Glucose-evolved populations
display a transient specialization: a pronounced cost early (peaking by ~600 generations) that
collapses by 900 generations and remains low thereafter, indicating reduced sensitivity to preculture
history. In contrast, galactose-evolved populations show a persistent, increasing cost across the
experiment, consistent with deepening specialization to galactose and strong history dependence
following glucose exposure.

Figure 6. Short-term history effects and recovery in home environment. For each evolutionary
background, we measured growth in the home environment after a brief exposure to the non-home
carbon source, with or without a recovery phase back in the home medium. Fitness is shown as a
ratio to the corresponding home-only control (for (A): divided by fitness of ggg; for (B): divided by
fitness of GGG), where "g/G" denote glucose/galactose and the string indicates evolution →
preculture → (optional recovery duration) → assay. Thus, g-G-g = glucose-evolved, precultured in
galactose, assayed in glucose; g-G-g7-g / g-G-g15-g include ~7 or ~15 generations of recovery in
glucose before assay. Analogously, G-g-G, G-g-G7-G, G-g-G15-G are for galactose-evolved lines.
Lines show the mean across six replicate populations at each evolutionary time point (0, 300, 600,
900, 1,200 generations). Composite fitness integrates lag and exponential growth from OD time
courses (Methods). **(A)** Glucose-evolved populations exhibit history dependence early: performance
in glucose after galactose preculture depends on recovery duration (g-G-g < g-G-g7-g ≈ g-G-g15-g)
through ~600 generations, after which the curves converge, indicating a loss of history dependence.
(B) Galactose-evolved populations (right) show a strong penalty when assayed immediately after
glucose preculture (G-g-G), whereas 7–15 generations of re-exposure to galactose (G-g-G7-G, G-g-
G15-G) restore fitness toward the home-only baseline across all evolutionary time points.

References

1. Bell, G. Fluctuating selection: the perpetual renewal of adaptation in variable environments. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **365**, 87–97 (2010).

2. Levin, S. A. The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur Award Lecture. *Ecology* **73**, 1943–1967 (1992).

3. Chevin, L.-M., Lande, R. & Mace, G. M. Adaptation, plasticity, and extinction in a changing environment: towards a predictive theory. *PLoS Biol.* **8**, e1000357 (2010).

4. Botero, C. A., Weissing, F. J., Wright, J. & Rubenstein, D. R. Evolutionary tipping points in the capacity to adapt to environmental change. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 184–189 (2015).

5. Lässig, M., Mustonen, V. & Walczak, A. M. Predicting evolution. *Nat. Ecol. Evol.* **1**, 77 (2017).

6. Stern, D. L. & Orgogozo, V. Is genetic evolution predictable? *Science* **323**, 746–751 (2009).

7. Orr, H. A. The genetic theory of adaptation: a brief history. *Nat. Rev. Genet.* **6**, 119–127 (2005).

8. Kryazhimskiy, S., Rice, D. P., Jerison, E. R. & Desai, M. M. Microbial evolution. Global epistasis makes adaptation predictable despite sequence-level stochasticity. *Science* **344**, 1519–1522 (2014).

9. Nosil, P., Flaxman, S. M., Feder, J. L. & Gompert, Z. Increasing our ability to predict contemporary evolution. *Nat. Commun.* **11**, 5592 (2020).

10. Ahlawat, N., Venkataraman, P., Brajesh, R. G. & Saini, S. Effects of resource packaging on the adaptative and pleiotropic consequences of evolution. *NPJ Syst. Biol. Appl.* **11**, 78 (2025).

11. Ahlawat, N., Mahilkar, A. & Saini, S. Resource presentation dictates genetic and phenotypic adaptation in yeast. *BMC Ecol. Evol.* **25**, 33 (2025).

12. Kawecki, T. J. *et al.* Experimental evolution. *Trends Ecol. Evol.* **27**, 547–560 (2012).

13. Elena, S. F. & Lenski, R. E. Evolution experiments with microorganisms: the dynamics and genetic bases of adaptation. *Nat. Rev. Genet.* **4**, 457–469 (2003).

14. Barrick, J. E. & Lenski, R. E. Genome dynamics during experimental evolution. *Nat. Rev. Genet.* **14**, 827–839 (2013).

15. Levy, S. F. *et al.* Quantitative evolutionary dynamics using high-resolution lineage tracking. *Nature* **519**, 181–186 (2015).

16. Dettman, J. R. *et al.* Evolutionary insight from whole-genome sequencing of experimentally evolved microbes. *Mol. Ecol.* **21**, 2058–2077 (2012).

17. Trumbly, R. J. Glucose repression in the yeast Saccharomyces cerevisiae. *Mol. Microbiol.* **6**, 15–

21 (1992).

18. Escalante-Chong, R. *et al.* Galactose metabolic genes in yeast respond to a ratio of galactose

and glucose. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 1636–1641 (2015).

19. Roop, J. I., Chang, K. C. & Brem, R. B. Polygenic evolution of a sugar specialization trade-off in

yeast. *Nature* **530**, 336–339 (2016).

20. Cerulus, B. *et al.* Transition between fermentation and respiration determines history-dependent

behavior in fluctuating carbon sources. *eLife* **7**, e39234 (2018).

21. Stockwell, S. R., Landry, C. R. & Rifkin, S. A. The yeast galactose network as a quantitative

model for cellular memory. *Mol. Biosyst.* **11**, 28–37 (2015).

22. Mahilkar, A., Nagendra, P., Venkataraman, P., Deshmukh, S. & Saini, S. Rapid evolution of pre-

zygotic reproductive barriers in allopatric populations. *Microbiol. Spectr.* **11**, e0195023 (2023).

23. Mahilkar, A., Nagendra, P., Alugoju, P., E, R. & Saini, S. Public good-driven release of

heterogeneous resources leads to genotypic diversification of an isogenic yeast population. *Evol.*

Int. J. Org. Evol. **76**, 2811–2828 (2022).

24. Rajeshkannan, null, Mahilkar, A. & Saini, S. GAL Regulon in the Yeast S. cerevisiae is Highly

Evolvable via Acquisition in the Coding Regions of the Regulatory Elements of the Network.

Front. Mol. Biosci. **9**, 801011 (2022).

25. Lang, G. I. *et al.* Pervasive genetic hitchhiking and clonal interference in forty evolving yeast

populations. *Nature* **500**, 571–574 (2013).

26. Payen, C. *et al.* High-Throughput Identification of Adaptive Mutations in Experimentally Evolved

Yeast Populations. *PLoS Genet.* **12**, e1006339 (2016).

27. Dunham, M. J. *et al.* Characteristic genome rearrangements in experimental evolution of

Saccharomyces cerevisiae. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 16144–16149 (2002).

28. Brown, C. J., Todd, K. M. & Rosenzweig, R. F. Multiple duplications of yeast hexose transport

genes in response to selection in a glucose-limited environment. *Mol. Biol. Evol.* **15**, 931–942

(1998).

29. Gresham, D. *et al.* The repertoire and dynamics of evolutionary adaptations to controlled nutrient-

limited environments in yeast. *PLoS Genet.* **4**, e1000303 (2008).

30. Gresham, D. *et al.* Adaptation to diverse nitrogen-limited environments by deletion or

extrachromosomal element formation of the GAP1 locus. *Proc. Natl. Acad. Sci. U. S. A.* **107**,

18551–18556 (2010).

31. Yona, A. H. *et al.* Chromosomal duplication is a transient evolutionary solution to stress. *Proc.*

Natl. Acad. Sci. U. S. A. **109**, 21010–21015 (2012).

32. Gorkovskiy, A. & Verstrepen, K. J. The Role of Structural Variation in Adaptation and Evolution of

Yeast and Other Fungi. *Genes* **12**, 699 (2021).

33. Hull, R. M., Cruz, C., Jack, C. V. & Houseley, J. Environmental change drives accelerated

adaptation through stimulated copy number variation. *PLoS Biol.* **15**, e2001333 (2017).

34. Gresham, D. *et al.* Adaptation to diverse nitrogen-limited environments by deletion or

extrachromosomal element formation of the GAP1 locus. *Proc. Natl. Acad. Sci. U. S. A.* **107**,

18551–18556 (2010).

35. Todd, R. T. & Selmecki, A. Expandable and reversible copy number amplification drives rapid

adaptation to antifungal drugs. *eLife* **9**, e58349 (2020).

36. Andersson, D. I. & Hughes, D. Gene amplification and adaptive evolution in bacteria. *Annu. Rev.*

Genet. **43**, 167–195 (2009).

37. Raj, N. & Saini, S. Increased privatization of a public resource leads to spread of cooperation in a

microbial population. *Microbiol. Spectr.* **12**, e0235823 (2024).

38. Fogel, S., Welch, J. W., Cathala, G. & Karin, M. Gene amplification in yeast: CUP1 copy number

regulates copper resistance. *Curr. Genet.* **7**, 347–355 (1983).

39. Onetto, C. A. *et al.* SO2 and copper tolerance exhibit an evolutionary trade-off in Saccharomyces

cerevisiae. *PLoS Genet.* **19**, e1010692 (2023).

40. Ning, Y., Dai, R., Zhang, L., Xu, Y. & Xiao, M. Copy number variants of ERG11: mechanism of

azole resistance in Candida parapsilosis. *Lancet Microbe* **5**, e10 (2024).

41. Vande Zande, P. *et al.* Step-wise evolution of azole resistance through copy number variation

followed by KSR1 loss of heterozygosity in Candida albicans. *PLoS Pathog.* **20**, e1012497

(2024).

42. Nair, S. *et al.* Adaptive copy number evolution in malaria parasites. *PLoS Genet.* **4**, e1000243

(2008).

43. Triglia, T., Foote, S. J., Kemp, D. J. & Cowman, A. F. Amplification of the multidrug resistance

gene pfmdr1 in Plasmodium falciparum has arisen as multiple independent events. *Mol. Cell.*

Biol. **11**, 5244–5250 (1991).

44. Hu, G. *et al.* Variation in chromosome copy number influences the virulence of Cryptococcus

neoformans and occurs in isolates from AIDS patients. *BMC Genomics* **12**, 526 (2011).

45. Abdul-Rahman, F. & Gresham, D. Copy number variation facilitates rapid toggling between

ecological strategies. *BioRxiv Prepr. Serv. Biol.* 2025.07.22.666191 (2025)

doi:10.1101/2025.07.22.666191.

46. Liu, H. & Zhang, J. Yeast Spontaneous Mutation Rate and Spectrum Vary with Environment. *Curr.*

Biol. CB **29**, 1584-1591.e3 (2019).

47. Zhu, Y. O., Siegal, M. L., Hall, D. W. & Petrov, D. A. Precise estimates of mutation rate and

spectrum in yeast. *Proc. Natl. Acad. Sci.* **111**, (2014).

48. Pankajam, A. V., Dash, S., Saifudeen, A., Dutta, A. & Nishant, K. T. Loss of Heterozygosity and

Base Mutation Rates Vary Among Saccharomyces cerevisiae Hybrid Strains. *G3 Bethesda Md*

10, 3309–3319 (2020).

49. Dutta, A., Dutreux, F. & Schacherer, J. Loss of Heterozygosity Spectrum Depends on Ploidy Level

in Natural Yeast Populations. *Mol. Biol. Evol.* **39**, msac214 (2022).

50. Zhao, Y. *et al.* Structures of naturally evolved CUP1 tandem arrays in yeast indicate that these

arrays are generated by unequal nonhomologous recombination. *G3 Bethesda Md* **4**, 2259–2269

(2014).

51. Yue, J.-X. *et al.* Contrasting evolutionary genome dynamics between domesticated and wild

yeasts. *Nat. Genet.* **49**, 913–924 (2017).

52. O'Donnell, S. *et al.* Telomere-to-telomere assemblies of 142 strains characterize the genome

structural landscape in Saccharomyces cerevisiae. *Nat. Genet.* **55**, 1390–1399 (2023).

53. Carlson, M., Celenza, J. L. & Eng, F. J. Evolution of the dispersed SUC gene family of

Saccharomyces by rearrangements of chromosome telomeres. *Mol. Cell. Biol.* **5**, 2894–2902

(1985).

54. Charron, M. J., Read, E., Haut, S. R. & Michels, C. A. Molecular evolution of the telomere-

associated MAL loci of Saccharomyces. *Genetics* **122**, 307–316 (1989).

55. Lyons, D. M., Zou, Z., Xu, H. & Zhang, J. Idiosyncratic epistasis creates universals in mutational

effects and evolutionary trajectories. *Nat. Ecol. Evol.* **4**, 1685–1693 (2020).

56. Bakerlee, C. W., Nguyen Ba, A. N., Shulgina, Y., Rojas Echenique, J. I. & Desai, M. M.

Idiosyncratic epistasis leads to global fitness-correlated trends. *Science* **376**, 630–635 (2022).

57. Bakerlee, C. W., Phillips, A. M., Nguyen Ba, A. N. & Desai, M. M. Dynamics and variability in the

pleiotropic effects of adaptation in laboratory budding yeast populations. *eLife* **10**, e70918 (2021).

58. Jerison, E. R. *et al.* Genetic variation in adaptability and pleiotropy in budding yeast. *eLife* **6**,

e27167 (2017).

59. Hassan, N. *et al.* CRISPR-PCDup: a novel approach for simultaneous segmental chromosomal

duplication in Saccharomyces cerevisiae. *AMB Express* **10**, 27 (2020).

60. Easmin, F. *et al.* CRISPR-PCD and CRISPR-PCRep: Two novel technologies for simultaneous

multiple segmental chromosomal deletion/replacement in Saccharomyces cerevisiae. *J. Biosci.*

Bioeng. **129**, 129–139 (2020).

61. James, T. Y. *et al.* Adaptation by Loss of Heterozygosity in Saccharomyces cerevisiae Clones

Under Divergent Selection. *Genetics* **213**, 665–683 (2019).

62. Smukowski Heil, C. S. *et al.* Loss of Heterozygosity Drives Adaptation in Hybrid Yeast. *Mol. Biol.*

Evol. **34**, 1596–1612 (2017).

63. Xu, H. *et al.* Chromosome drives via CRISPR-Cas9 in yeast. *Nat. Commun.* **11**, 4344 (2020).

64. Zimmer, A. *et al.* QTL dissection of Lag phase in wine fermentation reveals a new translocation

responsible for Saccharomyces cerevisiae adaptation to sulfite. *PloS One* **9**, e86298 (2014).

65. García-Ríos, E., Nuévalos, M., Barrio, E., Puig, S. & Guillamón, J. M. A new chromosomal

rearrangement improves the adaptation of wine yeasts to sulfite. *Environ. Microbiol.* **21**, 1771–

1781 (2019).

66. Selmecki, A., Gerami-Nejad, M., Paulson, C., Forche, A. & Berman, J. An isochromosome confers

drug resistance in vivo by amplification of two genes, ERG11 and TAC1. *Mol. Microbiol.* **68**, 624–

641 (2008).

67. Nguyen Ba, A. N. *et al.* High-resolution lineage tracking reveals travelling wave of adaptation in

laboratory yeast. *Nature* **575**, 494–499 (2019).

68. Jiang, T., Liu, S., Cao, S. & Wang, Y. Structural Variant Detection from Long-Read Sequencing

Data with cuteSV. *Methods Mol. Biol. Clifton NJ* **2493**, 137–151 (2022).

69. Chochinov, C. A. & Nguyen Ba, A. N. Bulk-Fitness Measurements Using Barcode Sequencing

Analysis in Yeast. *Methods Mol. Biol. Clifton NJ* **2477**, 399–415 (2022).

70. Dymond, J. S. Preparation of genomic DNA from Saccharomyces cerevisiae. *Methods Enzymol.*

529, 153–160 (2013).

71. Lang, G. I., Parsons, L. & Gammie, A. E. Mutation rates, spectra, and genome-wide distribution of

spontaneous mutations in mismatch repair deficient yeast. *G3 Bethesda Md* **3**, 1453–1465

(2013).

72. Johnson, M. S. *et al.* Phenotypic and molecular evolution across 10,000 generations in laboratory

budding yeast populations. *eLife* **10**, e63910 (2021).

73. Shumate, A. & Salzberg, S. L. Liftoff: accurate mapping of gene annotations. *Bioinforma. Oxf.*

Engl. **37**, 1639–1643 (2021).

74. Suvakov, M., Panda, A., Diesh, C., Holmes, I. & Abyzov, A. CNVpytor: a tool for copy number

variation detection and analysis from read depth and allele imbalance in whole-genome

sequencing. *GigaScience* **10**, giab074 (2021).