Alternate Splicing Directs PMCA2 to Lysosomes and is Linked to Neurodegeneration

Fernández-Suárez, M.E.[1,2], Bush, R.[1,3], Brenton, J.W.[2,4,5,6,7], Pereira, G.[8,9], Grant-Peters, M.[7,10],

Reynolds, R.[5,6], te Vruchte, D.[1,2], Shepherd, D.[1], Weng, Y.[1], Artaza-Fernández, E.[1], Lis, P.[2,11],

Sanchez-Pulido, L.[12], Morgan, A.[1], Davis, L.[1], Gómez-Coronado, D.[13], Eden, E.[14], Alessi,

D.[2,11], Ponting, C.P.[12], Galione, A [1], Newstead, S.[3,15], Hardy, J.[2,16,17], Patel, S.[8], Ryten,

M.[2,7,10,18], and Platt, F.M[*1,2].

1. Department of Pharmacology, University of Oxford, Oxford, UK.

2. Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy

Chase, MD, USA.

3. Department of Biochemistry, University of Oxford, Oxford, UK.

4. UCL Queen Square Institute of Neurology, Department of Neurodegenerative Diseases,

London, UK.

5. Department of Genetics and Genomic Medicine, Great Ormond Street Institute of Child

Health, University College London, London, UK.

6. NIHR Great Ormond Street Hospital Biomedical Research Centre, University College

London, London, UK.

7. UK Dementia Research Institute at The University of Cambridge, Cambridge, UK.

8. Department of Cell and Developmental Biology, University College London, London, UK.

9. Department of Pharmacology, Federal University of São Paulo (UNIFESP), São Paulo,

Brazil.

10. Department of Clinical Neurosciences, School of Clinical Medicine, University of

Cambridge, Cambridge, UK

11. Medical Research Council (MRC) Protein Phosphorylation and Ubiquitylation Unit,

School of Life Sciences, University of Dundee, Dundee, UK.

12. MRC Human Genetics Unit, Institute of Genetics & Cancer, University of Edinburgh,

Edinburgh, UK.

13. Department of Biochemistry-Research, Hospital Universitario Ramón y Cajal, IRYCIS,

Madrid, Spain.

14. Institute of Ophthalmology, University College London, London, UK.

15. Kavli Institute for Nanoscience Discovery, University of Oxford, Oxford, UK.

16. Department of Neurodegenerative Disease, UCL Institute of Neurology, UK.

17. UK Dementia Research Institute at UCL, London, UK.

18. Department of Medical Genetics, NIHR Cambridge Biomedical Research Centre,

University of Cambridge, Cambridge, UK.

* Corresponding author: frances.platt@pharm.ox.ac.uk

Abstract

Plasma membrane calcium ATPase (PMCAs) are believed to function exclusively at the plasma membrane where they expel calcium from the cytosol. We have unexpectedly identified a splice variant-dependent localisation of the PMCA isoform PMCA2 to the lysosome, where it forms an evolutionarily conserved complex with NPC1, the lysosomal membrane protein defective in the rare lysosomal storage disease Niemann-Pick disease type C (NPC). This interaction is involved in maintaining lysosomal calcium levels, and we have found its disruption contributes to the pathophysiology of both NPC and Parkinson's disease. Our findings uncover a previously unrecognised intracellular role for PMCA2, revealing a shared mechanism of lysosomal calcium and lipid regulation with direct relevance to neurodegeneration.

Introduction

Precise regulation of intracellular calcium (Ca^{2+}) homeostasis is essential for neuronal function and survival. Plasma membrane calcium ATPases (PMCAs) are high-affinity Ca^{2+} pumps that move cytosolic Ca^{2+} across the plasma membrane to the extracellular environment [1]. Of the four PMCA isoforms in humans (PMCA1–4), PMCA2 and PMCA3 are highly expressed in neurons in the brain, with PMCA2 particularly enriched in cerebellar Purkinje cells [1]. Loss-of-function mutations in PMCA2 or PMCA3 cause progressive cerebellar ataxia [2,3,4], and reduced PMCA2 expression has been associated with increased vulnerability of dopaminergic neurons in Parkinson's disease (PD) [5]. Moreover, soluble α-synuclein, implicated in PD pathogenesis, directly enhances PMCA1/2 activity, underscoring the relevance of PMCA regulation in neurodegeneration [6]. To date, all functional roles of PMCAs have been attributed to its localisation at the plasma membrane. However, the genes encoding the PMCAs are alternatively spliced, raising the possibility that they could have additional localisations and functions.

In a yeast screen, we previously identified PMC1, the single yeast orthologue of mammalian PMCAs, as a direct interactor of Ncr1, the yeast orthologue of mammalian NPC1 [7]. This interaction was conserved in mammalian cells [7]. NPC1 is a thirteen-transmembrane lysosomal protein encoded by *NPC1* that is mutated in the rare neurodegenerative lysosomal storage disorder, Niemann-Pick disease type C (NPC). NPC1 deficiency results in disrupted lysosomal lipid homeostasis, impaired lysosome: ER membrane contact site formation and, critically, lowered levels of lysosomal luminal Ca^{2+} [8], potentially due to a Ca^{2+} store filling defect, a feature also reported in PD [9]. However, the molecular mechanism underlying Ca^{2+} import into the lysosome remains incompletely understood.

Here, we show that alternative splicing directs specific PMCA2 isoforms to the lysosome, where they form a complex with NPC1 to regulate lysosomal Ca^{2+} levels. These findings have

76   uncovered a conserved calcium regulatory axis with broad implications for lysosome function

77   and neurodegenerative diseases. This study also highlights that splice variants can differentially

78   affect subcellular organelle targeting of proteins, expanding the function of proteins beyond

79   their canonical roles.

Results

PMCA2 colocalises with NPC1 in the lysosome.

We previously demonstrated that NPC1 and PMCA2 from rat cerebellum co-immunoprecipitate [7]. To determine whether this interaction occurs at the lysosomal membrane, we examined the localisation of endogenous PMCA2 in rat PC12 cells, a cell line reported to express PMCA2 and that can be differentiated into neurons [10] [11]. Immunocytochemistry with a pan anti-PMCA2 polyclonal antibody that recognises all splice variants, followed by confocal microscopy, revealed that endogenous PMCA2 colocalised with the lysosomal protein NPC1, with no detectable signal at the plasma membrane (Fig. 1a). Individual staining controls of PMCA2 and NPC1 showed no bleed-through between channels or non-specific staining when no primary antibody was present (Supplementary Figure 1). To test the specificity of the anti-PMCA2 antibody, we knocked down (KD) the expression of endogenous PMCA2 in PC12 cells by stably expressing a shRNA targeting *Atp2b2*, the gene encoding the PMCA2 protein. Immunocytochemistry showed a distinct punctate pattern consistent with lysosomal localisation of PMCA2 in wild type (WT) cells that was absent in the *Atp2b2* KD cells (Supplementary Figure 2).

To confirm these results, we used a complementary technique isolating lysosomes from the cerebellum of WT mice using Tagless LysoIP [12]. Lysosomes were immunoprecipitated using a monoclonal antibody bound to magnetic beads that binds a cytosolic epitope of the endogenous lysosomal membrane protein TMEM192. As a negative control, magnetic beads bound to an isotype-matched control antibody were used. The purified lysosomes were analysed using western blotting probing using the anti-PMCA2 polyclonal antibody, and a band of approximately 130 kDa was detected in the input fraction (Figure 1b). This band

corresponded to the predicted molecular weight of PMCA2 and was statistically significantly

enriched in the lysosomal fraction isolated from the cerebellum (Figure 1c).

Having confirmed the lysosomal localisation of PMCA2, we modelled the PMCA2 structure

and orientation in the limiting membrane of the lysosome and its direct interaction with NPC1

using AlphaFold 3 software. The resulting human NPC1-PMCA2 complex (Fig. 1d) models

both the N- and C-terminal domains of PMCA2 on the cytosolic face of the lysosomal

membrane, an orientation compatible with Ca^{2+} import into the lysosome (red arrow, Figure

1d and e). Strikingly, a corresponding AlphaFold 3 model of the yeast Pmc1/Ncr1 complex

(Fig. 1e) reveals a conserved topology, suggesting evolutionary conservation of the formation

and function of this complex.

Figure 1. PMCA2 is present in the lysosome. (a) Immunocytochemistry and confocal microscopy showed that endogenous PMCA2 colocalises with NPC1 in the lysosome of PC12 cells differentiated into neurons. NPC1 and PMCA2 were detected using primary antibodies directly labelled with CoraLite-555 and CoraLite-488, respectively. Representative image of five independent experiments, Pearson's coefficient r = 0.8636 ± 0.0437 standard deviation corresponding to the mean of five independent experiments, calculated using the mean of r from 10 fields per experiment. (b) Representative image of Tagless LysoIP followed by Western blot probing for PMCA2 in mouse cerebellar tissue, $n = 6$ mice. (c) Quantification of PMCA2 enrichment in lysosomal fraction from LysoIP experiments, $n = 6$ mice, lines represent the mean and standard deviation, one-way Anova test, $*p < 0.05$, $***p < 0.001$. (d) AlphaFold 3 model of the PMCA2-NPC1 complex in the lysosomal membrane. Directionality of Ca^{2+} transport is represented with a red arrow (e) AlphaFold 3 model of yeast orthologues Ncr1 and Pmc1 in the vacuolar membrane exhibiting a similar orientation and interaction.

PMCA2 depletion disrupts lysosomal Ca^{2+} signalling.

The discovery of the lysosomal localisation of PMCA2 and the orientation of PMCA2 in the complex with NPC1 raised the possibility that it could play a key role in filling the lysosome with Ca^2. To test this hypothesis, we used the PC12 *Atp2b2* KD cells described above. The shRNA construct used to generate these cells contained a GFP cassette as a positive control of transduction. These cells were differentiated into neurons and then loaded with Fura-2, a fluorescent probe that accumulates in the cytoplasm. Lysosomal Ca^{2+} levels were inferred by monitoring Ca^{2+} release into the cytoplasm. In these experiments, we used PC12 cells stably expressing a scramble shRNA and differentiated into neurons as controls. To specifically evaluate Ca^{2+} release from the lysosome, we stimulated the cells with a synergistic combination of the lysosomal TPC2 calcium channel agonists TPC2-A1N and TPC2-A1P [13] (both at 30 μM)

(Figures 2a and b). As shown in Figure 2c, lysosomal Ca^{2+} release in KD cells was statistically

significantly lower when compared to control cells, consistent with a reduction in lysosomal

Ca^{2+} store filling when PMCA2 was defective. There was no significant difference in basal

cytosolic Ca^{2+} levels in KD vs control cells (Figure 2d). As a positive control of Fura-2

response and to check neuronal excitability of differentiated PC12, cells were stimulated with

50 mM KCl. This causes a depolarisation of plasma membrane voltage-gated calcium channels

in excitable cells and consequently Ca^{2+} influx from outside the cell into the cytosol, which can

be measured with the Fura-2 probe. Both control and KD cells produced a significant response

to KCl stimulation (Figures 2e and f), confirming the suitability of the system and that

differentiation to the neuronal phenotype had occurred. There were no significant differences

in KD vs controls in response to KCl stimulation (Figure 2g). As all transduced cells expressed

GFP, we confirmed that GFP fluorescence did not leak into the Fura-2 channel and that there

was no correlation between GFP and Fura-2 fluorescence intensities (Supplementary Figure 3).

Figure 2. Lysosomal Ca^{2+} content is reduced in *Atp2b2* KD cells. PC12 cells stably

expressing scramble or *Atp2b2* targeting shRNA were preloaded with the Ca^{2+} probe Fura-2.

Ca^{2+} signal in response to a combination of the TPC2 agonists TPC2A1N (30 μM) and

TPC2A1P (30 μM) in scramble (a) and *Atp2b2* KD cells (b). Summary data (c) quantifying the

peak change in Ca^{2+} signal in scramble and *Atp2b2* KD cells stimulated with the TPC2 agonists.

Each point represents the mean response from > 47 cells per coverslip from three independent

experiments, two coverslips per experiment. (d) Basal Fura-2 signal in scramble and *Atp2b2*

KD cells. Each point represents the mean response from > 47 cells per coverslip from five

independent experiments, two to three coverslips per experiment. Ca^{2+} signal when scramble

(e) and *Atp2b2* KD cells (f) were depolarised with KCl (50 mM). Summary data (g) quantifying

the peak change in the Ca^{2+} signal in scramble and *Atp2b2* KD cells depolarised with KCl.

Each point represents the mean response from > 47 cells per coverslip from four independent

experiments, two coverslips per experiment. Lines represent the mean and standard deviation,

*$p < 0.05$ calculated by unpaired t-test.

Several PMCA splice variants contain a lysosomal targeting motif.

To understand how PMCA2 localises to the lysosome, we explored this and other PMCA

proteins in greater depth. The PMCA family in mammals comprises four genes *ATP2B1*,

ATP2B2, *ATP2B3* and *ATP2B4* that encode PMCA1, PMCA2, PMCA3 and PMCA4 proteins

respectively. Each contains two major splice sites, A and C, which generate multiple transcripts

[1]. Strikingly, we found that splicing at site C can generate transcripts that encode a canonical

tyrosine-based lysosomal targeting motif, YEGL. Tyrosine-based motifs are well-established

in mediating lysosomal trafficking, suggesting a potential mechanism for lysosomal PMCA

localisation [14]. Mapping the splicing patterns in human and rat revealed that several isoforms,

PMCA1b, 1c, 1d; PMCA2b; and PMCA3b, 3c, and 3d, retain this lysosomal targeting motif in

their translated sequences (Figure 3a and Supplementary Figure 4). In contrast, none of the

PMCA4 variants examined contained a YEGL motif, highlighting a possible isoform-specific

divergence in organelle targeting.

Additionally, we found that the human *ATP2B1* gene contains a Coordinated Lysosomal

Expression and Regulation (CLEAR) sequence in its promoter. CLEAR sequences are specific

E-box sites in the DNA recognised by the transcription factor TFEB to promote the coordinated

transcription of many lysosomal genes and plays a key role in regulating lysosome biogenesis

[15]. *ATP2B2* contains a CLEAR sequence between exons 1 and 2 (Figure 3a). These CLEAR

sequences are also present in rat with two sequences in *ATP2B1*, upstream of exon 1 and

between exon 1 and 2 and in *ATP2B2* between exon 1 and 2 and exon 2 and 3 (Supplementary

Figure 4). The CLEAR sequence in *ATP2B2* is therefore consistent with PMCA2 having a

lysosomal localisation.

We expanded our study from human and rat to other species and, interestingly, found that for

ATP2B1, of 100 vertebrate species' genomes collated and aligned by UCSC, all but one contain

a YEGL motif. The only exception was Lamprey, whose sequence is SLYEKLEKPEAR. For

ATP2B2, of the same 100 collated and aligned vertebrate species' genomes, all but two contain

a YEGL motif. The only exceptions were duck-billed platypus YEGI and, again, Lamprey

FEGF. An example of the alignment is shown in Figure 3b. The broad presence of YEGL

across species is indicative of a fundamental conserved role for this motif.

We also found a similar tyrosine vacuolar targeting motif, YYFL, in yeast PMC1. Interestingly,

in yeast and *Dictyostelium*, there is only one gene encoding p-type Ca^{2+} ATPases, *Pmc1* and

patA, respectively, with no differential splicing, that generates a vacuolar protein, suggesting

that PMCAs may have evolved first as lysosomal proteins [16] [17].

We then generated PC12 cells that stably expressed the different splice variants of PMCA2

fused to GFP under a tetracycline-inducible cassette. As predicted, the PMCA2wa splice

variant showed a clear plasma membrane-targeted GFP signal. Interestingly, although the

PMCA2wb, PMCA2xb, and PMCA2zb variants, which include the YEGL motif, showed some

intracellular GFP localisation, they also displayed a strong plasma membrane signal when

overexpressed, in contrast to the lysosomal localisation of endogenous PMCA2 in PC12 cells

(Supplementary Figure 5).

Figure 3. Specific PMCA splice variants contain the lysosomal targeting motif YEGL.

ATP2B1 **and** *ATP2B2* **contain CLEAR sequences.** (a) PMCA site C alternative splicing in

human can include or exclude the lysosomal targeting motif YEGL. The exon structure of the

different regions affected by alternative splicing is shown for each of the four PMCA genes.

Constitutively spliced exons are indicated as grey boxes. Alternatively-inserted exons are

shown in green. Lowercase letters represent resulting splice variants. Capital letters indicate

the position of the translated stop codons. In PMCA3 and PMCA4, splice variant "e" results

from a read-through of the exon into the following intron (indicated as small white boxes). The

sizes of alternatively spliced exons are given as nucleotide numbers. Splice variants shown

have supporting sequence data in the literature, RefSeq or Ensembl at the time of submission

of this work. (b) Example of the YEGL motif alignment in the *ATP2B2* gene from different

species.

PMCA2 knockdown induces multiple NPC cellular phenotypes.

Having established that PMCA2 can localise and function in the lysosome, and that *Atp2b2*

(PMCA2) KD in cells produces a defect in lysosomal Ca^{2+}, a hallmark of NPC, we studied

whether other biochemical characteristics of NPC (Figure 4a), such as cholesterol,

glycosphingolipid (GSL) and sphingosine accumulation, occur in cells defective in PMCA2.

To study cholesterol distribution, cells stably expressing a scramble (control) or *Atp2b2* shRNA

(KD) were differentiated into neurons and incubated in lipoprotein-deficient serum (LPDS)

supplemented with low-density lipoproteins (LDL). Cells were fixed and stained with filipin,

a fluorescent molecule that binds cholesterol. Using confocal microscopy, we observed a

difference in free cholesterol distribution between the control (scramble cells) and *Atp2b2* KD

cells (Figure 4b). The staining was diffuse in control cells whereas in KD cells we observed

intracellular staining in puncta compatible with lysosomal accumulation of cholesterol, a

hallmark of NPC. Next, PC12 cells were differentiated and sphingosine and GSLs were

measured by HPLC. No differences were found in sphingosine levels (Figure 4c). However,

GM1a-Gc, GD1a, GD1b and GlcCer accumulated in *Atp2b2* KD cells (Figure 4d), but no

differences were found for GT1b (Figure 4d).

Figure 4. *Atp2b2* deficiency induces multiple NPC phenotypes but not sphingosine

storage. (a) Diagram of the proposed NPC pathogenic cascade[8]. (b) Filipin staining of free

cholesterol in scramble and *Atp2b2* KD cells, representative image of 3 independent

experiments. Sphingosine (c) and GSLs (d) concentrations in scramble vs *Atp2b2* KD cells,

(n = 6 scramble and 5 KD for sphingosine, due to the exclusion of an outlier detected using

Grubbs' test; n = 6 for GSLs). Lines represent the mean and standard deviation, *$p < 0.05$, **p

< 0.01, ***$p < 0.001$ calculated by unpaired t test.

NPC1 KO leads to reduced *Atp2b2* (PMCA2) splice variant expression in the mouse brain.

We previously reported that *Atp2b2* (PMCA2) gene expression was reduced in the cerebellum

of the *Npc1⁻ᐟ⁻* mice an NPC model that recapitulates the biochemical and phenotypic features

of the human disease [7]. To further investigate the contribution of each *Atp2b2* splice variant to

this reduction, *Atp2b2* expression was measured in the cerebellum of the *Npc1⁻ᐟ⁻* mouse. First,

using PCR and gel electrophoresis, we found that the predominant splice variant in mouse

cerebellum is PMCA2b containing the YEGL lysosomal targeting motif, followed by PMCA2a

(Figure 5a). The expression of the transcript encoding PMCA2c was too low to be detected by

PCR. To accurately measure differences in splicing in *Npc1⁻ᐟ⁻* vs wild-type mice (WT), we used

quantitative PCR (qPCR). All *Atp2b2* splice variants were downregulated in *Npc1⁻ᐟ⁻* mice

compared to WT, with a 67.1 % overall reduction (Figure 5b).

Figure 5. Npc1 deficiency reduces *Atp2b2* splice variant expression. (a) Gel electrophoresis

of PCR products showing the relative abundance of the different *Atp2b2* splice variants in the

cerebellum of 8-week-old WT mice. (b) Relative gene expression measured by qPCR of

different splice variants of *Atp2b2* in WT and *Npc1⁻ᐟ⁻* (Hom) cerebellums of 8-week-old *Npc1⁻*

ᐟ⁻ mice, $n = 4$ mice. Lines represent the mean and standard deviation, ***$p < 0.001$, ****$p <$

0.0001 calculated by unpaired t test.

Many genes causative of lysosomal storage diseases, including *NPC1*, have been found in the

heterozygous state to be over-represented in Parkinson's disease [18]. We therefore investigated

whether mRNA levels of *NPC1* and its interacting partners (including PMCA2) were altered

in Lewy body diseases.

The splicing of PMCAs is altered in Lewy body diseases with implications for PMCA

lysosomal targeting and function.

To evaluate the expression and splicing of *ATP2B2* (PMCA2) across Lewy body diseases

(LBD), we analysed data from a study applying single-nucleus (snRNA) and bulk-tissue RNA-

sequencing to human post-mortem anterior cingulate cortex tissue from 28 individuals[19]. This

cohort included four groups consisting of healthy controls, Parkinson's disease (PD),

Parkinson's disease with dementia (PDD) and dementia with Lewy bodies (DLB), based on

clinical assessments of retrospectively reviewed case records ($n = 7$ per group). Using snRNA-

seq data we identified statistically significant reductions in gene expression in several cell types

in PD, PDD and DLB (Figure 6a). Notably, downregulation of *ATP2B2* expression was

consistent in excitatory neurons across the three disorders, with less consistent downregulation

across multiple glial cell types. These included reduced expression of *ATP2B2* in

oligodendrocyte precursor cells (control vs PD, FDR-adjusted $p = 0.031$, Log2FC = -1.1),

oligodendrocytes (control vs PDD FDR-adjusted $p < 0.05$, Log2FC = -0.68; control vs DLB,

FDR-adjusted $p < 0.001$, Log2FC = -3.5), and microglia (control vs DLB, FDR-adjusted $p <$

0.01, Log2FC = -3.1).

Given these findings, we extended the analysis to included *NPC1* and other genes encoding

confirmed or potential NPC1 interacting partners based on the literature [7,20,21,22]. We also

analysed the expression of other genes implicated in the NPC pathway (Figure 6a). Of the 21

genes of interest, we found that 13 were differentially expressed in at least one cell type or

disease state (Figure 6a). Consistent with the *ATP2B2* data, most changes were seen in

excitatory neurons and oligodendrocytes when focusing on DLB. Interestingly, the gene *FTH1,*

which is involved in iron metabolism showed downregulation in a broader range of cell types.

Next, we leveraged the availability of high-depth bulk RNA-seq data, and while we identified

no significant change in the expression of PMCA genes, given the importance of splicing for

the regulation of PMCA gene function/localisation, we investigated changes in transcript use

across disease states. We assessed differential splicing of all the PMCA genes as well as genes

identified as confirmed and potential interactors of NPC1 (Figure 6b). A gene was counted as

differentially spliced if it contained one or more differentially spliced clusters within its locus.

Interestingly, we found that 10 of the 21 genes of interest were significantly differentially

spliced when control samples were compared to PD, PDD and DLB sample sets. This gene set

included *ATP2B2*, *ATP2B1* and *NPC1*, which all showed evidence of significant differential

splicing across all Lewy body diseases.

However, we were particularly interested in splicing events that could impact on the inclusion

of the YEGL motif in members of the PMCA gene family. Specifically, we assessed changes

in *ATP2B2* splicing that would affect lysosomal targeting of the protein by calculating the

difference in the inclusion (delta percent spliced in (ΔPSI)) of specific 3' junctions found

uniquely in either the variant-a and -b transcripts. We found increased use of variant-b

junctions that would be predicted to include the lysosomal targeting sequence increased in use

in PD (FDR-adjusted p < 0.001; ΔPSI=37%). In DLB we noted a significant downregulation

of this junction type in *ATP2B2* (FDR-adjusted p < 0.01; ΔPSI=-6%; Figure 6c). These splicing

changes indicate there are dynamic changes in *ATP2B2* splicing across Lewy body diseases,

and this was further evidenced by splicing changes of *ATP2B1* and *ATP2B4* in these diseases,

indicating potential alterations in lysosomal targeting (Supplementary Table 1). Together,

these findings suggest that changes in PMCA splicing and lysosomal targeting play a role in

Lewy body diseases.

Figure 6. PMCA2, NPC1 and their wider interactome gene expression and splicing are

altered in Lewy body diseases. (a) Gene expression changes using single-nucleus RNA

sequencing in post-mortem human brain samples from PD, PDD and DLB cases and controls.

Fill of the tile indicates the significant (FDR < 0.05) log2(fold change), with non-significant

results (FDR > 0.05) coloured white. (b) Gene expression and differential splicing of *NPC1*

and potential NPC1 interacting partners using bulk RNA sequencing in PD, PDD and DLB

compared to controls. The fill colour of each tile represents the −log10(FDR) for the gene (left

panel) or the most significant spliced junction cluster within each gene (right panel), with FDR

significance <0.05 indicated by an *. (c) *ATP2B2* splice variants demonstrate differential

splicing profiles in Lewy body disorders. Sashimi plots showing delta percent spliced in

comparison to controls for junctions producing alternative *ATP2B2* splice variants a and b. (a)

n = 7 patients per group. (b and c) *n* = 5 controls, *n* = 7 PD, *n* = 6 PDD and *n* = 6 DLB. PD:

Parkinson's disease, PDD: PD with dementia, DLB: dementia with Lewy bodies.

Discussion

In this study, we surprisingly found that PMCA2 localises to the lysosome. A possible

explanation for this unexpected localisation of PMCA is the presence of a canonical tyrosine-

based lysosomal targeting motif (YEGL), which we identified in specific splice variants of

PMCA genes. In addition, both PMCA1 and PMCA2 harbour CLEAR (Coordinated

Lysosomal Expression and Regulation) elements, further supporting a lysosomal role for these

isoforms. Notably, the YEGL motif was conserved across 98% of the 100 species examined,

suggesting it plays a fundamental and evolutionarily conserved role. Intriguingly, orthologues

of PMCAs in yeast and *Dictyostelium* localise to the vacuole, the equivalent of the lysosome

in these organisms, raising the possibility that plasma membrane targeting of PMCAs may have

evolved secondarily to an ancestral lysosomal localisation and function.

Interestingly, when PMCA2 splice variants containing the lysosomal targeting motif were

overexpressed, they localised not only to intracellular compartments but also to the plasma

membrane. This dual localisation suggests that intracellular retention may require interaction

with specific partner proteins in the lysosome. In the absence of sufficient binding partners,

may, by default, insert in the plasma membrane, raising the possibility that stoichiometric

balance with interacting partners such as NPC1 also affects its subcellular distribution.

Our results are consistent with the hypothesis that the function of PMCA2 at the lysosomal

membrane is to complex with NPC1, which tethers the lysosome to the ER [20] and that, then,

PMCA2 fills the lysosome with Ca^{2+} at lysosome: ER contact sites. The lysosome is a regulated

Ca^{2+} store and signalling hub. Lysosomal Ca^{2+} release is needed for vesicular trafficking,

endosome/lysosome fusion, and phagocytosis [23]. Several proteins have been described to

facilitate calcium release from the lysosome, such as the TPC2 and TRPML1 channels [23] but

the mechanisms for lysosomal Ca^{2+} filling remain incompletely understood [24]. TMEM165 has

recently emerged as a candidate [25] [26]. However, it has long been speculated that both passive

and active Ca^{2+} transporters would be required [27]. Our study presents PMCA2 as an additional

candidate for lysosomal Ca^{2+} filling, specifically at NPC1-mediated contact sites between the

lysosome and ER.

After demonstrating its lysosomal localisation and interaction with NPC1 [7], we modelled its

orientation in the limiting membrane of the lysosome. Our results using AlphaFold favoured

an orientation compatible with transporting calcium into the lysosome. Consistent with this,

when PMCA2 was stably knocked down, the release of Ca^{2+} from the lysosome induced by the

TPC2 agonists TPC2-A1-N and TPC2-A1-P was significantly reduced compared to control

cells. These results are compatible with lower levels of lysosomal Ca^{2+} when PMCA2 is

defective, supporting a role for PMCA2 in mediating lysosomal Ca^{2+} filling. However, our

measurements were indirect, relying on monitoring cytosolic Ca^{2+} upon opening of a lysosomal

ion channel as opposed to direct measurements of Ca^{2+} within the lysosomal lumen. The latter

is challenging due to the acidic interior of lysosomes. Nevertheless, lysosomal calcium filling

by PMCA2 clearly merits further investigation.

When NPC1 is defective, the first event in the pathogenic cascade is the accumulation of

sphingosine that leads to the reduction of lysosomal Ca^{2+} levels [8]. The fact that in cells with

functional NPC1, knockdown of PMCA2 does not result in the accumulation of sphingosine is

consistent with NPC1-dependent sphingosine transport being upstream of the lysosomal Ca^{2+}

defect. Our discovery that PMCA2 expression is reduced in *Npc1*-null mice links this Ca^{2+}

transporter to the NPC pathway, explaining for the first time why lysosomal Ca^{2+} levels are

reduced when NPC1 is defective. This hypothesis is further supported by the fact that when we

knockdown PMCA2, lysosomal Ca^{2+} levels are also reduced. Another link between PMCA2

and the NPC pathway is that in cells with intact NPC1 but reduced levels of PMCA2, lipids

such as cholesterol and the majority of GSLs accumulate, as observed in NPC cells. These data

suggest that the NPC1/PMCA axis is a hub regulating lysosomal Ca^{2+} and lipid homeostasis at

NPC1 mediated contact sites between the lysosome and ER.

It is interesting that mutations in PMCA2 in humans and mouse models lead to ataxia [2,3],

another characteristic phenotype of NPC. Interestingly, PMCA2 is highly expressed in the

cerebellum, particularly in Purkinje neurons, a cell type that is uniquely susceptible to

degeneration in NPC [28].

Importantly, the reduction in PMCA2 gene expression was not limited to NPC but was also

observed in several cell types in PD, PDD, and DLB. This is highly relevant as it has been

reported that a reduction in *ATP2B2* (PMCA2) increases neuronal vulnerability in *in vitro* PD

models [5]. In this study, we found that a reduction in PMCA2 leads to GSL dyshomeostasis,

that has also been reported in PD [29] [30]. Furthermore, low lysosomal Ca^{2+} levels have also been

reported in PD [9] [24]. Since mutations in NPC1 are overrepresented in PD [18], taken together with

our findings in this study, we hypothesise that the NPC1-PMCA2 complex is part of a wider

interactome in the lysosome whose dysregulation plays a key role in NPC and Lewy body

pathologies. Supporting this hypothesis, gene expression of *NPC1, ATP2B2* and several of their

confirmed and potential binding partners are also dysregulated in different brain cell types in

Lewy body diseases.

A key finding in this study was a profound PD-related change in splice variant usage in several

members of the wider NPC1 interactome. Changes in the splice variant expressed could,

potentially, change the protein localisation, regulation or function in the disease state. This

result highlights the need for more comprehensive studies that include not only total expression

but also differential splice variant usage.

In conclusion, our study uncovers for the first time a previously unrecognised, differential

splicing-dependent role for PMCA2 involved in lysosomal Ca^{2+} filling, expanding the known

functions of the PMCA family beyond its canonical activity at the plasma membrane. We show

that alternative splicing directs specific PMCA2 isoforms to the limiting membrane of the

lysosome, where they interact with NPC1 to restore luminal Ca^{2+} levels. This mechanism is

essential for lysosomal function, and its disruption results in lysosomal Ca^{2+} depletion and lipid

accumulation, features common to Niemann-Pick disease type C and Parkinson's disease.

These findings define a novel function for the NPC1–PMCA2 complex in lysosomal Ca^{2+}

homeostasis and highlight how alternative splicing can diversify the functional landscape of

well-characterised genes. More broadly, our work underscores the importance of isoform-

specific regulation in organelle biology and opens new avenues for understanding and targeting

lysosomal dysfunction in neurodegenerative diseases.

Materials and Methods

Materials

RPMI 1640 culture medium was obtained from Sigma. Foetal bovine serum (FBS), horse

serum and Glutamax for supplementing the medium were obtained from Gibco. NGF-β from

rat (N2513), and the antibiotics penicillin and streptomycin were obtained from Sigma.

Low-density lipoprotein (LDL) and lipoprotein-deficient serum (LPDS) preparation

Human LDL (1.019–1.063 kg/l) was isolated by vertical rotor density gradient

ultracentrifugation of plasma from hypercholesterolemic patients, as previously described [31].

LPDS was prepared from FBS by ultracentrifugation at a density of 1.21 kg/l [31].

Cells

PC12 cells (ATCC CRL-1721) were maintained in RPMI 1640 supplemented with 10% horse

serum, 5% FBS, 2 mM L-alanyl-L-glutamine dipeptide in 0.85% NaCl (Glutamax), 100 U/ml

penicillin and 100 U/ml streptomycin at 37 °C, in humidified atmosphere with 5% CO_2. For

differentiation to neurons, PC12 cells were plated onto poly-D-lysine-coated coverslips or

flasks and incubated in RPMI 1640 supplemented with 1% horse serum, 100 ng/ml NGF-β and

the abovementioned additives for four days. Medium was replaced on the second day.

For cholesterol distribution experiments, after differentiation, PC12 neurons were pre-treated

with RPMI 1640 supplemented with 10% LPDS for 24 h. Subsequently, LDL (40 μg/ml

cholesterol) was added to the culture. After 24 h, cells were fixed with 4% paraformaldehyde

in PBS.

Generation of stable *Atp2b2* KD and scramble PC12 cells

Lentiviral particles were prepared by transfecting HEK293T cells with Gag-Pol and pMDG

packaging plasmids, and each of four rat *Atp2b2* or scrambled shRNA plasmid (TL709157,

Origene). PC12 cells were transduced with the lentiviral constructs for 24 h. Then, we

performed a kill curve with puromycin. After puromycin selection, *Atp2b2* expression was

assessed by qPCR. Cells transduced with the shRNA leading to the greatest reduction (> 80%)

of *Atp2b2* gene expression and scrambled shRNA cells were selected for further experiments.

Animals

All experiments were conducted using protocols approved by the UK Home Office Animal

Scientific Procedures Act, 1986. All animal work followed the ARRIVE guidelines.

BALB/cNctr-Npc1^{m1N}/J also named *Npc1^{nih}* mice, were generated by heterozygote

brother/sister matings obtained from Jackson Laboratory (Charles River, UK). Mice were bred

and housed in individually ventilated cages (IVCs) under non-sterile conditions. Mice were

culled at eight weeks of age with 800 mg/kg pentobarbital IP. For biochemical analysis, mice

were transcardially perfused with ice-cold PBS and their brains removed and dissected.

Microscopy

PC12 cells were plated onto poly-D-lysine-coated coverslips and differentiated into neurons

over four days. Then cells were fixed with 4 % paraformaldehyde for 10 min and permeabilised

with 0.1 % saponin in PBS for 1 hour at room temperature (RT). Cells were blocked with 10%

horse serum in PBS for 1 hour. Primary polyclonal PMCA2 (PA1-915, Invitrogen) and

monoclonal NPC1 (ab134113, abcam) antibodies were labelled with Flexable Coralite Plus-

555 (KFA502, proteintech) and Flexable Coralite Plus-488 (KFA501, proteintech) for rabbit

antibodies following the manufacturer's instructions. Cells were probed with the conjugated

antibodies overnight at 4°C, washed and mounted on slides using Prolong Gold mounting

medium. Alternatively, after differentiation and incubation with LPDS and LDL as described

above, PC12 neurons were fixed with 4% paraformaldehyde for 10 min and exposed to filipin

(35 μg/ml in PBS) for 1 h. After washing, coverslips were mounted on slides with ProLong

Gold. Confocal images were acquired using a Leica-SP8 confocal microscope with an LD 63X

objective.

All image analysis were performed using ImageJ (FIJI) software (version 2.16.0/1.54g,

ImageJ2). For quantification of colocalisation of NPC1 and PMCA2, the JACoP plugin [32] was

used.

Tagless LysoIP

Tagless LysoIP was performed as described by Saarela et al. [12] with minor modifications. Briefly, mouse cerebellum (whole cerebellum per data point) were collected and gently homogenised with 40 strokes of an electric homogeniser in 1 ml of potassium phosphate buffer saline (KPBS) supplemented with Halt protease inhibitor cocktail (Thermo Fisher Scientific) and PMFS (Cell Signaling Technology). Samples were centrifuged at 1000 g for 2 min at 4ºC. Supernatant was collected, and 50 μl was reserved as input (whole tissue extract). The rest of the supernatant was divided into two tubes, each containing 100 μl of magnetic beads bound to antibodies against murine TMEM192 or isotype IgG. Magnetic beads were previously washed with KPBS. Supernatants were incubated with the magnetic beads for 5 min at 4ºC on an orbital shaker. After incubation, the mixture was placed on a magnet for 1 min and the supernatant was removed. The beads were washed 3 times with ice cold KPBS supplemented with the protease inhibitors. Ripa buffer (Cell Signaling Technology) plus protease inhibitors was added to the beads and input. Samples with beads were placed on a magnet, and lysates were collected for western blot.

Western blotting

Cells and lysoIP fractions were lysed in Ripa Buffer (Cell Signaling Technology) supplemented with a Halt[TM] protease inhibitor cocktail (Thermo Fisher Scientific) and PMFS (Cell Signaling Technology). Protein expression was measured using the bicinchoninic acid method (BCA Protein Assay Kit, Pierce). Lysates were separated on 4–12% SDS–PAGE gels (Thermo Fisher Scientific) and then transferred to PVDF membranes. For Western blot normalisation, we used the Revert™ 700 Total Protein (LI-COR). After destaining, membranes were blocked with TBS-Intercept blocking buffer (LI-COR) and probed with primary polyclonal anti-PMCA2 (PA1-915, Invitrogen) antibody and secondary antibodies conjugated

to IRDye 800CW (LI-COR). Fluorescence was detected using an Odyssey M scanner (LI-

COR). Acquired images were visualised and quantified with the Empiria Studio Software (LI-

COR).

qPCR

Total RNA from mouse brain was isolated using the RNAeasy plus extraction kit (Qiagen)

following the manufacturer's protocol. cDNA was synthesised using the iScript cDNA

synthesis kit (BIO-RAD). Real-time PCR amplification was performed on a LightCycler 480

II, using the PowerUp SYBER Green (Thermo Fisher Scientific), and specific primers

(Supplementary table 2). Relative expression of each gene was determined by calculating the

increase in CP (ΔCP) and normalising by the housekeeping gene Rpl37 (ribosomal protein

L37).

GSLs analysis

Glucosylceramide (GlcCer) and downstream GSLs were analysed as previously described [33]

with minor modifications. Detailed protocols are available at protocols.IO for GlcCer and

GSLs [34] determination. Briefly, GSLs were extracted from cell homogenates (200 μg of

protein) in C:M 1:2 overnight. The mixture was centrifuged, chloroform and PBS were added

to the supernatant and centrifuged. The lower phase was dried and resuspended in a small

volume of C:M 1:3 and mixed with the upper phase. GSLs were recovered using C18 Isolute

columns (100 mg, Biotage), and three quarters of the column elutant was dried and resuspended

in ceramide glycanase buffer (50 mM sodium acetate pH 5.5, 1 mg/mL sodium

taurodeoxycholate). rECGase 1 (prepared by Genscript) was added and samples incubated

overnight. The other quarter of the column elutant was also dried and resuspended in ceramide

glycanase buffer, but Cerezyme® (Genzyme, Cambridge, MA) was added and incubated for

72 h. Released lipids were anthranilic acid (2-AA) labelled and purified on Discover DPA-6S

SPE columns (SUPELCO). Lipids were eluted in H_2O and loaded 60:140 H_2O: MeCN (v/v)

for normal phase high-performance liquid chromatography (NP-HPLC). The NP-HPLC

system consisted of a Waters Alliance 2695 separations module and an in-line Waters 2475

multi λ-fluorescence detector set at Ex λ 360 nm and Em λ 425 nm. The solid phase used was

a 4.6×250 mm TSK gel-Amide 80 column (Tosoh Bioscience). A 2AA-labelled glucose

homopolymer ladder (Ludger, UK) was included to determine the glucose unit values (GUs)

for the HPLC peaks. Individual GSL species were identified by their GU values and quantified

by comparison of integrated peak areas with a known amount of 2AA-labelled BioQuant

chitotriose standard (Ludger, UK). Results for cell homogenates were normalised to protein

content, determined by the bicinchoninic acid (BCA) assay.

GSL species identification and concentration quantification were performed using the software

HPLC-RS (https://github.com/ArchercatNEO/HPLC/tree/v0.9).

Sphingosine analysis by HPLC

Detailed protocol is available at protocols.IO [35]. Briefly, sphingoid bases were extracted from

homogenised cells (0.2 mg protein) in 100 μL H_2O and spiked with an internal standard (C20,

3 μL 0.03 mM). To the homogenate, 500 μL C:M 1:2 was added and sonicated for 10 minutes,

RT. To the samples, 500 μL 1 M NaCl, 500 μL chloroform, and 100 μL 3 M NaOH were added

and incubated for 15 minutes, RT, and vortexed every 5 minutes. Samples were then

centrifuged (16 000 g/10 minutes) and the lower phase purified on SPE NH2 columns (Biotage)

pre-equilibrated with 2×1 mL chloroform and eluted with 900 μL acetone. The column elutant

was then dried down under N_2 and resuspended in 50 μL pre-warmed HPLC-grade EtOH.

Sphingoid bases were labelled with 50 μL OPA labelling solution (o-phtalaldyhyde dissolved

in 20x EtOH, $1\times$ 2-mercaptoethanol and diluted in $2000\times$ 3% boric acid pH 10.5) and incubated

in the dark at 20ºC for 20 minutes, vortexing at 10 minutes intervals. Samples were buffered

in 100 μL MeOH:5 mM Tris pH 7 9:1, centrifuged (800 g / 2 min), and the supernatant was

loaded for reverse phase HPLC. Solvent A was MeOH, solvent B was H_2O, solvent C was

MeCN, and solvent D was MeCN:H_2O 1:4. The RP-HPLC system consisted of a VWR Hitachi

Elite LaChrom HPLC system with a L-2485 fluorescence detector set at Ex λ340nm and Em

λ455nm. The solid phase used was a Chromolith Performance RP-18e 100–4.6 HPLC column

(Merck, Darmstadt, Germany). Results were normalised to protein content.

Processing of human brain RNA-seq data

FASTQ files from a publicly available dataset on Lewy body diseases (PMID: 34309761),

containing samples from the frontal cortex of 7 PD, 6 PDD, 6 DLB and 5 age-matched non-

neurological control individuals, was run through a Nextflow [36] pipeline that performed quality

control (QC) of reads and aligned the samples to the human genome and transcriptome

(GRCh38 and Gencode v41 annotations [37]). Briefly, initial QC was performed using fastp

(v0.23.2 [38]), which removed low quality reads and bases, as well as the trimming of adapter

sequences. Filtered reads were aligned to the transcriptome using Salmon (correcting for

sequence-specific, fragment GC-content and positional biases; v1.9.0 [39]) using the entire

genome as a decoy sequence. For splicing analyses, reads were aligned to the genome using

STAR (v2.7.10a, with two-pass mode). FastQC (v0.11.9 [40]) was used to generate read QC data

before and after fastp filtering. RSEQC (v4.0.0 [41]), Qualimap (v2.2.2a [42]) and Picard (v2.27.5

[43]) were used to generate alignment QC metrics. MultiQC (v1.13 [44]) was used to visualise and

collate quality metrics from all pipeline modules. Full details for the pipelines and parameters

used for each module can be found in (Brenton [45]):

https://github.com/Jbrenton191/RNAseq_splicing_pipeline.

Transcriptome aligned reads generated from Salmon were used to identify covariates. Salmon

read files were imported into R (v4.3.0) and converted into gene-level count matrices using the

tximport (v1.30.0) and DESeq2 (v1.42.1) packages.

Sample and sequencing covariates were assessed for collinearity using a custom script to

reduce the number of covariates tested. This employed the stats (v4.2.3) and Hmisc (v5.2-0)

packages to generate Spearman correlations, Kruskal-Wallis and chi-square tests to assess

relationships between covariates. Of the numeric covariates that were collinear, those with the

lowest overall correlations to all other variables were kept. VariancePartition (v1.32.5) scoring

and correlations between remaining covariates and the top 10 principal components (PCs) were

then used to assess which covariates should be controlled for in the final analysis.

DESeq2 differential gene expression

Differential gene expression analysis was performed with DESeq (v1.42.1) using covariates generated as above. Genes with 0 counts in at least one sample in all of the case or control groups, as well as any gene overlapping ENCODE blacklist regions of the genome were removed prior to analysis. Genes were classified as significantly upregulated if their Benjamini and Hochberg FDR adjusted p value was ≤ 0.05 and Log2 fold change was > 0, and significantly downregulated if their FDR ≤ 0.05 and Log2 fold change was < 0.

Single nuclear gene expression analysis of PMCA family and NPC1 interactome

Publicly available processed single nuclear gene expression results from the same samples as used for above bulk gene expression (PMID: 34309761) were downloaded and interrogated for significant changes in genes in the PMCA and NPC1 family of genes and their interaction partners. All quality control, alignment and differential expression analysis measures are described in (PMID: 34309761).

Differential splicing of bulk RNAseq data from human brain

Leafcutter (0.2.9 [46]) was used to perform differential splicing. SJ.out files containing junction reads were first converted to junction files with junctions overlapping ENCODE blacklist regions removed using a custom R script. Junction files were then clustered using the leafcutter_cluster.py script from the leafcutter authors. The following options were used to remove junctions/introns with the following criteria: junctions supported by < 30 reads across all samples, supported by $< 0.1\%$ of the total number of junction read counts for the entire cluster and larger than 1 Mb.

Pairwise group comparisons, correcting for identified covariates, were made using Leafcutter's differential splicing.R function with the following options: intron clusters with more than 40

junctions and a coverage of ≤ 20 reads in ≤ 3 samples in per group were removed, as well as

individual introns that were not found in ≥ 5 samples. Clusters were considered significant if

they had an FDR-corrected p value of ≤ 0.05.

ATP2B2 splice variant analysis was performed by comparing the delta percent spliced in or

delta usage of specific junctions representing splice variants a and b in *ATP2B2* between cases

and controls. The coordinates for a variants are start:10329125 and end: 10331983 and

start:10329125 and end: 10338176 for b variants.

Plot generation

Plots were made using ggplot (v3.5.1) and ggtranscript (v0.99.9).

Conserved Sequence Alignment

Cross-species amino acid and nucleotide alignments were inspected via the UCSC Genome

Browser (https://genome.ucsc.edu/cgi-bin/hgTrackUi?g=cons100way) [47] using hg19 human

and rn6 rat genome assemblies.

In vitro **cytosolic calcium measurements using Fura-2**

PC12 cells plated on poly-D-lysine coverslips and differentiated to neurons were loaded with

the ratiometric fluorescent Ca^{2+} indicator Fura-2 by incubation for 1 h at RT with the

acetoxymethyl ester form of Fura-2 (Fura-2 AM, 2.5 μM) and 0.005% (vol/vol) pluronic acid

(both from Invitrogen) in HEPES-buffered saline (HBS) containing 10 mM HEPES, 1.25 mM

KH_2PO_4, 2 mM $MgSO_4$, 3 mM KCl, 156 mM NaCl, 2 mM $CaCl_2$, and 10 mM glucose (all

from Sigma-Aldrich) at pH 7.4. After labelling, cells were washed three times in HBS and

mounted in an imaging chamber (Bioscience Tools) with 1 ml HBS.

Epifluorescence images were captured every 3 s with a Megapixel monochrome cooled

coupled device camera attached to an Olympus IX73 inverted fluorescence microscope that

was fitted with a 20X objective, and a Cairn CoolLED multiple wavelength LED source under

the control of MetaFluor 7.10.3.279 software. The Fura-2 ratio was recorded using dual-

excitation wavelengths of 340 and 380 nm, and a 440-nm long-pass filter was used to collect

emitted fluorescence. GFP-tagged constructs were identified by excitation at 488 nm and

capturing emitted fluorescence using a 515 nm long-pass filter.

PC12 neurons were depolarised by replacing HBS with high-potassium buffer, containing 10

mM HEPES, 1.25 mM KH_2PO_4, 2 mM $MgSO_4$, 156 mM KCl, 3 mM NaCl, 2 mM $CaCl_2$, and

10 mM glucose (all from Sigma-Aldrich) at pH 7.4 such that the final KCl concentration was

50 mM. Cells were also acutely challenged with a combination of the TPC2 agonists TPC2-

A1N and TPC2-A1P, both at 30 μM.

AlphaFold3 structure predictions

All protein models were generated using AlphaFold 3 accessed via the AlphaFold Server with

default settings (https://alphafoldserver.com/) [48]. Proteins modelled included human NPC1

(UniProt ID: O15118-1), human PMCA2 (UniProt ID: Q01814-1), yeast Ncr1 (UniProt ID:

Q12200-1), and yeast Pmc1 (UniProt ID: P38929-1). Graphical representations of the predicted

protein complex structures were prepared using PyMOL (Schrödinger LLC).

Statistical analysis

To analyse PMCA2 enrichment in the lysosomal fraction following Tagless LysoIP

experiments, we performed a one-way Anova analysis, followed by a Tukey's multiple

comparisons test with a 95% confidence interval. Normality of the data was validated using

the Kolmogorov-Smirnov test. To evaluate the effects of knocking down the expression of

Atp2b2 in PC12 neurons on Ca^{2+} levels in basal conditions and in response to TPC2 agonists

and KCl, two-tailed unpaired t tests with 95% confidence interval were performed, degree of

freedom (df) = 24, df = 10 and df = 14, respectively, using the Prism software (version 10.4.1).

The same analysis was used to evaluate the effect of knocking down *Atp2b2* on sphingosine

(df = 9) and GSLs levels (df =10) and to compare the gene expression in *Npc1*-null mice to

WT mice (df = 6).

Acknowledgement and funding

This research has received funding from the Aligning Science Across Parkinson's [ASAP-000478 (M.E.F.S., J.W.B., D.T.V., J.H., M.R. and F.P.)] through the Michael J. Fox Foundation for Parkinson's Research (MJFF) and the Dolby Foundation (J.H.). Additional funding was provided by NPSuisse, the Australian NPC Disease Foundation (ANPDF), Niemann-Pick UK (NPUK), and the Niemann-Pick Research Fund (NPRF). M.E.F.S held a Royal Society Newton International Fellow (ref NF171121). F.M.P. is a Royal Society Wolfson Research Merit Award holder.

Biorender: Figure 3 and Supplementary Figure 4 were completely or partially created using Biorender.

Author Contributions:

M.E.F.S., R.B., J.W.B., G.P., M.G.P., R.R., D.T.V., D.S., Y.W., E.A.F., L. P., L.S.P., A.M., and L.D.: Acquisition and analysis of data. M.E.F.S., R.B., M.R. and F.P.: Drafting of the manuscript. M.E.F.S. and F.P.: Conception and design of the study. D.G.C., E.E., D.A., C.P.P., A.G., S.N., J.H., S.P. and M.R. supervised experiments, provided resources and contributed to methodology. F.M.P. supervised the entire study.

Data Availability

All data supporting the findings of this study are within the paper or available from the open repository Zenodo with the identifier DOI:10.5281/zenodo.17209111.

Code availability

Code generated on this study for HPLC data analysis (HPLC-RS) is available from:

https://github.com/ArchercatNEO/HPLC/tree/v0.9

Code used to analyse transcriptomic data is available from:

https://github.com/Jbrenton191/RNAseq_splicing_pipeline.

Conflict of interests

There are no conflicts of interest relevant to the data in this manuscript declared by any of the
authors.

References

Strehler, E. E. Plasma membrane calcium ATPases: From generic Ca(2+) sump pumps to versatile systems for fine-tuning cellular Ca(2.). *Biochem Biophys Res Commun* **460**, 26-33 (2015). https://doi.org:10.1016/j.bbrc.2015.01.121

Vicario, M. *et al.* A V1143F mutation in the neuronal-enriched isoform 2 of the PMCA pump is linked with ataxia. *Neurobiol Dis* **115**, 157-166 (2018). https://doi.org:10.1016/j.nbd.2018.04.009

Kurnellas, M. P. *et al.* Molecular alterations in the cerebellum of the plasma membrane calcium ATPase 2 (PMCA2)-null mouse indicate abnormalities in Purkinje neurons. *Mol Cell Neurosci* **34**, 178-188 (2007). https://doi.org:10.1016/j.mcn.2006.10.010

Vallese, F. *et al.* The ataxia-linked E1081Q mutation affects the sub-plasma membrane Ca(2+)-microdomains by tuning PMCA3 activity. *Cell Death Dis* **13**, 855 (2022). https://doi.org:10.1038/s41419-022-05300-y

Brendel, A., Renziehausen, J., Behl, C. & Hajieva, P. Downregulation of PMCA2 increases the vulnerability of midbrain neurons to mitochondrial complex I inhibition. *Neurotoxicology* **40**, 43-51 (2014). https://doi.org:10.1016/j.neuro.2013.11.003

Kowalski, A. *et al.* Monomeric alpha-synuclein activates the plasma membrane calcium pump. *EMBO J* **42**, e111122 (2023). https://doi.org:10.15252/embj.2022111122

Colaco, A. *et al.* Unbiased yeast screens identify cellular pathways affected in Niemann-Pick disease type C. *Life Sci Alliance* **3** (2020). https://doi.org:10.26508/lsa.201800253

Lloyd-Evans, E. *et al.* Niemann-Pick disease type C1 is a sphingosine storage disease that causes deregulation of lysosomal calcium. *Nat Med* **14**, 1247-1255 (2008). https://doi.org:10.1038/nm.1876

Kilpatrick, B. S. *et al.* Endoplasmic reticulum and lysosomal Ca(2)(+) stores are remodelled in GBA1-linked Parkinson disease patient fibroblasts. *Cell Calcium* **59**, 12-20 (2016). https://doi.org:10.1016/j.ceca.2015.11.002

Wiatrak, B., Kubis-Kubiak, A., Piwowar, A. & Barg, E. PC12 Cell Line: Cell Types, Coating of Culture Vessels, Differentiation and Other Culture Conditions. *Cells* **9** (2020). https://doi.org:10.3390/cells9040958

Boczek, T. *et al.* Plasma membrane Ca2+-ATPase isoforms composition regulates cellular pH homeostasis in differentiating PC12 cells in a manner dependent on cytosolic Ca2+ elevations. *PLoS One* **9**, e102352 (2014). https://doi.org:10.1371/journal.pone.0102352

Saarela, D. *et al.* Tagless LysoIP for immunoaffinity enrichment of native lysosomes from clinical samples. *J Clin Invest* **135** (2024). https://doi.org:10.1172/JCI183592

Gerndt, S. *et al.* Agonist-mediated switching of ion selectivity in TPC2 differentially promotes lysosomal function. *Elife* **9** (2020). https://doi.org:10.7554/eLife.54712

Staudt, C., Puissant, E. & Boonen, M. Subcellular Trafficking of Mammalian Lysosomal Proteins: An Extended View. *Int J Mol Sci* **18** (2016). https://doi.org:10.3390/ijms18010047

Palmieri, M. *et al.* Characterization of the CLEAR network reveals an integrated control of cellular clearance pathways. *Hum Mol Genet* **20**, 3852-3866 (2011). https://doi.org:10.1093/hmg/ddr306

Cunningham, K. W. & Fink, G. R. Calcineurin-dependent growth control in Saccharomyces cerevisiae mutants lacking PMC1, a homolog of plasma membrane Ca2+ ATPases. *J Cell Biol* **124**, 351-363 (1994). https://doi.org:10.1083/jcb.124.3.351

17 Moniakis, J., Coukell, M. B. & Janiec, A. Involvement of the Ca2+-ATPase PAT1 and
the contractile vacuole in calcium regulation in Dictyostelium discoideum. *J Cell Sci*
112 (Pt 3), 405-414 (1999). https://doi.org:10.1242/jcs.112.3.405

18 Robak, L. A. *et al.* Excessive burden of lysosomal storage disorder gene variants in
Parkinson's   disease.   *Brain*   **140**,   3191-3203   (2017).
https://doi.org:10.1093/brain/awx285

19 Feleke, R. *et al.* Cross-platform transcriptional profiling identifies common and distinct
molecular pathologies in Lewy body diseases. *Acta Neuropathol* **142**, 449-474 (2021).
https://doi.org:10.1007/s00401-021-02343-x

20 Hoglinger, D. *et al.* NPC1 regulates ER contacts with endocytic organelles to mediate
cholesterol egress. *Nat Commun* **10**, 4276 (2019). https://doi.org:10.1038/s41467-019-
12152-2

21 Peake, K. B. & Vance, J. E. Defective cholesterol trafficking in Niemann-Pick C-
deficient   cells.   *FEBS   Lett*   **584**,   2731-2739   (2010).
https://doi.org:10.1016/j.febslet.2010.04.047

22 van den Boomen, D. J. H. *et al.* A trimeric Rab7 GEF controls NPC1-dependent
lysosomal   cholesterol   export.   *Nat   Commun*   **11**,   5559   (2020).
https://doi.org:10.1038/s41467-020-19032-0

23 Morgan, A. J., Davis, L. C. & Galione, A. Choreographing endo-lysosomal Ca(2+)
throughout the life of a phagosome. *Biochim Biophys Acta Mol Cell Res* **1868**, 119040
(2021). https://doi.org:10.1016/j.bbamcr.2021.119040

24 Feng, X. & Yang, J. Lysosomal Calcium in Neurodegeneration. *Messenger (Los Angel)*
5, 56-66 (2016). https://doi.org:10.1166/msr.2016.1055

25 Chen, R. *et al.* TMEM165 replenishes lysosomal Ca^{2+} stores, protects
cells against Ca^{2+} overload, and mediates Ca^{2+}-induced
lysosomal H^{+} leakage. *bioRxiv*, 2024.2005.2009.593345 (2024).
https://doi.org:10.1101/2024.05.09.593345

26 Zajac, M. *et al.* A mechanism of lysosomal calcium entry. *Sci Adv* **10**, eadk2317 (2024).
https://doi.org:10.1126/sciadv.adk2317

27 Morgan, A. J., Platt, F. M., Lloyd-Evans, E. & Galione, A. Molecular mechanisms of
endolysosomal Ca2+ signalling in health and disease. *Biochem J* **439**, 349-374 (2011).
https://doi.org:10.1042/BJ20110949

28 Kim, S. *et al.* Disruptive lysosomal-metabolic signaling and neurodevelopmental
deficits that precede Purkinje cell loss in a mouse model of Niemann-Pick Type-C
disease. *Sci Rep* **13**, 5665 (2023). https://doi.org:10.1038/s41598-023-32971-0

29 Huebecker, M. *et al.* Reduced sphingolipid hydrolase activities, substrate accumulation
and ganglioside decline in Parkinson's disease. *Mol Neurodegener* **14**, 40 (2019).
https://doi.org:10.1186/s13024-019-0339-z

30 Te Vruchte, D. *et al.* Glycosphingolipid Changes in Plasma in Parkinson's Disease
Independent of Glucosylceramide Levels. *Mov Disord* **37**, 2129-2134 (2022).
https://doi.org:10.1002/mds.29163

31 Calvo, D., Gomez-Coronado, D., Suarez, Y., Lasuncion, M. A. & Vega, M. A. Human
CD36 is a high affinity receptor for the native lipoproteins HDL, LDL, and VLDL. *J*
Lipid Res **39**, 777-788 (1998).

32 Bolte, S. & Cordelieres, F. P. A guided tour into subcellular colocalization analysis in
light microscopy. *J Microsc* **224**, 213-232 (2006). https://doi.org:10.1111/j.1365-
2818.2006.01706.x

33 Neville, D. C. *et al.* Analysis of fluorescently labeled glycosphingolipid-derived
oligosaccharides following ceramide glycanase digestion and anthranilic acid labeling.
Anal Biochem **331**, 275-282 (2004). https://doi.org:10.1016/j.ab.2004.03.051

34 Priestman, D. A., Weng, Y., Sabir, M., Bush, R., Wallom, K., Fernández-Suárez, M.E.,
Platt, F.M., te Vruchte, D. Analysis of glycosphingolipids from cell lines. *protocols.io*
(2024). https://doi.org:https://dx.doi.org/10.17504/protocols.io.x54v9pn14g3e/v1
35 Danielle te Vruchte, Y. W., Mylene Huebecker, Kerri-Lee Wallom, Frances M Platt.
Analysis of sphingosine, sphinganine and glucosylsphingosine from cells, animal
tissues   and   plasma.   *protocols.io*   (2025).
https://doi.org:dx.doi.org/10.17504/protocols.io.bp2l6zkz1gqe/v1
36 Di Tommaso, P. *et al.* Nextflow enables reproducible computational workflows. *Nat*
Biotechnol **35**, 316-319 (2017). https://doi.org:10.1038/nbt.3820
37 Frankish, A. *et al.* GENCODE: reference annotation for the human and mouse genomes
in   2023.   *Nucleic   Acids   Res*   **51**,   D942-D949   (2023).
https://doi.org:10.1093/nar/gkac1071
38 Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ
preprocessor.   *Bioinformatics*   **34**,   i884-i890   (2018).
https://doi.org:10.1093/bioinformatics/bty560
39 Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast
and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419
(2017). https://doi.org:10.1038/nmeth.4197
40 Andrews, S. FastQC: A Quality Control tool for High Throughput Sequence Data.
Babraham Bioinformatics (2010).
41 Wang, L., Wang, S. & Li, W. RSeQC: quality control of RNA-seq experiments.
Bioinformatics **28**, 2184-2185 (2012). https://doi.org:10.1093/bioinformatics/bts356
42 Okonechnikov, K., Conesa, A. & Garcia-Alcalde, F. Qualimap 2: advanced multi-
sample quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292-
294 (2016). https://doi.org:10.1093/bioinformatics/btv566
43 Picard. Picard2019toolkit. *Broad Institute* **v2.27.5** (2019).
44 Ewels, P., Magnusson, M., Lundin, S. & Kaller, M. MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-3048
(2016). https://doi.org:10.1093/bioinformatics/btw354
45 Brenton, J. RNAseq_splicing_pipeline: Human RNAseq Alignment and QC for
Differential   Expression   and   Splicing   Analysis.   *Zenodo*   (2024).
https://doi.org:doi:10.5281/zenodo.13379191.
46 Li, Y. I. *et al.* Annotation-free quantification of RNA splicing using LeafCutter. *Nat*
Genet **50**, 151-158 (2018). https://doi.org:10.1038/s41588-017-0004-9
47 Perez, G. *et al.* The UCSC Genome Browser database: 2025 update. *Nucleic Acids Res*
53, D1243-D1249 (2025). https://doi.org:10.1093/nar/gkae974
48 Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493-500 (2024). https://doi.org:10.1038/s41586-024-07487-
w
880
881