Systems analysis reveals neuregulin-1 control of cardiomyocyte size and shape mediated by distinct PI3K and p38 pathways

Pichayathida Luanpaisanon[1], Karen A. Ryall[1], Philip M. Tan[1], James O'Hearn[1], Laura A. Woo[1], Bryana N. Harris[1], Bethany Wissmann[1], Alexander Paap[1], Matthew Rhoads[1], Jeffrey J. Saucerman[1,2*]

[1]Department of Biomedical Engineering, University of Virginia, Charlottesville VA, 22908
[2]Division of Cardiovascular Medicine, University of Virginia, Charlottesville VA, 22908
*Corresponding author. Email: jsaucerman@virginia.edu

Highlights
-Reverse-phase protein arrays capture distinct signatures of cellular signaling in response to diverse hypertrophic ligands.
-Partial least squares regression model maps proteomic signatures to diverse patterns of cell morphology and gene expression.
- Combinatorial ligand-inhibitor screen validates predicted causal regulators of mRNAs and cell morphology
-PI3K mediates both neuregulin-1-induced elongation and cell area, validating the PLSR model. In contrast, p38 regulates cell area but not elongation.
-Logic-based model demonstrates that the characterized mechanisms are sufficient to predict how distinct PI3K and p38 dynamics drive size and shape.

Abstract
       Pathological and physiological stresses induce diverse forms of cardiac hypertrophy, with distinct manifestations in cardiomyocyte size and shape regulated by still poorly understood signaling networks. Here, we combined high-content morphological profiling, phospho-protein arrays, and systems modeling to characterize the diverse forms of hypertrophy induced by angiotensin II, endothelin-1, insulin growth factor-1, and neuregulin-1. Reverse-phase protein array profiling and partial least squares regression modeling revealed that Akt, GSK3, and MAPK signaling are differentially regulated by hypertrophic agonists and are predictive of distinct phenotypic outcomes. Neuregulin-1 uniquely induced cardiomyocyte elongation in both neonatal rat and human iPSC-derived cardiomyocytes, in addition to increasing cell area. Pharmacological perturbations demonstrated that neuregulin1-induced elongation and area expansion both require PI3K activity, whereas p38 selectively mediates cell area. A logic-based network model incorporating dual-specificity phosphatases were sufficient to capture the amplifying PI3K and transient p38 signaling dynamics driving phenotypic changes. Together, these results identify distinct signaling cascades by which neuregulin-1 coordinates cardiomyocyte size and shape, providing mechanistic insight into how hypertrophic remodeling can be differentially regulated. This systems approach provides new insight into the pathways that drive distinct forms of cardiomyocyte hypertrophy, highlighting opportunities to selectively target maladaptive remodeling in heart failure.

Keywords: cardiomyocyte, morphology, computational model, elongation, proteomics, high-content microscopy

Abbreviations
Akt, protein kinase B; AngII, angiotensin II; ANOVA, analysis of variance; BNP, brain natriuretic peptide; CITED4, Cbp/p300-interacting transactivator 4; CM, cardiomyocyte; CTGF, connective tissue growth factor; ERK, extracellular related kinase; ET1, endothelin 1; GSK3, glycogen synthase kinase 3; IGF1, insulin-like growth factor; iPSC, inducible pluripotent stem cell; MAPK,

mitogen activated protein kinase; MEK; NRCM, Nrg1, neuregulin-1; p38, PI3K, PLSR, partial

least squares regression; RPPA, reverse phase protein array.

Introduction
Cardiac hypertrophy is a compensatory response of the heart to increased workload,
characterized by an increase in cardiomyocyte size and increased cardiac mass.
Cardiomyocytes account for approximately 30% of total cell number but contribute to 70–80%
of the heart's mass [1], [2]. Hypertrophy can be a response to either pathological stresses (e.g.
myocardial infarction, hypertension) or physiological stresses (e.g. exercise, pregnancy), in
which cardiomyocytes undergo considerable changes in cell size and shape [2]. While the
signaling pathways driving cardiomyocyte size have been studied intensely, the mechanisms
governing changes in cardiomyocyte shape remain poorly understood. For example, pathologic
hypertrophy such as hypertension causes a pressure overload that increases cross-sectional
area, resulting in concentric hypertrophy [3], [4]. Other pathologic stimuli such mitral
regurgitation cause volume overload, which elicits eccentric hypertrophy through a predominate
lengthening of individual cardiomyocytes, which is key to the transition to heart failure [5], [6]. At
the same time, physiological cues such as resistance and endurance exercise induce either
concentric or eccentric hypertrophy, driven by molecular pathways that are less characterized
[7], [8].
71
Cardiomyocyte hypertrophy is sensitive to both mechanical and chemical stimuli,
including cytokines, growth factors, catecholamines, vasoactive peptides, and hormones.
Among the most well-characterized hypertrophic growth factors are angiotensin II (AngII),
endothelin-1 (ET1), neuregulin-1 (Nrg1), and insulin growth factor-1 (IGF1) [9], [10], [11], [12],
[13]. Phenotypic screens comparing across stimuli have revealed ligand-specific regulation of
cardiomyocyte size, shape, and other properties. Bass et al. performed an early developed early
high-content image analyses that revealed that of 5 stimuli that increased cardiomyocyte area,
only phenylephrine and serum increased sarcomere organization associated with physiological
hypertrophy [14]. In a subsequent screen of 15 hypertrophic agonists, Ryall et al. found that
Nrg1 induced the greatest cardiomyocyte elongation and CITED4 mRNA. Knockdown of
CITED4 further accentuated elongation, revealing an incoherent feedforward loop [13].
However, in most cases it is still unclear how interconnected signaling pathways induce distinct
cell features associated with hypertrophy, especially for cell shape.
85
To identify pathways that distinctly regulate cardiomyocyte size, shape, and gene
expression, we performed a proteomic screen under AngII, ET1, Nrg1, and IGF1 conditions
where we previously found diverse cardiomyocyte phenotypes. Proteomic analysis revealed
distinct cell states depending on treatment and timepoint. We further mapped the phospho-
protein array data to the previous phenotypic screen to infer signaling modules and pathways.
Finally, we validated the predicted mediators of Nrg1-dependent cardiomyocyte size and shape
using rodent and human iPSC-derived cardiomyocytes.
93
94 **Results**
95
Hypertrophic ligands induce diverse phosphoprotein signatures predictive of
cardiomyocyte morphology and gene expression
Previously, we performed a phenotypic screen that revealed hypertrophic ligands that
induce diverse forms of cardiomyocyte morphology and gene expression [13]. That study found
a number of new ligand-specific relationships such as an incoherent feedforward with Nrg1-
and CITED4-regulated myocyte elongation regulated by CITED4, as well as CTGF regulating
AngII-induced Bax expression [13]. However, the signaling pathways driving these responses
were not characterized. To identify signatures of protein phosphorylation and abundance that
may drive these phenotypic responses, we treated neonatal rat cardiomyocytes with Nrg1 (10
ng/mL), AngII (1 µM), ET1 (100 nM), or IGF (10 nM) as previously, but here measuring 172 total

and phosphor-protein antibody probes by reverse phase protein array (RPPA) at 1 h and 48 h.
Heatmaps summarizing this data revealed both time- and ligand-specific proteomic responses
compared to serum-free controls **(Figure 1A)**. To highlight dynamic changes that are unique
among treatment groups, from each group we selected the top 20% proteins most expressed at
1 hr and 48 hr.
Venn diagrams revealed that many responses were largely shared across ligands at 1
hr, but proteomic responses became progressively more ligand-specific by 48 hr **(Figure 21)**.
For example, Nrg1 and AngII shared 22 differentially-expressed proteins at 1 hr and none at 48
hr. ET was the only condition with a substantial set of unique proteins already at 1 hr. Pathway
enrichment using Enrichr (Reactome database) [16, 17] identified enrichment of differentially
expressed proteins associated with AKT/mTOR signaling under Nrg1 treatment, receptor-
tyrosine-kinase–related pathways and MAPK signaling for ET treatment, infection/immune-
related pathways for AngII treatment, and developmental signaling and WNT signaling for IGF
treatment **(Figure 1C)**. Exploration of differentially expressed proteins associated within each of
these enriched pathways revealed that Nrg treatment induces early activation of the Ras/MAPK
signaling—elevated phosphorylation of ERK/MAPK, p38, JNK, c-Jun, and c-Raf—at 1 hr,
whereas AKT was significantly upregulated by 48 hr (**Figure 1D**).

Figure 1. Hypertrophic ligands induce diverse changes in protein abundance and phosphorylation. (A) Hierarchical clustering of RPPA data from 172 total/phospho probes following 1 hr and 48 hr treatment with Nrg1 (10 ng/mL, R&D #396-HB), Ang II (1 µM), ET1 (100 nM), or IGF1 (10 nM) (log2 fold-change compared to control). For each timepoint, data were collected from 2 independent isolations and were subsequently averaged. (B) Venn diagrams of top 20% most responsive proteins under each treatment at 1 hr vs 48 hr. (C) Enrichr Reactome pathway enrichment analysis, showing the

131   top 3 enriched pathways with overlap genes ≥ 10. (D) Differentially expressed or phosphorylated proteins,
132   annotated by Reactome pathways most enriched for each treatment.
133
The phospho-protein array measurements in **Figure 1** identified distinct signatures of
protein abundance and phosphorylation, but these measurements alone are insufficient to
determine which proteins most closely associate with cardiomyocyte morphology and gene
expression. To address this gap, we mapped proteomic responses (**Figure 1**) to our previous
screen of cardiomyocyte morphology and gene expression [13] using a supervised machine
learning model, partial least squares regression (PLSR). PLSR provides a powerful framework
for dimensionality reduction and visualization of correlations, thereby offering insight into the
organization of signaling networks [15].
This PLSR model was created by designating the RPPA data as the predictor block (172
probes across four hypertrophic ligands 1 hr and 48 hr) and the phenotypic outcomes as the
response block (11 mRNAs and four morphology metrics [13]) (**Figure 2A**). All outputs were
centered and scaled by z-score across ligands. The first two latent variables accounted for 82%
of the total variance, enabling visualization of RPPA and phenotypic loadings together with
ligand scores in a shared latent space **(Figure 2B)**. Consistent with trends observed in the
RPPA dataset alone (**Figure 1**), each treatment induced distinct contributions in the latent
space, reflecting ligand-specific signaling programs (**Figure 2B**).
Of note, Nrg1 aligned strongly with cardiomyocyte elongation and CITED4 expression,
consistent with our prior findings [13]. Further, unsupervised k-means clustering of protein
loadings revealed a distinct module of protein expression and phosphorylation (Cluster B) that
are associated with cardiomyocyte elongation. Indeed, Cluster B is composed of many
components of the PI3K pathway including phospho-PI3K, phospho-Akt, phospho-PDK1,
phospho-GSK3, phospho-mTOR, and phospho-p70S6K. Also, within Cluster B and correlated
with elongation are several MAPK superfamily proteins such as phospho-p38, phospho-MEK1,
phospho-MAPK, and phospho-p90RSK **(Figure 2B)**.
To further dissect proteins associated with cell area vs. elongation, we computed
corresponding alignment scores. While cell area was most aligned with abundance of proteins
such as tuberin and bRaf or phosphorylation of PKC, cardiomyocyte elongation was aligned
with phosphorylation of proteins close to PI3K/Akt signaling such as GSK3α, p70S6K, Akt,
mTor, and p38 (**Figure 2C**). To assess PLSR model performance, we assessed two
complementary metrics: (i) percent variance explained (R^2) and (ii) leave-one-out cross-
validation to estimate predictive value (Q^2) (**Figure 2D**). Observed-versus-fitted analyses for
individual phenotypes demonstrated high predictive accuracy, with R^2 values of 0.99 for cell
area and 0.84 for elongation **(Figure 2E)**. Taken together, these findings demonstrate that
Nrg1-induced changes in phenotype are most strongly associated with PI3K/Akt and MAPK
signaling modules, highlighting these pathways as potential regulators of cardiomyocyte
elongation.
170

Figure 2. Partial least squares regression (PLSR) model predicts unique signaling pathways associated with cardiomyocyte size and elongation. (A) Overview of experimental design for PLSR model, in which the goal is to identify predictive relationships from protein and phospho- protein expression (predictor block) to gene expression and morphological changes (response block). Protein levels were measured by RPPA 48 h after treatment (from Figure 1), gene expression was measured by qPCR 48 h after treatment, and morphology of individual cells was tracked by fluorescent microscopy from before treatment to 48 h after treatment (from [13]). (B) PLSR loadings are plotted for protein expression (predictor block; circle markers) and for gene expression and morphology (response block; square markers). Unique signaling pathways are grouped together via K-means cluster analysis. Middle arrows show orientation and relative magnitude of PLSR scores for ligand inputs at two timepoints (1hr and 48 h after treatment). (C) Alignment scores of cell area and elongation in PLS latent space shown in panel B. (D) PLSR model performance, including percent variance explained (R^2) and leave-one-out cross-validation to estimate predictive value (Q^2). (E) Observed-versus-fitted plots for area and elongation.

MEK, PI3K, and p38 mediate distinct ligand-specific phenotypes

Given the predicted unique associations of ERK, PI3K, or p38 with Cluster B (elongation) or Cluster C (cell area) (**Figure 2C**), we hypothesized that inhibiting these

pathways would modulate ligand-induced cardiomyocyte morphologies and gene expression.
Therefore, we performed a PCR screen of 8 mRNAs that were most responsive in our previous
ligand screen [13], but this time in response to 16 combinations of ligands Nrg1, ET, or AngII
with inhibitors of MEK (0.1 µM PD325901), p38 (10 µM SB203580), or PI3K (10 µM LY294002)
(**Figure 3A**). Broadly, mRNA abundances in response to ligands alone were consistent with the
previous study [13], for example strong BNP expression and CITED4 mRNAs in response to
Nrg1 and ET1 (**Figure 3B**). These responses were amplified by p38 inhibition, while on the
other hand Nrg1-induced CITED4 expression was suppressed by inhibition of MEK or PI3K.
These findings indicate that MEK, PI3K, and p38 may contribute to our previously characterized
Nrg1-CITED4-elongation feedforward loop [13]. P38 inhibition also amplified the effects of Nrg1
on VEGF and Bcl2 (**Figure 3B**).
We next asked whether MEK, p38, or PI3K inhibitors also modulate cardiomyocyte
morphology. We again used the live-cell morphology tracking assay (representative images in
Figure 3C), which images cells transfected with a plasmid expressing GFP under a troponin T
promoter [13]. We measured cardiomyocyte area, elongation, form factor, and perimeter in
response to the same 16 combinations of Nrg1/ET/AngII ligands and MEK/PI3K/p38 inhibitors.
Overall, phenotypic responses to ligands without MEK/PI3K/p38 inhibitors were consistent with
our previous study [13], with the strongest responses being ET1-induced cell area and
perimeter as well as Nrg1-induced elongation (**Figure 3D**). In conditions of Nrg1, ET1, or AngII
treatment, PI3K inhibition was most effective in reduced cell area and perimeter. In contrast,
Nrg1-dependent elongation was suppressed by MEK or PI3K inhibition but enhanced by p38
inhibition (**Figure 3E**). Overall, these results validate the predicted causal roles of MEK, PI3K,
and p38 anticipated from the PLSR model, and further show that correlated phospho-proteins
such as PI3K and p38 have distinct causal effects on phenotype (e.g. elongation).
214

215
Figure 3. MEK, p38, and PI3K inhibition distinctly mediate ligand-induced cardiomyocyte gene
expression and morphology. A) Heatmap showing overall mRNA responses in a ligand-inhibitor screen.
Myocytes were treated for 48 h with 10 ng/mL Nrg1 (R&D #396-HB), 10 nM ET1, 1 μM AngII, or serum
free negative control, in combination with inhibitors for MEK (0.1 μM PD0325901,) p38 (10 μM
SB203580), or PI3K (10 μM LY294002). B) Quantitative changes in mRNA abundance for BNP, CITED4,
VEGF, and Bcl2, with mRNA aggregated from 2 wells for each of 3 isolations. C) Representative images
of cardiomyocytes transiently transfected with GFP under the cTnT promoter, enabling live-cell tracking of
morphology. D) Heatmap showing overall morphologic responses in the ligand-inhibitor screen. E)
Quantitative changes in cardiomyocyte morphology metrics, where each dot represents the median
single-cell response from ~200 myocytes for each of 3 isolations. Significance was assessed using a
linear-mixed effects model (treatment group as fixed effect, Isolation as random intercept). # indicates the
ligand Control differs from SF Control; * indicates an inhibitor-treated condition differs from its own
ligand's Control. Significance levels for both symbols. * or # for p < 0.05, ** or ## for p < 0.01, *** or ###
for p < 0.001.

PI3K mediates Nrg1-induced elongation and cell area, while p38 mediates Nrg1-induced
increase in cell area

While the live-cell phenotypic screen in **Figure 3** is sensitive to single-cell changes, a
limitation is that it measures only from ~5% of GFP expressing cells which may not be
representative. Therefore, to determine whether Nrg1 also induced size and shape changes in a
broader population of cells, we treated rat neonatal cardiomyocytes and two commercial
preparations of human iPSC-derived cardiomyocytes (iCell hiPSC-CMs and hiPSC-CM[2] from
FujiFilm) (**Figure 4A**). At 48 h, we fixed and immune-stained the cells for α-actinin, followed by
high-content cell segmentation and extraction of cell morphological features. By high-content
analysis, we profiled 12 single-cell size and shape features for sensitivity to Nrg1 response. Of
these, the most sensitive metric to Nrg1 was the metric Feret Elongation (**Figure 4B**). The
geometric measure Feret Diameter was developed by L.R. Feret in the 1930s to quantify overall
extent of an object in a given direction [16]. We intentionally designed Feret Elongation to reflect
the ratio of a cell's overall extent of length (Max Feret Diameter) and overall extent of width (Min
Feret Diameter), analogous to conventional manual measurements of elongation for mature
cardiomyocytes (**Figure 4C**). We also showed that Feret Elongation is more sensitive in
measuring elongation of idealized shapes compared to other shape measures that compare
cells to circles (form factor) or ellipses (eccentricity) (**Supplemental Figure 1**). Nrg1
consistently increased Feret Elongation in this fixed-cell assay that captures all cardiomyocytes
(**Figure 4D**), consistent with the live-cell assay of GFP+ transfected cardiomyocytes (**Figure
3C**). The fixed cell assay additionally showed Nrg1-dependent increase in cell area (**Figure 4E**),
which was not seen in the live assay (**Figure 3C**) perhaps due to a lower sensitivity. We further
tested whether Nrg1 induces elongation of human iPSC-derived cardiomyocytes, which was
confirmed with two commercial products FujiFilm iCell Cardiomyocytes (hiPSC-CMs) and iCell
Cardiomyocytes[2] (hiPSC-CMs[2]) (**Supplemental Figure 2**). These results demonstrate that
Nrg1-dependent cardiomyocyte elongation is a robust finding across neonatal rat and human
iPSC-derived cardiomyocytes.

Feret Elongation $= \left(\dfrac{\text{Max Feret}}{\text{Min Feret}} \right) - 1$

258
Figure 4. Neuregulin induces cardiomyocyte elongation in a more inclusive and sensitive fixed-
cell assay. (A) Representative images of neonatal rat cardiomyocytes (NRCM) at 48 h after treatment
with control or 100 ng/mL Neuregulin-1 (Nrg1, R&D #377-HB). Cardiomyocyte marker (α-actinin) and
nuclei (DAPI) are shown. (B) Profiling 12 single-cell morphological features for sensitivity to Nrg1-
treatment compared to control NRCMs, quantified as fold-change from control minus 1. (C) Schematic of
how the Feret Elongation metric works for irregular geometries, using cartoons of animal ferrets for
illustration only. Application to idealized circles, ellipses and rectangles are shown in Supplemental Figure
1. (D) Quantitative change in Feret Elongation in response to Nrg treatment in NRCM. (E) Quantitative
change in cell area in response to Nrg treatment in NRCMs. Each point (dot, triangle, or square) reflects
the mean response of ~400 cells per well, and with bars showing mean response from across 3-4

wells/treatment from three cardiomyocyte isolations. Significance was assessed by linear-mixed effects
model (Condition as fixed effect, Isolation as random intercept), with * p < 0.05, ** p < 0.01, *** p < 0.001.
271
Phospho-protein array measurements and PLSR modeling from **Figure 2** indicated that
signatures of protein phosphorylation within the PI3K/AKT and MAPK superfamilies were
closely aligned with Nrg-dependent cell elongation, which was modulated by MEK, PI3K, and
p38 inhibitors using a live-cell phenotypic screen (**Figure 3**). We next asked whether the
opposing role of PI3K and p38 inhibitors on Nrg-dependent elongation would further reproduce
in the broader cell population accessible with fixed cell immunofluorescence. To test this
question, we treated NRCMs with 10 ng/mL Nrg1 and applied pharmacological inhibitors for
PI3K (10 µM LY294002) or p38 (10 µM SB203580) for 48 h, followed by fixation,
immunofluorescence, and high-content image analysis. Consistent with our prior NRCM
experiments in complementary live-cell and fixed-cell assays, Nrg1 treatment increased both
cardiomyocyte elongation and cell size (**Figure 5A-C**). PI3K inhibition significantly attenuated
Nrg1-induced elongation and reduced overall cell area, indicating a broad role in Nrg1-mediated
hypertrophy (**Figure 5A-C**). In contrast, p38 inhibition reduced the Nrg1-dependent increase in
cell area without significantly affecting elongation (**Figure 5A-C)**. Alone, PI3K inhibition
decreased Feret elongation, while p38 inhibition did not. Neither inhibitor alone affected cell
area.
288

Figure 5. Both PI3K and p38 inhibition mitigate Nrg1-dependent cell size, but only PI3K inhibition reduces cell elongation. (A) Representative images of neonatal rat cardiomyocytes (NRCMs) under four treatments: Control, 100 ng/mL Neuregulin-1 (Nrg1, R&D #377-HB), Nrg1 + 10 μM LY294002 (PI3Ki), and Nrg + 10 μM SB203580 (p38i). Cardiomyocyte marker (α-actinin) and nuclei (DAPI) are shown. (B) Quantification of Feret elongation and cell area (C) in response to PI3K and p38 inhibition, with and without Nrg1 treatment. Each point (dot, triangle, or square) reflects the mean response of ~200 cells per well, and with bars showing mean response from across 3-4 wells/treatment from two independent cardiomyocyte isolations. Significance was assessed by linear-mixed effects model (Condition as fixed effect, Isolation as random intercept), with * p < 0.05.

Network modeling predicts signaling cascades sufficient to regulate elongation and cell area following Nrg1 simulation

The above live-cell and fixed-cell experimental studies revealed that Nrg1 induces protein changes associated with cardiomyocyte elongation and that elongation and cell area, and that these phenotypes differentially require PI3K and p38 activity. We next asked whether our experimentally measured dynamics of PI3K and p38 are sufficient to predict the distinct network dynamics of elongation and cell area, with and without inhibitors of those pathways.

To address this question, we developed a logic-based computational model to formalize our network hypothesis, integrating data from RPPA measurements, phenotypic assays, and PLSR predictions. In this network model, Nrg1 was assumed to act through ERBB receptors, activating both PI3K/Akt/elongation/area and Ras/p38/area cascades. P38 activity was further constrained by negative regulation through Dual-specificity phosphatase (DUSP), which we hypothesized would be sufficient to predict the adaptive decline in p38 phosphorylation observed experimentally. We implemented this network structure in Netflux, a logic-based differential equation modeling platform [17].

Simulations of Nrg1 treatment predicted network-wide dynamic responses, including early activation of ERBB/Ras/p38 signaling, followed by a subsequent rise in PI3K signaling and DUSP-induced decline in p38 signaling (**Figure 6A-B**). As seen in the RPPA experiments, the model predicted that Akt phosphorylation rose slowly, while p38 phosphorylation was transient (**Figure 6C**). Perturbation simulations further validated the model. As measured experimentally, PI3K inhibition attenuated Nrg1-induced elongation and decreased cell area, while p38 inhibition selectively reduced cell area response without strongly impacting elongation **(Figure 6D).** Together, these findings demonstrate that simplified network model grounded in our experimental data recapitulates the differential pathway contributions of PI3K and p38 to both cell elongation and area.

Figure 6. Network model identifies mechanisms sufficient for distinct regulation of cell size and shape. (A) Logic-based network model of Nrg1 signaling, with nodes colored based on their normalized activity at 1 hr and 48 hr following Nrg1 treatment. (B) Simulated response of the network in response to Nrg1. (C) Comparison of phosphorylation levels between model predictions and experiments (Akt phosphorylation: average of pS473 and pT308; p38 phosphorylation: pT180/Y182), normalized by min/max values per experiment (D) Comparison of cell area and elongation between model predictions and experimental results, normalized by min/max values per experiment.

Discussion

A quantitative understanding of cardiomyocyte size and shape regulation is critical for elucidating the molecular pathways that govern cardiac hypertrophy. A wide range of studies have focused on molecular regulators of cardiomyocyte size, such the calcineurin/NFAT pathway [18]. In contrast, pathways that distinctly control cardiomyocyte shape, such cardiomyocyte elongation often seen after volume overload or endurance exercise, remain underexplored. Here, we show that Nrg1 regulates both size and shape across neonatal rat cardiomyocytes in complementary live-cell and fixed-cell assays, and two lines of hiPSC-derived cardiomyocytes. Using a new metric designed for sensitivity to length and width termed Feret elongation, we identified that in addition to increasing cell area, Nrg1 is a potent inducer of cardiomyocyte elongation.

To identify pathways associated with cardiomyocyte size and shape, we performed
proteomic screens in response to hypertrophic ligands ET1, AngII, Nrg1, and IGF1. Partial Least
Squares Regression modeling revealed patterns of protein expression and phosphorylation that
are predictive of gene regulation and cell morphology. Based on the predicted association of
cardiomyocyte elongation with MEK, PI3K, and p38, we performed an additional ligand-inhibitor
screen which identified distinct roles of these kinases on myocyte morphology and gene
expression. These predictions guided subsequent perturbation experiments with more sensitive
fixed-cell high-content microscopy, which identified that Nrg1 mediated PI3K-dependent cell
elongation while both PI3K and p38 contribute to cell area. Integrating these data into a logic-
based network model demonstrated sufficient mechanisms by which sustained PI3K drives cell
elongation, while negative feedback by dual-specificity phosphatases (DUSPs) [19], [20]
attenuates p38-driven regulation of cell area. Together, these results highlight how distinct
signaling cascades differentially regulate cardiomyocyte size versus shape.

Our proteomic-phenotypic analysis aligns with prior studies. For example, we found that
ET1 induced a distinct cluster of protein responses associated with receptor tyrosine kinase and
mitogen-activated protein kinase signaling pathways (**Figure 2D**) that were predictive of
pathological markers CTGF and Bax mRNAs as well as increased cell area [13]. In contrast to
cell area, only a few pathways have been shown to drive cardiomyocyte elongation (e.g.
CT1/LIF, MEK5, β_{IV}-spectrin, endurance exercise) or protect against it (CITED4, ERK1/2) [11],
[21], [22], [23], [4], [24], [25], [26], [27], [28]. Here by live-cell imaging [13] and fixed-cell
immunofluorescence, we find that Nrg1 most substantially increased elongation. Notably, in
neurons, Nrg1 is well characterized to promote neurite outgrowth and axonal elongation [29],
underscoring its broader role regulating cell shape. Together, these findings decouple "size"
from "shape" in cardiomyocytes: PI3K drives both, whereas p38 selectively regulates area.

Although Nrg1-mediated elongation was consistently observed across neonatal rat
cardiomyocytes, hiPSC-CMs, and hiPSC-CM[2] cells, *in vivo* studies are required to determine
the extent to which these cell phenotypes and proteomic signatures drive physiological vs.
pathological eccentric hypertrophy. Still, our data demonstrates that Nrg1 induces several
responses associated with physiological hypertrophy including PI3K/Akt activation [7], CITED4
expression [30], [13], and enhanced sarcomere organization associated with increased maturity
(**Supplemental Figure 3**) [7]. Further, we found Nrg1 downregulates pathological markers
C/EBPβ, CTGF, and Bax [13]. Indeed, a number of studies in rodents and humans have shown
that Nrg1 mediates beneficial aspects of endurance exercise, which induce a physiological
eccentric hypertrophy [25], [31], [32], [33].

Nrg1 has previously been implicated in cardiac development and disease [34], [35]. In
zebrafish, Nrg1 overexpression enhances cardiomyocyte proliferation and supports
trabeculation—an essential morphogenetic process required for myocardial growth prior to
coronary vessel formation. Disruption of trabeculation results in embryonic lethality or adult-
onset dilated cardiomyopathy [36], [37]. In mammals, Nrg1-induced hypertrophy also may be
accompanied by some degree of cardiomyocyte proliferation [37], [38], [39], [39], [40], [41]. In
humans, chronic exercise has been associated with elevated Nrg1 and physiological
hypertrophy [33]. Circulating Nrg1 also rises with heart-failure severity, consistent with a
compensatory response to stress [42]. Recombinant human Nrg1 (Neucardin) has completed
phase III trials as a treatment for chronic heart failure [43], although those results are not yet
available. Phase II trials showed structural and functional improvements [44]. However, not all
findings are favorable, with some evidence that Nrg1-induced hypertrophy and ejection fraction

may be accompanied by reduced cardiac output [12]. Together, these observations underscore
a context- and time-dependent role for Nrg1: acutely adaptive, but potentially maladaptive with
chronic exposure.

These findings of distinct regulation of size and shape signaling are currently limited to
an *in vitro* context, which motivates future studies. For example, do other more pathologically-
associated stimuli of myocyte elongation (LIF, MEK5 [21], [45]) induce distinct pathway
mechanisms from Nrg1, which appears more physiological? While we tested neonatal
cardiomyocytes and two human iPSC-cardiomyocyte lines, it is of interest to expand this study
to adult cardiomyocytes and subsequently to in vivo animal models. Future studies should test
Nrg1 and downstream pathways implicated here in vivo, incorporating longitudinal analyses to
distinguish adaptive from maladaptive remodeling. Examination of Nrg1-induced cell proteomic-
phenotype signatures and mechanisms should be tested in endurance exercise [31], which
involves physiological eccentric hypertrophy and Nrg1 secretion but incompletely characterized
mechanisms. Integration of single-cell transcriptomics or spatial proteomics with machine
learning image analysis may help identify how cell size and shape are coordinated at the tissue
level.

In summary, this study identifies Nrg1 as a potent regulator of both cardiomyocyte
elongation and area, mediated through distinct signaling pathways. PI3K signaling contributes to
both elongation and cell area expansion, whereas p38 selectively drives cell area growth. These
findings reveal that cardiomyocyte size and shape can be uncoupled mechanistically, advancing
our understanding of how hypertrophic remodeling is orchestrated. Leveraging systems-level
approaches to selectively modulate maladaptive elongation while preserving adaptive growth
may open new therapeutic opportunities for the treatment of cardiac hypertrophy.
418
Methods:

<u>Cell Culture</u>
All procedures were performed in accordance with the Guide for the Care and Use of
Laboratory Animals published by the US National Institutes of Health and approved by the
University of Virginia Institutional Animal Care and Use Committee. Neonatal rat
cardiomyocytes were isolated from 1 to 2 day old Sprague Dawley rats using the Neomyts
isolation kit (Cellutron, Baltimore, MD). Myocytes were cultured on 96-well Corning CellBIND
plates coated with SureCoat (a combination of collagen and laminin, Cellutron) and in plating
media (Dulbecco's modified Eagle media, 17% M199, 10% horse serum, 5% fetal bovine
serum, 100 U/mL penicillin, and 50 mg/mL streptomycin).
Human iPSC-derived cardiomyocytes (iCell hiPSC-CMs and hiPSC-CM^2 from FujiFilm)
were cultered on 96-well Corning CellBIND plates (10,000 cells/well) coated with SureCoat and
in iCell plating media (FujiFilm). After 48 h of incubation with daily media change (iCell
Maintenance Media, FujiFilm), the medium was replaced with Williams E Medium (Life
Technologies, A1217601) supplemented with cocktail B supplement (Life Technologies,
CM4000) serum-starved for at least 4 h before treatment.
Treatment of ligands and inhibitors were performed in comparison to equimolar DMSO
negative controls. Ligands were 10 nM ET1, 1 μM Ang II, 10 nM IGF1, or either 10 or 100 ng/mL
Nrg1. Figure 1 and 3 used 10 ng/mL Nrg1 R&D #396-HB, (in vitro EC50 0.06-0.3 ng/mL),
motivated by our previous concentration response [13]. However, Nrg1 #396-HB became
unavailable, so Figure 4, Figure 5, Supplemental Figures 2 and 3 used 100 ng/mL Nrg R&D
#377-hB (in vitro EC50 2.5-12.5 ng/mL) as noted in their figure legends. Inhibition of PI3K was
performed using 10 μM LY294002, which inhibits multiple isoforms of PI3K as well as off-targets
DNA-PK and CK2 with similar ~1 μM IC50. Inhibition of p38 was performed with 10 μM

SB203580, which inhibits p38 at IC50 ~0.5 µM, which is 100-500 greater potency than other
targets LCK and GSK3β. Inhibition of MEK was performed using 0.1 µM PD0325901, which
inhibits MEK1/MEK2 with IC50 0.33 nM.
446
Live-cell imaging and PCR
Myocytes isolated and plated as above, with a density of 100,000 myocytes/well. Two
days after isolation, myocytes were transfected with GFP under a cardiac myocyte specific
troponin T promoter using Lipofectamine 2000 (Invitrogen). Two days after transfection, 5x5
mosaic images of myocytes were collected using automated imaging scripts as described
previously [13]. After imaging, myocytes were rinsed and cultured in serum free media
containing the indicated ligands and/or inhibitors. 48 hours after stimulation with hypertrophic
agonists and inhibitors, follow-up images were recorded and total RNA was purified from the
myocytes using the RNeasy Mini kit (Qiagen). Complementary DNA was synthesized from 150
ng of total RNA using the iScript cDNA synthesis kit (Bio-Rad). mRNA lavels of nine genes (Bcl-
2, Bax, C/EBPβ, CITED4, VEGF, BNP, IκB, CTGF, and GAPDH) were measured using qPCR
(Bio-Rad CFX Connect) using iTaq Universal SYBR Green Supermix (Bio-Rad), 3.75 ng of
cDNA, and 400 nM of each primer set. Data were analyzed using the comparative CT method
with efficiency correction and measurements were taken from three independent myocyte
isolations.
462
Immunofluorescent and high-content imaging
Myocytes isolated, plated (at 30,000 myocytes/well), and cultured as above. To prepare
for immunofluorescent imaging, cardiomyocytes were first fixed with 4% paraformaldehyde for
20 min and then permeabilized with 0.1% Triton-X for 15 min. Cardiomyocytes were blocked
with 1% bovine serum albumin in PBS for 1 hr, then treated with mouse anti-α-actinin primary
antibody (Sigma-Aldrich Cat#A7811, RRID:AB_476766) at a concentration of 1:200 overnight.
Cardiomyocytes were blocked with 5% goat serum in PBS for 1 hr, then Alexa Fluor-568-
conjugated goat anti-mouse secondary antibody (Thermo Fisher Scientific Cat#A11031,
RRID:AB_144696) at a concentration of 1:200 was applied for 1 hr. The cells were stained with
DAPI prior to imaging.
High-content imaging was performed on the stained cardiomyocytes using an Operetta
CLS High Content Analysis System. These images were processed using CellProfiler using a
cellular segmentation algorithm developed previously and validated to within 5% of two
independent manual segmentations [14], [46]. Median cell area was used as a representative
measure of the cell population in each well, and cells with undetectable cytoplasm were not
counted.
479
Single-cell morphology features
To quantify cardiomyocyte shape features, single-cell morphological analysis was
performed on CellProfiler-segmented images. Myocytes were filtered based on integrated
intensity and percent of neighboring cells touching to ensure inclusion of only healthy, well-
isolated cells for accurate shape quantification.
For each retained cell, 15 morphological metrics were measured, including 10 size metrics and
5 shape metrics. From two size metrics, a new shape metric was derived:
487   Feret elongation = (Max Feret Diameter/Min Feret Diameter) - 1.
488
This metric (Feret Elongation) is based on the geometric feature Feret Diameter [16].
Feret Elongation captures the degree of myocyte elongation at all angles, where a value of 0
corresponds to a perfectly non-elongated cell, and higher values indicate greater elongation.
Feret elongation is well suited to quantify myocyte elongation because it is scale-independent,
rotation-independent, and sensitive to differences between maximum length and minimum

width. Scale independence means that the metric remains unchanged if a cell increases or
decreases in size proportionally. Additionally, subtracting 1 from the Feret ratio normalizes cells
to a baseline minimum width of 1, allowing for comparisons independent of size.
497
Reverse phase protein arrays
Reverse Phase Protein Array (RPPA) analysis was performed to measure the
phosphorylation state and changes in protein levels of 172 different proteins. RPPA is a high-
throughput antibody-based method for measuring protein concentration similar to Western blots.
Neonatal rat ventricular myocytes were cultured in 24-well plates (500,000
myocytes/well). Four days after isolation, myocytes were rinsed and cultured in serum-free
media containing 10 ng/mL Nrg1 (R&D #396-HB), 10 nM ET1, 1 µM Ang II, 10 nM IGF1, or
serum (10% horse serum and 5% fetal bovine serum). Myocyte protein was isolated at two time
points: 1 hour and 48 hours following administration of the ligands according to the protocol on
the MD Anderson Cancer Center RPPA Core Facility website. Protein concentration was
quantified using the Pierce 660 nM Protein Assay Kit (Thermo Scientific). Cell lysates were
submitted to the MD Anderson Cancer Center for analysis. Data was collected from 2
independent myocyte isolations per condition, and all data points were normalized relative to
serum-free condition for protein loading.
512
Partial Least Square Regression (PLSR) modeling
Partial Least Squares Regression (PLSR) analysis was performed using MATLAB to
infer relationships between RPPA protein expression data and phenotypic screen outputs,
which included gene expression and morphological features. The predictor block consisted of
172 proteins across 4 ligands, and the response block comprised 15 phenotypic outputs (11
mRNA transcripts and 4 morphology metrics) across the same ligands. Prior to constructing the
model, both RPPA and phenotype data were log-transformed and standardized by computing z-
scores across ligands to ensure comparability and eliminate scale differences. PLSR was
implemented using MATLAB's plsregress function. To determine the optimal number of latent
components, models with up to six components were tested. The model performance was
evaluated using both the cumulative coefficient of determination (**R^2**) and predictive ability (**Q^2**).
These metrics were calculated across components and visualized in a variance-explained plot,
allowing identification of the number of components that maximized model performance while
minimizing overfitting.

All MATLAB scripts used for data preprocessing, model fitting, cross-validation, and
visualization are available on GitHub at https://github.com/saucermanlab/neuregulin.

Network model construction
A knowledge-based computational model of the distinct cardiomyocyte size and shape
signaling network was developed from known direct molecular interaction from the literature,
RPPA measurements, phenotypic assays, and PLSR predictions. We employed the logic-based
differential equation (LDE) approach [47] to build a predictive framework to explore
cardiomyocyte cell area and elongation regulation. A normalized Hill function modeled activation
of each node by its upstream reactions. We modeled pathways crosstalk by continuous gates
representing "OR" and "AND" logic [47]. The OR gates were used for reactions that modify the
node regardless of others and the AND gates for reactions affecting each other. The default
reaction parameters in the model are reaction weight (W = 1), half-maximal effective
concentration (EC_{50} = 0.5), and Hill coefficient (n = 1.223). The time constant (τ =0.1), initial
activation (Yinit = 0), and maximal activation (Ymax = 1) regulate the dynamics of signaling
nodes. In conditions with Nrg1 treatment, its input reaction weight was increased (0.02 → 0.3).
Further, time constants (tau parameters) were manually calibrated to fit the dynamics of species

activation in the experimental data (Table S1). The Netflux software (available on GitHub at https://github.com/saucermanlab/Netflux) has been used to generate a system of LDEs.

Statistics

All data are presented as the mean ± SEM. Experiments had a hierarchical design of multiple wells (the experimental unit, randomly assigned to treatment conditions) for each cell isolation (from a neonatal litter composed of 12-16 pups) [48]. Thus, we used linear mixed models [49] accounting for this design with wells as fixed effects and isolations as random effects, with the number of wells and isolations provided in each figure legend. For single-cell measurements, individual cell measurements for each condition are provided in Supplemental Table 3. For all analyses, a P-value < 0.05 was considered statistically significant. For all analyses, a *P*-value < 0.05 was considered statistically significant. Statistical outputs from linear mixed effects models are shown in the Supplemental Text.

Network model, reverse phase protein array, and microscopy data are deposited in FigShare at: https://figshare.com/s/71baf954c2f3766cfc2c (private link to be replaced on publication)

Acknowledgments:
We thank Dr. Marieke Jones (University of Virginia School of Medicine) for statistical consultation and Dr. Alexander Clark (University of Virginia, Department of Biomedical Engineering) for feedback on the manuscript.

Disclosures:
None

Funding:
This study was supported by the National Institutes of Health R01HL162925, R01HL160665, R01HL172417 (to JJS), T32GM136615 (to PL), T32GM139787 (to PMT), and the National Science Foundation Predoctoral Fellowships (to KAR, BNH).

References:

[1] Y. K. Tham, B. C. Bernardo, J. Y. Y. Ooi, K. L. Weeks, and J. R. McMullen, "Pathophysiology of cardiac hypertrophy and heart failure: signaling pathways and novel therapeutic targets," *Arch Toxicol*, vol. 89, no. 9, pp. 1401–1438, Sep. 2015, doi: 10.1007/s00204-015-1477-x.

[2] P. Haftbaradaran Esfahani *et al.*, "Cell shape determines gene expression: cardiomyocyte morphotypic transcriptomes," *Basic Res Cardiol*, vol. 115, no. 1, p. 7, Jan. 2020, doi: 10.1007/s00395-019-0765-7.

[3] N. López, J. Díez, and M. A. Fortuño, "Differential hypertrophic effects of cardiotrophin-1 on adult cardiomyocytes from normotensive and spontaneously hypertensive rats," *Journal of Molecular and Cellular Cardiology*, vol. 41, no. 5, pp. 902–913, Nov. 2006, doi: 10.1016/j.yjmcc.2006.03.433.

[4] I. Kehat *et al.*, "Extracellular Signal-Regulated Kinases 1 and 2 Regulate the Balance Between Eccentric and Concentric Cardiac Growth," *Circulation Research*, vol. 108, no. 2, pp. 176–183, Jan. 2011, doi: 10.1161/CIRCRESAHA.110.231514.

[5] D. D. Hall *et al.*, "Ectopic expression of Cdk8 induces eccentric hypertrophy and heart failure," *JCI Insight*, vol. 2, no. 15, p. e92476, Aug. 2017, doi: 10.1172/jci.insight.92476.

[6] Y. Abe *et al.*, "Leptin induces elongation of cardiac myocytes and causes eccentric left ventricular dilatation with compensation," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 292, no. 5, pp. H2387–H2396, May 2007, doi: 10.1152/ajpheart.00579.2006.

[7] M. Nakamura and J. Sadoshima, "Mechanisms of physiological and pathological cardiac hypertrophy," *Nat Rev Cardiol*, vol. 15, no. 7, Art. no. 7, Jul. 2018, doi: 10.1038/s41569-018-0007-y.

[8] J. H. Van Berlo, M. Maillet, and J. D. Molkentin, "Signaling effectors underlying pathologic growth and remodeling of the heart," *J. Clin. Invest.*, vol. 123, no. 1, pp. 37–45, Jan. 2013, doi: 10.1172/JCI62839.

[9] J. D. Molkentin and G. W. Dorn Ii, "Cytoplasmic Signaling Pathways That Regulate Cardiac Hypertrophy," *Annu. Rev. Physiol.*, vol. 63, no. 1, pp. 391–426, Mar. 2001, doi: 10.1146/annurev.physiol.63.1.391.

[10] D. Zablocki and J. Sadoshima, "Solving the Cardiac Hypertrophy Riddle: The Angiotensin II–Mechanical Stress Connection," *Circulation Research*, vol. 113, no. 11, pp. 1192–1195, Nov. 2013, doi: 10.1161/CIRCRESAHA.113.302501.

[11] R. R. Baliga *et al.*, "NRG-1-induced cardiomyocyte hypertrophy. Role of PI-3-kinase, p70 ^{S6K}, and MEK-MAPK-RSK," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 277, no. 5, pp. H2026–H2037, Nov. 1999, doi: 10.1152/ajpheart.1999.277.5.H2026.

[12] M. Zurek *et al.*, "Neuregulin-1 Induces Cardiac Hypertrophy and Impairs Cardiac Performance in Post–Myocardial Infarction Rats," *Circulation*, vol. 142, no. 13, pp. 1308–1311, Sep. 2020, doi: 10.1161/CIRCULATIONAHA.119.044313.

[13] K. A. Ryall, V. J. Bezzerides, A. Rosenzweig, and J. J. Saucerman, "Phenotypic screen quantifying differential regulation of cardiac myocyte hypertrophy identifies CITED4 regulation of myocyte elongation," *Journal of Molecular and Cellular Cardiology*, vol. 72, pp. 74–84, Jul. 2014, doi: 10.1016/j.yjmcc.2014.02.013.

[14]   G. T. Bass *et al.*, "Automated image analysis identifies signaling pathways regulating distinct signatures of cardiac myocyte hypertrophy," *Journal of Molecular and Cellular Cardiology*, vol. 52, no. 5, pp. 923–930, May 2012, doi: 10.1016/j.yjmcc.2011.11.009.

[15]   K. A. Janes, J. G. Albeck, S. Gaudet, P. K. Sorger, D. A. Lauffenburger, and M. B. Yaffe, "A Systems Model of Signaling Identifies a Molecular Basis Set for Cytokine-Induced Apoptosis," *Science*, vol. 310, no. 5754, pp. 1646–1653, Dec. 2005, doi: 10.1126/science.1116598.

[16]   L. R. Feret, *La Grosseur des grains des matières pulvérulentes*. Eidgen. Materialprüfungsanstalt a. d. Eidgen. Technischen Hochschule, 19??

[17]   A. P. Clark, M. Chowkwale, A. Paap, S. Dang, and J. J. Saucerman, "Logic-based modeling of biological networks with Netflux," *PLoS Comput Biol*, vol. 21, no. 4, p. e1012864, Apr. 2025, doi: 10.1371/journal.pcbi.1012864.

[18]   B. J. Wilkins and J. D. Molkentin, "Calcineurin and cardiac hypertrophy: Where have we been? Where are we going?," *The Journal of Physiology*, vol. 541, no. 1, pp. 1–8, May 2002, doi: 10.1113/jphysiol.2002.017129.

[19]   M. Auger-Messier *et al.*, "Unrestrained p38 MAPK Activation in *Dusp1/4* Double-Null Mice Induces Cardiomyopathy," *Circulation Research*, vol. 112, no. 1, pp. 48–56, Jan. 2013, doi: 10.1161/CIRCRESAHA.112.272963.

[20]   A. Barajas-Espinosa, A. Basye, M. G. Angelos, and C.-A. Chen, "Modulation of p38 kinase by DUSP4 is important in regulating cardiovascular function under oxidative stress," *Free Radical Biology and Medicine*, vol. 89, pp. 170–181, Dec. 2015, doi: 10.1016/j.freeradbiomed.2015.07.013.

[21]   R. L. Nicol, "Activated MEK5 induces serial assembly of sarcomeres and eccentric cardiac hypertrophy," *The EMBO Journal*, vol. 20, no. 11, pp. 2757–2767, Jun. 2001, doi: 10.1093/emboj/20.11.2757.

[22]   K. C. Wollert *et al.*, "Cardiotrophin-1 Activates a Distinct Form of Cardiac Muscle Cell Hypertrophy," *Journal of Biological Chemistry*, vol. 271, no. 16, pp. 9535–9545, Apr. 1996, doi: 10.1074/jbc.271.16.9535.

[23]   M. Abdul-Ghani *et al.*, "Cardiotrophin 1 stimulates beneficial myogenic and vascular remodeling of the heart," *Cell Res*, vol. 27, no. 10, pp. 1195–1215, Oct. 2017, doi: 10.1038/cr.2017.87.

[24]   J. Davis *et al.*, "A Tension-Based Model Distinguishes Hypertrophic versus Dilated Cardiomyopathy," *Cell*, vol. 165, no. 5, pp. 1147–1159, May 2016, doi: 10.1016/j.cell.2016.04.002.

[25]   U. Wisløff, J. Helgerud, O. J. Kemi, and Ø. Ellingsen, "Intensity-controlled treadmill running in rats: V̇ $O_{2\ max}$ and cardiac hypertrophy," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 280, no. 3, pp. H1301–H1310, Mar. 2001, doi: 10.1152/ajpheart.2001.280.3.H1301.

[26]   M. H. Hastings *et al.*, "Animal Models of Exercise From Rodents to Pythons," *Circulation Research*, vol. 130, no. 12, pp. 1994–2014, Jun. 2022, doi: 10.1161/CIRCRESAHA.122.320247.

[27]   M. H. Hastings *et al.*, "Intrinsic and Extrinsic Contributors to the Cardiac Benefits of Exercise," *JACC: Basic to Translational Science*, vol. 9, no. 4, pp. 535–552, Apr. 2024, doi: 10.1016/j.jacbts.2023.07.011.

[28]   D. M. Nassal *et al.*, "The βIV-Spectrin/STAT3 Complex Regulates the Orientation of Cardiac Hypertrophic Growth," *Circ Res*, vol. 137, no. 11, pp. 1316–1332, Nov. 2025, doi: 10.1161/CIRCRESAHA.125.326378.

[29]   L. Mei and K.-A. Nave, "Neuregulin-ERBB Signaling in the Nervous System and Neuropsychiatric Diseases," *Neuron*, vol. 83, no. 1, pp. 27–49, Jul. 2014, doi: 10.1016/j.neuron.2014.06.007.

[30]   C. Lerchenmüller *et al.*, "CITED4 gene therapy protects against maladaptive cardiac remodeling after ischemia/reperfusion injury in mice," *Molecular Therapy*, vol. 32, no. 10, pp. 3683–3694, Oct. 2024, doi: 10.1016/j.ymthe.2024.07.018.

[31]   S. A. Khetarpal, H. Li, C. Lerchenmüller, J. Rhee, and A. Rosenzweig, "Molecular Mediators of the Cardiac Benefits of Exercise," *Circulation Research*, vol. 137, no. 2, pp. 163–183, Jul. 2025, doi: 10.1161/CIRCRESAHA.125.325762.

[32]   M.-X. Cai *et al.*, "Exercise training activates neuregulin 1/ErbB signaling and promotes cardiac repair in a rat myocardial infarction model," *Life Sciences*, vol. 149, pp. 1–9, Mar. 2016, doi: 10.1016/j.lfs.2016.02.055.

[33]   R. Mohammadzadeh, M. R. Zolfaghari, S. Nikanfar, K. Khademvatani, P. Zolfaghar Didani, and A. Fattahi, "Association of Neuregulin-1β with Physiological Cardiac Hypertrophy Following Acute and Chronic Exercise in Athlete and Non-Athlete Women," *Hum Physiol*, vol. 48, no. 1, pp. 102–107, Feb. 2022, doi: 10.1134/S036211972201011X.

[34]   O. Odiete, M. F. Hill, and D. B. Sawyer, "Neuregulin in Cardiovascular Development and Disease," *Circulation Research*, vol. 111, no. 10, pp. 1376–1385, Oct. 2012, doi: 10.1161/CIRCRESAHA.112.267286.

[35]   Y. Wang *et al.*, "Neuregulin-1, a potential therapeutic target for cardiac repair," *Front. Pharmacol.*, vol. 13, p. 945206, Aug. 2022, doi: 10.3389/fphar.2022.945206.

[36]   S. J. Rasouli and D. Y. R. Stainier, "Regulation of cardiomyocyte behavior in zebrafish trabeculation by Neuregulin 2a signaling," *Nat Commun*, vol. 8, no. 1, p. 15281, May 2017, doi: 10.1038/ncomms15281.

[37]   M. Gemberling, R. Karra, A. L. Dickson, and K. D. Poss, "Nrg1 is an injury-induced cardiomyocyte mitogen for the endogenous heart regeneration program in zebrafish," *eLife*, vol. 4, p. e05871, Apr. 2015, doi: 10.7554/eLife.05871.

[38]   G. D'Uva *et al.*, "ERBB2 triggers mammalian heart regeneration by promoting cardiomyocyte dedifferentiation and proliferation," *Nat Cell Biol*, vol. 17, no. 5, Art. no. 5, May 2015, doi: 10.1038/ncb3149.

[39]   K. Bersell, S. Arab, B. Haring, and B. Kühn, "Neuregulin1/ErbB4 Signaling Induces Cardiomyocyte Proliferation and Repair of Heart Injury," *Cell*, vol. 138, no. 2, pp. 257–270, Jul. 2009, doi: 10.1016/j.cell.2009.04.060.

[40]   B. D. Polizzotti *et al.*, "Neuregulin stimulation of cardiomyocyte regeneration in mice and human myocardium reveals a therapeutic window," *Sci. Transl. Med.*, vol. 7, no. 281, Apr. 2015, doi: 10.1126/scitranslmed.aaa5171.

[41]   C. L. Galindo *et al.*, "Anti‐Remodeling and Anti‐Fibrotic Effects of the Neuregulin‐1β Glial Growth Factor 2 in a Large Animal Model of Heart Failure," *JAHA*, vol. 3, no. 5, p. e000773, Sep. 2014, doi: 10.1161/JAHA.113.000773.

[42]   B. Ky *et al.*, "Neuregulin-1β Is Associated With Disease Severity and Adverse Outcomes in Chronic Heart Failure," *Circulation*, vol. 120, no. 4, pp. 310–317, Jul. 2009, doi: 10.1161/CIRCULATIONAHA.109.856310.

[43]   "Neucardin," *ClinicalTrials.gov*. U.S. National Library of Medicine. [Online]. Available:
https://clinicaltrials.gov/search?intr=neucardinClinicalTrials.gov
[44]   A. K. Geissler and D. B. Sawyer, "Neuregulin-1," *JACC: Basic to Translational Science*,
vol. 8, no. 9, pp. 1177–1179, Sep. 2023, doi: 10.1016/j.jacbts.2023.06.002.
[45]   N. Takahashi *et al.*, "Hypertrophic responses to cardiotrophin-1 are not mediated by
STAT3, but via a MEK5-ERK5 pathway in cultured cardiomyocytes," *Journal of Molecular*
and Cellular Cardiology, vol. 38, no. 1, pp. 185–192, Jan. 2005, doi:
10.1016/j.yjmcc.2004.10.016.
[46]   T. G. Eggertsen and J. J. Saucerman, "Virtual drug screen reveals context☐dependent
inhibition of cardiomyocyte hypertrophy," *British J Pharmacology*, vol. 180, no. 21, pp.
2721–2735, Nov. 2023, doi: 10.1111/bph.16163.
[47]   M. J. Kraeutler, A. R. Soltis, and J. J. Saucerman, "Modeling cardiac β-adrenergic
signaling with normalized-Hill differential equations: comparison with a biochemical
model," *BMC Syst Biol*, vol. 4, no. 1, p. 157, Dec. 2010, doi: 10.1186/1752-0509-4-157.
[48]   S. E. Lazic, C. J. Clarke-Williams, and M. R. Munafò, "What exactly is 'N' in cell
culture and animal experiments?," *PLOS Biology*, vol. 16, no. 4, p. e2005282, Apr. 2018,
doi: 10.1371/journal.pbio.2005282.
[49]   J. J. Faraway, *Extending the Linear Model with R: Generalized Linear, Mixed Effects and*
Nonparametric Regression Models, Second Edition, 2nd ed. New York: Chapman and
Hall/CRC, 2016. doi: 10.1201/9781315382722.
727

729
730 **Supplemental Figure 1.** Quantificationsusing cell morphological features in idealized shapes;
731 circle, ellipse, rectangle, and triangle
732

733
Supplemental Figure 2. Neuregulin-1 induces elongation of human iPSC-derived
cardiomyocytes. (A) Representative images of hiPSC iCell cardiomyocytes and hiPSC iCell
Cardiomyocytes² at 48 h after treatment with control or 100ng/mL Neuregulin (Nrg1, R&D #377-
HB). Cardiomyocyte marker (α-actinin) and nuclei (DAPI) are shown. (B) Change in Feret
Elongation in response to Nrg treatment in hiPSC-CM, and hiPSC-CM². (C) Change in cell area
in response to Nrg treatment in hiPSC-CM, and hiPSC-CM². Each point (dot or triangle) reflects
the mean of all cells per image, and the bar reflects mean across images. Asterisks denote p <
0.05 from Student's t-tests. "ns" denotes not significant.

Control   Neuregulin-1

742
743   **Supplemental Figure 3:** Nrg1-treated neonatal rat cardiomyocytes exhibited formation of
744   organized sarcomeres with visible myofibrils. 100 ng/mL Nrg1 (R&D #377-HB).

745