Integrated analysis reveals neuro-immune pathway in the central nervous system

that supports SGLT2i's protective effects in treatment of cardiac remodeling

First Author: Miao Yuan[1],

Author 1: Hanxue Wu[2],

Author 2: Jiawei Wang[2],

Author 3: Zihan Qiu[2],

Author 4: Kexin Li[1],

Author 5 (co-corresponding author): Jiaxi Xu[1,2,3]

Author 6 (corresponding author): Dengfeng Gao[1],

1 Department of Cardiology, Second Affiliated Hospital of Xi'an Jiaotong University,

Xi'an 710004, Shaanxi, China.

2 Department of Physiology and Pathophysiology, Xi'an Jiaotong University, Xi'an

710061, Shaanxi, China.

3 Key Laboratory of Neural and Vascular Biology, Ministry of Education, Hebei

Medical University, Shijiazhuang, Hebei, 050017, China.

Dengfeng Gao & Jiaxi Xu are corresponding author of this article. They are all

responsible for communicating with the other authors about progress, submissions of

revisions and final approval of proofs.

Dengfeng gao: gaomedic@mail.xjtu.edu.cn

Jiaxi Xu: xujiaxi@xjtu.edu.cn

Abstract

Background:

Growing evidence are showing beneficial effects of sodium glucose transport protein

2 inhibitors (SGLT2i) in treatment of heart failure, but underlying neurogenic

mechanism remains unclear. In this study the effect of empagliflozin (EM) on

sympatho-excitation and potential neurogenic mechanism for EM's therapeutic

effects on cardiac remodeling were studied.

Methods: Deoxycorticosterone acetate (DOCA)-salt and high-salt (8%) diet (HSD)

mouse models were utilized. Single-cell RNA sequencing was used to explore the

mechanism by which SGLT2 inhibitors improve cardiac remodeling in hypertension.

Meanwhile, blood samples were collected from hospitalized patients diagnosed with

heart failure to verify the results of animal studies.

Results: In DOCA-salt or HSD treated mice, EM was associated with a protective,

blood pressure-independent effect on cardiac remodeling. Both DOCA-salt and HSD

induced sympatho-excitation, together with neuronal hyper-activity in the pre-

autonomic regions of brain, and these were blunted in mice with EM co-treatment.

Additionally, single-nucleus RNA sequencing using hypothalami indicated that

cellular interplays among the vessels, microglia and inhibitory neurons were involved

in the disease- and EM-associated actions. Further analysis of microglia pinpointed a

close involvement of peripheral immune activation in disease-associated state

transformation of microglia, during DOCA-salt or HSD treatment, including

increased lymphocytes count and plasma level of interferon-γ. Differentially

expressed genes in neurons highlighted that EM abolished disease-associated

upregulation of protein ubiquitination, which might support imbalance of

presympathetic excitatory/inhibitory tones, and vasopressin production. In patients'

blood samples, EM was associated with significant elevation of hematocrit value in

all groups, and reduction of lymphocytes counts in the patients with high NT-proBNP

value (> 2550 pg/mL, no diuretic co-treatment).

Conclusions: Our data provide a neuro-immune pathway by which EM blunts

disease-associated cardiac sympathetic tone and hypertrophic remodeling.

Keywords: Hypertension; Sodium glucose transport protein 2 inhibitors;

Central nervous system; Cardiac remodeling

1. Background:

Cardiac remodeling refers to adaptive changes of the heart, in the perspective of

structure and systolic function, following physiological or pathological stimuli[1].

Recently, sodium-glucose co-transporter 2 inhibitors (SGLT2i) have shown beneficial

effect in the treatment of heart failure[2-4]. The mechanism of action of SGLT2i has

been extensively studied, including its impact on cardiac metabolism, energy

consumption, and direct effects on cardiac structure and function[5, 6].

Although multiple mechanisms of action of SGLT2i have been reported, the

possibility of SGLT2i acting through the autonomic nervous system, especially the

sympathetic nervous system, has not been fully dissected. The sympathetic nervous

system plays a key role in the occurrence and development of cardiac diseases[7]. It

affects heart contractility and heart rate by releasing catecholamines such as

epinephrine and norepinephrine. Beyond the heart, sympathetic nerves also innervate

other organs that contribute to pathological development of cardiac remodeling, such

as the kidney and spleen[8-13]. On one hand, elevation of renal sympathetic activity

reduces renal blood flow, induces release of renin and aldosterone, thus increases

water-sodium retention. On the other, elevation of sympathetic tone in the immune

organs, leads to deployment of lymphocytes from the spleen[13], and enhances

myeloid cell generation and leukocyte output in the bone marrow[14, 15]. Renin-

angiotensin-aldosterone system (RAAS) has been shown to induce pro-inflammatory

transformation of leukocytes and aggregation of T lymphocytes[16], to exacerbate

inflammatory responses. Within the brain, the paraventricular nucleus of

hypothalamus (PVN), a pre-autonomic center, has high capillary density, thin vessel

diameter, and complex vascular topology[11, 17-21], providing a possible interface

for immune cells and cytokines to further boost sympathetic outflow. Accordingly,

integrated effect of blunted sympatho-excitation to these organs could decrease the

cardiac afterload, support the reduction of cardiac energy consumption, and attenuate

inflammation, therefore blunting pathological development of cardiac remodeling.

Interestingly, SGLT2i increases urinary sodium excretion, reduces renal blood flow,

stimulates short-term renin secretion, but long-term impact on renin and aldosterone

is not observed[22, 23]. Clinically, SGLT2i is often applied with diuretics or RAAS

inhibitors[23]. Investigating how SGLT2i alone affect central regulation of

sympathetic tone would provide novel insight regarding its mechanism of action in

cardiac remodeling, and a new perspective on its clinical application.

In this study, via integrated analysis of deoxycorticosterone acetate (DOCA)-salt

and high-salt (8%) diet (HSD) mouse models, together with blood samples collected

from hospitalized patients diagnosed with heart failure, a pathway involving

communication between the peripheral immune system and central nervous system

(CNS), from leukocytes to neurons locating in the pre-autonomic center, was studied

as potential neurogenic mechanisms for SGLT2i's beneficial role in treatment of

cardiac remodeling.

2. Results:

3.1 Empagliflozin ameliorates cardiac remodeling

In this study, the therapeutic effect of empagliflozin (EM) was studied in both DOCA-salt model and 8% high-salt diet model (HSD). DOCA-salt model (3-week) was utilized for its character that induces salt-sensitive type of hypertension and does not elevate circulating renin and angiotensin levels[24-27], while HSD (8-week) was used because it increases salt-intake but cannot induce hypertension in C57BL6/J mice[28-30]. Meanwhile, hydralazine was provided to a subgroup of DOCA-salt-treated mice (DOCA+Hyd) to evaluate the impact of blood pressure. Firstly, there was no significant difference in body weight among the five groups of mice received treatment of DOCA-salt, DOCA+Hyd, DOCA+EM, sham, and sham+EM, respectively (Supplementary Figure1 A). Similarly, no significant difference in body weight was observed among the groups with HSD or normal salt diet (NSD) treatment (Supplementary Figure1 B). EM co-treatment did not affect drinking water consumption in DOCA-salt-treated mice but slightly increase water intake in mice with HSD (Supplementary Figure1 C-F). In DOCA-salt treated mice, co-treatment of EM did not alter either systolic or diastolic BP, mice with DOCA+EM treatment still exhibited hypertension. Unlike EM, co-treatment of hydralazine markedly lower BP in DOCA-salt-treated mice. HSD did not induce hypertension, and EM co-treatment did not alter BP either (Supplementary Figure1 G-N).

In both DOCA-salt and HSD models, cardiac remodeling was determined.

Firstly, increase in ratios of heart weight to body weight or tibia length were observed

in both models, indicating cardiac hypertrophy, and co-treatment of EM significantly

attenuate modeling-associated increase of heart weight ratios. Notably, co-treatment

of hydralazine reduced heart weight ratios, in comparison with control DOCA-salt-

treated mice, but their heart weight ratios still significantly higher than mice with EM

co-treatment, which showed similar levels as the sham controls (Figure 1A, B).

Meanwhile, HSD also induced increase in ratios of heart weight to body weight or

tibia length (Figure 1C, D), while co-treatment of EM significantly lowered these

ratios. Additionally, cardiac remodeling was also determined by tissue staining in left

ventricle (LV) and echocardiography. Both DOCA-salt and HSD treatments induced

hypertrophy of cardiomyocytes, increase of myocardial interstitial and perivascular

fibrosis in the LV (Figure 1E, F), heavier LV mass (Figure 1H, J), as well as

significant Collagen III deposition (Figure 1K, L). DOCA-salt treatment also

significantly impaired cardiac systolic function, showing reduced ejection fraction of

LV (Figure 1G). Co-treatment of EM significantly blunted LV hypertrophy, reduced

LV fibrosis and Collagen III deposition, and importantly preserved cardiac systolic

function in both models. Hydralazine did not exhibit similar protective effect as EM

co-treatment in the perspective of abolishing pathological cardiac remodeling (Figure

1G, I). To this end, we showed that EM co-treatment has protective effect in

pathological cardiac remodeling induced by DOCA-salt or HSD treatment, and this is

not dependent on potential anti-hypertensive effect of EM.

EM increases urinary sodium excretion, so we examined the plasma sodium

level and osmolality in each group of mice, to determine whether EM inhibits cardiac

remodeling by reducing blood sodium or osmolality. No marked difference was

detected regarding the plasma sodium level and osmolality among the

groups/treatments (Supplementary Figure2 A-D), but DOCA-salt treatment

significantly reduced the value of hematocrit (HCT). HCT values can be utilized to

represent changes in blood volume, and its reduction in DOCA-salt-treated mice

suggests increase of blood volume (Supplementary Figure2 E). Co-treatment of EM,

rather than hydralazine, recovered the HCT value to the baseline levels. we tested the

HCT of different groups. Unlike DOCA-salt model, HSD did not induce alteration in

the value of HCT (Supplementary Figure2 F).

3.2 Empagliflozin blunts sympatho-excitation

To evaluate sympatho-excitation, the levels of urinary norepinephrine (NE),

which had been normalized by creatinine (Cr) concentrations, were firstly determined.

Additionally, we analyzed urinary, which was used to standardize the urinary NE. As

reported previously, DOCA-salt treatment dramatically elevated urinary NE level

(Figure2 A), which was not affected by co-treatment of hydralazine, but significantly

lowered by co-treatment of EM. Similar to DOCA-salt treatment, HSD boosted

urinary NE level as well, and co-treatment of EM reduced its level (Figure2 B). In

addition to global index of sympatho-excitation, expression of tyrosine hydroxylase

(TH), a rate-limiting enzyme for NE biosynthesis, was determined to assess

sympathetic tone in the LV. DOCA-salt treatment significantly increase TH

expression in the LV (Figure2 C-D), which was significantly blunted by co-treatment

of EM. Similar as DOCA-salt treatment, HSD increased LV TH expression as well,

and this was attenuated by co-treatment of EM (Figure2 E-F).

In the pre-autonomic centers, the PVN and rostral ventrolateral medulla (RVLM),

immunofluorescence of c-Fos, the immediate early gene of neuronal activation, was

performed to determine modeling-associated neuronal hyper-activity. Co-treatment of

EM significantly blunted DOCA-salt- or HSD-induced increase in the number of c-

Fos-positive cells (Figure 3), indicating blunted neuronal activation. To this end, we

demonstrated that co-treatment of EM reduces neuronal activity in the pre-autonomic

centers and blunts modeling-induced sympatho-excitation.

3.3 DOCA-salt and EM co-treatment are associated with vascular immune response in

the PVN

To further dissect the potential mechanism by which EM reduces sympathetic

activation, PVN tissues from 5 mice/group, in DOCA, DOCA+EM and sham group,

were collected for the process of single nucleus RNA sequencing (snRNA-seq)

(Figure4 A). A total of 37, 828 cells were obtained in the three groups (Figure4 B).

We analyzed labels of the areas adjacent to the PVN, including the anterior

hypothalamic area (Pmch+), dorsomedial hypothalamic nucleus (Cck+), medial

preoptic area (Nts+), medial preoptic nucleus (Tac2+), suprachiasmatic nucleus

(Lhx1+), and ventromedial hypothalamus (Nr5a1+). The majority of the neurons

expressed markers for PVN-locating neurons (Sim+), confirming that we captured the

PVN area, and the presence of cells in the lateral, caudal, and anterior of PVN

confirmed that our architecture of the extended PVN captured the entire area

(Supplementary Figure 3).

Then, we merged cells from the 3 groups and extracted the top 3, 000 highly

variable genes for principal component analysis (PCA). The primary components of

PCA guided the unsupervised clustering process, resulting in 35 different cell clusters.

Linear dimensionality reduction was followed by graph-based unsupervised clustering

to achieve this outcome. Uniform manifold approximation and projection (UMAP)

was utilized for visualization purposes. We renamed the cell clusters with reference to

cellMarker 2.0 and classic marker genes for different cell types. Nine different types

of cells were identified from the 35 cell cluster: mature oligodendrocyte (5,854),

Excitatory neuron (11,225), Inhibitory neuron (11,390), Mix neuron (4,620),

Astrocyte (1,749), Microglial cell (1,444), Immature oligodendrocyte (1,264),

Endothelial cell & Pericyte (176), Ependymal cell (106) (Figure4 B, C).

To unveil interactions between these cells alter the excitability of neurons, cell-

chat analysis was firstly performed (Figure4 D). We found that DOCA-salt treatment

induced direct signaling regulation of microglia by the vascular cells (endothelial cell

& pericyte), and this was not observed in the sham control group (upper panel).

Regulation of inhibitory neurons by microglia was detected in both sham- and only

DOCA-salt-treated groups, but it was abolished by co-treatment of EM in DOCA-

salt-treated group (lower panel). This cell-chat analysis highlighted an important

involvement of vascular cells, which were further analyzed for DOCA-salt- or EM-

associated differentially expressed genes (DEGs) and the functional enrichment of

these DEGs (Figure4 E). Gene ontology (GO) enrichment analysis was performed

using DOCA-salt- and EM-associated DEGs, respectively. Both sets of DEGs were

found to be significantly related to cytokine signaling and adhesion of immune cells,

indicating DOCA-salt might induce immune response to the PVN vessels, therefore

increasing recruitment of circulating leukocytes, and this immune response could be

abolished by co-treatment of EM (Figure4 F). The PVN has high capillary density,

thin vessel diameter, and complex vascular topology[11, 17-21], providing a possible

interface for immune cells and cytokines to affect PVN cells. It has been shown that

cytokines from leukocytes, such as interferon, has direct impact on microglia[31], the

resident immune cell within the brain. The populations of microglia were then

extracted and analyzed to confirm the possible impact from peripheral immune cells.

2.4 EM co-treatment abolished reactive transformation of microglia in the PVN

We re-populated and divided the microglia from all groups into four sub-clusters.

Following analyzing the top genes of each sub-clusters, they were named according to

their major biological process, or reactive state. The 4 sub-clusters were homologous

homeostasis microglia (HOM), disease-associated microglia (DAM), interferon

activated microglia (IFN), and microglia associated with myelination (ORM) (Figure5

A, B, C). Next, DOCA-salt- or EM-associated DEGs were analyzed for the HOM,

DAM, and IFN clusters (Figure5 D, F, H).

Firstly, the HOM-DEGs that associates with DOCA-salt were mainly enriched in

microglial activation, such as G protein-coupled purinergic receptor activity and

calcium signaling, indicating upregulated microglial ATP/P_2Y_{12} receptor/Ca^{2+}

signaling in the PVN of DOCA-salt treated mice. Interestingly, the HOM-DEGs that

associates with EM co-treatment were mostly enriched in the pathway of cell

adhesion binding (Figure5 E), suggesting that EM co-treatment might not affect

microglia via reducing ATP in the PVN, but directly blunt microglia's interaction

with neurons. Secondly, the DAM-DEGs that associates with DOCA-salt were mainly

enriched in the pathways of epithelial invasion and phagocytosis, indicating further

induced reactive microglia by DOCA-salt, while the DAM-DEGs that associates with

EM co-treatment were functionally enriched in platelet activation and oxidative

phosphorylation (Figure5 G), indicating altered energy metabolism for microglia.

DOCA-salt-induced increase of reactive microglia and EM-related decrease of

reactive microglia were validated by reduced co-expression of Iba1 and CD68 in the

PVN (Figure6 A, B). Thirdly, the IFN-DEGs that associates with DOCA-salt were

mostly enriched in the pathway of efferocytosis, which is significantly influenced by

cytokines. The IFN-DEGs that associates with EM co-treatment is highly enriched in

phagocytosis and related pathways (Figure5 I), further confirming that microglia were

less reactive in the PVN of mice treated with EM.

Interferon-γ (IFN-γ) is a pro-inflammatory cytokine produced by NK cells, NK T

cells, and T cells, and it has been shown to induce disease-associated state

transformation of microglia[9, 12, 17, 32-40]. Accordingly, we identified several IFN-

DEGs that showed opposite pattern in DOCA-salt and EM co-treated group (Figure5

J), suggesting those IFN-DEGs could be directly related to the protective effect of EM.

Notably, *Txndc9*, *Ppia* are related to protection against neuroinflammation damage,

while *Adam17* is known for its role in signaling transduction of neuroinflammation

and efferocytosis.

Reactive state of microglia was validated by co-staining of Iba1 and CD68.

Increased number of reactive microglia was also found in HSD models, and co-

treatment of EM significantly blunted HSD-induced increase of microglial reactive

transformation (Figure6 C). Additionally, activation of peripheral immune system was

determined by lymphocyte counts and plasma level of IFN-γ. DOCA-salt or HSD

treatment significantly increased the cell counts of lymphocytes, while co-treatment

of EM blunted modeling-induced augment of circulating lymphocytes (Figure6 D, E).

Similarly, DOCA-salt or HSD treatment markedly elevated the concentration of IFN-

γ in peripheral blood, while this was normalized by co-treatment of EM (Figure6 F,

G). In addition to lymphocytes, co-treatment of EM also had impact on neutrophil

count caused by high sodium intake (Supplementary Figure3 G, H).

3.5 EM co-treatment was associated with overall blunted neuronal activity in the PVN

Next, snRNA-seq data of neuronal cells were analyzed. For excitatory neurons,

DOCA-salt increased the function of PVN neuro-endocrine neurons, meanwhile, it

upregulated the ubiquitination pathway of protein degradation (Figure7 A). Co-

treatment of EM showed effect on both mechanisms. As to inhibitory neurons, the

major effect of DOCA-salt treatment was upregulation of the ubiquitination pathway,

while EM co-treatment exhibited significant impact on DOCA-salt-induced protein

ubiquitination (Figure7 B), suggesting that EM could help to "calm down" the PVN

neuro-endocrine by blunting the pathway of protein ubiquitination. It has been

demonstrated that ACE2 ubiquitination impairs its enzymatic activity, thus

compromises its ability to support GABAergic inhibitory tone in the PVN[41, 42].

Accordingly, enzymatic activity of ACE2 was measured using hypothalami. As

previously reported, DOCA-salt treatment markedly reduced ACE2 activity. On the

other hand, EM co-treatment abolished DOCA-salt's effect on ACE2 enzymatic

activity (Figure7 C). Meanwhile, ACE2 activity was also impaired by HSD, and this

was prevented by co-treatment of EM (Figure7 D). Furthermore, expression level of

NEDD4L, which is responsible for ACE2 ubiquitination during hypertension, was

measured. In addition to DOCA-salt hypertension (Figure7 E, F), HSD induced

NEDD4L expression as well, and EM co-treatment significantly blunted DOCA-salt

or HSD induced upregulation of NEDD4L expression (Figure7 G, H).

ACE2 has showed robust expression in various types of PVN neurons, except for

glutamatergic neurons[43, 44]. Then we analyzed PVN neuronal markers to identify

which type of neurons response to EM co-treatment. Of note, expression levels of *Oxt*,

Avp, *Tac1*, *Trh*, and *Sst* were upregulated by DOCA-salt treatment while

downregulated by EM co-treatment (Figure7 I), indicating that neuro-endocrine and

neurosecretory neurons were involved in the protective effect of EM during DOCA-

salt hypertension. To confirm this, plasma co-peptin level was measure, and it showed

that DOCA-salt or HSD-related elevation of plasma co-peptin level, was markedly

blunted by co-treatment of EM (Figure7 J, K). To this end, we showed that EM co-

treatment abolished disease-associated reduction of ACE2 enzymatic activity via

affecting ubiquitin-signaling, and this could be result, also a mediator, for less

excitation of neuro-endocrine/neurosecretory neurons.

3.6 EM conditionally reduced blood lymphocyte count in patients with heart failure

In mice, we showed that EM conducted protection against cardiac remodeling in

DOCA-salt or HSD models possibly via reducing peripheral cytokine-mediated

activation of microglia, and reducing excitation of neuro-endocrine/neurosecretory

neurons. Next, we designed a cross-sectional observational clinical study, to further

validate the above findings in human. We recruited 1864 patients with heart failure

were admitted to our hospital and agreed to join this study from January 2018 to

January 2023, excluding 675 patients with incomplete hospitalization information or

failure to meet enrollment requirements. Our final study population was 1189 patients

with heart failure, divided into a non-treated group (n = 715) and a treated group (n =

474). After propensity score matching (PSM), the study population was divided into a

matched non-treated group (n = 437) and a matched treated group (n = 474). The flow

chart of this study is shown in (Figure8).

Table1 summarizes the clinical baseline between the two groups before and after

PSM. Although there were significant differences in some patient characteristics

between the non-treated group and the treated group before matching, all patient

characteristics except alcohol consumption, the level of NT-proBNP, and diuretic

using situation were similar between the matched non-treated group and the matched

treated group.

Table2 shows the results of comparing HCT, neutrophil count, lymphocyte count,

and monocyte count between groups after PSM. Compared to the matched non-

treated group, the matched treated group had higher HCT ($P < 0.001$) and lower

lymphocyte and monocyte counts (p = 0.02, p < 0.001). In addition, Supplementary

Table1 summarizes the results of HCT, neutrophil count, lymphocyte count, and

monocyte count between the two groups before PSM. The SGLT2i treated group had

lower HCT ($P < 0.001$) and higher lymphocyte count ($P < 0.001$). There were no

inter-group differences between monocyte counts and neutrophil counts ($P = 0.99$, P

$= 0.08$).

Due to alcohol consumption between the two groups after PSM, there are still

differences in the level of NT-proBNP and diuretic using situations. Therefore, we

used hierarchical analysis to explore the differences in HCT, neutrophil count,

lymphocyte count, and monocyte count between the matched treated group and the

matched non-treated group in populations with different alcohol consumption,

diuretic using situation, and NT-proBNP levels. Categorical variables are directly

stratified according to classification. Continuous variables (NT-proBNP) were divided

into four groups according to quartile positions. The results showed that the HCT in

the matched EM-treated group was higher than that in the matched non-treated group

regardless of the stratified situation. In the group who never drank alcohol (Alcohol1),

lymphocyte counts in the matched treated group were lower than in the matched non-

treated group. In the group with higher NT-proBNP (>2550), the lymphocyte count in

the matched treated group was lower than that in the matched non-treated group. In

the patients who do not use diuretic treated, the lymphocyte counts of the matched

treated group were lower than in the matched non-treated group ($P < 0.001$). In

patients treated with diuretics, neutrophil counts were lower in the matched treated

group ($P = 0.004$) (Figure8). SGLT2i may play a therapeutic role by regulating HCT

and lymphocyte count.

3. Discussion:

Emerging evidence highlights cardiovascular protective effects of SGLT2i,

potentially linked to sympathetic inhibition [45], yet their impact on autonomic brain

centers remains unexplored. Here, in two salt-related models (DOCA-salt and 8%-

HSD models), we showed that EM protected against cardiac remodeling partly by

suppressing central sympathetic outflow. For underlying mechanisms, we found that

EM could conduct its protective role via reducing peripheral cytokine-mediated

activation of microglia, and preventing over-excitation of neuro-

endocrine/neurosecretory neurons (Figure9).

Through clinical studies, we have found the effect of EM on lymphocyte count

and HCT. This links to what we found in the experimental models. NT-proBNP is an

important diagnostic biomarker for of acute heart failure with higher mortality risk.

Inflammation appears to be associated with natriuretic peptide (NP) release[46]. It has

been reported that BNP level is closely related to extent of LV sympathetic

overactivity in heart failure[47, 48]. According to the different affinity of NE to α-

and β-adrenergic receptor, increase of sympathetic outflow could induce higher

vascular resistance than its effects on heart. Also, elevated cardiac sympathetic tone

could further increase cardiac metabolism, reduce cardiac diastole and coronary blood

flow, therefore exacerbating the pathological progression. In patients with high NT-

proBNP value, EM treatment significantly reduced their number of circulating

lymphocytes. EM also significantly reduced lymphocyte count in patients without

diuretic treatment, while it markedly lowered neutrophil count in patients received

diuretic treatment. In mice, EM also reduced neutrophil count in both DOCA-salt and

HSD models (Supplementary Figure3 G, H), similar as their lymphocyte counts.

Notably, reduced level of circulating neutrophil might also contribute to less

advanced condition of pathological cardiac remodeling. However, for interaction with

the brain cells, it should be lymphocytes (especially the T cells) and associated

cytokines, because brain recruitment of neutrophil is very limited in non-infection

situation and we did not observe marked change in monocyte level following DOCA-

salt or HSD treatment. For T cells, we have shown that DOCA-salt dramatically

increase its ratio in peripheral blood mononuclear cells, and ablation of sympatho-

excitation, using transgenic mutation that targets the brain, significantly reduced the

ratio of both CD3+ and CD8+ T cells[44]. In current study, in both models, we found

that EM co-treatment reduced lymphocyte count and the plasma level of IFN-γ, which

is produced by lymphocytes. EM has been shown to reduce inflammation in diabetic

patients [49-52], but its relationship with lymphocytes and cytotoxic cytokines (*i.e.*

IFN-γ) is unknown. IFN-γ is a cytokine that closely correlates to the occurrence and

progression of hypertension [16, 53, 54], and it also acts on microglial located in the

CNS [55-57], thus promoting neuroinflammation. Reactive microglia in the PVN has

been demonstrated to promote excitation of PVN neurons and increase of sympathetic

outflow[11, 17-21].

EM co-treatment also increase HCT in the DOCA-salt treated mice

(Supplementary Figure3 E, F), similar as in the heart failure patients. Decrease of

HCT following DOCA-salt treatment might indicate increase of circulating blood

volume, which is similar as the results of some previous studies [58-60]. On the other

hand, SGLT2i reduces blood volume. This effect may be caused by the drug's diuretic

effect. This has also been confirmed in some clinical studies [61-63]. This suggests

that EM might blunt cardiac remodeling by promoting sodium ion excretion and

reducing circulating blood volume. But EM also significantly increased HCT, and

lowered neutrophil count, in the patients received diuretics, suggesting additional

effect of EM than its diuretic effect. To determine the effect of sodium excretion, we

also measured the levels of plasma sodium and osmolality in both DOCA-salt and

HSD models. Although the modeling mice had significantly more sodium intake than

baseline, their system showed a strong regulatory impact on blood sodium and

osmotic pressure (Supplementary Figure3 A-D). The body processes a series of

hormones that regulate water-sodium homeostasis, maintaining stable levels of blood

sodium and osmotic pressure. Among these hormones, vasopressin is secreted from

PVN neurosecretory AVP neurons, and plays an essential role in development of

DOCA-salt hypertension[24]. Unlike DOCA-salt model, HCT was not altered in mice

with HSD treatment, nevertheless, the plasma of level co-peptin, which represents

AVP level, was elevated by HSD and brought to the baseline in mice co-treatment of

EM, further supporting the involvement of CNS. Accordingly, we think that via

facilitating sodium excretion, EM reduced excitation of AVP (Arginine vasopressin)

neurons, and less excited AVP neurons secreted lower level of vasopressin, thus

further supporting the protective effect of EM.

This study attempted to verify the mechanism by which EM regulates the foot

sensation center through clinical samples, but there were several inherent limitations

because this study was a single-center retrospective design. First, selection bias may

exist because only patients with complete medical records are included and cases that

are more serious or lost to follow-up may be excluded, which may underestimate

adverse outcomes. Second, unmeasured confounding factors may affect the results,

although we adjusted for available variables. Despite these limitations, the

consistency of our findings with previous literature supports its validity.

This study used single-center retrospective study to explore how EM regulates

the sympathetic nervous system, acknowledging limitations like selection bias and

unmeasured confounders, though consistent findings with prior research support its

validity.

4. Conclusion:

In this study, we found that peripheral immune activation contributes to reactive

transformation of microglia in the PVN, meanwhile, upregulated ubiquitin-signaling

impairs the compensatory activity of ACE2, and this further promoting the hyper-

activity of PVN neurons. By targeting peripheral immune activation and over-

excitation of neuro-endocrine/neurosecretory neurons, EM reduced sympathetic

outflow and plasma vasopressin level, thus introducing protective effect on cardiac

pathological remodeling.

5. Methods:

5.1 Antibodies

antibodies	source	indentifier
Tyrosine hydroxylase	Abcam	ab31252
c-Fos	Abcam	c-Fos
IBA1	Abcam	ab208942
CD68	Thermo	MA5-16654
SGLT2	Proteintech	28683-1-AP
NEDD4L	Proteintech	13690-1-AP
GAPDH	CST	2118
Anti rabbit 488	Thermo	A-11008
Anti rat 555	Thermo	A-21434
Anti rabbit HRP	Proteintech	SA00001-2
Anti mouse HRP	Proteintech	SA00001-1

443 Stored data

SRA dataset	PRJNA1265702

444 Software and algorithms

Package	Version
R	4.2.3
Seurat	4.3.0.1

CellChat	1.6.1

5.2 Experimental Model and Subject Details

This study used male C57BL/6J mice of SPF level at 8 weeks old. They were

purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd. (Order

No. 44829700009980). The animals were raised in the Experimental Animal Center

of Xi'an Jiaotong University under SPF barrier conditions. The animal feed was

purchased from Jiangsu Xietong Pharmaceutical Bio-engineering Co., Ltd. (regular

feed: SWS9102, 8% NaCl feed: XTNacl-8C). Mice had free access to water and food.

All procedures were approved by the IACUC of Xi'an Jiaotong University

(XJTU2021-203), in agreement with the Guide for the Care and Use of Laboratory

Animals. Every effort was made to minimize the number of animals used and their

suffering. Institutional Animal Care and Use Committees in accordance with the

'Principles of Laboratory Animal Care by the National Society for Medical research

and the Guide for the Care and Use of Laboratory Animals' (National Institutes of

Health Publication No. 86-23, revised 1996).

5.3 Hypertension Model and High Salt Diet Model

All mice were adaptively fed for 1 week. The DOCA group, DOCA+EM group,

DOCA+Hyd group, Sham+EM group and Sham group all underwent right kidney

removal. The DOCA group, DOCA+EM group and DOCA+Hyd group underwent

Deoxycorticosterone acetate (DOCA) silicone implantation. Anesthesia was induced

with isoflurane (3% induction; 1-2% maintenance) using an oxygen flow of 0.5 L/min,

and the mice were placed on a heating pad to maintain body temperature at around

37.5□. The mouse was placed in a left lateral position on the operating table, and the

skin in the posterior peritoneal area was incised to expose the renal hilum of the right

kidney. The blood vessels were ligated with surgical suture at the renal hilum,

followed by removal of the right kidney. After surgery, subcutaneous injection of 0.9%

saline solution (1mL) was given via backpack skin. One week after recovery, either

DOCA, DOCA+EM or DOCA+Hyd groups received subcutaneous implantation of

DOCA silicone sheet (containing 1 mg of DOCA per gram body weight). Mice

implanted with DOCA silicone sheets had their drinking water replaced with 1%

NaCl solution.

HSD, HSD + EM and normal sodium diet (NSD) groups were fed adaptively for

one week. HSD, HSD + EM were fed with a diet containing 8% sodium for 12 weeks.

The NSD group was fed with a normal diet (sodium content was 0.45%) for 12 weeks.

All mice were given free access to food and water.

All mice treated with EM were administered by daily gavage at a dose of

10mg/kg/day. The dosages of the EM were selected based on the literature. All tissue

preparations were performed after euthanasia. Using a precision vaporizer with

induction chamber and waste gas scavenger, (indicate the gas anesthetic) will be

administered slowly up to [indicate: > 4.5 % (for Isoflurane)] in oxygen and

continued until respiratory arrest occurs for > 60 seconds. The chamber is flushed

with oxygen only, the animal is removed and rapid removal of (indicate tissues /

organs) is performed to assure euthanasia.

5.4 Single-cell RNA-sequencing

5 mice were sacrificed with a lethal dose of isoflurane, and the excess blood was

washed off in cold PBS, and the PVN region of each mouse was obtained according

to the mouse brain anatomy map. The tissue nucleus was extracted by SHBIO tissue

sample nucleus isolation kit (Cat# 52009-10).

Single-cell RNA-Seq libraries were prepared using SeekOne® Digital Droplet

(SeekOne® DD) Single Cell 3' library preparation kit (SeekGene Catalog No.

K00202-0201). The indexed sequencing libraries were cleanup with SPRI beads,

quantified by quantitative PCR (KAPA Biosystems KK4824) and then sequenced on

Illumina NovaSeq 6000 with PE150 read length or DNBSEQ-T7 platform with

PE150 read length.

We used the SeekSoul Tools pipeline to process the cleaned reads and generated the

transcript expression matrix. Quality control, clustering, and downstream analysis

were performed using Seurat v. 4.3.0.1 in R version 4.2.3 (2023-03-15). Briefly,

according to gene, UMI, and mitochondrial gene counts (genes > 100 and < 8000,

mitochondria < 5%).

For the clusters, data was normalized and scaled using the NormalizeData

function in Seurat with a factor of 10,000. The top 3000 variable genes were selected

using the FindVariableFeature function in Seurat, and the data was scaled using the

ScaleData function. Cells were clustered using the FindCluster function with 20

principal components and a resolution of 1.0. The cells were then plotted in a two-

dimensional space using the 'Unified Manifold Approximation and Projections'

(UMAP) technique, and the identities of the cell clusters were determined by their

overlap with known marker genes in the literature. 35 clusters were identified as 9 cell

types: Mature oligodendrocyte (5,854), Excitatory neuro (11,225), Inhibitory neuro

(11,390）, Mix neuro (4,620), Astrocyte (749), Microglial cell (1444), Immature

oligodendrocyte (1,264), Endothelial cell & Pericyte (176), Ependymal cell (106). In

order to better describe the neuron population, excitatory neurons, inhibitory neurons,

and mixed neurons were further separated and re-clustered. Cells were clustered using

the FindCluster function with 20 principal components and a resolution of 0.1.

We performed cell-cell communication analysis using CellChat v1.4.0, which is based

on the curated ligand-receptor interaction database (CellChatDB). To perform the

analysis, we provided normalized gene expression matrices for PVN tissues as input

to CellChat. The computeCommunProb function was used to calculate the total

numbers of interactions and interaction strengths, while the

computeCommunProbPathway function was used to calculate the communication

probabilities for each cell signaling pathway.

5.5 Echocardiography

Transthoracic echocardiography was performed 21 days after DOCA silicone

block implantation. Echocardiography was performed under anesthesia with

isoflurane (0.25 to 0.50%) supplemented with 100% O2 using a 30MHz probe on

Vevo®LAZR-X 3100. Left ventricular mass (LVM) and left ventricular ejection

fraction (LVEF) were analyzed from M-mode images.

5.6 Staining of the Heart

After the mice were directly sacrificed, the heart was removed, and the blood

washed off the surface in cold PBS. The heart was placed in 4% PFA for 48 hours and

then paraffin-embedded sections (10 µm) were performed. The H&E staining kit

(Servicebio Cat#: G1076-500ML) and MASSON staining kit (Servicebio Cat#:

G1006-20M) were used.

5.7 Immunocytochemistry

After antigen repair with sodium citrate, the paraffin sections of the heart were

washed twice with PBS, then 10min was incubated with 0.25%TritonX100 at room

temperature, and any non-specific binding was blocked by 10%NGS at room

temperature for 1 hour and incubated overnight with TH primary antibody (1:1000,

4 □). After washing with PBS for three times, the rabbits-488 were incubated at room

temperature for 1 hour and then washed again with PBS. Finally, anti-staining was

performed with DAPI (1-VOL1000-THERMOTIMO-Fisher Science).

For brain tissue, mice were fixed by cardiac perfusion with cold 4%PFA in PBS.

After the brain tissue was fixed by 4%PFA and embedded with OCT (SAKURA-

4583), the brain map skin of the control mice was sliced (25 μm) and specific parts

were taken for immunofluorescence staining. Wash with PBST twice, then incubate

10 min with 0.5%Triton-X100 at room temperature, block any non-specific binding

with 10% NGS at room temperature for 1 hour, dilute the primary antibody in

proportion and incubate overnight with 4□. After washing with PBS for three times,

the corresponding secondary antibody was used to incubate at room temperature for 1

hour, and then cleaned again with PBS. The cell nucleus was stained with DAPI

(Thermo Fisher Science Cat#: 62247). The main antibodies used in this study are

summarized in Supplementary Table 1. Finally, the fluorescence microscope

(Olympus BX43) was used to capture the image. Using Image J software for image

analysis

5.8 Norepinephrine Measurement

Urine samples were collected by bladder massage at the end of the experimental

protocol and noradrenaline concentrations were determined using a mouse

noradrenaline ELISA kit (Elabscience, Cat#: E-EL-0047) according to the

manufacturer's instructions. Noradrenaline levels are normalized by the corresponding

569 creatinine concentration (Elabscience, Cat#: E-BC-K188-M). Normalized

570 noradrenaline concentrations in urine are expressed as ng/ng creatinine.

572 5.9 Peripheral Blood Cell Counts and Plasma Sodium Detection

573  After the experimental protocol was completed, the mice were anesthetized with

574 isoflurane and blood was taken through the eyeball. Half of blood was anticoagulation

575 by EDTA anticoagulation. Then the blood was routinely examined by a fully

576 automated three-point population blood cell analyzer (Mindray BC-2800). The

577 remaining blood was anticoagulated by EDTA and centrifuged for 4 □. Follow the

578 manufacturer's instructions to use blood sodium detection kit (Elabscience, Cat#: E-

579 BC-K207-M), MCP-1 test kit (Elabscience, Cat#: E-EL-M3001), co-peptin test kit

580 (JINGMEI BIOTECHNOLOGY, Cat#: JM-12422M1), and IFN-γ test kit

581 (Elabscience, Cat#: E-EL-M0048) to determine the corresponding indicators. The

582 osmotic pressure was detected by an osmometer (Fiske Micro-Osmometer Model

583 210).

585 5.10 ACE2 Enzyme Activity Detection

586  Mice were sacrificed with a lethal dose of isoflurane, and the paraventricular

587 nucleus region of the inferior thalamus was obtained for each mouse according to the

588 mouse brain anatomy map. ACE2 enzyme activity was determined using the ACE2

589 enzyme activity detection kit (Beyotime, Cat#: P0319S) according to the

590 manufacturer's instructions.

592 5.11 Western Blotting

593  The tissue samples were placed in 8M urea solution (4.8g urea, 10% SDS 5mL

594 0.5mL 1M Tris PH 7.4 0.5mL, 0.5M EDTA 0.1mL plus H_2O to 10mL) supplemented

595 with a mixture of protease inhibitors (Roche, Cat#: 11697498001) and repeatedly

596 pumped with a 1ml needle. After quantifying the protein concentration by BCA

597 protein assay (Thermo Scientific, Cat#: A55860), the protein sample was mixed with

598 $4 \times$ SDS sample buffers (Thermo Scientific, Cat#: NP0007) and boiled at 65 °C for 10

599 minutes. 30 µg protein samples were separated on 4-12% MOPS gel (ACE, Cat#:

600 F15412LGel) and transferred to PVDF membrane (Millipore). Seal the imprint in 5%

601 skim milk for 1 hour. On the second day, the protein was detected with specific

602 antibody (diluted by 1purl 1000) and secondary antibody conjugated with HRP

603 (diluted by 1purl 2000). Aim at the rabbit secondary antibody conjugated with Ubr1

604 (sc-98882) Nedd4-2 and HRP conjugated with GAPDH.

605 The ECL luminescent solution (MISHUSHENGWU, Cat#: MI00607B) was dripped

606 evenly on the PVDF membrane (Millipore, Cat#: IPVH 00010) and the imaging was

607 carried out by the WB experimental chemiluminescence instrument (Bio-rad,

608 Cat#:0087170). The signal strength was quantified by ImageJ.

5.12 Clinical Study Design

This study is a retrospective single-center clinical study. All participants provided

written informed consent before participating in the study. The study was approved by

the Institutional Review Board at the Second Affiliated Hospital of Xi'an Jiaotong

University (IRB#2024-150). All methods were performed in accordance with the

relevant guidelines and regulations.

5.12.1 Study Participants

We reviewed patients diagnosed with heart failure and hospitalized in the Second

Affiliated Hospital of Xi'an Jiaotong University from January 2018 to January 2023.

The inclusion criteria are: 1) Signed informed consent and was older than 18 years old

at the time of signing informed consent; 2) Heart failure was diagnosed within 3

months before enrollment. The exclusion criteria are: 1) Cerebrovascular accident; 2)

Fever caused by various reasons, severe infections; 3) Moderate or severe anemia; 4)

Uncontrolled hypertension, systolic blood pressure greater than 160mmHg; 5) Renal

failure patients; 6) Hyperthyroid endocrine diseases; 7) Take SGLT2i but for less than

1 month.

A total of 1189 patients were included in the study, of which 474 were treated with

SGLT2i and 715 were not treated with SGLT2i. We performed PSM (propensity

score matching) to match background characteristics between treated and non-treated

group. Details of the PSM analysis are described in the Statistical Analysis section.

Clinical characteristics and outcomes of the 2 groups were compared before and after

propensity score matching. The primary interests of this study were the HCT,

neutrophil count, monocyte count, and lymphocyte count.

Blood pressure, heart rate, weight, and height were all collected on the day of

admission. Blood glucose, LDL, serum creatinine, serum uric acid, serum sodium,

serum potassium, and NT-proBNP were all measured on the day of admission or after

blood was collected on an empty stomach on the morning of the second day. LVEF is

the result of cardiac ultrasound during hospitalization. QRS time limit is the result of

electrocardiogram measurement on the day of admission.

5.12.2 Statistical Analysis

Data are expressed as a percentage of categorical variables, and continuous

variables were subjected to the Wilke-Shapiro test to determine whether the

continuous variables were normally distributed. The mean ± standard deviation (SD)

of continuously distributed variables is expressed, and non-normally distributed

variables are expressed as the median (quartile 1-quartile 3). Normal distributed

continuous variables were compared between the 2 groups using a student t-test.

Otherwise, continuous variables were compared using the Mann-Whitney U test.

Classification variables were compared using Fisher's exact test.

Clinical factors that may be relevant with outcome variables such as sex, age,

smoking status, alcohol consumption, diabetes, dyslipidemia, chronic kidney disease,

hypertension, angina pectoris, myocardial infarction, atrial fibrillation, renin

angiotensin system inhibitors were selected for PSM.ARNI/ACEI/ARB use, diuretics

use, beta-receptor blockers use, aldosterone receptor blockers use, QRS duration, NT-

proBNP, serum creatinine value, total cholesterol, LDL, hemoglobin, blood glucose ,

uric acid on admission were used as independent variables. For matching, the

tendency score value of each study subject was calculated, and the caliper value was

0.1. After PSM (1:1), the treated group was matched with the control group, and the

balance between the groups was good.

After PSM, HCT, monocyte count, neutrophil count and lymphocyte count in the

treated group and control group were tested by Mann-Whitney U test. If there are still

differences between the two groups after PSM matching, hierarchical analysis is

performed based on this variable and then HCT, monocyte count, neutrophil count

and lymphocyte count between groups will be compared.

A P value < 0.05 was considered statistically significant. We analyzed all data by

IBM SPSS statistics version 22 (Chicago, IL, USA).

6. Declarations

6.1 Ethics Approval and Consent to Participate

All procedures of anmial research were approved by the IACUC of Xi'an

Jiaotong University (XJTU2021-203), in agreement with the Guide for the Care and

Use of Laboratory Animals. Every effort was made to minimize the number of

animals used and their suffering. Institutional Animal Care and Use Committees in

accordance with the 'Principles of Laboratory Animal Care by the National Society

for Medical research and the Guide for the Care and Use of Laboratory Animals'

(National Institutes of Health Publication No. 86-23, revised 1996).

The clinical study of this study is a retrospective single-center clinical study. All

participants provided written informed consent before participating in the study. The

study was approved by the Institutional Review Board at the Second Affiliated

Hospital of Xi'an Jiaotong University (IRB#2024-150). All methods were performed

in accordance with the relevant guidelines and regulations.

6.2 Consent for publication

We confirm that the manuscript has been read and approved by all authors and

that there are no other persons who satisfied the criteria for authorship but are not

listed. We further confirm that the order of authors listed in the manuscript has been

approved by all of us.

6.3 Data Availability

All requests for raw data, analysed data and materials of this study will be

promptly reviewed by the corresponding authors (Dengfeng gao:

gaomedic@mail.xjtu.edu.cn) and the Second Affiliated Hospital of Xi'an Jiaotong

University.

The raw data of snRNA-seq are available in SRA with accession number

PRJNA1265702.

6.4 Competing Interests

692   All authors declare no competing interests.

693  6.5 Funding

694   This work was supported by the National Natural Science Foundation of China

695  (81872563 to Dengfeng Gao and 82100454 to Jiaxi Xu).

696  6.6 Acknowledgements

697   We would like to thank Shenglan Yuan for her guidance and assistance in the

698  animal experiment section. Thanks to the medical staff of the Department of

699  Cardiovascular Medicine of the Second Affiliated Hospital of Xi`an Jiaotong

700  University for their help in the clinical research part. Thanks to all the patients who

701  participated in the clinical research.

703  **Reference**：

704  [1] Gibb AA, Hill BG. Metabolic coordination of physiological and pathological cardiac

705  remodeling. Circulation Research, 2018, 123: 107-128

706  [2] Patel SM, Kang YM, Im K, *et al.* Sodium-glucose cotransporter-2 inhibitors and major

707  adverse cardiovascular outcomes: A smart-c collaborative meta-analysis. Circulation, 2024,

708  149: 1789-1801

709  [3] Udell JA, Yuan Z, Rush T, *et al.* Cardiovascular outcomes and risks after initiation of a

710  sodium glucose cotransporter 2 inhibitor results from the easel population-based cohort study

711  (evidence for cardiovascular outcomes with sodium glucose cotransporter 2 inhibitors in the

712  real world). Circulation, 2018, 137: 1450-1459

[4]   Kumar K, Kheiri B, Simpson TF, *et al.* Sodium-glucose cotransporter-2 inhibitors in heart

failure: A meta-analysis of randomized clinical trials. American Journal of Medicine, 2020, 133:

E625-E630

[5]   Dyck JRB, Sossalla S, Hamdani N, *et al.* Cardiac mechanisms of the beneficial effects of

sglt2 inhibitors in heart failure: Evidence for potential off-target effects. Journal of Molecular

and Cellular Cardiology, 2022, 167: 17-31

[6]   Cowie MR, Fisher M. Sglt2 inhibitors: Mechanisms of cardiovascular benefit beyond

glycaemic control. Nature Reviews Cardiology, 2020, 17: 761-772

[7] Zhang DY, Anderson AS. The sympathetic nervous system and heart failure. Cardiol Clin,

2014, 32: 33-45, vii

[8]   Allen WE, Blosser TR, Sullivan ZA, *et al.* Molecular and spatial signatures of mouse

brain aging at single-cell resolution. Cell, 2023, 186: 194-208 e118

[9]   Cao K, Qiu L, Lu X, *et al.* Microglia modulate general anesthesia through p2y(12)

receptor. Current biology : CB, 2023, 33: 2187-2200 e2186

[10] Li P, Ma S, Ma X, *et al.* Reversal of neurovascular decoupling and cognitive impairment

in patients with end-stage renal disease during a hemodialysis session: Evidence from a

comprehensive fmri analysis. Human brain mapping, 2023, 44: 989-1001

[11] Wang YL, Zhu H, Pan YT, *et al.* Dendritic cell mineralocorticoid receptor controls blood

pressure by regulating t helper 17 differentiation: Role of the plcbeta1/4-stat5-nf-kappab

pathway. Eur Heart J, 2025, 46: 1335-1351

[12] Zhu Q, Xiao L, Cheng G, *et al*. Self-maintaining macrophages within the kidney

contribute to salt and water balance by modulating kidney sympathetic nerve activity. Kidney

Int, 2023, 104: 324-333

[13] Perrotta M, Lori A, Carnevale L, *et al*. Deoxycorticosterone acetate-salt hypertension

activates placental growth factor in the spleen to couple sympathetic drive and immune

system activation. Cardiovasc Res, 2018, 114: 456-467

[14] Lu X, Crowley SD. Actions of immune cells in the hypertensive kidney. Curr Opin

Nephrol Hypertens, 2020, 29: 515-522

[15] Ye SJ, Huang H, Han X, *et al*. Dectin-1 acts as a non-classical receptor of ang ii to

induce cardiac remodeling. Circulation Research, 2023, 132: 707-722

[16] Benson LN, Liu Y, Wang X, *et al*. The ifnγ-pdl1 pathway enhances cd8t-dct interaction to

promote hypertension. Circulation Research, 2022, 130: 1550-1564

[17] Dong Z, Hou L, Luo W, *et al*. Myocardial infarction drives trained immunity of monocytes,

accelerating atherosclerosis. Eur Heart J, 2024, 45: 669-684

[18] McMoneagle E, Zhou J, Zhang S, *et al*. Neuronal k(+)-cl(-) cotransporter kcc2 as a

promising drug target for epilepsy treatment. Acta pharmacologica Sinica, 2024, 45: 1-22

[19] Santisteban MM, Schaeffer S, Anfray A, *et al*. Meningeal interleukin-17-producing t cells

mediate cognitive impairment in a mouse model of salt-sensitive hypertension. Nat Neurosci,

2024, 27: 63-77

[20] Wei B, Cheng G, Bi Q, *et al.* Microglia in the hypothalamic paraventricular nucleus sense

hemodynamic disturbance and promote sympathetic excitation in hypertension. Immunity,

2024,

[21] Yu Y, Weiss RM, Wei SG. Interleukin 17a contributes to blood-brain barrier disruption of

hypothalamic paraventricular nucleus in rats with myocardial infarction. Journal of the

American Heart Association, 2024, 13: e032533

[22] Nielsen SF, Duus CL, Buus NH, *et al.* Randomized, placebo-controlled trial on the renal

and systemic hemodynamic effects of empagliflozin. Kidney international reports, 2025, 10:

134-144

[23] Umanath K, Testani JM, Lewis JB. "Dip" in egfr: Stay the course with sglt-2 inhibition.

Circulation, 2022, 146: 463-465

[24] Basting T, Lazartigues E. Doca-salt hypertension: An update. Curr Hypertens Rep, 2017,

19: 32

[25] Tyler M Basting JX, Srinivas Sriramula, Snigdha Murkerjee, and Eric Lazartigues. Knock-

down of adam17 in glutamatergic or pre-autonomic neurons attenuates development of

neurogenic hypertension. The FASEB Journal, 2017, 31:

[26] Grobe JL, Buehrer BA, Hilzendeger AM, *et al.* Angiotensinergic signaling in the brain

mediates metabolic effects of deoxycorticosterone (doca)-salt in c57 mice. Hypertension,

2011, 57: 600-607

[27] Grobe JL, Buehrer BA, Hilzendeger AM, *et al.* Angiotensinergic signaling in the brain

mediates metabolic effects of deoxycorticosterone (doca)-salt in c57 mice. Hypertension,

2011, 57: 600-607

[28] Albo D, Akay CL, Marshall CL, *et al.* Neurogenesis in colorectal cancer is a marker of

aggressive tumor behavior and poor outcomes. Cancer, 2011, 117: 4834-4845

[29] Hao X, Chen J, Luo Z, *et al.* Trpv1 activation prevents high-salt diet-induced nocturnal

hypertension in mice. Pflügers Archiv - European Journal of Physiology, 2011, 461: 345-353

[30] Li KX, Lu YM, Xu ZH, *et al.* Neuregulin 1 regulates excitability of fast-spiking neurons

through kv1.1 and acts in epilepsy. Nat Neurosci, 2011, 15: 267-273

[31] Bal-Price A, Brown GC. Inflammatory neurodegeneration mediated by nitric oxide from

activated glia-inhibiting neuronal respiration, causing glutamate release and excitotoxicity.

The Journal of neuroscience : the official journal of the Society for Neuroscience, 2001, 21:

6480-6491

[32] Chen X, Firulyova M, Manis M, *et al.* Microglia-mediated t cell infiltration drives

neurodegeneration in tauopathy. Nature, 2023, 615: 668-677

[33] Chen X, Gao R, Song Y, *et al.* Astrocytic at1r deficiency ameliorates abeta-induced

cognitive deficits and synaptotoxicity through beta-arrestin2 signaling. Progress in

neurobiology, 2023, 228: 102489

[34] Lyu Y, Huo J, Jiang W, *et al.* Empagliflozin ameliorates cardiac dysfunction in heart

failure mice via regulating mitochondrial dynamics. European Journal of Pharmacology, 2023,

942: 175531

[35] Pongratz G, Straub RH. [role of the sympathetic nervous system in chronic inflammation].

Z Rheumatol, 2023, 82: 451-461

[36] Robert SM, Reeves BC, Kiziltug E, *et al.* The choroid plexus links innate immunity to csf

dysregulation in hydrocephalus. Cell, 2023, 186: 764-785 e721

[37] Seo DO, O'Donnell D, Jain N, *et al.* Apoe isoform- and microbiota-dependent

progression of neurodegeneration in a mouse model of tauopathy. Science, 2023, 379:

eadd1236

[38] Weng Y, Chen N, Zhang R, *et al.* An integral blood-brain barrier in adulthood relies on

microglia-derived pdgfb. Brain, behavior, and immunity, 2024, 115: 705-717

[39] Xu Z, Hu SW, Zhou Y, *et al.* Corticotropin-releasing factor neurones in the

paraventricular nucleus of the hypothalamus modulate isoflurane anaesthesia and its

responses to acute stress in mice. British journal of anaesthesia, 2023,

[40] Xue L, Cui SM, Yang FH, *et al.* Myocardial senescence induced by inactivation aldh2-

sirt1-gata4 pathway involves in heart remolding after acute infarction. Circulation, 2016, 134:

[41] Mazher M, Blessing O, Mona E, *et al.* Nedd4-2 up-regulation is associated with ace2

ubiquitination in hypertension. Cardiovasc Res, 2023, 119:

[42] Mohammed M, Ogunlade B, Elgazzaz M, *et al.* Nedd4-2 upregulation is associated with

ace2 ubiquitination in hypertension. Cardiovasc Res, 2023, 119: 2130-2141

[43] Mukerjee S, Gao H, Xu J, *et al.* Ace2 and adam17 interaction regulates the activity of

presympathetic neurons. Hypertension, 2019, 74: 1181-1191

[44] Xu J, Molinas AJR, Mukerjee S, *et al.* Activation of adam17 (a disintegrin and

metalloprotease 17) on glutamatergic neurons selectively promotes sympathoexcitation.

Hypertension, 2019, 73: 1266-1274

[45] Spallone V, Valensi P. Sglt2 inhibitors and the autonomic nervous system in diabetes: A

promising challenge to better understand multiple target improvement. Diabetes &

Metabolism, 2021, 47: 101224

[46] Fish-Trotter H, Ferguson JF, Patel N, *et al.* Inflammation and circulating natriuretic

peptide levels. Circ Heart Fail, 2020, 13: e006570

[47] Sakata K, Iida K, Mochiduki N, *et al.* Brain natriuretic peptide (bnp) level is closely

related to the extent of left ventricular sympathetic overactivity in chronic ischemic heart

failure. Intern Med, 2009, 48: 393-400

[48] Tsukuda K, Mogi M, Iwanami J, *et al.* Cognitive deficit in amyloid-beta-injected mice was

improved by pretreatment with a low dose of telmisartan partly because of peroxisome

proliferator-activated receptor-gamma activation. Hypertension, 2009, 54: 782-787

[49] Maayah ZH, Ferdaoussi M, Takahara S, *et al.* Empagliflozin suppresses inflammation

and protects against acute septic renal injury. Inflammopharmacology, 2021, 29: 269-279

[50] Gohari S, Reshadmanesh T, Khodabandehloo H, *et al.* The effect of empagliflozin on

markers of inflammation in patients with concomitant type 2 diabetes mellitus and coronary

artery disease: The empa-card randomized controlled trial. Diabetol Metab Syndr, 2022, 14:

170

[51] Benedetti R, Chianese U, Papulino C, *et al.* Unlocking the power of empagliflozin:

Rescuing inflammation in hyperglycaemia- exposed human cardiomyocytes through

comprehensive multi-level analysis. Eur J Heart Fail, 2025,

[52] Fu J, Xu H, Wu F, *et al.* Empagliflozin inhibits macrophage inflammation through ampk

signaling pathway and plays an anti-atherosclerosis role. Int J Cardiol, 2022, 367: 56-62

[53] Benson LN, Liu Y, Deck K, *et al.* Ifn-γ contributes to the immune mechanisms of

hypertension. Kidney360, 2022, 3: 2164-2173

[54] Shokoples BG, Comeau K, Higaki A, *et al.* Angiotensin ii-induced a steeper blood

pressure elevation in il-23 receptor-deficient mice: Role of interferon-γ-producing t cells.

Hypertension Research, 2023, 46: 40-49

[55] Ding X, Yan Y, Li X, *et al.* Silencing ifn-γ binding/signaling in astrocytes versus microglia

leads to opposite effects on central nervous system autoimmunity. J Immunol, 2015, 194:

4251-4264

[56] He Z, Yang Y, Xing Z, *et al.* Intraperitoneal injection of ifn-γ restores microglial

autophagy, promotes amyloid-β clearance and improves cognition in app/ps1 mice. Cell

Death & Disease, 2020, 11: 440

[57] Zhang H, Zhang T, Wang D, *et al.* Ifn-γ regulates the transformation of microglia into

dendritic-like cells via the erk/c-myc signaling pathway during cerebral ischemia/reperfusion in

mice. Neurochemistry International, 2020, 141: 104860

[58] Griffin M, Rao VS, Ivey-Miranda J, *et al.* Empagliflozin in heart failure. Circulation, 2020,

142: 1028-1039

[59] Shikuma J, Sakakura K, Sugiyama-Takahashi M, *et al.* Hematocrit elevation after sglt2

inhibitor administration may be associated with the degree of proximal tubular damage.

Medicine (Baltimore), 2022, 101: e31122

[60] Tian Q, Guo K, Deng J, *et al.* Effects of sglt2 inhibitors on haematocrit and haemoglobin

levels and the associated cardiorenal benefits in t2dm patients: A meta-analysis. J Cell Mol

Med, 2022, 26: 540-547

[61] Wilcox CS. Antihypertensive and renal mechanisms of sglt2 (sodium-glucose linked

transporter 2) inhibitors. Hypertension, 2020, 75: 894-901

[62] Omar M, Jensen J, Burkhoff D, *et al.* Effect of empagliflozin on blood volume

redistribution in patients with chronic heart failure and reduced ejection fraction: An analysis

from the empire hf randomized clinical trial. Circulation: Heart Failure, 2022, 15: e009156

[63] Sen T, Scholtes R, Greasley PJ, *et al.* Effects of dapagliflozin on volume status and

systemic haemodynamics in patients with chronic kidney disease without diabetes: Results

from dapasalt and diamond. Diabetes Obes Metab, 2022, 24: 1578-1587

A

PVN

RVLM

DOCA   DOCA+EM   Sham

C

PVN

RVLM

HSD   HSD+EM   NSD

B

the number of c-Fos(+) neuron

****
***
**

DOCA   DOCA+EM   Sham

the number of c-Fos(+) neuron

***
**
ns

DOCA   DOCA+EM   Sham

D

the number of c-Fos(+) neuron

**
*
ns

HSD   HSD+EM   NSD

the number of c-Fos(+) neuron

***
**
ns

HSD   HSD+EM   NSD

A

GO enrichment analysis of DEGs in excitatory neuron

DOCA VS Sham   DOCA+EM VS DOCA

GO terms (top to bottom):
- ubiquitin protein ligase binding
- ubiquitin ligase inhibitor activity
- ubiquitin−protein transferase inhibitor activity
- ubiquitin−protein transferase regulator activity
- neuropeptide receptor binding
- neuropeptide hormone activity
- neuropeptide receptor binding
- ubiquitin ligase inhibitor activity
- ubiquitin−protein transferase inhibitor activity
- ubiquitin protein ligase binding
- GABA receptor binding

$-\log_{10}(P\ \text{value})$

B

GO enrichment analysis of DEGs in inhibitory neuron

DOCA VS Sham   DOCA+EM VS DOCA

GO terms (top to bottom):
- response to endoplasmic reticulum stress
- ubiquitin protein ligase binding
- ubiquitin ligase inhibitor activity
- ubiquitin−dependent ERAD pathway
- GABA−ergic synapse
- negative regulation of proteasomal ubiquitin−dependent protein catabolic process
- ligand−gated ion channel signaling pathway
- ionotropic glutamate receptor signaling pathway

$-\log_{10}(P\ \text{value})$

C Relative activity of ACE2 — DOCA, DOCA+EM, Sham (* , **)

D Relative activity of ACE2 — HSD, HSD+EM, NSD (* , **)

E NEDD4L / GAPDH — Sham, DOCA+EM, DOCA

F Average Pixel (Arbitary Unit)$\times 10^{-4}$ — DOCA, DOCA+EM, Sham (* , **)

G NEDD4L / GAPDH — NSD, HSD, HSD+EM

H Average Pixel (Arbitary Unit)$\times 10^{-4}$ — HSD, HSD+EM, NSD (* , *)

I Heatmap: Oxt, Avp, Agtrap, Tac1, Trh, Agtr1a, Sst; group: DOCA, DOCA+EM, Sham

J the level of co-peptin (pg/ul) — DOCA, DOCA+EM, Sham (****, ****)

K the level of co-peptin (pg/ul) — HSD, HSD+EM, NSD (* , *)

A

Patients diagnosed with heart failure and hospitalized
from January 2018 to January 2023 (n=1864)

Exclusion
Missing clinical data such as medication usage
(n=410); Severe anemia or on dialysis or severe
infection(n=155); Repeated hospitalization
(n=110).

Final study population (n=1189)

Non-treated group
(n=715)

Treated group
(n=474)

Propensity score
matching

Non-treated group
(n=437)

Treated group
(n=474)

B

		HCT	Neutrophil count	Monocyte count	Lymphocyte count
Alcohol	1	***	ns	ns	***
	2	*	ns	ns	ns
	3	**	ns	ns	ns
	4	***	ns	ns	ns
NT-pro BNP (pg/mL)	≤1364	***	ns	ns	ns
	1364.1-2550	***	ns	ns	ns
	2550.1-5137	***	ns	ns	*
	>5137	***	ns	*	**
Diuretic	NO	***	ns	*	***
	YES	***	**	ns	ns

Microglial

EM

Activated microglia

Neuron

Ang-II → Ang-(1-7)

AT1R

ACE2

Ubiquitination

NEDD4L

Ubiquitin

Brain

PVN

Salt

Vascular

Endothelial cell

EM

IFN-γ

lymphocytes

Sympathetic nerve output increases

cardiomyocyte hypertrophy
cardiac fibrosis

lymphocytes	AT1R	Ang-(1-7)
IFN-γ	ACE2	NEDD4L
Ubiquitin	Ang-II	Empagliflozin (EM)

Table 1 Baseline comparison of the treated group and non-treated group before and after propensity score matching

		Non-treat group	treat group	p value	Matcher non-treat group	Matched treat group	p value
		n=715 (60.1%)	n=474 (39.9%)		n=437 (48%)	n=474 (52%)	
patient characteristics							
Sex (n)	M	479	341	0.71	299	341	0.246
	F	236	133		138	133	
Age(years)		68(59-75)	66(58-66)	0.032	67(57-74)	58(66-73)	0.701
Weight(kg)		65(60-75)	70(60-79.25)	0.04	67(60-77)	70(60-79.25)	0.286
Height(cm)		168(160-172)	170(162-174.25)	0.03	169(160-173)	170(162-174.25)	0.182
Hypertension	No	279	206	0.127	186	206	0.785
	Yes	436	268		251	268	
2 type diabetes	No	493	252	<0.001	270	252	0.009
	Yes	222	222		167	222	
Dyslipidemia	No	638	421	0.771	383	421	0.582
	Yes	76	53		54	53	
angina	No	551	390	0.03	363	390	0.754
	Yes	164	84		74	84	
MI	No	527	343	0.609	320	343	0.77
	Yes	188	131		117	131	
AF	No	464	315	0.579	285	315	0.694
	Yes	251	159		152	159	
Chronic kidney disease	No	141	386	0.621	363	386	0.52
	Yes	574	88		74	88	
Smoke[a]	1	460	298	0.859	286	298	0.677
	2	85	58		52	58	
	3	12	11		6	11	
	4	158	107		93	107	
Drink[b]	1	605	378	0.004	363	378	0.006
	2	11	13		2	13	
	3	16	28		13	28	
	4	83	55		59	55	
SBP (mmHg)		129(113-145)	129(110.75-142.25)	0.516	127(112-142.5)	78(69-88)	0.788
DBP (mmHg)		76(69-87)	78(69-88)	0.426	76(69-87)	78(69-88)	0.343
Heart rate (bmp)		80(70-94)	82(71-97)	0.088	80(70-95)	97(96-98)	0.277
SpO$_2$ (%)		97(96-98)	97(96-98)	0.479	97(96-98)	97(96-98)	0.418
QRS duration (ms)		96(86-108)	100(90-118)	<0.001	98(88-114)	100(90-118)	0.083
Blood glucose (mmol/L)		5.45(4.8-6.181)	6.181(4.99-6.838)	<0.001	5.81(5.005-6.49)	6.18(4.99-6.84)	0.137
Hemoglobin		135(120-147)	135.5(122-150)	0.229	136(122-148)	135.5(122-150)	0.767
NT-proBNP (pg/mL)		2095(1193-4153)	2896.5(1570.5-5595)	<0.001	2216(1248.5-4851)	2896.5(1570.5-5595)	0.002
Serum Na$^+$ (mmol/L)		140.1(138-142.3)	140.4(138-142.3)	0.385	140(137.8-142)	140.4(138-142.3)	0.07
Serum K$^+$ (mmol/L)		4.1(3.8-4.5)	4.1(3.8-4.4)	0.505	4.1(3.8-4.5)	4.1(3.8-4.4)	0.365

Serum Cr (mmol/L)		83.8(68.6-104.7)	83.65(70.25-100.925)	0.533	82.9(68-102)	83.65(70.25-100.93)	0.685
Total cholesterol (mmol/L)		3.5(2.86-4.2)	3.445(2.848-4.215)	0.638	3.55(2.85-4.285)	3.45(2.85-4.22)	0.392
LDL		2.01(1.46-2.58)	1.995(1.5-2.635)	0.552	2.1(1.5-2.65)	2.0(1.5-2.64)	0.55
Serum uric acid (mmol/L)		389.701(311-429)	389.701(329.75-434)	0.245	389.701(324-437.5)	389.70(329.75-434)	0.948
LVEF (%)		54(42-62)	47(37-58)	<0.001	49.95(38-60)	47(37-58)	0.057
ACEI/ARB/ARIN	No	19	3	0.011	6	3	0.259
	Yes	696	471		431	471	
MRA	No	217	69	<0.001	69	69	0.604
	Yes	498	405		368	405	
Beta blocker	No	31	11	0.065	12	11	0.683
	Yes	684	463		425	463	
Diuretic	No	434	352	<0.001	253	352	<0.001
	Yes	281	122		184	122	

[a] 1: Never or smoking occasionally;

2: Quit smoking more than half a year;

3: Quit smoking less than half a year;

4: Smoking every day.

[b] 1: Never;

2: Drinking occasionally (less than 3 times a week);

3: Regular alcohol consumption (about 3 times a week);

4: Quit drinking.

MI: myocardial infarction

AF: atrial fibrillation

SBP: systolic blood pressure.

DBP: diastolic blood pressure.

LDL: low density lipoprotein.

LVEF: left ventricular ejection fraction.

MRA: mineralcorticoid recept antagonis.

ACEI/ARB/ARIN: angiotensin-converting enzyme inhibitors/ Angiotensin Ⅱ receptor antagonist/ angiotensin receptor-neprilysin inhibitor.

Table 2 Comparison of HCT and peripheral immune cell counts between the treated group and non-treated group after propensity matching score

	n=911	Matched non-treat group n=437 (48%)	Matched treat group n=474 (52%)	*P* value
HCT		39.9(37-43.4)	43(39.63-46.1)	<0.001
Neutrophil count		4.29(3.34-5.58)	4.18(3.19-5.31)	0.22
Monocyte count		0.52(0.4-0.68)	0.49(0.39-0.63)	0.02
Lymphocyte count		1.52(1.19-1.91	1.39(1-1.77)	<0.001