SPP1[+]Macrophages promote the growth and metastasis of intrahepatic

cholangiocarcinoma via OSM and THBS1 signaling

Short title: **SPP1[+]**Macrophages promotes ICC progression and metastasis

Tingjie Wang[1,3*], Ruitao Long[2*], Zhongyu Cheng[2], Mi Zhang[2], Bing Wei[1,3], Cong

Lyu[1,3], Cuiyun Zhang[1,3], Guangbo Tang[4], Zihang Li[5], Lei,Li[2], Yida Li[6], Xiaofei Zhu[7],

Chao Shi[1,3], Ke Yang[1,3], Chuanrui Xu[2#]

*Tingjie Wang and Ruitao Long contributed equally to this work.

[1]Department of Molecular Pathology, The Affiliated Cancer Hospital of Zhengzhou

University, Henan Cancer Hospital, Zhengzhou, Henan, People's Republic of China.

[2]School of Pharmacy, Tongji Medical College, Huazhong University of Science and

Technology, Wuhan, China

[3]Henan Key Laboratory of Molecular Pathology, Zhengzhou, People's Republic of

China.

[4]Department of Clinical Medicine Research Center, The First Affiliated Hospital of

Gannan Medical University, Ganzhou, Jiangxi, People's Republic of China.

[5]Department of Hematology, The First Affiliated Hospital of Xi'an Jiaotong University,

No.277 Yanta West Road, Xi'an, Shaanxi, 710061, China.

[6]Department of Chemistry & Loker Hydrocarbon Research Institute, University of

Southern California, Los Angeles, California 90089, United States.

[7]Henan Key Laboratory of Immunology and Targeted Drugs, Xinxiang Key

Laboratory of Tumor Microenvironment and Immunotherapy, School of Medical

Technology, Xinxiang Medical University, Xinxiang, People's Republic of China.

Correspondence:

Address correspondence to: Chuanrui Xu, School of Pharmacy, Tongji Medical

College, Huazhong University of Science and Technology, Wuhan, Hubei, China.

Email: xcr@hust.edu.cn

Conflict of interest:

The authors declare no conflicts of interest to this work.

Abbreviations

TAMs, tumor-associated macrophages; ICC, intrahepatic cholangiocarcinoma; TME,

tumor microenvironment; EMT, epithelial mesenchymal transition; Mal, malignant

tumor cells region; Bdy, tumor boundary; nMal, adjacent non-malignant region; B, B

cells; DC, dendritic cells; CAF, cancer-associated fibroblasts; CHO, cholangiocyte;

HPE, hepatocyte; T_NK, T and NK cells; TEC, tumor endothelial cells. NEU,

neutrophils. EPI, epithelial cells. BMDMs, bone marrow–derived macrophages.

scRNAseq, single cell RNA sequencing; spRNAseq, spatial transcriptomics

sequencing; scATACseq, single cell ATAC sequencing.

Funding:

The research was supported by grants from National Natural Science Foundation of

China (82472988, 82403200, 82372667, 82273059 and 82573814), Henan Province

and Ministry of Health of Medical Science and Technology Program

(SBGJ202302028 for T.W. and SBGJ202402032 for C.L.), Henan Province Science

and Technology Research Project (242102310051), and Henan Province Clinical

Research-Oriented Physician Training Program (HNCRD202413).

Authors contributions:

C.X., T.W. designed the study. T.W., C.Z., analyzed the multi-omics data. R.L.

performed experiments. T.W., R.L. wrote the manuscript. C.X. revised the article. Z.C,

M.Z., C.L., B.W., L.L., X.Z. handled sample preparation. Y.L., G.T., C.S., K.Y., Z.L.

managed data collection, organization. All authors read and approved the manuscript.

Data availability:

The scRNA-seq data generated in this study are available from the GEO database

under accession code GSE308574. The remaining data are available within the

manuscript, supplementary information, or from the corresponding author upon

reasonable request.

Acknowledgements

This work was financially supported by the National Natural Science Foundation of

China (Grant No. 82472988, 82403200, 82372667, 82273059 and 82573814).

Abstract

Background & Aims: Tumor-associated macrophages (TAMs) are abundant in

intrahepatic cholangiocarcinoma (ICC), correlate with poor prognosis, but their

spatiotemporal behavior and non-immune-suppressive roles in ICC progression

remain unclear. This study sought to define transcriptional heterogeneity of TAMs and

map their interactions with ICC cells that promote malignancy beyond immune

suppression.

Methods: We employed three mouse ICC models, four clinical ICC patient datasets,

and three spatial transcriptomic approaches to chart the macrophage landscape over

time and space. Cell-cell communication analysis plus phosphoproteomics guided

functional validation via TAM-ICC co-culture, cytokine treatment, gene

silencing/knockout, orthotopic and metastatic ICC models, and three conditional

knockout mice. A prognostic ICC tissue microarray with multiplex

immunofluorescence and combinatorial blockade in mice tested efficacy of dual target

inhibition. Single-cell ATAC-seq provided epigenomic profiles of TAM subsets,

confirming M2 polarization networks via TCM treatment and gene silencing.

Results: In advanced ICC tissues, an M2-like subset Mφ_SPP1, defined by high

expression of *SPP1*,*VEGFA*, *ARG1* and *GPNMB*, emerged as the major macrophage

population, correlating with poor prognosis and spatially localizing to the tumor

boundary. Further, Mφ_SPP1 drove ICC cell proliferation via the

OSM-OSMR-STAT3-CCND1 and promoted invasion through

THBS1-CD47-mTOR-DNB1 signalling. OSM and THBS1 were co-elevated in ICC

patient samples; dual blockade of both targets in mouse models resulted in superior

inhibition of ICC progression. Lastly, MAFB directed M2 polarisation and

upregulated *OSM/THBS1* in Mφ_SPP1.

Conclusions: Mφ_SPP1 TAMs accumulate at the tumor boundary in late-stage ICC,

driving proliferation and invasion via *OSM* and *THBS1* signalling, underscoring the

potential of macrophage-targeted therapy in ICC.

Keywords: Intrahepatic cholangiocarcinoma, Spatiotemporal omics,

SPP1[+]Macrophage，OSM，THBS1

Introduction

Intrahepatic cholangiocarcinoma (ICC), originating from the intrahepatic bile duct

epithelium, shows rising global incidence. Its subtle early symptoms, aggressive

nature, and chemotherapy resistance drive a 5-year survival rate below 10%.[1]

Targeted therapies and immune checkpoint inhibitors benefit some solid tumors, but

responses in ICC remain inconsistent owing to its complex tumor microenvironment

(TME).[2] Tumor-associated macrophages (TAMs), the most abundant immune cells in

the ICC TME, promote immune suppression, angiogenesis, and stromal remodelling,[2,

3] fuelling progression and metastasis. Current TAM-targeting strategies—depletion,

reprogramming, recruitment control, and functional regulation—show variable

efficacy.[4, 5] Heterogeneity is a major barrier: CD163☐ macrophages have low

CSF-1R, reducing CSF-1R inhibitor response; CCL2 inhibitors are of limited use as

CCR2 is also expressed on T and NK cells; and prolonged macrophage depletion

disrupts immune homeostasis, causing adverse effects.[4] Thus, dissecting

communication cascades between ICC cells and TAMS is critical to identify

therapeutic targets that directly modulate ICC.

Distinct TAM phenotypes are associated with ICC progression. MARCO☐ TAMs

correlate with epithelial mesenchymal transition (EMT), angiogenesis, and hypoxia.[6]

S100P and *SPP1*—markers distinguishing perihilar and peripheral

cholangiocarcinoma—show differential macrophage infiltration tied to prognosis and

grade.[7] Moreover, TAM states are highly dynamic. Prior multi-omics studies largely

used patient samples, failing to track temporal polarisation across tumor stages.

Epigenetic regulation is also critically involved, but a temporal atlas of macrophage

epigenetics in ICC is lacking.

Spatial distribution further influences TAM functions.[8] Spatial-omics have

highlighted TAM heterogeneity.[9, 10] TAMs at invasive fronts mainly express

pro-angiogenic factors such as VEGFA, while core TAMs activate

immunosuppressive pathways.[6] Moreover, TAM plasticity is shaped by cues such as

extracellular matrix (ECM) gradients, lactate accumulation, and cytokines.[9] However,

the spatial dynamics of TAMs in ICC remain poorly defined.

Here, we developed a temporal ICC mouse model, tracked TAM evolution, and

identified shared human subpopulations, mapping their transcriptional and regulatory

landscapes. Using cell-cell communication analysis, co-culture, multiplex

immunofluorescence, and knockout models, we validated key TAM factors in ICC

progression. These findings are expected to support macrophage-targeted

immunotherapy for ICC.

Materials and methods

Animal Models：

Wild-type FVB and C57BL/6 mice were purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd. (Beijing, China). Genetically modified mice were provided by Shulaibao (Wuhan) Biotechnology Co., Ltd. (Wuhan, China). All laboratory mice were kept in a specific pathogen-free (SPF) environment. Experimental protocols involving animals were approved by the Ethics Committee of Huazhong University of Science and Technology. Detailed information on the animals used in this study is provided in Supplementary Table S1.

Cell source

The human cholangiocarcinoma cell lines HuCCT1, RBE, and QBC939, as well as the mouse fibroblast cell line L929 and macrophage cell line RAW 264.7, were used in this study. HuCCT1 was obtained from the Japanese Collection of Research Bioresources Cell Bank (JCRB, Osaka, Japan; accession number JCRB0425). RBE was acquired from the China Center for Type Culture Collection (CCTCC, Wuhan, China; accession number GDC0765). QBC939 was purchased from Shanghai Jinyuan Biological Technology Co., Ltd (Shanghai, China; product code JY-Y1457). The mouse cell lines L929 (ATCC® CCL-1™) and RAW 264.7 (ATCC® TIB-71™) were purchased directly from the American Type Culture Collection (ATCC, Manassas, VA, USA). All cells were cultured according to the providers' recommendations and were routinely tested to be free of mycoplasma contamination.

BR cells, isolated from BMI1/NRAS-driven orthotopic tumor models in FVB mice, were maintained in RPMI-1640 complete medium supplemented with 10% FBS and 1% penicillin–streptomycin at 37°C with 5% CO_2 in an incubator. These cells displayed consistent in vitro growth characteristics and were capable of forming tumors in both subcutaneous and orthotopic liver implantation models in wild-type FVB mice. Immunohistochemistry (IHC) and histopathological analysis confirmed the presence

of ICC features.

Establishment of the ICC mouse model

Hydrodynamic tail vein injection was performed in 6 – 8 weeks mice to establish

primary ICC mouse models. For AKT1/NICD1 ICC mouse model , a mixture of

plasmids, including 15 μg pT3-EF1aH-myr-AKT, 15 μg pT3-EF1a-NICD1, and 1.5

μg pCMV-SB, was diluted in 2 mL of sterile saline and filtered through a 0.22 μm

membrane to remove impurities. For AKT1/YAP1 ICC mouse model, a mixture of

plasmids, including 15 μg pT3-EF1aH-myr-AKT, 15 μg pT3EF1αH-YAPS127A, and

1.5 μg pCMV-SB, was diluted in 2 mL of sterile saline. For sgP19/KRAS

high-metastasis ICC model, a mixture of plasmids, including 10 μg

CRISPR-Cas9-sgP19, 25 μg pT3-EF1α-KRAS, and 5.0 μg pCMV-SB, was diluted in

2 mL of sterile saline. The injection volume for a 20 g mouse was 2 mL and adjusted

according to body weight in proportion. Mice were restrained on a tail vein

visualization platform, and the injection was completed within 5 – 7 seconds. Mice

were euthanized and dissected upon reaching predefined tumor endpoint criteria.

The plasmids were generously provided by Dr. Chen Xin (University of Hawai□i

Cancer Center). Bacterial cultures were grown in LB medium containing 100 μg/mL

ampicillin at 37°C with shaking at 180 rpm for 12 hours, followed by plasmid

extraction using the Omega plasmid purification kit.

Multi-omics analyses and bioinformatic analysis

scRNA-seq, spatial transcriptomics, scATAC-seq, phosphoproteomics were performed.

Detailed methods are provided in the Supplementary Methods.

Results

Mφ_SPP1 is dominant at the ICC boundary and linked to poor prognosis

To dissect macrophage subsets driving ICC malignancy, we established an

AKT1/NICD1 induced ICC mouse model via hydrodynamic tail vein injection and

collected liver tissue at day 7 (early ICC), day 15 (mid ICC), day 25 (mid/advanced

ICC), and day 31 (advanced ICC). Single cell RNA sequencing (scRNAseq) was

performed at all stages and spatial transcriptomics sequencing (spRNAseq) was

performed at day 31, with integrated analysis of public human datasets (Table S2, Fig.

S1A-F).

Four conserved TAM subsets were identified in mouse and human ICC:

Mφ_CXCL9(human)/Mφ_Cxcl9(mouse),Mφ_S100A/Mφ_S100a,Mφ_C1QA/Mφ_C1

qa, and Mφ_SPP1/Mφ_SPP1 (Fig. 1A-B, S1, Table S2). Mφ_Cxcl9 and Mφ_S100a

dominated early/mid-stage mouse ICC (Fig. 1B), while in humans, Mφ_CXCL9

localised to adjacent liver and elevated expression of *CXCL9*, *CXCL11*, and *CCL5*

(Fig. S1C, S1G), while Mφ_S100A was enriched in human ICC tissues with elevated

CD68, *VCAN*, *S100A4*, *S100A8* (Fig. 1C, S1C, S1G), suggesting an inflammatory

early ICC microenvironment. In contrast, Mφ_C1QA/Mφ_C1qa and

Mφ_SPP1/Mφ_SPP1 were enriched in human ICC and advanced mouse ICC, with

similar transcriptional signatures (Fig 1C). Mφ_SPP1 showed elevated *GPNMB*,

VEGFA, *ARG1*, *APOE*, *TREM2*, and *SPP1* levels (Fig 1C , S1C, S1G), consistent with

M2-like, tumor-promoting macrophages.[11, 12] Mφ_SPP1 showed significant

enrichment in cytokine secretion and tumor proliferation pathways (Fig. S1H). Similar

TAM subtypes and temporal patterns were confirmed in AKT1/YAP1 ICC mice

(Figure S2A–C). Kupffer cells (*Clec4f⁺Mrc1⁺ and Vegfa/Gpnmb/Spp1*-low), were

abundant in AKT1/NICD1 and AKT1/YAP1 ICC tissues, likely from liver damage

caused by hydrodynamic injection (Fig. 1C, S1G, S2A–D).

Analysis of dynamic macrophage temporal distribution using pseudo-time

analysis showed that Mφ_Cxcl9 andMφ_S100a predominated the early pseudo-time

branch, while Mφ_SPP1 progressively expanded in later branches (Fig 1D, S2E),

indicating that Mφ_SPP1 is the dominant population in advanced ICC tissues.

Prognostic analysis across three independent cohorts (GSE89749,[13] GSE107943,[14]

GSE244807[15]) confirmed the robust association of Mφ_SPP1 with poor outcome of

patients with ICC (Fig. 1E, S2F).

In mouse ICC allografts, established by co-injection of BMDMs with BR ICC

cells, tumor growth was significantly enhanced by macrophages, characterised by

F4/80□ and CD206□ (Fig. 1F-i, S3), while depletion of macrophages using

clodronate   liposome   (Fig.   S4)   suppressed   tumor   progression

(CD11b□F4/80□/CD45□), evidenced by increased liver-to-body weight ratio (Fig.

1F-ii). Co-culture of ICC cells with macrophages caused M2-like polarisation of

macrophages, with robust upregulation of M2 polarisation markers including *ARG1*,

VEGFA, *IL10*, and *MRC1* (Fig. S5), using non-contact Transwell co-cultures or

treatment with tumor-conditioned medium (TCM)(Fig. S5B-D). In Transwell

co-culture systems, EdU incorporation assay and migration assay demonstrated that

macrophage co-culture significantly enhanced ICC cell proliferation (Fig. S6A-D) and

migration (Fig. S6E).

Next, utilizing three spatial transcriptomics platforms and multiplex

immunofluorescence, we explored the spatial distribution of Mφ_Spp1 and its spatial

proximity to ICC cells across one public dataset[16] and four in-house ICC datasets

(Table S2). In the public human ICC dataset with the 10X Visium platform, we

observed that spatial clusters C0 and C4, located in the non-malignant region (nMal),

exhibited a high infiltration of hepatocytes (HPE) (Fig. 1G(i-ii)). In contrast, in the

malignant (Mal) and boundary (Bdy) regions, clusters C1, C2, and C3 were enriched

in ICC cells and $SPP1^+$ macrophages (Fig. 1Gii-iii, S7). Simultaneously, we

performed tumor niche analysis and confirmed the co-occurrence of $CK19^+$ epithelial

and macrophage cells (Fig. 1G-iv). Additionally, in the high-resolution spatial

transcriptome analysis of D31 AKT1/NICD1 ICC tissue using the BMKMANU

S1000 platform, we identified that the Mφ_Spp1 subtype was predominantly localized

to the tumor boundary, adjacent to $CK19^+$ ICC cells within the same tumor niche (Fig.

1H, S7). Similarly, in the in-house 10X HD transcriptome data from four human ICC

tissues, we conducted spatial domain analysis using BANKSY and RCTD, further

validating the spatial proximity between Mφ_SPP1 and $CK7^+CK19^+$ ICC cells (Fig.

1I i-iv). Specifically, we assessed whether the abundance of ICC cells could be

predicted by their spatially adjacent TME cells using mistyR in ICC patient #1. We

found that Mφ_Spp1 cells were the most predictive of the abundance of $CK7^+CK19^+$

ICC cells (Fig. 1I (v)), a finding also corroborated in the other three patients (Fig.

S7D-E). We also validated the spatial proximity between $SPP1^+CD68^+$ and $CK19^+$

cells in the other two ICC patient cohorts using multiplex immunofluorescence (mIF)

(Fig. 1J). The results showed that SPP1$^+$ CD68$^+$ macrophages were highly enriched in

the periphery of CK19$^+$ cells (0–20 μm), and their abundance gradually diminished

with increasing spatial distance(Fig. 1J).

Together, these findings indicate that M2-like Mφ_SPP1 accumulate at the ICC

boundary, driving proliferation and invasion through cytokine secretion.

Mφ_SPP1 promotes ICC proliferation via the OSM-OSMR-STAT3-CCND1 axis

To explore the mechanisms underlying Mφ_SPP1-mediated ICC progression, we

re-clustered and obtained seven epithelial cell subpopulations from human and

AKT1/NICD1 mouse ICC. Within human/mouse shared subtypes, E_Alb in mice or

E_ALB in human were normal hepatic epithelial cells with high *ALB* levels (Fig. 2A,

2B, S8A). E_Fxyd2 (mouse) or E_FXYD2 (human) were normal cholangiocytes with

high *FXYD2* levels (Fig. 2A, 2B, S8A). In contrast, E_Jun/E_Spp1 (mouse) and

E_CD74/E_IGFBP7 (human) showed high CNV scores (Fig. S8B) and were

identified as ICC cells. Specifically, E_Jun (mouse)/E_Cd74 (human) displayed high

expressions of *STAT3*, *JUN*, *HSPA1A* and *CCND1* (Fig. 2A, S8A), indicating their

stress-responsive and highly proliferative nature.[17, 18] Meanwhile, E_Jun/E_Spp1

(mouse) and E_IGFBP7 (human) exhibited high levels of the EMT-related genes *VIM,*

COL4A1, and *FN1* (Fig. 2A, S8A). In addition, E_Mki67 (mouse)/E_MKI67 (human)

displayed high expressions of MKI67 but low expressions of receptor genes.

Integration of three cell-cell communication approaches (CellChat, CellphoneDB,

NicheNet) to analyse interactions between Mφ_SPP1 and ICC cells with high

proliferation and EMT features and the downstream signalling pathways, revealed

that Mφ_SPP1 engages with ICC cells via *OSM* and *THBS1* ligands (Fig. 2C–D).

OSM, enriched in Mφ_Spp1 (mouse) and Mφ_SPP1 (human), interacted with ICC

cells of E_Jun in mice and E_CD74 in human via *OSMR* and *LIFR*, which in turn

activated the genes involved in cytokine response and epithelial proliferation (Fig. 2E,

S8C–D).

To test whether *OSMR* and *LIFR* are required for ICC progression, we generated

Osmr- or *Lifr-* liver conditional knockout mice Osmr$^{fl/fl}$;Alb-cre and Lifr$^{fl/fl}$;Alb-cre,

followed by hydrodynamic injection of AKT1/NICD1 via tail vein. Compared with

Lifr $^{fl/fl}$ mice, Lifr $^{fl/fl}$; Alb-Cre mice showed no significant difference in liver-to-body

weight ratio upon AKT1/NICD1 injection (Fig. 2F–G, S9A-B). However, Osmr

$^{fl/fl}$;Alb-Cre mice exhibited a marked reduction of liver-to-body weight ratio after

AKT1/NICD1 injection compared to Osmr $^{fl/fl}$ mice (Fig. 2F–G, S9A-B). qRT-PCR

revealed that *Osm* and *Osmr* levels increased with tumor progression (Fig. S9C).

Spatial interaction analysis (Fig. S10A-B), expression level in ICC scRNAseq cohorts

(GSE210066 and GSE201425, Table S2, Fig. S10C), and spatial mapping by

multiplex IF (Fig. 2H, S10G) all confirmed the presence of OSM□ macrophages in

the vicinity of OSMR□ tumor cells.

Functionally, RAW264.7 and QBC939 co-cultures increased *OSM*, *VEGFA*, and

IL10 in macrophages, and *OSMR* in QBC939 cells (Fig. 3A, S11A). Similarly,

co-culture with BMDMs increased OSMR protein levels in BR cells (Fig. S3B-i).

Recombinant OSM increased OSMR mRNA and protein levels, and viability of ICC

cells (Fig. 3B). OSMR knockdown (siRNA) significantly suppressed ICC cell

viability (Fig. S11C-D). Similarly, CRISPR-Cas9-mediated knockout of *OSMR*

resulted in significantly decreased proliferation, viability, and clonogenicity of ICC

cells (Fig. 3C, S11E).

Phosphoproteomics of QBC939 cells treated with recombinant OSM revealed

enrichment of PI3K-AKT, MAPK, and cell cycle-related pathways (Fig. 3D).

Moreover, *Osmr* in the tumor subtype E_Jun was involved in epithelial cell

proliferation, cytokine response, and MAPK pathways, and interacted with *Stat3* and

Ccnd1 (Fig. 3E). Thus, OSM-OSMR may activate STAT3-CCND1 signalling in

tumor cells. To test this hypothesis, we used Osmr$^{fl/fl}$;Alb-Cre mice injected with

AKT1/NICD1. H&E and IHC staining demonstrated that ICC lesions formed in

Osmr$^{fl/fl}$;Alb-Cre mice were significantly smaller than those in Osmr$^{fl/fl}$ mice (Fig. 3F,

S12A). The tumor-suppressive effect of *Osmr* knockout was similarly validated in a

hydrodynamic injection-constructed AKT1/YAP1 ICC mouse model, manifesting as

significantly reduced liver-to-body weight ratio and diminished tumor nodule

formation (Fig S12B, C). Notably, *Osmr* knockout did not alter F4/80□ and CD206□

macrophage infiltration levels (Fig. S12D). However, the levels of the downstream

proteins CCND1, BCL2, and p-STAT3 were markedly reduced (Fig. 3G), indicating

that Mφ_SPP1 promotes ICC growth via OSM-OSMR signalling and downstream

STAT3 phosphorylation, suggesting that deficiency of OSM-OSMR signalling leads

to retarded ICC growth in mice.

Mφ_SPP1 TAMs promote ICC invasion through the *THBS1-CD47-mTOR-DBN1*

axis

In addition to *OSM, THBS1* is secreted by Mφ_SPP1 which may promote ICC

development. Cell communication analysis showed that *THBS1* interacted with E_Jun

and E_Spp1 in AKT1/NICD mouse ICC tissues and with E_IGFBP7 in human ICC

via *CD47* (Fig. 4A, Fig. S13A). Using NicheNet, we found that *THBS1*-regulated

target genes in ICC cells were mainly enriched in EMT-related signalling (Fig. 4B,

S13B). Subsequently, we validated the THBS1-CD47 interaction both in vitro and in

vivo. In vitro, recombinant THBS1 exhibited moderate growth-promoting and

enhanced pro-migratory effects in ICC cells (Fig. S14A, B). Specifically, treatment

with 10 ng/mL recombinant THBS1 significantly enhanced wound-healing (Fig. 4C)

and migration of ICC cells (Fig. 4D). Correspondingly, *CD47* silencing significantly

impaired the stimulatory effects of THBS1 in promoting ICC cell migration (Fig.

4E–F, Fig. S14C, D). Stable knockdown of *Cd47* in BR cells significantly inhibited

the proliferation, colony formation capability and viability of ICC cells (Fig. 4G–J).

Co-culture with BMDMs promoted BR migration ability, whereas *Cd47* knockdown

blocked this effect (Fig. 4K). Similarly, *Cd47* silencing inhibited subcutaneous

allograft growth in mice (Fig. 4L, Fig. S14E, F). Moreover, *Cd47* knockdown

markedly suppressed ICC dissemination in a peritoneal metastasis mouse model, as

evidenced by reduced metastatic nodule formation and decreased ascites

accumulation (Fig. 4M(i-ii), Fig. S14G, H). Consistently, CD47 silencing attenuated

ICC growth and local invasion into the surrounding hepatic parenchyma in the

primary ICC mouse model (Fig. 4M-iii).

To elucidate the role of *THBS1*-mediated EMT in ICC, we performed western

blotting, which revealed that recombinant THBS1 induced rapid induction of

EMT-related proteins at 10 min with a peak at 2 h, accompanied with increased levels

of p-mTORC1 and downstream p-RPS6 (Fig. 5A, S15A). Phosphoproteomic analysis

of HuCCT1 cells treated with recombinant THBS1 at 0, 2, and 12 h revealed elevated

phosphorylation of PI3K-AKT and EMT-related signalling proteins at 2 h (Fig. 5B).

In comparison, differential protein phosphorylation was enriched in cell adhesion and

RNA metabolism pathways at 12 h (Fig. 5B, S15B). Specifically, ranking

differentially phosphorylated sites by fold change (2 h vs 0 h) identified DBN1

Ser142 as the most strongly upregulated phosphorylation site at 2 h (Fig. 5C).

Co-culture of ICC cells with BMDM led to increased *Arg1, Vegfa*, and *Thbs1* in

BMDMs, CD47, p-mTORC1, p-DBN1 and Vimentin in ICC cell lines (Fig. 5D, Fig.

S16A–B). The upregulation of *Thbs1* and *Cd47* was also detected by qRT-PCR in

AKT1/NICD1-induced liver tissues (Fig. S16C–D). Inhibition of mTORC1 by

rapamycin substantially suppressed the pro-migratory effect of BMDMs on ICC cells

(Fig. 5E) and completely abrogated the THBS1-induced migration (Fig. 5F),

indicating that THBS1 promotes ICC cell migration via mTORC1 activation.

Meanwhile, inhibition of DBN1 via its inhibitor BTP2 significantly blocked the

migration of ICC cells (Fig. 5G). Notably, BTP2 exerted its inhibitory effect by

decreasing phosphorylation at Ser142 of DBN1 (Fig. S16 E). Together, these results

demonstrate the critical roles of p-mTORC1 and DBN1 Ser142 phosphorylation in the

THBS1-mediated migration signalling pathway.

Finally, we examined the role of THBS1 in ICC progression by combing a highly

metastatic sgP19/KRAS ICC model with Thbs1$^{fl/fl}$;Lyz-Cre mice (Fig. 5I).

Myeloid-specific konckout of *Thbs1* significantly reduced liver-to-body weight ratios

and tumor burden, while also markedly suppressing peripheral metastasis to multiple

distant organs - including the intestines, lungs, lymph nodes, kidneys and diaphragm

(Fig. 5H-J，FigS16F). Notably, THBS1 knockout also demonstrated significant

efficacy in suppressing ICC within the AKT1/NICD1 model—which lacks prominent

peripheral metastasis (Fig.5 K, L). This finding underscores the crucial contribution of

local intrahepatic spread to ICC progression. Furthermore, Western blotting of ICC

tissues revealed that metastasis related CD47, DBN1, p-DBN1, p-mTORC1, and

Vimentin levels were significantly downregulated after *Thbs1* deletion (Fig. 5M).

Collectively, these findings corroborate that THBS1 promotes ICC metastasis via

PI3K-AKT-mTOR activation and DBN1 phosphorylation.

Combined blockade of OSM and THBS1 effectively inhibits ICC progression

Since *OSM* promotes proliferation and *THBS1* enhances invasiveness, we asked

whether dual inhibition could synergistically suppress ICC. We analysed *THBS1* and

OSM expression in 833 patients from 9 ICC/Cholangiocarcinoma cohorts (Table S2),

and calculated the mean expression levels of *THBS1* and *OSM* in each cohort. An ICC

sample was defined as double-positive if both genes had expression levels greater

than the mean value of the cohort. Using this criterion across 9 datasets, the

proportion of *THBS1/OSM* double-positive ICC samples ranged from 13.33% to

29.53%, and exceeded 20% in five datasets (Fig. 6A). Prognostic analyses across

three datasets showed that patients with high expression of both OSM and THBS1

had significantly shorter overall survival (OS) (Fig. 6B). Next, we genetically ablated

macrophage-derived *THBS1* using Thbs1[fl/fl]; Lyz-cre mice and pharmacologically

inhibited   OSM-OSMR   interaction   via   intraperitoneal   administration   of

OSM-SMI-10B, an OSMR inhibitor.[19] *Thbs1* deletion or *Osmr* inhibition alone

suppressed ICC progression moderately, while combined blockade attenuated ICC

development more robustly, as evidenced by further reduced tumor burden using

H&E staining and a decreased liver-to-body weight ratio (Fig. 6C–E). These results

collectively indicate that combined blockade of OSM and THBS1 may generate

synergetic therapeutic effects for ICC treatment. Finally, we validated the spatial

proximity relationship between OSM/THBS1 and CK19 cells using mIF (Fig. 6F).

The results showed that OSM[+]THBS1[+]CD68[+] cells were spatially adjacent to CK19[+]

cells (Fig. 6F i-ii), OSM and THBS1 were co-expressed in CD68[+] cells (Fig. 6F-iii).

Patients with high OSM/THBS1 co-expression was associated with significantly

worse survival (Fig. 6F-iv). These findings suggest that macrophage-derived OSM

and THBS1 represent promising therapeutic targets in a substantial subset of ICC

patients.

Mafb promotes M2 polarisation and enhances *OSM* and *THBS1* expression in

macrophages

We next examined epigenetic modification in macrophages. Liver tissues from

AKT1/NICD1 ICC mice at middle/advanced stages and vector controls were

subjected to time-series single cell ATAC sequencing (scATAC-seq) to map regulatory

elements in Mφ_SPP1 (Fig. S17A–C). Accessibility profile analysis of cell type

markers using ArchR (V1.0.3) and integrated AKT1/NICD1 scRNA-seq and

scATACseq data identified seven major cell types (CAF, TEC, EPI, B, NEU, T_NK,

and TAMs; Fig. 7A, S17A–B). Further, re-clustering of TAMs revealed three

subtypes—Mφ_Clec4f, Mφ_Spp1, and Mφ_S100a (Fig. 7A, S17C). ArchR to analyze

chromatin accessibility and transcription factor (TF) motifs across TAM subtypes

revealed that Mφ_Spp1 and Mφ_S100a subsets exhibited markedly distinct regulatory

landscapes (Fig. 7B). In Mφ_S100a, the Klf transcription factor family exhibited

specific activation (Fig 7B, S17D), with TF *Klf3* showing the highest expression and

regulating inflammation response genes including *S100a4* and *S100a8* (Fig. 7C, 7D,

S17D, S17E). In comparison, Mφ_Spp1 displayed preferential chromatin accessibility

of *Maf* transcription factor family (Fig 7B), with *Mafb* showing the highest expression

(Fig. 7C). Notably, *Mafb* regulated genes were enriched in cell differentiation

pathways—*Mafb* was predicted to regulate M2 polarisation markers (*Cd163*, *Mrc1*,

Il10) and *Osm* (Fig. 7D, S17F). *MAFB* was highly expressed at the tumor boundary in

mouse and human ICC tissues (Fig. 7E). *Mafb* also showed significantly increased

chromatin accessibility in Mφ_Clec4f (Kupffer cells). Mφ_Clec4f exhibited high

accessibility of *Cd163* and *Mrc1* but low expression of *Osm* and *Thbs1* (Fig. S17G),

indicating that the regulatory role of *Mafb* on *Osm/Thbs1* is specific to Mφ_Spp1.

Together, these data suggest that *Mafb* plays an important role in regulating

macrophage M2 polarisation.

To validate the role of Mafb in Mφ_Spp1, we combined macrophage pseudotime

analysis with immune infiltration profiling to identify key signals driving macrophage

polarization (Fig. S18). Pseudotime-associated genes were grouped into five clusters,

among which cluster 4 was enriched for mononuclear cell differentiation (Fig.

S18A–B). We then selected genes with an average expression > 1 in Mφ_Spp1 and a

positive correlation with M2 polarization (r > 0.5) in the TCGA-CHOL cohort using

TIMER2.0, yielding eight candidates, including the transcription factor *MAFB* and the

receptor *TGFBR2* (Fig. S18C–D).

We next performed cell–cell communication analysis between ICC cells and

TAM subtypes in the AKT1/NICD1 dataset, and found that the E_Jun ICC subcluster

interacts with Mφ_Spp1 via the Tgfb2–Tgfbr2 ligand–receptor pair (Fig. S18E–F). In

the Transwell co-culture system, *TGFB2* was upregulated in ICC cells, while *Tgfbr2*

was upregulated in BMDMs (Fig. S19A). As TGF-β2 has also been reported to

regulated M2 polarisation,[20] We found that TCM or recombinant TGF-β2

significantly upregulated the expression of M2 markers in BMDMs, including *Mrc1*

(Cd206), *Arg1*, and *Mafb*-regulated genes *Tgfbr2*, *Thbs1*, and *Osm* (Fig 7F, 7G, Fig

S19B, C). Crucially, siRNA knockdown of *Mafb* significantly reduced expression of

Arg1, Vegfa, Osm, and *Thbs1* (Fig. 7H). These results indicate that *MAFB* could

regulate macrophage M2 polarisation via *OSM* and *THBS1* signalling.

Discussion

The roles of TAMs in ICC have been linked to poor prognosis through promotion of tumor growth and invasiveness.[21, 22] Recent single-cell omics studies have revealed the molecular diversity and functions of TAMs in cancer,[23] but the specific subgroups driving ICC malignancy and its interactions with tumor cells remain unclear. Using a time-resolved ICC mouse model with integrated multi-omics, we identified Mφ_SPP1 as a poor-prognosis TAM subset in ICC. Spatial transcriptomics confirmed its boundary localisation and functional niche. Mφ_SPP1 displayed mainly M2-like traits with distinct marker genes separating it from other TAMs. Its prognostic significance was also observed in hepatocellular carcinoma (HCC),[11] underscoring relevance across liver cancer, highlighting Mφ_SPP1 as a therapeutic target for both ICC and HCC.[24] Mechanistically, Mφ_SPP1 promotes ICC proliferation and invasion through coordinated multiple pathway via OSM-OSMR signalling, activating STAT3 phosphorylation and CCND1 induction, similar to glioblastoma[25] This observation is consistent with the finding that OSM promotes cancer cell proliferation and differentiation in breast cancer and pancreatic ductal adenocarcinoma.[26] Its pro-invasive function depends on the secreted matricellular glycoprotein THBS1, which activates CD47 signalling in ICC cells. THBS1 derived from tumor-infiltrating monocyte-like cells reportedly induces an immunosuppressive TME and contributes to cancer metastasis.[27] Notably, we demonstrate that *CD47* activates mTOR in ICC cells but not IL6/STAT3, resulting in DBN1 Ser124 phosphorylation, as also observed

in oral squamous cell carcinoma.[28] While THBS1/CD47 mediate immune evasion,

our study reveals its direct role in ICC proliferation and invasion.

Our findings offer a new therapeutic strategy for ICC treatment aiming at

macrophages. OSM/OSMR and THBS1/CD47 axes mediate macrophage-ICC

interactions, suggesting their potential as therapeutic targets for ICC. Although

administration of single OSM or mTOR inhibitors (blocking CD47 signalling)

achieved limited inhibitory effect on ICC growth, combined blockade of OSM-OSMR

and THBS1/CD47 signalling generated synergetic anti-tumor effects in an ICC mouse

model. Consistently, a clinical trial showed that targeting OSM/OSMR signalling

alone led to poor outcome or drug resistance.[29] Thus, targeting dual modulators of

Mφ_SPP1 provides a potentially more robust therapeutic option. Further, OSM and

THBS1 were concurrently upregulated in 13–29% of patients with ICC and

OSM/OSMR and THBS1/CD47 signalling promoted ICC proliferation and migration,

respectively. Therefore, combination therapy targeting these two signalling pathways

may be effective in patients with ICC at different stages.

Lastly, our study indicates that Mφ_SPP1 polarisation is dependent on MafB and

demonstrates a metabolic profile with highly active lipid metabolism. MafB has been

reported with c-Maf functioning as a self-renewal blocker in monocytes and

macrophages, and deficiency of these two genes enables extended expansion of

mature monocytes and macrophages in culture while maintaining their differentiated

phenotype and function.[30] In addition, macrophages express acute inflammatory

response genes in a Maf-dependent manner upon bacteria exposure and

simultaneously downregulate Maf through accelerated degradation via the proteasome

pathway and microRNA-mediated silencing.[31] In this study, we further demonstrate

that MAF is required for macrophage polarisation in the TME and highlight the roles

of MAF in orchestrating macrophage activation and polarisation.

Supplementary Material

Note: To access the supplementary material accompanying this article, visit the online

version of *Gastroenterology* at www.gastrojournal.org,

References

[1] Moris D, Palta M, Kim C, et al. Advances in the treatment of intrahepatic cholangiocarcinoma: An overview of the current and future therapeutic landscape for clinicians. CA: a cancer journal for clinicians 2023;73:198-222. https://doi.org/10.3322/caac.21759.

[2] Greten TF, Schwabe R, Bardeesy N, et al. Immunology and immunotherapy of cholangiocarcinoma. Nat Rev Gastroenterol Hepatol 2023;20:349-365. https://doi.org/10.1038/s41575-022-00741-4.

[3] Zhou M, Wang C, Lu S, et al. Tumor-associated macrophages in cholangiocarcinoma: complex interplay and potential therapeutic target. EBioMedicine 2021;67:103375. https://doi.org/10.1016/j.ebiom.2021.103375.

[4] Guan F, Wang R, Yi Z, et al. Tissue macrophages: origin, heterogenity, biological functions, diseases and therapeutic targets. Signal Transduct Target Ther 2025;10:93. https://doi.org/10.1038/s41392-025-02124-y.

[5] Chen S, Saeed A, Liu Q, et al. Macrophages in immunoregulation and therapeutics. Signal Transduct Target Ther 2023;8:207. https://doi.org/10.1038/s41392-023-01452-1.

[6] Fan G, Tao C, Li L, et al. The co-location of MARCO+ tumor-associated macrophages and CTSE+ tumor cells determined the poor prognosis in intrahepatic cholangiocarcinoma. Hepatology 2025;82:25-41. https://doi.org/10.1097/HEP.0000000000001138.

[7] Song G, Shi Y, Meng L, et al. Single-cell transcriptomic analysis suggests two molecularly subtypes of intrahepatic cholangiocarcinoma. Nat Commun 2022;13:1642. https://doi.org/10.1038/s41467-022-29164-0.

[8] Li Z, Li Zhao, Chen Q, et al. Single-Cell and Spatial Transcriptomics Delineate the Microstructure and Immune Landscape of Intrahepatic Cholangiocarcinoma in the Leading-Edge Area. Ad Sci 2025;12:e2412740. https://doi.org/10.1002/advs.202412740.

[9] Yan J, Jiang Z, Zhang S, et al. Spatial☐temporal heterogeneities of liver cancer and the discovery of the invasive zone. Clin Transl Med 2025;15:e70224. https://doi.org/10.1002/ctm2.70224.

[10] Dong LQ, Shi Y, Ma LJ, et al. Spatial and temporal clonal evolution of intrahepatic cholangiocarcinoma. J Hepatol 2018;69:89-98. https://doi.org/10.1016/j.jhep.2018.02.029.

[11] Zhang Q, He Y, Luo N, et al. Landscape and Dynamics of Single Immune Cells in Hepatocellular Carcinoma. Cell 2019;179:829-845.e820. https://doi.org/10.1016/j.cell.2019.10.003.

[12] Su X, Liang C, Chen R, et al. Deciphering tumor microenvironment: CXCL9 and SPP1 as crucial determinants of tumor-associated macrophage polarity and prognostic indicators. Mol Cancer 2024;23:13. https://doi.org/10.1186/s12943-023-01931-7.

[13] Jusakul A, Cutcutache I, Yong CH, et al. Whole-Genome and Epigenomic Landscapes of Etiologically Distinct Subtypes of Cholangiocarcinoma. Cancer Discov 2017;7:1116-1135. https://doi.org/10.1158/2159-8290.CD-17-0368.

[14] Ahn KS, O'Brien D, Kang YN, et al. Prognostic subclass of intrahepatic cholangiocarcinoma by integrative molecular-clinical analysis and potential targeted approach. Hepatol Int 2019;13:490-500. https://doi.org/10.1007/s12072-019-09954-3.

[15] Lubuela G, Beaufrere A, Albuquerque M, et al. Prognostic impact of the tumor microenvironment in intrahepatic cholangiocarcinoma: identification of a peritumoral fibro-immune interface. Virchows Arch 2024;485:901-911. https://doi.org/10.1007/s00428-024-03922-5.

[16] Wu R, Guo W, Qiu X, et al. Comprehensive analysis of spatial architecture in primary liver cancer.
Sci Adv 2021;7:eabg3750. https://doi.org/10.1126/sciadv.abg3750.
[17] Low J, Altman R, Badolian A, et al. Heat-induced phosphatidylserine changes drive HSPA1A's
plasma membrane localization. Cell Stress Chaperones 2025;30.
https://doi.org/10.1016/j.cstres.2025.100092.
[18] Shashikanth N, Alaidi O, Basa L, et al. Role of C-Jun N-Terminal Kinases on a Stressed
Epithelium: Time for Testing Isoform Specificity. Biology 2025;14:649.
https://doi.org/10.3390/biology14060649.
[19] Wolf CL, Feci A, Tuccinardi JP, et al. Development of the First Small-Molecule Inhibitor
Targeting Oncostatin M for Treatment of Breast Cancer. J Med Chem 2025;68:15422-15445.

https://doi.org/10.1021/acs.jmedchem.4c03233

[20] Luo L, Wang S, Hu Y, et al. Precisely Regulating M2 Subtype Macrophages for Renal Fibrosis
Resolution. ACS Nano 2023;17:22508-22526. https://doi.org/10.1021/acsnano.3c05998.
[21] Hasita H, Komohara Y, Okabe H, et al. Significance of alternatively activated macrophages in
patients with intrahepatic cholangiocarcinoma. Cancer Sci 2010;101:1913-1919.
https://doi.org/10.1111/j.1349-7006.2010.01614.x.
[22] Hui Y, Lin Z, Liu Y, et al. Intrahepatic cholangiocarcinoma induced M2-polarized
tumor-associated macrophages facilitate tumor growth and invasiveness. Cancer Cell Int 2020;20:586.
https://doi.org/10.1186/s12935-020-01687-w.
[23] Ma RY, Black A, Qian BZ. Macrophage diversity in cancer revisited in the era of single-cell
omics. Trends Immunol 2022;43:546-563. https://doi.org/10.1016/j.it.2022.04.008.
[24] Okikawa S, Morine Y, Saito Y, et al. Inhibition of the VEGF signaling pathway attenuates
tumor□associated macrophage activity in liver cancer. Oncol Rep 2022;47:71.
https://doi.org/10.3892/or.2022.8282.
[25] Chen M, Ren R, Lin W, et al Exploring the oncostatin M (OSM) feed-forward signaling of
glioblastoma via STAT3 in pan-cancer analysis. Cancer Cell Int 2021;21:565.
https://doi.org/10.1186/s12935-021-02260-9.
[26] Polak KL, Tamagno I, Parameswaran N, et al. Oncostatin-M and OSM-Receptor Feed-Forward
Activation of MAPK Induces Separable Stem-like and Mesenchymal Programs. Mol Cancer Res
2023;21:975-990. https://doi.org/10.1158/1541-7786.MCR-22-0715.
[27] Omatsu M, Nakanishi Y, Iwane K, et al. THBS1-producing tumor-infiltrating monocyte-like cells
contribute to immunosuppression and metastasis in colorectal cancer. Nat Commun 2023;14:5534.
https://doi.org/10.1038/s41467-023-41095-y.
[28] You Y, Tian Z, Du Z, et al. M1-like tumor-associated macrophages cascade a
mesenchymal/stem-like phenotype of oral squamous cell carcinoma via the IL6/Stat3/THBS1 feedback
loop. J Exp Clin Cancer Res 2022;41:10. https://doi.org/10.1186/s13046-021-02222-z.
[29] Wolf CL, Pruett C, Lighter D, et al. The clinical relevance of OSM in inflammatory diseases: a
comprehensive review. Front Immunol 2023;14:1239732.
https://doi.org/10.3389/fimmu.2023.1239732.
[30] Aziz A, Soucie E, Sarrazin S, et al. MafB/c-Maf deficiency enables self-renewal of differentiated
functional macrophages. Science 2009;326:867-871. https://doi.org/10.1126/science.1176056.
[31] Kikuchi K, Iida M, Ikeda N, et al. Macrophages Switch Their Phenotype by Regulating Maf
Expression during Different Phases of Inflammation. J Immunol 2018;201:635-651.

https://doi.org/10.4049/jimmunol.1800040.

 Figure legends

 Fig. 1. Spatiotemporal heterogeneity of macrophages in mouse and human ICC tissues.(A) (i)

 UMAP (i) and stacked bar (ii) visualisation of ICC patient TAM subclusters profile (datasets

 GSE138709 and GSE189903). Malignant (Mal), boundary (Bdy), adjacent non-malignant (nMal).

 (B) UMAP(i), scatter plot of M2 polarisation scores (ii) and the temporal distribution (iii) of TAM

 subclusters in AKT1/NICD1 mice. WT (healthy, N=2), D7 (early stage, N=2), D15 (middle stage,

 N=3), D25 (middle to advanved stage, N=2), D31 (advanced stage, N=3). (C) Bubble plots

 displaying the expression of TAM marker genes in patients with ICC and the AKT1/NICD1 mice.

 Color indicates expression levels, bubble size reflects cell proportion. (D) Temporal dynamics of

 TAM subcluster scores (Mφ_S100a and Mφ_SPP1), red denotes mean expression above the 90th

 percentile. (E) Scatter–pie plots illustrating prognostic associations of distinct TAM subclusters.

 Pie chart illustrating cell prognosis outcomes in each TAM subtype. Blue, survival. Red, death.

 Grey, no significant association.(F) (i)Subcutaneous co-injection tumor model experiments were

 performed to validate the role of macrophage infiltration in ICC progression. BR group：injection

 of BR primary cells alone. BR+BMDM group: co-injection of BR primary cells with BMDM.

 From left to right, representative photographs of xenograft tumors, tumor volume growth curves

 (in units of 1,000 mm³), and tumor weights are shown. (ii) In the AKT1/NICD1-driven ICC mouse

 model, macrophage depletion was achieved via clodronate liposome injection to investigate its

 impact on ICC progression. The control group received neutral control liposomes. Left:

 representative liver images . Right: Macrophage infiltration levels (R1, CD11b□F4/80□ cells

 within CD45□ population; R2, F4/80□ cells within CD11b□ population) analyzed via flow

 cytometry and correlated with liver-to-body weight ratio using scatter plots. Correlation analysis

was performed using Spearman's rank correlation test. (G) (i) 10X Visium spatial transcriptomics

of ICC patient tissue identified seven cell populations.(ii) Scatter plot shows functional regions

predicted by Cottrazm (Mal = malignant, Bdy = boundary, nMal = adjacent non□malignant).(iii)

Stacked bar plots display cell-type proportions across spatial clusters from xCell analysis.(iv)

Heatmap shows scaled median cell-type compositions per niche; asterisks denote enrichment in a

niche (one-sided Wilcoxon rank□sum test, adj. $p < 0.05$). (H) (i)Spatial transcriptomics identified

12 clusters in the BMKMANU S1000 AKT1/NICD1 ICC tissue. Heatmap showing the

CK7$^+$CK19$^+$ ICC cells (ii) and Mφ_Spp1 (iii) using the RCTD cell type definition. (iv) Contour

maps highlighting spatial expression of mean expression of Mφ_Spp1 signature scores. (I) (i)

Spatial domains defined by BANKSY in ICC patient dataset using 10X HD platform. (ii) mIF

illustrated the spatial distribution of CD68$^+$ cells and their distances to CK19$^+$ cells. (iii) Cell-type

composition across the spatial domains. (iv) Spatial proximity between Mφ_SPP1 and

CK7$^+$CK19$^+$ cells. (v) Median importance of cell-type abundance in the prediction of abundances

of other cell types in the local niche view. (J) (i) Bar plots showing the proportion of SPP1$^+$CD68$^+$

cells at different distance bins from CK19$^+$ cells in two validation datasets. (ii) Heatmap of tissue

spatial infiltration patterns for CK19$^+$ cells (highlighted in yellow) in a representative sample. (iii)

Zoomed-in view demonstrating dense infiltration of SPP1$^+$CD68$^+$ macrophages surrounding

CK19$^+$ cells.

Fig. 2. Mφ_SPP1 promotes ICC proliferation via OSM-OSMR. (A) UMAP plot (left) and

bubble plot (right) illustrating the distribution of epithelial cells in AKT1/NICD1 mice and the cell

type features. (B) UMAP plots showing the ICC patients epithelial subtypes. (C) Venn diagram

showing overlap of ligand genes that are significantly upregulated in Mφ_SPP1 which exhibit

strong interactions with tumor cells as identified by three cell communication methods

(CellphoneDB, CpDB; CellChat, and Niche, NicheNet). (D) Heatmap showing the interaction

intensity between TAMs and ICC cells in AKT1/NICD1-mediated *OSM* signalling. (E) Bubble

chart visualising pathway enrichment results for downstream target genes of *OSM* in E_Jun

(AKT1/NICD1 mouse) and E_CD74 (Human) via NicheNet calculated by clusterprofiler.

Schematic of in vivo experimental design (F-i). Representative images (F-ii) of livers from Lifr$^{fl/fl}$

and Osmr$^{fl/fl}$ control mice and Lifr$^{fl/fl}$;Alb-Cre and Osmr$^{fl/fl}$;Alb-Cre mice injected with

AKT1/NICD1 and their liver-to-body weight ratios (G). Data are presented as mean ± SD.

Statistical significance was analyzed by one-way ANOVA ($p < 0.05$) followed by Dunnett's test

(ns, not significant, **$p < 0.01$, ***$p < 0.001$; ****$p < 0.0001$). (H) Multiplex

immunofluorescence showing spatial co-localisation between CD68□OSM□ TAMs and

OSMR□KRT19□ ICC cells in ICC patients. Scale bar, 200 μm.

Fig. 3. Mφ_SPP1 promotes ICC proliferation by upregulating p-STAT3 and CCND1 in ICC

cells. (A) Bar plot showing gene expression levels in co-culture. Statistical significance was

analyzed by one-way ANOVA ($p < 0.05$) followed by Dunnett's test (ns, not significant, **$p <$

0.01, ***$p < 0.001$; ****$p < 0.0001$). R, Raw264.7. Q, QBC939. (B) CCK-8 assay measuring cell

viability of HuCCT1 and QBC939 cells treated with 2.5 ng/mL or 5.0 ng/mL OSM, or left

untreated, for 5 consecutive days. (C) (i) EdU incorporation assay in QBC939 cells transduced

with control vector or OSMR-targeting sgRNAs (sgOSMR-1, sgOSMR-2). Scale bar, 200 μm. (ii)

CCK-8 assay measuring viability of HuCCT1 and QBC939 cells transduced with control vector or

sgOSMR-1 at days 1, 3, and 5 after seeding. (iii) Colony formation assay with crystal violet

staining of HuCCT1 and QBC939 cells transduced with control vector or sgOSMR-1. (D) Bar plot

showing KEGG pathway enrichment among phosphorylated proteins after OSM treatment in

QBC939 cells. Blue, downregulation; Red, upregulation. (E) Network showing OSMR related

pathways. H&E staining and IHC staining for OSMR, OSM, CK19, and Ki-67 in

paraffin-embedded liver sections(F) and western blot analysis of OSMR, CCND1, BCL2,

phosphorylated STAT3 (p-STAT3), phosphorylated mTORC1 (p-mTORC1), and OSM protein

levels , GAPDH served as a loading control (G) from $Osmr^{fl/fl}$ and $Osmr^{fl/fl}$; Alb-Cre mice injected

with AKT1/NICD1. Scale bar, 200 μm. Data are presented as mean ± SD. Significance was

determined by two-tailed unpaired t-test (***$p < 0.001$) for B and C.

Fig. 4. Mφ_SPP1 promotes ICC invasion via *THBS1-CD47* signaling. (A) (i) Heatmap of

interactions between TAMs and tumor subtypes in THBS1 signaling. Sender,ligand source.

Receiver, receptor. Color intensity indicates importance. (ii) Top, bar plot showing receive

contributions of ligand THBS1. Bottom row shows *Cd47* expression across AKT1/NICD1

epithelial subclusters. (iii) Spatial patterns of *THBS1-CD47* interaction. (B) Bubble chart

visualising pathway enrichment for downstream target genes of *THBS1* in E_Jun (AKT1/NICD1

mouse) and E_IGFBP7 (Human) via NicheNet. Bubble size depicts enrichment score and colour

showcasing significance (-log10(p.adj), clusterProfiler). (C) Representative scratch□wound

images of HuCCT1 and RBE cells with or without 10 ng/mL THBS1 or control. HuCCT1 at 0, 12

and 24 h; RBE at 0, 36 and 72 h. Scale bar, 150 μm. (D) Representative crystal violet–stained

Transwell migration images of primary BR cells and HuCCT1 cells treated with 10 ng/mL THBS1

or control. Scale bar, 150 μm. (E) qRT-PCR analysis of *CD47* levels in HuCCT1 cells transfected

with siScramble control, siCD47-1, or siCD47-2 for 48 h. Data are presented as mean ± SD.

Statistical significance was analyzed by one-way ANOVA ($p < 0.05$) followed by Dunnett's test

(****$p < 0.0001$).(F) Representative crystal violet–stained Transwell migration images of

HuCCT1 cells transfected with siScramble or siCD47-2. Scale bar, 150 μm. (G) qRT-PCR (i) and

western blot (ii) analyses of CD47 transcript and protein levels, respectively, in BR cells

transduced with LV3-shNC control lentivirus or LV3-shCD47. (H) EdU incorporation assay

assessing proliferation of BR cells transduced with LV3-shNC or LV3-shCD47 lentivirus. Scale

bar, 50 μm. (I) Colony formation assay comparing BR cells transduced with LV3-shNC or

LV3-shCd47 lentivirus. (J) CCK-8 assay measuring cell growth of BR cells transduced with

LV3-shNC or LV3-shCd47 lentivirus from day 1 to day 5 after seeding. (K) Representative crystal

violet–stained Transwell migration images and corresponding chamber images of BR cells

transduced with LV3-shNC or LV3-shCd47 co-cultured with BMDMs or cultured with complete

medium for 24 h. Scale bar, 150 μm. (L) Subcutaneous allograft ICC growth curves and

representative ICC tissue images (i) of BR cells transduced with LV3-shNC or LV3-shCd47

lentivirus (ii). (M) Representative images of mouse abdominal phenotypes at tumor endpoint after

intraperitoneal injection of BR cells transduced with LV3-shNC or LV3-shCd47; tumor sites

indicated by white arrows; quantification of metastatic nodules and volume of ascites collected.

Data are presented as mean ± SD. Significance was determined by two-tailed unpaired t-test (***p

< 0.001) for E, G, J and L.

Fig. 5. Mφ_SPP1 promotes ICC invasion via mTOR-mediated phosphorylation of DBN1. (A)

Western blotting showing the expression of p-mTOR, p-RPS6, EpCAM, N-cadherin, Vimentin,

and the internal control GAPDH in HuCCT1 cells after THBS1 protein stimulation. (B) Bar plot

displaying pathway enrichment analysis for HuCCT1 at 6 h and 2 h following the addition of

exogenous THBS1 protein. (C) Circos plot displaying differentially phosphorylated sites at

multiple time points after adding exogenous THBS1. From inner to outer rings:

phosphorylation-site proportions, phosphopeptide expression intensity (inner heatmap), sample

groups, and enriched signaling pathways; the outermost ring lists gene symbols and phosphosites

unique to each treatment. Colored links show shared phosphosites-gray, common to control and

treatments. Brown, mTOR-related. Red, fibrosis-related. Green, mesenchymal-related. Large and

small ticks denote 50 and 5 phosphopeptides, respectively. (D) Western blot analysis of DBN1,

phosphorylated DBN1 (p-DBN1), and Vimentin protein levels in BR cells cultured alone or

co-cultured with BMDMs for 24 h or 48 h; GAPDH served as a loading control. (E)

Representative crystal violet–stained images from Transwell migration assays of HuCCT1 and BR

cells co-cultured with BMDMs, with or without Rapamycin treatment. N = 5. Scale bar, 200 μm.

(F) Representative crystal violet–stained images from Transwell migration assays of HuCCT1 and

BR cells treated with THBS1, with or without Rapamycin blockade; groups include control,

THBS1 treatment, and THBS1 combined with Rapamycin. N = 5. Scale bar, 200 μm. (G)

Representative chamber and crystal violet–stained images of Transwell migration assays for

HuCCT1 cells (i) treated with 0, 5, 25, and 125 μM BTP2 inhibitor, and BR cells (ii) treated with

0, 25, and 125 μM BTP2 in the lower chamber. Scale bar, 200 μm. (H) Representative images of

metastatic ICC lesions from various organs in Thbs1$^{fl/fl}$;Lyz-Cre and Thbs1$^{fl/fl}$ mice injected with

sgP19/KRAS. Scale bars: 1 cm for liver; 5 mm for lung, spleen, kidney, peritoneum, and colon. (I).

H&E, CK19, and Ki67 staining of metastatic tumor nests in sections from different organs in

Thbs1$^{fl/fl}$;Lyz-Cre and Thbs1$^{fl/fl}$ mice injected with sgP19/KRAS. This lymph node is referred to

the largest lymph node adjacent to the spleen. Scale bar, 200 μm. (J) Western blot analysis of

DBN1, CD47, p-mTORC1(Ser2448), and Vimentin protein expression in tumor tissues from

Thbs1$^{fl/fl}$;Lyz-Cre and Thbs1$^{fl/fl}$ mice injected with sgP19/KRAS; GAPDH served as loading

control. (K) Representative images of liver phenotypes and quantification of liver-to-body weight

ratios in Thbs1$^{fl/fl}$;Lyz-Cre and Thbs1$^{fl/fl}$ mice injected with AKT1/NICD1. Data are presented as

mean ± SD. Significance was determined by two-tailed unpaired t-test (*$p < 0.05$), N = 6. (L)

H&E staining of liver sections showing whole-slide and magnified tumor areas in

Thbs1$^{fl/fl}$;Lyz-Cre and Thbs1$^{fl/fl}$ mice injected with AKT1/NICD1. Scale bar, 100 μm. (M) Western

blot analysis of THBS1, CD47, DBN1, p-DBN1(ser142), p-mTORC1(Ser2448), and Vimentin

protein expression in tumor tissues from Thbs1$^{fl/fl}$;Lyz-Cre and Thbs1$^{fl/fl}$ mice injected with

AKT1/NICD1; GAPDH served as loading control. AN, AKT1/NICD1.

Fig. 6. Combined intervention with OSM inhibitor and THBS1 knockout effectively delays

ICC progression. (A) Stacked bar plot illustrating the prevalence of OSM☐THBS1☐

dual-positive patients within distinct ICC cohorts. (B) High (red) and low (green) groups were

defined by the mean expression of OSM and THBS1 respectively; survival in three datasets was

compared using the log☐rank test. (C) Representative liver phenotype images from four

experimental groups in the *THBS1/OSM* combination blockade experiment: (1) Thbs1$^{fl/fl}$ +

vehicle control; (2) Thbs1$^{fl/fl}$ + OSM-SMI-10B; (3) Thbs1$^{fl/fl}$; Lyz2-Cre + vehicle control; (4)

Thbs1$^{fl/fl}$; Lyz2-Cre + OSM-SMI-10B. (C) Statistical analysis of liver-to-body weight ratio across

different treatment groups in the *THBS1/OSM* combination blockade experiment. Data are

presented as mean ± SD. Statistical significance was analysed by one-way ANOVA ($p < 0.05$)

followed by Dunnett's test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; ****$p < 0.0001$). (D) H&E and

IHC staining of liver sections from the *THBS1/OSM* combination blockade experiment. Ki67 and

Ck19 positively identifies ICC tumor nest regions. Scale bar, 200 μm. (F) (i) Proximity of

OSM$^+$THBS1$^+$CD68$^+$SPP1$^+$ cells to CK19$^+$ cells by mIF. yellow, CK19$^+$; orange□red/green/blue,

0–20, 20–40, 40–60 μm from CK19$^+$. (ii) mIF shows OSM$^+$THBS1$^+$CD68$^+$SPP1$^+$ macrophages

around CK19$^+$ cells. (iii) Spearman correlation between OSM/THBS1 and CD68. (iv) Patients

stratified by OSM/THBS1 intensity (red = high, green = low); survival compared by log□rank

test.

Fig. 7. Epigenetic regulatory profile of ICC TAM subtypes and key regulators in Mφ_SPP1.

(A) (i) UMAP of major subtypes identified by scATAC-seq in AKT1/NICD1. (ii) Chromatin

accessibility of representative marker genes for each cell type. (B) Volcano plot showing

differences in motif deviation scores (chromVAR) between Mφ_S100a and Mφ_SPP1

macrophage subclusters. X-axis: mean motif difference; Y-axis: -log10(FDR). Red dots mark the

top 10 transcription factors with the largest accessibility changes (mean difference > 0.05, FDR <

0.05; Wilcoxon test in ArchR). (C) (i) Venn diagram showing transcription factors with the most

distinct motifs between the two subclusters and their overlap with factors expressed >1. (ii)

UMAP scatter plots depicting chromatin accessibility distribution for representative differential

motifs. (D) Bubble plots depicting pathway enrichment of putative target genes regulated by

transcription factors *Klf3* in Mφ_S100a(i) and *Mafb* in Mφ_SPP1(ii). (E) Spatial expression

patterns of the key transcription factor *MAFB* in AKT1/NICD1 mice and human tissues. (F)

qRT-PCR analysis of *Tgfbr2*, *Mafb*, *Thbs1*, and *Osm* after BMDM polarising with TCM for 2, 6,

12, 24, 36, and 48 hours. (G) qRT-PCR evaluating *Tgfbr2, Mafb, Thbs1*, and *Osm* transcript levels

under different culture conditions. (H) qRT-PCR analysis of *Arg1, Vegfa, Osm*, and *Thbs1* in

siScramble non-polarised, siScramble TCM polarised, and siMafb-1 or siMafb-2 TCM treated

groups. Data are presented as mean ± SD. Statistical significance was analysed by one-way

ANOVA ($p < 0.05$) followed by Dunnett's t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; ****$p <$

0.0001) for F, G and H.

A AKT1/NICD1 mouse epithelial subtypes

Mean expression in group

Fraction of cells in group (%)

B Human epithelial subtypes

Adjacent   ICC

Cell Types
- E_ALB
- E_CD74
- E_FXYD2
- E_IGFBP7
- E_MKI67
- E_PROM1
- E_S100P

C Cell communication between Mφ_VEGFA and ICC subtypes

Intersection Size

178 165 59 28 19 16 16 14 10 7 6 5 4 4 4 3 2 2 2 1 1 1 1 1 1

OSM THBS1

Set size

Human
CePD
Cellchat
Mouse
Niche

250   0

D OSM interaction signaling network

Sources (Sender)

Communication Prob.
0.04
0.03
0.02
0.01
0

E Pathways enrichment of OSM target genes in ICC subtypes

Mouse   Human
E_Jun   E_CD74

- Tube development
- Response to cytokine
- Regulation of MAPK cascade
- Regulation of cell cycle
- Protein phosphorylation
- Epithelial cell proliferation

EnrichmentScore

Enrichment Score
- 9
- 12
- 15

$-\log_{10}(P.adj)$
6
5
4

F

Lifr^{fl/fl}   Osmr^{fl/fl}

Lifr^{fl/fl};Alb-Cre   Osmr^{fl/fl};Alb-Cre

→ Hydrodynamic injection of AKT1/NICD1 → Sacrifice mice 4 weeks post-injection or moribund

AKT1/NICD1   AKT1/NICD1

Osmr^{fl/fl}   Osmr^{fl/fl};Alb-Cre   Lifr^{fl/fl}   Lifr^{fl/fl};Alb-cre

G

Hepatosomatic index

ns   **

- Lifr^{fl/fl}
- Lifr^{fl/fl};Alb-Cre
- Osmr^{fl/fl}
- Osmr^{fl/fl};Alb-Cre

H

CK19   CD68

CK19 OSMR CD68 OSM   CK19 OSMR   CD68 OSM