Convergent evolution of aerobic fermentation through divergent mechanisms acting on

key shared glycolytic genes

Linda C. Horianopoulos[1,2]*, Antonis Rokas[3], Chris Todd Hittinger[1]*

[1] Laboratory of Genetics, J. F. Crow Institute for the Study of Evolution, Center for Genomic

Science Innovation, DOE Great Lakes Bioenergy Research Center, Wisconsin Energy Institute,

University of Wisconsin-Madison, Madison, WI 53726, USA

[2] Department of Food Science, University of Guelph, Guelph, ON, Canada, N1G 2W1

[3] Department of Biological Sciences, Vanderbilt University, Nashville, TN 37235, USA;

Evolutionary Studies Initiative, Vanderbilt University, Nashville, TN 37235, USA

*corresponding authors: lhoriano@uoguelph.ca and cthittinger@wisc.edu

Keywords: transcriptional rewiring, cis-regulatory evolution, gene duplication, aerobic

fermentation, glycolysis, Crabtree/Warburg Effect

ABSTRACT

As the tree of life becomes increasingly accessible to molecular investigations, describing mechanisms underlying evolutionary convergence and constraint will be crucial to understanding diversification. The lineage including the model yeast *Saccharomyces cerevisiae* evolved aerobic fermentation in part through an ancient whole genome duplication and retention of glycolytic genes. To evaluate the glycolytic rates across diverse yeasts, we developed and deployed an underline{extra}underline{c}ellular underline{a}cidification underline{r}ates (ECAR) assay on 299 species that span more than 400 million years of evolution and identified a clade in the genus *Saturnispora* that convergently evolved aerobic fermentation. Through comparative genomics and transcriptomics, we found that several glycolytic genes had higher expression and novel cis-regulatory elements in aerobically fermenting *Saturnispora* species. When the transcription factor required for their activation was deleted in *Saturnispora dispora,* the mutants had reduced glycolytic rates and increased respiration. Intriguingly, many of the upregulated genes are orthologous to duplicated glycolytic genes in *S. cerevisiae.* These divergent genetic mechanisms suggest that there are strong evolutionary constraints on how some traits like aerobic fermentation can arise convergently.

INTRODUCTION

The convergent evolution of complex traits provides insights into the mechanisms and constraints that govern their repeated emergence. Convergence can arise through distinct or common genetic mechanisms. One of the most compelling examples of convergent evolution in a complex metabolic trait is the evolution of C4 photosynthesis in more than 60 independent lineages (Sage *et al*, 2011). C4 plants have repeatedly and independently evolved mechanisms to increase photosynthesis efficiency by partitioning biochemical reactions and allowing the key photosynthesis enzyme, RuBisCO, to be in a high CO_2 environment (Brown *et al*, 2011; Williams *et al*, 2012; Lyu *et al*, 2025). Comparisons across C4 grasses have identified multiple underlying

evolutionary mechanisms, including gene family expansions of key enzymes, parallel amino

acid changes, and similar cis-regulatory elements in the promoters of C4 enzymes (Lyu *et al*,

2024). For example, a specific comparison between rice (C3) and sorghum (C4) revealed that

the promoters driving the C4 enzymes in sorghum co-opted an ancestral cis-regulatory element

for cell type specificity, thereby promoting this physical partitioning of biochemical reactions

(Swift *et al*, 2024). These innovations highlight how improved metabolic efficiency has been

rediscovered in many independent lineages through different mechanisms altering the

abundance, expression patterns, or activity of the same key C4 enzymes.

Within the yeasts of the subphylum Saccharomycotina, the Crabtree/Warburg Effect, in

which yeasts ferment even in the presence of oxygen, is a complex metabolic trait that has

evolved in at least two independent lineages, in *Saccharomyces* and relatives and some

Brettanomyces species (Rozpedowska *et al*, 2011). Comparative phenotyping of up to forty

species demonstrated that the Crabtree-positive phenotype likely evolved in a stepwise fashion

and that overflow metabolism, wherein cells maximize energy production rates through

glycolysis but produce fermentation products, is the fundamental mechanism driving aerobic

fermentation (Hagman & Piškur, 2015; Hagman *et al*, 2013). Some yeasts, including *S.*

cerevisiae, have undergone dramatic genetic changes, including preferential retention of

duplicated glycolytic genes from an ancient whole genome duplication (WGD), further

increasing their fermentative capacity (Conant & Wolfe, 2007; Escalera-Fanjul *et al*, 2019).

Brettanomyces species, which are also Crabtree-positive, do not have the high glucose

consumption rates of *S. cerevisiae*, but they do repress respiration when glucose is present

(Hagman *et al*, 2013; Rozpedowska *et al*, 2011). These independent occurrences of aerobic

fermentation suggest that there may be additional lineages of yeasts with independently evolved

preferences for fermentation.

Complex traits, such as the Crabtree/Warburg Effect, are difficult to assess using high

throughput assays. However, since a high glycolytic rate and overflow metabolism are thought

to precede the evolution of the Crabtree/Warburg Effect, we developed an assay to evaluate the

glycolytic rate of diverse species based on glucose-induced extracellular acidification rates

(ECAR). ECAR has been used to study mammalian cells, primarily cancer cells, which display

the Crabtree/Warburg Effect wherein they accumulate lactate as a byproduct of rapid aerobic

glycolysis (Schmidt *et al*, 2021; Zhang & Zhang, 2019). Acidification measurements have also

been applied in *S. cerevisiae* to assess fermentative capacity in industrial settings (Sigler, 2013;

M & K, 1982; Yamashoji et al, 2020; Sigler et al, 1981), and more recently, ECAR was

measured with the Seahorse Extracellular Flux analyzer to assess glycolytic rate in pathogenic

yeasts (Tucey *et al*, 2018; Yoo *et al*, 2024; Zhang *et al*, 2018). Assessing glycolytic rates across

a wide range of species could potentially uncover novel instances of high glycolytic rates and

provide insight into the genetic mechanisms underlying high overflow metabolism. Identification

of species with high glycolytic rates may also have important implications as increasing efforts

are being made to use non-conventional yeasts for biotechnological applications (Geijer *et al*,

2022; Wang *et al*, 2025; Radecka *et al*, 2015; Pyne *et al*, 2023; Patra *et al*, 2021).

Herein, we developed a microtiter plate-based assay to screen diverse yeast species for

ECAR. We confirmed that the species in the order Saccharomycetales, including the model

yeast *S. cerevisiae*, display rapid ECAR phenotypes, which are consistent with their rapid

glycolytic rates. However, we also identify multiple species in the genus *Saturnispora*, including

Saturnispora dispora, that have rapid ECAR compared to other species in the order Pichiales.

This order already contains two yeasts of considerable biotechnological interest: *Komagataella*

phaffii (previously known as *Pichia pastoris*), which is a leading chassis for protein production

(Heistinger *et al*, 2020; Gasser & Mattanovich, 2018), and *Pichia kudriavzevii* (also known as

Issatchenkia orientails, *Candida glycerinogenes*, and *Candida krusei*), which is emerging as an

important microbial cell factory for the production of organic acids (Tan *et al*, 2025; Suthers &

Maranas, 2022). Furthermore, additional species within the genus *Pichia* are actively being

developed for organic acid production (Pyne *et al*, 2023). This emerging interest in the industrial

applications of Pichiales warrants further work to understand the metabolic capabilities of this

order to guide their development as microbial cell factories. Moreover, characterizing the genetic

mechanism driving the strong fermentative capacity in *Saturnispora* can reveal the evolutionary

constraints (if any) on rapid fermentation and guide rational strategies to maximize rates of

organic acid production in Pichiales.

To understand how the unusually high glycolytic rate of *Sat. dispora* and its relatives

evolved, we used comparative genomics and transcriptomics and found that several genes

encoding glycolytic enzymes had increased expression at the transcript level in species with a

rapid ECAR phenotype. Intriguingly, genes orthologous to those whose duplications are key to

the high glycolytic rate of *S. cerevisiae* also had higher expression in the rapid ECAR

Saturnispora, which prompted us to investigate the promoter sequences of glycolytic genes for

binding sites of transcription factors (TFs) known to regulate carbon metabolism. We found that

there were more predicted Gal4p-binding sites ($CGGN_{11}CCG$) in promoters of some glycolytic

genes in the rapid ECAR species. Although Gal4p is named for its role in galactose metabolism

in *S. cerevisiae* (Johnston, 1987)*,* this role is evolutionarily derived and specific to

Saccharomycetales (Harrison *et al*, 2022). Gal4p homologs play important roles regulating

carbon metabolism in *Candida albicans* (Askew *et al*, 2009) and glycolysis in *K. phaffii*, although

these species do not natively exhibit aerobic fermentation (Ata *et al*, 2018). Using novel genetic

tools that we developed for *Sat. dispora*, we showed that *GAL4* was necessary for the induction

of the most highly expressed glycolytic gene, *TDH*. Altogether, we found an independently

evolved group of yeasts with a high glycolytic rate phenotype achieved through the recruitment

of cis-regulatory elements driving the direct transcriptional upregulation of orthologs of the same

glycolytic genes that retain duplicates in *S. cerevisiae*. This convergence suggests that the

evolution of increased glycolytic rate is constrained and requires high activity of key glycolytic

enzymes, which can be achieved by distinct molecular mechanisms.

RESULTS

Rapid ECAR in *Saccharomyces* and relatives and in the distantly related genus *Saturnispora*

We developed a high-throughput method to measure the glucose-dependent ECAR of yeast species as a proxy for glycolytic rate using a 96-well plate and a standard microplate reader (Fig. S1). We plotted the mean of three biological replicates per species as a colour gradient across the yeast phylogeny to identify clades with similar ECAR profiles. Accordingly, we found that many species in the order Saccharomycetales, particularly those yeasts that underwent an ancestral whole genome duplication (WGD), such as *S. cerevisiae*, had rapid acidification (Fig. 1A). Although this result was expected and validated the ability of our method to capture glycolytic rate, we were particularly interested in other clades that showed a rapid ECAR phenotype. For example, we were surprised to find a group in the order Pichiales in the genus *Saturnispora* that showed a rapid ECAR phenotype (as low as -0.032 pH/hr in *Sat. dispora*). These species have not been thoroughly characterized but are in the same order as multiple biotechnologically relevant species, including the Crabtree-negative species *K. pastoris* (low ECAR=0.022 pH/hr). The growing biotechnological interest in Pichiales yeasts prompted our specific interest in the metabolism of these species and in the evolution of the rapid ECAR trait in some *Saturnispora* species.

High glycolytic rate correlates with gene duplication in *S. cerevisiae* and relatives

The high glycolytic rate in *S. cerevisiae* is hypothesized to be, at least in part, due to an ancestral WGD and subsequent preferential retention of glycolytic genes (Conant & Wolfe, 2007; Hagman *et al*, 2013; Conant & Wolfe, 2008). To further validate the results of our screen and test the generality of the hypothesis that having a higher number of glycolytic genes is associated with rapid ECAR across yeasts, we compared the number of glycolytic genes and the measured ECAR from each yeast species in our dataset. We found a significant correlation

between the total number of glycolytic genes and ECAR across all yeast species after correcting

for phylogeny (Fig. 1B). To interrogate this pattern further, we examined the correlations

between the size of the gene families encoding each enzymatic step in glycolysis and ECAR.

Again, we found strong correlations when looking at all species (Fig. 1C). However, when we

focused on the taxonomic orders containing rapid ECAR species, we noticed that several of

these correlations held true in the order Saccharomycetales but were not significant within the

order Pichiales (Fig. 1C). When the correlations of individual genes were corrected for

phylogeny, we found that the significance values of their correlations were lost in the

Saccharomycetales, likely due to the limited power. Thus, the correlation between ECAR and

glycolytic gene family size may be driven mainly by the well-known WGD in the

Saccharomycetales (Fig. S2). Altogether, these results suggest that the *Saturnispora* species

with the rapid ECAR phenotype in the order Pichiales likely evolved independently from the

duplication of glycolytic genes observed in *S. cerevisiae*.

Figure 1. Variation of extracellular acidification rates (ECAR) across the yeast subphylum is associated with glycolytic genes. (A) The mean of the measured ECAR for each species is visualized as a circle around the yeast phylogeny with the lighter green blocks corresponding to more rapid ECAR. The model yeast *S. cerevisiae*, the whole genome duplication (WGD) in the Saccharomycetales (arrow), the genus *Saturnispora* (box), and the genus *Brettanomyces* (dashed box) are highlighted on the phylogeny. (B) A scatter plot showing the correlation between ECAR and the total number of glycolytic genes where each dot represents a single species. The correlation and p values are based on a phylogenetic generalized least squares model, which accounts for correlation stemming from shared phylogenetic history. (C) Pearson correlations between the number of copies of each glycolytic gene and the measured ECAR. The gene names in boxes have retained duplicates in *S. cerevisiae*; solid boxes indicate duplicates retained in all post-WGD yeasts, whereas the dashed boxes indicate duplicates retained in some but not all post-WGD yeasts (Conant & Wolfe, 2007). Significance of the Pearson correlation is indicated as * $p<0.05$, **$p<0.01$, ***$p<0.005$. The yeast phylogeny was pruned from Shen et al. (Shen *et al*, 2018).

Divergent phenotypes of rapid and low ECAR species in *Saturnispora*

To interrogate the Pichiales-specific rapid ECAR phenotype, we sought to identify closely related species with diverging phenotypes. Within the genus *Saturnispora*, we found that several species consistently showed a rapid ECAR phenotype, namely *Sat. dispora, Sat. hagleri, Sat. serradocipensis, Sat. zaruensis,* and *Sat. saitoi.* Conversely, *Sat. silvae* and *Sat. mendoncae* consistently displayed a low ECAR phenotype (Fig. 2A). Moving forward, we focused on four species, two rapid ECAR and the two low ECAR with the hypothesis that the rapid ECAR species would have metabolic phenotypes more similar to known Crabtree-positive yeasts. To confirm that the acidification reflected rapid glucose consumption, we measured glucose depletion in the media in a typical growth experiment. The rapid ECAR species consumed glucose more rapidly than the low ECAR species (Fig. 2B). The rapid ECAR species even produced ethanol and acetate from overflow metabolism, despite being grown under highly aerobic conditions in baffled flasks (Fig 2B). Accordingly, the rapid ECAR species also grew to a lower final OD compared to the low ECAR species (Fig. 2C). These divergent metabolic phenotypes show that these rapid ECAR yeasts are highly fermentative and indeed exhibit the Crabtree/Warburg Effect under aerobic conditions, which prompted us to interrogate potential genetic differences between the rapid ECAR and low ECAR species.

Figure 2. Species within the genus *Saturnispora* display divergent metabolic phenotypes.
(A) ECAR measurements identified species with consistently low or consistently rapid ECAR
within the genus *Saturnispora*. ECAR was also measured for *Martiniozyma abietophila* as a
closely related outgroup. The (B) extracellular metabolites in the supernatant and (C) growth of
two rapid ECAR and two low ECAR species were measured during growth in a 250-ml baffled
flask, which was highly aerobic. The error bars represent the standard deviation of four
biological replicates.

Transcriptional profiling of rapid and low ECAR species

Since there were no duplications in glycolytic genes in the rapid ECAR *Saturnispora*

species (Fig1C, Table S1), we used comparative transcriptomics to evaluate differences in gene

expression between the rapid and low ECAR species. To compare across species, we mapped

the reads to genes in each species, generated counts, summed counts across genes in the

same orthogroup, and calculated transcripts per million (TPM) for each orthogroup. Since our

primary interest was the glycolytic rate trait, we specifically retrieved the orthogroups encoding

central carbon metabolism genes. These data revealed that the rapid ECAR species had

significantly higher expression of several glycolytic genes, especially genes encoding

hexokinases and the enzymes of lower glycolysis (*TDH*, *GPM*, *ENO*, *PYK*) (Fig. 3). These

transcriptional differences are consistent with the rapid glucose consumption and aerobic

fermentation phenotype. Accordingly, we also found that the low ECAR species had higher

expression of genes encoding citric acid cycle enzymes, further supporting the conclusion that

the metabolisms of these closely related yeasts are highly divergent and favour aerobic

respiration (Fig. S3).

Figure 3. Divergent expression patterns of glycolytic genes between rapid ECAR and low ECAR *Saturnispora* species. The gene expression for orthologous glycolytic genes across two rapid ECAR (*Sat. dispora* = Sdis, *Sat. hagleri* = Shag) and two low ECAR (*Sat. mendoncae* = Smen, *Sat. silvae* = Ssil) are shown as transcripts per million (TPM). The bar height represents the average of four biological replicates, each replicate is shown as a black dot, and the error bars represent the standard error of the mean. The significance is relative to the TPM in *Sat. dispora* (ns = not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, **** $p < 0.001$). In the schematic of glycolysis, carbon backbones are white, and phosphates are yellow. Boxes around gene names are orthologous to genes that have retained duplicates in *S. cerevisiae* as in Fig. 1C, solid boxes indicate duplicates retained in all post-WGD yeasts, whereas the dashed boxes indicate duplicates retained in some but not all post-WGD yeasts (Conant & Wolfe, 2007).

The Gal4p transcription factor contributes to rapid glycolytic rate in rapid ECAR

Saturnispora

Since we observed such a striking difference in the expression of glycolytic genes

between rapid and low ECAR yeasts, we hypothesized that this difference was due to genetic

differences in the regulatory regions of their glycolytic genes. To test this, we retrieved the

nucleotide sequences 1 kb upstream of each glycolytic gene as the putative promoter

sequences of these species, as well as a few other Pichiales for comparison. We predicted the

conserved transcription factor (TF) binding sites in these putative promoters to identify TFs that

had more binding sites in our rapid ECAR species compared to the low ECAR species. We

found that the promoters of some glycolytic genes had additional Gal4p-binding sites in the

rapid ECAR species (Fig 4A, Table S2). In rapid ECAR *Saturnispora*, there was an extra

predicted Gal4p-binding site in the promoter of *TDH*, the most highly expressed glycolytic gene.

There were also uniquely predicted Gal4p-binding sites in the promoters of *HXK* and *ENO* in the

rapid ECAR species, which were absent from the low ECAR species (Fig. 4A, Table S2).

Interestingly, when we compared these promoters to several other Pichiales species, we found

that there was considerable variation, suggesting that they may be quite variable across

species.

To investigate the importance of Gal4p in the rapid ECAR phenotype, we performed a

phylogenetic analysis and found that, despite some lineage-specific duplications in this

orthogroup, *GAL4* is likely a 1:1 ortholog in *S. cerevisiae*, *K. phaffii*, *Sat. dispora*, and most

yeasts (Fig. S4). We then developed a system to genetically modify a rapid ECAR yeast, *Sat.*

dispora, based on methods previously described for *K. phaffii* (De Schutter & Callewaert, 2012)

and generated a deletion mutant by replacing this *GAL4* ortholog with a *kanMX* or *natMX*

resistance cassette using long homology to direct homologous recombination. Two independent

deletion mutants lacking *GAL4* had decreased glucose consumption, ethanol production, and a

low ECAR phenotype (Fig. 4). We also found that mutants lacking *GAL4* had a more respiratory

phenotype with a higher oxygen consumption rate (Fig. 4E, Fig. S5), further supporting the

hypothesis that the Gal4p transcription factor is required for the rapid ECAR metabolic

phenotype and aerobic fermentation in *Sat. dispora*.

Figure 4. The Gal4p transcription factor contributes to the rapid ECAR metabolic phenotype in *Sat. dispora*. (A) The number of predicted Gal4p-binding sites in the putative promoters 1kb upstream of each glycolytic gene in several Pichiales species. The genes marked with an asterisk have an additional binding site in the rapid ECAR species. The (B) growth and (C) extracellular metabolites for the *Sat. dispora* wild-type (WT) control and two independent *gal4Δ* deletion mutants. Two high ECAR and two low ECAR species were measured during growth in a highly aerobic 250-ml baffled flask. The error bars represent the standard deviation of four biological replicates. (D) The *gal4Δ* deletion mutants showed a low ECAR phenotype compared to the parent (WT) control. (E) The *gal4Δ* deletion mutants displayed higher oxygen consumption rates (OCR) at 90 minutes post inoculation compared to the parent (WT) control. The centre lines of boxplots represent the median, the bounds of the boxes represent the interquartile range, the whiskers represent the spread of the data, and each dot represents one of three biological replicates. The p values shown are the results of two-sided t-tests compared to the wild-type control.

To test whether there was a direct cis-regulatory effect of Gal4p, we also made a GFP

reporter construct driven by the *Sat. dispora TDH* promoter. Furthermore, we made a version of

this reporter in which the predicted Gal4p-binding sites were mutated using site-directed

mutagenesis. We inserted these reporters into a genomic safe haven, in which there were no

predicted genes, in both the wild-type background and the *gal4Δ::natMX* deletion mutants. The

expression of GFP was significantly reduced in the wild-type background when the predicted

binding sites were mutated. The expression was also significantly reduced when the reporter

construct with intact Gal4p-binding sites was transformed into the *gal4Δ::natMX* mutant.

Together these results show that, in the absence of either the trans-acting transcription factor or

the cis-regulatory binding sites, the promoter activity is greatly reduced (Fig. 5). Although the

GFP expression was very low in the *gal4Δ::natMX* deletion background, there was a small but

significant difference between the wild-type and mutant reporter constructs, suggesting that the

mutations may also have interfered with the ability of other transcription factors to bind.

Altogether, this result directly shows that Gal4p is necessary to activate the most highly

expressed glycolytic gene.

Figure 5. Gal4p is required for the induction of the *TDH* promoter. (A) A schematic diagram showing the location of the predicted Gal4p-binding sites in the *TDH* promoter, the reporter construct, and the mutations made in the predicted Gal4p-binding sites in the reporter construct promoter. (B) The fluorescence intensity of the wild-type and *gal4Δ* deletion mutants containing the reporter with the wild-type *TDH* promoter (P_{TDH}-*GFP*) or the mutated *TDH* promoter ($P_{tdh-UASG}$-*GFP*). The fluorescence of each reporter-expressing strain was normalized to its respective control strain (i.e. wild-type and *gal4Δ* mutants) lacking any reporter. The centre lines of boxplots represent the median, the bounds of the boxes represent the interquartile range, the whiskers represent the spread of the data, and each dot represents one of four biological replicates. The p values shown are the results of two-sided t-tests comparing the strains as indicated. (C) Representative images of the GFP fluorescence from the strains carrying the reporter constructs. The scale bar represents 10 μm. Uncropped images are available as Supplemental Data 1.

DISCUSSION

The discovery of shared or novel mechanisms to increase glycolytic rate can help us

understand the evolutionary constraints and guide manipulation of this trait across diverse yeast

species. In this study, we developed an assay to assess variation in glycolytic rate across the

yeast subphylum. Comparative genomics corroborated previous reports that duplications of

glycolytic genes were associated with increased glycolytic rates in yeasts, particularly in the

Saccharomycetales where a WGD occurred (Conant & Wolfe, 2007; Hagman & Piškur, 2015;

Dashko *et al*, 2014; Hagman *et al*, 2013; Lin & Li, 2011). However, there were no duplications or

expansions in glycolytic genes associated with the rapid ECAR phenotype in the Pichiales. This

contrast suggested that the rapid ECAR phenotype we observed in several *Saturnispora*

species had an independent mechanism. Using transcriptomics, we observed that several

glycolytic genes were upregulated in species with rapid ECAR. While these results were

consistent with the phenotypic observations, we were surprised to see how closely they

mirrored the gene duplications in *S. cerevisiae* (Conant & Wolfe, 2007). In particular, the genes

encoding enzymes in lower glycolysis, which were duplicated and retained in *S. cerevisiae*, had

much higher expression in the rapid ECAR species. The association of elevated expression of

lower glycolysis genes with aerobic fermentation is also consistent with previous evidence that

pull through lower glycolysis is key to fermentative capacity (Smits *et al*, 2000). Furthermore,

lower glycolysis, carries more flux than any biochemical alternative (Court *et al*, 2015),

suggesting that upregulating these existing genes may be the most effective strategy to

increase glycolytic rate.

Since the driving force between this phenotypic difference seemed to be at the

transcriptional level, we retrieved the putative promoter sequences upstream of glycolytic genes

to identify transcriptional activators that had more binding sites in the promoters in rapid ECAR

species. Several of the glycolytic gene promoters had more conserved binding sites for the

Gal4p transcription factor. Although Gal4p is well characterized for its role in activating the Leloir

pathway for galactose catabolism in *S. cerevisiae* (Johnston, 1987), it is also known to have

other roles in metabolism in different yeasts (Askew *et al*, 2009; Kim *et al*, 2024; Ata *et al*,

2018). Indeed, in another Pichiales yeast, the protein production chassis, *K. phaffii*,

overexpression of Gal4p has been shown to induce overflow metabolism (Ata *et al*, 2018). In

the case of *Sat. dispora,* this yeast natively has a rapid glycolytic rate and produces ethanol and

acetate as overflow metabolites. The promoters with additional Gal4p-binding sites in rapid

ECAR species compared to the low ECAR species in *Saturnispora* were upstream of *HXK,*

TDH, and *ENO.* This configuration again bore a striking parallel to modelling in *S. cerevisiae*

that predicted these three enzymes have a large impact on flux when their abundance

decreases (Conant & Wolfe, 2007). These results provide mechanistic insights into how the

transcriptional activation of glycolysis in *Saturnispora* evolved and suggest that the modelling

predicting the importance of these three key glycolytic enzymes extends to other yeasts beyond

S. cerevisiae.

To confirm the direct influence of Gal4p on the rapid ECAR phenotype, we developed a

transformation protocol for *Sat. dispora*. The deletion of *GAL4* decreases the glucose

consumption rate and renders the mutants more respiratory than fermentative. This result

confirms the importance of the Gal4p transcription factor in the rapid ECAR phenotype. To

ensure that *GAL4* deletion had a specific impact on the cis-regulation of glycolytic genes, we

developed a reporter system using the promoter of *TDH*, which was the most highly expressed

gene in both *Sat. dispora* and *Sat. hagleri*. Through monitoring the expression of GFP as a

reporter under the *TDH* promoter, we found that, if either the *GAL4* gene were missing or if the

Gal4p-binding sites were mutated, the GFP expression was greatly reduced, showing that both

the transcription factor Gal4p and its cis-regulatory binding sites are directly required for

induction. Although Gal4p is required for the high glucose consumption rate of *Sat. dispora*, we

can also infer that there are likely additional factors that contribute to the divergence in

phenotypes of the rapid and low ECAR species. For example, the *gal4Δ* mutants still

accumulated high levels of acetate and did not grow to the high biomass seen in the low ECAR

species.

The yeast phenotype of overflow metabolism is one that, on the surface, seems wasteful

as the organism produces fewer ATP molecules per glucose consumed. However, this strategy

has been proposed to deplete a limited resource and convert it into another product, which

would limit the growth of other microbes and could later be consumed. This "make-accumulate-

consume" phenotype has already been observed in two distinct yeast lineages (Rozpedowska

et al, 2011), and we now have evidence for a third independent lineage with a novel mechanism

driven by increased transcriptional activation of glycolysis. There are also interesting parallels in

the ecology of the rapid ECAR *Saturnispora* species and *S. cerevisiae*. *Sat. dispora* has

primarily been isolated from tree exudates, and *Sat. hagleri* has been isolated from fruits and

insects (Kurtzman *et al*, 2011; Harrison *et al*, 2024). Similar to *S. cerevisiae*, these yeasts are

isolated from environments that have occasional periods of rich sugar availability, which may

have prompted the evolution of rapid glycolysis (Liti, 2015; Jouhten *et al*, 2016). Although the

rapid ECAR phenotype of *Sat. dispora* arose from a distinct cis-regulatory mechanism, many of

the changes appear to lead to similar functional outcomes to those observed in *S. cerevisiae*.

Even though the precise molecular mechanism of evolution differed between *Sat. dispora* and

S. cerevisiae, the end result was still increased abundance of lower glycolytic enzymes,

glyceraldehyde-3-phosphate dehydrogenase, and hexokinase. Like the evolution of C4

photosynthesis in plants, the aerobic fermentation phenotype in yeasts represents another case

of convergent evolution in metabolic efficiency.

In the case of C4 plants, the abundance, activity, or localization of the same key C4

enzymes has been optimized through convergent evolution, whereas we find that, in aerobically

fermenting yeasts, the abundance of the same key glycolytic enzymes has been convergently

increased through gene duplications or through the recruitment of cis-regulatory elements. This

study provides new evidence supporting recent findings that trait evolution often employs the

same genes across divergent lineages of yeasts (David *et al*, 2025b; Aranguiz *et al*, 2025).

While previous studies have focused on gene presence and family size, we were able to identify

a mechanism of gene upregulation through the recruitment of cis-regulatory elements in the

promoters of congeneric species. These convergent adaptations suggest that there are likely

strong constraints on how a yeast can evolve a rapid glycolytic rate. Indeed, high glycolytic rates

and retained glycolytic duplicates may be a predictable occurrence, as suggested by multiple

recently discovered novel yeast WGDs (David *et al*, 2025a). Whether by gene duplication or cis-

regulatory evolution, it seems that an increase in the abundance of the same set of glycolytic

enzymes is required for yeasts to achieve high glycolytic rates. While organisms may deploy

divergent molecular mechanisms due to chance and the historical contingencies of their

lineages, these mounting examples suggest that evolutionary convergence operates within the

constraints of the pathway being manipulated by natural selection.

METHODS

Strain and culture conditions

All strains screened were previously described (Shen *et al*, 2018; Opulente *et al*, 2024)

and are listed in Table S1. Yeasts were routinely maintained on yeast extract peptone dextrose

(YPD) agar at room temperature. Fermentation and extracellular acidification assays were

conducted using synthetic complete (SCD) media containing yeast nitrogen base, 5 g/L

ammonium sulfate, complete dropout mix, and 20 g/L D-glucose (US Biologicals). All growth

conditions were at room temperature (RT, 22 °C) unless otherwise specified.

Extracellular acidification rate assay

All yeasts were grown for 24 hrs in 2 ml YPD, sub-cultured 1:4 in fresh SCD pH 6.8, and

grown for an additional 6 hrs. These actively growing cells were harvested by centrifugation,

washed twice in sterile water to remove excess media, and normalized by optical density. Cells

were inoculated into a 96-well plate at OD_{600nm} = 0.15 in SCD supplemented with 0.002%

phenol red. For each strain, 2% dextrose or a water control was added immediately before

beginning the assay. The absorbance at 560 nm was measured every 3 minutes for 90 minutes,

and the change in extracellular pH was calculated based on a standard curve (Fig. S1). The

dextrose-dependent acidification rate was determined as the difference between the

acidification rate in the presence of 2% dextrose and the acidification rate in the water control.

For each yeast species, three biological replicates were conducted on different days. The

average ECAR was plotted as a colour strip outside the phylogeny using iTOL v7 (Letunic &

Bork, 2024)

Comparative genomics

The orthogroups corresponding to genes encoding glycolytic enzymes were retrieved

from the published OrthoFinder analysis (Opulente *et al*, 2024). A correlation test between the

number of orthologs and the dextrose dependent acidification of each species was conducted

for each gene as well as the total sum of glycolytic genes. This analysis was also repeated for

the orders Saccharomycetales and Pichiales. To account for phylogeny, we used phylogenetic

generalized least squares (PGLS) using *R* v4.2.2 and the package caper v1.0.3 (Orme, 2013).

Data were visualized using ggpubr v0.6.0 (Kassambara, 2023) and corrplot v0.95 (Wei & Simko,

2024).

Growth and metabolite profiling

Yeast strains were grown overnight in YPD in biological quadruplicate. Overnight

cultures were collected, washed twice in sterile water, and inoculated in 50 ml fresh SCD pH 6.8

in 250-ml baffled flasks at a density of OD_{600nm} = 0.05. The flasks were incubated at RT while

shaking at 225 rpm. At each time point, OD_{600nm} was measured, and the supernatant was

analysed for glucose, ethanol, and acetate concentrations using high performance liquid

chromatography with a refractive index detector (HPLC-RID), as previously described

(Schwalbach *et al*, 2012). The mean and standard deviation across replicates was calculated in

R v4.2.2 and plots were made using the package ggplot2 v3.5.2 (Wickham, 2016).

RNA extraction and sequencing

RNA was collected from log-phase cells using acidic phenol as previously described for yeast (Collart & Oliviero, 1993). Briefly, cells were collected and mixed with 0.1 volumes of 5% acid phenol in 95% ethanol, harvested by centrifugation, and flash frozen. Cell pellets were lysed with glass beads, and RNA was extracted with acidic phenol twice, followed by a final chloroform extraction. The RNA was treated with RQ1 DNase (Promega) and subsequently cleaned up using an RNeasy column (Qiagen). Since the rapid and low ECAR yeasts grew at different rates, to obtain mid-log phase for all strains, the rapid ECAR yeasts were collected at 15 hrs post-inoculation, and the low ECAR yeasts were collected at 18 hrs post-inoculation.

We enriched for mRNA using the NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB), and libraries were prepared using the NEBNext Ultra II Directional RNA Library Kit for Illumina (NEB) following the manufacturer's instructions. Paired-end libraries were sequenced on a NovaSeq X Plus (Illumina) with 2x150 bp reads. Reads were trimmed using Trimmomatic v0.30 (Bolger *et al*, 2014) to remove adapters and subsequently checked in FastQC (Andrews S., 2010). Reads were mapped to the respective genome of each species using HiSat v2.1.0 (Kim *et al*, 2019). HTSeq-count v2.0.3 (Putri *et al*, 2022) was used to generate the counts of annotated genes using the non-unique fraction setting to avoid overcounting or discarding highly similar genes. To compare the transcriptome between species, we used OrthoFinder (Emms & Kelly, 2019) to identify groups of orthologous genes within the *Saturnispora* species. Subsequently, the sum of counts and gene lengths for each orthogroup was calculated using a custom python script (https://github.com/Linda-Horianopoulos/Aerobic_fermentation_scripts). The counts and total gene lengths were then used to determine the TPM for each orthogroup in each species using EdgeR (Robinson *et al*, 2009). The orthogroups encoding glycolytic and citric acid cycle enzymes were retrieved, and plots were made to visualize the expression in each species in *R* v4.2.2 using ggplot2 v3.5.2 (Wickham, 2016).

Identification of putative transcription factor binding sites

Putative promoter sequences for each of the glycolytic genes were retrieved from the

annotation files using gffread v0.12.7 (Pertea & Pertea, 2020). Yeastract (Teixeira *et al*, 2023)

was used to identify all conserved *S. cerevisiae* transcription factor binding sites in the 1000 bp

upstream of the coding sequence of each glycolytic gene from *Sat. dispora, Sat. hagleri, Sat.*

mendoncae, Sat. silvae, *Martiniozyma abietophilia*, *P. kudriavzevii*, and *K. phaffii*. The number of

predicted Gal4p-binding sites (exact consensus matches to $CGGN_{11}CCG$ (Johnston, 1987)) in

the promoter of each gene for each species was visualized using iTOL v7 (Letunic & Bork,

2024).

Phylogenetic analysis of *GAL4* orthologs

The genes in the same orthogroup as the canonical *S. cerevisiae GAL4* (YPL248C)

were retrieved for all species included in our ECAR screen. The translated amino acid

sequences were retrieved, aligned using MAFFT v7.222 (Katoh & Standley, 2013), and the

alignment was trimmed using trimal v1.4.1 applying the gappyout option (Capella-Gutiérrez *et*

al, 2009). The phylogeny was inferred using IQtree v1.6.8 with automatic substitution model

selection and 1000 bootstraps (Nguyen *et al*, 2015). The phylogeny was visualized in iTOL v7

(Letunic & Bork, 2024) and manually pruned to remove an outgroup on a disproportionately long

internal branch, which encode divergent C6 zinc transcription factors.

Gene deletion

The coding sequence of *GAL4* in *Sat. dispora* was replaced with a *kanMX* marker using

an electroporation protocol adapted from *K. phaffii* (De Schutter & Callewaert, 2012). Briefly, ~1-

kb regions upstream and downstream of the *GAL4* coding sequence were amplified from

genomic DNA using primers with homology to the *kanMX* marker and the pRS426 backbone

(Table S3). The *kanMX* marker was amplified from pUG6 (Goldstein & McCusker, 1999) and the

pRS426 backbone (Christianson *et al*, 1992) was also amplified using primers specified in Table

S3. These fragments were assembled using HiFi assembly (NEB) and transformed into *E. cloni*

(Lucigen). The full construct was subsequently amplified from the plasmid, and 1 μg was

transformed into *Sat. dispora*. Electroporation was performed in 80 μL of ice cold 1 M sorbitol

using 1.5 kV. After electroporation, cells were recovered in 1 mL of YPD + 1 M sorbitol and

plated on YPD plates supplemented with 1 M sorbitol and 300 μg/ml G418. Transformants were

screened for stable resistance against G418, and correct replacement of the coding sequence

was confirmed with colony PCR. To allow the *kanMX* marker to be used for additional

modifications, the *kanMX* cassette was replaced with a *natMX* cassette, which was amplified

from pAG25, using the homology of the promoter and terminator. For selection of colonies

integrating the *natMX* marker, colonies were plated on YPD plates supplemented with 1 M

sorbitol and 25 μg/ml nourseothricin. To ensure that the phenotype was not due to any spurious

background mutations, two independent colonies were used in all experiments. The strains

constructed for this study are listed in Table S4.

Oxygen consumption measurements

The Resipher system (Lucid Scientific) was used to monitor the oxygen levels and

calculate the oxygen consumption rates (OCR). Three biological replicates of overnight cultures

were washed twice in sterile water and normalized to OD600 = 0.1 in 95 μl in a 96-well plate in

SC medium lacking a carbon source. The plate was centrifuged at 300 *x g* for 2 minutes to

pellet the cells to the bottom of the plate, and immediately before beginning reading, 5 ul of 40%

dextrose was added for a final concentration of 2% dextrose. The plate was incubated in the

dark at room temperature while the oxygen concentration was measured. The oxygen

consumption rates were retrieved at the 90-minute timepoint for consistency with the ECAR

assays. The data was visualized in *R* using ggpubr v0.6.0 (Kassambara, 2023) and the

significance of differences between strains were determined using a two-sided t-test.

Evaluating the cis-regulatory elements in the promoter of *TDH*

An insertion construct was designed in a genomic safe haven to allow heterologous

proteins to be expressed from a region of the genome with no risk of gene disruption. A plasmid

containing ~1kb regions to direct homologous recombination to the genomic safe haven site

flanking the sequence for GFP driven by the *TDH* promoter from *Sat. dispora* was synthesized

by the Joint Genome Institute (JGI). Since this promoter had two predicted Gal4p-binding sites,

we mutated them both using site directed mutagenesis as previously described (Zheng *et al*,

2004) with the modification that Phusion polymerase was used to amplify the construct. Binding

site mutants were confirmed by whole plasmid sequencing (Plasmidsaurus). Both versions of

the reporter construct were transformed into the *Sat. dispora* wild-type strain and *gal4Δ::natMX*

deletion mutant using electroporation. Transformants were screened using colony PCR. Four

biological replicates of each strain, including controls lacking any reporter were grown overnight

in YPD, washed twice in sterile water, and inoculated in fresh SC+D pH 6.8 at OD_{600nm} = 0.1.

After growing for 40 hrs, cells were normalized to OD_{600nm} = 1 and the fluorescence intensity

was measured using a BMG Labtech FLUOstar plate reader. The background from control

strains without a reporter was subtracted and the fluorescence intensity was plotted in *R* using

the package ggpubr v0.6.0 (Kassambara, 2023). The statistical significance was determined

using two-sided t-tests.

Images of cells expressing GFP in the WT and *gal4Δ* background with the intact and

mutated promoter grown under the same conditions (SC+D pH 6.8 for 40 hours) were captured

with an EVOS FL microscope with 100x oil immersion plan apochromat objective lens. Images

were uniformly processed and cropped in FIJI (Schindelin *et al*, 2012). Unedited images are

available as Supplemental Data 1.

Data availability

The RNA-Sequencing data has been deposited to NCBI's GEO and the accession number will

be available upon publication. All other data are provided as supplemental tables. Custom code

is available on Github (https://github.com/Linda-Horianopoulos/Aerobic_fermentation_scripts) or

as a release on Zenodo (https://doi.org/10.5281/zenodo.17185390).

Acknowledgements

The authors thank members of the Hittinger Lab and Y1000+ Project (http://y1000plus.org) team members for helpful discussions. This project was supported by the National Science Foundation under Grants No. DEB-2110403 (to C.T.H.); DEB-2110404 (to A.R.); in part by the Great Lakes Bioenergy Research Center, U.S. Department of Energy, Office of Science, Biological and Environmental Research Program under Award Number DE-SC0018409 (of which C.T.H. is a co-investigator); and the National Institute of Food and Agriculture, United States Department of Agriculture, Hatch project 7005101 (to C.T.H.). Research in A.R.'s lab is also supported by the National Institutes of Health/National Institute of Allergy and Infectious Diseases (R01 AI153356). The work (proposal: 10.46936/10.25585/60008615) conducted by the U.S. Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy operated under Contract No. DE-AC02-05CH11231. We specifically thank Ian Blaby for helping design and Robert Evans for synthesizing the safe haven insertion plasmid as part of Yasuo Yoshikuni's team. L.C.H. was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) postdoctoral fellowship.

Author contributions

L.C.H. and C.T.H. conceptualized this project. L.C.H. developed methodology, performed experiments, and analysed data with the input and guidance of A.R. and C.T.H. L.C.H. and C.T.H. wrote the manuscript, and A.R. reviewed, edited, and approved the manuscript. L.C.H., C.T.H., and A.R. acquired funding for this project.

Conflict of Interest

A.R. is a scientific consultant of LifeMine Therapeutics, Inc. All other authors declare no competing interests.

REFERENCES

Andrews S. (2010) FastQC: a quality control tool for high throughput sequence data. *http://www.bioinformatics.babraham.ac.uk/projects/fastqc*

Aranguiz K, Horianopoulos LC, Elkin L, Abá KS, Jordahl D, Overmyer KA, Wrobel RL, Coon JJ, Shiu SH, Rokas A, *et al* (2025) Machine learning reveals genes impacting oxidative stress resistance across yeasts. *Nature Communications 2025 16:1* 16: 1–15

Askew C, Sellam A, Epp E, Hogues H, Mullick A, Nantel A & Whiteway M (2009) Transcriptional Regulation of Carbohydrate Metabolism in the Human Pathogen *Candida albicans*. *PLoS Pathog* 5: e1000612

Ata Ö, Rebnegger C, Tatto NE, Valli M, Mairinger T, Hann S, Steiger MG, Çalık P & Mattanovich D (2018) A single Gal4-like transcription factor activates the Crabtree effect in *Komagataella phaffii*. *Nat Commun* 9: 1–10

Bolger AM, Lohse M & Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30: 2114

Brown NJ, Newell CA, Stanley S, Chen JE, Perrin AJ, Kajala K & Hibberd JM (2011) Independent and parallel recruitment of preexisting mechanisms underlying C4 photosynthesis. *Science (1979)* 331

Capella-Gutiérrez S, Silla-Martínez JM & Gabaldón T (2009) trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25: 1972

Christianson TW, Sikorski RS, Dante M, Shero JH & Hieter P (1992) Multifunctional yeast high-copy-number shuttle vectors. *Gene* 110: 119–122

Collart MA & Oliviero S (1993) Preparation of Yeast RNA. *Curr Protoc Mol Biol* 23: 13.12.1-13.12.5

Conant GC & Wolfe KH (2007) Increased glycolytic flux as an outcome of whole-genome duplication in yeast. *Mol Syst Biol* 3

Conant GC & Wolfe KH (2008) Turning a hobby into a job: How duplicated genes find new functions. *Nature Reviews Genetics 2008 9:12* 9: 938–950

Court SJ, Waclaw B & Allen RJ (2015) Lower glycolysis carries a higher flux than any biochemically possible alternative. *Nature Communications 2015 6:1* 6: 1–8

Dashko S, Zhou N, Compagno C & Piškur J (2014) Why, when, and how did yeast evolve alcoholic fermentation? *FEMS Yeast Res* 14: 826–832

David KT, Horianopoulos L, Gonçalves C, Steenwyk JL, Pontes A, Gonçalves P, Hittinger CT, Pennell M & Rokas A (2025a) Discovery of additional ancient genome duplications in yeasts. *bioRxiv*: 2025.08.31.673279

David KT, Schraiber JG, Crandall JG, Labella AL, Opulente DA, Harrison MC, Wolters JF, Zhou X, Shen XX, Groenewald M, *et al* (2025b) Convergent expansions of keystone gene families drive metabolic innovation in Saccharomycotina yeasts. *Proc Natl Acad Sci U S A* 122: e2500165122

Emms DM & Kelly S (2019) OrthoFinder: Phylogenetic orthology inference for comparative
genomics. *Genome Biol* 20

Escalera-Fanjul X, Quezada H, Riego-Ruiz L & González A (2019) Whole-Genome Duplication
and Yeast's Fruitful Way of Life. *Trends in Genetics* 35

Gasser B & Mattanovich D (2018) A yeast for all seasons – Is *Pichia pastoris* a suitable chassis
organism for future bioproduction? *FEMS Microbiol Lett* 365: 181

Geijer C, Ledesma-Amaro R & Tomas-Pejo E (2022) Unraveling the potential of non-
conventional yeasts in biotechnology. *FEMS Yeast Res* 22

Goldstein AL & McCusker JH (1999) Three new dominant drug resistance cassettes for gene
disruption in *Saccharomyces cerevisiae*. *Yeast* 15

Hagman A & Piškur J (2015) A Study on the Fundamental Mechanism and the Evolutionary
Driving Forces behind Aerobic Fermentation in Yeast. *PLoS One* 10: e0116942

Hagman A, Säll T, Compagno C & Piskur J (2013) Yeast '"Make-Accumulate-Consume"' Life
Strategy Evolved as a Multi-Step Process That Predates the Whole Genome Duplication.

Harrison MC, LaBella AL, Hittinger CT & Rokas A (2022) The evolution of the GALactose
utilization pathway in budding yeasts. *Trends in Genetics* 38: 97–106

Harrison MC, Opulente DA, Wolters JF, Shen XX, Zhou X, Groenewald M, Hittinger CT, Rokas A
& LaBella AL (2024) Exploring Saccharomycotina Yeast Ecology Through an Ecological
Ontology Framework. *Yeast* 41: 615–628

Heistinger L, Gasser B & Mattanovich D (2020) Microbe Profile: *Komagataella phaffii*: a
methanol devouring biotech yeast formerly known as *Pichia pastoris*. *Microbiology (N Y)*
166: 614–616

Johnston M (1987) A model fungal gene regulatory mechanism: the *GAL* genes of
Saccharomyces cerevisiae. *Microbiol Rev* 51

Jouhten P, Ponomarova O, Gonzalez R & Patil KR (2016) *Saccharomyces cerevisiae*
metabolism in ecological context. *FEMS Yeast Res* 16: 80

Kassambara A (2023) ggpubr: 'ggplot2' Based Publication Ready Plots. *R package version 060*

Katoh K & Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol Biol Evol* 30: 772–780

Kim D, Paggi JM, Park C, Bennett C & Salzberg SL (2019) Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37: 907–915

Kim M-J, White AM & Mitchell AP (2024) Strain variation in *Candida albicans* glycolytic gene
regulation. *mSphere* 9

Kurtzman CP, Fell JW & Boekhout T (2011) The Yeasts: A Taxonomic Study, Volume 1 Elsevier

Letunic I & Bork P (2024) Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool. *Nucleic Acids Res* 52

Lin Z & Li WH (2011) Expansion of hexose transporter genes was associated with the evolution
of aerobic fermentation in yeasts. *Mol Biol Evol* 28: 131–142

Liti G (2015) The fascinating and secret wild life of the budding yeast *S. cerevisiae*. *Elife* 4

Lyu H, Yim WC & Yu Q (2024) Genomic and Transcriptomic Insights into the Evolution of C4
Photosynthesis in Grasses. *Genome Biol Evol* 16

Lyu MJA, Du H, Yao H, Zhang Z, Chen G, Huang Y, Ni X, Chen F, Zhao YY, Tang Q, *et al* (2025)
A dominant role of transcriptional regulation during the evolution of C4 photosynthesis in
Flaveria species. *Nature Communications 2025 16:1* 16: 1–17

Opekarova M & Sigler K (1982) Acidification power: indicator of metabolic activity and autolytic
changes in *Saccharomyces cerevisiae*. *Folia Microbiol (Praha)* 27: 395–403

Nguyen LT, Schmidt HA, Von Haeseler A & Minh BQ (2015) IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol* 32:
268–274

Opulente DA, LaBella AL, Harrison MC, Wolters JF, Liu C, Li Y, Kominek J, Steenwyk JL,
Stoneman HR, VanDenAvond J, *et al* (2024) Genomic factors shape carbon and nitrogen
metabolic niche breadth across Saccharomycotina yeasts. *Science* 384: eadj4503

Orme D (2013) The caper package : comparative analysis of phylogenetics and evolution in R.
R package version 05, 2

Patra P, Das M, Kundu P & Ghosh A (2021) Recent advances in systems and synthetic biology
approaches for developing novel cell-factories in non-conventional yeasts. *Biotechnol Adv*
47 doi:10.1016/j.biotechadv.2021.107695

Pertea G & Pertea M (2020) GFF Utilities: GffRead and GffCompare. *F1000Res* 9

Putri GH, Anders S, Pyl PT, Pimanda JE & Zanini F (2022) Analysing high-throughput
sequencing data in Python with HTSeq 2.0. *Bioinformatics* 38: 2943–2945

Pyne ME, Bagley JA, Narcross L, Kevvai K, Exley K, Davies M, Wang Q, Whiteway M & Martin
VJJ (2023) Screening non-conventional yeasts for acid tolerance and engineering *Pichia*
occidentalis for production of muconic acid. *Nature Communications 2023 14:1* 14: 1–13

Radecka D, Mukherjee V, Mateo RQ, Stojiljkovic M, Foulquié-Moreno MR & Thevelein JM
(2015) Looking beyond *Saccharomyces*: The potential of non-conventional yeast species
for desirable traits in bioethanol fermentation. *FEMS Yeast Res* 15
doi:10.1093/femsyr/fov053

Robinson MD, McCarthy DJ & Smyth GK (2009) edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* 26: 139

Rozpedowska E, Hellborg L, Ishchuk OP, Orhan F, Galafassi S, Merico A, Woolfit M, Compagno
C & Piškur J (2011) Parallel evolution of the make–accumulate–consume strategy in
Saccharomyces and *Dekkera* yeasts. *Nature Communications 2011 2:1* 2: 1–7

Sage RF, Christin PA & Edwards EJ (2011) The C4 plant lineages of planet Earth. *J Exp Bot* 62:
3155–3169

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
C, Saalfeld S, Schmid B, *et al* (2012) Fiji: An open-source platform for biological-image
analysis. *Nat Methods* 9: 676–682

Schmidt CA, Fisher-Wellman KH & Darrell Neufer P (2021) From OCR and ECAR to energy:
Perspectives on the design and interpretation of bioenergetics studies. *Journal of Biological*
Chemistry 297: 101140–101141

De Schutter K & Callewaert N (2012) *Pichia* surface display: A tool for screening single domain
antibodies. *Methods in Molecular Biology* 911: 125–134

Schwalbach MS, Keating DH, Tremaine M, Marner WD, Zhang Y, Bothfeld W, Higbee A, Grass
JA, Cotten C, Reed JL, *et al* (2012) Complex physiology and compound stress responses
during fermentation of alkali-pretreated corn stover hydrolysate by an *Escherichia coli*
ethanologen. *Appl Environ Microbiol* 78: 3442–3457

Shen XX, Opulente DA, Kominek J, Zhou X, Steenwyk JL, Buh K V., Haase MAB, Wisecaver
JH, Wang M, Doering DT, *et al* (2018) Tempo and Mode of Genome Evolution in the
Budding Yeast Subphylum. *Cell* 175: 1533-1545.e20

Sigler K (2013) Acidification power (AP) test and similar methods for assessment and prediction
of fermentation activity of industrial microorganisms. *Kvasny Prumysl* 59: 204–208

Sigler K, Knotková A & Kotyk A (1981) Factors governing substrate-induced generation and
extrusion of protons in the yeast *Saccharomyces cerevisiae*. *BBA - Biomembranes* 643:
572–582

Smits HP, Hauf J, Müller S, Hobley TJ, Zimmermann FK, Hahn-Hägerdal B, Nielsen J & Olsson
L (2000) Simultaneous overexpression of enzymes of the lower part of glycolysis can
enhance the fermentative capacity of *Saccharomyces cerevisiae*. *Yeast* 16

Suthers PF & Maranas CD (2022) Examining organic acid production potential and growth-
coupled strategies in *Issatchenkia orientalis* using constraint-based modeling. *Biotechnol*
Prog 38

Swift J, Luginbuehl LH, Hua L, Schreier TB, Donald RM, Stanley S, Wang N, Lee TA, Nery JR,
Ecker JR, *et al* (2024) Exaptation of ancestral cell-identity networks enables C4
photosynthesis. *Nature 2024 636:8041* 636: 143–150

Tan SI, Liu Z, Tran VG, Martin TA & Zhao H (2025) *Issatchenkia orientalis* as a platform
organism for cost-effective production of organic acids. *Metab Eng* 89: 12–21

Teixeira MC, Viana R, Palma M, Oliveira J, Galocha M, Mota MN, Couceiro D, Pereira MG,
Antunes M, Costa I V., *et al* (2023) YEASTRACT+: a portal for the exploitation of global
transcription regulation and metabolic model data in yeast biotechnology and
pathogenesis. *Nucleic Acids Res* 51: D785–D791

Tucey TM, Verma J, Harrison PF, Snelgrove SL, Lo TL, Scherer AK, Barugahare AA, Powell DR,
Wheeler RT, Hickey MJ, *et al* (2018) Glucose Homeostasis Is Important for Immune Cell
Viability during *Candida* Challenge and Host Survival of Systemic Fungal Infection. *Cell*
Metab 27: 988-1006.e7

Wang Y, Wang Y, Cui J, Wu C, Yu B & Wang L (2025) Non-conventional yeasts: promising cell
factories for organic acid bioproduction. *Trends Biotechnol* 43: 1609–1626

Wei T & Simko V (2024) R package 'corrplot': Visualization of a Correlation Matrix. *(Version
095)*

Wickham H (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York

Williams BP, Aubry S & Hibberd JM (2012) Molecular evolution of genes recruited into C4
photosynthesis. *Trends Plant Sci* 17: 213–220

Yamashoji S, Al Mamun A & Bari L (2020) Visual and simple determination of glucose-induced
acidification by yeast cells: application to rapid cytotoxicity test. *Heliyon* 6: e03924

Yoo K, Bhattacharya S, Oliveira NK, Pereira de Sa N, Matos GS, Del Poeta M & Fries BC
(2024) With age comes resilience: how mitochondrial modulation drives age-associated
fluconazole tolerance in *Cryptococcus neoformans*. *mBio* 15

Zhang J & Zhang Q (2019) Using Seahorse machine to measure OCR and ECAR in cancer
cells. *Methods in Molecular Biology* 1928: 353–363

Zhang P, Li H, Cheng J, Sun AY, Wang L, Mirchevska G, Calderone R & Li D (2018) Respiratory
stress in mitochondrial electron transport chain complex mutants of *Candida albicans*
activates Snf1 kinase response. *Fungal Genetics and Biology* 111: 73–84

Zheng L, Baumann U & Reymond JL (2004) An efficient one-step site-directed and site-
saturation mutagenesis protocol. *Nucleic Acids Res* 32: e115–e115

Supplemental Figure 1.Optimization of the Extracellular Acidification Rate (ECAR) assay.
(A) A standard curve of the absorbance at 560nm generated from media between pH 6.5 and
7.5. (B) Optimization assay using *S. cerevisiae* showing that the control with no carbon source
(noC) did not acidify the media, whereas the cells with 2% glucose added (dex) consistently
acidified the media across six biological replicates. (C) The slopes calculated from the six
biological replicates showing the dextrose-dependent pH change.

757
Supplemental Figure 2. The gene family size of glycolytic genes correlates with ECAR
across all yeasts. (A-C) The correlation between ECAR and the number of homologs of each
glycolytic gene showing the Pearson's correlation and p-value for (A) all phenotyped yeasts, (B)
the yeasts in the order Saccharomycetales, and (C) the yeasts in the order Pichiales. (D-F) The
phylogenetically corrected correlations between ECAR and the number of homologs of each
glycolytic gene for (D) all phenotyped yeasts, (E) the yeasts in the order Saccharomycetales,
and (F) the yeasts in the order Pichiales. The R-value from the phylogenetic generalized least
squares is in the box and the p-value is indicated as * p<0.05 and ** p<0.01. Data points are
colour coded according to taxonomic order as in Figure 1.

Supplemental Figure 3. Overview of comparative transcriptomics in rapid and low ECAR
Saturnispora **species.** (A) A Principal Component Analysis (PCA) showing the similarity of
expression profiles for the four species evaluated. The relative expression of (B) glycolytic
genes and (C) citric acid cycle encoding genes across all four replicates for each species (Sdis
= *Sat. dispora,* Shag = *Sat. hagleri*, Smen = *Sat. mendoncae,* Ssil = *Ssilvae*).
773

Supplemental Figure 4. Unrooted phylogeny of the Gal4p orthologs in the budding yeasts used in this study. (A) A phylogeny including all translated amino acid sequences of genes in the orthogroup containing canonical Gal4p. A black arrow indicates the location of a branch, which was pruned in (B) due to the length of the branch and the topological incongruence with the species phylogeny in this part of the tree, suggesting that these are likely other related C6 zinc transcription factors. The Newick files for these trees are available as Supplemental Data 2 & 3. The location of the Gal4p orthologs from *S. cerevisiae* and *Sat. dispora* are highlighted on both trees and the branches are coloured according to the phylogenetic order.

785
786 **Supplemental Figure 5. The respiration profile of *Sat. dispora* and *gal4Δ* deletion mutants**
787 **over 48 hours of growth**. For each strain, the black line represents the average oxygen
788 consumption rate (OCR), and the shaded area represents the standard error of the mean. Note
789 that both mutants (Gal4_1, Gal4_19) have considerably higher OCR at several phases of the
790 experiment, including the beginning and end.
791
792 **Supplemental Tables:**
793
794 **Table S1.** The species used in this study. For each species used, information about the strain,
795 the phylogenetic order, and naming conventions relating this to the Y1000+ Project genome
796 information (Opulente *et al*, 2024) are provided to improve usability. The data underlying Figure
797 1 including measured extracellular acidification rates, the number of orthologs of each glycolytic
798 gene, and the sum of all of the glycolytic genes are also given for each species.
799
800 **Table S2.** The systematic identification of transcription factor binding sites in the putative
801 promoters of glycolytic genes for the four *Saturnispora* species used for transcriptomics. For
802 each gene, the transcription factor, binding site, and position are noted.
803
804 **Table S3.** The primers used in this study.
805
806 **Table S4.** Genetically modified strains made and used for this study.
807
808 **Supplemental Data:**
809
810 **Supplemental data 1.** Uncropped microscopy images of GFP expressing *Sat. dispora*.
811
812 **Supplemental data 2.** Newick tree file of the full unrooted Gal4p tree.
813
814 **Supplemental data 3.** Newick tree file of the pruned unrooted Gal4p tree.