Magnesium isoglycyrrhizinate alleviates alcohol-associated liver disease through targeting

HSD11B1

Lu Xiao[1,2,†], Lu Li[3,†] , Shasha Wu[4], Zhaoyi Che[1], Yuyang Du[4], Jingyi Zheng[5], Jingsong Yan[4], Hao

Wang[1], Hong Zhang[6], Yan Li[5,*], Jia Xiao[1,7,8,*]

[1]Clinical Medicine Research Institute and Department of Anesthesiology, The First Affiliated

Hospital of Jinan University, Guangzhou 510632, China; [2]Department of Gastroenterology,

People's Hospital of Guangming District, Shenzhen 518107, China; [3]Department of

Gastroenterology, The First Affiliated Hospital of Jinan University, Guangzhou 510632, China

[4]Department of Systems Biology, School of Life Sciences, Southern University of Science and

Technology, Shenzhen 518055, China; [5]Shenzhen Hospital of Southern Medical University,

Shenzhen 518101, China; [6]Department of Interventional Radiology and Vascular Surgery, The

Sixth Affiliated Hospital of Jinan University, Dongguan 523573, China; [7]Guangdong Provincial

Key Laboratory of Tumor Interventional Diagnosis and Treatment, Zhuhai Hospital Affiliated with

Jinan University, Zhuhai 519000, China; [8]Department of Gastroenterology, Qingdao Central

Hospital, University of Health and Rehabilitation Sciences, Qingdao, 266071, China

*For correspondence: edwinsiu@connect.hku.hk (JX), liyan181@smu.edu.cn (Y L)

[†] These authors have contributed equally.

Abbreviations

ALD, alcohol-associated liver disease; ALT, alanine aminotransferase; AST, aspartate

aminotransferase; DEG, differentially expressed gene; DILI, drug-induced liver injury; GL,

generation glycyrrhizin; H&E, hematoxylin and eosin; HSD11B1, hydroxysteroid 11-beta

dehydrogenase 1; IDI1, isopentenyl diphosphate delta isomerase 1; IL-6, interleukin-6;

KEGG, Kyoto Encyclopedia of Genes and Genomes; LDH, lactate dehydrogenase; MASLD,

metabolic dysfunction-associated steatotic liver disease; MgIG, magnesium isoglycyrrhizinate;

MST, microscale thermophoresis; NAS, nonalcoholic fatty liver disease activity score;

NIAAA, National Institute on Alcohol Abuse and Alcoholism; n-Srebp2, nuclear SREBP2;

PA, palmitic acid; PBS, phosphate-buffered saline; p-Srebp2, precursor SREBP2; SREBP2,

sterol regulatory element binding protein 2; TC, total cholesterol; TG, triglyceride; TNF-α,

tumor necrosis factor-alpha; ULM, univariate linear model.

Abstract

While magnesium isoglycyrrhizinate (MgIG) is a clinically approved therapy for alcohol-associated liver disease (ALD), its precise molecular targets and mechanisms remain uncharacterized. This study aimed to define MgIG's hepatoprotective actions in chronic-binge ALD mouse models and ethanol/palmitic acid-exposed AML-12 hepatocytes. Through an integrated strategy encompassing RNA sequencing, molecular docking, and microscale thermophoresis, we discovered that MgIG directly binds to hydroxysteroid 11-beta dehydrogenase 1 (HSD11B1) at residue 187, a finding corroborated by molecular dynamics simulations. *In vivo*, MgIG markedly attenuated alcohol-induced liver injury, evidenced by ameliorated histological damage, reduced hepatic steatosis, and normalized liver-to-body weight ratios. *In vitro*, it effectively reduced lipid accumulation, inflammation, and apoptosis. Mechanistically, RNA sequencing identified isopentenyl diphosphate delta isomerase 1 (IDI1) as a key downstream effector. Hepatocyte-specific genetic manipulations confirmed that MgIG modulates the SREBP2-IDI1 axis, thereby suppressing lipogenesis, inflammatory responses, and apoptotic pathways. We reveal HSD11B1 as a novel direct molecular target of MgIG and elucidate its therapeutic mechanism through the HSD11B1-SREBP2-IDI1 signaling axis, which profoundly impacts ALD pathogenesis. These findings not only validate MgIG's clinical utility but also highlight a promising new therapeutic target for ALD.

Keywords: MgIG, ALD, HSD11B1, SREBP2, Pharmacological intervention

Introduction

Alcohol-associated liver disease (ALD) encompasses a spectrum of hepatic pathologies ranging from steatosis to fibrosis, cirrhosis, and ultimately, hepatocellular carcinoma, contributing significantly to the global burden of disease and premature mortality [1,2]. With alcohol use disorder affecting approximately 283 million individuals worldwide (5.1% of the population), alcohol emerges as a leading cause of cirrhosis, conferring a 260-fold increased risk of liver-related death [3]. Excessive alcohol consumption, the primary etiological factor in ALD, triggers a cascade of pathological events, including oxidative stress, inflammation, and lipid metabolism dysregulation [4]. Current ALD management primarily focuses on alcohol abstinence and supportive care. While potential therapeutic targets, such as oxidative stress pathways, inflammatory cascades, and the gut-liver axis, have been extensively investigated, the identification of druggable hepatic signaling pathways remains elusive [5,6].

Hydroxysteroid 11-beta dehydrogenase 1 (HSD11B1), a cortisol-regenerating enzyme predominantly expressed in the liver and kidneys, elevates endogenous glucocorticoids and has been implicated in the pathogenesis of metabolic dysfunction-associated steatotic liver disease (MASLD) and liver fibrosis [7,8]. Similarly, isopentenyl diphosphate delta isomerase 1 (IDI1), a cytoplasmic enzyme involved in lipid metabolism, has been identified as a key risk factor for hepatocellular carcinoma [9] and intrahepatic cholestasis [10]. Emerging evidence suggests that exercise may ameliorate MASLD by downregulating IDI1 expression [11]. However, the precise roles of HSD11B1 and IDI1 in ALD progression remain to be elucidated.

Magnesium isoglycyrrhizinate (MgIG), a fourth-generation glycyrrhizin (GL) derivative and a magnesium salt of a single stereoisomer (Fig. 1A), exhibits enhanced pharmacological properties compared to its predecessors. As an 18α-GL, MgIG possesses greater lipophilicity than the β-isomer, facilitating superior binding to target cell receptors and steroid hormones [12]. This enhanced binding capacity translates to potent anti-inflammatory effects, hepatoprotective properties, and improved liver function [13]. MgIG has demonstrated therapeutic efficacy in various liver diseases, including drug-induced liver injury (DILI) [14], MASLD [15], and liver fibrosis [16]. Notably, MgIG has been preliminarily shown to inhibit ethanol-induced activation of the hedgehog signaling pathway in hepatocytes [17] and attenuate acute alcohol-induced hepatic

steatosis in a zebrafish model by modulating lipid metabolism-related gene expression [18].

Furthermore, MgIG has been recognized by the Chinese Society of Hepatology for its clinical

utility in ALD treatment [19]. Despite its therapeutic promise and proven safety, the precise

molecular targets of MgIG in the liver remain poorly defined, hindering the development of

optimized treatment strategies for ALD. This study employed *in vivo* and *in vitro* ALD models to

investigate the hepatoprotective mechanisms of MgIG, revealing its direct interaction with

HSD11B1 as a key mediator of its therapeutic effects and the involvement of downstream

signaling of the sterol regulatory element binding protein 2 (SREBP2)-IDI1 axis.

Methods

Chemicals and reagents

MgIG was generously provided by Chia-Tai Tianqing Pharmaceutical Group Co., Ltd. (Nanjing,

China) in the form of a magnesium isoglycyrrhizinate injection (5 mg/ml). High-purity ethanol

was obtained from MACKLIN (Shanghai, China). Palmitic acid (PA) and fatty acid-free bovine

serum albumin (BSA) were purchased from Sigma-Aldrich (St. Louis, MO, USA). All cell culture

reagents and consumables were obtained from Gibco (Carlsbad, CA, USA) or Corning

Incorporated (Corning, NY, USA). Primary antibodies against tumor necrosis factor-α (TNF-α,

#11948), interleukin-6 (IL-6, #12912), glyceraldehyde-3-phosphate dehydrogenase (GAPDH,

#2118S), Bax (#2772S), Bcl-2 (#3498S), and hemagglutinin (HA, #3724S) were purchased from

Cell Signaling Technology (Shanghai, China). Antibodies against IDI1 (#11166-2-AP) and

SREBP2 (#28212-1-AP) were obtained from Proteintech (Wuhan, China). An antibody against

HSD11B1 (#AF3397) was purchased from R&D Systems (Minneapolis, MN, USA). A Flag

antibody (F1804) was purchased from Sigma-Aldrich. All antibody validation experiments were

performed according to the manufacturers' protocols. Protein A/G Magarose beads (#80105G)

were purchased from Invitrogen (Carlsbad, CA, USA).

Animal experiments

Male C57BL/6 wild-type mice (7 weeks old; 19-21 g) were purchased from Guangzhou Qingle

Biosciences Co., Ltd. (Guangzhou, China). Mice were acclimated to the housing environment for

one week prior to experimental procedures. Alcoholic liver injury was induced using a modified

National Institute on Alcohol Abuse and Alcoholism (NIAAA) model [20]. Briefly, mice (n = 4 per

group) were initially fed *ad libitum* with the control (Ctrl) Lieber-DeCarli liquid diet for 5 days to

facilitate acclimation. Subsequently, the ALD groups (designated as 'A-Ctrl' and 'MgIG' in figures)

received the Lieber-DeCarli diet containing 5% (v/v) ethanol for 10 days, while the control group

(designated as 'Ctrl' in figures) received an isocaloric control diet. Body weight and food intake

were monitored daily. On day 11, ethanol-fed mice and their pair-fed controls were administered a

single dose of ethanol (5 g/kg body weight) or an isocaloric maltose dextrin solution, respectively,

via oral gavage in the early morning. Mice were euthanized 9 h later (Fig. 1B). A vehicle-MgIG

control group was not included in this study because: (1) MgIG is a clinically approved drug with

an established safety profile; (2) the dosage used in this study was calculated based on human

clinical dosages using a standard conversion formula; and (3) the vehicle safety of MgIG with

similar dose has been previously confirmed in an animal study [21]. For viral infections, mice (n = 5

per group) received tail vein injections of 5×10^{11} genome copies of either AAV8 control or

AAV-shRNA/plasmid. After 14 days, mice were fasted for 4 h at the end of the dark cycle and then

euthanized to confirm hepatic upregulation or downregulation of *Hsd11b1*, *Srebp2*, or *Idi1*. These

mice were then subjected to the NIAAA model and/or MgIG administration as described above.

Target gene expression in the liver was verified using quantitative PCR and Western blotting. All

experimental procedures were approved by the Ethical Committee of Jinan University, China

(IACUC-20220708-04).

RNA sequencing and functional enrichment analysis

Total RNA was extracted from liver tissue samples using TRIzol reagent (Invitrogen) following

the manufacturer's instructions. RNA quality was assessed using a Nanodrop ONE

spectrophotometer (Thermo Fisher, Waltham, MA, USA). Samples with A260/A280 ratios above

1.8 and A260/A230 ratios above 2.0 were deemed acceptable for further processing. RNA

integrity was evaluated using an Agilent 2200 TapeStation (Agilent Technologies, Santa Clara, CA,

USA), and only samples with an RNA integrity number above 7.0 were used for library

preparation. Ribosomal RNA (rRNA) was depleted from total RNA using the Epicentre Ribo-Zero

rRNA Removal Kit (Illumina) and fragmented to an average size of 200 bp. First-strand and

second-strand cDNA synthesis, followed by adapter ligation and low-cycle enrichment, were

performed using the NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs,

Ipswich, MA, USA). Library quality was assessed using the Agilent 2200 TapeStation and Qubit

2.0 fluorometer (Thermo Fisher). Libraries were diluted to 10 pmol/L and subjected to cluster

generation on a pair-end flow cell, followed by sequencing (2×150 bp) on an Illumina HiSeq

3000. Raw sequencing reads were processed to remove adapter sequences, low-quality reads, and

reads containing poly-N stretches. Clean reads were aligned to the mouse reference genome using

HISAT2 with default parameters. Gene expression levels were quantified using HTSeq, and

differential expression analysis was performed using DESeq with read counts as input. The

Benjamini-Hochberg method was used to adjust p-values for multiple testing. Differentially

expressed genes (DEGs) were defined as those with a fold change > 2 and an adjusted p-value $<$

0.05. Heatmap visualization and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway

enrichment analysis were performed using the identified DEGs. A p-value < 0.05 was considered

statistically significant for KEGG pathway enrichment.

Biochemical and cytokine analyses of mice serum and cell supernatants

Serum levels of alanine aminotransferase (ALT), aspartate aminotransferase (AST), triglyceride

(TG) and total cholesterol (TC) were measured using BK2800 (BIOBASE, Jinnan, China). Cell

supernatants cytokine levels were determined by using corresponding ELISA kits from R&D

Systems (TNF-α: #VAL609; IL-6: #VAL604G).

Liver tissue histology

Liver tissue samples were fixed in 10% neutral buffered formalin and subsequently embedded in

paraffin. Tissue sections (5 μm thickness) were prepared and stained with hematoxylin and eosin

(H&E) and Oil Red O for histological examination. Images were acquired using a LEICA Qwin

Image Analyzer (Leica Microsystems, Milton Keynes, UK). The nonalcoholic fatty liver disease

activity score (NAS) was determined for each group as previously described [22].

Cell culture and transfection

The AML-12 mouse normal hepatocyte cell line was obtained from the Cell Bank of Type Culture Collection, Chinese Academy of Sciences (Shanghai, China). Cells were cultured in Dulbecco's modified Eagle's medium supplemented with 10% (v/v) fetal bovine serum, incubated at 37 °C with 5% CO_2 in a cell incubator. To induce an ALD-like phenotype in AML-12 cells, cells were exposed to a combination of 25 mmol/L ethanol and 0.1-1 mmol/L palmitic acid (PA; Sigma-Aldrich; #P0500) for 24 h to determine the optimal concentration for inducing cellular injury, as previously described (12). We chose the combination of ethanol and palmitic acid in the *in vitro* experiments because ethanol-only treatment could not phenocopy alcoholic cell death, inflammation, oxidative stress, and the dysregulated lipid metabolism in hepatocytes exhibited in early-stage ALD [23-26]. Following this initial treatment, varying concentrations of MgIG were added to the culture medium for an additional 24-hour to assess its effects and determine the optimal dose. For gene knockdown and overexpression experiments, Lipofectamine 3000 reagent (Thermo Fisher; #L3000001) was used to transfect AML-12 cells with *Hsd11b1* or *Idi1*-targeting constructs in 6-well plates.

Cell lipid accumulation, viability, and apoptosis assays

Intracellular lipid droplets were visualized by staining AML-12 cells with Nile Red. Briefly, cells were washed with phosphate-buffered saline (PBS), fixed with 4% formaldehyde for 10 min, and then stained with Nile Red solution (0.1 mg/mL; #N8440, Solarbio, Beijing, China) for 10 min at room temperature in the dark. Cells were then counterstained with DAPI (5 μg/mL; Sigma-Aldrich; D9542) for 5 min. Cell viability and apoptosis were assessed in 96-well plates using a CCK8 cell viability assay and a lactate dehydrogenase (LDH) cytotoxicity assay kit, respectively, according to the manufacturer's instructions (MedChem Excess, Shanghai, China; HY-K1090 and HY-K0301).

Quantitative PCR

Total RNA was extracted from liver tissue or cultured cells using TRIzol reagent. The first strand cDNA was synthesized from total RNA using a PrimeScript RT Reagent Kit (Takara, Shiga, Japan;

#RR047A) following the manufacturer's instructions. Quantitative PCR reaction was performed

with the SYBR Premix Taq Quantitative PCR Kit (Takara) according to the manufacturer's

instructions, on a qTOWER3 machine (Analytik Jena AG, Jena, Germany). Primer information is

as listed in Supporting Information Table S1. β-actin was used as an internal control. The relative

quantification of mRNA expression levels was calculated using the $2^{-\Delta\Delta Ct}$ method. All qPCR

experiments were conducted in compliance with the MIQE guidelines [27].

Protein extraction and Western blot

Samples were lysed using Pierce RIPA buffer (Thermo Fisher) and their protein concentrations

were measured with the Bio-Rad Protein Assay (Bio-Rad, Hercules, CA). Western blot analyses of

all proteins were conducted with β-actin serving as the internal control.

Coimmunoprecipitation (Co-IP)

Co-IP buffer (50 mM Tris-HCl, 5 mM EDTA, 150 mM NaCl, and 1% NP-40 pH 7.6) mixed with

protease inhibitor cocktail (Roche, Bern, Switzerland) were used to lyse the cells. The samples

were then incubated with the corresponding antibodies and Protein A/G Magarose beads at 4°C

overnight.

Protein structure modeling and molecular docking

We first employed DecoupleR's [28] built-in Univariate Linear Model (ULM) to compute the

transcription factor activity perturbed by MgIG, based on the expression of differentially

expressed genes. The findings revealed that *Srebf2* was the transcription factor most significantly

reduced by MgIG. We then identified potential SREBP2/IDI1-interacting proteins using the

STRING and PubMed databases. The 3D structures of these proteins were obtained from the

Protein Data Bank (PDB) or UniProt, while the 3D structure of MgIG was retrieved from

PubChem. Molecular docking simulations were performed using Schrödinger software to predict

the binding affinity (Glide score) of MgIG to each protein. The lowest-energy docking

conformations were visualized using PyMOL.

Microscale thermophoresis (MST) assay

The binding interaction between MgIG and HSD11B1 was assessed using a Monolith NT.115 Blue/Green instrument (NanoTemper Technologies, Munich, Germany). Initially, we evaluated the affinity and labeling efficiency of the dye with the His-tagged protein. Subsequently, HSD11B1 proteins (100 nM) were labeled and incubated with MgIG at a concentration gradient ranging from 50 μM to 1.53 nM for 30 min at room temperature. The samples were then analyzed at 60% MST power and a temperature of 25°C. Dissociation constants (Kd) were calculated based on a 1:1 binding stoichiometry.

Molecular dynamics (MD) simulations

Three molecular systems were constructed in this study: first, Compound-HSD11B complex was established through molecular docking. Second, the last frame from 100 ns MD simulation was used to generate APO-HSD11B1 by removing the ligand. Finally, based on this conformation and reported portal vein ethanol concentrations in ALD, five ethanol molecules (0.1 g/dL) were incorporated to construct the EtOH-HSD11B1 system, simulating the ALD environment [29-31]. For each system, receptor was prepared and missing atoms of residues were fixed by the advanced PDB-Preparation tool in Yinfo Cloud Computing Platform using PDBFixer and the tLEaP module in AmberTools 20. The AM1-BCC charges were calculated for compound magnesium isoglycyrrhizinate by the Amber antechamber program [32]. MD simulations was performed using AmberTools 20 package with AMBER ff19SB [33] and GAFF [34] force field.

The system was solvated by a truncated octahedron water box using OPC water model with a margin of 10 Å. Periodic boundary condition (PBC) was used and the net charge neutralized by Na^+ ions. Nonbonded van der Waals interactions were calculated using the Lennard-Jones 12-6 potentials with a 10 Å cutoff, while long-range electrostatics were treated using the Particle Mesh Ewald (PME) algorithm. The SHAKE algorithm was applied to constrain bonds involving hydrogen atoms [35]. To removed improper atom contacts, the structure was first minimized by (1) 2500 steps of steepest descent and 2500 steps of conjugate gradient, under a harmonic constraint of 10.0 kcal/(mol·Å2) on heavy atoms; (2) relaxing the entire system by 10000 steps of steepest descent and 10000 steps of conjugate gradient. And then the system was gradually heated up to

300 K by a 20 ps NVT simulation. Subsequently, two steps of equilibration phases were carried

out: (1) a 200 ps NPT simulation with constraints on heavy atoms followed by (2) a 1 ns NVT

simulation without restraint. The temperature was maintained at 300 K using the Berendsen

thermostat with 1 ps coupling constant and the pressure at 1 atm using Monte Carlo barostat with

1 ps relaxation time. Finally, the system was subjected to a 100 ns NVT simulation with a time

step of 2 fs. The root-mean-square deviation (RMSD), root-mean-square fluctuation (RMSF),

radius of gyration (RG), solvent-accessible surface area (SASA) and hydrogen bonds were

analyzed by the CPPTRAJ module, while principal component analysis (PCA) and dynamic

cross-correlation matrix (DCCM) were analyzed by R package Bio3D [36]. The binding free

energies were calculated using the Molecular Mechanics Generalized Born (MM/PBSA) method

for the 100 ns MD trajectory [37].

HSD11B1 enzyme activity assays

The enzyme activity of HSD11B1 in cells was evaluated according to its capability to convert

cortisone to cortisol [38]. AML-12 cells were seeded into 6-well culture plates and induced ALD as

above. Different concentrations of MgIG (0.1, 0.25, and 0.5 mg/mL) were added 24 h prior to the

co-treatment, with a control group included. AML-12 cells were exposed to 160 nmol/L cortisone

for 24 h. The reaction mixtures were collected and analyzed for cortisol levels using an ELISA kit

(R&D Systems, Minneapolis, MN, USA), according to the manufacturer's instructions.

Statistical analysis

The data from each group are presented as the mean ± standard deviation (SD). For data that

followed a normal distribution, statistical comparisons between two groups were conducted using

an unpaired two-tailed Student's t-test; for comparisons involving three or more groups, a two-way

ANOVA followed by a Student-Newman-Keuls post hoc test was applied (Prism 5.0, Graphpad

Software, Inc., San Diego, CA). A P-value of less than 0.05 was deemed statistically significant.

Results

MgIG alleviates liver injury in a mouse model of ALD

To investigate the hepatoprotective effects of MgIG *in vivo*, we employed a well-established

chronic-binge NIAAA mouse model of ALD (Fig. 1B). Based on the drug's clinical instructions,

previous studies [21,39], and our preliminary experimental results (Fig. S1), we determined an

optimal MgIG dose of 50 mg/kg. Compared to mice fed a normal diet (Ctrl group), mice in the

alcohol-fed control group (A-Ctrl) exhibited characteristic histological features of ALD, including

steatosis and inflammatory cell infiltration, resulting in an elevated NAS. MgIG treatment

effectively attenuated these pathological changes (Fig. 1C and 1D). Consistent with these

histological findings, the liver-to-body weight ratio, a marker of liver injury, was significantly

increased in the A-Ctrl group, and this increase was markedly reduced by MgIG treatment (Fig.

1E). Notably, serum levels of liver enzymes (ALT and AST) and triglycerides (TG), indicators of

liver function and lipid metabolism, were significantly different between the A-Ctrl and

MgIG-treated groups (Fig. 1F). In line with the observed histological and physiological

improvements, MgIG treatment also modulated the expression of genes involved in lipid

metabolism (*Srebp-1c*, *Srebp2*, *Acc1*, and *Scd1*), inflammation (*Tnf-α* and *Il-6*), and apoptosis

(*Bax* and *Bcl2*) in the liver tissue of A-Ctrl mice (Fig. 1G).

MgIG protects against ethanol-induced hepatocyte injury via IDI1

To further elucidate the molecular mechanisms underlying the hepatoprotective effects of MgIG,

we performed RNA sequencing analysis on liver tissue from the NIAAA mouse model. DEG

analysis revealed a significant upregulation of *Idi1* in the A-Ctrl group compared to the Ctrl group

(Fig. 2A). Furthermore, KEGG pathway enrichment analysis identified metabolic pathways

among the top five downregulated pathways in the MgIG-treated group compared to the A-Ctrl

group, with *Idi1* consistently ranked among the top two genes in these pathways (Fig. 2C). To

further investigate the role of IDI1 and its potential interaction with MgIG, we established an *in*

vitro model of ethanol/palmitic acid (EtOH/PA)-induced hepatocyte injury using AML-12 cells.

We first optimized the EtOH/PA concentrations to induce significant cell death and lipid

accumulation (Fig. S2), identifying 250 mmol/L ethanol and 0.2 mmol/L PA as the optimal

combination. MgIG treatment (0.1-1.0 mg/mL) demonstrated a dose-dependent protective effect

against EtOH/PA-induced injury, as evidenced by improved cell viability, reduced lipid

accumulation, and decreased apoptosis (Fig. 2D and 2E). Consistent with the *in vivo* findings,

EtOH/PA treatment increased the expression of genes involved in lipid metabolism (*Acc1* and

Scd1) and inflammation (*Tnf-α* and *Il-6*), and MgIG treatment effectively reversed these effects

(Fig. 2F). Based on these results, we selected 0.25 mg/mL as the optimal MgIG concentration for

subsequent *in vitro* experiments.

In line with the *in vivo* observations, *Idi1* expression was significantly increased in AML-12

cells following EtOH/PA treatment, and this effect was attenuated by MgIG treatment (Fig. 3A and

S3A-B). As shown in Fig. 3B and 3C, transfection with siRNA and plasmid effectively decreased

or increased *Idi1* mRNA levels without affecting cell viability, respectively. Knockdown of *Idi1*

significantly ameliorated EtOH/PA-induced cell injury, reducing apoptosis, inflammatory cytokine

levels, and lipid accumulation, while *Idi1* overexpression exacerbated these effects (Fig. 3D-3G).

Taken together, these findings suggest that MgIG protects against ethanol-induced hepatocyte

injury, at least in part, through the modulation of IDI1.

MgIG directly binds to HSD11B1

To identify potential direct binding partners of MgIG, we initially performed molecular docking

simulations between MgIG and IDI1. However, the resulting Glide score of -2.74 indicated a lack

of favorable binding interactions, which implied that there should be an immediate binding protein

in the upstream of IDI1. To identify potential upstream regulators of *Idi1* expression, we utilized

the ULM within the DecoupleR package to analyze transcription factor activity. This analysis

identified SREBP2 as the most significantly affected transcription factor (Fig. 4A). We then

expanded our search to include upstream regulators of IDI1 identified using the STRING database.

Among these proteins, HSD11B1 exhibited the highest binding affinity for MgIG, with a Glide

score of -8.75 (Fig. 4B). Given the homology between human and mouse *Hsd11b1* and the need

for future translational research, we focused on validating the interaction between human

HSD11B1 and MgIG. We conducted 100 ns MD simulations for Compound-HSD11B1

(MgIG-HSD11B1 binding in physiological condition) and EtOH-HSD11B1 (MgIG-HSD11B1

binding in alcoholic condition), with free HSD11B1 as a control. The results revealed that

HSD11B1 exhibited high flexibility and dynamic motion in its unbound state but became

significantly more rigid and stable upon binding to MgIG, particularly at residues 219–233 (Fig.

4C-4E). This structural stabilization was further supported by the radius of gyration curves, which

indicated a more compact conformation upon binding (Fig. S4A and SB). Further analyses,

including PCA and DCCM, demonstrated alterations in motion direction, dynamic patterns, and

residue correlations. Specifically, upon binding, HSD11B1 exhibited reduced motion amplitude

and weaker residue correlations, with its dynamic patterns becoming more disordered (Fig.

S4C-4E). In the Compound-HSD11B1 system (Fig. 4E), key interactions included hydrogen

bonds (Tyr177, Tyr183), a salt bridge (Lys187) and hydrophobic interactions (Leu126, Leu217,

Ala223, Ala226, and Ile230). In contrast, the EtOH-HSD11B1 system (Fig. 4F) exhibited

hydrogen bonds (Thr124, Ser170, Tyr183, Leu217, and Arg269) along with an additional salt

bridge (Tyr183 and Arg269). Notably, the EtOH-HSD11B1 system demonstrated stronger

hydrogen bonding and more favorable binding free energy compared to the Compound-HSD11B1

system (Fig. 4G and S4F), suggesting enhanced stability and binding affinity in an ethanol

environment. Based on molecular docking and MD results, we further validated the binding

between MgIG and the human full-length wild-type (WT) HSD11B1 protein as well as its

site-directed mutants (Tyr177, M1; Tyr183, M2; Lys187, M3) using MST. The Kd value of the

WT protein was 6.35 μM (Fig. 4H), while those of M1, M2, and M3 were 48.6 μM, 11.5 μM, and

135.6 μM, respectively (Fig. 4I). These data indicate that MgIG primarily exerts its anti-ALD

effect by inhibiting HSD11B1 through the critical Lys187 site.

To further investigate the functional significance of this interaction, we modulated *Hsd11b1*

expression in AML-12 cells using siRNA- and plasmid-mediated knockdown and overexpression,

respectively. As shown in Fig. 4A-4C, transfection with siRNA and plasmid effectively decreased

or increased *Hsd11b1* mRNA levels without affecting cell viability, respectively. Knockdown of

Hsd11b1 significantly attenuated EtOH/PA-induced inflammation, apoptosis, and lipid

accumulation, even in the absence of MgIG treatment. Conversely, overexpression of *Hsd11b1*

exacerbated these effects. Importantly, MgIG treatment partially reversed the detrimental effects

of *Hsd11b1* overexpression (Fig. 5D-5F). To further investigate whether the binding of MgIG to

HSD11B1 affects its enzymatic activity, we added different concentrations of MgIG (0.1, 0.25,

and 0.5 mg/mL) to the cellular model. ELISA results showed that HSD11B1 activity was

significantly elevated in the ALD group, whereas increasing concentrations of MgIG led to a

dose-dependent reduction in HSD11B1 activity (Fig. 5G). Considering the quite low basal

expression of *Hsd11b1* in normal AML-12 cells, we further examined cortisol levels in the

supernatant of AML-12 cells overexpressing *Hsd11b1*, with or without MgIG co-treatment (Fig.

5H). These findings suggested that MgIG exerted its protective effects against EtOH/PA-induced

hepatocyte injury, at least in part, by directly binding to and modulating the activity of HSD11B1.

MgIG protects hepatocyte partially through the HSD11B1–SREBP2–IDI1 axis

To investigate the relationship between SREBP2 and HSD11B1, we conducted co-IP experiments,

which revealed a direct interaction between the precursor form of SREBP2 (p-SREBP2) and

HSD11B1. This interaction was weakened following MgIG treatment (Fig. 6A and 6B). To further

explore this interaction, we modulated *Hsd11b1* expression in EtOH/PA induced AML-12 cells

with or without MgIG treatment (Fig. 6C and 6D). Knockdown of *Hsd11b1* resulted in an

accumulation of p-SREBP2, while *Hsd11b1* overexpression increased the levels of nuclear

SREBP2 (n-SREBP2), an effect further enhanced by the EtOH/PA treatment. These findings

suggest that HSD11B1 may regulate the processing and nuclear translocation of SREBP2.

Immunofluorescent analysis confirmed these observations (Fig. 6E). As shown in Fig. 6F, the

co-IP results indicated that HSD11B1 also interacted with its downstream protein IDI1. In addition,

the dual-luciferase reporter assay demonstrated that *Srebp2* positively regulates *Idi1* expression

(Fig. 6G). In *in vivo* experiments, it was confirmed that in normal liver tissue, the expression of

Srebp2 and *Idi1* followed a similar trend in response to overexpression or knockdown of *Hsd11b1*

(Fig. 6H and 6I). Knockdown of *Srebp2* resulted in a decrease in *Idi1* expression, without

affecting *Hsd11b1* levels. Conversely, neither knockdown nor overexpression of *Idi1* influenced

the expression of the upstream genes *Hsd11b1* or *Srebp2*. In liver tissue from ALD, the expression

changes of upstream and downstream genes upon overexpression or knockdown of *Hsd11b1* or

Idi1 mirrored those seen in normal liver tissue. However, *Srebp2* knockdown induced a negative

feedback effect, reducing *Hsd11b1* expression, which was not observed in normal liver tissue (Fig.

6J and 6K).

To further investigate the roles of HSD11B1, SREBP2, and IDI1 in the *in vivo* ALD model, we

used AAV8-mediated shRNA delivery to knock down their expression or employed

overexpression plasmids to increase their expression specifically in hepatocytes. As shown in Fig.

S6A-6C, knockdown or overexpression of *Hsd11b1* or *Idi1* in mice fed a normal control diet did

not result in significant changes in liver histology, lipid accumulation, or serum ALT and AST

levels. However, in the ALD model, knockdown of *Hsd11b1*, *Srebp2*, or *Idi1* attenuated

ethanol-induced liver injury, as evidenced by improved histological NAS scores, reduced steatosis,

and decreased inflammation. MgIG treatment further enhanced these beneficial effects.

Conversely, overexpression of *Hsd11b1* or *Idi1* abolished the therapeutic effects of MgIG on

ethanol-induced liver injury (Fig. 7A-7F and S7). Interestingly, modulation of *Idi1* expression did

not affect the expression of *Hsd11b1* or *Srebp2*, suggesting that IDI1 acts downstream of

HSD11B1 and SREBP2. In contrast, modulation of *Hsd11b1* expression altered the levels of

n-SREBP2 and IDI1 (Fig. S7E-G), confirming that HSD11B1 regulates the SREBP2-IDI1

pathway. Collectively, these data indicate that MgIG protects against ALD, at least in part, by

modulating the HSD11B1-SREBP2-IDI1 axis.

Discussion

While previous research using a zebrafish model demonstrated that MgIG alleviates

alcohol-induced liver injury by modulating lipid metabolism-related gene expression [18], the

precise molecular mechanisms and direct targets of MgIG remained unclear. This lack of

mechanistic understanding hindered the development of optimized therapeutic strategies for ALD.

In this study, we provide compelling evidence that MgIG effectively ameliorates key features of

ALD, including hepatic steatosis, inflammation, and apoptosis, in both *in vivo* and *in vitro* models.

Our findings shed light on the complex interplay of molecular pathways involved in ALD

pathogenesis and highlight the therapeutic potential of targeting the HSD11B1-SREBP2-IDI1

axis.

Our RNA sequencing analysis identified IDI1 as a key player in ALD pathogenesis and a

critical mediator of MgIG's hepatoprotective effects. IDI1, an enzyme in the mevalonate pathway,

is involved in the biosynthesis of isoprenoids, which are essential for various cellular processes,

including cholesterol synthesis. Dysregulation of cholesterol metabolism is a hallmark of ALD,

contributing to the accumulation of lipids in the liver. Our findings demonstrate that *Idi1*

expression is significantly upregulated in the livers of mice subjected to the NIAAA model of

ALD, suggesting that increased IDI1 activity may contribute to the development of hepatic

steatosis. Genetic manipulation of *Idi1* expression further confirmed its role in ALD. Specifically,

Idi1 knockdown protected against alcohol-induced liver injury, an effect further enhanced by

MgIG treatment. This suggests that reducing IDI1 activity can ameliorate the detrimental effects

of alcohol on the liver and that MgIG may exert its therapeutic benefits, at least in part, by

suppressing IDI1. Conversely, *Idi1* overexpression abolished the therapeutic benefits of MgIG,

both *in vivo* and *in vitro*, highlighting the importance of IDI1 in mediating MgIG's effects. These

findings are consistent with recent reports highlighting the role of IDI1 in hepatic cholesterol

accumulation and lipid toxicity [40][41] and its involvement in hepatocellular carcinoma progression

via inhibition of the cGAS-STING pathway [9]. Furthermore, our study identified SREBP2, a

master regulator of lipid metabolism, as an upstream regulator of IDI1. SREBP2 belongs to a

family of transcription factors that control the expression of genes involved in cholesterol and

fatty acid biosynthesis. SREBP2 has been implicated in various aspects of liver pathology,

including NLRP3 inflammasome activation [42], lipid accumulation [43], liver cancer progression [44],

with its target protein IDI1 upregulated in parallel [30]. Our findings demonstrate that both SREBP2

and IDI1 are upregulated in response to EtOH/PA treatment and that MgIG effectively inhibits

their expression, consistent with the notion that SREBP2 contributes to the development of

MASLD [45]. The identification of SREBP2 as a key regulator of IDI1 in the context of ALD

provides a mechanistic link between lipid metabolism and the pathogenesis of this disease.

To identify the direct molecular target of MgIG, we employed a combination of molecular

docking, MD, MST analysis, and genetical manipulation studies. These approaches converged on

HSD11B1, a key enzyme in glucocorticoid metabolism, as a high-affinity binding partner of

MgIG. HSD11B1 catalyzes the conversion of inactive cortisone to active cortisol in the liver,

thereby increasing intracellular glucocorticoid levels. Glucocorticoids are steroid hormones that

play a critical role in regulating glucose and lipid metabolism, and their dysregulation has been

implicated in the development of metabolic disorders, including type 2 diabetes [46]. As a potential

inhibitor of HSD11B1 [47], GL can improve patients' insulin and lipid levels [48]. In addition, the

increased expression of *Hsd11b1* enhances the enzymatic reducing activity, promoting a

fat-forming phenotype in cell models through elevated cortisol levels [49,50]. Meanwhile, *Hsd11b1*

deficiency prevents hepatic steatosis in high-fat diets-fed mice [51]. Our findings demonstrate that

MgIG directly binds to the Lys187 of HSD11B1 and that modulation of *Hsd11b1* expression in

hepatocytes significantly influences the development of ALD.

While some studies have suggested a potential pro-fibrotic role for HSD11B1 inhibition [8], our

findings, along with those of other recent studies [7,51,52], support a protective role for *Hsd11b1*

knockdown in the context of alcoholic liver injury. These discrepancies may be attributed to

differences in experimental models and the specific cell types targeted. For instance, the study by

Zou *et al* [8] examined the effects of global *Hsd11b1* deficiency, which could trigger compensatory

activation of the hypothalamic-pituitary-adrenal axis or paracrine regulatory mechanisms,

potentially affecting the pro-fibrotic response. In contrast, our study focused on

hepatocyte-specific *Hsd11b1* knockdown, which may avoid these confounding factors.

Importantly, our study provides evidence for a novel HSD11B1-SREBP2-IDI1 axis in the

pathogenesis of ALD. We demonstrate that HSD11B1 regulates the processing and nuclear

translocation of SREBP2, which in turn controls the expression of IDI1. Notably, this is consistent

with previous studies showing that increased SREBP2 expression and nuclear translocation

activate the NLRP3 inflammasome and the mevalonate pathway, thereby promoting inflammation

and tumor progression [53,54]. This regulatory axis appears to be a key target of MgIG's therapeutic

effects. By directly binding to HSD11B1, MgIG may inhibit its activity, leading to reduced

SREBP2 activation and subsequent downregulation of IDI1. This, in turn, could contribute to the

observed improvements in hepatic steatosis, inflammation, and apoptosis [55-57].

This study provides mechanistic evidence that MgIG directly targets HSD11B1 and modulates

the HSD11B1−SREBP2−IDI1 axis to attenuate ALD. However, several limitations should be

acknowledged. First, our findings are derived from preclinical mouse models and hepatocyte

systems, and pharmacokinetics, biodistribution, and long-term safety of MgIG remain to be

comprehensively evaluated in larger and more diverse populations. Second, although molecular

docking and thermophoresis confirmed direct binding of MgIG to HSD11B1, high-resolution

structural studies such as X-ray crystallography or cryo-electron microscopy are needed to

delineate the precise binding interface and conformational changes. Third, other signaling

pathways potentially influenced by MgIG have not been systematically explored and may

contribute to its therapeutic effects. Future work should focus on validating these findings in

human tissues, optimizing MgIG dosing strategies, and conducting multicenter clinical trials to

assess its translational potential and establish HSD11B1 as a druggable target for precision therapy

in ALD.

Acknowledgements

JX is supported by National Natural Science Foundation of China (U23A20401) and Science

and Technology Projects in Guangzhou (202201020066). YL is supported by National Natural

Science Foundation of China (82372768). HZ is supported by Medical Research Fund of The

Six Affiliated Hospital of Jinan University.

Authorcontributions

Jia Xiao: Supervision, Conceptualization, Funding acquisition, Writing– review & editing, Data curation. **Yan Li**: Funding acquisition, Conceptualization, Data curation, Writing– review & editing. **Lu Xiao**: Writing original draft, Methodology, Data curation, Visualization. **Lu Li**: Writing original draft, Methodology, Data curation, Visualization. **Shasha Wu**: Data curation, Methodology. **Zhaoyi Che**: Data curation, Methodology. **Yuyang Du**: Data curation, Methodology. **Jingyi Zheng**: Data curation, Methodology. **Jingsong Yan**: Data curation. **Hua Wang:** Wrote and revised the manuscript. **Hong Zhang:** Funding acquisition, Wrote and revised the manuscript.

Data availability

All data is available in the manuscript or the supplementary materials. RNA-seq data from

this study is available from: GSE284311. All materials generated as part of this study will be

made available upon request to the corresponding authors.

Declaration ofinterests

The authors declare no competing interests.

REFERENCES

1.   Aberg, F., Jiang, Z.G., Cortez-Pinto, H., and Mannisto, V. (2024). Alcohol-associated liver disease-Global epidemiology. Hepatology. https://doi.org/10.1097/HEP.0000000000000899.

2.   Gan, C., Yuan, Y., Shen, H., Gao, J., Kong, X., Che, Z., Guo, Y., Wang, H., Dong, E., and Xiao,
J. (2025). Liver diseases: epidemiology, causes, trends and predictions. Signal Transduct
Target Ther 10, 33. https://doi.org/10.1038/s41392-024-02072-z.

3.   Devarbhavi, H., Asrani, S.K., Arab, J.P., Nartey, Y.A., Pose, E., and Kamath, P.S. (2023).
Global burden of liver disease: 2023 update. J Hepatol 79, 516-537.
https://doi.org/10.1016/j.jhep.2023.03.017.

4.   Díaz, L.A., Arab, J.P., Louvet, A., Bataller, R., and Arrese, M. (2023). The intersection
between alcohol-related liver disease and nonalcoholic fatty liver disease. Nat Rev
Gastroenterol Hepatol 20, 764-783. https://doi.org/10.1038/s41575-023-00822-y.

5.   Wu, X., Fan, X., Miyata, T., Kim, A., Cajigas-Du Ross, C.K., Ray, S., Huang, E., Taiwo, M.,
Arya, R., Wu, J. et al. (2023). Recent Advances in Understanding of Pathogenesis of
Alcohol-Associated Liver Disease. Annu Rev Pathol 18, 411-438.
https://doi.org/10.1146/annurev-pathmechdis-031521-030435.

6.   Gao, H., Jiang, Y., Zeng, G., Huda, N., Thoudam, T., Yang, Z., Liangpunsakul, S., and Ma, J.
(2024). Cell-to-cell and organ-to-organ crosstalk in the pathogenesis of alcohol-associated
liver disease. eGastroenterology 2. https://doi.org/10.1136/egastro-2024-100104.

7.   Li, C., Li, M., Sheng, W., Zhou, W., Zhang, Z., Ji, G., and Zhang, L. (2024). High dietary
Fructose Drives Metabolic Dysfunction-Associated Steatotic Liver Disease via Activating
ubiquitin-specific peptidase 2/11beta-hydroxysteroid dehydrogenase type 1 Pathway in Mice.
Int J Biol Sci 20, 3480-3496. https://doi.org/10.7150/ijbs.97309.

8.   Zou, X., Ramachandran, P., Kendall, T.J., Pellicoro, A., Dora, E., Aucott, R.L., Manwani, K.,
Man, T.Y., Chapman, K.E., Henderson, N.C. et al. (2018). 11Beta-hydroxysteroid
dehydrogenase-1 deficiency or inhibition enhances hepatic myofibroblast activation in murine
liver fibrosis. Hepatology 67, 2167-2181. https://doi.org/10.1002/hep.29734.

9.   Fu, L., Ding, H., Bai, Y., Cheng, L., Hu, S., and Guo, Q. (2024). IDI1 inhibits the cGAS-Sting
signaling pathway in hepatocellular carcinoma. Heliyon 10.
https://doi.org/10.1016/j.heliyon.2024.e27205.

10.   Yan, J.B., Nie, Y.M., Chen, Z., Yao, J.M., Zhang, S., and Chen, Z.Y. (2023). The
IDI1/SREBP2 axis drives intrahepatic cholestasis and is a treatment target of
San-Huang-Cai-Zhu formula identified by sequencing and experiments. Front Pharmacol 14.
https://doi.org/ARTN 109393410.3389/fphar.2023.1093934.

11.   Fan, X., Wang, H., Wang, W., Shen, J., and Wang, Z. (2024). Exercise training alleviates
cholesterol and lipid accumulation in mice with non-alcoholic steatohepatitis: Reduction of
KMT2D-mediated histone methylation of IDI1. Exp Cell Res 442, 114265.
https://doi.org/10.1016/j.yexcr.2024.114265.

12.   Wang, L., Yang, R., Yuan, B., Liu, Y., and Liu, C. (2015). The antiviral and antimicrobial
activities of licorice, a widely-used Chinese herb. Acta Pharm Sin B 5, 310-315.
https://doi.org/10.1016/j.apsb.2015.05.005.

13.   Ma, D., Zhang, J., Zhang, Y., Zhang, X., Han, X., Song, T., Zhang, Y., and Chu, L. (2018).
Inhibition of myocardial hypertrophy by magnesium isoglycyrrhizinate through the
TLR4/NF-κB signaling pathway in mice. Int Immunopharmacol 55, 237-244.
https://doi.org/10.1016/j.intimp.2017.12.019.

14.   Liu, M., Zheng, B., Liu, P., Zhang, J., Chu, X., Dong, C., Shi, J., Liang, Y., Chu, L., Liu, Y. et
al. (2021). Exploration of the hepatoprotective effect and mechanism of magnesium

isoglycyrrhizinate in mice with arsenic trioxide-induced acute liver injury. Mol Med Rep 23.

https://doi.org/10.3892/mmr.2021.12077.

15.   Lai, X., Zhou, H., Wan, Y., Kuang, J., Yang, Y., Mai, L., Chen, Y., and Liu, B. (2024).

Magnesium isoglycyrrhizinate attenuates nonalcoholic fatty liver disease by strengthening

intestinal mucosal barrier. Int Immunopharmacol 128, 111429.

https://doi.org/10.1016/j.intimp.2023.111429.

16.   Yang, Y.-Z., Zhao, X.-J., Xu, H.-J., Wang, S.-C., Pan, Y., Wang, S.-J., Xu, Q., Jiao, R.-Q., Gu,

H.-M., and Kong, L.-D. (2019). Magnesium isoglycyrrhizinate ameliorates high

fructose-induced liver fibrosis in rat by increasing miR-375-3p to suppress JAK2/STAT3

pathway and TGF-β1/Smad signaling. Acta Pharmacol Sin 40, 879-894.

https://doi.org/10.1038/s41401-018-0194-4.

17.   Lu, C., Xu, W., Shao, J., Zhang, F., Chen, A., and Zheng, S. (2017). Blockade of hedgehog

pathway is required for the protective effects of magnesium isoglycyrrhizinate against

ethanol-induced hepatocyte steatosis and apoptosis. IUBMB Life 69, 540-552.

https://doi.org/10.1002/iub.1639.

18.   Dai, W., Wang, K., Zhen, X., Huang, Z., and Liu, L. (2022). Magnesium isoglycyrrhizinate

attenuates acute alcohol-induced hepatic steatosis in a zebrafish model by regulating lipid

metabolism and ER stress. Nutr Metab (Lond) 19, 23.

https://doi.org/10.1186/s12986-022-00655-7.

19.   National Workshop on Fatty, L., Alcoholic Liver Disease, C.S.o.H.C.M.A., and Fatty Liver

Expert Committee, C.M.D.A. (2018). [Guidelines of prevention and treatment for alcoholic

liver disease: a 2018 update]. Zhonghua Gan Zang Bing Za Zhi 26, 188-194.

https://doi.org/10.3760/cma.j.issn.1007-3418.2018.03.007.

20.   Bertola, A., Mathews, S., Ki, S.H., Wang, H., and Gao, B. (2013). Mouse model of chronic

and binge ethanol feeding (the NIAAA model). Nat Protoc 8, 627-637.

https://doi.org/10.1038/nprot.2013.032.

21.   Wang, Y., Zhang, Z., Wang, X., Qi, D., Qu, A., and Wang, G. (2017). Amelioration of

Ethanol-Induced Hepatitis by Magnesium Isoglycyrrhizinate through Inhibition of Neutrophil

Cell Infiltration and Oxidative Damage. Mediators Inflamm 2017, 3526903.

https://doi.org/10.1155/2017/3526903.

22.   Brunt, E.M., Kleiner, D.E., Wilson, L.A., Belt, P., Neuschwander-Tetri, B.A., and Network,

N.C.R. (2011). Nonalcoholic Fatty Liver Disease (NAFLD) Activity Score and the

Histopathologic Diagnosis in NAFLD: Distinct Clinicopathologic Meanings. Hepatology 53,

810-820. https://doi.org/10.1002/hep.24127.

23.   Che, Z., Song, Y., Xu, C., Li, W., Dong, Z., Wang, C., Ren, Y., So, K.-F., Tipoe, G.L., Wang, F.

et al. (2023). Melatonin alleviates alcoholic liver disease via EGFR-BRG1-TERT axis

regulation. Acta Pharm Sin B 13, 100-112. https://doi.org/10.1016/j.apsb.2022.06.015.

24.   Yi, H.W., Ma, Y.X., Wang, X.N., Wang, C.F., Lu, J., Cao, W., and Wu, X.D. (2015). Ethanol

promotes saturated fatty acid-induced hepatoxicity through endoplasmic reticulum (ER) stress

response. Chin J Nat Med 13, 250-256. https://doi.org/10.1016/s1875-5364(15)30011-x.

25.   Qiu, F., Zeng, R., Li, D., Ye, T., Xu, W., Wang, X., Yan, X., Li, H., and Hu, X. (2023).

Establishment and bioinformatics evaluation of the ethanol combined with palmitic

acid-induced mouse hepatocyte AFLD model (the Hu-Qiu Model). Heliyon 9, e19359.

https://doi.org/10.1016/j.heliyon.2023.e19359.

26. Sharma, R.R., Rashid, H., Bhat, A.M., Sajeeda, A., Gupta, R., and Abdullah, S.T. (2023).
Glabridin ameliorates intracellular events caused by palmitic acid and alcohol in mouse
hepatocytes and fast food diet and alcohol -induced steatohepatitis and fibrosis in C57BL/6J
mice model. Food Chem Toxicol 180, 114038. https://doi.org/10.1016/j.fct.2023.114038.

27. Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M., Mueller, R.,
Nolan, T., Pfaffl, M.W., Shipley, G.L. et al. (2009). The MIQE guidelines: minimum
information for publication of quantitative real-time PCR experiments. Clin Chem 55,
611-622. https://doi.org/10.1373/clinchem.2008.112797.

28. Badia, I.M.P., Velez Santiago, J., Braunger, J., Geiss, C., Dimitrov, D., Muller-Dott, S., Taus,
P., Dugourd, A., Holland, C.H., Ramirez Flores, R.O. et al. (2022). decoupleR: ensemble of
computational methods to infer biological activities from omics data. Bioinform Adv 2,
vbac016. https://doi.org/10.1093/bioadv/vbac016.

29. Meijnikman, A.S., Davids, M., Herrema, H., Aydin, O., Tremaroli, V., Rios-Morales, M.,
Levels, H., Bruin, S., de Brauw, M., Verheij, J. et al. (2022). Microbiome-derived ethanol in
nonalcoholic fatty liver disease. Nature Medicine 28, 2100-2106.
https://doi.org/10.1038/s41591-022-02016-6.

30. Norrmen, C., Figlia, G., Lebrun-Julien, F., Pereira, J.A., Trotzmuller, M., Kofeler, H.C.,
Rantanen, V., Wessig, C., van Deijk, A.L., Smit, A.B. et al. (2014). mTORC1 controls PNS
myelination along the mTORC1-RXRgamma-SREBP-lipid biosynthesis axis in Schwann cells.
Cell Rep 9, 646-660. https://doi.org/10.1016/j.celrep.2014.09.001.

31. https://www.niaaa.nih.gov/alcohols-effects-health/alcohol-drinking-patterns

32. Wang, J., Wang, W., Kollman, P.A., and Case, D.A. (2006). Automatic atom type and bond
type perception in molecular mechanical calculations. J Mol Graph Model 25, 247-260.
https://doi.org/10.1016/j.jmgm.2005.12.005.

33. Tian, C., Kasavajhala, K., Belfon, K.A.A., Raguette, L., Huang, H., Migues, A.N., Bickel, J.,
Wang, Y., Pincay, J., Wu, Q. et al. (2019). ff19SB: Amino-Acid-Specific Protein Backbone
Parameters Trained against Quantum Mechanics Energy Surfaces in Solution. Journal of
Chemical Theory and Computation 16, 528-552. https://doi.org/10.1021/acs.jctc.9b00591.

34. Wang, J., Wolf, R.M., Caldwell, J.W., Kollman, P.A., and Case, D.A. (2004). Development
and testing of a general amber force field. J Comput Chem 25, 1157-1174.
https://doi.org/10.1002/jcc.20035.

35. Song, X., Bao, L., Feng, C., Huang, Q., Zhang, F., Gao, X., and Han, R. (2024). Accurate
Prediction of Protein Structural Flexibility by Deep Learning Integrating Intricate Atomic
Structures and Cryo-EM Density Information. Nat Commun 15, 5538.
https://doi.org/10.1038/s41467-024-49858-x.

36. Zheng, N., Cai, Y., Zhang, Z., Zhou, H., Deng, Y., Du, S., Tu, M., Fang, W., and Xia, X.
(2025). Tailoring industrial enzymes for thermostability and activity evolution by the machine
learning-based iCASE strategy. Nat Commun 16, 604.
https://doi.org/10.1038/s41467-025-55944-5.

37. Valdés-Tresanco, M.S., Valdés-Tresanco, M.E., Valiente, P.A., and Moreno, E. (2021).
gmx_MMPBSA: A New Tool to Perform End-State Free Energy Calculations with
GROMACS. J Chem Theory Comput 17, 6281-6291.
https://doi.org/10.1021/acs.jctc.1c00645.

38. Xu, Z., Liu, D., Liu, D., Ren, X., Liu, H., Qi, G., Zhou, Y., Wu, C., Zhu, K., Zou, Z. et al.

(2022). Equisetin is an anti-obesity candidate through targeting 11β-HSD1. Acta Pharm Sin B 12, 2358-2373. https://doi.org/10.1016/j.apsb.2022.01.006.

39. Lai, X., Zhou, H., Wan, Y., Kuang, J., Yang, Y., Mai, L., Chen, Y., and Liu, B. (2024). Magnesium isoglycyrrhizinate attenuates nonalcoholic fatty liver disease by strengthening intestinal mucosal barrier. Int Immunopharmacol 128, 111429. https://doi.org/10.1016/j.intimp.2023.111429.

40. Lei, Z., Rong, H., Yang, Y., Yu, S., Zhang, T., Chen, L., Nie, Y., Song, Q., Hu, Q., and Guo, J. (2022). Loperamide induces excessive accumulation of bile acids in the liver of mice with different diets. Toxicology 477, 153278. https://doi.org/10.1016/j.tox.2022.153278.

41. Lu, L., Hao, K., Hong, Y., Liu, J., Zhu, J., Jiang, W., Zhu, Z., Wang, G., and Peng, Y. (2021). Magnesium Isoglycyrrhizinate Reduces Hepatic Lipotoxicity through Regulating Metabolic Abnormalities. Int J Mol Sci 22. https://doi.org/10.3390/ijms22115884.

42. Wang, X., Wu, F.P., Huang, Y.R., Li, H.D., Cao, X.Y., You, Y., Meng, Z.F., Sun, K.Y., and Shen, X.Y. (2023). Matrine suppresses NLRP3 inflammasome activation via regulating PTPN2/JNK/SREBP2 pathway in sepsis. Phytomedicine 109, 154574. https://doi.org/10.1016/j.phymed.2022.154574.

43. Wang, S., Zhou, Y., Yu, R., Ling, J., Li, B., Yang, C., Cheng, Z., Qian, R., Lin, Z., Yu, C. et al. (2023). Loss of hepatic FTCD promotes lipid accumulation and hepatocarcinogenesis by upregulating PPARgamma and SREBP2. JHEP Rep 5, 100843. https://doi.org/10.1016/j.jhepr.2023.100843.

44. Chen, W., Wen, L., Bao, Y., Tang, Z., Zhao, J., Zhang, X., Wei, T., Zhang, J., Ma, T., Zhang, Q. et al. (2022). Gut flora disequilibrium promotes the initiation of liver cancer by modulating tryptophan metabolism and up-regulating SREBP2. Proc Natl Acad Sci U S A 119, e2203894119. https://doi.org/10.1073/pnas.2203894119.

45. Foster, C., Gagnon, C.A., and Ashraf, A.P. (2024). Altered lipid metabolism and the development of metabolic-associated fatty liver disease. Curr Opin Lipidol 35, 200-207. https://doi.org/10.1097/MOL.0000000000000933.

46. Devang, N., Adhikari, P., Nandini, M., Satyamoorthy, K., and Rai, P.S. (2021). Effect of licorice on patients with HSD11B1 gene polymorphisms- a pilot study. J Ayurveda Integr Med 12, 131-135. https://doi.org/10.1016/j.jaim.2020.06.006.

47. Leyla Galandarli1*, C.M., Asli Javadzade1, Afat Mammadova1, , and Amrahov1, N. (2022). MOLECULAR DOCKING OF GLYCYRRHIZIN WITH 11β-HSD1 PROTEIN RESEARCH IN: AGRICULTURAL & VETERINARY SCIENCES 6, 141-146.

48. Hughes, K.A., Webster, S.P., and Walker, B.R. (2008). 11-Beta-hydroxysteroid dehydrogenase type 1 (11beta-HSD1) inhibitors in type 2 diabetes mellitus and obesity. Expert Opin Investig Drugs 17, 481-496. https://doi.org/10.1517/13543784.17.4.481.

49. Honma, T., Shinohara, N., Ito, J., Kijima, R., Sugawara, S., Arai, T., Tsuduki, T., and Ikeda, I. (2012). High-fat diet intake accelerates aging, increases expression of Hsd11b1, and promotes lipid accumulation in liver of SAMP10 mouse. Biogerontology 13, 93-103. https://doi.org/10.1007/s10522-011-9363-2.

50. Blaschke, M., Koepp, R., Streit, F., Beismann, J., Manthey, G., Seitz, M.T., Kragl, A., and Siggelkow, H. (2021). The rise in expression and activity of 11beta-HSD1 in human mesenchymal progenitor cells induces adipogenesis through increased local cortisol synthesis. J Steroid Biochem Mol Biol 210, 105850. https://doi.org/10.1016/j.jsbmb.2021.105850.

51.   Morton, N.M., Holmes, M.C., Fievet, C., Staels, B., Tailleux, A., Mullins, J.J., and Seckl, J.R.
(2001). Improved lipid and lipoprotein profile, hepatic insulin sensitivity, and glucose
tolerance in 11beta-hydroxysteroid dehydrogenase type 1 null mice. J Biol Chem 276,
41293-41300. https://doi.org/10.1074/jbc.M103676200.

52.   Lee, S.Y., Kim, S., Choi, I., Song, Y., Kim, N., Ryu, H.C., Lim, J.W., Kang, H.J., Kim, J., and
Seo, H.R. (2022). Inhibition of 11beta-hydroxysteroid dehydrogenase 1 relieves fibrosis
through depolarizing of hepatic stellate cell in NASH. Cell Death Dis 13, 1011.
https://doi.org/10.1038/s41419-022-05452-x.

53.   Maier, C.R., Hartmann, O., Prieto-Garcia, C., Al-Shami, K.M., Schlicker, L., Vogel, F.C.E.,
Haid, S., Klann, K., Buck, V., Münch, C. et al. (2023). USP28 controls SREBP2 and the
mevalonate pathway to drive tumour growth in squamous cancer. Cell Death Differ 30,
1710-1725. https://doi.org/10.1038/s41418-023-01173-6.

54.   Guo, C., Chi, Z., Jiang, D., Xu, T., Yu, W., Wang, Z., Chen, S., Zhang, L., Liu, Q., Guo, X. et
al. (2018). Cholesterol Homeostatic Regulator SCAP-SREBP2 Integrates NLRP3
Inflammasome Activation and Cholesterol Biosynthetic Signaling in Macrophages. Immunity
49, 842-856.e847. https://doi.org/10.1016/j.immuni.2018.08.021.

55.   Hu, M., Han, T., Pan, Q., Ni, D., Gao, F., Wang, L., Ren, H., Zhang, X., Jiao, H., Wang, Y. et
al. (2022). The GR-gp78 Pathway is involved in Hepatic Lipid Accumulation Induced by
Overexpression of 11β-HSD1. Int J Biol Sci 18, 3107-3121.
https://doi.org/10.7150/ijbs.42376.

56.   Zhang, Y., Zhu, Z., Sun, L., Yin, W., Liang, Y., Chen, H., Bi, Y., Zhai, W., Yin, Y., and Zhang,
W. (2023). Hepatic G Protein-Coupled Receptor 180 Deficiency Ameliorates High Fat
Diet-Induced Lipid Accumulation via the Gi-PKA-SREBP Pathway. Nutrients 15.
https://doi.org/10.3390/nu15081838.

57.   Xu, D., Wang, Z., Xia, Y., Shao, F., Xia, W., Wei, Y., Li, X., Qian, X., Lee, J.H., Du, L. et al.
(2020). The gluconeogenic enzyme PCK1 phosphorylates INSIG1/2 for lipogenesis. Nature
580, 530-535. https://doi.org/10.1038/s41586-020-2183-2.

Figure Legends

Fig. 1. MgIG alleviates liver damage in a mouse model of chronic-binge alcoholic liver disease (the NIAAA model). (A) Chemical structure of MgIG. (B) Flowchart illustrating the modeling process for the NIAAA model. (C) Representative results of H&E and Oil Red O staining from the livers of mice in Ctrl, A-Ctrl and MgIG groups (n = 5). (D) Alterations in NAS (NAFLD activity score) and Oil Red O quantification (a.v.: arbitrary value) (n = 5). (E) Ratios of liver weight to body weight (LW/BW) in mice (n = 5). (F) Alterations in serum biochemical parameters (ALT, AST and TG) in mice in three groups (n = 5). (G) Alterations in mRNA

expression of lipid metabolism genes (*Srebp-1c*, *Srebp2*, *Acc1*, and *Scd1*), systemic inflammation

markers (*Tnf-α*, *Il-6, Il-6,* and *Il-16*), and apoptosis-related genes (*Bax* and *Bcl2*) in the mice liver.

The data are presented as mean ± SD. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$.

Scale bar: 50 μm.

Fig. 2. MgIG protects ethanol-induced hepatocyte injury in a cell model. (A) Volcano plot

analysis showing differentially expressed genes between the MgIG group (ALD+MgIG) and the

A-Ctrl (ALD only) group after RNA-seq in liver of mice. (B) The top 5 regulated pathways (both

up- and down-regulated) and the top 3 genes within each pathway are shown. (C) Volcano plot analysis showing differentially expressed genes between the A-Ctrl group and the Ctrl group after RNA-seq in liver of mice. (D) Changes in AML-12 cell viability, apoptosis, and Nile Red staining signals after ethanol/PA treatment, with or without co-treatment with different MgIG doses (0, 0.1, 0.25, 0.5, and 1 mg/ml) (n = 4). (E) Representative images of Nile Red staining in AML-12 cells treated with ethanol/PA, with or without co-treatment with different MgIG doses. (F) Changes in mRNA expression of lipid metabolism genes (*Acc1* and *Scd1*) and systemic inflammation markers (*Tnf-α* and *Il-6*) in AML-12 cells treated with ethanol/PA, with or without co-treatment with different MgIG doses. Data are expressed as mean ± SD. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$. Scale bar: 20 μm.

Fig. 3 IDI1 is involved in MgIG-mediated hepatocyte protection against ethanol. (A) Quantitative PCR validation of RNA-seq results revealed changes in *Idi1* gene expression in AML-12 cells treated with ethanol/PA, with or without 0.25 mg/ml MgIG co-treatment (n = 4). (B) Quantitative PCR results confirmed the knockdown efficiency of *Idi1* genes by siRNA in AML-12 cells. The changes in AML-12 cell viability following *Idi1* siRNA transfection (24 h) were also

assessed (n = 4). (C) Quantitative PCR results confirmed the overexpression efficiency of *Idi1* genes via plasmid transfection in AML-12 cells. Changes in AML-12 cell viability following *Idi1* plasmid transfection (24 h) were also evaluated (n = 4). (D) Changes in cell viability and apoptosis ratios were assessed in ethanol/PA-treated AML-12 cells, with or without 0.25 mg/ml MgIG, following *Idi1* knockdown/overexpression (n = 4). (E, F) Western blot results for TNF-α, IL-6, Bax, and Bcl-2 and cell supernatant results for TNF-α and IL-6 in AML-12 cells treated with ethanol/PA and ethanol/PA + MgIG, with or without *Idi1* knockdown/overexpression (n = 3). (G, H) Nile Red staining area (%) with corresponding representative cell staining images. Data are expressed as mean ± SD. For Western blot quantification: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$. Scale bar: 20 μm.

Fig. 4. HSD11B1 is the direct binding protein of MgIG for the protection of ethanol-induced hepatocytes injury. (A) Bar chart displaying the top 10 upregulated and downregulated transcription factor activity scores in the MgIG (ALD+MgIG) group compared to the A-Ctrl (ALD only) group. (B) The glide score of MgIG binding to the protein structure as determined by molecular docking analysis. (C, D) RMSD and RMSF analysis of the three systems: APO-HSD11B1 (blue) represents unbound HSD11B1 in a physiological saline system, Compound-HSD11B1 (red) represents HSD11B1 bound to MgIG in a physiological saline system, and EtOH-HSD11B1 (green) represents HSD11B1 bound to MgIG in a 0.1 mg/mL ethanol solvent system. (E, F) Molecular modeling analysis of MgIG binding at the HSD11B1 domain in normal

saline and ethanol systems. Left: Cartoon view of MgIG at the HSD11B1 binding site. Right: Close-up surface view of MgIG at the HSD11B1 binding sites. (G) Hydrogen bond analysis of HSD11B1-MgIG interactions in normal saline and ethanol systems. (H, I) The microscale thermophoresis (MST) assay demonstrated direct binding between varying doses of MgIG and human HSD11B1 protein at residues 187. WT: wild-type HSD11B1; M1, M2, M3: point mutations at Tyr177, Tyr183, and Lys187, respectively.

Fig. 5. HSD11B1 is involved in MgIG-mediated hepatocyte protection against ethanol. (A) Quantitative PCR validation of RNA-seq results revealed changes in *Hsd11b1* gene expression in AML-12 cells treated with ethanol/PA, with or without 0.25 mg/ml MgIG co-treatment (n = 4). (B, C) Quantitative PCR confirmed the knockdown efficiency of Hsd11b1 (siRNA) and the overexpression efficiency of Idi1 (plasmid) in AML-12 cells, and cell viability changes were assessed 48 h after transfection (n = 4). (D) Nile Red staining area (%) with corresponding representative cell staining images (n=4). (E, F) Western blot results for TNF-α, IL-6, Bax, and Bcl-2 and cell supernatant results for TNF-α and IL-6 in AML-12 cells treated with ethanol/PA and ethanol/PA + MgIG, with or without *Hsd11b1* knockdown/overexpression (n = 3). Data are expressed as mean ± SD. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. Scale bar: 20 μm.

Fig. 6. MgIG exerts its protective effect via the HSD11B1-SREBP2-IDI1 axis in ALD model.

(A, B) Co-IP was used to verify the direct binding between HSD11B1 and SREBP2 with or without MgIG treatment. (C, D) Alterations in protein levels of HSD11B1, p-SREBP2, n-SREBP2, and IDI1 in AML-12 cells induced by EtOH/PA, following knockdown or overexpression of *Hsd11b1*, with and without MgIG treatment (n = 3). (E) Immunofluorescent staining was conducted to visualize the distribution and expression of SREBP2 (red) in EtOH/PA-induced AML-12 cells, with or without MgIG, following *Hsd11b1* knockdown or overexpression. (F) Co-IP was used to verify the direct binding between HSD11B1 and IDI1. (G) Effects of *Srebp2* on

Idi1 transcriptional regulation were measured by luciferase assays in AML-12 and 293T cell lines. Idi1-wild-type (WT) or Idi1-mutant (Mut), plasmids with WT promoter cDNA clone of *Idi1* or with mutant promoter cDNA clone plasmid; pRL-TK, an internal control reporter plasmid. (H, I) The expression levels of *Hsd11b1*, *Srebp2*, and *Idi1* in normal liver were measured following the knockdown of *Hsd11b1*, *Srebp2*, and *Idi1*, or the overexpression of *Hsd11b1* and *Idi1*, respectively (n = 4). (J, K) The expression levels of *Hsd11b1*, *Srebp2*, and *Idi1* in ALD liver were measured following the knockdown of *Hsd11b1*, *Srebp2*, and *Idi1*, or the overexpression of *Hsd11b1* and *Idi1*, respectively (n = 5). Data are expressed as mean ± SD. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$. $****P < 0.0001$. Scale bar: 10 μm.

Fig. 7. MgIG alleviates ALD-induced liver injury via the HSD11B1-SREBP2-IDI1 axis. (A-D) Representative liver H&E and Oil Red O staining results from ALD mice with *Hsd11b1*, *Srebp2*, or *Idi1* knockdown or overexpression, with and/or without MgIG co-treatment. Changes in quantitative NAS (NAFLD activity score) and Oil Red staining (area %) were calculated and analyzed (n = 5). (E-F) Changes of serum ALT and TNF-α from ALD mice with *Hsd11b1*, *Srebp2*,

or *Idi1* knockdown or overexpression, with and/or without MgIG co-treatment (n = 5). (G) Alterations in protein levels of HSD11B1, p-SREBP2, n-SREBP 2, and Idi1 in ALD mice, following knockdown or overexpression of *Hsd11b1*, with and without MgIG co-treatment (n = 3). Data are expressed as mean ± SD. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. Scale bar: 50 μm.

Drug	Target	PDB ID /UniProt ID	Glide score
MgIG	Hsd11b1	1Y5R	-8.75
	Gp78	Q9R049	-6.95
	Nfkb	8TKN	-6.35
	Insig1	Q8BGI3	-6.16
	Srebp1	Q9WTN3	-6.13
	Tlr4	7MLM	-5.95
	Tbx3	P70324	-5.74
	Irf3	7JFM	-5.73
	Insig2	Q91WG1	-5.69
	Klf10	O89091	-5.51
	Ep300	B2RWS6	-5.43
	Gpr55	Q3UJF0	-5.41
	Tgfb1/1	P04202	-5.36
	Ets2	P15037	-5.36
	Mecp2	Q9Z2D6	-5.35
	Pparg	P37238	-5.32
	Scap	Q6GQT	-5.29
	sp1	O89090	-5.23
	Nr1h3	3F6C	-5.22
	Stat2	Q9WVL2	-5.2
	Insr	7SL7	-5.18
	Atf6	F6VAN0	-5.17
	Mbtps2	Q8CHX6	-5.06
	Ppara	P23204	-4.99
	Nfya	P23708	-4.96
	Tp53	1SSF	-4.77
	Zeb1	Q64318	-4.72
	Zbtb7a	O88939	-4.64
	Foxl2	O88470	-4.45
	Grp78	P20029	-4.36
	Nr1d1	Q3UV55	-4.35
	Smad1	P70340	-4.28
	Hdac1	O09106	-4.19
	Npc1l1	8EUS	-4.18
	Med15	Q924H2	-4.14
	Kpnb1	1UKL(A/B)	-4.12
	Creb	4I9O	-4.05
	Srebp2	Q3U1N2	-3.95
	Ctnnb1	4E8V	-3.89
	Hey1	Q9WV93	-3.69
	Pbx1	1LFU	-3.44
	Ppargc1b	Q8VHJ7	-3.07
	Lmna	1UFG	-2.99
	Lats2	2COS	-2.46