Authors

Kaleigh E. Davis[1], Tess N. Grainger[1], Po-Ju Ke[2], Patrick L. Thompson[3], Mary I. O'Connor[3], Joey R. Bernhardt[1]

Affiliations

1. University of Guelph, Department of Integrative Biology; kaleigh.davis@uoguelph.ca; tgrainge@uoguelph.ca; joey.bernhardt@uoguelph.ca
2. Institute of Ecology and Evolutionary Biology, National Taiwan University; pojuke@ntu.edu.tw
3. University of British Columbia, Department of Zoology and Biodiversity Research Centre; oconnor@zoology.ubc.ca; patrick.thompson@dfo-mpo.gc.ca

Running Title: Effects of warming on competition

Keywords: warming, competition, coexistence, global change, metabolic theory, coexistence theory

Article Type: Letter

Word Counts
Abstract: 150
Main text: 4985
Text boxes: N/A

Number of References: 64

Number of Figures, Tables, and Text Boxes: 5

Data Accessibility Statement: Data and code for analyses are available at DOI: 10.5281/zenodo.17253649.

Corresponding Author Contact
Kaleigh Davis
Mailing Address: 50 Stone Road, Guelph, ON, N1G 2W1
Email: kaleigh.davis@uoguelph.ca
Phone: +1 (778)839-0351

<u>Abstract</u>

Understanding the effects of climate change on ecological communities has been

limited by a lack of general theory for how temperature affects competition. To fill this

knowledge gap, we integrated Modern Coexistence Theory and the Metabolic Theory of

Ecology by incorporating empirically-derived temperature sensitivities into Modern

Coexistence Theory's central model. We then simulated warming in consumer-resource

systems and found that warming reduced both niche and fitness differences, making

species more ecologically similar and competitive interactions more neutral. The

greatest shifts in competition occurred when temperature sensitivities among species

were highly asymmetrical. Effects of warming on competition via niche differences were

comparable to those on fitness differences, suggesting that the emphasis on vital rates

in global change research may overlook key biodiversity drivers. This general theory

expands the domains of two prominent ecological theories and provides predictions for

how warming may alter competition even in benign regions of species' thermal niches.

 <u>Introduction</u>

Understanding the effects of climate warming on biodiversity is a central challenge of

modern ecology (Parmesan & Yohe 2003; Peters & Lovejoy 1992; Román-Palacios &

Wiens 2020). Species interactions, and competition in particular, influence species'

responses to the environment, and are thus critical to understanding the effects of

warming on biodiversity (Godoy 2019; Tilman 1994). While a considerable body of

empirical work has demonstrated shifts in competition with changing environments (e.g.,

Alexander *et al.* 2015; Germain *et al.* 2018; Pérez-Ramos *et al.* 2019), the nature of

these shifts has been variable, and explanations usually context specific. Whether there

are any general patterns in how warming affects competition, or whether these changes

are inherently species-specific is a major open question. In the face of rapid

environmental change, new theory is needed to address this question.

Modern Coexistence Theory has become the leading framework for understanding

competition and coexistence (Chesson 2000; Godoy & Levine 2014). This theory has

the advantage of being agnostic to the specific mechanisms or context of competitive

interactions, which allows it to be used to investigate general principles of competition

and to be applied across a wide range of systems (Godwin *et al.* 2020; Grainger *et al.*

2019). Within Modern Coexistence Theory, in the absence of environmental

fluctuations, species' competitive differences are described along two axes: fitness

differences and niche differences (Chesson 2000). Fitness differences are interspecific

differences that give one species a competitive advantage and stem from differences

between competing species in intrinsic growth rate and sensitivity to competition. Niche

differences are differences that reduce the strength of interspecific competition relative

to intraspecific competition, for example, differences in resource use or susceptibility to

predators (Chesson 2000; Chesson & Kuang 2008). The balance of these differences

determines whether two species can coexist, one species will exclude the other, or they

are competitively neutral (Adler *et al.* 2007). In global change research, work to

understand the effects of warming on species abundance and distribution has

predominantly emphasized shifts in vital rates (Spaak *et al.* 2021; fitness differences;

e.g. Merow et al 2014, Ehrlen and Morris 2015; but see Adler et al., 2012; Usinowicz &

Levine 2018), while the potential effects of warming on niche differences remain

underexplored.

One criticism of Modern Coexistence Theory is that, because it is phenomenological, it

is not well-suited to generate a priori predictions about how a given factor will affect

competition and coexistence (Terry 2025). Indeed, the parameters used to estimate

niche and fitness differences in classical Modern Coexistence Theory models describe

the outcomes of competitive interactions but are not related explicitly to underlying

mechanisms (Chesson 2000). However, niche and fitness differences can be

alternatively calculated from their constituent mechanistic processes (Chesson & Kuang

2008; Letten *et al.* 2017), for example, parameters related to resource competition

(Song *et al.* 2019). Fortunately, the effects of temperature on many of these mechanistic

processes have also been investigated empirically (Dell *et al.* 2011; Rall *et al.* 2010). To

evaluate how these temperature effects shift competitive dynamics with warming, the

logical next step is to integrate empirically-grounded temperature dependencies into

Modern Coexistence Theory models.

The Metabolic Theory of Ecology is a widely used framework for understanding the

effects of temperature on ecological processes. A central premise of this framework is

that ecological processes, such as population growth and species interactions, all stem

from metabolic processes, which control the uptake and transformation of resources

into maintenance, growth and reproduction (Brown *et al.* 2004). In this way, metabolism

fundamentally and predictably constrains higher order processes across systems,

leading to a predictable "scaling" of temperature effects across levels of organization.

From this principle of metabolic scaling, the Metabolic Theory of Ecology has been used

to predict the effect of temperature on population growth rates (Savage *et al.* 2004),

population carrying capacities (Bernhardt *et al.* 2018a), consumer-resource interactions

(O'Connor 2009; O'Connor *et al.* 2011), food webs (Gibert *et al.* 2022; Gilbert *et al.*

2014; Vasseur & McCann 2005), and ecosystem productivity (Enquist *et al.* 2003;

Lopez-Urrutia *et al.* 2006; Welter *et al.* 2015). Despite this broad success, extending the

metabolic scaling approach to competitive interactions has proven challenging, due to a

conflict in focus between competition theory and the Metabolic Theory of Ecology.

While competition theory has focused primarily on changes in ecological outcomes

driven by interspecific variation in common underlying traits (here, intra-process

asymmetry, Fig. 1) (e.g., Kraft *et al.* 2008; Weiher *et al.* 1998), the Metabolic Theory of

Ecology has largely ignored interspecific variation in thermal traits, and focused instead

on average differences in thermal responses across metabolic processes (here, inter-

process asymmetry, Fig. 1, but see Clegg & Pawar 2024; Gibert *et al.* 2022). Since our

focal traits are thermal traits, we will refer to these intra- and inter-process thermal

asymmetries (Fig. 1). Progress to incorporate interspecific variability in thermal traits

into metabolic scaling studies of competition has been made primarily by exploring

consequences of high, above optimal, temperature, and generalist-specialist trade-offs

in thermal tolerance breadth (e.g., Amarasekare & Johnson 2017; Simon &

Amarasekare 2024; Smith & Amarasekare 2018). However, most species live, on

average, well below their thermal optimum in nature (Bernhardt *et al.* 2018b; Martin &

Huey 2008), and whether interspecific trait differences in the non-stressful (sub-optimal)

temperature range, alone, can produce general patterns in competitive dynamics under

warming is an open question. Incorporating temperature-dependence at sub-optimal

temperatures into the governing processes of Modern Coexistence Theory enables us

to answer this question.

Here we draw from the Metabolic Theory of Ecology and Modern Coexistence Theory to

develop theory for how warming affects competition for shared resources at suboptimal

temperatures. We use this theory to address three questions: 1) How do the

temperature sensitivities of each process underlying competition contribute to changes

in niche and fitness differences with warming? 2) How do inter- and intra-process

thermal asymmetries, affect these changes? 3) Are there general effects of warming on

competition that emerge from the combined effects of warming on its suite of underlying

mechanistic processes? To answer these questions, we analyzed a MacArthur

consumer–resource model (MacArthur 1970) using the canonical two-species Lotka–

Volterra formulation of Modern Coexistence Theory that expresses phenomenological

competition coefficients with mechanistic parameters (Chesson 1990, 2011; Chesson &

Kuang 2008; Song *et al.* 2019). We modeled the effects of warming on competition by

making each parameter in the MacArthur consumer–resource model temperature-

sensitive. We synthesized temperature sensitivity data for each model parameter from

the published literature to generate an empirical distribution of temperature sensitivities

and then used this distribution to simulate effects of warming on competition between

species pairs. In doing so, we identified a novel set of general predictions for how

competition is affected by warming, and the conditions under which we expect to see

more extreme shifts in competition.

<u>Materials and Methods</u>

Resource competition model

We modeled competition in a two-consumer system (N_1 and N_2) with two substitutable

resources (R_a and R_b) following the classic MacArthur consumer–resource model

(MacArthur 1970):

169
$$\frac{dN_i}{dt} = \sum_{k\in\{a,b\}} v_{ik} c_{ik} R_k N_i - m_i N_i \qquad\qquad \text{Eq. 1}$$

170
$$\frac{dR_k}{dt} = r_k R_k \left(1 - \frac{R_k}{K_k}\right) - \sum_{i=1}^{2} c_{ik} R_k N_i \, , \qquad\qquad \text{Eq. 2}$$

with *i = 1* or *2* and *k = a* or *b* (see Fig. S1 for simulations with >2 resources). Here, the

resource follows logistic growth with intrinsic growth rate r_k and carrying capacity K_k.

Consumer species *i* uses resource *k* following a Holling type-I functional response, with

per capita consumption rate (hereafter "consumption rate") parameter c_{ik} and

conversion efficiency parameter v_{ik}. While the type-I functional response is sometimes

considered unrealistic, recent work suggests that it may be appropriate in more cases

than previously thought (Coblentz *et al.* 2023; Novak *et al.* 2025). Finally, consumers

experience per-capita mortality m_i. We converted this into a two-species Lotka–Volterra

competition model by separation of timescales. Specifically, by assuming fast resource

dynamics, the quasi-equilibrium of R_k can be solved as:

183
$$\widehat{R_k} = \frac{K_k}{r_k}\left(r_k - \sum_{i=1}^{2} c_{ik}\, N_i\right),$$
Eq. 3

185   which is a function of consumer abundances. Substituting this expression back into Eq.

186   1, we arrive at the following:

188
$$\frac{dN_i}{dt} = N_i\left[\left(\sum_{k\in\{a,b\}} v_{ik}\, c_{ik} K_k - m_i\right) - \left(\sum_{k\in\{a,b\}} v_{ik}\frac{K_k}{r_k} c_{ik}^2\right)N_i - \left(\sum_{k\in\{a,b\}} v_{ik}\frac{K_k}{r_k} c_{ik} c_{jk}\right)N_j\right]$$

189
$$= g_i N_i\left[1 - \sum_{j=1}^{2} \alpha_{ij}\, N_j\right],$$

191   Eq. 4

192   with

194
$$g_i = \sum_{k\in\{a,b\}} v_{ik}\, c_{ik} K_k - m_i$$
Eq. 5

196
$$\alpha_{ij} = \frac{\sum_{k\in\{a,b\}} v_{ik}\frac{K_k}{r_k}c_{ik}c_{jk}}{g_i} = \frac{\sum_{k\in\{a,b\}} v_{ik}\frac{K_k}{r_k}c_{ik}c_{jk}}{\sum_{k\in\{a,b\}} v_{ik}c_{ik}K_k - m_i} \quad .$$
Eq. 6

Here, g_i represents consumer i's intrinsic growth rate, which equals the maximum

energy gain by consumption minus the consumer's per-capita mortality rate. On the

other hand, α_{ij} represents the relative reduction in consumer species i's intrinsic growth

rate caused by competition from species j (i.e., relative competition coefficient; sensu

Song *et al.* 2019).

Modern Coexistence Theory then defines niche overlap (ρ) and fitness ratio ($\frac{f_2}{f_1}$) as

follows:

207
$$\rho = \sqrt{\frac{\alpha_{12}\alpha_{21}}{\alpha_{11}\alpha_{22}}}$$
Eq. 7

209
$$\frac{f_2}{f_1} = \sqrt{\frac{\alpha_{11}\alpha_{12}}{\alpha_{22}\alpha_{21}}} \quad .$$
Eq. 8

Here, ρ quantifies the relative strength of intraspecific and interspecific competition, and

$\frac{f_2}{f_1}$ quantifies species' relative sensitivity to competition. Finally, we define niche

differences (ND) as $-log(\rho)$ and fitness differences (FD) as $log\left(\frac{f_2}{f_1}\right)$. While other

definitions of ND and FD are more flexible (Spaak *et al.* 2023) these definitions allow us

to compare ND and FD on the same scale, and to recover the ecologically intuitive

condition for species coexistence, i.e., the ND between a pair of species must be

greater than their FD for them to coexist.

Adding temperature sensitivity to model parameters

For each mechanistic process from the MacArthur model (i.e., r_k, K_k, c_{ik}, v_{ik}, m_i; Eq. 4),

we modeled temperature sensitivity using an exponential Arrhenius-form function:

$$Y(T) \; = \; Y_{T_{amb}} \; e^{\left(\frac{E_Y}{k_B T_{amb}} - \frac{E_Y}{k_B T}\right)} \, , \qquad\qquad \text{Eq. 9}$$

where Y is any parameter of the consumer-resource model (Eq. 4), T is the

environmental temperature in Kelvin (K), T_{amb} is the ambient temperature, here set to

10 °C (= 283.15 K), $Y_{T_{amb}}$ is the rate of the process at the ambient temperature, E_Y is

the exponential rate of increase in process rate with increasing temperature (Fig. 1;

hereafter temperature sensitivity), and k_B is Boltzmann's constant (eV/K). The

temperature sensitivity parameter, E_Y, is equivalent to the activation energy parameter

from chemistry's Boltzmann-Arrhenius model, which has been applied to ecological

rates within the Metabolic Theory of Ecology (Brown *et al.* 2004).

By standardizing environmental temperature, T, relative to ambient temperature, T_{amb},

we isolate the effects of 15 °C warming, rather than the particular range of temperatures

employed for our analyses (10 – 25 °C). While it is known that most individual- and

population-level temperature responses are unimodal (Amarasekare & Savage 2012;

Kingsolver 2009) and that exponential models typically overpredict the temperature

sensitivity of processes (Knies & Kingsolver 2010; Michaletz & Garen 2024; Pawar *et al.*

2016), many studies have focused on the rate of increase in the exponential portion of

the temperature-response curve (Fig. 1B, 1C) to model scaling effects of temperature

(Bernhardt *et al.* 2018a; Gibert *et al.* 2022; Gilbert *et al.* 2014). As such, we used an

exponential model to describe warming-induced shifts in competition when both species

were in this exponentially increasing portion of their thermal niche, and where the

effects of temperature were non-stressful (sub-optimal). We obtained temperature

dependence estimates for each parameter from posterior distributions (Fig. 2)

generated from an empirical data synthesis for each parameter (see *Supplemental*

Methods).

Simulating effects of warming on competition

Model starting conditions

We answered our three questions by simulating competition across a 15-degree

temperature gradient (10°C to 25°C) between consumer species pairs that co-occur or

coexist under ambient temperatures (T_{amb} = 10°C). All simulations presented in the

main text had the same starting conditions at T_{amb} , captured by the intercepts in the

Arrhenius models ($Y_{T_{amb}}$; Eq. 9, Table S1) that comprise the temperature-dependent

consumer-resource model (Eq. 4): at T_{amb}, species were predicted to coexist and

consumer species pairs have symmetrical resource preferences, so that each consumer

specialized to the same degree on one of two substitutable resources (given by c_{ik} and

v_{ik}; Table S1). At T_{amb} the preferred resource of N_2 (here, r_a) had a faster growth rate

than the preferred resource of N_1 (here, r_b). This set of starting conditions placed our

species pairs on the boundary between coexistence and competitive exclusion of N_1 by

N_2 (Fig. 3). Other starting conditions for competing consumers that co-exist under

ambient temperature conditions were explored in supplementary analyses (Table S1,

Figs. S2, S3, S4). Although 15°C of warming is unrealistic for changes in average

temperatures from a climate change perspective, we used this large thermal gradient to

demonstrate our results clearly. Applying a more realistic thermal gradient (5 °C)

generated qualitatively similar results, but with less contrast (Fig. S5).

Simulation: Questions 1 and 2

To examine the effects of each parameter's temperature sensitivity on ND and FD

(Question 1) and the effects of thermal asymmetries on competition (Question 2), we

ran 500 warming simulations, varying one focal parameter at a time. For each

simulation we randomly drew temperature sensitivity, E_Y, values for the focal parameter

from its empirical posterior distribution, and we set all other parameters' temperature

sensitivities equal to 0. After each temperature sensitivity for the focal parameter was

drawn, we simulated 15°C warming and calculated ND and FD every 0.1°C, according

to Eqs. 7 and 8. For resource growth rate, r_k, resource carrying capacity, K_k, and

consumer mortality rate, m_i, temperature sensitivities for each consumer or resource

were drawn independently for each simulation. For consumption rate, c_{ik}, we drew

temperature sensitivities independently for each species' consumption rate on each

resource (See Fig. S6 for other draw configurations). Finally, for resource conversion

efficiency, v_{ik}, we assumed that the temperature sensitivity primarily reflects the

nutritional composition of the resource, and therefore was the same for both consumers

of a given resource (i.e. $E_{v_{1a}} = E_{v_{2a}}$).

To explore the effects of inter- and intra-process thermal asymmetries on competition

(Question 2), we quantified the magnitude of the thermal asymmetries from the

temperature sensitivities drawn for each simulation and examined the associated shift in

ND–FD space (See *Supplemental Methods*). We quantified the magnitude of thermal

asymmetry as the absolute difference between the values drawn for the focal parameter

in each simulation (e.g. $|E_{r_a} - E_{r_b}|$). We then quantified the shift in ND–FD space as the

Euclidean distance between the model's starting position and the position after 15°C

simulated warming.

Simulation: Question 3

To identify general effects of warming on competition (Question 3), we used the same

set of starting conditions as in Questions 1 and 2 (Table S1), but imposed temperature

sensitivity on all model parameters simultaneously, then simulated warming over a 15°C

thermal gradient. For each simulation (N = 500), we drew temperature sensitivity values,

E_Y, for all parameters from their respective distributions, then calculated ND and FD at

each 0.1°C step. We quantified effects of warming on competition based on the

distance between the species' starting ND and FD values (at T_{amb}, 10°C) and their

median ND and FD values at the final temperature (25 °C).

Data and code for analyses are available at DOI: 10.5281/zenodo.17253649.

Results

Empirically-derived temperature sensitivity of parameters

We synthesized consumer-resource parameters' temperature sensitivity estimates from

58 studies, representing 79 ectothermic consumer or resource species. Represented

species included heterotrophic and autotrophic as well as unicellular and multicellular

organisms (Table S2, Fig. S7). We found considerable variation within and across

temperature sensitivity estimates, E_Y (Fig. 2; Table S3), suggesting both inter-process

and intra-process thermal asymmetries were present in our synthesized dataset.

Posterior distributions and 95% credible intervals for the temperature sensitivities of

resource growth rate, r_k, consumption rate, c_{ik}, and consumer mortality rate, m_i, were

positive, indicating that these rates increase with temperature. Posterior means and

95% credible intervals were: E_{r_k} = 0.79 eV (0.62, 0.96 eV; 30 observations); $E_{c_{ik}}$ = 0.56

eV (0.50, 0.63 eV; 105 observations); E_{m_i} = 0.49 eV (0.35, 0.62 eV; 8 observations; Fig.

2A, 2B, 2E). Overlapping 95% credible intervals across these three parameters

suggests no inter-process thermal asymmetry among them (Fig. 2F). Contrastingly,

resource carrying capacity, K_k, had a negative temperature sensitivity, E_{K_k} = -0.65 eV (-

0.98, -0.33 eV; 8 observations; Fig. 2C), indicating that carrying capacities decrease

with warming. This negative temperature sensitivity was statistically distinct from that of

consumption rate, resource growth rate, and mortality rate (Fig. 2F). Finally, the

posterior distribution for the temperature sensitivity of conversion efficiency had a 95%

credible interval that spanned both negative and positive values and was centered near

0: $E_{v_{ik}}$ = 0.0036 eV (-0.41, 0.42 eV; 5 observations; Fig. 2D). Credible intervals suggest

inter-process thermal asymmetries between conversion efficiency, consumption rate

and resource growth rate, but not between resource carrying capacity and conversion

efficiency or mortality rate (Fig. 2F).

Question 1: How do temperature sensitive processes underlying competition

affect niche and fitness differences?

For competing consumer species that co-exist under ambient temperature conditions,

warming the system caused every process except mortality to have large effects on

niche and/or fitness differences (ND, FD; Fig. 3). Effects of warming on resource

consumption rate, c_{ik}, drove variation in both ND and FD (Fig. 3A), whereas effects of

warming on resource growth rate, r_k, primarily drove variation in FD (Fig. 3B). We found

that temperature sensitivity in these two processes had the largest potential to affect

competitive outcomes in our model, because their temperature sensitivity can move the

species pair along a trajectory that is far from parallel (and in some cases close to

perpendicular) to the coexistence boundary as the system warms. In contrast, effects of

warming on resource carrying capacity, K_k, and conversion efficiency, v_{ik}, drove

variation parallel to the coexistence boundary (Fig. 3C, 3D, S3). In our focal analysis,

where species pairs underwent warming from a starting position on the coexistence

boundary, temperature-dependent carrying capacity and conversion efficiency drove the

species towards increased functional similarity (i.e. toward the neutral point).

In supplemental analysis of co-existing species pairs with different resource preferences

at ambient temperature, and therefore different starting points in ND—FD space (Table

S1), effects of temperature-sensitivity in resource consumption rate, c_{ik}, resource growth

rate, r_k, and mortality, m_i, were similar, while effects of temperature-sensitivity in

resource carrying capacity, K_k, and resource conversion efficiency, v_{ik}, differed (Fig.

S2C-D, S3C-D). Specifically, temperature-sensitive resource carrying capacity and

conversion efficiency had qualitatively different effects on ND and FD within the 15°C

temperature gradient when consumers' resource preferences were sufficiently different

at the ambient temperature. That is, when warming amplified the competitive

dominance of one consumer species over the other (e.g. when N_1 had a stronger

preference for its preferred resource than N_2, and conversion efficiency of that preferred

resource increased with warming), those consumers increased in ND over the 15°C

temperature gradient, and the species' competitive trajectory pointed away from the

neutral point (Fig. S3D).

Across all analyses temperature-sensitive of consumer mortality rate only affected FD.

Competitive dynamics were highly insensitive to variation in the temperature sensitivity

of mortality, as indicated by the absence of visible shifts in Fig. 3E (see also Fig. 4E).

Question 2: How do thermal asymmetries contribute to shifts in competition with

warming?

We found that competing species that responded identically to warming (i.e. had

identical temperature sensitivities for all parameters) experienced no change in

competitive dynamics with warming (Fig. S8). That is, without intra-process thermal

asymmetries, inter-process thermal asymmetries did not affect competitive dynamics.

However, when species differed in their temperature sensitivities for a given process,

competitive dynamics shifted with warming. Within each parameter, larger differences in

temperature sensitivities between competing species or between their resources (intra-

process thermal asymmetries) were associated with larger shifts in competition, i.e.

larger Euclidean distances between the species' "start" point at the ambient temperature

and the "end" point after 15°C warming (Fig. 4, S9). For consumer mortality rate, m_i,

changes in position with warming were too small relative to that of other parameters to

be visible in our analysis (Fig. 4E).

For each parameter except consumption rate, warming created two distinct trajectories

that moved in opposite directions away from the ambient temperature point (Fig. 4B-E).

Which trajectory a species pair moved along depended on the relative value of the two

focal temperature sensitivities, or the directionality of the intra-process thermal

asymmetry (Fig. 4B-E). For resource growth rate and consumer mortality rate, these

trajectories moved monotonically toward the region of NFD space where a consumer's

preferred resource was more abundant with warming, or where consumer mortality was

minimized, though the latter effect was too small to observe for most species pairs. For

example, if the temperature sensitivity for growth rate was higher for resource a than for

resource b, then, when the system was warmed, the coexistence trajectory moved

monotonically toward the region of competitive dominance for the species that

preferentially consumes resource a, here N_2 (e.g. Fig. 3B, 4B). Interestingly, the two

trajectories for carrying capacity and conversion efficiency comprised one monotonic

trajectory and one non-monotonic trajectory, both of which pointed ultimately toward

neutrality (Fig. 3C-D, 4C-D, S3). However, large differences in ambient-temperature

resource preferences between competing species drove qualitative changes to these

trajectories over the 15 °C warming gradient (Table S1, Fig. S3).

Question 3: How does warming affect competition via effects on niche and fitness

differences?

For consumer species existing at the boundary of coexistence and competitive

exclusion, the temperature sensitivity of all processes underlying competition generally

drove decreases in both ND and FD with warming (Fig. 5A, yellow dot to magenta dot).

The majority of species pairs decreased in ND (58%) or FD (58%) and the median

decreases in ND and FD were similar in magnitude (5.5% decrease and 9.7%

decrease, respectively; Fig. 5C-D). Declines in ND and FD with warming were

associated with decreasing resource carrying capacity and resulting declines in all

competition coefficients (Fig. S10). This led to a general trend of competitive dynamics

shifting toward neutrality with warming (Fig. 5A). The distribution of shifts in competition

with warming was skewed, with very large shifts being rare (Fig. 5B). The magnitude of

shift in competition was positively related to the magnitude of intra-process thermal

asymmetries (Table S4). This result is reinforced by analysis of competition between

species at the extremes of the observed temperature sensitivity distributions (top and

bottom 5%), which showed much larger shifts in competition with 15°C warming (Fig

S11).

While some competitive dynamics became less neutral (i.e. increased in ND and FD)

under 15°C warming, simulating extreme warming revealed that trajectories that initially

moved away from neutrality bent back toward neutrality with sufficient warming (Fig.

S10, S12). These results were robust to different consumer and resource traits and

numbers of resource species, but the amount of warming required to shift the median

effect toward neutrality under these different conditions varied (Figs. S1-S4).

Discussion

In this study, we drew from two prominent theories in ecology to address a central

question in global change biology: how does warming affect competition for shared

resources? Using empirically-derived temperature sensitivities of the processes

underlying competition, we found three general patterns in how simulated competition

responded to warming. First, we found that warming tended to reduce both fitness

differences and niche differences. This resulted in a general trend of warming causing

species pairs to move toward neutrality. Next, we found that warming-induced increases

in mortality had surprisingly little effect on these shifts in competition with warming.

Finally, we found that species pairs that underwent the largest shifts in competition with

warming had highly asymmetric temperature sensitivities in the processes underlying

competition. Together, these findings suggest that understanding how communities will

change with warming requires an understanding of how both population vital rates and

resource use traits change with temperature, and how variable these traits are within a

community.

While much of the work seeking to understand biodiversity effects of climate change

has focused on temperature-induced changes in fitness differences (e.g. growth rates;

Ehrlén & Morris 2015; Kordas *et al.* 2011; Merow *et al.* 2014), here we show that

understanding the effects of warming on competition requires an equal consideration of

niche differences. As expected, resource growth rate had a large effect on competition

in our analysis. This effect was near orthogonal to the coexistence boundary, occurring

primarily via shifts in fitness differences and causing resource growth rate to affect

competitive hierarchies (by moving species from one region of ND—FD space to

another) under warming. However, warming is unlikely to affect resource growth rate in

isolation, and when warming was allowed to affect the full suite of processes underlying

competition, species pairs generally also decreased in niche differences with warming

(Fig. 5). These shifts were driven primarily by the effects of warming on consumption

rate, resource carrying capacity, and consumer conversion efficiency – all of which

determine patterns of relative resource use among species. Of these processes,

warming effects on consumption rates also had the capacity to alter competitive

hierarchies (Fig. 3A). Together, these results add to a limited body of literature that

suggests that temperature effects on niche differences play a key role in how warming

affects communities (Adler *et al.* 2012).

Warming-induced changes to mortality rates have been a focal point of climate

modelling and thermal physiology (Deutsch *et al.* 2008; Sunday *et al.* 2014), and have

been shown to have effects on competition and, specifically, on fitness differences

(Song *et al.* 2019). In our analysis, despite exponentially increasing consumer mortality

rates across the thermal gradient (Amarasekare & Savage 2012; Simon & Amarasekare

2024) effects of warming on mortality rate had almost no effect on competition. This is

due to the independence of consumer mortality rate from effects of interspecific

competition in consumer-resource models (Eq. 6; Song *et al.* 2019), and is consistent

with other work suggesting that mortality rates alone cannot drive shifts in coexistence

(Chesson 2000; Smith & Amarasekare 2018). This suggests that, when species are

living at temperatures far below their thermal optimum, and biological processes

underlying competition are increasing exponentially with warming, quantifying effects of

warming on mortality may be less important than previously thought.

Within Modern Coexistence Theory, competition is considered neutral when there is an

absence of both niche differences and fitness differences; biologically this means that

species are functionally equivalent (Adler *et al.* 2007). In our analysis, warming had a

general tendency to shift competing species towards neutrality, regardless of species

thermal traits under ambient temperatures (Fig. 5, S1-S5). In our simulations, this

tendency to move toward neutrality as temperatures increased was related to the

exponential decrease in resource carrying capacity with warming (Fig. 2, S10). As

consumers became increasingly limited by reduced resource supply, intra- and inter-

specific competition weakened, and consumers thus became more ecologically similar

to one another. This result is consistent with theoretical (Hubbell 2001; Lande 1993) and

empirical (Schreiber *et al.* 2022; Siepielski *et al.* 2010) work showing that smaller

population sizes are associated with increased neutrality and stochasticity. Additionally,

our finding that systems trend toward neutrality regardless of competitors' traits under

ambient conditions is consistent with recent work suggesting that that there may be

characteristic trajectories for how competition changes under extrinsic influences, rather

than bespoke shifts based on interspecific trait differences (Pastore *et al.* 2021).

While our finding that warming generally makes competition more neutral is

independent of the model's starting conditions, we found that the specific resource-use

and thermal traits of species influence the amount of warming required to shift

competition toward neutrality (Fig. 5, Figs. S1-S4). That is, for some consumer pairs, an

increase in neutrality may not be realized within a biologically plausible range of

temperatures because the amount of warming required to create this effect might be

large. In addition, if warming exceed species' thermal optima for one or more of the

processes underlying competition, then competitive trajectories may diverge from those

predicted here (Fig. S13). This is relevant in systems where species' thermal optima are

close to mean environmental temperatures, and may also be relevant in systems with

highly variable environmental temperatures, which at times exceed thermal optima

(Amarasekare & Johnson 2017; Casas Goncalves & Amarasekare 2021; Smith &

Amarasekare 2018).

Under changing thermal regimes, which types of species pairs are likely to experience

large competitive shifts remains an open question. While most species simulated in our

study experienced relatively small shifts in competitive dynamics with warming, species

pairs that experienced large shifts had large intra-process thermal asymmetries (Fig. 4,

Table S4, Fig. S9). While a considerably body of work has described how inter-process

thermal asymmetries can drive cross-scale responses to warming (e.g., Dell *et al.* 2014;

Gibert *et al.* 2022; Gilbert *et al.* 2014; Lopez-Urrutia *et al.* 2006), in our analysis, inter-

process thermal asymmetries required accompanying intra-process thermal

asymmetries in order to affect competition. For example, a higher temperature

sensitivity of resource growth rate than conversion efficiency cannot, on its own, cause

a shift in competition between species (Fig. S8): there must also be interspecific

variation in the temperature sensitivities of the growth rates or conversion efficiencies of

the different resources to affect competition. More generally, we found that consumer

species (or their resources) that have the most divergent thermal traits will undergo the

largest shifts in competition with warming. Previous studies have demonstrated the

effects of warming on competition, usually based on differences in thermal optima

among species to generate effects of warming on competition (Simon & Amarasekare

2024; Smith & Amarasekare 2018). Here, we show for the first time that general effects

of warming on competition can emerge from interspecific differences in suboptimal

thermal traits alone, demonstrating how warming changes competition even in

conditions that are considered thermally benign for all species considered.

Conclusions

Making general predictions for the effects of warming on communities has been a major

challenge in ecology. The Metabolic Theory of Ecology, with its focus on the

temperature sensitivity of metabolic processes across biological scales, and Modern

Coexistence Theory, through its conceptual framework for species competition, have

both advanced our understanding of ecosystem responses to environmental change.

However, differences in the domains of these two fields, one of which focuses on

interspecific differences and the other which focuses on macroecological differences in

key metabolic processes as drivers of change, have prevented their integration to date.

Here we integrated concepts from both fields and combined this with empirical thermal

trait data to develop new, general theory for how competition changes with warming and

advance a more coherent understanding of global change. The next step will be to test

these predictions empirically. Doing so will facilitate better linkages between general

patterns in biological responses to warming and context-specific stewardship strategies

under rapid environmental change.

<u>Acknowledgements</u>

JRB, TNG, and MIO were supported by Discovery Grants from the Natural Science and

Engineering Research Council of Canada, and KD was supported by a Centre for

Ecosystem Management Postdoctoral Fellowship. We thank Megan Szojka for helpful

feedback on the manuscript.

<u>References</u>

Adler, P.B., Dalgleish, H.J. & Ellner, S.P. (2012). Forecasting plant community impacts of

climate variability and change: when do competitive interactions matter? *J. Ecol.*, 100,

478–487.

Adler, P.B., HilleRisLambers, J. & Levine, J.M. (2007). A niche for neutrality. *Ecol. Lett.*, 10, 95–

104.

Alexander, J.M., Diez, J.M. & Levine, J.M. (2015). Novel competitors shape species' responses

to climate change. *Nature*, 525, 515–518.

Amarasekare, P. & Johnson, C. (2017). Evolution of Thermal Reaction Norms in Seasonally

Varying Environments. *Am. Nat.*, 189, E31–E45.

Amarasekare, P. & Savage, V. (2012). A Framework for Elucidating the Temperature

Dependence of Fitness. *Am. Nat.*, 179, 178–191.

Bernhardt, J.R., Sunday, J.M. & O'Connor, M.I. (2018a). Metabolic theory and the temperature

size rule explain the temperature dependence of population carrying capacity. *Am. Nat.*,

In press.

Bernhardt, J.R., Sunday, J.M., Thompson, P.L. & O'Connor, M.I. (2018b). Nonlinear averaging

of thermal experience predicts population growth rates in a thermally variable

environment. *Proc. R. Soc. B Biol. Sci.*, 285, 20181076.

Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). Toward a Metabolic

Theory of Ecology. *Ecology*, 85, 1771–1789.

Casas Goncalves, G. & Amarasekare, P. (2021). Persistence of tri-trophic interactions in

seasonal environments. *J. Anim. Ecol.*, 90, 298–310.

Chesson, P. (1990). MacArthur's consumer-resource model. *Theor. Popul. Biol.*, 37, 26–38.

Chesson, P. (2000). General Theory of Competitive Coexistence in Spatially-Varying

Environments. *Theor. Popul. Biol.*, 58, 211–237.

Chesson, P. (2011). Ecological niches and diversity maintenance. *Res. Biodivers. Models Appl.*

InTech Rij. Croat., 43–60.

Chesson, P. & Kuang, J.J. (2008). The interaction between predation and competition. *Nature*,

456, 235–238.

Clegg, T. & Pawar, S. (2024). Variation in thermal physiology can drive the temperature-

dependence of microbial community richness. *eLife*, 13, e84662.

Coblentz, K.E., Novak, M. & DeLong, J.P. (2023). Predator feeding rates may often be

unsaturated under typical prey densities. *Ecol. Lett.*, 26, 302–312.

Dell, A.I., Pawar, S. & Savage, V.M. (2011). Systematic variation in the temperature

dependence of physiological and ecological traits. *Proc. Natl. Acad. Sci.*, 108, 10591–

10596.

Dell, A.I., Pawar, S. & Savage, V.M. (2014). Temperature dependence of trophic interactions

are driven by asymmetry of species responses and foraging strategy. *J. Anim. Ecol.*, 83,

70–84.

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., *et al.*

(2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proc. Natl.*

Acad. Sci., 105, 6668–6672.

Ehrlén, J. & Morris, W.F. (2015). Predicting changes in the distribution and abundance of

species under environmental change. *Ecol. Lett.*, 18, 303–314.

Enquist, B.J., Economo, E.P., Huxman, T.E., Allen, A.P., Ignace, D.D. & Gillooly, J.F. (2003).

Scaling Metabolism from organisims to ecosystems. *Nature*, 423, 639–642.

Germain, R.M., Mayfield, M.M. & Gilbert, B. (2018). The 'filtering' metaphor revisited:

competition and environment jointly structure invasibility and coexistence. *Biol. Lett.*, 14,

20180460.

Gibert, J.P., Grady, J.M. & Dell, A.I. (2022). Food web consequences of thermal asymmetries.

Funct. Ecol., 36, 1887–1899.

Gilbert, B., Tunney, T.D., McCann, K.S., DeLong, J.P., Vasseur, D.A., Savage, V., *et al.* (2014).

A bioenergetic framework for the temperature dependence of trophic interactions. *Ecol.*

Lett., 17, 902–914.

Godoy, O. (2019). Coexistence theory as a tool to understand biological invasions in species

interaction networks: Implications for the study of novel ecosystems. *Funct. Ecol.*, 33,

1190–1201.

Godoy, O. & Levine, J.M. (2014). Phenology effects on invasion success: insights from coupling

field experiments to coexistence theory. *Ecology*, 95, 726–736.

Godwin, C.M., Chang, F.-H. & Cardinale, B.J. (2020). An empiricist's guide to modern

coexistence theory for competitive communities. *Oikos*, 129, 1109–1127.

Grainger, T.N., Levine, J.M. & Gilbert, B. (2019). The Invasion Criterion: A Common Currency

for Ecological Research. *Trends Ecol. Evol.*, 34, 925–935.

Hubbell, S.P. (2001). The unified neutral theory of biodiversity and biogeography. In: *The unified*

neutral theory of biodiversity and biogeography. Princeton University Press.

Kingsolver, J.G. (2009). The Well-Temperatured Biologist: (American Society of Naturalists

Presidential Address). *Am. Nat.*, 174, 755–768.

Knies, J.L. & Kingsolver, J.G. (2010). Erroneous Arrhenius: Modified Arrhenius Model Best

Explains the Temperature Dependence of Ectotherm Fitness. *Am. Nat.*, 176, 227–233.

Kordas, R.L., Harley, C.D.G.G. & O'Connor, M.I. (2011). Community ecology in a warming

world: The influence of temperature on interspecific interactions in marine systems. *J.*

Exp. Mar. Biol. Ecol., 400, 218–226.

Kraft, N.J.B., Valencia, R. & Ackerly, D.D. (2008). Functional Traits and Niche-Based Tree

Community Assembly in an Amazonian Forest. *Science*, 322, 580–582.

Lande, R. (1993). Risks of Population Extinction from Demographic and Environmental

Stochasticity and Random Catastrophes. *Am. Nat.*, 142, 911–927.

Letten, A.D., Ke, P.J. & Fukami, T. (2017). Linking modern coexistence theory and

contemporary niche theory. *Ecol. Monogr.*, 87, 161–177.

Lopez-Urrutia, A., San Martin, E., Harris, R.P. & Irigoien, X. (2006). Scaling the metabolic

balance of the oceans. *Proc. Natl. Acad. Sci.*, 103, 8739–8744.

MacArthur, R. (1970). Species packing and competitive equilibrium for many species. *Theor.*

Popul. Biol., 1, 1–11.

Martin, T.L. & Huey, R.B. (2008). Why "Suboptimal" Is Optimal: Jensen's Inequality and

Ectotherm Thermal Preferences. *Am. Nat.*, 171, E102–E118.

Merow, C., Latimer, A.M., Wilson, A.M., McMahon, S.M., Rebelo, A.G. & Silander Jr, J.A.

(2014). On using integral projection models to generate demographically driven

predictions of species' distributions: development and validation using sparse data.

Ecography, 37, 1167–1183.

Michaletz, S.T. & Garen, J.C. (2024). Hotter is not (always) better: Embracing unimodal scaling

of biological rates with temperature. *Ecol. Lett.*, 27, e14381.

Novak, M., Coblentz, K.E. & DeLong, J.P. (2025). In Defense of Type I Functional Responses:

The Frequency and Population Dynamic Effects of Feeding on Multiple Prey at a Time.

Am. Nat., 206, 347–361.

O'Connor, M.I. (2009). Warming Strengthens an Herbivore: Plant Interaction. *Ecology*, 90, 388–

398.

O'Connor, M.I., Gilbert, B. & Brown, C. (2011). Theoretical Predictions for How Temperature

Affects the Dynamics of Interacting Herbivores and Plants. *Am. Nat.*, 178, 626–638.

Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts

across natural systems. *Nature*, 421, 37–42.

Pastore, A.I., Barabás, G., Bimler, M.D., Mayfield, M.M. & Miller, T.E. (2021). The evolution of

niche overlap and competitive differences. *Nat. Ecol. Evol.*, 5, 330–337.

Pawar, S., Dell, A.I., Savage, V.M. & Knies, J.L. (2016). Real versus Artificial Variation in the

Thermal Sensitivity of Biological Traits. *Am. Nat.*, 187, E41–E52.

Pérez-Ramos, I.M., Matías, L., Gómez-Aparicio, L. & Godoy, Ó. (2019). Functional traits and

phenotypic plasticity modulate species coexistence across contrasting climatic

conditions. *Nat. Commun.*, 10, 2555.

Peters, R.L. & Lovejoy, T.E. (1992). *Global warming and biological diversity*. Yale University

Press.

Rall, B.C., Vucic-Pestic, O., Ehnes, R.B., Emmerson, M. & Brose, U. (2010). Temperature,

predator-prey interaction strength and population stability: TEMPERATURE AND

INTERACTION STRENGTH. *Glob. Change Biol.*, 16, 2145–2157.

Román-Palacios, C. & Wiens, J.J. (2020). Recent responses to climate change reveal the

drivers of species extinction and survival. *Proc. Natl. Acad. Sci. U. S. A.*, 117, 4211–

4217.

Savage, V.M., Gillooly, J.F., Brown, J.H., West, G.B. & Charnov, E.L. (2004). Effects of body

size and temperature on population growth, 429–441.

Schreiber, S.J., Levine, J.M., Godoy, O., Kraft, N.J.B. & Hart, S.P. (2022). Does deterministic

coexistence theory matter in a finite world? *Ecology*, 104, e3838.

Siepielski, A.M., Hung, K.-L., Bein, E.E.B. & McPeek, M.A. (2010). Experimental evidence for

neutral community dynamics governing an insect assemblage. *Ecology*, 91, 847–857.

Simon, M.W. & Amarasekare, P. (2024). Predicting the fundamental thermal niche of

ectotherms. *Ecology*.

Smith, D.J. & Amarasekare, P. (2018). Toward a Mechanistic Understanding of Thermal Niche

Partitioning. *Am. Nat.*, 191, E57–E75.

Song, C., Barabás, G. & Saavedra, S. (2019). On the Consequences of the Interdependence of

Stabilizing and Equalizing Mechanisms. *Am. Nat.*, 194, 627–639.

Spaak, J.W., Carpentier, C. & De Laender, F. (2021). Species richness increases fitness

differences, but does not affect niche differences. *Ecol. Lett.*, 24, 2611–2623.

Spaak, J.W., Ke, P.-J., Letten, A.D. & De Laender, F. (2023). Different measures of niche and

fitness differences tell different tales. *Oikos*, 2023, e09573.

Sunday, J.M., Bates, A.E., Kearney, M.R., Colwell, R.K., Dulvy, N.K., Longino, J.T., *et al.*

(2014). Thermal-safety margins and the necessity of thermoregulatory behavior across

latitude and elevation. *Proc. Natl. Acad. Sci.*, 111, 5610–5615.

Terry, J.C.D. (2025). An Experimental Validation Test of Ecological Coexistence Theory to

Forecast Extinction Under Rising Temperatures. *Ecol. Lett.*, 28, e70047.

Tilman, D. (1994). Competition and Biodiversity in Spatially Structured Habitats. *Ecology*, 75, 2–

16.

Usinowicz, J. & Levine, J.M. (2018). Species persistence under climate change: a geographical

scale coexistence problem. *Ecol. Lett.*, 21, 1589–1603.

Vasseur, D.A. & McCann, K.S. (2005). A Mechanistic Approach for Modeling Temperature-

Dependent Consumer-Resource Dynamics. *Am. Nat.*, 166, 184–198.

Weiher, E., Clarke, G.D.P. & Keddy, P.A. (1998). Community Assembly Rules, Morphological

Dispersion, and the Coexistence of Plant Species. *Oikos*, 81, 309–322.

Welter, J.R., Benstead, J.P., Cross, W.F., Hood, J.M., Huryn, A.D., Johnson, P.W., *et al.* (2015).

Does N2 fixation amplify the temperature dependence of ecosystem metabolism?

Ecology, 92, 603–610.

Figure 1

A

B

Inter-process thermal asymmetry

C

Intra-process thermal asymmetry

Figure 2

Figure 3

Figure 4

Figure 1. Schematic of a consumer-resource model with temperature dependent

processes, yielding inter- and intra-process thermal asymmetries. In panel A, two

unique consumers are shown as boxes (N_1 and N_2) and two substitutable resources as

circles (R_a and R_b). Arrows show directional energy flow processes and are labeled with

the model parameters they represent: m_i is the mortality of consumer i, c_{ik} is

consumption rate of resource k by consumer i, and r_k is the intrinsic growth rate of

resource k. Parameters K_k, resource carrying capacity, and v_{ik}, conversion efficiency,

are not shown, but mediate the abundance of available resource, R_k, and the transfer of

energy from R_k to N_i, respectively. Panels B and C show how temperature sensitivity, E,

or the increase in the rate of a process with temperature below the temperature where

the curve peaks (central panels), is incorporated into the model via biological rates of

the consumer and resource, and how this can generate two types of thermal

asymmetries depending on where in the consumer-resource system a difference in

temperature sensitivity occurs (left panels). The ambient temperature is given as T_{amb},

and the rates of a process at this temperature, corresponding to $Y_{T_{amb}}$ in Eq. 9, are

given. The thickness of the arrows in the left panels corresponds to the temperature

sensitivity depicted in the central panels. The right panels in B and C relate a single

observation of temperature sensitivity (dashed line) for a parameter to the broader

empirical distribution of temperature sensitivities for that parameter constructed from our

data synthesis. In an inter-process thermal asymmetry (Panel B), the two temperature

sensitivities come from independent distributions of temperature sensitivity in each

parameter, whereas in an intra-process thermal asymmetry (Panel C), the two

temperature sensitivities come from the same distribution of temperature sensitivity.

Figure 2. Posterior distributions for temperature sensitivity of consumer-resource

model parameters. Panels A-E show published empirical observations for each

parameter in the consumer resource model (Eq. 4) in orange circles and posterior

distributions informed by these observations in grey density plots. Red lines indicate the

mean of the posterior distribution. Panel F shows the means and 95% credible intervals

(CIs) for each parameter.

Figure 3. Effects of temperature sensitivity in each model parameter on

competition with warming. In each panel, temperature sensitivities of the focal

parameter are drawn randomly from the parameter's empirical distribution (Fig. 2) and

then the system is warmed 15°C, while temperature sensitivities for all other parameters

are held at 0. This is repeated 500 times and each simulated warming path is plotted.

All simulations begin at the same ambient temperature, 10°C (black point). For all

parameters except for consumption rate (i.e. panels B-E), the number of temperature

sensitivities drawn for each simulation is two -- one for each consumer or resource

species (Fig. 1A). This results in perfectly overlain trajectories for each simulation: note

different lengths of the temperature gradient path, which ends with a dark red color. For

consumption rate, in panel A, there are four temperature sensitivities drawn for each

simulation, corresponding to one temperature sensitivity per resource, for each species

(Fig. 1A). As a result, warming trajectories do not fall perfectly on top of one another.

The shaded region in the plots shows where species can coexist, i.e., where $ND >$

$|FD|$, and the unshaded regions show where one species competitively excludes the

other, i.e., $ND < |FD|$.

Figure 4. Shifts in competition with warming are related to intra-process thermal

asymmetries. Panels A-E show displacement of species pairs with warming as a

function of thermal asymmetries in each parameter in the consumer-resource model,

and panel F depicts displacement of species pairs with warming on canonical niche

difference-fitness difference axes. Thermal asymmetries are given as the absolute value

of the difference between two draws of temperature sensitivities (one for each

consumer or resource species) in each simulation and displacement is given as the

Euclidean distance between a species pair's starting position and its position after 15°C

warming. For each panel (A-E), only the focal temperature sensitivity parameter was

allowed to vary; all other temperature sensitivity parameters were fixed at 0. Data points

correspond to the difference in temperature sensitivities drawn for each of 500

simulations, and the color of the points indicates the direction of the thermal asymmetry,

i.e. which E_Y value is larger. For consumption rate (panel A), we vary only two of the

four possible temperature sensitivities, here species N_1's consumption rate of each

resource, to display the effect of this thermal asymmetry most clearly.

Figure 5. Effects of warming on competition. All parameters are drawn from their

distributions and then warmed 15°C, and this simulation is repeated 500 times. A)

Simulation path for each draw of parameter values from the ambient temperature, 10°C

(yellow dot), to 25°C (15°C warming). The median position along both axes after 15°C

warming is plotted as a magenta dot. B) Distribution of Euclidean distances between the

position at the ambient temperature and after 15°C warming across 500 simulations. C)

Shifts in niche differences with warming. The median niche difference values (magenta

dot) and the niche difference values for each simulation (grey dots) after 15°C warming

are shown compared to the value at the ambient temperature (yellow dot). D) Shifts in

fitness differences with warming, displayed as in panel C.