ETEC heat-labile toxin promotes β-catenin stabilization and transcriptional reprogramming to disrupt

intestinal epithelial differentiation

Alaullah Sheikh[1*], Bipul Setu[1], Nikhilesh Joardar[1,2], John C. Martin[1], Bruce A. Rosa, [1] Makedonka

Mitreva[1,3,4], James M. Fleckenstein[1,5*]

[1]Division of Infectious Diseases, Department of Medicine, Washington University in Saint Louis School
of Medicine, Saint Louis, Missouri, USA; [2]Department of Bioengineering, School of Engineering and
Technology, Dr. Vishwanath Karad MIT World Peace University, Maharashtra, India, [3]Department of
Genetics, Washington University in Saint Louis School of Medicine, Saint Louis, Missouri, USA;
[4]McDonnell Genome Institute, Washington University in Saint Louis School of Medicine, Saint Louis,
Missouri, USA; [5]Medicine Service, Infectious Disease Section, Veterans Affairs Health Care System, Saint
Louis, Missouri, USA.
12

*Co-corresponding authors

Alaullah Sheikh
Division of Infectious Diseases
Department of Medicine
Washington University School of Medicine
Campus Box 8051
660 South Euclid Avenue
Saint Louis Missouri, USA 63110
p 314-362-9219
asheikh@wustl.edu
23
James M. Fleckenstein
Division of Infectious Diseases
Department of Medicine
Washington University School of Medicine
Campus Box 8051
660 South Euclid Avenue
Saint Louis Missouri, USA 63110
p 314-362-9218
jfleckenstein@wustl.edu
33

Abstract

Enterotoxigenic *Escherichia coli* (ETEC), defined by their production of heat-labile (LT) and heat-stable (ST) enterotoxins, are a common cause of acute diarrheal illness in children from low- and middle-income countries, and are also linked to long-term sequelae such as malnutrition, and growth impairment. While the mechanisms underlying toxin-mediated acute diarrhea are known, the molecular events involved in ETEC related sequelae remain unclear. Here, we demonstrate that the ETEC heat-labile toxin (LT) profoundly remodels intestinal epithelial composition and function through modulation of WNT/β-catenin signaling. Using human ileal enteroids, we demonstrate that LT stabilizes β-catenin independently of WNT ligands, promotes its nuclear accumulation, and enhances TCF/LEF-driven transcription. Single-cell transcriptomic analyses reveal that LT increases intestinal proliferation by enhancing cell cycle activity across all epithelial lineages, thus disrupting epithelial composition by expanding proliferative progenitor populations at the expense of absorptive enterocytes. Simultaneously, LT impairs epithelial maturation and suppresses transcriptional programs required for nutrient absorption and differentiation. Together, these findings identify LT as a potent driver of intestinal epithelial reprogramming, providing mechanistic insight into how ETEC infection may drive long-term consequences beyond acute diarrhea and may inform strategies to prevent major sequelae, including malnutrition, that affect millions of children worldwide.

Introduction

The enterotoxigenic *E. coli* (ETEC) are a diverse group of diarrheal pathogens defined by their production of heat-labile (LT) and/or heat-stable (ST) enterotoxins[1]. The ETEC pathovar is thought to be responsible for hundreds of millions of cases of diarrheal illness each year, primarily among young children of low-middle income countries (LMICs)[2]. Children of LMICs frequently suffer multiple episodes of ETEC diarrhea before in the first few years of life[3], and ETEC are a leading cause of death due to diarrheal illness in children under the age of five years[4]. While the death rate from acute diarrheal illness has declined with advances in oral rehydration therapy and other measures, children with ETEC diarrhea are also at increased risk of death after resolution of the initial episode[5]. Moreover, children with moderate-severe diarrhea[6] as well as less severe diarrhea[7] due to ETEC inexplicably suffer long-term sequelae including from a condition referred to as environmental enteric dysfunction (EED) characterized histologically by atrophy of small intestinal villi accompanied by expansion of crypts[8,9], and clinically by poor growth and malnutrition. Unfortunately, in the absence of a licensed vaccine, morbidity due to acute diarrheal illness as well as long-term sequelae of impaired growth and malnutrition have continued unabated[7].

The fundamental molecular pathogenesis of acute diarrheal illness caused by ETEC is well-established. Delivery of LT and ST activates production of cAMP and cGMP cyclic nucleotides, respectively. Increases in intracellular cAMP leads to activation of protein kinase A (PKA) while cGMP activates protein kinase G (PKG). Kinase-mediated phosphorylation of cellular ion channels in enterocyte membranes leads to the net export of salt and water into the intestinal lumen, culminating in profuse watery diarrhea characteristic of ETEC.

In contrast, the sequelae linked to ETEC are still poorly understood. Recently, however we demonstrated that heat-labile toxin modulates the transcription of hundreds of genes in target intestinal epithelial cells, including those responsible for the biogenesis of enterocyte microvilli responsible

nutrient absorption[10]. Moreover, repeated challenge of infant mice with toxigenic ETEC leads to impaired

growth[10] similar to observations in children. These studies provided the first suggestion that ETEC toxins

might drive molecular events that culminate in sequelae.

Nutrient absorption in the small intestine depends on tightly regulated epithelial differentiation and

maturation processes that generate functionally specialized cell types. Intestinal stem cells (ISCs),

residing at the base of crypts, undergo proliferation and lineage specification to give rise to absorptive

enterocytes and secretory lineages including goblet, Paneth, and enteroendocrine cells[11]. As these cells

migrate along the crypt-villus axis, they progressively differentiate, acquiring mature phenotypes and

functional attributes[12]. In particular, enterocytes undergo a stepwise transcriptional and morphological

maturation program culminating in the expression of solute carrier (SLC) transporters, and other

molecules essential for nutrient uptake and barrier function[13]. Theoretically, disruption of this

developmental trajectory would compromise nutrient absorption and contribute to malnutrition among

children from low-resource settings[8,14-17].

Intestinal epithelial maturation is tightly regulated. The luminal surface of the small intestine is

comprised of millions of villi, tiny (~0.5 mm long) fingerlike projections that are surrounded at their base

by invaginations known as crypts. As epithelial cells covering the villi are propagated from intestinal stem

cells (ISC) residing in the crypts they quickly differentiate into mature epithelial cells, with absorptive

enterocytes covered in microvilli making up the majority. These mature cells on the villous surface

undergo continual renewal over a period of 3-5 days[18] as they are constantly replenished by the

population of crypt stem cells. Differentiation of ISC into enterocytes and other cell types along the

crypt-villus axis is controlled by a number of signaling molecules including WNT[19-21].

High concentrations of WNT generated by adjacent Paneth cells[22] favor the maintenance and

proliferation of crypt (ISC) populations. In the absence of WNT, a "destruction complex" comprised of

GSK-3 β, CK1α, APC, and Axin2 engages β-catenin. GSK-3 β phosphorylation of β-catenin targets it for

ubiquitination[23] and removal from the cytoplasm. Binding of WNT to its cognate LRP5/6-Frizzled co-

receptors on the cell surface activates Dishevelled, a cytoplasmic protein that antagonizes the

destruction complex[24]. This permits accumulation of monomeric β-catenin in the cytoplasm[25] and

ultimately its translocation to the nucleus where it binds to T-cell factor (TCF) and lymphoid enhancer

factor (LEF) transcription factors[26], leading to activation of WNT/TCF target genes responsible for

maintenance of the crypt progenitor cells. In addition to canonical WNT signaling, β-catenin can be

stabilized both directly and indirectly by modulating features of the destruction complex. Here, we

demonstrate that heat-labile enterotoxin of ETEC enhances β-catenin stabilization to activate

downstream signaling in small intestinal epithelia, disrupting differentiation and maturation required for

efficient nutrient uptake.

Results

Heat-labile toxin activates transcription of multiple WNT-responsive genes

Bioinformatic analysis (CompBio v2.9) of earlier bulk RNAseq data[10] from small intestinal enteroids

treated heat-labile toxin revealed that transcription of genes governed by the WNT-β-catenin pathway

were significantly upregulated in response to LT (figure 1a). Upregulated genes included Axin2, RNF43,

MYC, SOX9, cyclin D1 (CCND1), EphB2, LGR5, PROM1, DKK1, CLDN1, TSPAN8, along with a number of

other genes governing the intestinal cell differentiation including OLFM4, CD24, REG4, XBP1, and

ATOH1[21,27,28]. Using CEACAM6 as a positive control[29] for LT-induced gene expression, we validated impact

on genes in this pathway by qRT-PCR (Figure 1b). In keeping with these results, we also observed cellular

accumulation of β-catenin following treatment of polarized small intestinal enteroids with LT (Figure

1c,d).

To functionally assess transcriptional modulation by β-catenin, we stably transfected small intestinal

enteroids with a lentivirus vector construct (7TFP)[30] which contains 7 copies of the TCF/LEF transcription

factor DNA-binding site 5'-CCTTTGATC-3' upstream of firefly luciferase[31]. Treatment of 7TFP carrying

reporter enteroids with LT resulted in significant increases in TCF/LEF-mediated transcriptional activity

(Figure 1e). These effects were abrogated when the enzymatically inactive mutant form of LT (mLT) was

used, indicating that catalytic activity of LT is required for this activation.

Heat-labile toxin stabilizes cytosolic β-catenin to facilitate nuclear translocation

In canonical WNT signaling pathways, β-catenin accumulation in the nucleus permits engagement of

TCF/LEF transcription factors to activate transcription of downstream genes (Figure 2a). Interestingly,

activation of cyclic nucleotide pathways has the potential to interfere with degradation of β-catenin. PKA

phosphorylation of β-catenin at position Ser_{675} inhibits β-catenin ubiquitination, thereby permitting its

accumulation[32], while activation of PKA can also result in phosphorylation of GSK-3β at Ser_9 resulting in

its inhibition[33-37]. Similarly, cGMP activation of PKG also results in phosphorylation of GSK-3β at Ser_9[38,39]

to inhibit enzymatic activity. Accordingly, we observed time-dependent increases in total β-catenin in

both cytoplasmic and nuclear compartments (Figure 2b-e). In parallel, LT induced phosphorylation of β-

catenin at Ser_{675} (Figure 2b-e, Supplementary figure 1), a known activating modification that promotes β-

catenin nuclear localization and transcriptional activity[40], further supporting functional stabilization.

In addition, we found that LT modulated phosphorylation of GSK-3β, a central negative regulator of β-

catenin stability. Although total protein levels remained relatively unchanged, GSK-3β phosphorylation at

the Ser_9 residue increased significantly in LT-treated samples over time (Figure 2b, d).

Collectively, these data demonstrate that heat-labile toxin acts to stabilize β-catenin independent of

WNT by directed post translational modification of β-catenin and elements of the destruction

complex[32,41]. Interestingly, we also noted that transcription of the *CTNNB1* gene encoding β-catenin was

itself upregulated in response to LT, potentially related to the presence of cAMP response elements[42] in

the CTNNB1 promoter[43].

Heat-labile toxin disrupts epithelial cell composition

Given the central role of WNT/β-catenin signaling in regulating intestinal stem cells[20] and their

differentiation to different epithelial cell types[18], we next investigated whether LT-induced stabilization of

β-catenin impacts epithelial cell dynamics. In the intestine, proliferative expansion without coordinated

differentiation can shift the cellular composition and impair epithelial function[44]. To investigate whether

heat-labile toxin perturbs this balance, we used single-cell RNA sequencing to examine the impact of LT

on differentiation of human small intestinal enteroids. Control (∅) and LT-treated enteroid cells were

harvested and subjected to Illumina NovaSeq X Plus single-cell RNA sequencing, followed by analysis

with Seurat (v5.0)[45]. After quality control filtering, 1,963 cells represented by an average of 14,342 reads

and 3,266 genes were sequenced in the control sample, and 2207 cells, an average of 14,342 reads and

3283 genes in the LT sample. To assess changes in epithelial cell composition, we then performed

unsupervised graph-based clustering of single-cell transcriptomes from control and LT-treated enteroids

(resolution setting of 0.92, defining 14 clusters), followed by annotation using canonical lineage

markers[11,13], confidently identifying all expected major cell types (Figure 3a; Table 1; **Supplementary**

dataset 1).

LT treatment induced striking alterations in enteroid cell composition. All major intestinal epithelial

cell types, including absorptive enterocytes, goblet cells, Paneth cells, and enteroendocrine cells, along

with intermediate populations likely representing transitional states along the differentiation axis, were

identified in both control and LT-treated samples (Figure 3a–c). However, a highly proliferative cluster,

largely absent in controls, was markedly expanded following LT exposure, while a differentiated

population present only in control samples was lost (Figure 3d–e). Differential abundance analysis

confirmed enrichment of early-stage progenitor and transit-amplifying populations (Clusters 0, 7, and 8)

in LT-treated samples, alongside significant depletion of mature enterocytes (Clusters 4 and 5) and

BEST4[+] absorptive cells (Cluster 13) (Figure 3e; Table 1).

To further explore the identity of these transcriptionally distinct or depleted populations (Clusters 7,

and 13, respectively) in LT-treated enteroids, we performed over-representation analysis (ORA). ORA of

cluster 7 revealed strong enrichment for mitotic cell cycle pathways, including G2/M transition, spindle

checkpoint, and DNA replication, indicating a highly proliferative state (Supplementary figure 2a).

Consistently, transcription factor analysis identified dominant enrichment of E2F family regulators (E2F1,

E2F2, E2F3) and their partners DP1/DP2, which are key drivers of S-phase entry and DNA synthesis

(Supplementary figure 2b). Additional enriched motifs included MYC, NFY, and YY1, supporting enhanced

biosynthetic and metabolic activity consistent with transit-amplifying cells. In contrast, Cluster 13 was

enriched for pathways associated with terminal epithelial differentiation and absorptive function. These

included mitochondrial respiration, oxidative phosphorylation, the TCA cycle, and lipid metabolism

(Supplementary figure 2a). The absence of proliferative transcription factor signatures in Cluster 13

further supports its identity as a mature enterocyte population (Supplementary figure 2b). Together,

these findings reveal that LT exposure promotes the expansion of transcriptionally active proliferating

cells while depleting functionally mature absorptive cells, disrupting the balance between proliferation

and differentiation in the intestinal epithelium.

To gain deeper insight into the transcriptional alterations underlying these shifts, we conducted

differential gene expression analysis on all cell clusters, identifying thousands of differentially expressed

genes (DEGs) significantly upregulated or downregulated in LT-treated enteroids compared to control

(Table 2, **Supplementary dataset 2**). To assess functional implications, we performed over-representation

analysis (ORA) of differentially expressed genes. Upregulated DEGs were enriched for pathways in RNA

metabolism, cell cycle regulation, and stress response, consistent with a proliferative epithelial state

(Supplementary figure 3). Downregulated DEGs were enriched for bile acid transport, lipid metabolism,

cholesterol biosynthesis, and other absorptive functions, indicating impaired maturation and loss of

differentiation (Supplementary figure 3). Gene ontology independent analysis (CompBio v2.9) likewise

highlighted suppression of metabolic and transport functions, including intestinal absorption, lipid and

bile metabolism, fatty acid uptake, amino acid metabolism, and cholesterol processing (Supplementary

figure 4, also see **Supplementary dataset 2**). Taken together, these data indicate that LT exposure drives

a shift in epithelial composition by promoting proliferation and suppressing differentiation.

Heat-labile toxin promotes intestinal epithelial cell proliferation and reshapes cell cycle dynamics

across epithelial cell clusters

Because proliferation kinetics directly influence epithelial cell-type composition[44], we next examined

how LT affects the cell cycle status of intestinal epithelial cells. Using canonical gene sets for G1, S, and

G2/M phases[46] (Supplementary figure 5a), we assigned cell cycle stages to individual cells in LT-treated

and control enteroids. UMAP projections revealed a marked shift in overall cell cycle distribution upon LT

exposure (Figure 4a). Specifically, LT-treated cells showed a significant expansion of S-phase (p = 2.21 x

10^{-5}, χ^2) and G2/M-phase populations (p <1 x 10^{-16}), accompanied by a reduction in G1-phase cells (Table

3).

Cell cycle phase scores further confirmed these shifts, with elevated expression of S-phase and G2/M

gene sets in LT-treated enteroids (Figure 4b). Consistent with this, we observed differential expression of

key phase-specific cell cycle regulators, including RB1 and CDKN1A (G1-phase), PCNA and MCM3 (S-

phase), and CCNB2 and CDK1 (G2/M-phase) (Figure 4c). LT exposure also upregulated stem and

progenitor markers such as CD24, CD44, and OLFM4, reinforcing the expansion of the undifferentiated,

proliferative compartment (Supplementary figure 5b).

Compartmentalization of intestinal cell proliferation is critical for epithelial maintenance and function

with higher proliferative activity at the crypt zone and no proliferation at the tips of villi. Analysis of cell

cycle dynamics across transcriptionally defined clusters revealed a widespread reduction in G1-phase

cells and robust enrichment of G2/M-phase cells across epithelial subpopulations (Figure 5a; Table 3),

indicating that LT alters cell cycle dynamics across the entire epithelial landscape.

To further support the observed shift in cell cycle dynamics, we analyzed the expression of canonical

proliferation markers. LT-treated enteroids exhibited strong upregulation of MKI67 and TOP2A, two well-

established indicators of active proliferation[47] (Figure 5b). To validate these transcriptomic findings at the

protein level, we performed phenotypic assays using polarized epithelial monolayers. Enteroids were

cultured on Transwell filters to establish apical–basolateral polarity and treated with LT during the

differentiation phase. Confocal immunofluorescence microscopy revealed a significant increase in

MKI67$^+$ nuclei in LT-treated monolayers compared to controls (Figure 5c-d), confirming enhanced

proliferative activity at the protein level.

To functionally validate these observations, we utilized 5-ethynyl-2'-deoxyuridine (EdU) incorporation

assay in 3D culture of enteroids treated with LT. Following differentiation, both control and LT-treated

enteroids were incubated with EdU. LT treatment led to significantly higher incorporation of EdU

(p<0.01), indicating enhanced DNA synthesis and entry into S phase (Figure 5e–f). Collectively, these

findings indicate that LT skews epithelial homeostasis by promoting a proliferative, progenitor-dominated

state across the intestinal epithelium.

Heat-labile toxin suppresses epithelial differentiation and impairs functional maturation

Given this marked shift away from differentiated lineages, we next investigated whether LT-induced

disruption of differentiation affects epithelial maturation. Villin, a structural component of apical

microvilli, and intestinal alkaline phosphatase (ALPI), a brush-border enzyme, serve as canonical markers

of mature enterocytes[12,48]. To assess the functional consequences of LT-induced differentiation defects,

we examined villin and ALPI protein expression in membrane and cytosolic fractions, respectively, from

control and LT-treated enteroids. Immunoblot analysis showed significantly reduced expression of villin,

and alkaline phosphatase (ALPI), in LT-treated enteroids compared to controls (Figure 6a–b), suggesting

impaired epithelial maturation at the protein level.

To further investigate the impact of LT on the functional maturation of the epithelia, we examined the

expression of key solute carrier (SLC) genes responsible for nutrient transport. Although we observed

robust expression of multiple SLC family members in control enteroids, LT-treated enteroids exhibited

marked repression of these genes across the epithelial landscape (Supplementary figure 6). Notably, LT

diminished the expression of SLC28A1 (nucleoside), SLC23A1 (vitamin C), SLC30A1/30A4 (zinc), SLC7A7

(amino acids), SLC5A11 (myo-inositol), SLC19A3 (thiamine), SLC11A2 (iron), SLC17A4 (glutamate),

SLC15A1 (di/tripeptides), and other transporters, consistent with a depletion of functionally mature

enterocytes.

Because nutrient absorption is a key functional hallmark of mature enterocytes, we next analyzed the

transcriptional consequences of LT treatment on nutrient transporter expression across five

transcriptionally defined enterocyte subclusters (EC1–EC5). LT exposure led to a global reduction in

transporter gene expression scores across multiple categories, including sugar, amino acid, water, bile

acid, lipid, and vitamin transport (Figure 6c, left). This was further supported by reduced average

expressions and cell-level prevalence of multiple solute carrier (SLC) family genes[12] in LT-treated cells

compared to controls (Figure 6c, right).

Consistent with these observations, prior bulk RNA-seq studies have shown that LT suppresses

SLC19A3, a thiamine transporter, along with its upstream regulator SP1, leading to impaired thiamine

uptake[49]. Our single-cell RNA-seq data recapitulated this finding, with decreased expression of both

SLC19A3 and SP1 in LT-treated enteroids (Figure 6d). Additionally, differential gene expression analysis of

enterocyte clusters revealed significant downregulation of other key absorptive genes, including SLC5A1

(sodium/glucose cotransporter 1), SLC10A2 (ileal sodium/bile acid cotransporter), SLC28A1

(sodium/nucleoside cotransporter 1), FABP6 (fatty acid transporter), and APOB (fatty acid transporter),

confirming suppression of essential nutrient acquisition pathways (Figure 6e).

Altogether, these studies suggest that LT exerts a profound effect on the developing intestinal

epithelia. The LT-driven proliferative response coincided with transcriptional activation of WNT/β-catenin

target genes and nuclear accumulation of β-catenin, consistent with canonical WNT pathway activation.

By impacting major transcriptional pathways, including those orchestrated by β-catenin, LT disrupts

epithelial differentiation and subsequent maturation.

Discussion

ETEC infections have repeatedly been linked to long-term sequelae including malnutrition and impaired growth, significantly compounding the morbidity burden beyond the acute infection [5,8,15,16,50,51]. While LT-producing ETEC have specifically been linked to malnutrition among young children[50], the precise molecular mechanisms underlying links between ETEC and nutritional sequelae remain incompletely understood. The present studies suggest that LT may alter the trajectory of epithelial cell maturation, stalling the development of mature absorptive enterocytes. This disruption of normal epithelial homeostasis parallels earlier observations that LT also interferes with transcriptional programs that govern the biogenesis of enterocyte microvilli, comprising the major surface for nutrient absorption, along with LT modulation of a multitude of nutrient transporters[52]. The collective impacts of LT on the architecture and function of absorptive intestinal epithelia likely contribute significantly to the nutritional deficits that accompany ETEC infections in young children.

Here we show that the heat-labile enterotoxin reprograms intestinal epithelia by subversion of the fundamental regulation of stem cell maintenance, proliferation, and differentiation[18,53,54] governed by β-catenin. While LT is classically known for inducing diarrhea via cAMP-mediated chloride secretion[55,56], its broader impact on intestinal development has remained poorly characterized. Here, we show that LT activates β-catenin signaling independent of WNT by stabilizing β-catenin through PKA-mediated phosphorylation of β-catenin and components of its degradation complex[32,57,58]. The resulting enhanced and aberrant β-catenin signaling ultimately reprograms homeostatic maturation of intestinal epithelia.

The intricate nature of cAMP and PKA signaling is reflected in complex array of changes that follow exposure to heat-heat labile toxin. Mechanistically, LT modulates transcriptional programs by PKA-mediated phosphorylation of a family of cAMP-responsive nuclear factors including the cAMP response element binding protein (CREB) and the cAMP response element modulator (CREM), permitting their interaction with transcriptional co-activators including CREB-binding protein (CBP) as they bind

cooperatively to short DNA consensus sequences known as cAMP response elements (CRE)[59] in the

promoter regions of target genes. Intriguingly, genetic studies in Bangladeshi children identified CREM

polymorphisms associated with increased susceptibility to diarrhea and reduced weight-for-age[60],

potentially supporting the involvement of cAMP signaling in nutritional sequelae.

The cAMP pathway is also at the center of complex regulatory cascades involving multiple

transcription factors in addition to CREB and CREM that are also modulated by PKA[59]. These include the

transcriptional regulators HNF4A and SMAD4, both essential for enterocyte differentiation and known to

be sensitive to PKA-dependent phosphorylation[61,62 63-65]. Beyond direct transcriptional effects, cAMP-

driven activation of co-regulators such as CBP and PHF2 can induce histone acetylation and methylation,

also raising the possibility that LT reinforces impaired epithelial maturation through epigenetic

remodeling[62].

Our data supports the view that LT activates β-catenin signaling to impair terminal enterocyte

differentiation. Under homeostatic conditions, β-catenin regulates the balance between stem cell

renewal and lineage commitment in the crypt-villus axis[21,86-88]. Disruption of this axis, such as through

loss of WNT antagonists like APC, leads to crypt expansion and impaired epithelial maturation [21,66,67]. Our

findings align with this paradigm, showing that LT mimics hyperactive WNT signaling to expand

progenitor and transit-amplifying populations while depleting differentiated lineages, including mature

enterocytes and BEST4[+] absorptive cells[68]. This imbalance is characteristic of a poorly differentiated

epithelium and closely mirrors the pathological features of environmental enteric dysfunction (EED),

characterized by villus blunting, crypt expansion, and malabsorption[69,70].

The depletion of mature absorptive cells and downregulation of nutrient transporter genes in LT-

treated enteroids recapitulate key features of immature epithelia. Epithelial maturation involves

functional enrichment of nutrient absorption capacity in villus tip cells, where genes required for brush

border formation and fat absorption are upregulated in differentiated villus enterocytes[12]. However, LT-

treated epithelia exhibit reduced expression of ALPI, and VIL1, canonical markers of enterocyte

maturation, as well as transcriptional repression of multiple solute carrier family genes essential for

absorptive functions of nutrient, bile acid, and vitamin transport. The failure to achieve mature

enterocyte identity is particularly relevant in the context of EED, where altered bile acid metabolism and

nutrient malabsorption are linked to stunted growth. Notably, a study in undernourished Pakistani

infants found that disrupted bile acid homeostasis correlated with EED histopathology and growth

faltering[71]. LT-induced repression of bile acid transporters (SLC10A2, FABP6) may offer a plausible

mechanistic link to these clinical observations.

This concept is further supported by recent animal studies, where bile acid accumulation and NAD^+

deficiency induced by a low-protein diet led to villus atrophy and epithelial loss, phenotypes that were

reversed by dual supplementation with NAD^+ precursors and bile acid sequestrants [72]. Our findings raise

the possibility that LT could act synergistically with nutritional stressors to maintain an immature

epithelial state, exacerbating malabsorption and stunting in vulnerable children. The ability of LT to

disrupt intestinal maturation and impair nutrient uptake underscores the pathogenic potential of ETEC

beyond acute diarrhea.

From a broader perspective, this work positions LT as a potential molecular link between ETEC

infection and long-term intestinal dysfunction. The sustained shift toward a proliferative,

undifferentiated epithelial state, driven by toxin-mediated signaling, mirrors intestinal phenotypes

observed in chronically infected or malnourished children, where impaired differentiation contributes to

long-term functional deficits[8,16,17,73]. These findings are especially concerning in the context of repeated

or chronic ETEC exposure, which is common in low-resource settings and has been linked to growth

faltering in children even in the absence of overt diarrhea[15,50,51,74-78].

While this study provides mechanistic insight into LT-induced epithelial reprogramming, several

limitations should be noted. First, the findings are based on *in vitro* enteroid models, which lack stromal,

immune, and microbial components that shape epithelial responses *in vivo.* Second, although we

identify β-catenin signaling as a key mediator, the specific upstream effectors connecting LT-induced

cAMP production to transcriptional reprogramming remain to be fully defined. In addition to LT, many

ETEC make heat-stable toxins (ST) which engage guanylate cyclase C (GC-C) to stimulate production of

cGMP and activation of protein kinase G (PKG). Although ST has not been thoroughly evaluated in the

context of EED, ST and other GC-C agonists are known to impact cell cycle progression and modulate

epithelial proliferation and differentiation[79-81].

Understanding how these common enterotoxins contribute to pathogenesis of intestinal dysfunction

that follow ETEC infections can inform the rational design of vaccines that protect against acute illness as

well as the tremendous burden of sequelae[52,82]. This work advances our understanding of how ETEC

contributes to long-term intestinal dysfunction and malnutrition and offers a framework for developing

strategies to mitigate the persistent sequelae of enteric infections in vulnerable populations.

Materials and methods

All experiments were conducted in accordance with relevant ethical guidelines, and study protocols were approved by the Institutional Biosafety Committee (IBC), and Institutional Review Board (IRB) at Washington University School of Medicine in St. Louis.

Growth and maintenance of small intestinal enteroid cell lines

Hu235D derived from human small intestinal (ileal) stem cells were obtained from the Precision Animal Models and Organoids Core (PAMOC) maintained by the Washington University Digestive Diseases Research Core Center (DDRCC), and were propagated as previously described[83]. Briefly, crypt intestinal stem cells maintained in liquid nitrogen were thawed, resuspended in Cultrex (Cultrex BME Type II, Catalog# 3532-010-02, Biotechne) and transferred to wells in a 48-well plate. Cells were grown for 3-4 days in growth media containing 50% L-WRN conditioned media derived from the L-WRN cell line (provided by the POMAC), a 1:1 mixture of primary media [Advanced DMEM/F12, Invitrogen 12634028, supplemented with fetal bovine serum (FBS, 20%), 2 mM L-glutamine, 100 U/ml penicillin, 0.1 mg/ml streptomycin], 10 µM ROCK inhibitor Y-27632 (Tocris Bioscience) and 10 µM TGFBR1 inhibitor SB 431542 (Tocris Bioscience) [83][84].

For RNA-seq analysis, cells were grown in Cultex for 4 days for enteroid formation, diluted 1:2 in Cultrex to reduce enteroid density, replated and grew for 3 more days in growth media. To allow differentiation, enteroids were then cultured for additional 3 days in differentiation media (5% L-WRN conditioned media supplemented with 10 µM ROCK inhibitor). Enteroids were treated with heat-labile toxin (100ng/ml) either during the differentiation period (LT), or PBS treated (control). For polarized enteroid monolayers, following trypsinization cells were seeded onto Transwell filters (Corning) coated with type IV human collagen (Sigma) and grew in growth media for confluent monolayer. After confluency growth, media was replaced with differentiation media to promote differentiation.

Quantitative RT-PCR

Quantitative real-time PCR was conducted on a QuantStudio 3 detection system (Applied Biosystems) using Fast SYBR Green master mix (Applied Biosystems/Thermo Fisher). Specificity of amplification for each sample was confirmed by dissociation curve analysis, and PCR product size was verified through agarose gel electrophoresis. For transcript amplification from small intestinal enteroids, previously described methods were followed, with relative gene expression normalized to GAPDH. All primers used are provided in Supplementary table 1.

Fluorescence microscopy

For fluorescence microscopy, polarized enteroid monolayers were fixed in 2% paraformaldehyde for 30 minutes at 37°C, followed by an additional 30 minutes at room temperature. After fixation, the monolayers were washed three times with PBS and blocked with 2% BSA-PBS to minimize non-specific binding. The primary antibodies (as detailed in Supplementary table 2) used for labeling specific proteins were diluted in 2% BSA-PBS and incubated with the monolayers for 1 hour at room temperature. Fluorescent dye-conjugated secondary antibodies were then diluted in 2% BSA-PBS and applied to the monolayers for 45 minutes at room temperature. Cell membranes and nuclei were stained using CellMask (1:2000 dilution, Invitrogen) and DAPI (1:1000 dilution, Invitrogen), respectively. Slides were washed 3x with PBS and coverslipped with Prolong Gold antifade reagent (Invitrogen). Images were captured and analyzed using a Nikon C2 confocal microscope equipped with NIS-Elements AR 5.11.01 software (Nikon).

Construction of TCF/LEF reporter enteroid lines.

HEK293T cells (ATCC CRL-3216) were propagated at 37°C with 5% CO_2 in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% heat-inactivated fetal bovine serum (FBS) and 2 mM L-glutamine in T25 flasks. For 7TFP containing lentivirus production, cells were co-transfected with 5 µg of

p7TFP, 5 µg of psPAX2, and 3 µg of pMD2.G plasmids (see Supplementary table 3) in a 500 µl of culture

medium with 39 µl of FuGENE HD transfection reagent (Promega). Media was changed following 18 h

post transfection. Two days post transfection, the 7TFP-lentivirus–containing supernatant was collected,

clarified by centrifugation at 3,000 rpm for 15 minutes, and concentrated by adding polyethylene glycol

(PEG) 8000 at a 1:3 ratio (PEG:supernatant). The mixture was rotated overnight at 4°C and centrifuged at

1,600 × g for 1 hour. The resulting viral pellet was resuspended in 50% conditioned medium and used to

transduce Hu235D human small intestinal enteroids, in the presence of polybrene (final concentration:

8 µg/ml). After 8 hours at 37°C, cells were pelleted, resuspended in fresh Cultrex, and cultured for 3

days. Transduced cells were selected using puromycin (2.5 µg/ml) for p7TFP-expressing enteroids

(Hu235D::7TFP) and sorted in liquid nitrogen.

7TCF/LEF Luciferase assays

The Hu235D::7TFP cell line was resuspended in 50% L-WRN conditioned media enumerated

(Countess FL, ThermoFisher Scientific), and 35,000 live cells added in a total volume of 100 µl per well of

a 96 well plate (TPP, flat bottom, Cat# 92096) pre-coated with 3% Cultrex. After 48 hours growth at 5%

CO_2, 37°C, media was replaced with 5% L-WRN + Y-27632 to encourage differentiation. After 2 days, cells

in respective wells (~95% confluency) were treated with toxins as indicated in primary media only. After

equilibration to room temperature, 5'-fluoroluciferin reagent (ONE-Glo EX, E8110 Promega) was added

to wells and plates shaken at 200 rpm for 10 minutes. 120 µl from each well was then transferred to

corresponding wells of an opaque white 96 well plate (Microlite 1+, ThermoFisher 7417) and

luminescence measured (Synergy H1, BioTek). All treatments were performed at least in triplicate,

repeated at least three times.

Subcellular fractionation and immunoblotting

To isolate nuclear and cytoplasmic fractions, subcellular fractionation of human ileal enteroids was

performed using the NE-PER™ Nuclear and Cytoplasmic Extraction Kit (Thermo Fisher Scientific, #78833),

following the manufacturer's protocol with minor adaptations. Enteroids (~1×10^6 cells per condition)

were collected after treatment (e.g., with heat-labile toxin or PBS control), washed twice in ice-cold PBS,

and pelleted by centrifugation at 500 × g for 5 min at 4°C. Cytoplasmic and nuclear extracts were

sequentially isolated using the provided CER I, CER II, and NER buffers supplemented with Halt™

protease and phosphatase inhibitor cocktail (Thermo Fisher Scientific, #78440). After incubation with

CER I and CER II to lyse the cytoplasmic membrane, lysates were centrifuged at 16,000 × g for 5 min to

separate the supernatant for cytoplasmic fraction. The pellet was then resuspended in NER buffer and

incubated with vigorous vortexing every 10 min over a 40-min period. The nuclear fraction was collected

by centrifugation at 16,000 × g for 10 min. Protein concentrations were measured using a BCA assay

(Thermo Fisher Scientific), and fraction purity was confirmed by immunoblotting for cytoplasmic

(GAPDH) and nuclear (Lamin b1) markers. For immunoblotting, 10–30 μg of protein per sample was

mixed with Laemmli sample buffer containing 100 mM DTT, boiled at 95°C for 10 min, and resolved on

4–12% SDS-PAGE gels. Proteins were transferred to nitrocellulose membranes (100 V, 1 h) and blocked in

5% non-fat milk in PBS-Tween (0.1%). Membranes were incubated overnight at 4°C with primary

antibodies at 1:1000 dilutions (Supplementary table 2). After washing three times with PBST, HRP-

conjugated secondary antibodies (1:5,000) were applied for 1 h at room temperature. Signals were

developed using Clarity ECL substrate (BioRAD) and visualized with a c600 imager (azure Biosystems).

Band intensities were quantified using ImageJ and normalized to total protein across biological

replicates.

Single nuclei isolation for RNA sequencing

Following treatment, enteroids were collected in ice-cold PBS supplemented with 0.5 mM EDTA, resuspended, and centrifuged at 200g for 5 minutes. For single-cell preparation, the cell pellets were resuspended in TrypLE (Invitrogen) and incubated at 37°C for 8 minutes to obtain a single-cell suspension. Cell debris and large clumps were removed by passing the single-cell suspension through a 40 µm filter, followed by centrifugation at 250g for 5 minutes. Prior to nuclei isolation, the cells were washed once with ice-cold PBS supplemented with 0.04% BSA and counted to obtain 1 million cells. Single nuclei were isolated following the protocol for nuclei isolation from fresh cells (Protocol CG000365 Rev C, 10X Genomics). Briefly, 100 µl of chilled Lysis Buffer was added to the cell pellet, mixed by pipetting 10 times, incubated for 5 minutes on ice, and washed with 1 ml of Wash Buffer. Nuclei were pelleted by centrifugation at 500g for 5 minutes at 4°C, washed two more times, then resuspended in Nuclei Buffer and immediately processed for sequencing.

cDNA library preparation and sequencing

For single-cell RNA sequencing together with single-cell ATAC sequencing, Multiome 3v3.1 GEX and ATAC libraries were prepared by using the Chromium Next GEM Single Cell Multiome ATAC + Gene Expression Reagent Kits following the recommended protocol (User Guide, CG000338, 10X Genomics). Samples were prepared on the 10x Genomics platform using the Chromium Next GEM Single Cell Multiome ATAC + Gene Expression Reagent Bundle (16 rxns, PN-1000283), Chromium Next GEM Chip J Single Cell Kit (48 rxns, PN-1000234), Single Index Kit N Set A (96 rxns, PN-1000212, for ATAC), and Dual Index Kit TT Set A (96 rxns, PN-1000215, for 3v3.1 GEX). The concentration of each library was determined by qPCR using the KAPA Library Quantification Kit (qPCR MasterMix cat#7959478001, and Library Quantification Kit, Cat# 7960085001, Roche Diagnostics System), following the manufacturer's protocol, to achieve appropriate cluster counts for the Illumina NovaSeq6000 instrument. GEX libraries were pooled and run on 0.0192 of a NovaSeq X Plus 25B flow cell using a 28x10x10x150 sequencing

recipe, as per the manufacturer's protocol, with a target coverage of 500M read-pairs per sample. ATAC

libraries were pooled and run on 0.0115 of a NovaSeq X Plus 25B flow cell with a 51x8x16x51 sequencing

recipe, following the manufacturer's protocol, aiming for a target coverage of 300M read-pairs per

sample.

Data processing

Single-cell RNA-seq and ATAC-seq data were processed using the 10x Genomics Cell Ranger ARC

pipeline (version 2.0.2). For alignment and quantification of gene expression data, raw sequencing reads

were processed using the default parameters provided by the 10x Genomics software suite. The

standard reference genome and annotations used were based on the refdata-cellranger-arc-GRCh38-

2020-A-2.0.0 build provided by 10x Genomics. Briefly, the Illumina sequencer's base call files (BCLs) for

each flow cell directory (ATAC or Gene Expression) were demultiplexed using the Illumina bcl2fastq

software to generate FASTQ files. These reads were then aligned to the GRCh38 human reference

genome, and unique molecular identifier (UMI) counts were obtained using the 10x Cell Ranger toolkit.

Barcode selection and filtering for single-cell droplets were performed according to the Cell Ranger

pipeline[11,27]

Data filtering and normalization

Single-cell Multiome libraries were used for initial quality control and cell filtering. For the present

study, downstream analyses were performed exclusively on the RNA-seq data, which is the focus of this

manuscript. Analyses of the ATAC-seq data will be reported separately. UMI count matrices for single

cells were analyzed in Seurat (v5.0)[45]. To ensure data quality, cells with fewer than 1,000 RNA counts or

more than 25,000 RNA counts were removed, along with cells where more than 50% of reads were

mapped to mitochondrial genes, thereby excluding potential low-quality cells, debris, and likely

multiplets. To further refine the dataset, doublets were identified and removed using DoubletFinder [85]

based on the expected doublet rate and model homotypic interactions, ensuring more accurate cell

classification. Following these steps, we applied the SCTransform method, a regularized negative

binomial regression approach that corrects for sequencing depth and other technical variations while

preserving biological heterogeneity in the dataset[86]. The output from SCTransform was used to identify

highly variable features, which were then used for downstream analysis.

Marker gene identification for cell clustering

To identify marker genes across all clusters, we used the FindAllMarkers function in Seurat. Genes

were designated as marker genes for a given cluster if they were significantly upregulated in that cluster

(Wilcoxon rank-sum test $p < 0.05$), expressed in at least 25% of cells within the cluster, and showed a

minimum log fold change of 0.25 (indicating approximately 19% higher expression) relative to other

clusters. Expression heatmaps of the top 10 signature genes for each cluster are shown in Figure 3b, and

all marker genes names and statistics for their enrichment are provided for every cluster in

Supplementary dataset 1.

Differential gene expression analysis

To identify gene expression variability and cell-specific responses to treatment, differential gene

expression within individual cell clusters was analyzed. Differentially expressed genes (DEGs) were

identified using Seurat's FindMarkers function to evaluate transcriptional changes across cell clusters

following LT treatment. For each cluster, DEGs were determined using a log2-fold-change threshold of

±0.25 to capture biologically relevant differences, and considering only genes expressed in at least 30%

of cells within each group. The Wilcoxon rank-sum test was used for statistical comparisons, allowing

identification of both up- and down-regulated genes and providing a comprehensive view of treatment-

induced transcriptional changes. The list of all DEGs is provided in **Supplementary dataset 2**.

Cell cycle analysis

Each cell was annotated for its cell cycle stage using the CellCycleScoring function and the updated G1/S and G2/M cell cycle genes from the `cc.genes.updated.2019` dataset in Seurat[46,87], which evaluates the expression of stage-specific marker genes. Specifically, this function assigns each cell a quantitative score based on the expression of 54 genes related to the S phase and 43 genes associated with the G2/M phase[27106]. By comparing these scores, cells were categorized as being in G1, S, or G2/M phases. Cells with low scores in both S and G2/M markers are inferred to be in the G1 phase, indicating a quiescent or resting state, while cells with high S or G2/M scores are identified as actively cycling. Composition of cell cycle phase across different clusters is provided in supplementary source data for tables (see Supplementary table 4). For visualization, the distribution of cell cycle phases per sample and cluster was plotted using ggplot2.

Labeling of proliferating cells

Actively proliferating cells were detected using the Click-iT EdU imaging kit (Molecular Probes, Life Technologies) according to the manufacturer's protocol. Briefly, following toxin treatments, organoids were incubated for three hours at 37°C in tissue culture media supplemented with 10 µM EdU. Following incubation, organoids were fixed with 4% paraformaldehyde as previously described and permeabilized with 0.5% Triton X-100 for 20 minutes at room temperature. The EdU-labeled cells were detected with the Click-iT® Plus reaction cocktail, followed by counterstaining with DAPI. For EdU analysis, twenty random fields were imaged at 20× magnification, and the numbers of EdU-positive and DAPI-positive nuclei were counted. The percentage of EdU-positive nuclei was calculated as a proportion of the total nuclei count per enteroid. Each experiment was conducted in duplicate.

Functional enrichment analysis

Ontology-independent algorithms in CompBio 2.9 (COmprehensive Multi-omics Platform for Biological InterpretatiOn) (https://gtac.wustl.edu/bioinformatics/) were used to analyze existing

transcriptome data in RNAseq datasets. Enrichment analysis was carried out using WebGestalt

(https://www.webgestalt.org/) against reference gene sets curated from Reactome pathways and

network databases. Over-representation analysis (ORA) was performed to identify biological pathways

and functional categories significantly enriched among differentially expressed genes. Genes with an

adjusted *p*-value < 0.05 were selected as input. Statistical significance was determined using a

hypergeometric test, with p-values adjusted for multiple testing by the Benjamini–Hochberg method

(FDR < 0.05). Redundancy among enriched categories was reduced using the weighted set cover method,

and results were visualized to highlight key biological processes associated with the gene sets.

Statistical analysis

All statistical analyses were conducted using GraphPad Prism (version 10.5.0) and RStudio (version

2023.12.1), unless otherwise specified. For comparisons between two groups, the non-parametric

Mann–Whitney U test was used. For comparisons among more than two groups, the Kruskal–Wallis test

followed by Dunn's multiple comparisons test was applied. To assess differences in cell proportions or

categorical distributions across experimental conditions, Chi-square (χ^2) tests were used. When

applicable, exact *P*-values were calculated and reported. In figures, statistical significance is indicated as:

P < 0.05 (*), P < 0.01 (**), P < 0.001 (***), and P < 0.0001 (****).

Data availability

Single-cell RNA-seq data have been deposited in the NCBI Sequence Read Archive (SRA) under the

BioProject accession PRJNA1330728 (https://www.ncbi.nlm.nih.gov/bioproject/). Primary source data

and original images are available on Figshare, as detailed in Supplementary table 4. All other data

supporting the findings of these studies are included in the manuscript as supplementary dataset and

are also available from the corresponding author upon reasonable request.

Figures

Figure 1

Figure 1. LT activates WNT/β-catenin signaling in human intestinal epithelial cells.

a) Heatmap showing differential expression (z-score) of WNT/β-catenin signaling components and target

genes in control (∅) and LT-treated enteroids based on bulk RNA-seq. Closed green symbols signify genes

known to be modulated by canonical WNT-β-catenin pathway as profiled

(https://web.stanford.edu/group/nusselab/cgi-bin/wnt/target_genes). Open green symbols indicate

genes possessing an upstream 3'UTR CTTTGT motif for the LEF1: lymphoid enhancer-binding factor

http://www2.stat.duke.edu/~sayan/genesets/Jan2006/cards/C3/CTTTGT_V_LEF1_Q2.html. **b**) RT-qPCR

validation of selected genes showing significantly increased expression following LT treatment. Gene

expression was measured at 24 and 72 hours post-treatment. Data are presented as expression levels

normalized to GAPDH and shown relative to the untreated control. Inset: CEACAM6 expression is

included as a positive control for LT-mediated induction. Statistical comparisons were made using

Kruskal-Wallis; ns = not significant, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. **c**)

Representative confocal immunofluorescence images of β-catenin (yellow) and nuclei (blue) in LT-treated

and control enteroids showing increased nuclear β-catenin accumulation in LT-treated samples. Scale

bar, 25 μm. **d**) Quantification of nuclear β-catenin intensity per cell from images shown in panel (c). Each

dot represents an individual nucleus. ****p<0.0001 Mann-Whitney (two-tailed). **e**) Luciferase reporter

activity (in Relative light units, RLU) following treatment of Hu235D::7TFP cells with heat-labile toxin (LT)

or mutant, catalytically inactive toxin (mLT, E112K). The data shown are combined from 3-7 independent

experiments. Each dot represents signal per well and each Bar represents geometric mean. Statistical

analysis by one-way ANOVA (Kruskal-Wallis), ****p<0.0001.

Figure 2

Figure 2. Heat-labile toxin stabilizes β-catenin signaling independent of WNT.

a) Diagram illustrating canonical WNT–β-catenin signaling (left and middle panels). Panel at right

depicts anticipated impacts of PKA-mediated, WNT-independent signaling. **b)** Representative

immunoblots of cytoplasmic and **c)** nuclear fractions from LT-treated enteroids at indicated time points

(4, 24, 72 hours) or untreated controls (Ø). Blots were probed for total β-catenin, phospho-β-catenin

(Ser$_{675}$), total GSK3β, and phospho-GSK3β (Ser$_9$). GAPDH and lamin B1 served as loading controls for

cytoplasmic and nuclear fractions, respectively. **d-e)** Graphs represent densitometry measurements of

601   immunoblots of cytoplasmic (**d**) and nuclear (**e**) fractions, normalized to respective loading controls,

602   summarizing 4 independent experiments. Bars represent geometric means ± SD.

Figure 3

Figure 3. LT treatment reshapes intestinal epithelial cell composition.

a) UMAP projection of single-cell transcriptomes from control and LT-treated ileal enteroids, colored

by cluster identity. Unsupervised clustering identified 14 transcriptionally distinct epithelial populations,

which were annotated based on lineage marker gene expression. These include enterocyte progenitors,

intermediate cells, secretory lineages (goblet, enteroendocrine, and Paneth-like), stem cells, and

multiple stages of differentiated enterocytes. **b**) Heatmap showing the relative expression levels of the

top 10 marker genes per cluster, showing distinct transcriptional signatures corresponding to absorptive,

secretory, intermediate, and proliferative compartments. Expression values represent SCTransform-

normalized Pearson residuals. **c**) UMAP feature plots showing expression of known cell type specific

markers. Enterocyte markers: ANPEP, SI, ALPI; goblet cell markers: MUC2, SPINK4, TFF3; enteroendocrine

cell markers: CHGA, CHGB, THP1; transit-amplifying cell markers: MKI67, TOP2A, PCNA. Red dotted

circles highlight specific cell clusters with high expression values. **d**) UMAP showing the distribution of

cells from control (orange) and LT-treated (blue) samples, revealing a shift in cell type representation.

Inset: Stacked bar plot showing the proportion of cells from each sample (control vs LT) within each

cluster. **e**) Differential abundance analysis for epithelial cell clusters in LT-treated versus control

enteroids. Each point represents an epithelial cell cluster. The x-axis shows the $\log_2$ ratio of observed to

expected cell counts in LT-treated enteroids, and the y-axis shows the $-\log_{10}$(FDR-adjusted p-value).

Vertical and horizontal dashed lines indicate the direction of change (assigned based on whether the

observed LT count exceeded or fell below the expected value) and statistical significance thresholds,

respectively. Clusters to the right of the vertical line are enriched in LT-treated samples, whereas clusters

to the left are reduced. Significantly altered clusters are labeled. Statistical analysis was performed per

cluster using a Chi-square test with FDR correction (significant if FDR <0.05). BEST4+EC, BEST4 positive

enterocyte; EC, enterocyte; EC prog.- enterocyte progenitor; EEC, enteroendocrine cell; GC, goblet cell;

ISC, intestinal stem cell; int., intermediate; sec. prog., secretory progenitor; TA, transit amplifying.

Figure 4

631
632
Figure 4: LT alters epithelial cell cycle dynamics.

a) UMAP projections of single-cell transcriptomes from control and LT-treated human ileal enteroids,

colored by assigned cell cycle phase based on canonical marker gene expression (G1, S, G2/M). **b**) XY

scatterplots of individual cell S-phase and G2/M-phase scores derived from gene expression signatures.

c) UMAP feature plots highlighting expression of representative phase-specific marker genes across

conditions. G1-associated genes (RB1, CDKN1A), S-phase associated genes (PCNA, MCM3) and G2/M-

phase associated genes (CCNB2, CDK1). Color scales represent normalized expression.

Figure 5

Figure 5: LT affects cell cycle progression across intestinal epithelial lineages and promotes

epithelial proliferation.

a) Donut charts represent the global distribution of cells across cell cycle phases in control and LT-

treated enteroids. Bar plots below show the proportion of cells in G1 (green), S (blue), and G2/M (red)

phases across transcriptionally defined cell clusters (0–13) in control and LT-treated enteroids. **b**) UMAP

feature plots showing expression of proliferation markers MKI67 (top) and TOP2A (bottom) in control

(left) and LT-treated (right) human ileal enteroids. The color scale indicates normalized expression levels.

c) Representative confocal images of MKI67 (white) and nuclei (DAPI, blue) in control (left) and LT-treated

(right) enteroids. Scale bar, 10 μm. **d**) Quantification of MKI67$^+$ cells per field of view. Each point

represents the number of MKI67$^+$ nuclei in an individual field; the horizontal line denotes the mean.

Data are from two independent experiments (≥50 fields of view total). Statistical significance was

determined by a two-tailed, nonparametric Mann–Whitney test (****p < 0.0001). **e**) Representative

confocal images of control (left) and LT-treated (right) enteroids labeled with EdU (green) for DNA

synthesis, DAPI (blue) for nuclei, and a membrane stain (red). White asterisks mark the luminal space

(equivalent to intestinal lumen) in enteroids. Off note, LT-treated enteroids showed a marked increase in

luminal area, consistent with LT-induced fluid secretion. Scale bar, 25 μm. **f**) Quantification of EdU$^+$

nuclei per enteroid (n = 25 enteroids per treatment). Each point represents the number of EdU$^+$ nuclei

per individual enteroid; the horizontal line denotes the mean. Statistical significance was determined by

a two-tailed, nonparametric Mann–Whitney test (**p < 0.01).

Figure 6

Figure 6. LT affects enterocyte maturation and disrupts nutrient transporter expression.

a) Western blot analysis of ALPI (intestinal alkaline phosphatase enzyme) expression in control and LT-treated enteroids. Representative blot and corresponding densitometric histograms generated using ImageJ (left). Quantification of normalized ALPI expression (right). Each data point represents an individual replicate (n = 7 per group). Horizontal lines on the plot indicate mean. Statistical significance was determined using a non-parametric Mann-Whitney test (*p < 0.05, two-tailed). **b**) Western blot analysis of Villin (structural protein) expression in control and LT-treated enteroids. Representative blot and corresponding densitometric histograms generated using ImageJ (left). Quantification of normalized Villin expression (right). Each data point represents an individual replicate (n = 6 per group). Horizontal lines on the plot indicate mean. Statistical significance was determined using a non-parametric Mann-Whitney test (*p < 0.05, two-tailed). **c**) Analysis of nutrient transporter genes across enterocyte subclusters (EC-1 to EC-5). Data were derived from scRNA-seq of control and LT-treated enteroids. Violin plots showing distributions of mean expression of major nutrient transporter gene families (nucleotide, metal, sugar, organic, inorganic, amino acid, water, vitamin, bile, lipid). Dot plot illustrating expression patterns of specific genes involved in nutrient absorption and transport. The color scale indicates the average expression, and the dot size indicates proportion of cells expressing individual transporter genes across five enterocyte subclusters (EC-1 to EC-5). **d**) UMAP feature plots showing SLC19A3 thiamine transporter and SP1 regulator gene expression across cell clusters. **e**) Volcano plots showing differential expression of nutrient transporter genes within enterocyte subclusters EC-1 through EC-5. The x-axis indicates $\log_2$ fold-change (LT vs. control), and the y-axis indicates $-\log_{10}$ adjusted p-value. Genes significantly upregulated in LT-treated enteroids are shown in red, significantly downregulated genes in blue, and non-significant changes in gray. Selected transporters with significant changes are labeled. Statistical testing was performed using a Wilcoxon rank-sum test with Benjamini–Hochberg correction for multiple comparisons; adjusted p < 0.05 was considered significant. EC, enterocyte.

Tables

Table-1. Transcriptionally-defined epithelial cell cluster compositions in ileal enteroids.

Cluster	Identity		Treatment conditions		Chi2 Statistic	*P* Value	Change with LT
			Control (%)	LT (%)			
0	EC prog.	Enterocyte progenitor	270 (13.8)	461 (20.9)	36.079	1.90E-09	Expanded
1	int.-1	Intermediate-1	229 (11.7)	217 (9.8)	3.467	6.26E-02	Not Significant
2	int. -2	Intermediate-2	242 (12.3)	212 (9.6)	7.658	5.65E-03	Depleted
3	GC	Goblet cell	234 (11.9)	246 (11.1)	0.538	4.63E-01	Not Significant
4	EC-1	Enterocyte-1	264 (13.4)	163 (7.4)	40.897	1.60E-10	Depleted
5	EC-2	Enterocyte-2	239 (12.2)	116 (5.3)	62.978	2.09E-15	Depleted
6	EC-3	Enterocyte-3	147 (7.5)	179 (8.1)	0.475	4.91E-01	Not Significant
7	TA	Transit amplifying	17 (0.9)	243 (11.0)	181.147	2.72E-41	Expanded
8	Sec. prog.	Secretory progenitor	33 (1.7)	133 (6.0)	50.188	1.40E-12	Expanded
9	EC-4	Enterocyte-4	80 (4.1)	78 (3.5)	0.693	4.05E-01	Not Significant
10	ISC	Intestinal stem cell	84 (4.3)	62 (2.8)	6.216	1.27E-02	Depleted
11	EC-5	Enterocyte-5	76 (3.9)	67 (3.0)	1.947	1.63E-01	Not Significant
12	EEC	Enteroendocrine cell	17 (0.9)	28 (1.3)	1.223	2.69E-01	Not Significant
13	BEST4+EC	BEST4+ enterocyte	31 (1.6)	2 (0.1)	27.458	1.61E-07	Depleted

Table-2. Total number of differentially expressed genes per cluster following LT treatment

Description		Cell counts		Number of differentially expressed genes (DEGs)*	
Cluster #	Identity	control	LT	Higher in LT	Lower in LT
0	Enterocyte progenitor	270	461	634	901
1	Intermediate-1	229	217	455	545
2	Intermediate-2	242	212	447	323
3	Goblet cell	234	246	236	509
4	Enterocyte-1	264	163	256	274
5	Enterocyte-2	239	116	310	310
6	Enterocyte-3	147	179	354	420
7	Transit amplifying	17	243	N/A	N/A
8	Secretory progenitor	33	133	85	305
9	Enterocyte-4	80	78	N/A	N/A
10	Intestinal stem cell	84	62	157	267
11	Enterocyte-5	76	67	148	132
12	Enteroendocrine cell	17	28	N/A	N/A
13	BEST4+ enterocyte	31	2	N/A	N/A

* Differentially expressed genes across epithelial clusters in LT-treated vs. control enteroids. Genes with minimal log2FC of +/-0.25, and expressed in at least 10% of cells with FDR adjusted P value of <0.05 were considered significant.

Table-3. Cell cycle phase distribution in control and LT-treated enteroids

Clusters	Phase	Cell count (%)		Chi²_statistic	P value
		Control	LT		
all combined	G1	1303 (66.4)	1031 (46.8)	162.2	< 1e-16
	S	602 (33.7)	546 (24.7)	18	2.21E-05
	G2/M	58 (2.9)	630 (28.5)	492.01	< 1e-16
#0	G1	197 (73.0)	221 (47.9)	61.06	<0.00001
	G2M	10 (3.7)	107 (23.2)		
	S	63 (23.3)	133 (28.9)		
#1	G1	144 (62.9)	104 (47.9)	43.74	<0.00001
	G2M	6 (2.6)	51 (23.5)		
	S	79 (34.5)	62 (28.6)		
#2	G1	137 (56.6)	139 (65.6)	37.596	<0.00001
	G2M	7 (2.9)	32 (15.1)		
	S	98 (40.5)	41 (19.3)		
#3	G1	187 (79.9)	147 (59.8)	53.1702	<0.00001
	G2M	3 (1.3)	57 (23.2)		
	S	44 (18.8)	42 (17.1)		
#4	G1	176 (66.7)	82 (50.3)	23.0313	<0.00001
	G2M	1 (0.4)	12 (7.4)		
	S	87 (33.0)	69 (42.3)		
#5	G1	165 (69.0)	67 (57.8)	21.4174	0.000022
	G2M	2 (0.8)	13 (11.2)		
	S	72 (30.1)	36 (31.0)		
#6	G1	95 (64.6)	110 (61.5)	10.847	0.004412
	G2M	7 (4.8)	28 (15.6)		
	S	45 (30.6)	41 (22.9)		
#7	G1	7 (41.2)	1 (0.4)	90.4198	<0.00001
	G2M	7 (41.2)	214 (88.1)		
	S	3 (17.6)	28 (11.5)		
#8	G1	18 (54.5)	37 (27.8)	11.3691	0.0034
	G2M	5 (15.2)	58 (43.6)		
	S	10 (30.3)	38 (28.6)		
#9	G1	48 (60.0)	41 (52.6)	5.5985	0.061
	G2M	4 (5.0)	13 (16.7)		
	S	28 (35.0)	24 (30.8)		
#10	G1	68 (81.0)	22 (35.5)	38.4988	<0.00001
	G2M	3 (3.6)	25 (40.3)		
	S	13 (15.5)	15 (24.2)		
#11	G1	44 (57.9)	39 (58.2)	19.4254	0.000061
	G2M	1 (1.3)	15 (22.4)		
	S	31 (40.8)	13 (19.4)		
#12	G1	15 (88.2)	20 (71.4)	1.4768	0.47787
	S	NF*	5 (17.9)		
	G1	2 (11.8)	3 (10.7)		
#13	G1	2 (6.5)	1 (50.0)	5.4403	0.065863
	G2M	2 (6.5)	NF*		
	S	27 (87.1)	1 (50.0)		

 *NF-not found

Supplementary data

Supplementary figures

Supplementary figure 1

Supplementary figure 1. Phophorylated β-catenin (Ser$_{675}$) accumulates in target intestinal cells

following treatment with heat-labile toxin.

a. Top panels show phospho-β-catenin accumulation in representative field of LT-treated Hu235D

cells compared to untreated cells (**b**). **c.** Violin plots at right summarize p-β-catenin Ser$_{675}$ signal per

cell obtained from n=7088 cells in control, and n=7176 cells in LT-treated group. Data was

accumulated over 20 randomly selected fields. Heavy dashed lines in each group represent median

values, thin lines are quartiles. ****p<0.0001 (Mann-Whitney).

Supplementary figure 2

Supplementary figure 2. Over-representation analysis (ORA) of cluster 7 (TA) and cluster 13 marker

genes (BEST4+EC). Dot plots displaying significantly enriched (FDR < 0.05) pathways (a) and transcription

factors (b) along the Y-axis, among cluster marker genes for clusters 7 and 13. Dot size indicates the

number of marker genes contributing to each pathway, and color reflects the adjusted statistical

significance (−log10 FDR), with yellow indicating stronger enrichment and blue indicating lower

significance.

 Supplementary figure 3

Supplementary figure 3. Over-representation analysis (ORA) of significantly up- and down-

regulated genes between LT and control samples.

The dot plot displays significantly enriched biological pathways (Y-axis) across 14 epithelial cell

clusters (X-axis) among differentially expressed genes in LT-treated vs. control intestinal organoids (FDR

<0.05). Circles represent upregulated pathways, while triangles represent downregulated pathways. Dot

size indicates the number of DEGs contributing to each pathway, and color reflects the adjusted

statistical significance (−log10 FDR), with yellow indicating stronger enrichment and blue indicating lower

significance.

Supplementary figure 4. CompBio analysis of differentially expressed genes (DEGs) in LT-treated

enteroids.

a) Bar plot showing themes significantly enriched among downregulated genes (FDR-adjusted p < 0.05).

b) Bar plot showing themes significantly enriched among upregulated genes (FDR-adjusted p < 0.05).

Pathway enrichment was conducted using CompBio v2.9 with default parameters. NEScore represents

the normalized enrichment score derived from CompBio theme association analysis. Themes were

prioritized based on total score. See Supplementary dataset for full theme descriptions and gene lists.

Supplementary figure 5

Supplementary figure 5: Expression of cell-cycle phase and stem/progenitor genes.

a) Ridge plots showing expression of cell cycle phase marker genes. Normalized expression of S-phase

genes (left) and G2/M-phase genes (right) in control and LT-treated enteroids. b) Feature plots displaying

expression of CD44, CD24, and OLFM4 markers of stem or undifferentiated states across cell clusters in

control (left) and LT-treated (right) enteroids.

Supplementary figure 6

Supplementary figure 6. LT suppresses expression of multiple nutrient transporter genes in human

ileal enteroids.

UMAP feature plots showing the expression of select solute carrier (SLC) family genes involved in

nutrient absorption across control (left panels) and LT-treated (right panels) human ileal enteroids. Each

pair of panels displays normalized expression levels of a specific SLC gene projected onto the epithelial

cell landscape. Gene function annotations are summarized in the accompanying table.

Supplementary tables

Supplementary table 1: Primer sequences used in Quantitative RT-PCR experiments

Gene	PrimerBank ID*	direction	amplicon size	sequence	Source*
CTNNB1	148233337c2	F	150	CATCTACACAGTTTGATGCTGCT	PrimerBank
		R		GCAGTTTTGTCAGTTCAGGGA	PrimerBank
SOX9	182765453c1	F	85	AGCGAACGCACATCAAGAC	PrimerBank
		R		CTGTAGGCGATCTGTTGGGG	PrimerBank
OLFM4	325053703c2	F	135	ACTGTCCGAATTGACATCATGG	PrimerBank
		R		TTCTGAGCTTCCACCAAAACTC	PrimerBank
MYC	239582723c1	F	119	GGCTCCTGGCAAAAGGTCA	PrimerBank
		R		CTGCGTAGTTGTGCTGATGT	PrimerBank
CCND1	77628152c1	F	135	GCTGCGAAGTGGAAACCATC	PrimerBank
		R		CCTCCTTCTGCACACATTTGAA	PrimerBank
CD24	73623396c1	F	166	CTCCTACCCACGCAGATTTATTC	PrimerBank
		R		AGAGTGAGACCACGAAGAGAC	PrimerBank
PROM1	224994190c1	F	99	AGTCGGAAACTGGCAGATAGC	PrimerBank
		R		GGTAGTGTTGTACTGGGCCAAT	PrimerBank
CEACAM6	NA	F	113	CACCGTCGGCATCACGA	[29]
	NA	R		GAAGAATTCAGGGTCTGGTCCA	[29]
*https://pga.mgh.harvard.edu/primerbank/index.html					

 Supplementary table 2: Antibody descriptions and sources for immunoblot and fluorescence

 microscopy experiments

Protein, antigen	clone[1]	host	isotype	conjugation	source	RRID*
GAPDH, C terminal peptide	m |14C10	Rabbit	IgG		Cell Signaling 2118S	AB_561053
GSK-3a/b (synthetic peptide surrounding Gln269 of human GSK-3a protein)	D75D3	Rabbit	IgG		Cell Signaling 5676	AB_10547140
GSK-3b Ser$_9$ phospho-peptide	polyclonal	Rabbit	IgG		Cell Signaling 9336S	AB_331405
GSK-3a/b (Ser21/9) synthetic phophopeptides	Polyclonal	Rabbit			Cell Signaling 9331	AB_329830
b-catenin (synthetic peptide circa Pro174 of human protein)	monoclonal (D10A8)	Rabbit	IgG		Cell Signaling 8480	AB_11127855
b-catenin Ser$_{675}$ C-terminal phospho-peptide	polyclonal	Rabbit	IgG		Invitrogen PA5-77933	AB_2736355
MKI67	polyclonal	Rabbit	IgG		Novus NB500-170	AB_10001977
Villin (BDID2C3)	monoclonal	Mouse	IgG1		Santa Cruz sc-66022	AB_2216247
ALPI	polyclonal	Rabbit	IgG		Invitrogen PA5-22210	AB_11153471
GAPDH (14C10)	monoclonal	Rabbit			Cell Signaling 2118S	AB_561053
Lamin B1	monoclonal (EPR22165-121)	Rabbit	IgG		Abcam ab229025	AB_3083735
Anti-rabbit IgG	polyclonal	Goat	IgG (H+L)	Alexa Fluor 488	ThermoFisher A-11070	AB_143165
Anti-rabbit IgG	polyclonal	Goat	IgG	HRP	Cell Signaling 7074S	AB_2099233
Anti-mouse IgG	polyclonal	Horse	IgG	HRP	Cell Signaling 7076S	AB_330924

 * https://rrid.site/

762 **Supplementary table 3: Plasmids used in the construction of TCF/LEF reporter enteroid lines**

plasmid	description	source	reference
psPAX2	2nd generation lentiviral packaging plasmid	addgene[1]	
pMD2.G	VSV-G envelope expressing plasmid for lentiviral packaging	addgene[2]	
p7TFP	7X copies of the TCF/LEF transcriptional response element GATCAAAGG linked to firefly luciferase ; puromycin R	addgene[3]	30

[1]psPAX2 was a gift from Didier Trono (https://www.addgene.org/12260/; RRID:Addgene_12260)
[2]pMD2.G was a gift from Didier Trono (https://www.addgene.org/12259/ ; RRID:Addgene_12259)
[3]p7TFP was a gift from Roel Nusse (https://www.addgene.org/24308/; RRID:Addgene_24308)
[4]p7TGC was a gift from Roel Nusse (https://www.addgene.org/24304/); RRID:Addgene_24304)

763
764
765

766 **Supplementary table 4: links to the original images and datasets**

Type	Description	DOI
Figure 2b-c.	Original blots of β-catenin, phospho-β-catenin- Ser$_{675}$, GSK3β, phospho-GSK3β-Ser$_9$, Lamin-b1, and GAPDH	10.6084/m9.figshare.30100273
Figure 6a.	Original blot of intestinal alkaline phosphatase (ALPI)	10.6084/m9.figshare.30104512
Figure 6b.	Original blot of Villin	10.6084/m9.figshare.30104635
Data	Data associated with graphs	10.6084/m9.figshare.30198553
Data	Data associated with tables	10.6084/m9.figshare.30201280

Acknowledgements

JMF was supported by funding from the Department of Veterans Affairs (VA) (5I01BX001469-05), and

the National Institute of Allergy and Infectious Diseases (NIAID) of the National Institutes of Health (NIH)

R01 AI170949, R01 AI089894, and support by the NIH Washington University DDRCC Grant NIDDK P30

DK052574**.** AS was supported by National Institute of Allergy and Infectious Diseases of the National

Institutes of Health under Award Number T32AI007172 and JIT1052 (supported by the Washington

University Institute of Clinical and Translational Sciences grant UL1TR002345 from the National Center

for Advancing Translational Sciences (NCATS) of the NIH. We thank the Genome Technology Access

Center at the McDonnell Genome Institute at Washington University School of Medicine for help with

genomic analysis. The Center is partially supported by NCI Cancer Center Support Grant #P30 CA91842

to the Siteman Cancer Center from the National Center for Research Resources (NCRR), a component of

the NIH, and NIH Roadmap for Medical Research. This publication is solely the responsibility of the

authors and does not necessarily represent the official view of NCRR, NIH, or VA.

References

1   Fleckenstein, J. M. & Sheikh, A. Emerging Themes in the Molecular Pathogenesis of
Enterotoxigenic Escherichia coli. *J Infect Dis* (2021). https://doi.org/10.1093/infdis/jiab359
2   Khalil, I. A. *et al.* Morbidity and mortality due to shigella and enterotoxigenic Escherichia coli
diarrhoea: the Global Burden of Disease Study 1990-2016. *Lancet Infect Dis* **18**, 1229-1240 (2018).
https://doi.org/10.1016/S1473-3099(18)30475-4
3   Qadri, F. *et al.* Disease burden due to enterotoxigenic Escherichia coli in the first 2 years of life in
an urban community in Bangladesh. *Infection and immunity* **75**, 3961-3968 (2007).
https://doi.org/10.1128/IAI.00459-07
4   Kotloff, K. L. *et al.* Burden and aetiology of diarrhoeal disease in infants and young children in
developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control
study. *Lancet* **382**, 209-222 (2013). https://doi.org/10.1016/S0140-6736(13)60844-2
5   Levine, M. M. *et al.* Diarrhoeal disease and subsequent risk of death in infants and children
residing in low-income and middle-income countries: analysis of the GEMS case-control study and
12-month GEMS-1A follow-on study. *Lancet Glob Health* **8**, e204-e214 (2020).
https://doi.org/10.1016/S2214-109X(19)30541-8
6   Nasrin, D. *et al.* Pathogens associated with linear growth faltering in children with diarrhea and
impact of antibiotic treatment: The Global Enteric Multicenter Study. *J Infect Dis* (2021).
https://doi.org/10.1093/infdis/jiab434

7   Kotloff, K. L. *et al.* The incidence, aetiology, and adverse clinical consequences of less severe
diarrhoeal episodes among infants and children residing in low-income and middle-income
countries: a 12-month case-control study as a follow-on to the Global Enteric Multicenter Study
(GEMS). *Lancet Glob Health* **7**, e568-e584 (2019). https://doi.org/10.1016/S2214-109X(19)30076-
2

8   Amadi, B. *et al.* Adaptation of the small intestine to microbial enteropathogens in Zambian
children with stunting. *Nat Microbiol* **6**, 445-454 (2021). https://doi.org/10.1038/s41564-020-
00849-w

9   Kelly, P. *et al.* Histopathology underlying environmental enteric dysfunction in a cohort study of
undernourished children in Bangladesh, Pakistan, and Zambia compared with United States
children. *Am J Clin Nutr* **120 Suppl 1**, S15-S30 (2024).
https://doi.org/10.1016/j.ajcnut.2024.02.028

10   Sheikh, A. *et al.* Enterotoxigenic Escherichia coli heat-labile toxin drives enteropathic changes in
small intestinal epithelia. *Nature communications* **13**, 6886 (2022).
https://doi.org/https://doi.org/10.1038/s41467-022-34687-7

11   Haber, A. L. *et al.* A single-cell survey of the small intestinal epithelium. *Nature* **551**, 333-339
(2017). https://doi.org/10.1038/nature24489

12   Mariadason, J. M. *et al.* Gene expression profiling of intestinal epithelial cell maturation along the
crypt-villus axis. *Gastroenterology* **128**, 1081-1088 (2005).
https://doi.org/10.1053/j.gastro.2005.01.054

13   Moor, A. E. *et al.* Spatial Reconstruction of Single Enterocytes Uncovers Broad Zonation along the
Intestinal Villus Axis. *Cell* **175**, 1156-1167 e1115 (2018).
https://doi.org/10.1016/j.cell.2018.08.063

14   Yu, J. *et al.* Environmental Enteric Dysfunction Includes a Broad Spectrum of Inflammatory
Responses and Epithelial Repair Processes. *Cell Mol Gastroenterol Hepatol* **2**, 158-174 e151 (2016).
https://doi.org/10.1016/j.jcmgh.2015.12.002

15   George, C. M. *et al.* Enteric Infections in Young Children are Associated with Environmental
Enteropathy and Impaired Growth. *Trop Med Int Health* **23**, 26-33 (2018).
https://doi.org/10.1111/tmi.13002

16   Guerrant, R. L., Oria, R. B., Moore, S. R., Oria, M. O. & Lima, A. A. Malnutrition as an enteric
infectious disease with long-term effects on child development. *Nutr Rev* **66**, 487-505 (2008).
https://doi.org/10.1111/j.1753-4887.2008.00082.x

17   Rosenberg, I. H., Solomons, N. W. & Schneider, R. E. Malabsorption associated with diarrhea and
intestinal infections. *Am J Clin Nutr* **30**, 1248-1253 (1977). https://doi.org/10.1093/ajcn/30.8.1248

18   van der Flier, L. G. & Clevers, H. Stem cells, self-renewal, and differentiation in the intestinal
epithelium. *Annu Rev Physiol* **71**, 241-260 (2009).
https://doi.org/10.1146/annurev.physiol.010908.163145

19   Fevr, T., Robine, S., Louvard, D. & Huelsken, J. Wnt/beta-catenin is essential for intestinal
homeostasis and maintenance of intestinal stem cells. *Molecular and cellular biology* **27**, 7551-
7559 (2007). https://doi.org/10.1128/MCB.01034-07

20   Van der Flier, L. G. *et al.* The Intestinal Wnt/TCF Signature. *Gastroenterology* **132**, 628-632 (2007).
https://doi.org/10.1053/j.gastro.2006.08.039

21   Vanuytsel, T., Senger, S., Fasano, A. & Shea-Donohue, T. Major signaling pathways in intestinal stem
cells. *Biochim Biophys Acta* **1830**, 2410-2426 (2013).
https://doi.org/10.1016/j.bbagen.2012.08.006

22   Farin, H. F. *et al.* Visualization of a short-range Wnt gradient in the intestinal stem-cell niche.
Nature **530**, 340-343 (2016). https://doi.org/10.1038/nature16937

23 Winston, J. T. *et al.* The SCFbeta-TRCP-ubiquitin ligase complex associates specifically with
phosphorylated destruction motifs in IkappaBalpha and beta-catenin and stimulates IkappaBalpha
ubiquitination in vitro. *Genes Dev* **13**, 270-283 (1999). https://doi.org/10.1101/gad.13.3.270

24 Taipale, J. & Beachy, P. A. The Hedgehog and Wnt signalling pathways in cancer. *Nature* **411**, 349-
354 (2001). https://doi.org/10.1038/35077219

25 Papkoff, J., Rubinfeld, B., Schryver, B. & Polakis, P. Wnt-1 regulates free pools of catenins and
stabilizes APC-catenin complexes. *Molecular and cellular biology* **16**, 2128-2134 (1996).
https://doi.org/10.1128/MCB.16.5.2128

26 Korinek, V. *et al.* Constitutive transcriptional activation by a beta-catenin-Tcf complex in APC-/-
colon carcinoma. *Science* **275**, 1784-1787 (1997). https://doi.org/10.1126/science.275.5307.1784

27 Wang, Y. *et al.* Single-cell transcriptome analysis reveals differential nutrient absorption functions
in human intestine. *J Exp Med* **217** (2020). https://doi.org/10.1084/jem.20191130

28 Beumer, J. & Clevers, H. Cell fate specification and differentiation in the adult mammalian
intestine. *Nat Rev Mol Cell Biol* **22**, 39-53 (2021). https://doi.org/10.1038/s41580-020-0278-0

29 Sheikh, A. *et al.* CEACAMs serve as toxin-stimulated receptors for enterotoxigenic Escherichia coli.
Proc Natl Acad Sci U S A **117**, 29055-29062 (2020). https://doi.org/10.1073/pnas.2012480117

30 Fuerer, C. & Nusse, R. Lentiviral vectors to probe and manipulate the Wnt signaling pathway. *PLoS*
One **5**, e9370 (2010). https://doi.org/10.1371/journal.pone.0009370

31 Cadigan, K. M. & Waterman, M. L. TCF/LEFs and Wnt signaling in the nucleus. *Cold Spring Harb*
Perspect Biol **4** (2012). https://doi.org/10.1101/cshperspect.a007906

32 Hino, S., Tanji, C., Nakayama, K. I. & Kikuchi, A. Phosphorylation of beta-catenin by cyclic AMP-
dependent protein kinase stabilizes beta-catenin through inhibition of its ubiquitination.
Molecular and cellular biology **25**, 9063-9072 (2005). https://doi.org/10.1128/MCB.25.20.9063-
9072.2005

33 Li, M. *et al.* Cyclic AMP promotes neuronal survival by phosphorylation of glycogen synthase kinase
3beta. *Molecular and cellular biology* **20**, 9356-9363 (2000).
https://doi.org/10.1128/MCB.20.24.9356-9363.2000

34 Fang, X. *et al.* Phosphorylation and inactivation of glycogen synthase kinase 3 by protein kinase A.
Proc Natl Acad Sci U S A **97**, 11960-11965 (2000). https://doi.org/10.1073/pnas.220413597

35 Tanji, C. *et al.* A-kinase anchoring protein AKAP220 binds to glycogen synthase kinase-3beta (GSK-
3beta) and mediates protein kinase A-dependent inhibition of GSK-3beta. *The Journal of*
biological chemistry **277**, 36955-36961 (2002). https://doi.org/10.1074/jbc.M206210200

36 Suzuki, A. *et al.* PTH/cAMP/PKA signaling facilitates canonical Wnt signaling via inactivation of
glycogen synthase kinase-3beta in osteoblastic Saos-2 cells. *J Cell Biochem* **104**, 304-317 (2008).
https://doi.org/10.1002/jcb.21626

37 Wang, S. *et al.* Quantitative Phosphoproteomic Study Reveals that Protein Kinase A Regulates
Neural Stem Cell Differentiation Through Phosphorylation of Catenin Beta-1 and Glycogen
Synthase Kinase 3beta. *Stem Cells* **34**, 2090-2101 (2016). https://doi.org/10.1002/stem.2387

38 Kawasaki, Y. *et al.* Phosphorylation of GSK-3beta by cGMP-dependent protein kinase II promotes
hypertrophic differentiation of murine chondrocytes. *The Journal of clinical investigation* **118**,
2506-2515 (2008). https://doi.org/10.1172/JCI35243

39 Zhao, X., Zhuang, S., Chen, Y., Boss, G. R. & Pilz, R. B. Cyclic GMP-dependent protein kinase
regulates CCAAT enhancer-binding protein beta functions through inhibition of glycogen synthase
kinase-3. *The Journal of biological chemistry* **280**, 32683-32692 (2005).
https://doi.org/10.1074/jbc.M505486200

40 Taurin, S., Sandbo, N., Yau, D. M., Sethakorn, N. & Dulin, N. O. Phosphorylation of beta-catenin by
PKA promotes ATP-induced proliferation of vascular smooth muscle cells. *Am J Physiol Cell Physiol*
294, C1169-1174 (2008). https://doi.org/10.1152/ajpcell.00096.2008

41   Taurin, S., Sandbo, N., Qin, Y., Browning, D. & Dulin, N. O. Phosphorylation of beta-catenin by cyclic
AMP-dependent protein kinase. *The Journal of biological chemistry* **281**, 9971-9976 (2006).
https://doi.org/10.1074/jbc.M508778200

42   Zambon, A. C. *et al.* Gene expression patterns define key transcriptional events in cell-cycle
regulation by cAMP and protein kinase A. *Proc Natl Acad Sci U S A* **102**, 8561-8566 (2005).
https://doi.org/10.1073/pnas.0503363102

43   Hua, X. *et al.* A Ctnnb1 enhancer transcriptionally regulates Wnt signaling dosage to balance
homeostasis   and   tumorigenesis   of   intestinal   epithelia.   *Elife*   **13**   (2024).
https://doi.org/10.7554/eLife.98238

44   Sanman, L. E. *et al.* Transit-Amplifying Cells Coordinate Changes in Intestinal Epithelial Cell-Type
Composition. *Dev Cell* **56**, 356-365 e359 (2021). https://doi.org/10.1016/j.devcel.2020.12.020

45   Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of single-cell
gene expression data. *Nat Biotechnol* **33**, 495-502 (2015). https://doi.org/10.1038/nbt.3192

46   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic
data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420 (2018).
https://doi.org/10.1038/nbt.4096

47   Scholzen, T. & Gerdes, J. The Ki-67 protein: from the known and the unknown. *J Cell Physiol* **182**,
311-322   (2000).   https://doi.org/10.1002/(SICI)1097-4652(200003)182:3<311::AID-
JCP1>3.0.CO;2-9

48   Olsen, L., Bressendorff, S., Troelsen, J. T. & Olsen, J. Differentiation-dependent activation of the
human intestinal alkaline phosphatase promoter by HNF-4 in intestinal cells. *American journal of*
physiology.   Gastrointestinal   and   liver   physiology   **289**,   G220-226   (2005).
https://doi.org/10.1152/ajpgi.00449.2004

49   Sheikh, A. *et al.* Enterotoxigenic Escherichia coli heat-labile toxin drives enteropathic changes in
small   intestinal   epithelia.   *Nature   communications*   **13**,   6886   (2022).
https://doi.org/10.1038/s41467-022-34687-7

50   Platts-Mills, J. A. *et al.* Association between enteropathogens and malnutrition in children aged 6-
23 mo in Bangladesh: a case-control study. *Am J Clin Nutr* **105**, 1132-1138 (2017).
https://doi.org/10.3945/ajcn.116.138800

51   Rogawski, E. T. *et al.* Use of quantitative molecular diagnostic methods to investigate the effect of
enteropathogen infections on linear growth in children in low-resource settings: longitudinal
analysis of results from the MAL-ED cohort study. *Lancet Glob Health* **6**, e1319-e1328 (2018).
https://doi.org/10.1016/S2214-109X(18)30351-6

52   Sheikh, A. *et al.* Enterotoxigenic Escherichia coli heat-labile toxin drives enteropathic changes in
small   intestinal   epithelia.   *Nature   communications*   **13**,   6886   (2022).
https://doi.org/10.1038/s41467-022-34687-7

53   Valenta, T., Hausmann, G. & Basler, K. The many faces and functions of beta-catenin. *EMBO J* **31**,
2714-2736 (2012). https://doi.org/10.1038/emboj.2012.150

54   MacDonald, B. T., Tamai, K. & He, X. Wnt/beta-catenin signaling: components, mechanisms, and
diseases. *Dev Cell* **17**, 9-26 (2009). https://doi.org/10.1016/j.devcel.2009.06.016

55   Fleckenstein, J. M. & Kuhlmann, F. M. Enterotoxigenic Escherichia coli Infections. *Curr Infect Dis*
Rep **21**, 9 (2019). https://doi.org/10.1007/s11908-019-0665-x

56   Fleckenstein, J. M. *et al.* Molecular mechanisms of enterotoxigenic Escherichia coli infection.
Microbes Infect **12**, 89-98 (2010). https://doi.org/S1286-4579(09)00235-4 [pii]
10.1016/j.micinf.2009.10.002

57   Chen, Y. *et al.* Ser9 phosphorylation of GSK-3beta promotes aging in the heart through suppression
of autophagy. *J Cardiovasc Aging* **1** (2021). https://doi.org/10.20517/jca.2021.13

58   Fang, D. *et al.* Phosphorylation of beta-catenin by AKT promotes beta-catenin transcriptional
activity. *J Biol Chem* **282**, 11221-11229 (2007). https://doi.org/10.1074/jbc.M611871200

59   Sassone-Corsi, P. The cyclic AMP pathway. *Cold Spring Harb Perspect Biol* **4** (2012).
https://doi.org/10.1101/cshperspect.a011148

60   Brown, A. C. *et al.* The cAMP responsive element modulator (CREM) transcription factor influences
susceptibility to undernutrition and infection. *mBio*, e0139025 (2025).
https://doi.org/10.1128/mbio.01390-25

61   Vemuri, K., Radi, S. H., Sladek, F. M. & Verzi, M. P. Multiple roles and regulatory mechanisms of the
transcription factor HNF4 in the intestine. *Front Endocrinol (Lausanne)* **14**, 1232569 (2023).
https://doi.org/10.3389/fendo.2023.1232569

62   Sands, W. A. & Palmer, T. M. Regulating gene transcription in response to cyclic AMP elevation.
Cell Signal **20**, 460-466 (2008). https://doi.org/10.1016/j.cellsig.2007.10.005

63   Alaullah Sheikh, B. T., Tim J Vickers, John C Martin, Bruce A Rosa, Subrata Sabui, Supratim Basu,
Rita D Simoes, Makedonka Mitreva, Chad Storer, Erik Tyksen, Richard D Head, Wandy Beatty,
Hamid M Said, James M Fleckenstein. Enterotoxigenic Escherichia coli heat-labile toxin drives
enteropathic changes in small intestinal epithelia. *Nature communications* **Nov 12;13(1):6886**
(2022). https://doi.org/https://doi.org/10.1038/s41467-022-34687-7

64   Chen, L. *et al.* A reinforcing HNF4-SMAD4 feed-forward module stabilizes enterocyte identity. *Nat*
Genet **51**, 777-785 (2019). https://doi.org/10.1038/s41588-019-0384-0

65   Chen, L. *et al.* HNF4 factors control chromatin accessibility and are redundantly required for
maturation of the fetal intestine. *Development* **146** (2019). https://doi.org/10.1242/dev.179432

66   Sansom, O. J. *et al.* Loss of Apc in vivo immediately perturbs Wnt signaling, differentiation, and
migration. *Genes Dev* **18**, 1385-1390 (2004). https://doi.org/10.1101/gad.287404

67   Andreu, P. *et al.* Crypt-restricted proliferation and commitment to the Paneth cell lineage following
Apc loss in the mouse intestine. *Development* **132**, 1443-1451 (2005).
https://doi.org/10.1242/dev.01700

68   Malonga, T., Vialaneix, N. & Beaumont, M. BEST4(+) cells in the intestinal epithelium. *Am J Physiol*
Cell Physiol **326**, C1345-C1352 (2024). https://doi.org/10.1152/ajpcell.00042.2024

69   Kelly, P. *et al.* Gene expression profiles compared in environmental and malnutrition enteropathy
in Zambian children and adults. *EBioMedicine* **70**, 103509 (2021).
https://doi.org/10.1016/j.ebiom.2021.103509

70   Mulenga, C. *et al.* Epithelial Abnormalities in the Small Intestine of Zambian Children With
Stunting. *Front Med (Lausanne)* **9**, 849677 (2022). https://doi.org/10.3389/fmed.2022.849677

71   Zhao, X. *et al.* Bile Acid Profiling Reveals Distinct Signatures in Undernourished Children with
Environmental Enteric Dysfunction. *J Nutr* **151**, 3689-3700 (2021).
https://doi.org/10.1093/jn/nxab321

72   Malique, A. *et al.* NAD(+) precursors and bile acid sequestration treat preclinical refractory
environmental enteric dysfunction. *Science translational medicine* **16**, eabq4145 (2024).
https://doi.org/10.1126/scitranslmed.abq4145

73   Guerrant, R. L., Schorling, J. B., McAuliffe, J. F. & de Souza, M. A. Diarrhea as a cause and an effect
of malnutrition: diarrhea prevents catch-up growth and malnutrition increases diarrhea frequency
and duration. *Am J Trop Med Hyg* **47**, 28-35 (1992). https://doi.org/10.4269/ajtmh.1992.47.28

74   Owino, V. *et al.* Environmental Enteric Dysfunction and Growth Failure/Stunting in Global Child
Health. *Pediatrics* **138** (2016). https://doi.org/10.1542/peds.2016-0641

75   Vonaesch, P. *et al.* Identifying the etiology and pathophysiology underlying stunting and
environmental enteropathy: study protocol of the AFRIBIOTA project. *BMC Pediatr* **18**, 236 (2018).
https://doi.org/10.1186/s12887-018-1189-5

76   Rogawski, E. T. & Guerrant, R. L. The Burden of Enteropathy and "Subclinical" Infections. *Pediatr*
Clin North Am **64**, 815-836 (2017). https://doi.org/10.1016/j.pcl.2017.03.003

77   Kosek, M. N. & Investigators, M.-E. N. Causal Pathways from Enteropathogens to Environmental
Enteropathy: Findings from the MAL-ED Birth Cohort Study. *EBioMedicine* **18**, 109-117 (2017).
https://doi.org/10.1016/j.ebiom.2017.02.024

78   Black, R. E., Brown, K. H. & Becker, S. Effects of diarrhea associated with specific enteropathogens
on the growth of children in rural Bangladesh. *Pediatrics* **73**, 799-805 (1984).

79   Li, P. *et al.* Homeostatic control of the crypt-villus axis by the bacterial enterotoxin receptor
guanylyl cyclase C restricts the proliferating compartment in intestine. *The American journal of*
pathology **171**, 1847-1858 (2007). https://doi.org/10.2353/ajpath.2007.070198

80   Pitari, G. M. *et al.* Bacterial enterotoxins are associated with resistance to colon cancer. *Proc Natl*
Acad Sci U S A **100**, 2695-2699 (2003). https://doi.org/10.1073/pnas.0434905100

81   Pitari, G. M., Di Guglielmo, M. D., Park, J., Schulz, S. & Waldman, S. A. Guanylyl cyclase C agonists
regulate progression through the cell cycle of human colon carcinoma cells. *Proc Natl Acad Sci U*
S A **98**, 7846-7851 (2001). https://doi.org/10.1073/pnas.141124698

82   Pasetti, M. F. *et al.* Safety and immunogenicity in humans of enterotoxigenic Escherichia coli
double mutant heat-labile toxin administered intradermally. *NPJ Vaccines* **10**, 23 (2025).
https://doi.org/10.1038/s41541-025-01071-7

83   VanDussen, K. L. *et al.* Development of an enhanced human gastrointestinal epithelial culture
system to facilitate patient-based assays. *Gut* **64**, 911-920 (2015). https://doi.org/10.1136/gutjnl-
2013-306651

84   Miyoshi, H. & Stappenbeck, T. S. In vitro expansion and genetic modification of gastrointestinal
stem cells in spheroid culture. *Nat Protoc* **8**, 2471-2482 (2013).
https://doi.org/10.1038/nprot.2013.153

85   McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-Cell RNA
Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337 e324 (2019).
https://doi.org/10.1016/j.cels.2019.03.003

86   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression. *Genome Biol* **20**, 296 (2019).
https://doi.org/10.1186/s13059-019-1874-1

87   Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-
seq. *Science* **352**, 189-196 (2016). https://doi.org/10.1126/science.aad0501

1023