Title: A load-bearing function for the cytoplasmic membrane of *Escherichia coli*

Authors: Jiawei Sun[1], Miriam Min-Chen Cheng[1], Ellen van Wijngaarden[2],

Christopher J. Hernandez[3], Thomas G. Bernhardt[4,5], Kerwyn Casey Huang[1,6,7,†]

Affiliations:

[1]Department of Bioengineering, Stanford University, Stanford, CA 94305, USA

[2]Sibley School of Mechanical and Aerospace Engineering, Cornell University, Ithaca, NY 14850, USA

[3]Departments of Orthopaedic Surgery and Bioengineering and Therapeutic Sciences, UCSF

[4]Department of Microbiology, Harvard University Medical School, Boston, MA 02115, USA

[5]Howard Hughes Medical Institute, Harvard University Medical School, Boston, MA 02115, USA

[6]Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA 94305, USA

[7]Chan Zuckerberg Biohub, San Francisco, CA 94158, USA

21 †To whom correspondence should be addressed: kchuang@stanford.edu

23 Keywords: MreB; cytoplasmic membrane; cell wall; peptidoglycan; mechanical

24 coupling; osmotic shocks; membrane stiffness; elongasome; transglycosylase;

25 MltG

Abstract

The structural integrity of bacterial cells is traditionally attributed to the peptidoglycan cell wall, and more recently to the outer membrane, with the cytoplasmic membrane assumed to be mechanically passive. Cells lacking filaments of the actin homolog MreB are more bendable, suggesting a role for the cytoskeleton in cell stiffness. Here, we show that MreB does not stiffen the envelope directly, but instead mechanically couples the cell wall to the cytoplasmic membrane through its role in peptidoglycan synthesis, increasing resistance to bending. Under hyperosmotic stress, MreB relocalized to the poles, forming linkages that prevent membrane detachment from the cell wall and attenuate cytoplasmic contraction. Disruption of MreB filament formation, nutrient starvation, or inactivation of glycan elongation factors abolished or reduced this coupling, revealing that peptidoglycan biosynthesis actively mediates stress distribution across surface layers. Our findings redefine the bacterial envelope as a mechanically integrated composite, with the cytoplasmic membrane having substantial load-bearing capacity.

Introduction

The bacterial cell envelope is a versatile and dynamic structure that acts as a chemical barrier and supports cell growth and division. The envelope bears mechanical stresses from both inside the cell and the external environment[1], including forces due to fluid flow, solid substrates, and osmolarity changes[2]. In particular, a major function of the envelope is to withstand the turgor pressure generated by high solute concentrations in the cytoplasm[3,4]. In Gram-negative bacteria, the envelope consists of a peptidoglycan cell wall sandwiched by the cytoplasmic and outer membranes. The outer membrane is an asymmetric bilayer with phospholipids in the inner leaflet and lipopolysaccharides in the outer leaflet that have important roles in chemical permeability[5]. The cell wall is a rigid macromolecule that has well-recognized importance as an antibiotic target[6] and in resisting cell expansion due to turgor pressure[7]. Robust envelope growth and division requires coordination by a host of enzymes[8]. In many rod-shaped bacteria, an essential component of the elongation machinery is the actin homolog MreB, which forms filaments that bind to the inner surface of the cytoplasmic membrane and interact with the cell-wall synthesis machinery to control cell shape through the insertion pattern of peptidoglycan precursors[9].

The cytoplasmic membrane is often ignored in discussions of bacterial

mechanics. Nonetheless, previous studies hinted at an underappreciated load-

bearing capacity of the cytoplasmic membrane in resisting expansion due to

turgor pressure. The gain-of-function mutation *mlaA** in *E. coli* results in rapid

transfer of phospholipids from the cytoplasmic membrane to the outer

membrane during starvation, and this transfer causes the cytoplasmic membrane

to shrink by ~20% and detach from the cell wall[10,11]. Once the cytoplasmic

membrane of starved *mlaA** cells ruptures, its contents expand in volume to

reach approximately the original cell length pre-plasmolysis[10,11]. The ability of

these cells to persist in a shrunken state for several minutes, together with the

subsequent expansion, suggests that the cytoplasmic membrane alone can bear

turgor pressure upon material loss and hence has substantial load-bearing

capacity.

Recent studies have revealed that many other components of the envelope

beyond the cell wall can impact cellular mechanical integrity[12]. The outer

membrane has stiffness properties similar to those of the cell wall and is critical

for maintaining cell viability during rapid changes in external osmolarity[13].

Mechanical linkages between the cell wall and outer membrane such as Braun's

lipoprotein (Lpp) and OmpA strengthen the cell envelope and determine

periplasmic width[14-17]. Mechanosensitive channels in the cytoplasmic membrane

enable rapid osmolyte efflux to protect cells against hypoosmotic shocks[17,18].

Beyond the envelope itself, deletion of a number of other proteins can impact the

stiffness of *E. coli* cells[19]. In particular, disruption of MreB filaments by the

antibiotic A22 decreased the resistance of *E. coli* cells to bending[20]. One

interpretation of this finding is that MreB forms an ultrastructure that directly

increases the mechanical stiffness of the envelope, mimicking the function of its

eukaryotic counterpart actin[21]. However, the mechanism by which MreB confers

increased bending rigidity could involve its interaction partners within the cell-

wall synthesis machinery including MreC, MreD[22], RodA, and RodZ[23,24]. During

wall synthesis, MreB filaments translate approximately along the circumferential

direction[25-28]. Given this motion, it is currently unclear how mechanical forces

could be transmitted from MreB, bound to the inner surface of the cytoplasmic

membrane, to the rest of the cell envelope.

In this study, we investigate the mechanical contribution of MreB to the cell

envelope by quantifying cell wall deformation during hyperosmotic shocks.

Under moderate shocks that are sufficient to cause plasmolysis,

depolymerization of MreB by A22 did not significantly increase cell wall

deformation, suggesting that MreB does not directly contribute to envelope

stiffness. However, under large-magnitude oscillatory hyperosmotic shocks,

cytoplasmic contraction gradually decreased in cells with intact MreB, whereas

A22-treated cells with depolymerized MreB maintained the same degree of

plasmolysis in each successive shock, indicating that MreB filaments are required

to prevent cytoplasmic contraction from the cell wall. Fluorescence microscopy

revealed that MreB relocalizes to the cell poles during hyperosmotic conditions,

where it drives the formation of structural linkages between the cytoplasmic

membrane and the cell wall that enable polar wall synthesis. Perturbations of

certain cell-wall synthesis factors altered the coupling, indicating that the

elongation machinery forms a physical linkage between the cytoplasmic

membrane and the peptidoglycan. Using a mutant with a weak outer membrane,

we show that wall-less spheroplasts can transiently remain viable even after the

outer membrane ruptures, indicating that the cytoplasmic membrane can bear

substantial mechanical stress. Based on these results, we propose that rather than

directly strengthening the cell wall, MreB-directed cell-wall synthesis

mechanically couples the cytoplasmic membrane and the cell wall, enabling the

intrinsic stiffness of the cytoplasmic membrane to contribute to the overall

strength of the envelope.

Results

Disruption of MreB polymerization does not alter cell-wall stiffness

The previous observation that A22 treatment decreases the bending rigidity of *E.*

coli cells[20] on a short time scale before transcriptional reprogramming could occur

suggested that MreB might directly impact envelope stiffness. To test this

hypothesis, we queried whether MreB polymerization and membrane binding

affects other morphological responses to perturbations beyond cell bending.

First, we reasoned that if MreB filaments were stress bearing, then their

depolymerization would alter the force balance across the cell envelope and

hence cell size would change. We quantified the dimensions of cells during

steady-state growth in a microfluidic flow cell (Methods) with high temporal

resolution before and after addition of 100 µg/mL A22, which rapidly disrupted

MreB polymerization as expected (Movie S1). Cell width and length were

unaffected (Fig. 1a), indicating that MreB was not bearing levels of longitudinal

or circumferential stress relevant to cell morphology.

Nonetheless, it remained possible that MreB contributes to envelope stiffness by

bearing stress when cells experience mechanical perturbations that disrupt

steady-state growth. To investigate this possibility, we applied oscillatory

osmotic shocks in a microfluidic flow cell and measured the deformation of the

cell wall upon each hyperosmotic shock in the absence and presence of A22

(Methods). In this assay, log-phase cells were alternately subjected to

hyperosmotic (LB+400 mM sorbitol) and normal (LB) osmotic conditions (Fig.

1b). Each hyperosmotic shock led to contraction of the cell wall and was

sufficiently large in magnitude to reduce turgor pressure to zero, as evidenced by

plasmolysis of the cytoplasm (Fig. 1c). The stiffness of the cell envelope is related

to the degree of deformation upon removal of turgor[12]. As observed in previous

studies with 100-mM shocks[7], the magnitude of length and width contractions

was consistent across shock cycles (Fig. 1d). We found that the degree of

contraction was not significantly different between cells without or with 10 min

of treatment with 100 µg/mL A22 prior to the start of the shocks (Fig. 1d). Thus,

disruption of MreB polymerization does not affect the mechanical response of

the cell envelope to forces that induce contraction/extension.

Taken together, these results indicate that MreB does not contribute directly to

cell envelope stiffness, despite the significant impact of A22 on bending rigidity[20].

MreB promotes a physical coupling between the cytoplasmic membrane to the

cell wall

Since MreB and its associated components of the cell-wall synthesis machinery

form a transmembrane structure connecting the cytoplasmic membrane to the

cell wall[29], we reasoned that MreB may prevent detachment of the cytoplasmic

membrane from the cell wall during osmotic shocks. To accentuate the impact of

A22 treatment, we increased the magnitude of the osmotic shock oscillations to 1

M (Methods), which substantially inhibited *E. coli* cell growth (Fig. S1). To

counteract the induction of cell division by hyperosmotic shocks[30], we also

treated cells with the antibiotic cephalexin (50 μg/mL), which targets the key

division protein PBP3[31], in the microfluidic flow cell for 40 min prior to the

shocks (Methods). In cells pre-treated with A22 for 10 min, the degree of

cytoplasmic contraction was consistently large over the course of 10 osmotic

shock cycles (Methods, Fig. 1e). In contrast, in untreated cells, the degree of

cytoplasmic contraction gradually decreased over the cycles (Fig. 1e). During the

last few shocks, many untreated cells exhibited little plasmolysis (Fig. 1f),

suggesting that polymerized MreB gradually inhibited detachment of the

cytoplasm from the cell wall. In addition to inhibiting detachment of the

cytoplasm from the cell wall, the linkages mediated by MreB also caused the cell

wall to buckle under the negative pressure generated by cytoplasmic contraction

for 1 M shocks, resulting in thinning of the cylindrical region around midcell

(Fig. S2). Treatment with 80 μg/mL MP265, a structural analog of A22, led to

similar effects as A22 (Fig. S3, Methods), arguing against potential off-target

effects of A22. These findings indicate that MreB promotes physical coupling of

the cytoplasmic membrane to the rest of the cell envelope.

Plasmolysis results in relocalization of MreB to the cell poles

To probe the dependence of plasmolysis dynamics on oscillatory osmotic shock

magnitude, we tracked the subcellular localization of MreB in a strain expressing

an MreB-msfGFP sandwich fusion that complements *mreB* deletion[32,33] (Table S1).

During steady-state growth, MreB filaments localized primarily to the cylindrical

region of the cell (Fig. 2a)[33], as expected, where they guide the insertion of new

cell wall material[27]. During 1 M oscillatory osmotic shocks, the first few

hyperosmotic shocks induced cytoplasmic contraction mostly near the cell poles

(Fig. 2a), indicating the lack of coupling between the cytoplasmic membrane and

the rest of the envelope at the poles, where MreB was largely absent. However, in

subsequent osmotic shock cycles, MreB partially relocalized to the cell poles,

resulting in a more homogeneous distribution of MreB across the cell surface

(Fig. 2a-c). This gradual relocalization coincided with the decrease in cytoplasmic

contraction (Fig. 1e). In contrast, lower-magnitude (400 mM) osmotic shocks for

which cytoplasmic contraction remained constant over shock cycles (Fig. 1e)

resulted in less relocalization of MreB (Fig. 2c). Thus, we reasoned that repeated,

large-magnitude osmotic shocks led to the formation of MreB-mediated linkages

between the cytoplasmic membrane and the other envelope layers at the cell

poles that are stable over the 4-min period of the shock cycles, which limited

contraction during subsequent shock cycles.

To test whether hyperosmotic shocks themselves trigger MreB relocalization, we

varied the duration of hyperosmolarity and observed nearly identical

relocalization trajectories over assay time regardless of the number of shock

cycles (Fig. S4). These results indicate that MreB filaments relocalize

continuously during plasmolysis and recovery, reflecting a process that depends

on the time under osmotic perturbation rather than an acute effect of the shock.

Taken together, these findings suggest that hyperosmolarity drives relocalization

of MreB to the poles, where they appear to establish new physical linkages

between the cytoplasmic membrane and cell wall that limit cytoplasmic

contraction during subsequent osmotic shocks (Fig. 2d).

Peptidoglycan elongation couples the cytoplasmic membrane and cell wall

Since MreB resides solely on the inner surface of the cytoplasmic membrane, we

surmised that any physical connection to the cell wall must rely on other cellular

components. While many proteins interact with both the cytoplasmic membrane

and cell wall, presumably only a subset are critical for maintaining the overall

physical coupling between layers. We hypothesized that the most likely

candidates would be components of the cell-wall synthesis machinery.

Supporting this hypothesis, when we performed oscillatory osmotic shocks with

1 M sorbitol in a nutrient-free buffer osmotically balanced with LB (0.85X PBS,

Methods), regardless of A22 treatment, cells exhibited plasmolysis without a

reduction in cytoplasmic contraction over time (Fig. 3a). These data suggest that

linkage between the membrane and cell wall requires cell growth. Repeated

osmotic shocks in the nutrient-free media still resulted in MreB relocalization to

the poles (Fig. S5), indicating that MreB relocalization is not dependent on wall

synthesis and is not sufficient for linking the cytoplasmic membrane to the cell

wall.

To test whether MreB relocalization to the poles leads to polar wall synthesis,

rather than the side-wall synthesis typical of normal growth, we performed

pulse-chase experiments in cells labeled with 3-[[(7-Hydroxy-2-oxo-2H-1-

benzopyran-3-yl)carbonyl]amino]-D-alanine hydrocholoride (HADA), which is

inserted into the peptidoglycan at locations of growth[33,34]. During subsequent

growth in HADA-free medium (Methods), the ratio of fluorescence at the poles

relative to the cell body decayed more rapidly in cells undergoing hyperosmotic

shocks, indicating increased polar wall synthesis (Fig. 3b). Thus, polar

localization of MreB leads to relocalization of elongation machinery function.

We next hypothesized that proteins involved in cell-wall synthesis[29] contribute to

the membrane-cell wall connection. To identify such factors, we performed

oscillatory shocks in combination with genetic or chemical perturbation of

elongation-associated targets. First, we overexpressed RodA from an inducible

plasmid (Table S1) prior to osmotic shocks (Methods). Opposite to

depolymerization of MreB by A22, cytoplasmic contraction was reduced during

repeated plasmolysis compared with no induction (Fig. 3c). Overexpression of a

dominant-negative mutant $rodA^{D262N}$ (Table S1), which blocks the

glycosyltransferase activity of RodA[35], led to a similar phenotype (Fig. S6). These

results suggest that increasing the copy number of RodA is sufficient to enhance

the coupling between MreB and the cell envelope, regardless of RodA activity.

To assess the contribution of other elongation-associated PBPs, we inhibited their

activity using β-lactam antibiotics, which specifically target transpeptidase

function. Inhibition of PBP1A/B by cefsulodin, PBP2 by mecillinam, or all PBPs

by ampicillin had no effect on cytoplasmic contraction during the shocks (Fig.

S7), suggesting that transpeptidase function is not required for establishing the

cytoplasmic membrane–cell wall connection.

To probe the role of glycan strand synthesis, we focused on a deletion mutant of

mltG (Table S1), which encodes a cytoplasmic membrane-anchored enzyme

implicated in the termination of peptidoglycan synthesis[36]. Deletion of *mltG* is

predicted to enhance the association of the elongation machinery with the cell

wall and results in longer glycan strands[36]. Under oscillatory osmotic shocks,

Δ*mltG* cells exhibited a slower transition to the state of lower cytoplasmic

contraction compared with wild-type cells (Fig. 3f). Given the role of MltG in

termination[37], we reasoned that the accumulation of synthases at the cell poles

necessary for reducing cytoplasmic contraction was delayed in Δ*mltG* cells.

Taken together, these findings highlight the functional duality of the elongation

machinery in both cell-wall synthesis and as a mechanically robust connection

between the cytoplasmic membrane and the cell wall.

The cytoplasmic membrane has substantial load-bearing capacity

Our findings collectively suggest that MreB does not directly contribute to

envelope stiffness, and instead promotes physical coupling between the

cytoplasmic membrane to the cell wall via its role in cell-wall synthesis. This

linkage is unlikely to provide substantial stiffness in the longitudinal or

circumferential directions along the surface of the cell, since the direction of

forces exerted to link the layers would be predominantly perpendicular to the

surface. However, if the cytoplasmic membrane itself were load bearing, such a

coupling could explain the substantial decrease in bending rigidity under A22

treatment[20], wherein MreB (and the coupling it promotes) is required for bending

to engage the mechanical response of the cytoplasmic membrane (Fig. 4a). While

it is typically assumed that the cytoplasmic membrane is relatively flexible

compared to the cell wall by virtue of it being a lipid bilayer[12,38], previous

molecular dynamics simulations suggested that a protein-rich membrane patch

mimicking the cytoplasmic membrane can exhibit substantial stiffness

comparable to that of the cell wall and the outer membrane[39].

To directly test the capacity of the inner membrane to bear stress, we sought to

reduce or remove the stress-bearing capacity of the cell wall and outer

membrane. Cell wall-free spheroplasts can be generated with lysozyme[40].

However, the outer membrane is thought to be essential, potentially due to

osmotic balancing in the periplasm[17], and no mechanism currently exists for

robustly removing the outer membrane while retaining cell viability[12]. Thus, we

leveraged our previous finding that deletion of *bamD* results in a highly

mechanically compromised outer membrane due to a drastic reduction in outer

membrane protein levels[41]. We successfully generated wall-free Δ*bamD*

spheroplasts using 20 μg/mL lysozyme, indicating that even without a cell wall,

at least some cells with a highly mechanically compromised OM can persist in a

rich medium without osmoprotection (Fig. 4b).

To monitor changes to the cell envelope during and after cell-wall digestion, we

tracked Δ*bamD* cells expressing cytoplasmic GFP and periplasmic mCherry in a

microfluidic flow cell (Methods). Upon lysozyme introduction, cells lost their rod

shape and became spherical (Fig. 4c). Without the stiff cell wall that typically

bears turgor pressure, the mechanical load shifted to the two membranes,

resulting in a ~20% increase in the surface area of the cytoplasmic membrane

(Fig. 4d, Methods). Approximately half of Δ*bamD* spheroplasts persisted for >10

min with an expanded cytoplasmic membrane before lysis, highlighting the

ability of the membrane to bear turgor pressure. The spheroplasts survived for a

shorter period compared to in a liquid environment (Fig. 4b), likely due to the

mechanical confinement of the thin microfluidic flow cell. During the process of

lysis, Δ*bamD* spheroplasts lost the periplasmic mCherry signal earlier than or

simultaneously with the loss of cytoplasmic GFP (Fig. 4c), indicating that the

outer membrane ruptured before or at the same time as the cytoplasmic

membrane. After outer membrane rupture, ~50% of spheroplasts persisted with

an intact cytoplasmic membrane for >5 min. In these cells, outer membrane

rupture caused almost no change in the cytoplasm area (Fig. 4e), suggesting that

the turgor pressure was borne solely by the cytoplasmic membrane rather than

the mechanically compromised outer membrane. These findings provide direct

evidence of the load-bearing capability of the cytoplasmic membrane, such that

when coupled to the cell wall by the process of cell-wall synthesis, the

cytoplasmic membrane can increase the stiffness of the cell envelope.

Discussion

In this study, we provide a revised perspective on the origins of mechanical resilience of the cell envelope, and demonstrate that the cell-wall synthesis machinery has a dual catalytic–structural function in integrating growth with envelope mechanics. We found that MreB does not contribute directly to cell wall stiffness under repeated plasmolysis (Fig. 1a,d) but instead functions as a platform for the components of the cell-wall synthesis machinery to link the cytoplasmic membrane and the cell wall sufficiently strongly as to prevent plasmolysis. Fluorescently labeled MreB filaments relocalized to the cell poles during large-magnitude osmotic shocks (Fig. 2), which in turn limited cytoplasmic contraction in these regions during periods of hyperosmolarity and led to peptidoglycan synthesis at the poles (Fig. 3b) as long as nutrients were available (Fig. 3a). Among the many components involved in cell-wall elongation, overexpression of RodA (Fig. 3c) and removal of MltG (Fig. 3e) impacted the connection between the cytoplasmic membrane and cell wall. Motivated by these findings and our evidence of substantial load-bearing capacity of the cytoplasmic membrane (Fig. 4), we propose a biophysical model in which MreB and active peptidoglycan synthesis are necessary for mechanically coupling the cytoplasmic membrane to the rest of the cell envelope. This model provides an explanation for the previously observed changes in

bending rigidity during A22 treatment[20]: with intact MreB filaments, the

cytoplasmic membrane can contribute to resisting bending. In fact, the ~30%

decrease in bending rigidity during A22 treatment is in line with expectations

based on detaching a single layer from an envelope with three layers of roughly

equal mechanical importance.

The proteins that physically link bacterial surface layers are often critical to

cellular mechanical integrity. Braun's lipoprotein (Lpp) and certain β-barrels

connect the OM to the cell wall and impact the ability of the outer membrane to

bear load under compression[13]. Lpp also regulates periplasmic width and the

stiffness of *E. coli* cells[14]. However, the linkages between the cell wall and the

cytoplasmic membrane have largely been overlooked, in part due to the lack of

suitable assays for interrogating this connection. Our findings highlight the

dynamic nature of the cytoplasmic membrane-peptidoglycan linkage. High-

magnitude oscillatory osmotic shocks induced linkage formation at otherwise

uncoupled cell poles, a process dependent on the relocalization of MreB and

repositioning of at least certain components of the elongation machinery. The

linkages, which are likely transient rather than persisting as static structures,

may rely on several factors. While an obvious candidate is crosslinked glycans,

our results suggest that, although they may contribute, they are not required.

RodA does not extend far enough into the periplasm to bind peptidoglycan,

suggesting that other factors such as RodZ or MreC mediate the connection.

Potential redundancy or decoupling of the activity and binding of the factors

involved may complicate elucidating the precise makeup and dynamics of the

coupling. Nonetheless, it is striking that A22 treatment alone is sufficient to alter

cytoplasmic contraction under repeated osmotic shocks (Fig. 1E), highlighting

the key role of MreB in organizing the peptidoglycan synthesis machinery.

One open question raised by our work is the mechanism by which MreB

relocalizes to the cell poles during large hyperosmotic shocks. Previous studies

found that MreB is enriched in log-phase cells at regions of the membrane with

negative mean curvature[33,42], which has been hypothesized to underlie a feedback

control mechanism for maintaining rod-like shape[9]. Plasmolysis generates

regions of negative mean curvature near the poles, as indicated by the

cytoplasmic membrane dye MitoTracker Green[43,44] (Fig. S8a), and cytoplasmic

contraction could generate wrinkles in the membrane with large absolute

curvature that could either bias the localization of MreB filaments based on

geometric preference or by disrupting their circumferential motion[45]. However,

MreB relocalized to cell poles at a rate independent of whether obvious

plasmolysis occurred at that pole (Fig. S8b), indicating that at least large

deformations are not necessary for relocalization. Regardless of the mechanism,

the ability of MreB to target distinct regions of the cell depending on

environmental cues such as osmolarity may provide bacterial cells with

morphological flexibility. This flexibility is apparent during the transition into

and out of stationary phase, during which MreB also exhibits polar localization[24],

which our study suggests could be due to osmolarity changes.

In addition to mechanically strengthening the cell envelope, linkages between

surface layers control the distance between layers[14], which can be critical for the

function of proteins that span the space[46-48]. In the case of MreB, the cell-wall

synthesis-mediated linkages are mostly present only along the cylindrical cell

body during log-phase growth. Thus, during a small osmotic perturbation, the

cell poles can function as a temporary buffer for water efflux, while the distance

between the cytoplasmic membrane and the cell wall along the rest of the

cylindrical body is not affected, allowing for continued growth even in the

absence of turgor[13,49]. In the future, it will be interesting to determine whether

MreB-directed cell-wall synthesis plays a similar mechanical function in other

species. Regardless, a deeper understanding of the mechanical interactions

between the cytoplasmic membrane and the cell wall provides new insights into

the structural stability of the cell envelope and reveals potential targets and

mechanisms for antimicrobial intervention.

Methods

Bacterial strains and growth conditions

All strains and plasmids used in this study are listed in Table S1. *E. coli* cells were

grown overnight in lysogeny broth (LB, Lennox, Fisher Scientific, Cat. #BP9722-

500) at 37 °C, diluted into fresh LB, and grown for 2 h into log phase for

microfluidic measurements. Strains were grown with 50 µg/mL kanamycin

(Sigma-Aldrich, Cat. #K4000) or 50 µg/mL chloramphenicol (Calbiochem, Cat.

#220551) as necessary.

The Δ*mltG* strain expressing cytoplasmic GFP was generated by λ-Red

recombination using pSIM6. A chloramphenicol resistance cassette (cmR)

flanked by FRT sites was PCR-amplified from pKD3 with primers containing the

same 50-nt homology overhangs upstream and downstream of *mltG* as the Keio

collection[50] (forward primer: 5'-

ACGTTATATGAATATTTAGCCCCACTTTGTGAGCGCCCGAATTAGTCATGt

ccatatgaatatcctccttagttcctattcc-3'; reverse primer: 5'-

CGCCTTCCAGCCCCTCAATGACGATATACTTACTGCGCATTTTTTTCCTTtgt

aggctggagctgcttcg-3'). The PCR product was electroporated into BW25113

carrying pSIM6, and recombinants were selected on LB agar with 50 µg/mL

chloramphenicol. The strain was subsequently transformed with pZS21-GFP

(Table S1) to drive constitutive GFP expression.

Microfluidic hyperosmotic shock assays

Cells from a saturated culture grown overnight were diluted 1:200 into fresh LB

and grown at 37 °C for 2 h. These log-phase cells were cultured with 25 μg/mL

wheat germ agglutinin (WGA) conjugated to AlexaFluor-555 (AF555, Invitrogen

W32464) to label the outer membrane. Cells were then loaded into CellASIC

ONIX microfluidic flow cells (Sigma-Aldrich, Cat. #B04A-03-5PK), in which

media were exchanged using the CellASIC ONIX2 microfluidic platform (Sigma-

Aldrich, Cat. #CAX2-S0000). Oscillatory hyperosmotic shock assays were

performed at room temperature.

Cells were first perfused with LB for 30 min with 50 μg/mL cephalexin (MP

Biomedicals, Cat. #150585) to inhibit cell division and then treated with 100

μg/mL A22 (gift from Douglas Weibel), 80 μg/mL MP265 (MedChemExpress,

Cat. #HY-131583), a beta-lactam antibiotic (100 μg/mL ampicillin (RPI, Cat.

#A40040), 50 μg/mL mecillinam (Sigma-Aldrich, Cat. #33447), or 30 μg/mL

cefsulodin (MP Biomedicals, Cat. #198677), or 0.85X PBS (Gibco, Cat. #70011044)

for 10 min to perturb cell-wall synthesis. Cells were then subjected to oscillatory

hyperosmotic shocks involving 400-, 600-, or 1000-mM sorbitol (Sigma-Aldrich,

Cat. #S1876, dissolved in LB or 0.85X PBS) and recovery without sorbitol, with a

period of 4 min (2 min shock and 2 min recovery). Pre-treatment compounds

(A22, MP265, beta-lactam, or 0.85x PBS) were maintained during osmotic shocks.

Cells were stained with 25 μg/mL WGA-AF555 throughout incubation in the

flow cell.

To overexpress RodA or RodA^{D262N} (Table S1), cells were induced with 1 mM

IPTG (Sigma-Aldrich, Cat. #I6758) for 1 h in liquid culture before loading to the

microfluidic flow cell. After loading, cells were incubated with 1 mM IPTG for an

additional 30 min, and IPTG was maintained throughout subsequent osmotic

shock assays.

Fluorescent labeling of peptidoglycan and cytoplasmic membrane

A saturated culture grown overnight in LB medium was diluted 1:200 into LB

medium containing 1 mM HADA (MedChemExpress, Cat. #HY-131045/CS-

0124027) and grown at 37 °C for 2 h to ensure homogenous peptidoglycan

labeling. Cells with fluorescently labeled peptidoglycan were then loaded into

microfluidic flow cells and grown in LB with 1 mM HADA at room temperature,

followed by hyperosmotic shocks with 1 mM HADA. Cells were then switched

to HADA-free LB medium to track peptidoglycan synthesis as indicated by the

decay of HADA signals.

To visualize the cytoplasmic membrane during osmotic shocks, cells were

labeled with 2 µg/mL MitoTracker Green (Invitrogen, Cat. #M7514) in liquid

culture and in the microfluidic flow cell.

Single-cell imaging

Cells undergoing oscillatory hyperosmotic shocks were monitored using a Nikon

Eclipse Ti-E inverted epifluorescence microscope with a 100X (NA 1.40) oil-

immersion objective (Nikon Instruments). Phase-contrast and fluorescence

images were collected on a Prime BSI Express sCMOS camera (Teledyne

Photometrics) using *μManager* v. 2.0[51].

Image analysis

Time-lapse images were segmented with the machine learning-based software

DeepCell[52] trained on WGA-stained cells, and the segmented images were

analyzed using *Morphometrics*[52] to obtain cell contours at sub-pixel resolution.

Using these contours, single-cell lineages were tracked. Cells that were

segmented or tracked incorrectly or that lysed during osmotic shocks were

filtered out via manual inspection. Cell length and width were calculated using

the *MicrobeTracker*[53] meshing algorithm. Cytoplasmic area was inferred using

cytoplasmic GFP fluorescence, and cytoplasmic contraction during plasmolysis

was computed based on the fraction of the total area enclosed by the cell contour

represented by the cytoplasmic area.

To determine the spatial localization of MreB filaments in cells expressing MreB-

GFP, in Fig. 2b, fluorescence profiles along cell contours were calculated as

previously described[54] and the degree of fluorescence along the cylindrical region

and the cell poles was compared. In Fig. 2c, S4, and S6b, the average fluorescence

intensity was calculated for the projected cylindrical region and both cell poles,

where the highly curved polar regions were defined based on a curvature

threshold.

Cell shape analysis

Cell wall area contraction in Fig. 1d was calculated as (area plasmolysis

contraction)/(area after plasmolysis)–1. Cytoplasmic area expansion in Fig. 4e

was calculated as (area after OM rupture)/(area before OM rupture)–1. Cell

width was extracted from the pill mesh generated by the *MicrobeTracker*[53]

meshing algorithm and defined as the maximum width measured perpendicular

to the cell's long axis. Cell width increase in Fig. S3b was calculated as (width

under shock)/(width prior to shock)–1.

The cytoplasmic membrane area in Fig. 4d was calculated separately for rod-

shaped cells prior to wall digestion and for spheroplasts after digestion using the

same approximation as in ref. [13]. Cell contours were generated using

Morphometrics[52] from cytoplasmic GFP fluorescence. For rod-shaped cells, the

membrane area was estimated from the contour assuming rotational symmetry

around the long axis. For spheroplasts, which adopted amorphous shapes within

the thin flow cell, the flow cell height was assumed equal to the cell width prior

to wall digestion, w. The projected area, a, and perimeter, s, were calculated from

the contour, and the cytoplasmic membrane area was then estimated as $2a+ws$.

Spheroplast generation

To generate cell wall-free spheroplasts with a compromised OM, Δ*bamD* cells

(Table S1) were grown in LB for 2 h and treated with 100 μg/mL lysozyme

(Sigma-Aldrich, Cat. #L6876) in LB. To generate and monitor spheroplasts in a

microfluidic flow cell, Δ*bamD* cells were treated with 100 μg/mL lysozyme in LB.

Expression of periplasmic mCherry was induced with 1 mM IPTG.

Data and code availability

530   All data used for generating figures in this study are available at the Stanford

531   Digital Repository: https://purl.stanford.edu/gg063dh4581.

Figures

Figure 1: MreB couples the cytoplasmic membrane to the cell wall.

535   a) Disruption of MreB filaments with 100 μg/mL A22 in LB medium did

536   not alter cell length or width, suggesting that MreB filaments are not

537   under stress during exponential growth. A22 treatment started at *t* = 10

538   min.

539   b) Schematic of the oscillatory hyperosmotic shock assay in which cells

were periodically subjected to hyper- and normal osmolarity

(Methods). Under normal osmolarity, the cell wall is under tension due

to turgor pressure; under hyperosmolarity, the cytoplasm loses water

and detaches from the cell wall.

c) Exposure to 400 mM hyperosmotic shock cycles led to cytoplasmic

contraction, typically seen at the cell poles (red/white arrows highlight

one example). The OM and the cytoplasm are labeled with wheat germ

agglutinin (WGA) AlexFluor-555 and GFP expressed from a plasmid,

respectively. Scale bar: 5 µm.

d) Disruption of MreB filaments via pre-treatment with 100 µg/mL A22

did not alter cell wall contraction (Methods) under 400 mM oscillatory

hyperosmotic shocks. **: $p < 0.01$, n.s.: $p > 0.01$; Welch's t-test was

performed for each osmotic shock cycle.

e) Inhibition of cytoplasmic contraction by MreB in plasmolyzed cells

was dependent on the number and magnitude of the oscillatory

osmotic shocks. The impact of A22 pre-treatment on cytoplasmic

contraction was prominent only under large hyperosmotic shocks (600

mM and 1 M).

f) The presence of MreB filaments resulted in mechanical coupling

between the cytoplasmic membrane and the cell wall (red arrows);

560   disruption of MreB by A22 allowed unhampered plasmolysis (white

561   arrows). Cells are labeled with WGA AF-555 and cytoplasmic GFP

562   expressed from a plasmid. Scale bar: 5 μm.

Figure 2: Hyperosmolarity causes relocalization of MreB filaments to the poles.

a) 1-M oscillatory osmotic shocks resulted in relocalization of MreB-

    msfGFP filaments to cell poles (red arrows), which are generally

    depleted of MreB-msfGFP during normal growth (white arrows). Scale

    bar: 5 µm.

b) MreB-msfGFP profiles of individual cell poles, averaged across the

    contours of *n*=348 cells, demonstrate the increasingly homogenous

    distribution of MreB after more 1-M hyperosmotic shocks. The profiles

    span the pole and surrounding region (left schematic).

c) The increase in the ratio of MreB-msfGFP intensity at the cell poles compared to the cylindrical cell body (left schematic) demonstrates the relocalization of MreB to the poles. The degree of relocalization was dependent on the shock magnitude. The sawtooth behavior is due to repeated plasmolysis and subsequent recovery.

d) Schematic of MreB relocalization under hyperosmotic shocks and formation of new mechanical linkages at the cell poles.

Figure 3: Inactivation of cell-wall synthesis alters the coupling between the cytoplasmic membrane and the peptidoglycan.

a) In 0.85X PBS, wild-type cells exhibited plasmolysis without any reduction in cytoplasmic contraction over repeated 1-M osmotic shocks, regardless of A22 treatment.

b) Pulse-chase experiment in cells pre-labeled with 1 mM HADA. Compared to cells grown without shocks, cells exposed to shocks exhibited greater decreases in fluorescence at the poles, when

evaluated at the same level of fluorescence decay in the cell body,

indicating increased polar peptidoglycan synthesis following osmotic

shock-induced MreB relocalization. Top: schematic of the HADA

labeling and hyperosmotic shock protocol. Cells were tracked in a

microfluidic flow cell during growth without HADA starting at t=0.

Each data point corresponds to an individual cell pole. $N \geq 25$ cells were

tracked for each condition. **: $p < 0.01$, ***: $p < 0.001$, n.s.: $p > 0.01$;

Welch's t-test was performed on binned data points.

c) Overexpression of RodA reduced cytoplasmic contraction during 600-

mM osmotic shocks.

d) $\Delta mltG$ cells exhibited a slower transition to the coupled state compared

to wild type during 1-M shocks. The degree of contraction was

significantly different (Welch's t-test: $p < 0.01$) in the first 3 shocks and

similar after $n>3$ shocks.

Figure 4: The cytoplasmic membrane has the capacity to bear substantial

mechanical stress.

  a) MreB-mediated coupling between the cytoplasmic membrane and the

    cell wall may explain the decrease in bending rigidity of *E. coli* cells

    upon A22 treatment[20] if the cytoplasmic membrane is sufficiently stiff.

  b) Spheroplasts without a cell wall can be generated from Δ*bamD* cells

    with weak outer membranes using 20 μg/mL lysozyme in LB medium

    without additional osmoprotection. Cells were imaged 3 h after

    lysozyme treatment (Methods). The panel shows a composite of

    representative snapshots of individual spheroplasts. Scale bar: 5 μm.

  c) Schematics and images during the generation of Δ*bamD* spheroplasts,

rupture of the weakened outer membrane, and lysis due to inner

membrane rupture. Δ*bamD* cells were labeled with cytoplasmic GFP

and periplasmic mCherry and tracked in a microfluidic flow cell

(Methods). Loss of periplasmic mCherry signal indicates rupture of the

outer membrane, which always occurred prior to or at the same time

as cytoplasmic membrane rupture. Scale bar: 5 μm.

d) The surface area of the cytoplasmic membrane increased when the cell

wall was degraded, indicating that the remaining envelope layers

(cytoplasmic and outer membranes) bore more of the stress due to

turgor pressure in the absence of the wall. *n*=35 cells.

e) Rupture of the outer membrane in Δ*bamD* spheroplasts resulted in

only a slight increase in cytoplasmic area, suggesting that the outer

membrane was not bearing substantial stresses in the spheroplasts.

n=17 cells.

629 **Supplementary Figures**

632 **Figure S1: Cell growth is inhibited under large hyperosmotic shocks.**

633 During a hyperosmotic shock with 1 M sorbitol in LB, cell growth was inhibited

634 for an extended period. Cells were monitored in a microfluidic flow cell

635 (Methods). *n*=22 cells.

Figure S2: When mechanically coupled to the cytoplasm, the cell wall exhibits

large deformations and increases in cell width during large hyperosmotic

shocks.

a) After several osmotic shock cycles (1 M sorbitol in LB), when the cell wall

is mechanically coupled with the cytoplasmic membrane by MreB

filaments, the cell wall becomes flattened by the negative osmotic pressure

from inside the cytoplasm under high environmental osmolarity, resulting

in increased cell width around midcell. These large deformations only

occurred under large (600 mM or 1 M) osmotic shocks.

b) Cell width increased after several 1-M hyperosmotic shock cycles. This

widening was abolished by A22 pre-treatment, with cells contracting in

width upon loss of turgor.

c) Schematic of cell flattening caused by the mechanical linkages and

osmotic pressure. Red and blow arrows indicate extension and

compression along different axes of the cross-section.

Figure S3: Disruption of MreB filaments with MP265 eliminates the mechanical coupling between the cytoplasmic membrane and the cell wall during oscillatory osmotic shocks. Wild-type cells were treated with 80 µg/mL MP265 10 min prior to and during 1-M osmotic shocks. The lack of coupling is similar to the effects of A22 treatment (Fig. 3e).

Figure S4: MreB relocalization to the poles is determined by the time under

osmolarity oscillation rather than the number of shock cycles. Hyperosmotic

shocks with extended periods (4 min, 10 min) of hyperosmolarity (upper

schematic) resulted in similar degrees of MreB relocalization as with a 2-min

664   period for a given assay time. Data points represent individual osmotic shock

665   cycles. Fluorescence was evaluated during times of normal osmolarity to avoid

666   bias caused by plasmolysis.

Figure S5: MreB relocalizes to the poles during oscillatory osmotic shocks in a nutrient-free buffer. In 0.85X PBS, MreB relocalization due to osmotic shocks progressed at a similar rate as in LB.

Figure S6: Overexpression of the dominant negative mutant $rodA^{D262N}$ reduced

674   **cytoplasmic contraction during 600-mM osmotic shocks.**

Figure S7: Inhibition of PBPs by 50 μg/mL mecillinam, 30 μg/mL cefsulodin, or 100 μg/mL ampicillin did not impact the dynamics of cytoplasmic contraction during 1-M shocks.

Figure S8: MreB relocalization to each cell poles during hyperosmotic shocks

is not dependent on the magnitude of cytoplasmic contraction at the pole.

a) Plasmolysis resulted in clear regions with negative mean curvature near

cell poles. The cytoplasmic membrane was stained with MitoTracker

Green. Scale bar: 5 μm.

b) During plasmolysis, many cells exhibited cytoplasmic contraction at only

one cell pole. No significant difference was observed between MreB

relocalization at the two poles after repeated shocks. Fluorescence was

evaluated during times of normal osmolarity to avoid bias caused by

plasmolysis. n=11 cells. **: $p < 0.01$, n.s.: $p > 0.01$; Welch's t-test was

performed for each osmotic shock cycle.

 Supplementary Table

Strain	Genotype	Source/reference
MG1655	F⁻, lambda⁻, *rph-1*	CGSC #6300
KC1016	MG1655 pZS21-GFP	Gift from Thomas Silhavy
KC508	MG1655 *mreB-msfGFP*	[33]
KC496	MG1655 Δ*lacIZYA*::frt Δ*ponA*::frt attλ::pTB309(*Para*::*ponA*)	[55]
pPR1	MG1655 *Ptac*::*pbpA-rodA*^{D262N}	[35]
pPR6	MG1655 *Ptac*::*pbpA-rodA*	[35]
BW25113	F⁻ λ⁻ Δ(*araD-araB*)567 Δ*lacZ4787*(::*rrnB-3*) *rph-1* Δ(*rhaD-rhaB*)568 *hsdR514* pZS21-GFP	Gift from Jie Xiao
Δ*mltG*	BW25113 Δ*mltG*::*cmR* pZS21-GFP	This study
IMB1056	MC4100 *ara*^{R} *bamA*^{E470K} Δ*bamD*	[41]
IMB1144	IMB1056 *attHK*::*Plac-mCherry* pZS21-GFP	[41]
Plasmids	**Genotype**	**Source/reference**
pZS21-GFP	*pZS21kan*::*GFP*	Gift from Thomas Silhavy
pPR1	*Ptac*::*pbpA-rodA*^{D262N}	[35]

| pPR6 | *Ptac::pbpA-rodA* | [35] |

Table S1: *E. coli* strains and plasmids used in this study.

Supplemental Movies

Supplemental Movie 1: A22 treatment (100 μg/mL) depolymerizes MreB

filaments within 10 min in LB medium. Scale bar: 5 μm.

Acknowledgements

The authors thank the Huang lab for helpful discussions. The authors acknowledge support from a Stanford Interdisciplinary Graduate Fellowship (to J.S.), NIH RM1 Award GM135102 (to K.C.H.), and NSF Award EF-2125383 (to K.C.H.). K.C.H. is a Chan Zuckerberg Biohub Investigator. This work was also supported in part by the National Science Foundation under Grant PHYS-1607611 and the hospitality of the Aspen Center for Physics.

References

Mitchell, A. M. & Silhavy, T. J. Envelope stress responses: balancing damage repair and toxicity. *Nature Reviews Microbiology* **17**, 417-428 (2019).

Persat, A. *et al.* The mechanical world of bacteria. *Cell* **161**, 988-997 (2015).

Rojas, E. R. & Huang, K. C. Regulation of microbial growth by turgor pressure. *Current Opinion in Microbiology* **42**, 62-70 (2018).

Deng, Y., Sun, M. & Shaevitz, J. W. Direct measurement of cell wall stress stiffening and turgor pressure in live bacterial cells. *Physical review letters* **107**, 158101 (2011).

Silhavy, T. J., Kahne, D. & Walker, S. The bacterial cell envelope. *Cold Spring Harbor perspectives in biology* **2**, a000414 (2010).

Hȯltje, J.-V. Growth of the stress-bearing and shape-maintaining murein sacculus of Escherichia coli. *Microbiology and molecular biology reviews* **62**, 181-203 (1998).

Rojas, E., Theriot, J. A. & Huang, K. C. Response of Escherichia coli growth rate to osmotic shock. *Proceedings of the National Academy of Sciences* **111**, 7807-7812 (2014).

Egan, A. J., Errington, J. & Vollmer, W. Regulation of peptidoglycan synthesis and remodelling. *Nature Reviews Microbiology* **18**, 446-460 (2020).

9   Shi, H., Bratton, B. P., Gitai, Z. & Huang, K. C. How to build a bacterial

cell: MreB as the foreman of E. coli construction. *Cell* **172**, 1294-1305 (2018).

10   Sutterlin, H. A. *et al.* Disruption of lipid homeostasis in the Gram-negative

cell envelope activates a novel cell death pathway. *Proceedings of the*

National Academy of Sciences **113**, E1565-E1574 (2016).

11   Grimm, J. *et al.* The inner membrane protein YhdP modulates the rate of

anterograde phospholipid flow in Escherichia coli. *Proceedings of the*

National Academy of Sciences **117**, 26907-26914 (2020).

12   Sun, J., Rutherford, S. T., Silhavy, T. J. & Huang, K. C. Physical properties

of the bacterial outer membrane. *Nature Reviews Microbiology* **20**, 236-248

(2022).

13   Rojas, E. R. *et al.* The outer membrane is an essential load-bearing element

in Gram-negative bacteria. *Nature* **559**, 617-621 (2018).

14   Mathelié-Guinlet, M., Asmar, A. T., Collet, J.-F. & Dufrêne, Y. F.

Lipoprotein Lpp regulates the mechanical properties of the E. coli cell

envelope. *Nature communications* **11**, 1789 (2020).

15   Park, J. S. *et al.* Mechanism of anchoring of OmpA protein to the cell wall

peptidoglycan of the gram-negative bacterial outer membrane. *The FASEB*

Journal **26**, 219 (2012).

16   Sandoz, K. M. *et al.* β-Barrel proteins tether the outer membrane in many

Gram-negative bacteria. *Nature microbiology* **6**, 19-26 (2021).

17   Deghelt, M. *et al.* Peptidoglycan–outer membrane attachment generates

periplasmic pressure to prevent lysis in Gram-negative bacteria. *Nature*

Microbiology, 1-12 (2025).

18   Haswell, E. S., Phillips, R. & Rees, D. C. Mechanosensitive channels: what

can they do and how do they do it? *Structure* **19**, 1356-1369 (2011).

19   Auer, G. K. *et al.* Mechanical genomics identifies diverse modulators of

bacterial cell stiffness. *Cell systems* **2**, 402-411 (2016).

20   Wang, S., Arellano-Santoyo, H., Combs, P. A. & Shaevitz, J. W. Actin-like

cytoskeleton filaments contribute to cell mechanics in bacteria. *Proceedings*

of the National Academy of Sciences **107**, 9182-9185 (2010).

21   Fletcher, D. A. & Mullins, R. D. Cell mechanics and the cytoskeleton.

Nature **463**, 485-492 (2010).

22   Liu, X., Biboy, J., Consoli, E., Vollmer, W. & den Blaauwen, T. MreC and

MreD balance the interaction between the elongasome proteins PBP2 and

RodA. *PLoS Genetics* **16**, e1009276 (2020).

23   Shiomi, D. *et al.* Mutations in cell elongation genes mreB, mrdA and mrdB

suppress the shape defect of RodZ‐deficient cells. *Molecular microbiology*

87, 1029-1044 (2013).

24   Colavin, A., Shi, H. & Huang, K. C. RodZ modulates geometric

localization of the bacterial actin MreB to regulate cell shape. *Nature*

communications **9**, 1280 (2018).

25   Garner, E. C. *et al.* Coupled, circumferential motions of the cell wall

synthesis machinery and MreB filaments in B. subtilis. *Science* **333**, 222-225

(2011).

26   Domínguez-Escobar, J. *et al.* Processive movement of MreB-associated cell

wall biosynthetic complexes in bacteria. *Science* **333**, 225-228 (2011).

27   Van Teeffelen, S. *et al.* The bacterial actin MreB rotates, and rotation

depends on cell-wall assembly. *Proceedings of the National Academy of*

Sciences **108**, 15822-15827 (2011).

28   Lee, T. K. *et al.* A dynamically assembled cell wall synthesis machinery

buffers cell growth. *Proceedings of the National Academy of Sciences* **111**,

4554-4559 (2014).

29   Garner, E. C. Toward a mechanistic understanding of bacterial rod shape

formation and regulation. *Annual Review of Cell and Developmental Biology*

37, 1-21 (2021).

30   Sun, J., Shi, H. & Huang, K. C. Hyperosmotic Shock Transiently

Accelerates Constriction Rate in Escherichia coli. *Frontiers in Microbiology*

12, 718600 (2021).

31   Eberhardt, C., Kuerschner, L. & Weiss, D. S. Probing the catalytic activity

of a cell division-specific transpeptidase in vivo with β-lactams. *Journal of*

bacteriology **185**, 3726-3734 (2003).

32   Ouzounov, N. *et al.* MreB orientation correlates with cell diameter in

Escherichia coli. *Biophysical Journal* **111**, 1035-1043 (2016).

33   Ursell, T. S. *et al.* Rod-like bacterial shape is maintained by feedback

between cell curvature and cytoskeletal localization. *Proceedings of the*

National Academy of Sciences **111**, E1025-E1034 (2014).

34   Kuru, E. *et al.* In situ probing of newly synthesized peptidoglycan in live

bacteria with fluorescent D‐amino acids. *Angewandte Chemie International*

Edition **51**, 12519-12523 (2012).

35   Cho, H. *et al.* Bacterial cell wall biogenesis is mediated by SEDS and PBP

polymerase families functioning semi-autonomously. *Nature microbiology*

1, 1-8 (2016).

36   Yunck, R., Cho, H. & Bernhardt, T. G. Identification of MltG as a potential

terminase for peptidoglycan polymerization in bacteria. *Molecular*

microbiology **99**, 700-718 (2016).

37   Bohrhunter, J. L., Rohs, P. D., Torres, G., Yunck, R. & Bernhardt, T. G.

MltG activity antagonizes cell wall synthesis by both types of

peptidoglycan polymerases in Escherichia coli. *Molecular Microbiology* **115**,

1170-1180 (2021).

38   Bloom, M., Evans, E. & Mouritsen, O. G. Physical properties of the fluid

lipid-bilayer component of cell membranes: a perspective. *Quarterly*

reviews of biophysics **24**, 293-397 (1991).

39   Hwang, H., Paracini, N., Parks, J. M., Lakey, J. H. & Gumbart, J. C.

Distribution of mechanical stress in the Escherichia coli cell envelope.

Biochimica et Biophysica Acta (BBA)-Biomembranes **1860**, 2566-2575 (2018).

40   Ranjit, D. K. & Young, K. D. The Rcs stress response and accessory

envelope proteins are required for de novo generation of cell shape in

Escherichia coli. *Journal of bacteriology* **195**, 2452-2462 (2013).

41   Mikheyeva, I. V., Sun, J., Huang, K. C. & Silhavy, T. J. Mechanism of outer

membrane destabilization by global reduction of protein content. *Nature*

communications **14**, 5715 (2023).

42   Bratton, B. P., Shaevitz, J. W., Gitai, Z. & Morgenstein, R. M. MreB

polymers and curvature localization are enhanced by RodZ and predict E.

coli's cylindrical uniformity. *Nature communications* **9**, 2797 (2018).

43   Vadia, S. *et al.* Fatty acid availability sets cell envelope capacity and

dictates microbial cell size. *Current Biology* **27**, 1757-1767. e1755 (2017).

44   Shi, H. *et al.* Starvation induces shrinkage of the bacterial cytoplasm.

Proceedings of the National Academy of Sciences **118**, e2104686118 (2021).

45   Wong, F., Garner, E. C. & Amir, A. Mechanics and dynamics of

translocating MreB filaments on curved membranes. *Elife* **8**, e40472 (2019).

46   Miller, S. I. & Salama, N. R. The gram-negative bacterial periplasm: Size

matters. *PLoS biology* **16**, e2004935 (2018).

47   Miguel, A. *et al.* Modulation of bacterial cell size and growth rate via

activation of a cell envelope stress response. *bioRxiv*, 2022.2007.

2026.501648 (2022).

48   Zietek, M. *et al.* Bacterial cell widening alters periplasmic size and

activates envelope stress responses. *BioRxiv*, 2022.2007. 2026.501644 (2022).

49   Al-Mosleh, S., Gopinathan, A., Santangelo, C. D., Huang, K. C. & Rojas, E.

R. Feedback linking cell envelope stiffness, curvature, and synthesis

enables robust rod-shaped bacterial growth. *Proceedings of the National*

Academy of Sciences **119**, e2200728119 (2022).

50   Baba, T. *et al.* Construction of Escherichia coli K‐12 in‐frame, single‐

gene knockout mutants: the Keio collection. *Molecular systems biology* **2**,

2006.0008 (2006).

51   Edelstein, A. D. *et al.* Advanced methods of microscope control using

μManager software. *Journal of biological methods* **1**, e10 (2014).

52   Ursell, T. *et al.* Rapid, precise quantification of bacterial cellular

dimensions across a genomic-scale knockout library. *BMC biology* **15**, 1-15

(2017).

53   Sliusarenko, O., Heinritz, J., Emonet, T. & Jacobs‐Wagner, C. High‐

throughput, subpixel precision analysis of bacterial morphogenesis and

intracellular spatio‐temporal dynamics. *Molecular microbiology* **80**, 612-

627 (2011).

54   Shi, H. *et al.* Deep Phenotypic Mapping of Bacterial Cytoskeletal Mutants

Reveals Physiological Robustness to Cell Size. *Curr Biol* **27**, 3419-3429

e3414 (2017). https://doi.org:10.1016/j.cub.2017.09.065

55   Paradis-Bleau, C. *et al.* Lipoprotein cofactors located in the outer

membrane activate bacterial cell wall polymerases. *Cell* **143**, 1110-1120

(2010).

855