1 **PAK4 promotes vertex remodeling to maintain epithelial integrity and barrier**
2 **function**

3 Babli Adhikary[1], Alexander Chang[1], Atsuko Y. Higashi[2], Hideki Chiba[3], Ann L.
4 Miller[1], Tomohito Higashi[3]*
5
6 [1]Department of Molecular, Cellular, and Developmental Biology (MCDB),
7 University of Michigan, Ann Arbor, MI 48109-1085, USA,
8 [2]Department of Nephrology and Hypertension, and [3]Department of Basic
9 Pathology, Fukushima Medical University, Fukushima 960-1295, Japan.
10
*Address correspondence to:
Tomohito Higashi, Ph.D.
Associate Professor
Department of Basic Pathology, Fukushima Medical University
1 Hikariga-oka, Fukushima, 960-1295, Japan.
tohigash@fmu.ac.jp
+81.24.547.1169

18
19 Short title: PAK4 regulates vertex remodeling
20
21 Abbreviations: AJ, adherens junction; PAK4, p21-activated kinase 4; PBD, p21-
22 binding domain; TJ, tight junction; tAJ, tricellular adherens junction; tTJ, tricellular
23 tight junction; TER, transepithelial electrical resistance; ZnUMBA, Zinc-Based
24 Ultrasensitive Microscopic Barrier Assay.
25
26
Summary

Adhikary et al. show that PAK4, recruited to epithelial junctions by Afadin, drives
dynamic vertex remodeling in cells and embryos. Inhibiting PAK4 disrupts tissue
packing and compromises junctional continuity and barrier function, implicating
PAK4 and Afadin as critical partners in vertex remodeling to preserve tissue
integrity.

Abstract
Cell-cell junctions are essential for epithelial integrity and barrier function, but the
mechanisms regulating their remodeling remain unclear. Here, we investigated the
role of the junctional kinase PAK4 in vertex remodeling. PAK4 localized to apical
junctions and accumulated at multicellular vertices in MCDK cells and *Xenopus*
embryos. Inhibition or knockout of PAK4 increased higher-order vertices, caused
junctional discontinuities, and impaired barrier function in MDCK cells. PAK4
recruitment required the scaffolding protein Afadin. *Afdn*-KO cells exhibited severe
junctional defects and reduced barrier function. Artificial targeting of PAK4 in *Afdn*-
KO cells partially restored junctional continuity and barrier function. In *Xenopus*
embryos, live imaging showed dynamic PAK4 accumulation at remodeling vertices,
and PAK4 inhibition hindered resolution of multicellular vertices. Expression of an
amino-terminal fragment (PAK4-NT) impaired remodeling and induced cytokinetic
failure. Live imaging in *Xenopus* revealed barrier leakage at multicellular vertices
upon PAK4 inhibition. These findings indicate that PAK4 and Afadin cooperate to
maintain epithelial integrity and barrier function by promoting vertex remodeling.

Introduction

Epithelial tissues maintain their architecture and contribute to the homeostasis of multicellular organisms by forming well-ordered sheets of tightly connected epithelial cells. Dynamic rearrangement of epithelial sheets is essential not only during morphogenesis processes in development but also for the maintenance of adult tissues (Guillot and Lecuit, 2013b; Heisenberg and Bellaïche, 2013). In many epithelia, constituent cells are continuously turned over – old cells are removed from the sheet, while new cells are added either through cell division within the sheet or by insertion from outside the sheet. To preserve epithelial integrity and barrier function, the tissue must accommodate cell rearrangements such as intercalation, positional exchange between neighboring cells, and displacement, while maintaining intercellular adhesion and overall junctional organization (Varadarajan et al., 2019; Paré and Zallen, 2020; Mira-Osuna and Le Borgne, 2024; Higashi et al., 2024). Disruption of these processes compromises epithelial barrier integrity, which is critical for preventing uncontrolled paracellular flux and protecting against inflammation and pathogen invasion (Krug et al., 2014; Luissint et al., 2016; Saha et al., 2025).

Cell-cell junctions, composed of tight junctions (TJs), adherens junctions (AJs), and desmosomes, mediate intercellular adhesion in epithelial cells (Farquhar and Palade, 1963). TJs are crucial for maintaining the epithelial barrier (Van Itallie and Anderson, 2014; Otani and Furuse, 2020; Balda and Matter, 2023; Citi et al., 2024), whereas AJs and desmosomes provide mechanical strength (Meng and Takeichi, 2009; Rübsam et al., 2018; Eckert et al., 2025). To accommodate frequent cellular rearrangements, these junctional complexes undergo remodeling in response to epithelial reorganization, cell signaling, and cytoskeletal rearrangements, thereby supporting both tissue renewal and morphogenesis while preserving adhesion. Of particular interest is the remodeling that occurs at cell vertices – the points where multiple cell-cell junctions converge. In this study, we use the term "vertex" to describe such sites and specify them according to the number of participating cells: tricellular vertex (three cells), tetracellular or 4-way vertex (four cells), and rosette (five or more cells). We collectively refer to 4-way vertices and rosettes as higher-order vertices.

At these multicellular vertices of vertebrate epithelial cells, specialized junctional structures are assembled. Among them, tricellular tight junctions (tTJs) are well characterized, with distinct molecular components such as angulin family proteins and tricellulin, and tTJs play a crucial role in sealing the paracellular space of vertices (Ikenouchi et al., 2005; Masuda et al., 2011; Higashi et al., 2013; Sugawara et al., 2021). By contrast, the existence of tricellular adherens junctions (tAJs) has been proposed (Yonemura, 2011) and their components and regulation have recently begun to be understood in *Drosophila* (Lye et al., 2014; Finegan et al., 2019; Letizia et al., 2019; Uechi and Kuranaga, 2019), but the precise molecular composition and architecture of vertebrate tAJs remains less well defined.

Multicellular vertices are dynamically remodeled during tissue rearrangement (Sugimura and Otani, 2024). Several different modes of vertex remodeling have been characterized. One is vertex sliding, in which a tricellular vertex shifts position along a bicellular junctions as it shrinks or elongates (Vanderleest et al., 2018). Another is the T1 transition (Weaire and Rivier, 1984;

Bertet et al., 2004), a topological transition in which two neighboring tricellular
vertices converge as the intervening bicellular junction shortens, transiently forming
a 4-way vertex. This intermediate state then resolves by elongation of a new
junction in the perpendicular direction, giving rise to two new tricellular vertices with
exchanged neighbors. A third example occurs during the final stages of cytokinesis
in epithelial cells. As the cleavage furrow ingresses, a transient 4-way vertex is
formed at the division plane. This intermediate state resolves either by establishing
stable adhesion between the two daughter cells or by allowing the neighboring
cells to insert between the daughters, thereby generating a new neighbor-neighbor
interface (Gibson et al., 2006; Herszterg et al., 2013; Guillot and Lecuit, 2013a;
Founounou et al., 2013; Higashi et al., 2016; Firmino et al., 2016; Landino et al.,
2025). Such cytokinesis-associated remodeling closely resembles the later phase
of a T1 transition and can thus be regarded as a specialized mode of vertex
remodeling in proliferating epithelia. In addition, more complex rearrangements
involve the formation of rosette structures, in which five or more cells converge at
single vertex before being resolved into simpler configurations (Blankenship et al.,
2006). These events are coordinated by adhesion complexes and the actomyosin
cytoskeleton, under the control of Rho-family GTPases and their effectors (Lecuit
and Yap, 2015; Arnold et al., 2017; Perez-Vale and Peifer, 2020; Huebner et al.,
2021; Cavanaugh et al., 2022; Campàs et al., 2024). However, the molecular
mechanisms that ensure barrier integrity during these highly dynamic vertex
remodeling processes remain incompletely understood.
120   PAK4 is a serine/threonine kinase originally identified as a Cdc42 effector
(Abo et al., 1998) and is essential for embryonic viability in mice (Qu et al., 2003).
PAK4 has been implicated in apical junction formation (Wallace et al., 2010) and
cell polarity (Selamat et al., 2015). Although PAK4 is localized at cell-cell junctions
(Hou et al., 2013; Selamat et al., 2015) and its preferential localization at
multicellular vertices has been observed (Baskaran et al., 2021), its roles in
regulating junctional remodeling are not fully understood. Recently, proximity
labeling studies identified Afadin, a scaffold protein essential for junctional actin
organization (Mandai et al., 1997; Ikeda et al., 1999; Sakakibara et al., 2018, 2020;
Kuno et al., 2025; Gong et al., 2025), as a possible binding partner of PAK4
(Baskaran et al., 2021; Choi et al., 2025). Afadin is required for the localization of
PAK4 at cell-cell junctions (Baskaran et al., 2021) and for the maintenance of
epithelial integrity (Choi et al., 2016; Sakakibara et al., 2020). Furthermore, the
Drosophila homolog of Afadin, Canoe (Cno), preferentially localizes to epithelial
vertices and has been linked to their dynamic remodeling events (Sawyer et al.,
2009; Choi et al., 2013; Yu and Zallen, 2020), suggesting that PAK4 may
contribute to the regulation of vertex remodeling in coordination with Afadin.
137   In this study, we investigated the function and dynamics of PAK4 at
138   epithelial vertices using gene-edited MDCK II cells and live imaging of *Xenopus*
139   *laevis* embryos. We further examined the role of Afadin in recruiting PAK4 to
140   junctions and demonstrate that both proteins are critical for maintaining epithelial
141   integrity and barrier function specifically at cell vertices.
142
143

Results
PAK4 preferentially accumulates at higher-order vertices

144
145
To examine the localization of PAK4 in epithelia that undergo dynamic
remodeling *in vivo*, frozen sections of mouse small intestine were stained with an
anti-PAK4 antibody. PAK4 was detected at cell-cell junctions and was particularly
enriched at 4-way (tetracellular) vertices and at some tricellular vertices (Fig. 1A).
Quantitative analysis revealed that PAK4 accumulation at tricellular vertices was
increased compared to the average intensity at bicellular junctions (1.45 ± 0.65-
fold) (Fig. 1B). At 4-way vertices, PAK4 was enriched to approximately twice the
level observed at bicellular junctions (2.23 ± 1.13-fold) (Fig. 1B). The TJ-associated
protein ZO-1 was also enriched at tricellular vertices (1.48 ± 0.65-fold) and 4-way
vertices (1.83 ± 0.54-fold) (Fig. 1C), although its accumulation at 4-way vertices
was generally less prominent than that of PAK4 (Fig. 1A, right panels).
In cultured *cldn2*-KO MDCK II (Ctrl) cells (Saito et al., 2021), PAK4 was
localized at bicellular junctions, as previously reported (Hou et al., 2013; Selamat
et al., 2015; Baskaran et al., 2021) and showed enrichment at 4-way
vertices/rosettes and many tricellular vertices (Fig. 1D). Quantification of
fluorescence intensity revealed that PAK4 accumulated at tricellular vertices to
varying degrees, in most cases ranging from levels similar to those at bicellular
junctions to as much as fivefold higher (2.45 ± 1.20-fold) (Fig. 1E). Additionally,
some tricellular vertices showed marked enrichment (Fig. 1D, arrowheads; Fig. 1E).
Moreover, at 4-way vertices and rosettes (Fig. 1D, arrows), PAK4 was consistently
enriched on average to approximately four times the level observed at bicellular
junctions (4.22 ± 1.48-fold) (Fig. 1E). By comparison, ZO-1 showed relatively
uniform accumulation, reaching about threefold the level of bicellular junctions at
both tricellular (3.11 ± 0.76-fold) and 4-way vertices/rosettes (3.21 ± 0.64-fold) (Fig.
1F).
These results indicate that PAK4 is enriched at multicellular vertices of cell-
cell junctions in mammalian epithelial cells.
173

PAK4 accumulates at remodeling vertices in *Xenopus* gastrula-stage epithelia

174
175
To examine the dynamics of PAK4 at remodeling vertices in a developing
organism, we turned to gastrula-stage *Xenopus laevis* embryos. Immunostaining
with an anti-PAK4 antibody detected PAK4 at cell-cell junctions, and PAK4 was
particularly enriched at higher-order vertices (4-way vertices and rosettes) as well
as at some tricellular vertices (Fig. 2A). We then live imaged PAK4-mNeon with TJ
(TagBFP-ZO-1)- and AJ (PLEKHA7-mCherry)-localizing proteins (Fig. 2B).
Quantitative analysis of live images revealed that PAK4 accumulation at 4-way
vertices and rosettes (2.74 ± 0.21) was significantly higher than at tricellular
vertices (1.64 ± 0.11) (Fig. 2C). ZO-1 was also enriched at 4-way vertices and
rosettes (3.47 ± 0.25) compared to tricellular vertices (1.84 ± 0.13) (Fig. 2D). To
examine the lateral localization of PAK4, we quantified the relative fluorescence
intensity along the *z*-axis (Fig. 2E) (Higashi et al., 2019). The PAK4-mNeon peak
closely overlapped with the ZO-1 peak, while it was slightly offset from the
PLEKHA7 peak, indicating that PAK4 co-localizes with apical junctional proteins
(Fig. 2E).

To understand dynamic changes in PAK4 localization during junction
remodeling, we performed live imaging of PAK4-mNeon at remodeling and non-
remodeling vertices in the gastrula-stage *Xenopus* epithelium. We defined a
"remodeling vertex" as a tricellular vertex that has an acute angle to start off, and
then actively remodels over several minutes, resulting in a vertex angle greater
than 90⁰. In contrast, we defined a "non-remodeling vertex" as a tricellular vertex
where the starting angle is >90⁰, and the change in angle is ± 20⁰. PAK4 appeared
to be enriched at actively remodeling vertices, whereas PAK4 intensity remained
more stable at non-remodeling vertices (Fig. 2F and Video 1). Indeed,
quantification showed that early in vertex remodeling, PAK4 intensity was similar at
remodeling vs. non-remodeling vertices. Then, midway through vertex remodeling
PAK4 at remodeling vertices was significantly enriched (1.56 ± 0.07-fold) over
PAK4 at non-remodeling vertices (Fig. 2G). PAK4 intensity at remodeling vertices
then returned to baseline levels as remodeling completed (Fig. 2G).
These results indicate that PAK4 is enriched at multicellular vertices in
developing epithelial tissue, and PAK4 especially accumulates at remodeling
tricellular vertices.
208
Inhibition of PAK4 disrupts vertex geometry in MDCK II epithelial sheets
To assess the function of PAK4, Ctrl MDCK II cells were treated with
different concentrations of the ATP-competitive PAK4 inhibitor, PF-3758309.
Treatment with PAK4 inhibitor led to an increased number of higher-order vertices
compared to DMSO-treated cells (Fig. 3A). We performed three types of
quantitative analyses to evaluate this effect. First, the ratio of the total number of
cell vertices to the total number of cells in the monolayer was calculated. In an
ideal hexagonally packed epithelial sheet, each vertex is formed where three cells
meet, and each cell contributes to six such vertices. Given this geometry, the total
number of vertices divided by the total number of cells is expected to be 2. When
cells deviate from hexagonal packing and higher-order junctions (involving four or
more cells) are present, this ratio decreases. For example, in tightly packed arrays
of rectangular and triangular cells, the vertex-to-cell ratios become 1 and 0.5,
respectively. Discontinuities in cell-cell junctions can also lead to a further
reduction in the number of detectable vertices. In this analysis, monolayers treated
with DMSO or 500 nM PAK4 inhibitor exhibited vertex-to-cell ratios close to 2 (1.82
± 0.05 [DMSO] and 1.80 ± 0.09 [500 nM PAK4 inhibitor]) (Fig. 3B). In contrast,
treatment with 5 µM PAK4 inhibitor resulted in a significant reduction in this ratio
(1.71 ± 0.08) (Fig. 3B), suggesting a disturbance in normal junctional organization.
Second, all vertices were classified according to the number of converging cells,
and the proportions of 4-way vertices and rosettes were quantified. At 500 nM
PAK4 inhibitor, a significant increase in 4-way vertices was observed, and at 5 µM,
both 4-way vertices and rosettes were significantly increased (Fig. 3C). Third, the
number of discontinuities in cell-cell junctions was quantified. A slight increase was
noted in the 5 µM PAK4 inhibitor-treated cells, although the change was not
statistically significant (Fig. 3D).
These results suggest that kinase activity of PAK4 is required for the
maintenance of tricellular vertices as well as for the resolution of 4-way vertices
and rosettes.
238

PAK4 inhibition hinders resolution of higher-order vertices in *Xenopus* epithelium

To assess the function of PAK4 during vertex remodeling in development, *Xenopus* embryos expressing TagBFP2-ZO-1, Angulin-1-3xGFP, and the F-actin probe LifeAct-miRFP were treated with the PAK4 inhibitor, PF-3758309. PAK4 inhibition led to abnormal clumpy signal around multicellular vertices for ZO-1, Angulin-1, and F-actin compared with control DMSO-treated embryos (Fig. 4A). To investigate the resolution of multicellular vertices in developing embryos, we examined tissue packing following cytokinesis. Daughter cells in control (DMSO-treated) and PAK4 inhibitor-treated embryos were categorized based on the geometry of the newly formed interface: daughter-daughter interface (Type I), 4-way vertex (Type II), or neighbor-neighbor interface (Type III) (Fig. 4B) (Gibson et al., 2006; Higashi et al., 2016). Control embryos displayed a 4-way vertex (Type II, 100%) immediately after cytokinesis, and after 60 minutes, some of these rearranged into daughter-daughter interfaces (Type I, 68.4%), while others rearranged into neighbor-neighbor interfaces (Type III, 10.5%) (Fig. 4C-D). PAK4 inhibitor-treated embryos also showed a high percentage of 4-way vertices (Type II, 95.7%) immediately after cytokinesis, but the majority of these failed to rearrange over time; 69.6% remained as 4-way vertices 60 minutes post cytokinesis, while 13.0% rearranged into daughter-daughter interfaces (Type I), and 17.4% rearranged into neighbor-neighbor interfaces (Type III) (Fig. 4C-D). Therefore, PAK4 inhibition hinders 4-way vertex resolution following cytokinesis, affecting cell packing patterns in the developing *Xenopus* epithelium.

Next, we examined whether PAK4 inhibition causes defects in rosette resolution in developing *Xenopus* epithelia. Rosette resolution was defined as the rearrangement of a multicellular rosette – where five or more cells meet – into a simpler configuration, in which newly adjoining neighbors extend their bicellular junctions and generate tricellular or 4-way vertices. PAK4 inhibition causes a significant delay in rosette resolution compared to DMSO-treated control embryos (Fig. 4E and Video 2). In control embryos, the majority of rosettes resolve within 30 minutes (66.67%), with only a small percentage (6.67%) not resolved after 60 minutes of imaging (Fig. 4F). In contrast, in PAK4 inhibitor-treated embryos, the majority of rosettes (61.54%) were not resolved after 60 minutes of observation (Fig. 4F). In the population of rosettes that were resolved, the average time for resolution was significantly higher in PAK4 inhibitor-treated embryos (median = 35.83 min) compared to controls (median = 17.13 min) (Fig. 4G). A limitation of this rosette resolution analysis is that we quantified rosettes from the start of imaging rather than from the time of their actual formation, which leads to an underestimation of resolution times. However, this systematic bias was applied equally to both control and PAK4 inhibitor-treated embryos.

Taken together, these results suggest that PAK4 kinase activity is required for proper organization of junctional proteins and F-actin at multicellular vertices and plays a crucial role in resolution of 4-way vertices and rosettes in developing *Xenopus* epithelia.

PAK4-knockout epithelial sheets exhibit discontinuities in cell-cell junctions and an increased number of multicellular vertices

To further examine the function of PAK4, we established two independent
PAK4-knockout (KO) MDCK II cell clones (*PAK4*-KO#1 and *PAK4*-KO#2) with
CRISPR/Cas9-based genome editing (Fig. S1A-C) (Saito et al., 2024) using Ctrl
cells as a parental cell line. Immunostaining and immunoblotting analyses
confirmed complete loss of PAK4 signal in both clones (Fig. S1G-H).
Using these *PAK4*-KO cell clones, we examined the roles of PAK4 at cell-
cell junctions and vertices. When cultured on a glass surface, Ctrl cells formed a
well-ordered monolayer by 40 hours after plating. In contrast, *PAK4*-KO cells
formed disorganized sheets, with both clones showing a significant increase in the
number of discontinuities in ZO-1 signal (Fig. 5A). In *PAK4*-KO cells, the vertex-to-
cell ratio was significantly decreased to less than 1.0 (Fig. 5B), and the proportion
of 4-way vertices and rosettes among all cell vertices was increased compared to
Ctrl cells, although the increase reached statistical significance only in one clone
(*PAK4*-KO#1) (Fig. 5C). Additionally, the number of junction discontinuities was
around 0.5 per cell, which was much higher than the Ctrl (Fig. 5D). Stable
expression of Flag-tagged wild-type PAK4 in the *PAK4*-KO#1 clone restored both
vertex organization and junctional continuity (Fig. S2A-E). In contrast, expression
of a kinase-dead mutant (Flag-PAK4 K351R), which is equivalent to a K299R
mutation in PAK1 (Zhang et al., 1995; Sells et al., 1997), failed to rescue the
discontinuity phenotype (Fig. S2A-E), suggesting that kinase activity is required for
maintaining epithelial integrity. Vertex organization was only partially restored by
Flag-PAK4 K351R, implicating a kinase activity-independent contribution of PAK4
to vertex patterning.
These results with *PAK4*-KO cells complement our experiments with the
PAK4 inhibitor, further supporting the role of PAK4 in maintaining junctional
integrity in epithelial cells.
312
PAK4 knockout results in formation of sarcomere-like MyoIIB structures
along cell-cell junctions
We noticed that the cell-cell junctions of *PAK4*-KO MDCK II cells cultured on
glass coverslips appear straighter than Ctrl cells, and this effect was more obvious
when the cells were cultured on polycarbonate filters (Fig. S3D). Since the shape
of cell-cell junctions is determined by the actomyosin cytoskeleton, we next
examined the Myosin II organization in *PAK4*-KO cells. Whereas Ctrl cells exhibit
faint staining for MyoIIB along cell-cell junctions, *PAK4*-KO cells show dotted
MyoIIB signal flanking both sides of cell-cell junctions (Fig. 5E), which closely
resembles the sarcomere-like structure reported in inner ear cells (Ebrahim et al.,
2013), ZO-1/2-double-deficient MDCK II cells (Yamazaki et al., 2008; Fanning et al.,
2012; Choi et al., 2016), and *claudin/JAM-A*-KO MDCK II cells (Nguyen et al.,
2024). MyoIIA also exhibited a punctate pattern at both sides of *PAK4*-KO cell-cell
junctions (Fig. S3A) similar to MyoIIB (Fig. S3B). To examine whether junctional
Myosin is activated, we stained for S18/T19-double phosphorylated myosin light
chain (2P-MLC). 2P-MLC was localized at cell-cell junctions around the tricellular
vertices in *PAK4*-KO cells (Fig. S3C), suggesting that the junctional tension is
increased. Vinculin, an actin-binding protein recruited to AJs under tension
(Yonemura et al., 2010; le Duc et al., 2010; Huang et al., 2017; Higashi et al.,
2016; van den Goor et al., 2024), also accumulated at the tricellular region of

PAK4-KO cells (Fig. S3D), while other TJ and AJ proteins did not exhibit notable
changes in localization at junctions in *PAK4*-KO cells (Fig. S3E-H).
These results suggest that PAK4 suppresses the formation of a high
tension actomyosin array flanking cell-cell junctions, thus regulating the linearity of
cell-cell junctions.
338
Afadin is required for the junctional localization of PAK4 and maintenance of
epithelial integrity
It was reported that Afadin is responsible for localization of PAK4 at cell-cell
junctions (Baskaran et al., 2021). To assess Afadin's role in PAK4-driven vertex
remodeling, we established Afadin knockout cell clones using CRISPR/Cas9
genome editing (*Afdn*-KO#1 and *Afdn*-KO#2) (Fig S1D-H) using Ctrl cells as a
parental cell line. PAK4 was no longer localized at cell-cell junctions in *Afdn*-KO
cells (Fig. S1G), consistent with the previous report (Baskaran et al., 2021).
However, Afadin was still localized at cell-cell junctions and cell vertices of *PAK4*-
KO cells (Fig. S1G), suggesting that Afadin recruits PAK4 to cell-cell junctions. It
was reported that knockdown of Afadin in ZO-1/ZO-2-double knockdown MDCK II
cells resulted in discontinuities at cell-cell junctions (Choi et al., 2016). To assess
whether Afadin (independent of ZO-1/ZO-2 double knockdown) is required for
maintenance of epithelial integrity, *Afdn*-KO cells were cultured on glass coverslips.
Similar to *PAK4*-KO cells, both clones of *Afdn*-KO cells exhibited discontinuities in
cell-cell junctions and an increased frequency of 4-way junctions (Fig. 6A-D).
These phenotypes were largely rescued by exogenous expression of FLAG-Afadin
in the *Afdn*-KO#1 cell clone (Fig. S2F-J).
These results suggest that Afadin is required for the recruitment of PAK4 to
cell-cell junctions and the maintenance of epithelial integrity and vertex
organization.
360
Artificial targeting of PAK4 to adherens junctions partially rescues Afadin
knockout phenotypes
Afadin is known for its role as an actin-binding protein that regulates the
actin cytoskeleton at cell-cell junctions (Mandai et al., 1997). Additionally, we and
others have shown that Afadin is required for PAK4 localization at cell-cell
junctions (Fig. S1G) (Baskaran et al., 2021). Thus, we next examined how Afadin's
ability to recruit PAK4 to cell-cell junctions affects the maintenance of epithelial
integrity. To test this, we assessed whether artificial recruitment of PAK4 to
junctions is sufficient to maintain epithelial integrity in the absence of Afadin. We
established an *Afdn*-KO cell clone stably expressing PAK4 conjugated with Nectin2,
an Afadin-binding AJ adhesion protein (Takahashi et al., 1999). The Nectin2-
FLAG-PAK4 construct was localized at cell-cell boundaries (Fig. 6E), but its
localization was not restricted to AJs. Remarkably, the epithelial integrity defects of
Afdn-KO cells were partially rescued in Nectin2-FLAG-PAK4-expressing cells (Fig.
6F-I). The vertex-to-cell ratio was significantly increased to around 1.0 (Fig. 6G),
and the frequency of 4-way junctions was significantly decreased compared to the
Afdn-KO cells (Fig. 6H). In addition, the frequency of junctional discontinuities was
significantly reduced (Fig. 6I). Notably, the kinase-dead version of PAK4
conjugated with Nectin2 failed to rescue the phenotypes (Fig. 6F-I).

Together, these data suggest that junctional localization of PAK4 is
important for the maintenance of epithelial integrity and that one of Afadin's key
roles is to recruit PAK4 to cell-cell junctions and cell vertices.
383
Afadin interacts with the N-terminal region of PAK4
To gain insights into the binding interface between PAK4 and Afadin, we
used AlphaFold3 (Abramson et al., 2024) to predict the 3D structure of the PAK4-
Afadin complex (Fig. S4A). Although a large part of the PAK4 and Afadin
polypeptides appeared unstructured, the amino-terminal region of PAK4 near the
p21-binding domain (PBD) (PAK4-NT; aa 68-89) and the amino-terminal part of
Afadin including the Ras association 1 (RA1) domain exhibited low predicted
aligned error (PAE) scores (Fig. S4B), suggesting that these regions may interact
directly. Further, the AlphaBridge algorithm (Álvarez-Salmoral et al., 2024)
suggested possible interaction of these regions (Fig. S4C,F). The interaction of the
predicted Afadin-interacting region of PAK4 with Afadin appears to affect the
conformation of the adjacent PAK4 autoinhibitory domain (AID), which is
associated with the active site groove of the kinase domain in the inactive state
(Fig. S4B,D-E). Interestingly, the amino-acids involved in the interaction in the
Afadin-PAK4 complex model are at least partially conserved among the group 2
PAK family including PAK4, PAK5, PAK6, and *Drosophila* mbt, but are not
conserved in the group 1 PAK family including PAK1, PAK2, and PAK3 (Fig. S4G).
To examine whether the identified domains are responsible for interaction
of PAK4 and Afadin, HEK293T cells were transfected with a GFP-tagged RA1
fragment of Afadin (GFP-Afdn-RA1) and mCherry-tagged full-length PAK4 (mChe-
PAK4), and GFP-Afdn-RA1 was immunoprecipitated. A small amount of mChe-
PAK4 was co-immunoprecipitated (Fig. S4H), suggesting that the Afadin RA1
domain can interact with PAK4. FLAG-tagged full-length Afadin could also be co-
immunoprecipitated with GFP-tagged PAK4-NT (Fig. S4H).
These results implicate that PAK4 and Afadin may interact through the N-
terminal region of PAK4 and the RA1 domain of Afadin.
410
Inhibiting PAK4 recruitment to junctions prevents vertex remodeling and
causes cytokinetic failure in the *Xenopus* epithelium
Based on the finding that PAK4-NT is responsible for interaction with
Afadin, we examined whether PAK4-NT competes with endogenous PAK4 for
binding Afadin and localizing to AJs. We first established a Ctrl MDCK II cell clone
stably expressing GFP-PAK4-NT. In these cells, GFP-PAK4-NT accumulated at
multicellular vertices as well as cell boundaries (Fig. S4I). Mixed culture of these
cells with Ctrl cells showed that the endogenous full-length PAK4 was no longer
localized at cell-cell boundaries in the GFP-PAK4-NT-expressing cells. This
suggests that GFP-PAK4-NT can be used as a tool to interfere the localization of
endogenous PAK4 at AJs (Fig. S4J), although we cannot exclude the possibility
that PAK4-NT prevents PAK4 localization through different mechanisms, such as
preventing interaction with active Cdc42 or other binding partners.
We then tested the effects of expressing PAK4-NT (Fig. 7A) on vertex
remodeling in the *Xenopus* epithelium. Inhibiting the recruitment of endogenous
PAK4 to junctions led to an increased number of higher-order vertices (Fig. 7B).
We quantified the ratio of the number of tricellular vertices, 4-way vertices, or

rosettes to the total number of cell vertices in control and PAK4-NT embryos.
Notably, PAK4-NT expression resulted in a significant increase in ratio of 4-way
vertices compared to control (Fig. 7C). Likewise, a significant increase in rosettes
was observed in PAK4-NT embryos (Fig. 7C). We conclude that inhibiting
endogenous PAK4 recruitment to junctions via expression of the PAK4-NT
fragment prevents multicellular vertex remodeling, leading to an increased
proportion of higher-order vertices in the developing *Xenopus* epithelium.
Live imaging of PAK4-NT-expressing embryos revealed that inhibiting
PAK4 junctional localization also caused defects in epithelial cytokinesis. First, the
majority of PAK4-NT cell divisions formed a 4-way vertex. In contrast to controls,
where these 4-way vertices remodel into daughter-daughter or neighbor-neighbor
interfaces (Fig. 4D), the neighboring cells in PAK4-NT embryos often retracted,
leading to re-fusion of the daughter cells and ultimately failed cytokinesis (Fig. 7D
and Video 3). Even in cases where cytokinesis was able to complete, other defects
like asymmetric cell division were observed (Fig. 7D). Quantification of cytokinesis
completion showed that in PAK4-NT-expressing embryos, 46.88% of cytokinesis
events failed, whereas only 2.17% failed in controls (Fig. 7E). Additionally,
inhibiting PAK4 recruitment to junctions via expression of PAK4-NT also caused
junctional discontinuities (Fig. 7F and Video 3).
Taken together, we show that inhibition of endogenous PAK4 recruitment to
junctions causes failure in multicellular vertex remodeling and cytokinesis, as well
as junctional discontinuities in the *Xenopus* epithelium.
450
PAK4 and Afadin are required for establishment of the epithelial barrier
Loss of epithelial integrity often results in impaired barrier function. To
investigate the roles of PAK4 and Afadin in epithelial barrier maintenance, we
measured transepithelial electrical resistance (TER) of *PAK4*-KO and *Afdn*-KO
MDCK II cells cultured on transwell filters. Since the Ctrl cells are *Cldn2*-KO MDCK
II cells, in which cation-permeable *claudin-2* was knocked out, the TER of Ctrl cells
reaches 3.9-4.1 $k\Omega\bullet cm^2$ on day 7 after seeding (Fig. 8A). In contrast, *PAK4*-KO
cells exhibited impaired barrier function (1.5-2.0 $k\Omega\bullet cm^2$) on day 7 (Fig. 8B),
indicating that PAK4 is required for establishment of a tight barrier. Furthermore,
the TER values of *Afdn*-KO cells remain even lower (0.06-0.21 $\Omega\bullet cm^2$) than those
of *PAK4*-KO cells (Fig. 8C), suggesting that Afadin contributes to the establishment
of epithelial barrier not only by recruiting PAK4 to the junctions and vertices, but
also through other molecular mechanism(s).
To examine whether PAK4 and Afadin function in a cell-autonomous
manner, we cultured Ctrl and KO cells together and evaluated the barrier function
of mixed cultures. To distinguish Ctrl and KO cells, we used a cell clone of Ctrl
cells stably expressing mCherry conjugated with nuclear localization signals (NLS)
(Ctrl mChe-NLS). When two cell clones with different barrier properties are mixed
together and each cell clone does not affect the barrier function of the other clone,
the combined TER can be predicted by harmonic mean of TER values of the two
cell clones cultured separately ($R_{pred} = 2 / ((1 / R_{clone1}) + (1 / R_{clone2}))$). For example,
the *angulin-1*-knockout cell clone (*Ang1*-KO) has leaky barrier properties (0.22-
0.25 $k\Omega\bullet cm^2$) (Fig. 8A). The TER of a mixed culture of Ctrl mChe-NLS and *Ang1*-
KO cells on day 7 was 0.31-0.35 $k\Omega\bullet cm^2$, which was close to the predicted value
(0.44 $k\Omega\bullet cm^2$) (Fig. 8D). In contrast, when *PAK4*-KO cells were mixed with Ctrl

mChe-NLS cells, the TER increased much higher (3.5-3.9 kΩ•cm^2) than the
predicted values (2.2 kΩ•cm^2 [#1] and 2.6 kΩ•cm^2 [#2]) and reached a level almost
comparable with Ctrl cells (Fig. 8E). The discrepancy was more evident in a mixed
culture of Ctrl mChe-NLS and *Afdn*-KO cells (measured, 3.5-3.9 kΩ•cm^2; predicted,
0.14 kΩ•cm^2 [#1] and 0.37 kΩ•cm^2 [#2]). Again, the TER of a mixed culture of Ctrl
mChe-NLS and *Afdn*-KO cells was similar to that of Ctrl cells (Fig. 8F). We
confirmed that the ratio of *PAK4*-KO or *Afdn*-KO cells to Ctrl mChe-NLS cells
remained 1:1 throughout the experimental period (Fig. S3I), indicating that KO cells
are not eliminated by mChe-NLS cells. Of interest, the sarcomere-like structure of
MyoIIB in the *PAK4*-KO and *Afdn*-KO cells was not observed in the KO cells within
the mixed culture (Fig. 8G-H), suggesting that neighboring Ctrl cells may convey a
mechanical or chemical signal to *PAK4*-KO or *Afdn*-KO cells and suppress the
formation of the sarcomere-like MyoII structure. To examine how far the
suppressive effect of Ctrl mChe-NLS cells on myosin assembly extends into the
neighboring *PAK4*-KO clone, we co-cultured Ctrl mChe-NLS and *PAK4*-KO cells
side by side and analyzed the boundary region. The suppression of the sarcomere-
like structure of MyoIIB was restricted to the first two layers of PAK4-KO cells
adjacent to Ctrl mChe-NLS cells (Fig. 8I).
Since artificial targeting of PAK4 to junctions using the Nectin2-FLAG-PAK4
construct partially rescued the epithelial integrity defects, we speculated that
artificial PAK4 targeting could also improve the barrier properties of *Afdn*-KO cells.
Expressing Nectin2 alone in *Afdn*-KO cells did not affect barrier function (Fig. 8J).
As predicted, *Afdn*-KO cells stably expressing Nectin2-FLAG-PAK4 exhibited much
higher TER values compared to the parental *Afdn*-KO cell clone (Fig. 8K), although
the barrier function was not fully rescued to the Ctrl level. The kinase-dead version
(Nectin2-FLAG-PAK4 K351R) failed to rescue the barrier defect (Fig. 8K),
indicating that kinase activity of PAK4 is important to support barrier function.
These data suggest that PAK4 recruitment by Afadin is important for the
establishment of the epithelial barrier. They also suggest that Ctrl NLS-mChe cells
affect the cell structure and mechanical properties of neighboring *PAK4*-KO and
Afdn-KO cells and can rescue the barrier defects of KO cells.
507
PAK4 inhibition disrupts epithelial barrier function in *Xenopus* embryos
In PAK4 inhibitor-treated embryos, small openings were regularly observed
at multicellular rosettes (Fig. S5A). Thus, we wanted to investigate the barrier
function of the *Xenopus* epithelium upon PAK4 inhibition. To assess changes in
barrier function, we used the Zinc-Based Ultrasensitive Microscopic Barrier Assay
(ZnUMBA) to visualize dynamic barrier leaks, which appear as increases in the
fluorescence intensity of a zinc ion indicator, FluoZin3 (Stephenson et al., 2019;
Higashi et al., 2023b) (Fig. 9A). Whole-field and junctional FluoZin3 intensity were
not significantly different in control vs. PAK4 inhibitor-treated embryos (Fig. S5B-C),
indicating that overall barrier function is not severely affected. In controls, there
were occasional local barrier leaks (increases FluoZin3 signal) (Fig. 9B). In
contrast, PAK4 inhibitor-treated embryos exhibited increased leaks – particularly at
vertices (Fig. 9B and Video 4). Indeed, PAK4 inhibitor-treated embryos had a
significantly higher percentage of barrier leaks at vertices (32.74±4.51%) compared
to controls (16.83±1.84%) (Fig. 9C). Additionally, PAK4 inhibitor-treated embryos
showed a significant increase in FluoZin3 intensity at higher-order vertices (4-way

or higher) compared to controls (Fig. 9D). Of note, the barrier leaks observed in
controls were quickly repaired, and FluoZin3 signal returned to baseline (Fig. 9E).
In contrast, in PAK4 inhibitor-treated embryos, repeated or persistent leaks were
observed at some vertices (Fig. 9E).
Next, we wanted to test whether inhibiting endogenous PAK4 recruitment to
junctions via expressing PAK4-NT would disrupt epithelial barrier function. In
control embryos, baseline levels of FluoZin3 signal at cell-cell junctions were low,
with occasional local barrier leaks that were rapidly repaired, indicating that barrier
function is intact (Fig. 10A). In contrast, PAK4-NT-expressing embryos exhibited
striking barrier defects. PAK4-NT embryos had a higher number of leaks
specifically localized to the multicellular vertices (Fig. 10A). In 50% of PAK4-NT-
expressing embryos, the whole-field intensity of FluoZin-3 increased over time (Fig.
10A-B), indicating global barrier disruption (Video 5). Significant differences
between control and PAK4-NT-expressing embryos were observed for both whole-
field and junctional FluoZin3 intensity (Fig. 10C-D). Additionally, PAK4-NT-
expressing embryos exhibited a significantly higher percentage of leaks at vertices
(63.85±9.23%) compared to controls (16.83±1.84%) (Fig. 10E). Quantification of
the intensity of FluoZin3 signal at higher-order vertices (4-way or higher) showed
that PAK4-NT-expressing embryos exhibited a significant increase in FluoZin3
intensity at higher-order vertices compared to controls (Fig. 10F). The barrier leaks
observed in controls were rapidly repaired (Fig. 10G). However, PAK4 N-term-
expressing embryos, often failed to effectively repair their leaks, and persistent or
repeated leaks were common (Fig. 10G). The global barrier disruption observed in
these embryos could be a result of the failure to repair local leaks.
Taken together, these results suggest that PAK4 is required for maintaining
epithelial barrier function *in vivo*. Pharmacological inhibition of PAK4 or expression
of PAK4-NT causes increased leaks – both in number and severity – at
multicellular vertices, challenging the barrier integrity of the developing epithelium.
552
553

Discussion

Vertex remodeling plays a fundamental role in epithelial tissue organization and dynamics

Vertex remodeling drives directional cell rearrangements, contributing to processes like convergent extension during development in multicellular organisms. In adult tissues like the intestine, similar vertex remodeling events are believed to occur during the steady-state renewal of epithelial sheets, where cells derived from stem cell niches migrate across the epithelial plane. In both contexts, vertex remodeling must meet two seemingly opposing requirements: 1) maintaining epithelial integrity and barrier function at tTJs and 2) enabling flexible reorganization of cell neighbors. Our study demonstrates that PAK4 helps fulfill both of these roles during passive vertex remodeling events, as shown by gene knockouts in cultured MCDK II epithelial monolayers and live imaging of the *Xenopus* embryonic epithelium. However, whether PAK4 is also required for active, developmentally-driven remodeling during convergent extension remains an open question. Of interest, improper folding of the caudal portion of the neural tube in PAK4-null mouse embryos (Qu et al., 2003) suggests a role for PAK4 in convergent extension.

PAK4 localizes to remodeling vertices and is involved in the resolution of higher-order vertices

Vertex remodeling at higher-order vertices can be divided into three stages: contraction of bicellular junctions leading to vertex convergence, formation of multicellular vertices in which four or more cells share a vertex, and resolution of such multicellular vertices into new configuration of lower-order vertices. We observed that PAK4 preferentially localizes to vertices undergoing remodeling, particularly to 4-way vertices and rosettes, and that PAK4 knockout or inhibition leads to a significant accumulation of unresolved rosettes. These findings suggest that PAK4 is primarily involved in the resolution phase. While Afadin is required for the recruitment of PAK4 to junctions, the vertex- and rosette-specific enrichment of PAK4 is more prominent than that of Afadin, implying that additional signals are needed. Candidates include local activation of Cdc42 (Oda et al., 2014), phosphorylation of Afadin (Yu and Zallen, 2020), and liquid-liquid phase separation (LLPS)-mediated condensate formation by Afadin (Kuno et al., 2025). Previous studies have identified additional regulators such as Sidekick (Lye et al., 2014; Finegan et al., 2019; Letizia et al., 2019; Uechi and Kuranaga, 2019; Malin et al., 2022), M6 (Wittek et al., 2020; Esmangart de Bournonville and Le Borgne, 2020; Ikawa et al., 2023), PTEN (Bardet et al., 2013; Malin et al., 2024), Ajuba (Razzell et al., 2018; Ikawa et al., 2023), and Vinculin (van den Goor et al., 2024; Landino et al., 2025) as modulators of vertex dynamics. Interestingly, PAK4 was reported to interact with Vinculin in the context of podosome formation in myeloid cells (Foxall et al., 2019), suggesting a possible link between mechanical remodeling and actin-cytoskeleton regulation. Recently, it was reported that the *Drosophila* homolog of PAK4, mbt, regulates Sidekick localization at tAJs, raising a possibility that PAK4 regulates the structure of tAJs through Sidekick (Gandhi et al., 2025), although it is still unclear whether the vertebrate homologs of Sidekick, Sdk1 and Sdk2, are involved in tAJ architecture and function. Whether such interactions also cooperate

601 at vertebrate epithelial vertices, and how they contribute to rosette resolution and
602 junctional rearrangement, are important topics for future investigation.
603
PAK4's kinase activity is essential for its function in vertex remodeling
Vertex remodeling was impaired by pharmacological inhibition of PAK4
(Figs. 3 and 4) as well as by genetic ablation of PAK4 (Fig. 5), and this phenotype
was not rescued by expression of a kinase-dead mutant (Fig. S2), highlighting the
importance of its kinase activity in this process. However, the direct
phosphorylation targets of PAK4 in this context remain unknown. It was reported
that U2OS cells treated with a PAK4 inhibitor did not exhibit changes in Afadin
phosphorylation, but several candidate proteins proximal to PAK4 were identified,
including cingulin, p120-catenin, DLG5, Scrib, ARHGEF11 (also known as PDZ-
Rho GEF), and ZO-1/2 (Baskaran et al., 2021). Some of these proteins may act
downstream of PAK4 to regulate membrane plasticity, cell polarity, and
cytoskeletal reorganization during rosette resolution. Notably, ARHGEF11 binds to
PAK4 (Barac et al., 2004) and has been reported as a phosphorylation target of
PAK4 in invadosome regulation in cancer cells (Nicholas et al., 2016), raising the
possibility of shared pathways in epithelial and invasive structures. PAK4 is known
to undergo autophosphorylation at K350 (K351 in dog PAK4) within the activation
loop of the kinase domain, and this modification is required for its activation (Abo et
al., 1998). Although this site is generally considered constitutively phosphorylated
(Callow et al., 2002), long-term kinase inhibition or a loss-of-function mutation may
disrupt this phosphorylation and thereby alter the conformation of PAK4. In this
scenario, PAK4 may act not only as a kinase but also as a structural scaffold at
remodeling vertices, influencing junctional and cytoskeletal architecture.
Supporting this view, we observed repositioning of PAK4-localizing vertices during
cell rearrangements (Fig. 2F-G), suggesting a transient shift in junctional material
properties from solid-like to liquid-like states. LLPS-prone proteins, such as Afadin
(Kuno et al., 2025), are promising candidates for mediating such transitions.
630
Epithelial integrity maintenance by Afadin and recruitment of PAK4
Afadin has been recognized as a key component in stabilizing actin
filaments at AJs through its interactions with nectins and α-catenin (Mandai et al.,
1997; Takahashi et al., 1999; Tachibana et al., 2000; Pokutta et al., 2002;
Sakakibara et al., 2020), a mechanism that is central to the maintenance of
epithelial integrity. Building on this framework, our study uncovers an additional
function: Afadin recruits PAK4 to cell-cell junctions where it is important for
maintaining epithelial integrity. This is evidenced by the findings that PAK4 and
Afadin interact (Fig. S4), and artificial targeting of PAK4 to cell-cell junctions in
Afdn-KO cells partially rescued epithelial integrity defects (Fig. 6E-I). The
incomplete rescue strongly suggests that Afadin contributes to epithelial stability
through mechanisms beyond PAK4 recruitment, in line with its canonical role in
actin anchoring (Mandai et al., 1997; Sawyer et al., 2009; Sakakibara et al., 2020;
Gong et al., 2025; Jensen et al., 2025). A recent report further refines this view by
indicating that mature AJs (named *zonula adherens matura*) can be partitioned into
two belt-like zones: and apical Afadin/nectin-populated zone and a basal
cadherin/catenin-populated zone, with F-actin preferentially tethered at the
Afadin/nectin zone (Mangeol et al., 2024). The configuration thought to be

important for assembling AJs capable of withstanding mechanical tension. At the same time, Afadin is also reported to cooperate with α-catenin and Vinculin to stabilize actin bundles (Gong et al., 2025), implying crosstalk between Afadin/nectin- and cadherin/catenin-based zones. How Afadin-recruited PAK4 interfaces with this zonal architecture and with classical actin regulators remain unclear, although our data suggest that PAK4 itself localizes to the Afadin/nectin-based zone (Fig. 2E). Resolving these relationships will be an important direction for future work.

The N-termini of both Afadin and its *Drosophila* homolog Cno harbor tandem RA domains (RA1 and RA2), which have been regarded as modules for binding Ras family GTPases (Kuriyama et al., 1996; Boettner et al., 2000) such as Rap1. This model has explained the genetic interaction and overlapping functions of Afadin/Cno and Rap1 (Boettner et al., 2003; Sawyer et al., 2009; Choi et al., 2013). However, a recent study revealed that the sequence of RA1, but not that of RA2, is essential for epithelial integrity (McParland et al., 2024). This difference is puzzling given that both domains are capable of binding Rap1, albeit with different affinities (Linnemann et al., 1999; Wohlgemuth et al., 2005; McParland et al., 2024). Our structural predictions suggesting that RA1 also provides a platform for recruitment of PAK4 (or its *Drosophila* homolog mbt) would explain the unique requirement of the RA1 domain.

The role of epithelial vertices in maintaining barrier function

This study demonstrated that loss of PAK4 or inhibition of its activity leads to impaired epithelial barrier function, with the ZnUMBA assay revealing particularly pronounced barrier defects at tricellular and multicellular vertices. These sites are known to be mechanically vulnerable because they concentrate tensile forces, and the observed defects highlight the importance of PAK4 in maintaining junctional integrity under mechanical stress. At tricellular junctions within vertebrate epithelia, barrier properties are normally ensured by specialized tTJs, composed of proteins such as tricellulin and angulins (Higashi and Miller, 2017; Higashi and Chiba, 2020). Upon PAK4 inhibition, however, angulin-1 localization at multicellular vertices exhibited gap-like openings at the center (Fig. S5A), suggesting that PAK4 is essential for preserving the architecture of tTJs. Previous studies have shown that epithelial barriers are dynamically maintained by mechanisms including localized and transient Rho activation ("Rho flares") (Stephenson et al., 2019; Varadarajan et al., 2022; Chumki et al., 2022; Craig et al., 2025) and claudin-7 polymerization following its release from EpCAM or Trop2 by proteases (Higashi et al., 2023a; Cho et al., 2025). Rho flares frequently occur at multicellular vertices (Stephenson et al., 2019), and polymerized claudin-7 also accumulates at vertices (Higashi et al., 2023a). Together, these observations indicate that multiple vertex-centered mechanisms act in concert to sustain barrier function and that PAK4 likely interfaces with these repair systems. Elucidating the phosphorylation targets of PAK4 will therefore provide important insight into the molecular pathways governing barrier maintenance at tricellular vertices.

New insights into the sarcomere-like structure at bicellular cell-cell junctions

In *PAK4*-KO and *Afdn*-KO cells, we observed that myosin IIA and IIB exhibit a punctate, sarcomere-like organization along cell-cell junctions. Although the

molecular mechanism underlying the generation of these sarcomere-like arrays
remains unclear, their formation does not require either PAK4 or Afadin. Mixed-
culture experiments with Ctrl cells demonstrated that these proteins do not
suppress sarcomere-like structure formation through a direct, cell-autonomous
mechanism but rather influence junctional actomyosin patterning in a cell-
nonautonomous manner. This phenotype may reflect an imbalance in mechanical
forces acting at cell-cell junctions, where tensile forces perpendicular and parallel
to the junction axis normally cooperate to support epithelial morphology and
mechanoresponsiveness (Lecuit and Yap, 2015; Tang, 2018; Angulo-Urarte et al.,
2020; Lenne et al., 2021). Among these, perpendicular forces, in particular, drive
the zigzag morphology of cell-cell junctions (Miyazaki et al., 2023), which is
prominent in Ctrl MDCK II cells. Attenuation of perpendicular forces may allow
parallel actomyosin arrays to stabilize, thereby promoting sarcomere-like patterning
along bicellular junctions and increasing tension at vertices. This raises the
question of which molecules are responsible for generating perpendicular forces at
cell-cell junctions. ZO proteins are prime candidates, as ZO-1/ZO-2 double
depletion induces sarcomere-like structures (Yamazaki et al., 2008; Fanning et al.,
2012; Choi et al., 2016; Otani et al., 2019), whereas ZO-1 overexpression
exaggerates junctional curvature (Tokuda et al., 2014). ZO-1 may exert these
perpendicular forces through mechanosensitive conformational changes (Spadaro
et al., 2017), which are reduced in *claudins/JAM-A*-KO cells that also display
junctional myosin bundles (Nguyen et al., 2024). Notably, both ZO-deficient cells
and Afadin-deficient cells failed to develop sarcomere-like myosin arrays when
surrounded by control cells (Choi et al., 2016) (Fig. 8H), implicating a similar cell-
nonautonomous mechanism. For example, Afadin and PAK4 might regulate ZO-1
conformation similar to claudins/JAM-A. Alternatively, they might generate
perpendicular tensile forces on bicellular junctions independently of ZO proteins,
although this possibility appears less likely, since junctions are straightened in ZO-
1/2 double-knockdown cells despite the increased junctional accumulation of
Afadin (Choi et al., 2016), suggesting that perpendicular force generation does not
simply correlate with the amount of Afadin at junctions. Of interest, the *Drosophila*
homolog of ZO-1, Polychaetoid, also works in parallel with Afadin homolog, Canoe,
to maintain epithelial integrity during development (Manning et al., 2019). Future
studies should address how Afadin/PAK4 and ZO proteins cooperatively maintain
epithelial integrity.
732
733 **Concluding remarks**
734   Here, we demonstrate a novel role for PAK4 and Afadin in the maintenance
735 of epithelial integrity and barrier function during vertex remodeling. Defining
736 phosphorylation targets of PAK4 and downstream signaling pathways is an
737 important future direction.
738

Materials and methods
Cell culture
MDCK II and HEK293T cells were cultured in Dulbecco's Modified Eagle Medium (DMEM; D5796; Sigma-Aldrich, St. Louis, MO, USA) supplemented with 5% fetal bovine serum (FBS; F7524; Sigma-Aldrich). Cells were maintained in a 5% CO_2 incubator at 37ºC. All KO clones of MDCK II cells were established from a parental *cldn2*-KO MDCK II cell clone (Saito et al., 2021). Thus, the *cldn2*-KO cell line is referred to as the "Ctrl" cell line throughout this manuscript. The *cldn2*-KO cell clone expressing mCherry conjugated with 3xnuclear localization signal (mChe-NLS) was described previously (Higashi et al., 2023b).

Antibodies used in cell culture
The following primary antibodies were used: Rabbit anti-PAK4 monoclonal antibody (mAb) (clone E7H7R; #52694), rabbit anti-phospho-Myosin light chain 2 (Thr18/Ser19) polyclonal antibody (pAb) (#3674), and rabbit anti-EpCAM mAb (clone E6V8Y; #93790) (Cell Signaling Technology, Danvers, MA, USA); rabbit anti-l/s-Afadin pAb (#A0224) and mouse anti-vinculin mAb (clone VIN-11-5; #V4505) (Sigma-Aldrich); rabbit anti-non-muscle Myosin heavy chain II-A (MyoIIA) pAb (#909801) and rabbit anti-non-muscle Myosin heavy chain II-B (MyoIIB) pAb (#909901) (BioLegend, San Diego, CA, USA); rat anti-ZO-1 (alpha+) mAb (clone R40.76; sc-33725), mouse anti-claudin-1 mAb (clone XX7; sc-81796), and mouse anti-beta-catenin mAb (clone E-5; sc-7963) (Santa Cruz Biotechnology, Dallas, TX, USA); mouse anti-DYKDDDDK tag (FLAG-tag) mAb (clone 1E6; #012-22348) (Wako-Fujifilm, Osaka, Japan); mouse anti-p120-catenin (CTNND1) mAb (clone 2F7H8; #66208-1-Ig), mouse anti-GFP tag mAb (clone 1E10H7; #66002-1-Ig), and rabbit anti-mCherry pAb (#26765-1-AP) (Proteintech, Rosemont, IL, USA); rabbit anti-claudin-3 pAb (#34-1700), mouse anti-claudin-4 mAb (clone 3E2C1; #32-9400), and Alexa Flour 488-conjugated mouse anti-ZO-1 mAb (Clone ZO1-1A12; #339188) (Thermo Fisher Scientific, Waltham, MA, USA); rabbit anti-claudin-7 pAb (#18875) (Immuno-Biological Laboratories, Fujioka, Gunma, Japan); rabbit anti-Partitioning-defective 3 (Par3) pAb (#07-330) (Merck Millipore, Burlington, MA, USA).
For immunofluorescence imaging, the following secondary antibodies were used: Alexa Fluor 488-conjugated AffiniPure donkey anti-mouse IgG (H+L) pAb (#715-545-151), Alexa Fluor 488-conjugated AffiniPure donkey anti-rabbit IgG (H+L) pAb (#711-545-152), Alexa Fluor 488-conjugated AffiniPure donkey anti-rat IgG (H+L) pAb (#712-545-153), Cy3-conjugated AffiniPure donkey anti-mouse IgG (H+L) pAb (#715-165-151), Cy3-conjugated AffiniPure donkey anti-rabbit IgG (H+L) pAb (#711-165-152), Cy3-conjugated AffiniPure donkey anti-rat IgG (H+L) pAb (#712-165-153), Alexa Fluor 647-conjugated AffiniPure donkey anti-rabbit IgG (H+L) pAb (#711-605-152), and Alexa Fluor 647-conjugated AffiniPure donkey anti-mouse IgG (H+L) pAb (#715-605-151) (Jackson ImmunoResearch Laboratories, West Grove, PA, USA).
For immunoblotting, the following secondary antibodies were used: Horseradish peroxidase (HRP)-linked sheep anti-mouse IgG pAb (#NA931V) (GE Healthcare, Chicago, IL, USA); HRP-linked goat anti-rabbit IgG pAb (#7074P) (Cell Signaling Technology); HRP-linked anti-β-actin mAb (clone C4; sc-47778HRP; Santa Cruz Biotechnology).

787
Generation of KO MDCK II cell lines
Cldn2-KO cells were described previously (Saito et al., 2021). To generate *PAK4*-
KO and *Afdn*-KO cells, DNA oligonucleotides encoding single-guide RNAs
(sgRNAs) were synthesized (Integrated DNA Technologies; Coralville, IA, USA),
and the annealed DNA oligonucleotides were cloned into pSpCas9 (BB)-2A-Puro
(PX459) plasmids (plasmid #62988; Addgene, MA, USA) (Ran et al., 2013) at the
Bbs I cleavage sites. Two gRNAs were designed for each gene to induce an
excision of exons encoding the entire coding region (*PAK4*) or the initiation codon
(*Afdn*) (Saito et al., 2024). Parental cells were transiently transfected with a pair of
plasmids encoding sgRNAs using PEI-max (Polysciences, PA, USA) and selected
using 3 µg/mL of puromycin (Sigma-Aldrich) for 1 d. The cells were then sparsely
seeded onto 10-cm dishes and cell clones were obtained by scraping off the
colonies. Genomic DNA was extracted from cell clones and screened by PCR
using GoTaq DNA polymerase (#M7123; Promega, Madison, WI, USA) and
specific primers. PCR products of KO clones were sequenced (Macrogen, Tokyo,
Japan) to determine the precise deletion sites. After the establishment of cell
clones, we confirmed that the cells were no longer resistant to puromycin.

DNA constructs used in cell culture
A cDNA library of MDCK II cells was synthesized using the PrimeScript II 1st
strand cDNA synthesis kit (#6210A; Takara Bio). cDNAs encoding canine *PAK4*,
AFADIN (*Afdn*), and *NECTIN-2* and their fragments were amplified using
PrimeSTAR GXL DNA polymerase (#R050A; Takara Bio) from the cDNA library of
MDCK II cells. Primer sets used are 5'-
AT*GGATCC*GGCACC<u>ATG</u>TTTGGGAAGAAGAAG-3' and 5'-
AT*GAATTC*ATCTGGTGCGGTTCTGGCGC-3' (*PAK4*); 5'-
ATG*AGATCT*GGGAGG<u>ATG</u>TCGGCGGGCGGC-3' and 5'-
ATT*GAATTC*ACTTGGTGTTCAGTTCGTTTTC-3' (*AFADIN*); 5'-
AATAAT*GCTAGC*<u>ATG</u>GCCCGGGCCGCAGCCCTCC-3' and 5'-
AATAAT*GCGGCCGC*CCACATAAATCGCCCGCCGTGAAATGAGG-3' (*NECTIN-*
2). Restriction sites (BamH I and EcoR I for *PAK4*; Bgl II and EcoR I for *AFADIN*;
Nhe I and Not I for *NECTIN-2*) are italicized, and the start and termination codons
are underlined. Five restriction sites in the *AFADIN* cDNA sequence were disrupted
by introducing a silent mutation using specific primer sets (5'-
GATGACATTGA*GAA<u>C</u>TC*CCGACTGGCTGC-3' and 5'-
GCAGCCAGTCGG*GA<u>G</u>TTC*TCAATGTCATC-3';
GGAGGATGCAG*GA<u>G</u>TTC*CGCAGCTCGGATG-3' and 5'-
CCGAGCTGCG*GAA<u>C</u>TC*CTGCATCCTCCTCTC-3'; 5'-
CGGATCAG*GAAT<u>C</u>C*AGCCCCACCACTGC-3' and 5'-
GTGGTGGGGCT*<u>G</u>GATTC*CTGATCCGAAC-3'; 5'-
GCAAGCATT*GA<u>G</u>TTC*AGGGAAAGTTCTG-3' and 5'-
CTTTCCCT*GAA<u>C</u>TC*AATGCTTGCAGGC-3'; 5'-
GCAGGACCG*AGA<u>C</u>CT*TAGTCGAATCAC-3' and 5'-
GTGATTCGACTA*A<u>G</u>GTCT*CGGTCCTGC-3'). The fragments were digested with
restriction enzymes, purified with Nucleospin Gel and PCR Clean-up kit (#U0609C;
Takara Bio), and cloned into the pCAG vector with 1xFlag (DYKDDDDK) and
2xStrep II (WSHPQFEK) tags (pCAG-FS2 vector) or pCAG-nGFP (Saito et al.,

2021). The DNA sequences were verified (Macrogen, Eurofin Genomics [Tokyo,
Japan], or Azenta [Burlington, MA, USA]).
837
Immunofluorescence microscopy of MDCK II cells
For fluorescence microscopy of MDCK II cells, 2×10^5 cells/mL of cells were
seeded onto glass coverslips placed in wells of 12-well plates. In Fig. 3 experiment,
PF-3758309 (#6005; Tocris) dissolved in DMSO (#D2550; Sigma-Aldrich) was
added 4 hrs before fixation. After 36-44 hrs, the cells were fixed with 100% ethanol
(#057-00456; Wako) for 15 min at -20°C or 2.7% formalin (#064-00406; Wako)
(1% formaldehyde) in PBS for 20 min at 20-28°C and permeabilized with 0.2%
Triton X-100 (#168-11805; Wako) in PBS.
For staining of established epithelial sheets of MDCK II cells, 1.0×10^5 cells/mL of
cells were seeded onto polycarbonate Transwell filters with 0.4-μm pore size
(#3401, Corning). Culture medium was replaced daily from the day after seeding.
After 6 days, the cells were fixed with 100% ethanol for 15 min at -20°C or 2.7%
formalin in PBS for 20 min at 20-28°C and permeabilized with 0.2% Triton X-100 in
PBS. For side-by-side culture (Fig. 8I), 80 μL suspensions of Ctrl and *PAK4*-KO#1
cells at 5.0×10^5 cells/mL density were placed on each half of the Transwell filter,
carefully avoiding contact between the two droplets. On the day of seeding, the
plate containing filters were gentry returned to the incubator to prevent the two cell
populations from merging. From the following day, once cells had adhered to the
filter, the culture medium was replaced daily in the same manner as for standard
culture.
Coverslips or filter membranes excised from the filter cups were washed with PBS
thrice. After blocking with 2% bovine serum albumin (BSA) (#A7030; Sigma-
Aldrich) in PBS, the samples were incubated with primary antibodies diluted in PBS
containing 0.2% BSA for 1 h at 20-28°C. After washing with PBS, the samples
were incubated with fluorescently labelled secondary antibodies diluted in PBS
containing 0.2% BSA for 1 h at 20-28°C. After washing, the cells were embedded
in FLUORO-GEL II with DAPI (Electron Microscopy Sciences, PA, USA). The
samples were observed using a laser scanning confocal microscope (FV1000;
Olympus, Tokyo, Japan) with a 60x oil-immersion objective lens (UPlanSApo 60x,
Olympus) at laser wavelengths of 405, 488, 559, and 635 nm, or a fluorescence
microscope (BX61, Olympus) with a 40x objective lens (UPlanSApo 40x, Olympus)
equipped with a mercury lamp, dichroic filter sets (NIBA, WIG and WU) and a
cooled CCD camera (DP71, Olympus). Images were acquired using the FluoView
ver. 4.2b (Olympus) or CellSens ver. 1.14 (Olympus) and processed using ImageJ
(NIH) and Photoshop (Adobe, CA, USA).
873
Immunofluorescence microscopy of mouse small intestine
All experiments were carried out strictly adhered to the Japanese Guidelines for
Proper Conduct of Animal Experiments. The protocol for the experiments using
animals (#2024025) was reviewed by the Animal Care and Use Committee of
Fukushima Medical University and approved by the university's president. The 8-
week-old female C57/BL6 mice were euthanized by cervical dislocation, and
dissected tissue blocks were fixed in 4% paraformaldehyde (PFA) at 4°C for 2 hr,
serially soaked in 15% and 30% sucrose in PBS, and embedded in OCT
compound (Sakura Finetek Japan, Tokyo, Japan). The frozen tissues were cut into

883 6-μm-thick sections in a cryostat (Leica CM1950; Leica Biosystems, Nussloch,
884 Germany) at −20ºC, soaked in 0.5% Antigen activator Immunosaver (#333; Nisshin
885 EM, Tokyo, Japan) solution at 70ºC for overnight. The sections were permeabilized
886 with PBS containing 0.1%(w/v) Triton X-100 for 15 min at 20-28ºC, blocked with
887 2% BSA in PBS, stained with primary and secondary antibodies, and observed
888 with FV1000 confocal microscope as described above.
889

Quantification of fluorescence intensity at bicellular junctions and
890 
891 **multicellular vertices in mouse tissues and MDCK II cells**
892 Z-stacks of confocal fluorescence images were used for quantification.
893 Fluorescence intensities were measured using ImageJ software. Circular regions
894 of interest (ROIs) with a diameter of 5 pixels were placed manually at each vertex,
895 which were classified as tricellular vertices or tetracellular vertices/rosettes (>4
896 cells). For bicellular junctions, linear ROIs with a thickness of 5 pixels were drawn
897 along the entire length of each junction. Background values were determined for
898 each image by placing five circular ROIs with100-pixel diameter in a cell-free
899 region (for Fig. 1A-C) or ROIs with 50-pixel diameter in a junction-free region (for
900 Fig. 1D-F). The mean background intensity was subtracted from vertex and
901 bicellular junction measurements. For each image, a length-weighted mean
902 fluorescence intensity of bicellular junctions was calculated and used for
903 normalization. Vertex intensities were then expressed relative to this bicellular
904 reference (bicellular = 1). Data were visualized as bubble plots, in which the area
905 of each bubble corresponded to the length of the traced bicellular junction, while
906 tricellular and tetracellular/rosette vertices were plotted as normalized values with a
907 uniform bubble size.
908

Quantification of epithelial integrity of MDCK II cells
909 
910 Fluorescence images of ZO-1 and DAPI from 4 coverslips for each cell clone or
911 each treatment were used for quantification. For each coverslip, five non-
912 overlapping fields of 1360 x 1024 pixels (corresponding to 216 x 163 μm) were
913 collected, resulting in total of 20 images per cell clone or treatment. To assess
914 epithelial integrity, ZO-1 staining was used to identify and manually mark all
915 vertices in each image using ImageJ software. Vertices were classified as
916 tricellular, tetracellular, or rosette-type (>5 cells), and the proportion of each
917 category was calculated. Discontinuities in cell-cell junctions were also counted
918 from ZO-1 images. Cell numbers were determined from the DAPI channel. Nuclei
919 touching the image border were counted as 0.5 to correct for partial cells. The total
920 number of vertices was divided by the number of nuclei to obtain the vertex-to-cell
921 ratio. Similarly, the number of junctional discontinuities was normalized to the
922 number of nuclei to yield the number of discontinuities per cell. Each image was
923 treated as one sample (n = 20 per clone).
924

Immunoblotting
925 
926 Cells or bead-bound proteins were boiled in the SDS sample buffer for 5 min.
927 Proteins were separated by SDS-PAGE using 5-20% gradient gels (Fujifilm Wako,
928 Osaka, Japan), and transferred to a polyvinylidene fluoride (PVDF) membrane
929 (Immobilon, Merck). The membranes were blocked with 5% non-fat dried milk in
930 Tric-buffered saline (TBS) containing 0.1% Tween-20 (TBS-T) for 30 min at 20-

28ºC, followed by incubation with a primary antibody diluted in TBS-T overnight at
4ºC. After washing with TBS-T, the membranes were incubated with an HRP-
conjugated secondary antibody in TBS-T for 1 h at 20-28ºC. After washing with
TBS-T, the membranes were incubated with ECL Prime (GE Healthcare) and
developed using LAS4000 (GE Healthcare). The images were processed using
Photoshop.
937
Co-immunoprecipitation
To examine the interaction between PAK4 and Afadin, HEK293T cells were co-
transfected with the expression vectors using PEI-Max in a 6-well plate and
cultured for 2 d. For FLAG-IP, cells were lysed with IP-lysis buffer (50 mM Tris-HCl,
pH 7.4, 150 mM NaCl, 0.5% Nonidet P-40 [NP-40, #25223-04, Nacalai Tesque,
Kyoto, Japan], 1 mM EDTA, 1 mM DTT, cOmplete protease inhibitor cocktail
[#11836145001; Sigma-Aldrich]), precleared with Protein G-Sepharose
(#17061801; Cytiva, Tokyo, Japan), and centrifuged at 15,000 x g for 20 min at
4ºC. Supernatants were mixed with mouse-anti-DYKDDDDK mAb-bound Protein
G-Sepharose and rotated for 2 hr at 4ºC. Beads were washed five times using IP-
wash buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 0.05% NP-40, 1 mM EDTA, 1
mM DTT) and bead-bound proteins were eluted with SDS sample buffer. For GFP-
trap, cells were lysed with GFP-trap lysis buffer (10 mM Tris-HCl, pH 7.5, 150 mM
NaCl, 0.5% NP-40, 0.5 mM EDTA, cOmplete protease inhibitor cocktail) and left
on-ice for 30 min with pipetting every 10 min. Lysates were centrifuged at 15,000 x
g for 20 min at 4ºC and diluted with 1.5x volume of dilution buffer (10 mM Tris-HCl,
pH 7.5, 150 mM NaCl, 0.5 mM EDTA). Diluted lysates were mixed with GFP-trap
magnetic agarose (#GTMA-20; Proteintech) and rotated for 1 hr at 4ºC. Beads
were washed five times with GFP-trap wash buffer (10 mM Tris-HCl, pH 7.5, 150
mM NaCl, 0.05% NP-40, 0.5 mM EDTA) using a magnet. Bead-bound proteins
were eluted with SDS sample buffer.
959
Transepithelial electrical resistance (TER) measurement
Cells were seeded onto 12-well polycarbonate Transwell filters with 0.4-µm pore
size at 1.0×10^5 cells/mL and cultured for 7 d. The transepithelial electrical
resistance (TER) of the cell sheet was measured each day using a volt-ohm meter
(Millicell ERS-2; EMD Millipore). All measurements were subtracted by a blank
measurement of a Transwell filter with medium alone and then multiplied by the
culture area of the Transwell filter to calculate the unit area resistance.
967
AlphaFold3 prediction of the structure of PAK4-Afadin complex
The structures of the PAK4-Afadin complex, PAK4 alone, and Afadin alone were
predicted using the AlphaFold server (https://alphafoldserver.com) (Abramson et
al., 2024) and visualized using Open-Source PyMOL (The PyMOL Molecular
Graphics System, Version 3.0 Schrödinger, LLC., Available at:
http://www.pymol.org/pymol). PAE values were visualized with PAE viewer
(https://pae-viewer.uni-goettingen.de) (Elfmann and Stülke, 2023). Possible
interacting region was visualized with AlphaBridge (https://alpha-bridge.eu)
(Álvarez-Salmoral et al., 2024).
977
DNA constructs used in *Xenopus* experiments

Xenopus PAK4.L sequence (*X. laevis* v10.1) was synthesized by TWIST
Bioscience (South San Francisco, CA, USA) and cloned into pCS2+/mNeon.
Untagged *Xenopus* PAK4-NT was cloned by PCR amplifying the N-term fragment
(amino acids:
MFAKKKKRVEISAPSNFEHRVHTGFDQQEQKFTGLPRQWQSLIEESAKRPKPLV
DPSYITTIKHVPQKTIVRGNKMSLDGSLAWLLDEFDDMSVCRSNS) from
pCS2+/mNeon-PAK4 plasmid (described above) and inserting it into a pCS2+
vector digested with EcoR I and Xho I using Gibson cloning (Gibson et al., 2009)
with the following primers: Forward 5′-
CTTGTTCTTTTTGCAGGATCCCATCGATTCGAATTCACCATGTTTGCTAAGAAG
AAGAAGC-3'; Reverse 5′-
GTTTGCCGGTCGAACTCTTAACTCGAGCCTCTAGAACTATAGTGAGTCGTATT
ACGT-3'. Constructs were verified by sequencing using Plasmidsaurus (Louisville,
KY, USA). pCSf107mT/Lifeact-miRFP703 (Yamamoto et al., 2021) was a gift from
Dr. Kazuhiro Aoki's lab (Kyoto University, Japan). All other DNA constructs were
previously described: pCS2+/TagBFP2-ZO-1 (Stephenson et al., 2019),
pCS2+/PLEKHA7-mCherry (Higashi et al., 2019), and pCS2+/Angulin-1-3xGFP
(Higashi et al., 2016).
997
mRNA preparation
DNA was linearized using either Not I or Kpn I (in the case of ZO-1 construct).
Linearized or non-linearized (in the case of Lifeact-miRFP703 construct) DNA was
then transcribed using the mMessage mMachine SP6 Transcription Kit (#AM1340,
Thermo Fisher Scientific). Transcribed mRNA was purified using the RNeasy Mini
Kit (#74104, Qiagen, Germatown, MD, USA) and stored at -80ºC until use.
1004
***Xenopus laevis* embryos and microinjections**
All experiments conducted using *Xenopus laevis* embryos were carried out in
compliance with the University of Michigan Institutional Animal Care and Use
Committee and the U.S. Department of Health and Human Services Guide for the
Care and Use of Laboratory Animals. The day prior to egg collection, female
Xenopus laevis frogs (Xenopus 1 or National *Xenopus* Resource [NXR]) were
induced to hyperovulate using human chorionic gonadotrophin (#198591, MP
Biomedicals, Irvine CA, USA). The following day, eggs were collected from female
frogs and fertilized *in vitro*. Fertilized embryos were dejellied approximately 30 min
after fertilization and transferred to 0.1× MMR (10 mM NaCl, 0.2 mM KCl, 0.2 mM
$CaCl_2$, 0.1 mM $MgSO_4$, and 0.5 mM HEPES, pH 7.4-7.6). Once embryos
developed to the 2- or 4-cell stage, they were microinjected with the desired
mRNAs. Cells were injected once per cell (for embryos at the 4-cell stage) or twice
per cell (for embryos at the 2-cell stage). For PAK4, ZO-1, and PLEKHA7
colocalization data, each 5 nl of injection volume contained: 25 pg of PAK4-mNeon,
70 pg of TagBFP2-ZO-1, and 28 pg of PLEKHA7-mCherry. For PAK4 inhibitor
experiments, each 5 nl of injection volume contained: 70 pg of TagBFP2-ZO-1, 25
pg of Angulin-1-3xGFP, and 150 pg of LifeAct-miRFP. For PAK4-NT experiments,
each 5 nl of injection volume contained: 70 pg of TagBFP2-ZO-1, 150pg of LifeAct-
miRFP, and 150-200pg of PAK4-NT. For ZnUMBA imaging, albino *Xenopus laevis*
embryos were used. Embryos were injected with the following mRNA constructs:
70 pg of TagBFP2-ZO-1 and 150 pg of PAK4-NT, per 5 nl injection volume.

Injected embryos were then incubated in 0.1× MMR and allowed to develop for 16-
24 h at 15°C or 13°C until they reached gastrula stage (Nieuwkoop and Faber
stages 10-11).
1030
Zinc-based Ultrasensitive Microscopic Barrier Assay (ZnUMBA)
For ZnUMBA experiments, gastrula-stage embryos were microinjected one time in
the blastocoel with 10 nl of a mixture of 100 µM $CaCl_2$, 100 µM EDTA, and 1 mM
FluoZin-3 (#F24193, Thermo Fisher Scientific). FluoZin-3–injected embryos were
allowed to recover from microinjection for 5-10 min in 0.1× MMR. After recovery,
embryos were mounted in 1 mM $ZnCl_2$ in 0.1× MMR before imaging (Higashi et al.,
2023b).
1038
Immunofluorescence staining of *Xenopus* embryos
PFA fixation was performed as follows: gastrula-stage albino embryos were placed
in a solution of 1.5% PFA, 0.25% glutaraldehyde, 0.2% Triton X-100, and Alexa
Fluor 568 phalloidin (#A12380, Thermo Fisher Scientific) (1:100) in 0.88× MT
buffer (80 mM K-PIPES, 5 mM EGTA, 1mM $MgCl_2$, pH 6.8) and allowed to fix on a
shaker overnight at room temperature. Fixed embryos were quenched in 100 mM
sodium borohydride in PBS on a shaker for 1 hr at room temperature. Embryos
were then bisected to keep the animal cap and blocked with blocking solution (10%
FBS, 5% DMSO, 0.1% NP-40 in 1× TBS) overnight on a nutator at 4°C. The animal
caps were then incubated overnight on a nutator at 4°C in the blocking solution
with mouse anti-PAK4 mAb (#sc-390507, Santa Cruz Biotechnology) (1:50). Next,
they were washed three times in blocking solution and incubated overnight on a
nutator at 4°C in the blocking solution with goat Alexa Fluor 488 anti-mouse pAb
(A-11001, Invitrogen) (1:200), and Alexa Fluor 568 phalloidin (A12380, Invitrogen)
(1:100). The next day, embryos were washed in 0.1% NP-40 in TBS (TBSN), then
incubated in 1:1000 DAPI (10 mg/ml stock) diluted in TBSN for 30 min at room
temperature on the nutator. Next, they were washed for 10 min with TBSN five
times. Finally, embryos were washed and mounted in PBS before imaging.
1057
Live imaging of *Xenopus* embryos
Live imaging videos of gastrula-stage *Xenopus* embryos were acquired using an
Olympus FluoView 3000, and 60X supercorrected Plan Apo N 60X OSC objective
(NA = 1.4, working distance = 0.12 mm), with mFV10-ASW software. Embryos
were mounted as described previously (Reyes et al., 2014). Embryos were imaged
with a scanning speed of 2 or 4 µs/pixel, a 512 × 512 field of view, 1× zoom,
capturing 6-10 of the most apical slices of the animal cap of the embryo, with a
step size of 0.5 µm. For live imaging of PAK4 with ZO-1 and PLEKHA7 (Fig. 2E),
embryos were imaged with a scanning speed of 4 µs/pixel, a 512 × 512 field of
view, 1× zoom, capturing 50 of the most apical optical slices of the animal cap of
the embryo, with a step size of 0.5 µm (Higashi et al., 2019). Fixed embryos were
imaged with a scanning speed of 4 µs/pixel, a 512 × 512 field of view, 1× zoom,
capturing 40 optical slices of the embryo, with a step size of 0.5 µm.
1071
Drug treatment of *Xenopus* embryos
The PAK4 inhibitor, PF-3758309 (S7094, Selleck, Houston, TX, USA), was
resuspended in DMSO as a 10 mM stock solution, aliquoted and stored in -80°C.

Following mRNA injection at the 2- or 4-cell stages, embryos were incubated
overnight at 15°C in either 0.1% DMSO (Control) or 10 µM PAK4 inhibitor in 0.1×
MMR.
1078
Image analysis of *Xenopus* embryos
Quantification of the localization of PAK4 at cell-cell junctions
Gastrula-stage embryos expressing PAK4-mNeon together with PLEKHA7-
mCherry and TagBFP2-ZO-1 were live imaged as described above. Square
images (50 pixels × 50 pixels) of straight junctions were excised from the original
images, and 50 z-slices across the junctions were averaged using ImageJ.
Fluorescence intensity profiles along x- and z-axes were determined by multiple
line scans using ImageJ (four line scans from each embryo) (Higashi et al., 2019).
This process was repeated for 16 junctions from 5 embryos and aligned using the
highest ZO-1 intensity; mean and SD was calculated. For measuring the intensity
of PAK4 and ZO-1 at the vertices, a circular ROI of size 10 x 10 pixels was used to
measure the intensity at 16 bicellular junction (BCJ), tricellular vertices (TCJ) or
higher-order vertices (4-way and rosettes) from maximum z-projected images of 4
different embryos using Fiji. The intensity at TCJs and higher-order vertices were
normalized by their counterpart BCJ intensity.
1094
Quantification of PAK4 intensity at remodeling vertices
For measuring the change in PAK4 intensity, a circular ROI of size 3 x 3 pixels
(1.24 µm x 1.24 µm) was used to measure the intensity at remodeling vertices,
non-remodeling vertices, and the cytoplasm over 60-90 mins using Fiji. The
intensity was normalized as follows: ((Intensity at vertex - Cytoplasmic Intensity) /
Cytoplasmic Intensity). To measure the change in angle, the angle tool in Fiji was
used with the lines spanning 5 µm along each junction. The early stage of
remodeling was defined by time point when the vertex starts actively moving. The
late stage is towards the end as the vertex stabilizes. The middle stage is the time
point exactly halfway. A non-remodeling vertex was chosen in the same field of
view as a remodeling vertex and the measurements were taken for all the
timepoints as its remodeling vertex counterpart. PAK4 intensity at the remodeling
vertices at early, middle, and late stages was normalized as follows :(Intensity at
remodeling vertex/ Average intensity of non-remodeling vertices at that stage).
1109
Quantification of cell packing upon PAK4 inhibitor treatment
Distribution of cell-cell interfaces after cytokinesis: Using cells expressing
TagBFP2-ZO-1, the distribution of Type I (daughter-daughter interface), Type II (4-
way junction), and Type III (neighbor-neighbor interface) interfaces was determined
for each embryo. Movies of both control and PAK4-inhibited embryos were
analyzed from 13 to 60 minutes after ingression began.
Rosette resolution: Using cells expressing TagBFP2-ZO-1, the number of rosettes
(where 5 or more cells meet) at the start of imaging was determined for each
embryo. The marked rosettes were then analyzed for both control and PAK4
inhibited embryos for 60-80 minutes. The time when the 5-way vertex starts to
resolve from the beginning of the movie was noted as the time of rosette resolution.
1121
Quantification of cytokinesis failure

Using cells expressing TagBFP2-ZO-1, cell division events were followed for 13-96
minutes after ingression began in 5 controls (46 cell divisions) and 8 PAK4-NT-
expressing embryos (32 cell divisions). Formation of a 4-way vertex with clear ZO-
1 intensity was marked as cytokinesis completion, whereas dissolution of the
cytokinetic ring, inability to form a 4-way vertex with clear ZO-1 intensity, or
breaking off of an already formed daughter-daughter interface were considered
cytokinetic failure.
1130
ZnUMBA
Whole field intensity analysis: All movies were imaged using the same imaging
parameters. The images were maximum z-projected and contrast adjusted to 0-
4095 in Fiji. Whole field intensity was measured by Fiji at 0 min, ~7 min, and ~15
min since the start of imaging. If the junctions were not visible at 15 mins in the
FluoZin3 channel, it was considered a global barrier leak.
Junctional intensity analysis: Cell-cell junctions were traced using the TagBFP2-
ZO-1 channel and threshold option in Fiji. Post thresholding, the image was made
binary and an ROI was created tracing the cell-cell junction. The ROI was
superimposed on the FluoZin3 channel, and the intensity was measured in Control,
PAK4 inhibitor-treated embryos, and PAK4-NT-expressing embryos.
Quantification of leaky vertices: Barrier leaks were counted manually by identifying
vertices where FluoZin3 intensity increased over baseline intensity, as revealed by
a change in color using the FIRE LUT applied in ImageJ. This count was then
normalized by the number of vertices present in the field of view, and the
percentage of leaky vertices was calculated. In the case of PAK4-NT quantification,
embryos with global barrier leaks were excluded from this analysis.
Quantification of FluoZin3 intensity at higher-order vertices: A circular ROI of size 7
x 7 pixels (2.90 µm x 2.90 µm) was used to measure intensity at higher-order
vertices (4-way or higher) and in the cytoplasm at 0 min, ~7min, and ~15 min using
Fiji, and the intensity was normalized as follows: (Intensity at higher-order vertex -
Cytoplasmic Intensity) / (Cytoplasmic Intensity).
1153
1154   **Video editing**
1155   Videos were initially generated and concatenated using FiJi 2.16.0/1.5p. Overlay
1156   animations were added using Adobe Premiere pro 25.5.
1157
1158
Statistical analysis and plotting
Data shown in Figs. 1B-C, 1E-F, 3B-D, 5B-D, 6B-D, 6G-I, S2B-D, and S2G-I are
presented as the mean ± standard error (S.E.M.) using Microsoft Excel 365. Data
in Figs 8A-F and 8J-K are presented as the mean ± standard deviation using
Microsoft Excel 365. Statistical significance of the differences was evaluated using
a two-tailed unpaired Welch's t-test (Figs. 1B-C, 1E-F, 3B-D, 5B-D, 6B-D, 6G-I,
S2B-D, S2G-I). Bonferroni correction was used to address the multiple comparison
problem in Figs. 3B-D, 5B-D, 6B-D, 6-I, S2B-D, S2-I. All tests were performed
using Microsoft Excel 365. Weighted average intensity at bicellular junctions in Figs.
1B and 1E was calculated using R software (https://www.r-project.org) using
weighted.mean() function.

For data presented in Figs. 2, 4, 7, 9, 10, and S5, all graphs were plotted using
python 3.8.10 using the following packages: matplotlib (3.7.5), numpy (1.24.4),
openpyxl (3.1.5), pandas (2.0.3), pip (25.0.1), scipy (1.10.1), seaborn (0.13.2),
statsmodels (0.14.1), and statannotations (0.7.2). Some of the basic data handling
was performed on Microsoft Excel 365. A two-tailed unpaired Student's *t*-test was
performed for Figs. 2C, 2D, 2G, 4G, 7C, 9D, 10C-D, 10F, and S5B-C. A two-tailed
unpaired Welch's *t*-test was used for Figs. 9C and 10E.
1177
1178
1179

Acknowledgements
We would like to express our gratitude to Misa Satoh, Seiko Watanabe, Keiko
Watari, Chiaki Ozaki, and Joji Kai for their technical assistance, Ryutaro Shirakawa
(Japan Institute for Health Security) for the critical review of our manuscript, and
Janis F. Miroll for assistance with statistical analysis and plotting with Python. This
work was funded by the JSPS (Japan Society for the Promotion of Science)
KAKENHI (Grant Number 24K09456), the 2024 Naito Grant for the advancement
of natural science from Naito Foundation, the 2024 Narishige Zoological Science
Award from Narishige Fund to T.H., and NIH R35GM153204 to A.L.M. The authors
declare no competing financial interests.
Author contributions: This project was designed by T.H. All the experiments and
data analyses using MDCK II cells were performed by T.H. with input from H.C. All
the experiments and data analyses using *Xenopus* embryos were performed by
B.A. A.C. and A.Y.H made DNA constructs and performed immunostaining of
mouse tissues, respectively. A.L.M. and T.H. supervised the project. A.L.M. and
T.H. were responsible for project administration. B.A. and T.H. wrote the
manuscript with input from other authors, and all authors reviewed and revised the
manuscript.

1198

 Figure legends
Figure 1. PAK4 is enriched at higher-order cell-cell junctions in the mouse
small intestine and MDCK II cells.
(A,D) Immunofluorescence staining of mouse small intestine fixed with
formaldehyde (A) and of EtOH-fixed Ctrl MDCK II cells (D). Samples were labeled
with rabbit anti-PAK4 mAb and mouse anti-ZO-1 mAb (A) or rabbit anti-PAK4 mAb
(green), rat anti-ZO-1 mAb (red) and DAPI (blue) (D). White arrowhead, tricellular
vertices with PAK4 signal; yellow arrows, 4-way vertices/rosettes. Scale bars, 20
μm. (B, C, E, F) Quantification of fluorescence intensities. Background-subtracted
fluorescence intensities of PAK4 (B, E) and ZO-1 (C, F) at tricellular vertices and 4-
way vertices/rosettes were normalized to the junction-length-weighted average
intensity at bicellular junctions. Individual measurements at bicellular junctions are
shown as circles, with the areas proportional to junction length. Individual
measurements at tricellular vertices and 4-way vertices/rosettes are shown as
circles and the bars indicate mean ± sd. The values of tricellular vertices and 4-way
vertices/rosettes were compared using a two-tailed unpaired Welch's t-test (ns, p >
0.05; **, p < 0.01; ****, p < 0.0001).
1216
Figure 2. PAK4 is enriched at remodeling higher-order cell-cell vertices in
***Xenopus* epithelia**
(A) Immunofluorescence staining of the animal hemisphere of gastrula-stage (NF
stage 10) *Xenopus* embryos with mouse anti-PAK4 mAb (green), phalloidin
(magenta), and DAPI (blue). Zoomed-in panels highlight PAK4 recruitment at
tricellular vertices (blue boxes) and 4-way vertex/rosette (orange boxes). Scale
bars, 20 μm (upper), 10 μm (lower). (B) Live confocal images of epithelial cells in
gastrula-stage *Xenopus* embryo expressing mNeon-PAK4 (PAK4), TagBFP2-ZO-1
(ZO-1), and PLEKHA7-mCherry (PLEKHA7). Zoomed-in panels highlight
localization of PAK4, ZO-1, and PLEKHA7 at bicellular junction (magenta),
tricellular vertex (blue), and higher-order vertex (orange). Scale bar, 20 μm. (C,D)
Quantification of PAK4 (C) and ZO-1 (D) fluorescence intensities at tricellular
vertices and 4-way vertices/rosettes, normalized to bicellular junction (BCJ)
fluorescence. Individual measurements (circles) and mean ± SEM (bars) are
shown. Values were compared using a two-tailed unpaired Student's t-test (****, p
< 0.0001; n = 16 vertices, 4 embryos). (E) Representative side view of a bicellular
junction. PAK4 (green) is localized apically along with TJ protein ZO-1 (blue). It
was a slight offset from AJ protein PLEKHA7 (magenta). Scale bar, 10 μm. Graph
shows intensity profiles of PAK4, ZO-1, and PLEKHA7 along the apico-basal axis
(mean ± SD. n = 16 junctions, 5 embryos). (F) Montage of non-remodeling (top)
and remodeling (middle and bottom) vertices labeled with PAK4-mNeon, shown
using the green Fire blue lookup table (LUT). PAK4 intensity at the vertex (green
line on graphs) and vertex angle (magenta line on graphs) were measured over
time. White arrows, vertex movement; yellow arrowheads, PAK4 accumulation; red
arrowheads, end of remodeling. Scale bars, 10 μm. (G) Quantification of PAK4
intensity at remodeling vertices at early, middle, and late stages of remodeling,
normalized to the average intensity of non-remodeling vertices at the same time
points. Individual measurements (circles) and mean ± SEM (bars) are shown.
Values were compared using a two-tailed unpaired Student's t-test (****, p <
0.0001; n = 14 vertices, 7 embryos, 3 independent experiments).

1247

Figure 3. PAK4 inhibition leads to an increase in higher-order cell-cell
vertices.
(A) Immunofluorescence staining of DMSO- or PF-3758309 (PAK4 inhibitor)-
treated Ctrl MDCK II (*cldn2*-KO) cells with rat anti-ZO-1 mAb (green) and DAPI
(blue). Arrows, 4-way junctions; circle, rosettes. Scale bar, 20 μm. (B-D)
Quantification of vertex-to-cell ratio (B), proportions of 4-way vertices and rosettes
(C), and number of discontinuities in cell-cell junctions (D). The values in PAK4
inhibitor-treated monolayers (n = 20) were compared with those of DMSO-treated
monolayers (n = 20) using a two-tailed unpaired Welch's *t* test with Bonferroni's
correction (ns, $p > 0.05$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; $p < 0.0001$). S.E.M.
(bars) and individual measurements (circles) are shown.
1259

Figure 4. PAK4 inhibition impairs higher-order junction remodeling in
***Xenopus* epithelia.**
(A) Live confocal images of epithelial cells in the animal hemisphere of gastrula-
stage *Xenopus* embryos expressing TagBFP2-ZO-1 (ZO-1), Angulin-1-3xGFP
(Angulin-1), and LifeAct-miRFP (F-actin) treated with 0.1% DMSO (control) or 10
μM PAK4 inhibitor PF-3758309. Zoomed-in panels (orange boxes) highlight the
junctional localization pattern of ZO-1, Angulin-1, and F-actin in control versus
PAK4 inhibitor-treated cells. Scale bars, 20 μm. (B) Schematic of 4-way vertex
resolution following cytokinesis. (C) Montage showing cells before cleavage furrow
ingression (gray) and at 13 and 60 min after ingression. Dividing cells are color
coded for type I (orange) and type II (blue) interfaces. Scale bars, 20 μm. (D)
Quantification of types I, II, and III interfaces at 13, 30, 45, and 60 min after
ingression. (control: 3 embryos; inhibitor: 4 embryos; 3 independent experiments).
(E) Montage of control and PAK4 inhibitor-treated embryos expressing TagBFP2-
ZO-1, showing rosettes before (yellow) and after resolution (green). Scale bars, 20
μm. (F) Quantification of rosette resolution at 15, 30, 45, and 60 min after the onset
of imaging (control: 5 embryos, 15 rosettes; inhibitor: 6 embryos, 26 rosettes; 3
independent experiments). (G) Quantification of time required for rosette resolution
(control: 5 embryos, 14 rosettes; inhibitor: 6 embryos, 14 rosettes; 3 independent
experiments). Data were compared using a two-tailed unpaired Student's *t*-test (**,
$p < 0.01$).
1281

Figure 5. Loss of PAK4 impairs epithelial integrity in cultured cells.
(A) Immunofluorescence staining of Ctrl and *PAK4*-KO cell clones (#1 and #2) with
anti-ZO-1 mAb (green) and DAPI (blue). Arrows, 4-way vertices; circle, rosettes;
yellow arrowheads, discontinuities in cell-cell junctions. Scale bar, 20 μm. (B-D)
Quantification of the vertex-to-cell ratio (B), proportions of 4-way vertices and
rosettes (C), and number of discontinuities in cell-cell junctions (D). Values from
PAK4-KO monolayers (n = 20) were compared with those of Ctrl monolayers (n =
20) using a two-tailed Welch's *t*-test with Bonferroni's correction (ns, $p > 0.05$; *, p
< 0.05; **, $p < 0.01$; ****, $p < 0.0001$). S.E.M. (bars) and individual measurements
(circles) are shown. (E) Immunofluorescence staining of Ctrl and *PAK4*-KO cells
with anti-MyoIIB pAb (green), anti-ZO-1 mAb (red), and DAPI (blue). Scale bar, 20
μm.
1294

Figure 6. Loss of Afadin compromises epithelial integrity, partially rescued
by junctional targeting of PAK4.
(A) Immunofluorescence staining of Ctrl and *Afdn*-KO cell clones (#1 and #2) using
anti-ZO-1 mAb (green) and DAPI (blue). Arrows, 4-way vertices; circle, rosette;
yellow arrowheads, discontinuities in cell-cell junctions. Scale bar, 20 µm. (B-D)
Quantification of the vertex-to-cell ratio (B), proportions of 4-way vertices and
rosettes (C), and number of discontinuities in cell-cell junctions (D). Values from
Afdn-KO monolayers (n = 20) were compared with those of Ctrl monolayers (n =
20) using a two-tailed Welch's *t*-test with Bonferroni's correction (ns, $p > 0.05$; *, p
< 0.05; ****, $p < 0.0001$). S.E.M. (bars) and individual measurements (circles) are
shown. (E) Immunofluorescence staining of *Afdn*-KO #1 cells expressing Nectin2-
FLAG-PAK4 WT or KR mutant using anti-ZO-1 (red) and anti-FLAG (green) mAbs
and DAPI (blue). Scale bar, 20 µm. (F) Immunofluorescence staining of *Afdn*-KO
#1 cells and *Afdn*-KO #1 cells expressing Nectin2-FLAG-PAK4 WT or KR mutant
labeled as in (A). Scale bar, 20 µm. (G-I) Quantification of the vertex-to-cell ratio
(G), proportions of 4-way vertices and rosettes (H), and number of discontinuities
in cell-cell junctions (I). Values from Nectin2-FLAG-PAK4-expressing Afdn-KO #1
cell monolayers (n = 20) were compared with those of *Afdn*-KO #1 monolayers (n =
20) using a two-tailed Welch's *t*-test with Bonferroni's correction (ns, $p > 0.05$; *, p
< 0.05; ***, $p < 0.001$; ****, $p < 0.0001$). S.E.M. (bars) and individual measurements
(circles) are shown.
1316
Figure 7. Inhibition of PAK4 kinase activity compromises junctional integrity
in *Xenopus* epithelia.
(A) Domain map of PAK4. N-terminal fragment of PAK4 (PAK4-NT) was expressed
in the embryos to act as a dominant-negative. (B) Live confocal images of epithelial
cells in the animal hemisphere of gastrula-stage *Xenopus* embryos expressing
TagBFP2-ZO-1 (ZO-1) +/- PAK4-NT. Zoomed-in panels show examples
highlighting 4-way (magenta) and higher-order (5 way or more) (yellow) vertices.
Scale bars, 50 µm (upper), 25µm (lower). (C) Quantification of 3-way, 4-way, and
higher-order vertices (control: 12 embryos; PAK4-NT: 25 embryos; 3 independent
experiments). Data were compared using a two-tailed unpaired Student's *t*-test (***,
$p < 0.001$; **, $p < 0.01$) (D) Montage of control and PAK4-NT cells expressing
TagBFP2-ZO-1 (ZO-1) and LifeAct-miRFP (F-actin), showing changes before and
during ingression (gray) and post-cytokinesis (green). Scale bar, 25 µm. (E)
Quantification of cytokinetic failure (controls: 46 cell division events from 5
embryos; PAK4-NT: 32 cell division events from 8 embryo; 3 independent
experiments). (F) Montages of control and PAK4-NT embryos expressing
TagBFP2-ZO-1 (ZO-1) and LifeAct-miRFP (F-actin), showing junctional
discontinuities (blue arrowheads). Scale bars, 25 µm.
1335
Figure 8. PAK4 and Afadin contribute to epithelial barrier function in cultured
cells.
(A-H,J-K) TER measurements for the following cell types: (A) Ctrl, Ctrl cells
expressing mCherry-NLS (mChe-NLS), and *Ang1*-KO cells;(B) *PAK4*-KO cells; (C)
Afdn-KO cells; (D) a mixed culture of mChe-NLS and *Ang1*-KO cells; (E) a mixed
culture of mChe-NLS and *PAK4*-KO cells; (F) a mixed culture of mChe-NLS and
Afdn-KO cells; (J) *Afdn*-KO #1 cells expressing Nectin2-FLAG; (K) *Afdn*-KO #1

cells expressing Nectin2-FLAG-PAK4 WT or kinase-dead (KR) mutant. Values are
shown as mean ± SD. In D-F, solid lines represent TER values measured in mixed
cultures, while dotted lines indicate estimated combined resistance values
calculated from the separately measured TER values of mChe-NLS and KO cell
cultures. (G-H) Immunofluorescence staining of the *PAK4*-KO #1 (G) or *Afdn*-KO
#1 (H) cells cultured with (lower) or without (upper) mChe-NLS cells on the filter
using anti-MyoIIB pAb (left) and anti-ZO-1 mAb (right). Scale bars, 20 µm. (I)
Immunofluorescence staining at the boundary between mChe-NLS (asterisks) and
PAK4-KO #1 cells grown side by side on a filter using anti-MyoIIB pAb (green),
anti-ZO-1 mAb (red), and DAPI (blue). Double arrows indicate the distance
between Ctrl cells and sarcomere-like MyoII structure in the *PAK4*-KO cells. Scale
bar, 20 µm.
1355
Figure 9. PAK4 inhibition impairs epithelial barrier integrity at higher-order
vertices in *Xenopus* embryos.
(A) Schematic of the ZnUMBA assay. Disruption of TJ allows Zn^{2+} to bind FluoZin-
3, resulting in increased FluoZin-3 fluorescence, which indicates barrier leakage.
(B) Live confocal images of FluoZin-3 signal (FIRE LUT) in control and PAK4
inhibiter-treated *Xenopus* embryos. Yellow arrowheads indicate increased signal at
higher-order vertices. Zoomed-in panels (green boxes) highlight barrier leaks at
higher-order vertices. Scale bars, 20 µm (upper), 10 µm (lower). (C) Quantification
of leaky vertices (control: 13 embryos, 1375 vertices; PAK4 inhibitor: 10 embryos,
796 vertices; 3 independent experiments). Error bars, SEM. Data were compared
using a two-tailed unpaired Welch's *t*-test (**, $p < 0.01$). (D) Quantification of
FluoZin-3 intensity at higher-order vertices (4-way and above), normalized to non-
leaky bicellular junctions at 0, 7, and 15 min after the onset of imaging (control: 12
embryos, 20 vertices; PAK4 inhibitor: 11 embryos, 26 vertices; 3 independent
experiments). Violin plots show the median (solid line) and 25th/75th percentiles
(dotted lines), with individual measurements overlaid (circles). Data are compared
using a two-tailed unpaired Student's *t*-test (*, $p < 0.05$, **, $p < 0.01$). (E) Montage
of FluoZin-3 signal (FIRE LUT) at representative higher-order vertices in control
and PAK4 inhibitor-treated embryos, showing leaks (yellow arrowheads). Scale
bars, 5 µm.
1376
Figure 10. PAK4-NT impairs epithelial barrier function at higher-order
vertices in *Xenopus* embryos.
(A) Live confocal images of FluoZin-3 signal (FIRE LUT) in control and PAK4-NT-
expressing *Xenopus* embryos. Yellow arrowheads indicate elevated signal at
higher-order vertices. Zoomed-in panels (green boxes) highlight local barrier leaks.
Scale bars, 20 µm (upper), 10 µm (lower). (B) Percentage of embryos exhibiting
global barrier leaks. n = 12 (control, black), 10 (PAK4-NT, orange). (C)
Quantification of whole-field FluoZin-3 intensity at 0, ~7, and ~15 min after the
onset of imaging (control: 13 embryos; PAK4-NT: 10 embryos; 3 independent
experiments). (D) Quantification of junctional FluoZin-3 intensity at 0, ~7, and ~15
min after the onset of imaging (control: 13 embryos, PAK4-NT: 6 embryos; 3
independent experiments). (E) Quantification of vertices exhibiting barrier leaks
(control: 13 embryos, 1375 vertices; PAK4-NT: 7 embryos, 815 vertices; 3
independent experiments). (F) Quantification of FluoZin-3 intensity at higher-order

vertices (4-way and above), normalized to the non-leaky bicellular junctions
(control: 12 embryos, 20 vertices; PAK4-NT: 6 embryos, 15 vertices; 3 independent
experiments). (G) Montage of FluoZin-3 signal (FIRE-LUT) at representative
higher-order vertices showing repeated and persistent leaks (yellow arrowheads) in
PAK4-NT-expressing embryos. Scale bars, 5 µm. For violin plots (C, D, F), solid
and dotted lines indicate the median and the 25th/75th percentiles. Data were
analyzed using a two-tailed unpaired Student's *t*-test (C, D, F) or Welch's *t*-test (E).
*, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001.
1399
1400
Video legends
Video 1. Time-lapse confocal imaging of gastrula-stage *Xenopus* embryos
showing increased PAK4 localization at remodeling vertex. Video shows PAK4
(PAK4-mNeon, green Fire Blue LUT) intensity at a non-remodeling vertex (left) and
remodeling vertex (right). Yellow arrows show PAK4 accumulation during
remodeling, while red arrows point out the non-remodeling vertex as well as the
end of remodeling. Time interval, 24 sec; video frame rate, 10 fps. Related to Fig.
2F.
1409
Video 2. Time-lapse confocal imaging of gastrula-stage *Xenopus* embryo
showing impaired higher-order vertex resolution remodeling upon PAK4
inhibition. Video illustrates that the PAK4 inhibitor treated embryo (right) has
higher number of rosettes (yellow circules) and these take longer to resolve
compared to control (DMSO treated) embryo (left). Time interval, 34 sec (left video),
32 sec (right video); video frame rate, 15 fps. Related to Fig. 4.
1416
Video 3. Time-lapse confocal imaging showing cytokinetic failure and
junctional discontinuities in PAK4-NT-expressing gastrula-stage *Xenopus*
embryos. First video illustrates that inhibiting the recruitment of endogenous PAK4
via expressing PAK4-NT causes cytokinetic failure. The cell starts dividing marked
by cytokinetic furrow ingression (white arrow). Eventually the daughter-daughter
cell interface breaks (cyan arrow) leading to cytokinesis failure. Time interval, 37
sec; video frame rate, 7 fps. Related to Figure 7D, E. Second video illustrates
junction discontinues in PAK4-NT embryos. The yellow arrowheads show some of
the breaks and discontinues in the junctions. Time interval, 37 sec; video frame
rate, 7 fps. Related to Figure 7F.
1427
Video 4. Time-lapse confocal imaging of gastrula-stage *Xenopus* embryos
showing barrier leaks upon PAK4 inhibition. PAK4 inhibitor-treated embryos
(right) have increased baseline FluoZin3 intensity (FIRE LUT) and an increased
frequency of tight junction leaks compared with controls (left). Time interval, 2:44
(min:sec); video frame rate, 3 fps. Brightness and contrast were adjusted to the
same levels for both videos. Related to Figure 9B.
1434
Video 5. Time-lapse confocal imaging of gastrula-stage *Xenopus* embryos
showing barrier leaks upon PAK4-NT expression. PAK4 NT-expressing
embryos (right) have increased baseline FluoZin3 intensity (FIRE LUT), global
barrier disruption (#2), and an increased frequency of tight junction leaks at

tricellular vertices (#1) compared with controls (left). Time interval, 2:44 (min:sec);
video frame rate, 3 fps. Brightness and contrast were adjusted to the same levels
for all the videos. Related to Figure 10A.

References

Abo, A., J. Qu, M.S. Cammarano, C. Dan, A. Fritsch, V. Baud, B. Belisle, and A. Minden. 1998. PAK4, a novel effector for Cdc42Hs, is implicated in the reorganization of the actin cytoskeleton and in the formation of filopodia. *EMBO J.* 17:6527–6540. doi:10.1093/emboj/17.22.6527.

Abramson, J., J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L. Willmore, A.J. Ballard, J. Bambrick, S.W. Bodenstein, D.A. Evans, C.-C. Hung, M. O'Neill, D. Reiman, K. Tunyasuvunakool, Z. Wu, A. Žemgulytė, E. Arvaniti, C. Beattie, O. Bertolli, A. Bridgland, A. Cherepanov, M. Congreve, A.I. Cowen-Rivers, A. Cowie, M. Figurnov, F.B. Fuchs, H. Gladman, R. Jain, Y.A. Khan, C.M.R. Low, K. Perlin, A. Potapenko, P. Savy, S. Singh, A. Stecula, A. Thillaisundaram, C. Tong, S. Yakneen, E.D. Zhong, M. Zielinski, A. Žídek, V. Bapst, P. Kohli, M. Jaderberg, D. Hassabis, and J.M. Jumper. 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*. 630:493–500. doi:10.1038/s41586-024-07487-w.

Álvarez-Salmoral, D., R. Borza, R. Xie, R.P. Joosten, M.L. Hekkelman, and A. Perrakis. 2024. AlphaBridge: tools for the analysis of predicted macromolecular complexes. doi:10.1101/2024.10.23.619601.

Angulo-Urarte, A., T. van der Wal, and S. Huveneers. 2020. Cell-cell junctions as sensors and transducers of mechanical forces. *Biochim. Biophys. Acta Biomembr.* 1862:183316. doi:10.1016/j.bbamem.2020.183316.

Arnold, T.R., R.E. Stephenson, and A.L. Miller. 2017. Rho GTPases and actomyosin: Partners in regulating epithelial cell-cell junction structure and function. *Exp. Cell Res.* 358:20–30. doi:10.1016/j.yexcr.2017.03.053.

Balda, M.S., and K. Matter. 2023. Tight junctions. *Curr. Biol. CB.* 33:R1135–R1140. doi:10.1016/j.cub.2023.09.027.

Barac, A., J. Basile, J. Vázquez-Prado, Y. Gao, Y. Zheng, and J.S. Gutkind. 2004. Direct interaction of p21-activated kinase 4 with PDZ-RhoGEF, a G protein-linked Rho guanine exchange factor. *J. Biol. Chem.* 279:6182–6189. doi:10.1074/jbc.M309579200.

Bardet, P.-L., B. Guirao, C. Paoletti, F. Serman, V. Léopold, F. Bosveld, Y. Goya, V. Mirouse, F. Graner, and Y. Bellaïche. 2013. PTEN controls junction lengthening

and stability during cell rearrangement in epithelial tissue. *Dev. Cell*. 25:534–546.

doi:10.1016/j.devcel.2013.04.020.

Baskaran, Y., F.P.-L. Tay, E.Y.W. Ng, C.L.F. Swa, S. Wee, J. Gunaratne, and E.

Manser. 2021. Proximity proteomics identifies PAK4 as a component of Afadin-

Nectin junctions. *Nat. Commun*. 12:5315. doi:10.1038/s41467-021-25011-w.

Bertet, C., L. Sulak, and T. Lecuit. 2004. Myosin-dependent junction remodelling

controls planar cell intercalation and axis elongation. *Nature*. 429:667–671.

doi:10.1038/nature02590.

Blankenship, J.T., S.T. Backovic, J.S.P. Sanny, O. Weitz, and J.A. Zallen. 2006.

Multicellular rosette formation links planar cell polarity to tissue morphogenesis.

Dev. Cell. 11:459–470. doi:10.1016/j.devcel.2006.09.007.

Boettner, B., E.E. Govek, J. Cross, and L. Van Aelst. 2000. The junctional

multidomain protein AF-6 is a binding partner of the Rap1A GTPase and

associates with the actin cytoskeletal regulator profilin. *Proc. Natl. Acad. Sci. U. S.*

A. 97:9064–9069. doi:10.1073/pnas.97.16.9064.

Boettner, B., P. Harjes, S. Ishimaru, M. Heke, H.Q. Fan, Y. Qin, L. Van Aelst, and

U. Gaul. 2003. The AF-6 homolog canoe acts as a Rap1 effector during dorsal

closure of the Drosophila embryo. *Genetics*. 165:159–169.

doi:10.1093/genetics/165.1.159.

Callow, M.G., F. Clairvoyant, S. Zhu, B. Schryver, D.B. Whyte, J.R. Bischoff, B.

Jallal, and T. Smeal. 2002. Requirement for PAK4 in the anchorage-independent

growth of human cancer cell lines. *J. Biol. Chem.* 277:550–558.

doi:10.1074/jbc.M105732200.

Campàs, O., I. Noordstra, and A.S. Yap. 2024. Adherens junctions as molecular

regulators of emergent tissue mechanics. *Nat. Rev. Mol. Cell Biol.* 25:252–269.

doi:10.1038/s41580-023-00688-7.

Cavanaugh, K.E., M.F. Staddon, T.A. Chmiel, R. Harmon, S. Budnar, A.S. Yap, S.

Banerjee, and M.L. Gardel. 2022. Force-dependent intercellular adhesion

strengthening underlies asymmetric adherens junction contraction. *Curr. Biol. CB*.

32:1986-2000.e5. doi:10.1016/j.cub.2022.03.024.

Cho, Y., A. Taniguchi, A. Kubo, and J. Ikenouchi. 2025. Rho-ROCK liberates

sequestered claudin for rapid de novo tight junction formation. *eLife*. 13:RP102794.

doi:10.7554/eLife.102794.

Choi, W., B.R. Acharya, G. Peyret, M.-A. Fardin, R.-M. Mège, B. Ladoux, A.S. Yap,

A.S. Fanning, and M. Peifer. 2016. Remodeling the zonula adherens in response

to tension and the role of afadin in this response. *J. Cell Biol.* 213:243–260.

doi:10.1083/jcb.201506115.

Choi, W., D. Goldfarb, F. Yan, M.B. Major, A.S. Fanning, and M. Peifer. 2025.

Proximity proteomics provides a new resource for exploring the function of Afadin

and the complexity of cell-cell adherens junctions. *Biol. Open*. 14:bio061811.

doi:10.1242/bio.061811.

Choi, W., N.J. Harris, K.D. Sumigray, and M. Peifer. 2013. Rap1 and Canoe/afadin

are essential for establishment of apical-basal polarity in the Drosophila embryo.

Mol. Biol. Cell. 24:945–963. doi:10.1091/mbc.E12-10-0736.

Chumki, S.A., L.M. van den Goor, B.N. Hall, and A.L. Miller. 2022. p115RhoGEF

activates RhoA to support tight junction maintenance and remodeling. *Mol. Biol.*

Cell. 33:ar136. doi:10.1091/mbc.E22-06-0205.

Citi, S., M. Fromm, M. Furuse, L. González-Mariscal, A. Nusrat, S. Tsukita, and J.R.

Turner. 2024. A short guide to the tight junction. *J. Cell Sci*. 137:jcs261776.

doi:10.1242/jcs.261776.

Craig, Z., T.R. Arnold, K. Walworth, A. Walkon, and A.L. Miller. 2025. Anillin tunes

contractility and regulates barrier function during Rho flare-mediated tight junction

remodeling. *Mol. Biol. Cell*. 36:ar31. doi:10.1091/mbc.E24-11-0513.

le Duc, Q., Q. Shi, I. Blonk, A. Sonnenberg, N. Wang, D. Leckband, and J. de Rooij.

2010. Vinculin potentiates E-cadherin mechanosensing and is recruited to actin-

anchored sites within adherens junctions in a myosin II-dependent manner. *J. Cell*

Biol. 189:1107–1115. doi:10.1083/jcb.201001149.

Ebrahim, S., T. Fujita, B.A. Millis, E. Kozin, X. Ma, S. Kawamoto, M.A. Baird, M.

Davidson, S. Yonemura, Y. Hisa, M.A. Conti, R.S. Adelstein, H. Sakaguchi, and B.

Kachar. 2013. NMII forms a contractile transcellular sarcomeric network to regulate

apical cell junctions and tissue geometry. *Curr. Biol. CB.* 23:731–736.

doi:10.1016/j.cub.2013.03.039.

Eckert, J., V. Viasnoff, and A.S. Yap. 2025. New directions in epithelial

mechanoadaptation. *Curr. Opin. Cell Biol.* 95:102536.

doi:10.1016/j.ceb.2025.102536.

Elfmann, C., and J. Stülke. 2023. PAE viewer: a webserver for the interactive

visualization of the predicted aligned error for multimer structure predictions and

crosslinks. *Nucleic Acids Res.* 51:W404–W410. doi:10.1093/nar/gkad350.

Esmangart de Bournonville, T., and R. Le Borgne. 2020. Interplay between

Anakonda, Gliotactin, and M6 for Tricellular Junction Assembly and Anchoring of

Septate Junctions in Drosophila Epithelium. *Curr. Biol. CB.* 30:4245-4253.e4.

doi:10.1016/j.cub.2020.07.090.

Fanning, A.S., C.M. Van Itallie, and J.M. Anderson. 2012. Zonula occludens-1 and

-2 regulate apical cell structure and the zonula adherens cytoskeleton in polarized

epithelia. *Mol. Biol. Cell.* 23:577–590. doi:10.1091/mbc.E11-09-0791.

Farquhar, M.G., and G.E. Palade. 1963. Junctional complexes in various epithelia.

J. Cell Biol. 17:375–412. doi:10.1083/jcb.17.2.375.

Finegan, T.M., N. Hervieux, A. Nestor-Bergmann, A.G. Fletcher, G.B. Blanchard,

and B. Sanson. 2019. The tricellular vertex-specific adhesion molecule Sidekick

facilitates polarised cell intercalation during Drosophila axis extension. *PLoS Biol.*

17:e3000522. doi:10.1371/journal.pbio.3000522.

Firmino, J., D. Rocancourt, M. Saadaoui, C. Moreau, and J. Gros. 2016. Cell

Division Drives Epithelial Cell Rearrangements during Gastrulation in Chick. *Dev.*

Cell. 36:249–261. doi:10.1016/j.devcel.2016.01.007.

Founounou, N., N. Loyer, and R. Le Borgne. 2013. Septins regulate the

contractility of the actomyosin ring to enable adherens junction remodeling during

cytokinesis of epithelial cells. *Dev. Cell.* 24:242–255.

doi:10.1016/j.devcel.2013.01.008.

Foxall, E., A. Staszowska, L.M. Hirvonen, M. Georgouli, M. Ciccioli, A. Rimmer, L.

Williams, Y. Calle, V. Sanz-Moreno, S. Cox, G.E. Jones, and C.M. Wells. 2019.

PAK4 Kinase Activity Plays a Crucial Role in the Podosome Ring of Myeloid Cells.

Cell Rep. 29:3385-3393.e6. doi:10.1016/j.celrep.2019.11.016.

Gandhi, D., B.P. Griffin, M.E. Bustillo, F. Chelemen, G. Jang, A. Hairston, M.

Llimargas, and J.E. Treisman. 2025. Mushroom bodies tiny regulates Sidekick

localization to tricellular adherens junctions. *Dev. Camb. Engl.* dev.204777.

doi:10.1242/dev.204777.

Gibson, D.G., L. Young, R.-Y. Chuang, J.C. Venter, C.A. Hutchison, and H.O.

Smith. 2009. Enzymatic assembly of DNA molecules up to several hundred

kilobases. *Nat. Methods.* 6:343–345. doi:10.1038/nmeth.1318.

Gibson, M.C., A.B. Patel, R. Nagpal, and N. Perrimon. 2006. The emergence of

geometric order in proliferating metazoan epithelia. *Nature.* 442:1038–1041.

doi:10.1038/nature05014.

Gong, R., M.J. Reynolds, X. Sun, and G.M. Alushin. 2025. Afadin mediates

cadherin-catenin complex clustering on F-actin linked to cooperative binding and

filament curvature. *Sci. Adv.* 11:eadu0989. doi:10.1126/sciadv.adu0989.

van den Goor, L., J. Iseler, K.M. Koning, and A.L. Miller. 2024. Mechanosensitive

recruitment of Vinculin maintains junction integrity and barrier function at epithelial

tricellular junctions. *Curr. Biol. CB.* 34:4677-4691.e5.

doi:10.1016/j.cub.2024.08.060.

Guillot, C., and T. Lecuit. 2013a. Adhesion disengagement uncouples intrinsic and

extrinsic forces to drive cytokinesis in epithelial tissues. *Dev. Cell.* 24:227–241.

doi:10.1016/j.devcel.2013.01.010.

Guillot, C., and T. Lecuit. 2013b. Mechanics of epithelial tissue homeostasis and

morphogenesis. *Science.* 340:1185–1189. doi:10.1126/science.1235249.

Heisenberg, C.-P., and Y. Bellaïche. 2013. Forces in tissue morphogenesis and

patterning. *Cell.* 153:948–962. doi:10.1016/j.cell.2013.05.008.

Herszterg, S., A. Leibfried, F. Bosveld, C. Martin, and Y. Bellaiche. 2013. Interplay

between the dividing cell and its neighbors regulates adherens junction formation

during cytokinesis in epithelial tissue. *Dev. Cell.* 24:256–270.

doi:10.1016/j.devcel.2012.11.019.

Higashi, T., T.R. Arnold, R.E. Stephenson, K.M. Dinshaw, and A.L. Miller. 2016.

Maintenance of the Epithelial Barrier and Remodeling of Cell-Cell Junctions during

Cytokinesis. *Curr. Biol. CB*. 26:1829–1842. doi:10.1016/j.cub.2016.05.036.

Higashi, T., and H. Chiba. 2020. Molecular organization, regulation and function of

tricellular junctions. *Biochim. Biophys. Acta Biomembr.* 1862:183143.

doi:10.1016/j.bbamem.2019.183143.

Higashi, T., and A.L. Miller. 2017. Tricellular junctions: how to build junctions at the

TRICKiest points of epithelial cells. *Mol. Biol. Cell*. 28:2023–2034.

doi:10.1091/mbc.E16-10-0697.

Higashi, T., A.C. Saito, and H. Chiba. 2024. Damage control of epithelial barrier

function in dynamic environments. *Eur. J. Cell Biol.* 103:151410.

doi:10.1016/j.ejcb.2024.151410.

Higashi, T., A.C. Saito, Y. Fukazawa, M. Furuse, A.Y. Higashi, M. Ono, and H.

Chiba. 2023a. EpCAM proteolysis and release of complexed claudin-7 repair and

maintain the tight junction barrier. *J. Cell Biol.* 222:e202204079.

doi:10.1083/jcb.202204079.

Higashi, T., R.E. Stephenson, and A.L. Miller. 2019. Comprehensive analysis of

formin localization in Xenopus epithelial cells. *Mol. Biol. Cell*. 30:82–95.

doi:10.1091/mbc.E18-02-0133.

Higashi, T., R.E. Stephenson, C. Schwayer, K. Huljev, A.Y. Higashi, C.-P.

Heisenberg, H. Chiba, and A.L. Miller. 2023b. ZnUMBA - a live imaging method to

detect local barrier breaches. *J. Cell Sci.* 136:jcs260668. doi:10.1242/jcs.260668.

Higashi, T., S. Tokuda, S. Kitajiri, S. Masuda, H. Nakamura, Y. Oda, and M.

Furuse. 2013. Analysis of the "angulin" proteins LSR, ILDR1 and ILDR2--tricellulin

recruitment, epithelial barrier function and implication in deafness pathogenesis. *J.*

Cell Sci. 126:966–977. doi:10.1242/jcs.116442.

Hou, A., L.X. Toh, K.H. Gan, K.J.R. Lee, E. Manser, and L. Tong. 2013. Rho

GTPases and regulation of cell migration and polarization in human corneal

epithelial cells. *PloS One*. 8:e77107. doi:10.1371/journal.pone.0077107.

Huang, D.L., N.A. Bax, C.D. Buckley, W.I. Weis, and A.R. Dunn. 2017. Vinculin
forms a directionally asymmetric catch bond with F-actin. *Science*. 357:703–706.
doi:10.1126/science.aan2556.

Huebner, R.J., A.N. Malmi-Kakkada, S. Sarıkaya, S. Weng, D. Thirumalai, and J.B.
Wallingford. 2021. Mechanical heterogeneity along single cell-cell junctions is
driven by lateral clustering of cadherins during vertebrate axis elongation. *eLife*.
10:e65390. doi:10.7554/eLife.65390.

Ikawa, K., S. Ishihara, Y. Tamori, and K. Sugimura. 2023. Attachment and
detachment of cortical myosin regulates cell junction exchange during cell
rearrangement in the Drosophila wing epithelium. *Curr. Biol. CB*. 33:263-275.e4.
doi:10.1016/j.cub.2022.11.067.

Ikeda, W., H. Nakanishi, J. Miyoshi, K. Mandai, H. Ishizaki, M. Tanaka, A. Togawa,
K. Takahashi, H. Nishioka, H. Yoshida, A. Mizoguchi, S. Nishikawa, and Y. Takai.
1999. Afadin: A key molecule essential for structural organization of cell-cell
junctions of polarized epithelia during embryogenesis. *J. Cell Biol*. 146:1117–1132.
doi:10.1083/jcb.146.5.1117.

Ikenouchi, J., M. Furuse, K. Furuse, H. Sasaki, S. Tsukita, and S. Tsukita. 2005.
Tricellulin constitutes a novel barrier at tricellular contacts of epithelial cells. *J. Cell
Biol*. 171:939–945. doi:10.1083/jcb.200510043.

Jensen, C.C., N.J. Gurley, A.J. Mathias, L.R. Wolfsberg, Y. Xiao, Z. Zhou, K.C.
Slep, and M. Peifer. 2025. A key role for Canoe's intrinsically disordered region in
linking cell junctions to the cytoskeleton. *BioRxiv Prepr. Serv. Biol.*
2025.05.10.653259. doi:10.1101/2025.05.10.653259.

Krug, S.M., J.D. Schulzke, and M. Fromm. 2014. Tight junction, selective
permeability, and related diseases. *Semin. Cell Dev. Biol*. 36:166–176.
doi:10.1016/j.semcdb.2014.09.002.

Kuno, S., R. Nakamura, T. Otani, and H. Togashi. 2025. Multivalent afadin
interaction promotes IDR-mediated condensate formation and junctional separation
of epithelial cells. *Cell Rep.* 44:115335. doi:10.1016/j.celrep.2025.115335.

Kuriyama, M., N. Harada, S. Kuroda, T. Yamamoto, M. Nakafuku, A. Iwamatsu, D.
Yamamoto, R. Prasad, C. Croce, E. Canaani, and K. Kaibuchi. 1996. Identification

of AF-6 and canoe as putative targets for Ras. *J. Biol. Chem.* 271:607–610.

doi:10.1074/jbc.271.2.607.

Landino, J., E. Misterovich, L. van den Goor, B. Adhikary, S. Chumki, L.A.

Davidson, and A.L. Miller. 2025. Neighbor cells restrain furrowing during Xenopus

epithelial cytokinesis. *Dev. Cell.* 60:2139-2148.e6.

doi:10.1016/j.devcel.2025.03.010.

Lecuit, T., and A.S. Yap. 2015. E-cadherin junctions as active mechanical

integrators in tissue dynamics. *Nat. Cell Biol.* 17:533–539. doi:10.1038/ncb3136.

Lenne, P.-F., J.-F. Rupprecht, and V. Viasnoff. 2021. Cell Junction Mechanics

beyond the Bounds of Adhesion and Tension. *Dev. Cell.* 56:202–212.

doi:10.1016/j.devcel.2020.12.018.

Letizia, A., D. He, S. Astigarraga, J. Colombelli, V. Hatini, M. Llimargas, and J.E.

Treisman. 2019. Sidekick Is a Key Component of Tricellular Adherens Junctions

that Acts to Resolve Cell Rearrangements. *Dev. Cell.* 50:313-326.e5.

doi:10.1016/j.devcel.2019.07.007.

Linnemann, T., M. Geyer, B.K. Jaitner, C. Block, H.R. Kalbitzer, A. Wittinghofer,

and C. Herrmann. 1999. Thermodynamic and kinetic characterization of the

interaction between the Ras binding domain of AF6 and members of the Ras

subfamily. *J. Biol. Chem.* 274:13556–13562. doi:10.1074/jbc.274.19.13556.

Luissint, A.-C., C.A. Parkos, and A. Nusrat. 2016. Inflammation and the Intestinal

Barrier: Leukocyte-Epithelial Cell Interactions, Cell Junction Remodeling, and

Mucosal Repair. *Gastroenterology.* 151:616–632. doi:10.1053/j.gastro.2016.07.008.

Lye, C.M., H.W. Naylor, and B. Sanson. 2014. Subcellular localisations of the CPTI

collection of YFP-tagged proteins in Drosophila embryos. *Dev. Camb. Engl.*

141:4006–4017. doi:10.1242/dev.111310.

Malin, J., C. Rosa Birriel, S. Astigarraga, J.E. Treisman, and V. Hatini. 2022.

Sidekick dynamically rebalances contractile and protrusive forces to control tissue

morphogenesis. *J. Cell Biol.* 221:e202107035. doi:10.1083/jcb.202107035.

Malin, J., C. Rosa-Birriel, and V. Hatini. 2024. Pten, PI3K, and PtdIns(3,4,5)P3

dynamics control pulsatile actin branching in Drosophila retina morphogenesis.

Dev. Cell. 59:1593-1608.e6. doi:10.1016/j.devcel.2024.03.035.

Mandai, K., H. Nakanishi, A. Satoh, H. Obaishi, M. Wada, H. Nishioka, M. Itoh, A.

Mizoguchi, T. Aoki, T. Fujimoto, Y. Matsuda, S. Tsukita, and Y. Takai. 1997.

Afadin: A novel actin filament-binding protein with one PDZ domain localized at

cadherin-based cell-to-cell adherens junction. *J. Cell Biol.* 139:517–528.

doi:10.1083/jcb.139.2.517.

Mangeol, P., D. Massey-Harroche, M. Sebbagh, F. Richard, A. Le Bivic, and P.-F.

Lenne. 2024. The zonula adherens matura redefines the apical junction of

intestinal epithelia. *Proc. Natl. Acad. Sci. U. S. A.* 121:e2316722121.

doi:10.1073/pnas.2316722121.

Manning, L.A., K.Z. Perez-Vale, K.N. Schaefer, M.T. Sewell, and M. Peifer. 2019.

The Drosophila Afadin and ZO-1 homologues Canoe and Polychaetoid act in

parallel to maintain epithelial integrity when challenged by adherens junction

remodeling. *Mol. Biol. Cell.* 30:1938–1960. doi:10.1091/mbc.E19-04-0209.

Masuda, S., Y. Oda, H. Sasaki, J. Ikenouchi, T. Higashi, M. Akashi, E. Nishi, and M.

Furuse. 2011. LSR defines cell corners for tricellular tight junction formation in

epithelial cells. *J. Cell Sci.* 124:548–555. doi:10.1242/jcs.072058.

McParland, E.D., N.J. Gurley, L.R. Wolfsberg, T.A. Butcher, A. Bhattarai, C.C.

Jensen, R.I. Johnson, K.C. Slep, and M. Peifer. 2024. The dual Ras-association

domains of Drosophila Canoe have differential roles in linking cell junctions to the

cytoskeleton during morphogenesis. *J. Cell Sci.* 137:jcs263546.

doi:10.1242/jcs.263546.

Meng, W., and M. Takeichi. 2009. Adherens junction: molecular architecture and

regulation. *Cold Spring Harb. Perspect. Biol.* 1:a002899.

doi:10.1101/cshperspect.a002899.

Mira-Osuna, M., and R. Le Borgne. 2024. Assembly, dynamics and remodeling of

epithelial cell junctions throughout development. *Dev. Camb. Engl.* 151:dev201086.

doi:10.1242/dev.201086.

Miyazaki, S., T. Otani, K. Sugihara, T. Fujimori, M. Furuse, and T. Miura. 2023.

Mechanism of interdigitation formation at apical boundary of MDCK cell. *iScience.*

26:106594. doi:10.1016/j.isci.2023.106594.

Nguyen, T.P., T. Otani, M. Tsutsumi, N. Kinoshita, S. Fujiwara, T. Nemoto, T. Fujimori, and M. Furuse. 2024. Tight junction membrane proteins regulate the mechanical resistance of the apical junctional complex. *J. Cell Biol.* 223:e202307104. doi:10.1083/jcb.202307104.

Nicholas, N.S., A. Pipili, M.S. Lesjak, S.M. Ameer-Beg, J.L.C. Geh, C. Healy, A.D. MacKenzie Ross, M. Parsons, F.O. Nestle, K.E. Lacy, and C.M. Wells. 2016. PAK4 suppresses PDZ-RhoGEF activity to drive invadopodia maturation in melanoma cells. *Oncotarget.* 7:70881–70897. doi:10.18632/oncotarget.12282.

Oda, Y., T. Otani, J. Ikenouchi, and M. Furuse. 2014. Tricellulin regulates junctional tension of epithelial cells at tricellular contacts through Cdc42. *J. Cell Sci.* 127:4201–4212. doi:10.1242/jcs.150607.

Otani, T., and M. Furuse. 2020. Tight Junction Structure and Function Revisited. *Trends Cell Biol.* 30:805–817. doi:10.1016/j.tcb.2020.08.004.

Otani, T., T.P. Nguyen, S. Tokuda, K. Sugihara, T. Sugawara, K. Furuse, T. Miura, K. Ebnet, and M. Furuse. 2019. Claudins and JAM-A coordinately regulate tight junction formation and epithelial polarity. *J. Cell Biol.* 218:3372–3396. doi:10.1083/jcb.201812157.

Paré, A.C., and J.A. Zallen. 2020. Cellular, molecular, and biophysical control of epithelial cell intercalation. *Curr. Top. Dev. Biol.* 136:167–193. doi:10.1016/bs.ctdb.2019.11.014.

Perez-Vale, K.Z., and M. Peifer. 2020. Orchestrating morphogenesis: building the body plan by cell shape changes and movements. *Dev. Camb. Engl.* 147:dev191049. doi:10.1242/dev.191049.

Pokutta, S., F. Drees, Y. Takai, W.J. Nelson, and W.I. Weis. 2002. Biochemical and structural definition of the l-afadin- and actin-binding sites of alpha-catenin. *J. Biol. Chem.* 277:18868–18874. doi:10.1074/jbc.M201463200.

Qu, J., X. Li, B.G. Novitch, Y. Zheng, M. Kohn, J.-M. Xie, S. Kozinn, R. Bronson, A.A. Beg, and A. Minden. 2003. PAK4 kinase is essential for embryonic viability and for proper neuronal development. *Mol. Cell. Biol.* 23:7122–7133. doi:10.1128/MCB.23.20.7122-7133.2003.

Razzell, W., M.E. Bustillo, and J.A. Zallen. 2018. The force-sensitive protein Ajuba

regulates cell adhesion during epithelial morphogenesis. *J. Cell Biol.* 217:3715–

3730. doi:10.1083/jcb.201801171.

Reyes, C.C., M. Jin, E.B. Breznau, R. Espino, R. Delgado-Gonzalo, A.B.

Goryachev, and A.L. Miller. 2014. Anillin regulates cell-cell junction integrity by

organizing junctional accumulation of Rho-GTP and actomyosin. *Curr. Biol. CB.*

24:1263–1270. doi:10.1016/j.cub.2014.04.021.

Rübsam, M., J.A. Broussard, S.A. Wickström, O. Nekrasova, K.J. Green, and C.M.

Niessen. 2018. Adherens Junctions and Desmosomes Coordinate Mechanics and

Signaling to Orchestrate Tissue Morphogenesis and Function: An Evolutionary

Perspective. *Cold Spring Harb. Perspect. Biol.* 10:a029207.

doi:10.1101/cshperspect.a029207.

Saha, K., Y. Zhou, and J.R. Turner. 2025. Tight junction regulation, intestinal

permeability, and mucosal immunity in gastrointestinal health and disease. *Curr.*

Opin. Gastroenterol. 41:46–53. doi:10.1097/MOG.0000000000001066.

Saito, A.C., T. Higashi, and H. Chiba. 2024. Protocol for establishing knockout cell

clones by deletion of a large gene fragment using CRISPR-Cas9 with multiple

guide RNAs. *STAR Protoc.* 5:103179. doi:10.1016/j.xpro.2024.103179.

Saito, A.C., T. Higashi, Y. Fukazawa, T. Otani, M. Tauchi, A.Y. Higashi, M. Furuse,

and H. Chiba. 2021. Occludin and tricellulin facilitate formation of anastomosing

tight-junction strand network to improve barrier function. *Mol. Biol. Cell.* 32:722–

738. doi:10.1091/mbc.E20-07-0464.

Sakakibara, S., T. Maruo, M. Miyata, K. Mizutani, and Y. Takai. 2018. Requirement

of the F-actin-binding activity of l-afadin for enhancing the formation of adherens

and tight junctions. *Genes Cells Devoted Mol. Cell. Mech.* 23:185–199.

doi:10.1111/gtc.12566.

Sakakibara, S., K. Mizutani, A. Sugiura, A. Sakane, T. Sasaki, S. Yonemura, and Y.

Takai. 2020. Afadin regulates actomyosin organization through αE-catenin at

adherens junctions. *J. Cell Biol.* 219:e201907079. doi:10.1083/jcb.201907079.

Sawyer, J.K., N.J. Harris, K.C. Slep, U. Gaul, and M. Peifer. 2009. The Drosophila

afadin homologue Canoe regulates linkage of the actin cytoskeleton to adherens

junctions during apical constriction. *J. Cell Biol.* 186:57–73.

doi:10.1083/jcb.200904001.

Selamat, W., P.-L.F. Tay, Y. Baskaran, and E. Manser. 2015. The Cdc42 Effector

Kinase PAK4 Localizes to Cell-Cell Junctions and Contributes to Establishing Cell

Polarity. *PloS One.* 10:e0129634. doi:10.1371/journal.pone.0129634.

Sells, M.A., U.G. Knaus, S. Bagrodia, D.M. Ambrose, G.M. Bokoch, and J.

Chernoff. 1997. Human p21-activated kinase (Pak1) regulates actin organization in

mammalian cells. *Curr. Biol. CB.* 7:202–210. doi:10.1016/s0960-9822(97)70091-5.

Spadaro, D., S. Le, T. Laroche, I. Mean, L. Jond, J. Yan, and S. Citi. 2017.

Tension-Dependent Stretching Activates ZO-1 to Control the Junctional

Localization of Its Interactors. *Curr. Biol. CB.* 27:3783-3795.e8.

doi:10.1016/j.cub.2017.11.014.

Stephenson, R.E., T. Higashi, I.S. Erofeev, T.R. Arnold, M. Leda, A.B. Goryachev,

and A.L. Miller. 2019. Rho Flares Repair Local Tight Junction Leaks. *Dev. Cell.*

48:445-459.e5. doi:10.1016/j.devcel.2019.01.016.

Sugawara, T., K. Furuse, T. Otani, T. Wakayama, and M. Furuse. 2021. Angulin-1

seals tricellular contacts independently of tricellulin and claudins. *J. Cell Biol.*

220:e202005062. doi:10.1083/jcb.202005062.

Tachibana, K., H. Nakanishi, K. Mandai, K. Ozaki, W. Ikeda, Y. Yamamoto, A.

Nagafuchi, S. Tsukita, and Y. Takai. 2000. Two cell adhesion molecules, nectin

and cadherin, interact through their cytoplasmic domain-associated proteins. *J.*

Cell Biol. 150:1161–1176. doi:10.1083/jcb.150.5.1161.

Takahashi, K., H. Nakanishi, M. Miyahara, K. Mandai, K. Satoh, A. Satoh, H.

Nishioka, J. Aoki, A. Nomoto, A. Mizoguchi, and Y. Takai. 1999. Nectin/PRR: an

immunoglobulin-like cell adhesion molecule recruited to cadherin-based adherens

junctions through interaction with Afadin, a PDZ domain-containing protein. *J. Cell*

Biol. 145:539–549. doi:10.1083/jcb.145.3.539.

Tang, V.W. 2018. Cell-cell adhesion interface: orthogonal and parallel forces from

contraction, protrusion, and retraction. *F1000Research.* 7:F1000 Faculty Rev-1544.

doi:10.12688/f1000research.15860.1.

Tokuda, S., T. Higashi, and M. Furuse. 2014. ZO-1 knockout by TALEN-mediated

gene targeting in MDCK cells: involvement of ZO-1 in the regulation of

cytoskeleton and cell shape. *PloS One*. 9:e104994.

doi:10.1371/journal.pone.0104994.

Uechi, H., and E. Kuranaga. 2019. The Tricellular Junction Protein Sidekick

Regulates Vertex Dynamics to Promote Bicellular Junction Extension. *Dev. Cell*.

50:327-338.e5. doi:10.1016/j.devcel.2019.06.017.

Van Itallie, C.M., and J.M. Anderson. 2014. Architecture of tight junctions and

principles of molecular composition. *Semin. Cell Dev. Biol.* 36:157–165.

doi:10.1016/j.semcdb.2014.08.011.

Vanderleest, T.E., C.M. Smits, Y. Xie, C.E. Jewett, J.T. Blankenship, and D.

Loerke. 2018. Vertex sliding drives intercalation by radial coupling of adhesion and

actomyosin networks during Drosophila germband extension. *eLife*. 7:e34586.

doi:10.7554/eLife.34586.

Varadarajan, S., S.A. Chumki, R.E. Stephenson, E.R. Misterovich, J.L. Wu, C.E.

Dudley, I.S. Erofeev, A.B. Goryachev, and A.L. Miller. 2022. Mechanosensitive

calcium flashes promote sustained RhoA activation during tight junction remodeling.

J. Cell Biol. 221:e202105107. doi:10.1083/jcb.202105107.

Varadarajan, S., R.E. Stephenson, and A.L. Miller. 2019. Multiscale dynamics of

tight junction remodeling. *J. Cell Sci.* 132:jcs229286. doi:10.1242/jcs.229286.

Wallace, S.W., J. Durgan, D. Jin, and A. Hall. 2010. Cdc42 regulates apical

junction formation in human bronchial epithelial cells through PAK4 and Par6B. *Mol.*

Biol. Cell. 21:2996–3006. doi:10.1091/mbc.E10-05-0429.

Weaire, D., and N. Rivier. 1984. Soap, cells and statistics—random patterns in two

dimensions. *Contemp. Phys.* 25:59–99. doi:10.1080/00107518408210979.

Wittek, A., M. Hollmann, R. Schleutker, and S. Luschnig. 2020. The

Transmembrane Proteins M6 and Anakonda Cooperate to Initiate Tricellular

Junction Assembly in Epithelia of Drosophila. *Curr. Biol. CB*. 30:4254-4262.e5.

doi:10.1016/j.cub.2020.08.003.

Wohlgemuth, S., C. Kiel, A. Krämer, L. Serrano, F. Wittinghofer, and C. Herrmann.

2005. Recognizing and defining true Ras binding domains I: biochemical analysis.

J. Mol. Biol. 348:741–758. doi:10.1016/j.jmb.2005.02.048.

Yamamoto, K., H. Miura, M. Ishida, Y. Mii, N. Kinoshita, S. Takada, N. Ueno, S.

Sawai, Y. Kondo, and K. Aoki. 2021. Optogenetic relaxation of actomyosin

contractility uncovers mechanistic roles of cortical tension during cytokinesis. *Nat.*

Commun. 12:7145. doi:10.1038/s41467-021-27458-3.

Yamazaki, Y., K. Umeda, M. Wada, S. Nada, M. Okada, S. Tsukita, and S. Tsukita.

2008. ZO-1- and ZO-2-dependent integration of myosin-2 to epithelial zonula

adherens. *Mol. Biol. Cell.* 19:3801–3811. doi:10.1091/mbc.e08-04-0352.

Yonemura, S. 2011. Cadherin-actin interactions at adherens junctions. *Curr. Opin.*

Cell Biol. 23:515–522. doi:10.1016/j.ceb.2011.07.001.

Yonemura, S., Y. Wada, T. Watanabe, A. Nagafuchi, and M. Shibata. 2010. alpha-

Catenin as a tension transducer that induces adherens junction development. *Nat.*

Cell Biol. 12:533–542. doi:10.1038/ncb2055.

Yu, H.H., and J.A. Zallen. 2020. Abl and Canoe/Afadin mediate

mechanotransduction at tricellular junctions. *Science.* 370:eaba5528.

doi:10.1126/science.aba5528.

Zhang, S., J. Han, M.A. Sells, J. Chernoff, U.G. Knaus, R.J. Ulevitch, and G.M.

Bokoch. 1995. Rho family GTPases regulate p38 mitogen-activated protein kinase

through the downstream mediator Pak1. *J. Biol. Chem.* 270:23934–23936.

doi:10.1074/jbc.270.41.23934.

1857

Fig. 1

Fig. 2

Fig. 3

A MDCK II (*cldn2*-KO) cells

PAK4 inhibitor

DMSO   500 nM   5 µM

ZO-1

DAPI
ZO-1

20 µm

B
Total vertices / total cells

**** p = 2.33e-05
ns

DMSO   500 nM   5 µM
PAK4 inhibitor

C
Ratio of vertices (%)

4-way   rosettes
✕   ✶ ✹

** p = 0.0035
*** p = 1.57e-09
ns
* p = 0.047

DMSO   500 nM   5 µM
PAK4 inhibitor

D
Junction discontinuities / cell

ns
ns

DMSO   500 nM   5 µM
PAK4 inhibitor

Fig. 4

A *Xenopus* gastrula-stage embryo

ZO-1 Angulin-1 F-actin ZO-1 Angulin-1 F-actin

DMSO

PAK4 inhibitor

20 μm

B Resolution of 4-way vertex post-cytokinesis

Furrowing starts

~13 min

Cytokinesis completes

60 min

Type I   Type II   Type III

C

00:00 13:00 60:00

Control

PAK4 inhibitor

20 μm

ZO-1

D Cell packing post-cytokinesis

Type I
Type II
Type III

Control
PAK4 inhibitor

Time post-furrow ingression begins (min)

E

Rosette resolution

ZO-1

Control

00:00 30:00 43:49

PAK4 inhibitor embryo #1

00:00 30:48 78:15

PAK4 inhibitor embryo #2

00:00 30:00 53:42

20 μm

F

Control
PAK4 inhibitor

% of Rosettes resolved

Time (min)

No resolution

G Time taken for rosette resolution

** p=0.0037

Time (min)

Control   PAK4 Inhibitor

Fig. 5

A MDCK II (*cldn2*-KO) cells

Control   *PAK4*-KO#1   *PAK4*-KO#2

ZO-1

DAPI ZO-1

20 µm

B

Total vertices / total cells

**** p = 1.23e-14
**** p = 6.11e-15

Ctrl   #1   #2
PAK4-KO

C

Ratio of vertices (%)

4-way ✕
rosettes ✭ ✶

ns
ns
** p = 0.0024
* p = 0.0210

Ctrl   #1   #2
PAK4-KO

D

Junction discontinuities / cell

**** p = 1.44e-10
**** p = 2.28e-11

Ctrl   #1   #2
PAK4-KO

E

Control   *PAK4*-KO#1   *PAK4*-KO#2

MyoIIB

MyoIIB ZO-1 DAPI

20 µm

Fig. 6

Fig. 7

A

PAK4 WT — CRIB/PBD, Predicted Afadin-interacting region, Kinase

PAK4 N-Term (NT)

B *Xenopus* gastrula-stage embryo

Control #1 | Control #2 | PAK4-NT #1 | PAK4-NT #2

ZO-1

50 µm (each)

25 µm (each inset)

C

Ratio of vertices/embryo

*** p=0.0003 (Tricellular vertices)
*** p=0.0001 (4-way vertices)
** p=0.0011 (Rosettes)

Control, PAK4-NT

D PAK4-NT causes cytokinetic defects

ZO-1 F-actin

Control: 00:00, 04:52, 08:33, 18:52 — 25 µm

PAK4-NT #1: 00:00, 19:29, 30:27, 57:15 — 25 µm
Cytokinesis failure

PAK4-NT #2: 00:00, 06:33, 15:16, 17:27 — 25 µm
Asymmetric cell division

E

Percentage of cytokinetic events

Control: 45, 1
PAK4-NT: 17, 15

Cytokinesis failure
Cytokinesis completed

F PAK4-NT causes junction breaks

Example #1: ZO-1 — 00:00, 05:29, 15:50
F-actin ZO-1

Example #2: ZO-1 — 00:00, 04:59, 09:58
F-actin ZO-1 — 25 µm

Fig. 8

Fig. 9

Fig. 10

A *Xenopus* gastrula-stage embryo

Control · FluoZin3 · 20 μm
PAK4-NT #1 · FluoZin3 · 20 μm
PAK4-NT #2 · FluoZin3 · 20 μm

10 μm (×6)

High / Low

B Global barrier leak

embryos with global barrier leak (%)

Control · PAK4-NT

C Whole-field FZ3 intensity

Control / PAK4-NT

Raw intensity (A.U.)

** p=0.0084
* p=0.0238
* p=0.0160

Time (min): 0 · ~7 · ~15

D Junctional FZ3 intensity

Control / PAK4-NT

Normalized intensity (A.U.) (normalized to-whole-field intensity)

** p=0.0071
** p=0.0047
** p=0.0016

Time (min): 0 · ~7 · ~15

E Leaky vertices

** p=0.0020

vertices that experience leaks (%)

Control · PAK4-NT

F FZ3 intensity at higher order vertices

Control / PAK4-NT

Normalized intensity (A.U.) (normalized to non-leaky BCJ)

*** p=2.22e-04
*** p=1.17e-04
**** p=1.42e-07

Time (min): 0 · ~7 · ~15

G Impaired barrier repair in embryos expressing PAK4-NT

Time (min:sec): 00:00 · 02:42 · 05:23 · 08:05 · 10:47 · 13:29

Control · FluoZin3 · 5 μm

Persistent leak · PAK4-NT embryo #1 · FluoZin3 · 5 μm

Repeated leak · PAK4-NT embryo #2 · FluoZin3 · 5 μm

High / Low