1   **Dual Function of Mitochondrial Complex III in *Plasmodium falciparum***

2   River S. Rell, Anurag Shukla, Joanne M. Morrisey, Ijeoma C. Okoye, Michael W. Mather, and Akhil

3   B. Vaidya

4   Center for Molecular Parasitology, Department of Microbiology and Immunology, Drexel

5   University College of Medicine, Philadelphia, PA 19129

Abstract

Complex III of the malaria parasite mitochondrial electron transport chain (mtETC) has been validated as an attractive target for currently used antimalarials. We previously showed that the main function of mtETC in blood stage *Plasmodium falciparum* is to regenerate ubiquinone, which serves as an obligatory co-substrate of dihydroorotate dehydrogenase (DHOD), an essential mitochondrial enzyme for pyrimidine biosynthesis. *P. falciparum* can be rendered resistant to all mtETC inhibitors by provision of a bypass mediated by cytosolic yeast DHOD, a fumarate-reducing enzyme. Malaria parasite mitochondrial DNA (mtDNA) encodes only 3 proteins, each a component of mtETC. However, attempts to eliminate mtDNA in transgenic parasites expressing yDHOD have been unsuccessful, suggesting the possibility that essential function(s) other than the canonical redox reactions of the mtETC also require mtDNA maintenance. Here we have tested the hypothesis that Complex III serves the dual functions of processing imported mitochondrial proteins, as well as ubiquinone regeneration. We have generated transgenic lines that conditionally express mitochondrial processing peptidase a (MPPα), which is also a component of Complex III. Using these parasites, we have determined that MPPα is essential even when the need for mitochondrial electron transport is bypassed. MPPα knockdown also resulted in hypersensitivity of the parasites to proguanil, a drug that synergizes with mtETC inhibitors such as atovaquone. Pulldown with MPPα followed by proteomics revealed the association of multiple mitochondrially targeted proteins, in addition to all components of Complex III. These results are consistent with the suggestion that Complex III in *P. falciparum* serves both mtETC and protein processing functions in mitochondrial physiology.

Introduction

Mitochondrial Complex III in Apicomplexan parasites is a clinically validated drug target, and inhibitors of it are in therapeutic use. Divergence of the parasite Complex III from the host underlies the selectivity of these potent drugs (1, 2). Working through a modified Q cycle (3, 4), Complex III catalyzes the transfer of electrons from ubiquinol to cytochrome *c* while transporting protons into the mitochondrial intermembrane space. Inhibition of this step in mtETC results in blockage of all mitochondrial dehydrogenases that require ubiquinone as the electron acceptor. In previous studies, we have shown that in the blood stage *P. falciparum* the only critical dehydrogenase for parasite survival is dihydroorotate dehydrogenase (DHOD), an essential enzyme that is required for pyrimidine biosynthesis by the parasites (5). When the parasites are genetically modified to express cytosolic yeast DHOD (yDHOD), they are rendered independent of mtETC, since yDHOD uses fumarate as the electron acceptor, rather than ubiquinone (5). This metabolic bypass rendered the parasites resistant to all mtETC inhibitors such as atovaquone. Mitochondrial DNA (mtDNA) in malaria parasites is highly reduced in size and encodes only three proteins, each a component of mtETC (cytochrome *b* and subunits I and III of cytochrome c oxidase) (1, 6, 7). Therefore, we reasoned that it should be possible to eliminate mtDNA from mtETC-independent parasites expressing yDHOD. However, our multiple attempts to achieve this by disrupting genes encoding mitochondrial functions required for mtDNA maintenance and mitochondrial translation have failed in transgenic parasites expressing yDHOD (8-10). Thus, it is likely that the mtETC complexes serve one or more functions in addition to respiration in blood stage malaria parasites. At present the nature of these functions remains unknown.

Except for the 3 proteins encoded by the parasite mtDNA, all mitochondrial proteins are

encoded in the nuclear genome, translated in the cytoplasm and imported into the

mitochondrion. Many of these proteins contain targeting peptides at their N-termini. These

peptides are cleaved off by mitochondrial processing peptidase (MPP) (11, 12). MPPs in

mammalian and yeast mitochondria consist of heterodimers of α and β subunits that are

localized within the mitochondrial matrix (13, 14). The catalytic sites are present within the β

subunit, which is a metalloprotease with a Zn^{2+} ion at the active site (15). The α subunit does

not have catalytic property but is essential for substrate recognition and proper presentation of

the peptide within the catalytic site in multiple steps (16, 17). In addition to the matrix-localized

MPP in fungi and mammals, Complex III in eukaryotes also contain Core1 and Core2 subunits

that have high sequence homology to the MPP subunits α and B, respectively (17-19). These

subunits in yeast have lost their catalytic activity, whereas in mammals they serve to process

just one protein, the Rieske 2Fe2S subunit of Complex III (20, 21). Hence, the majority of

mitochondrial proteins in these organisms have to be processed by the matrix located MPP. In

stark contrast, plant mitochondria do not contain matrix located MPPs, which led Braun to

suggest that Complex III associated Core proteins fulfilled MPP function, as later confirmed, and

thus Complex III has dual functions in plants: ubiquinol oxidation to serve mtETC and processing

of mitochondrially targeted proteins (22, 23). We hypothesize that in malaria parasites correct

assembly of Complex III is essential not just for its function in mtETC but also for processing

most imported mitochondrial proteins. Thus, even when mtETC has been rendered redundant,

Complex III would still need to be assembled to process mitochondrial proteins. In this study we

have used conditional expression of MPPα to investigate the potential dual role of Complex III in

blood stage *P. falciparum*.

Materials and Methods

P. falciparum **culture and transfection**

We used *P. falciparum* D10_VP2KO, in which the pfVP2 locus encoding a type 2 H^+ pumping

pyrophosphatase was disrupted with the yDHOD selectable maker, for transfections (24). These

parasites are resistant to all mtETC inhibitors, as well as inhibitors of parasite DHOD.

D10_VP2KO and Dd2attb asexual blood stage parasites were cultured in human (O+) RBC, at

2.5% hematocrit in RPMI1640 medium supplemented with 0.5% Albumax, 2 g/liter $NaHCO_3$, 15

mM HEPES pH 7.4 , 1 mM hypoxanthine and 50 mg/liter gentamicin in 90% N_2, 5% O_2 and 5%

CO_2. Testing for Mycoplasma was done on a regular basis to ensure the cultures were free of

any such infection.

For transfection, parasite cultures with 5-7% ring stage parasites were washed three

times with Cytomix. The parasites were resuspended to 50% hematocrit in Cytomix. 50 µg of

linearized pMG75 MPPα plasmid isolated using a Qiagen Maxikit (Qiagen, Valencia, CA, USA)

and precipitated in ethanol was centrifuged down and resuspended in 100 µl Cytomix. The

plasmid and 250 µl of the parasite cytomix suspension were combined and transferred to a 0.2

cm electroporation cuvette on ice. Electroporations were carried out using a BioRad GenePulser

set at 0.31kV, 960uF. The electroporated cells were immediately transferred to a T-25 flask

containing 0.2 ml uninfected 50% RBCs and 10 ml medium. Parasites were selected using 2.5

µg/ml Blasticidin and 250 nM anhydrotetracycline (aTc).

Transfection vector generation

Optimal gRNA sequences were identified using CRISPOR software to generate a double strand

break within the endogenous MPPα locus (25). gRNA sequences were cloned into a Cas9-sgRNA

vector (26)(S1A and B).

pMG75 (27) plasmid backbone was used to construct the repair template vector used to help

generate D10_VP2KO_MPPα, the MPPα conditional knockdown line in the D10_VP2KO

background. (24). A synthetic gene block was ordered from GenScript containing 584 base pair

MPPα 5' homologous region (nucleotide 1591-2175) in the coding sequence, containing silent

blocking mutations in the selected gRNA recognition site, and 550 base pair 3'homologous

region (nucleotide 2178-2728) in the 3'UTR, separated by a linker region containing restriction

enzyme sites for SacII and BspEII for linearization of the plasmid. The homologous regions are

flanked by BstEII and BssHII for ligation into the pMG75 backbone, which contains the TetR-

DOZI-aptamer regulatory system, as well as a BSD selectable marker. The synthetic gene was

inserted into the pMG75 plasmid backbone via T4 DNA ligase. All plasmids were sequenced

using Plasmidsaurus (South San Francisco) nanopore sequencing. The plasmid was linearized

with SacII and BspEII before transfection.

114 **Growth Assay**

115 Parasites were enriched for late trophozoite and early schizont stages using a layered step

116 gradient of 70%, 60%, and 35% Percoll. Equal amounts of synchronized parasites were split into

117 a 6 well plate containing Human (O+) RBC at 2.5% hematocrit in 10ml of complete RPMI

118 medium. Triplicate cultures were grown in the presence or absence of aTc, with daily medium

119 change. Cumulative parasitemias were counted over the span of 8 days using flow cytometry

120 after staining with SYBR Green. Cumulative parasitemias were plotted using GraphPad Prism.

121 **Immunofluorescence assays.**

122 50 µL of packed volume parasite cultures were fixed with 4% v/v paraformaldehyde and

123 0.0075% v/v glutaraldehyde with slow mixing in a rotor for one hour at room temperature. The

124 cells were washed 3x with PBS, followed by permeabilization with 0.125% v/v Triton X-100 in

125 PBS. The cells were treated with $NaBH_4$ (0.1 mg/ml) for five minutes at room temperature while

126 mixing on a rotor. Samples were washed twice with 1X PBS and then blocked with 3% w/v BSA

127 for 1 hour. Anti-HA (Lot#12923, Santa Cruz Biotechnology) and anti HSP60 (Lot# PP487392

128 Novus Bio) antibodies were added to the parasites at 1:500 dilution, and the samples were

129 incubated overnight at 4°C in a rotor. Following three PBS washes, Alexa Fluor 488 conjugated

130 goat anti-mouse IgG (Invitrogen Lot# SG251135) and Alexa Flour 568 conjugated goat anti-

131 rabbit (Invitrogen lot #2447870) secondary antibodies were added at 1:250 dilution, followed

132 by overnight incubation at 4°C. After three PBS washes, the treated parasite samples were

133 resuspended in anti-fade buffer (S2828, ThermoFisher Scientific) and imaged using a Nikon Ti

134 microscope.

Live Cell Microscopy

35 mm glass bottom culture dishes were coated with 0.1% poly-D-lysine overnight and washed

3 times with PBS. 250 µL of 2.5% hematocrit parasite culture was added to the dishes and

incubated for 30 min to allow attachment. Culture dishes were washed 3 times with 1XPBS

followed by addition of phenol red-free RPMI. Parasites were stained with 62.5 nM

MitoTracker Red (Life Technologies by ThermoFisher Scientific) for 20 minutes, followed by 0.5

mg/ml Hoechst nuclear staining for 10min. Imaging was done using the Nikon Ti microscope.

Parasites were temperature controlled at 37 ℃. Hoechst-stained parasites were visualized

with the DAPI filter set, while MitoTracker red was visualized using the tetramethyl rhodamine

isocyanate (TRITC) filter set. Images were 2-dimentionally deconvoluted and processed using

Nikon NIS Elements (5.30.02) Imaging Software. The presence of mitochondrial membrane

potential or parasite plasma membrane potential was visualized following incubation with

MitoTracker red. For statistical analysis of the membrane potential visualization, 30 parasites

were counted and imaged. The results were graphed using GraphPad Prism.

Western blotting

 Saponin lysed parasite pellets were suspended in approximately ten volumes of 2% w/v SDS,

5% β-mercaptoethanol, and 1% bromophenol blue. Samples were centrifuged for 5 minutes at

13,000g, and the supernatant was subjected to SDS-PAGE. The separated proteins on the gel

were transferred to PVDM membranes at 20 volts overnight. Powdered milk solution was used

to block the membranes. Blots were then incubated at a dilution ratio of 1:10,000 with primary

antibody. After probing with primary antibody (Mouse monoclonal anti-HA Lot#12923, Santa

Cruz Biotechnology, Rabbit anti- *Plasmodium* aldolase Lot#GR289448-4 ABCAM, Rabbit anti

HSP60 Lot# PP487392 Novus Bio) blots were washed with Tris-buffered saline with tween

(TBST) and incubated with secondary antibody (Goat anti mouse IgG-HRP Lot#GR3219575-4

ABCAM, Mouse anti rabbit IgG-HRP Lot#L1819 Santa Cruz Biotechnology) conjugated with

horseradish peroxidase at a concentration of 1:10,000.

Immunopulldown and Proteomics

100 µL parasite pellets from both wildtype and D10_VP2KO_MPPα were solubilized overnight

at 4^{o}C in solubilization buffer containing 2% digitonin, 200 mM 6- aminocaproic acid, 50 mM

Bis-tris, pH 7, 1 mM EDTA, 1 mM AEBSF, and protease inhibitor cocktail (Sigma-Aldrich). The

solubilized sample was centrifuged at 14,000 rpm for 15 min at 4 °C to remove debris, and the

supernatant was incubated with anti HA antibody coated magnetic Dynabeads (Pierce Anti-HA,

Clone 2-2.2.14, Clone 9E10; Thermo Fisher Scientific). Beads were washed 3 times in 1x PBS,

and solubilized total protein was added to 50 µl of the beads, followed by overnight mixing by

rotation at 4 °C. Protein-bound beads were washed three times with 1x PBS + 0.02% digitonin.

Beads with bound proteins were used for LC/MS on bead proteomics at the University of South

Florida Mass Spectrometry & Proteomic Core Facility. Protein samples were prepared for LC-

MS/MS using S-Traps (Protifi). 5% SDS in 50 mM triethylammonium bicarbonate (TEAB) was

added to the IP beads. Protein samples were reduced with 20 mM dithiothreitol (DTT) at 95 °C

for 10 min and alkylated using 40 mM iodoacetamide (IAA) in the dark at room temperature for

30 min. The samples were acidified with 12% phosphoric acid (final concentration 1.2%) before

adding 6x volumes of S-trap buffer (90% Methanol, 100 mM TEAB). The protein solution was

loaded onto micro S-traps, and centrifuged at 4000xg for 30 sec. Three washes with 150 µl S-

trap buffer were performed. 0.5 µg of Trypsin/LysC (Promega) was added to the S-trap filter

and digested overnight (~16 hrs) at 37 °C. Peptides were eluted with sequential additions of 50

mM TEAB, water+0.1% formic acid, and 50/50 acetonitrile/water+0.1% formic acid, centrifuged

at 4000xg for 1 min each. Peptides were completely dried in a vacuum-centrifuge concentrator

before resuspension in water+0.1% formic acid.

Peptides were separated using a Vanquish Neo UHPLC (Thermo) with a 50 cm EASY-Spray

PepMap Neo C18 column (Thermo) and analyzed on a Q Exactive Plus hybrid quadrupole mass

spectrometer (Thermo). Peptides were separated over a 1hr gradient (Mobile Phase A:

water+0.1% Formic Acid, Mobile Phase B: Acetonitrile+0.1% Formic Acid; 2% B-32% B), and data

was acquired using data-dependent acquisition (DDA) with a Top 30 method. Full MS scans

were acquired at 70k resolution, MS/MS scans were acquired at 17.5k resolution. Raw data files

were searched using MaxQuant (v2.4.9) against the current *Plasmodium falciparum* proteome

sequences downloaded from PlasmoDB.org, as well as MaxQuant generated databases of

reverse protein sequences and common contaminant protein sequences. Digestion settings

specified Trypsin/P specific enzyme digestion, with a maximum of 2 missed cleavages and a

minimum peptide length of 7. Both PSM (peptide spectral match) and protein FDR cutoffs were

set at 1%. Cysteine carbamidomethylation was set as a fixed modification. Methionine

oxidation and acetylation (protein N-term) were set as variable modifications. The MS/MS

match mass tolerance was set at 20ppm. Label Free Quantification intensities of enriched

proteins were compared across the replicates to determine fold change and statistical

significance using pair wise Student t-test. Volcano plots were visualized using Graphpad Prism.

Growth inhibition assay

All parasite growth inhibition assays were performed in triplicate in 96-well plates as described by Desjardin et al. (28). *P. falciparum*-infected erythrocytes had aTc removed 48 hours prior to the start of the assay. At 1.0% initial parasitemia and 1.5% hematocrit, these parasites were exposed to various concentrations of the indicated drug/inhibitor for 48h and then pulsed with 0.5 µCi of ^{3}H-hypoxanthine for an additional 24h. Untreated parasitized and unparasitized red blood cells were incubated concurrently with the treated parasites as controls for growth and radioactive precursor incorporation. The 96 well plates were then frozen at -80°C for 24 h to disrupt erythrocyte and parasite membrane integrity. Following freezing, the plates were warmed to 37°C, and parasites were harvested onto EasyTab^{TM}-C Self-Aligning Glass Fiber Filters (Packard, Meridian, CT, USA). The filters were then dried completely and placed into an Omnifilter^{TM} 96-well plate filter case (Packard, Meridian, CT, USA). 30µl of OmniScint^{TM} (Packard, Meridian, CT, USA) scintillation fluid was added to each well and the radiation was counted with a TopCount^{TM} counter. The incorporation of radioactivity into nucleic acids served as a measure of cell proliferation and is reported by TopCount as counts per minute (cpm). The total cpm of the non-parasitized red blood cells was subtracted from the cpm of all other wells. The cpm of each sample was converted to percent by dividing it by the cpm of the untreated control. The percent growth versus drug/inhibitor concentration was plotted using GraphPad Prism software. A nonlinear regression dose-response curve was calculated in Prism after adjusting the lowest percent to equal zero and the highest percent to 100.

Results

Generation of parasite lines conditionally expressing MPPα

A parasite line expressing yDHOD (D10_VP2KO) was chosen as the parental line to

prepare the MPPα conditional knockdown line. These parasites are independent of mtETC

activity and are resistant to all mtETC inhibitors (5, 29). We modified the endogenous MPPα

locus (PF3D7_0523100) in D10_VP2KO parasites using CRISPR-Cas9 to generate parasites

conditionally expressing 3HA-tagged MPPα under regulation of the aptamer-TetR-DOZI system

(Figure S1).   PCR confirmed the integration of the expected portion of the donor plasmid within

the modified MPPα locus (Figure 1A and B). Localization of MPPα -3HA was determined by

immunofluorescence using anti-HA antibody (Figure 1C). We used HSP-60 antibody as a

mitochondrial marker to demonstrate colocalization of MPPα at the parasite mitochondrion, as

shown in Fig. 1C.

Figure 1: Generation of parasite lines conditionally expressing MPPα (A) Schematic of MPPα
locus modification using CRISPR-Cas9. Vertical arrow indicates position of sgRNA binding. Also
indicated are the 5' and 3' homologous regions (HR), 5' HR coming from MPPα CDS and the 3'
HR coming from 3'UTR. (B) Using primers indicated by PCR1, amplification of the forward and
reverse primers from the endogenous locus shows that the locus is disrupted in the MPPα
transfected line. Using primers indicated for PCR2, amplification of the forward primer from the
endogenous locus and the reverse primer from the HA tag present in the plasmid determines
correct integration of the donor DNA into the parasite genome. (C) Representative
immunofluorescence images of a parasite conditionally expressing MPPα, showing its
mitochondrial localization.

MPPα is essential for parasite viability

We assessed expression of MPPα in the presence and absence of aTc by carrying out

western blot analysis of parasite lysates collected over 8 days of parasite cultures; HSP60

expression was used as a loading control. As shown in Fig. 2A and B, expression of MPPα was

below detection level by day 4 of parasite growth in absence of aTc. We assessed the effects of

MPPα knockdown on parasite growth by removing aTc at the trophozoite stage of the parasite.

As shown in Fig. 2C, parasite growth ceased in the second cycle following aTc removal. No

viable parasites were detected via Giemsa stain in MPPα (-) condition by day 6 following aTc

removal. Since these studies were carried out using a parasite line that was independent of

mtETC activity by having the metabolic bypass provided by yDHOD, suggesting that MPPα

serves a critical function in addition to supporting mtETC activity as a subunit of complex III.

Figure 2: MPPα is essential for parasite viability. (A) Western blot analysis using anti-HA
and anti-HSP60 antibodies. (B) Quantification of Western blot normalized to intensities of anti-
HSP60. Pixel intensities were measured via Image J.⬚(C) Assessment of growth of MPPα (+) and
MPPα(-) parasites. Growth assay data was conducted in triplicate by flow cytometry using SYBR
green stained parasites.

Mitochondrial membrane potential is disrupted in knockdown parasites.

Maintenance of electrochemical potential across the inner mitochondrial membrane is

critical for the survival of all eukaryotes. We have previously shown that blood stages of

Plasmodium falciparum have two independent means to maintain mitochondrial membrane

potential: one dependent on mtETC and the other independent of mtETC (5). Therefore, we

next examined mitochondrial membrane potential in parasites with and without MPPα

expression.   Using live cell microscopy, the presence of mitochondrial membrane potential was

visualized using Mitotracker red staining in both knockdown and wild type parasites. After one

cycle of MPPα knockdown, 40% of the parasites were observed to have a diffused Mitotracker

staining, which is indicative of collapsed mitochondrial membrane potential. Since parasite

plasma membrane potential remains intact (initially), the positively charged probe accumulates

within the cytoplasm, but dissipation of the mitochondrial membrane potential would prevent

its accumulation in the mitochondrion. After two cycles, 73% of knockdown parasites displayed

diffuse staining. By Day 6 of MPPα knockdown, no parasites with mitochondrial staining were

visible (Figure 3A and B). While MPPα depletion may directly affect mtETC, the secondary

pathway for mitochondrial membrane potential should remain operative. Thus, these data

indicate that removal of MPPα affects both the primary and secondary pathways for generating

membrane potential.

Figure 3: Mitochondrial membrane potential is disrupted in knockdown parasites. (A)
Live cell microscopy images of MPPα (+) and MPPα (−) parasites. Mitotracker staining (red)
revealed tubular morphology in MPPα(+) parasites but was diffused in varying number of
MPPα(-) parasites starting as early as Day 2 following the knockdown (B) Quantitation of
morphological profiles of 30 parasites at each time points was assessed, showing gradual
increase in parasites showing diffuse Mitotracker staining resulting from disruption of
mitochondrial membrane potential.

Knockdown of MPPα expression causes hypersensitivity to proguanil.

A commonly used antimalarial drug, Malarone, is a synergistic combination of atovaquone and

proguanil (30, 31). We have previously shown that the synergistic property of proguanil results

from its inhibition of the secondary pathway for mitochondrial membrane potential generation

(5). Since MPPα depletion affects both mtETC dependent and independent pathways of

generating membrane potential (Figure 4A), we wanted to investigate whether the absence of

MPPα influences proguanil sensitivity of the parasites. We carried out growth inhibition assays

using [3]H-hypoxanthine incorporation by parasites treated with proguanil in presence and

absence of MPPα. While MPPα(+) parasites showed EC50 of 1585 nM for proguanil, MPPα(-)

parasites were approximately 200-fold hypersensitive to proguanil with EC50 of 8.4 nM (Figure

4B). In contrast, there were no changes in response of the parasites to other inhibitors (DSM1,

atovaquone and PA21A092) in presence or absence of MPPα (Figure S3). We have previously

shown that ubiquinone analogue, decyl-ubiquinone, is able to partially rescue parasite growth

when subjected to Complex III inhibition (32). Thus, we assessed the ability of decyl-

ubiquinone to reverse hypersensitivity to proguanil in MPPα(-) parasites. Addition of 25 μM

decyl-ubiquinone had only a relatively minor effect on hypersensitivity to proguanil in MPPα(-)

parasites (Figure 4C).

Figure 4: Knockdown of MPPα expression causes hypersensitivity to proguanil. (A) Parasite
growth inhibition assessed by [3]H-hypoxanthine incorporation following proguanil treatment.
For MPPα- parasites, aTc was removed for 48 h followed by [3]H-hypoxanthine addition for 72hr.
MPPα(+) parasites were also grown for 72hr followed by [3]H-hypoxanthine addition. (B) EC$_{50}$
values for DSM1, atovaquone, PA21A092, and proguanil in MPPα+ and MPPα- parasites. While
the EC50 values for Complex III, DHOD and PfATP4 inhibitors were similar for MPPα(+) and
MPPα(-) parasites, MPPα(-) parasites were almost 200-fold hypersensitive to proguanil. (C)
Parasite growth inhibition assessed by [3]H-hypoxanthine incorporation following proguanil
treatment and addition of 25 μM decyl-ubiquinone where indicated. For MPPα(-) parasites, aTc
was removed for 48 h followed by [3]H-hypoxanthine addition for 72hr.

Proteomic analysis of MPPα immunoprecipitation

To gain insight into interactions of MPPα with other mitochondrial proteins, we carried

out immunoprecipitation from both MPPα tagged and wildtype parasites using anti-HA beads

followed by LC-MS/MS analysis. Overall, 93 proteins were identified as significantly enriched in

the MPPα immunoprecipitated protein samples, using a cutoff of two-fold change compared to

the control with p-value <0.05 (Figure 5A). Of the putative 12 subunits of Complex III, 11

subunits, including mitochondrially encoded cytochrome *b,* were enriched greater than a

million-fold, based on the label free quantification (LFQ) intensity of the peptides observed.

QCR13 (Pf3D7_0817800) was the only putative Complex III protein that was absent, likely due

to its small size of just 54 amino acids. Remarkably, there were 45 additional predicted or

experimentally confirmed mitochondrial proteins that were equally significantly enriched. This

suggests their interactions with MPPα are of similar magnitude as MPPα's association with

Complex III. We postulate the presence of these mitochondrial proteins with MPPα being by

virtue of their transient association during processing of their mitochondrial targeting peptides.

The significance of additional proteins that were enriched but not predicted to be

mitochondrial remains unclear. It is interesting to note, that cytochrome *c* oxidases subunits 1

and 3 were not enriched. This is likely due to these proteins being encoded by mtDNA and not

requiring processing of mitochondrial targeting sequences (1,6) . These data suggest that

Complex III is engaged in the dual functions of mtETC integrity, as well as processing imported

mitochondrial proteins.

Figure 5: Proteomic analysis of MPPα immunoprecipitation (A) Volcano plot of
enriched proteins with a $\log_2$-fold change of label free quantification (LFQ) intensity and p-
values of < 0.05 found in immunoprecipitated MPPα samples. The red dots represent proteins
present in Complex III. The purple dots represent proteins that are predicted to be targeted to
the mitochondria. The blue dots represent the other proteins that were significantly enriched in

the immunoprecipitated samples. (B) Pie chart depicting the number of proteins that fall into
the categories of Complex III, likely mitochondrially targeted, and other enriched proteins.

Discussion

344   Although mitochondria in all extant organisms originated from the symbiotic

arrangement between an archaeon and an α-proteobacterium that led to the emergence of

eukaryotes, mitochondrial genomes and physiology are as divergent as the organisms in which

they reside (33, 34). Oxidative phosphorylation to generate ATP, while a major function of

mitochondria in many organisms, is not essential in some organisms, especially those that live in

anaerobic environments (35). Indeed, such organisms often lack mtDNA but continue to

possess a remnant mitochondrion called a mitosome (36, 37). In addition, certain yeasts (38),

cultured vertebrate cells (39) and bloodstream-forms of trypanosomes (40) can be manipulated

to lose their mtDNA while maintaining the mitochondrion itself. These are called rho^0 cells, and

investigations with such rho^0 cells have helped to establish that the absolute essential function

of their retained mtDNA-negative mitochondria is to synthesize and export inorganic Fe-S

clusters for a variety of iron-sulfur proteins required for cell survival (41, 42). To permit this

essential function, mitochondria need to maintain an electropotential across their inner

membrane, which is required for protein and metabolite transport. In mitochondria with an

active mtETC the membrane potential is normally established through proton pumping by the

respiratory complexes. A lack of mtDNA will lead to the absence of mtETC, so then the

electropotential will need to be generated through alternative means, generally by reverse

action of ATP synthase and/or by electrogenic exchange of ATP/ADP (43).

Asexual blood stage *P. falciparum* rely on glycolysis for ATP generation with little

contribution from mitochondrial oxidative phosphorylation (44, 45). Yet the parasite maintains

an active mtETC, inhibition of which by antimalarials such as atovaquone results in parasite

death. We have previously shown that the critical function of mtETC in blood stage *P. falciparum*

is to re-oxidize ubiquinol to support mitochondrial DHOD, which is needed for pyrimidine

biosynthesis, an essential pathway in the parasites (5). Parasites engineered to bypass

mitochondrial DHOD become mtETC independent and resistant to drugs such as atovaquone.

We also showed that generation of electropotential across the mitochondrial inner membrane

was still required in these mtETC-independent parasites, and that the generation of this

potential could be inhibited by proguanil, resulting in parasite death when combined with

mtETC inhibitors (5). Since mtDNA in *P. falciparum* encodes just three components of mtETC, we

initially reasoned that it should be possible generate rho^0 parasites. However, various attempts

failed to generate such parasites (8-10). It is of interest to note that all organisms where rho^0

cells can be generated contain matrix-localized MPP (e.g., yeast, trypanosomes, and mammals),

whereas organisms with Complex III associated MPP, such as plants (and *P. falciparum*), cannot

be rendered rho^0. Because cytochrome *b* of Complex III is always encoded by mtDNA,

maintenance of mtDNA remains essential in these organisms for proper assembly of MPP within

Complex III, even when mtETC is made non-essential.

A previous study by Espino-Sanchez et al. (46) assessed effects of conditionally knocking

down expression of cytochrome c_1, a component of Complex III, in *P. falciparum*. They showed

that the parasites lacking this component of Complex III failed to survive but could be rescued

by addition of decyl-ubiquinone, consistent with Complex III being the source of ubiquinone to

support DHOD. Furthermore, parasites lacking cytochrome c_1 continued to maintain

mitochondrial membrane potential, which could be collapsed by treatment with proguanil (46).

These results are consistent with prior inferences that Complex III is essential for mtETC activity.

In contrast to cytochrome c_1 knockdown, however, we show here that knockdown of MPPα not

only results in parasite demise but also a collapsed mitochondrial membrane potential.

Parasites with MPPα knockdown cannot be rescued by either the presence of yDHOD or

addition of decyl-ubiquinone. Since MPPα is a component of Complex III, our study establishes

a secondary role of MPP in addition to maintaining the structure and function of Complex III of

mtETC. We posit that this secondary role is to process mitochondrially targeted proteins. In

support of this, we observed peptides from multiple mitochondrially targeted proteins in

proteomic analyses of MPPα pulldown samples (Figure 5). We suggest that these peptides are

generated from proteins transiently associated with Complex III as they are being processed by

the MPP subunits.

References

1.   Vaidya AB, Lashgari MS, Pologe LG, Morrisey J. Structural features of Plasmodium
cytochrome b that may underlie susceptibility to 8-aminoquinolines and
hydroxynaphthoquinones. Molecular and Biochemical Parasitology. 1993;58(1):33-42.
2.   Fry M, Pudney M. Site of action of the antimalarial hydroxynaphthoquinone, 2-[trans-4-
(4'-chlorophenyl) cyclohexyl]-3-hydroxy-1,4-naphthoquinone (566C80). Biochem Pharmacol.
1992;43(7):1545-53.
3.   Crofts AR. The Q-cycle - A Personal Perspective. Photosynth Res. 2004;80(1-3):223-43.
4.   Mitchell P. The protonmotive Q cycle: A general formulation. FEBS Letters.
1975;59(2):137-9.
5.   Painter HJ, Morrisey JM, Mather MW, Vaidya AB. Specific role of mitochondrial
electron transport in blood-stage Plasmodium falciparum. Nature. 2007;446(7131):88-91.
6.   Vaidya AB, Akella R, Suplick K. Sequences similar to genes for two mitochondrial
proteins and portions of ribosomal RNA in tandemly arrayed 6-kilobase-pair DNA of a malarial
parasite. Mol Biochem Parasitol. 1989;35(2):97-107.
7.   Lamb IM, Okoye IC, Mather MW, Vaidya AB. Unique Properties of Apicomplexan
Mitochondria. Annual Review of Microbiology. 2023;77(Volume 77, 2023):541-60.
8.   Ke H, Morrisey JM, Ganesan SM, Mather MW, Vaidya AB. Mitochondrial RNA
polymerase is an essential enzyme in erythrocytic stages of Plasmodium falciparum. Mol
Biochem Parasitol. 2012;185(1):48-51.
9.   Ke H, Dass S, Morrisey JM, Mather MW, Vaidya AB. The mitochondrial ribosomal
protein L13 is critical for the structural and functional integrity of the mitochondrion in
Plasmodium falciparum. J Biol Chem. 2018;293(21):8128-37.
10.   Dass S, Mather MW, Morrisey JM, Ling L, Vaidya AB, Ke H. Transcriptional changes in
Plasmodium falciparum upon conditional knock down of mitochondrial ribosomal proteins
RSM22 and L23. PLOS ONE. 2022;17(10):e0274993.
11.   Chacinska A, Koehler CM, Milenkovic D, Lithgow T, Pfanner N. Importing
Mitochondrial Proteins: Machineries and Mechanisms. Cell. 2009;138(4):628-44.
12.   Wiedemann N, Pfanner N. Mitochondrial Machineries for Protein Import and Assembly.
Annual Review of Biochemistry. 2017;86(Volume 86, 2017):685-714.
13.   Pollock RA, Hartl FU, Cheng MY, Ostermann J, Horwich A, Neupert W. The processing
peptidase of yeast mitochondria: the two co-operating components MPP and PEP are structurally
related. Embo j. 1988;7(11):3493-500.
14.   Böhni PC, Daum G, Schatz G. Import of proteins into mitochondria. Partial purification
of a matrix-located protease involved in cleavage of mitochondrial precursor polypeptides. J Biol
Chem. 1983;258(8):4937-43.
15.   Taylor AB, Smith BS, Kitada S, Kojima K, Miyaura H, Otwinowski Z, et al. Crystal
Structures of Mitochondrial Processing Peptidase Reveal the Mode for Specific Cleavage of
Import Signal Sequences. Structure. 2001;9(7):615-25.
16.   Dvořáková-Holá K, Matušková A, Kubala M, Otyepka M, Kučera T, Večeř J, et al.
Glycine-Rich Loop of Mitochondrial Processing Peptidase α-Subunit Is Responsible for
Substrate Recognition by a Mechanism Analogous to Mitochondrial Receptor Tom20. Journal of
Molecular Biology. 2010;396(5):1197-210.

17.   Nagao Y, Kitada S, Kojima K, Toh H, Kuhara S, Ogishima T, et al. Glycine-rich region of
mitochondrial processing peptidase alpha-subunit is essential for binding and cleavage of the
precursor proteins. J Biol Chem. 2000;275(44):34552-6.
18.   Iwata S, Lee JW, Okada K, Lee JK, Iwata M, Rasmussen B, et al. Complete structure of
the 11-subunit bovine mitochondrial cytochrome bc1 complex. Science. 1998;281(5373):64-71.
19.   Berry EA, Guergova-Kuras M, Huang L-s, Crofts AR. STRUCTURE AND FUNCTION
OF CYTOCHROME bc COMPLEXES. Annual Review of Biochemistry. 2000;69(1):1005-75.
20.   Brandt U, Yu L, Yu CA, Trumpower BL. The mitochondrial targeting presequence of the
Rieske iron-sulfur protein is processed in a single step after insertion into the cytochrome bc1
complex in mammals and retained as a subunit in the complex. Journal of Biological Chemistry.
1993;268(12):8387-90.
21.   Yang M, Jensen RE, Yaffe MP, Oppliger W, Schatz G. Import of proteins into yeast
mitochondria: the purified matrix processing protease contains two subunits which are encoded
by the nuclear MAS1 and MAS2 genes. Embo j. 1988;7(12):3857-62.
22.   Brumme S, Kruft V, Schmitz UK, Braun HP. New insights into the co-evolution of
cytochrome c reductase and the mitochondrial processing peptidase. J Biol Chem.
1998;273(21):13143-9.
23.   Braun HP. The two roles of complex III in plants. Elife. 2021;10.
24.   Ganesan SM, Morrisey JM, Ke H, Painter HJ, Laroiya K, Phillips MA, et al. Yeast
dihydroorotate dehydrogenase as a new selectable marker for Plasmodium falciparum
transfection. Mol Biochem Parasitol. 2011;177(1):29-34.
25.   Concordet JP, Haeussler M. CRISPOR: intuitive guide selection for CRISPR/Cas9
genome editing experiments and screens. Nucleic Acids Res. 2018;46(W1):W242-w5.
26.   Spillman NJ, Beck JR, Ganesan SM, Niles JC, Goldberg DE. The chaperonin TRiC forms
an oligomeric complex in the malaria parasite cytosol. Cell Microbiol. 2017;19(6).
27.   Ganesan SM, Falla A, Goldfless SJ, Nasamu AS, Niles JC. Synthetic RNA-protein
modules integrated with native translation mechanisms to control gene expression in malaria
parasites. Nat Commun. 2016;7:10727.
28.   Desjardins RE, Canfield CJ, Haynes JD, Chulay JD. Quantitative assessment of
antimalarial activity in vitro by a semiautomated microdilution technique. Antimicrob Agents
Chemother. 1979;16(6):710-8.
29.   Ganesan SM, Morrisey JM, Ke H, Painter HJ, Laroiya K, Phillips MA, et al. Yeast
dihydroorotate dehydrogenase as a new selectable marker for Plasmodium falciparum
transfection. Mol Biochem Parasitol. 2011 (in press).
30.   Looareesuwan S, Chulay JD, Canfield CJ, Hutchinson DB. Malarone (atovaquone and
proguanil hydrochloride): a review of its clinical development for treatment of malaria. Malarone
Clinical Trials Study Group. Am J Trop Med Hyg. 1999;60(4):533-41.
31.   Srivastava IK, Vaidya AB. A mechanism for the synergistic antimalarial action of
atovaquone and proguanil. Antimicrob Agents Chemother. 1999;43(6):1334-9.
32.   Ke H, Morrisey JM, Ganesan SM, Painter HJ, Mather MW, Vaidya AB. Variation among
Plasmodium falciparum strains in their reliance on mitochondrial electron transport chain
function. Eukaryot Cell. 2011;10(8):1053-61.
33.   Lane N, Martin W. The energetics of genome complexity. Nature. 2010;467(7318):929-3
34.   Martin W, Muller M. The hydrogen hypothesis for the first eukaryote. Nature.
1998;392(6671):37-41.

35.   Stairs CW, Leger MM, Roger AJ. Diversity and origins of anaerobic metabolism in
mitochondria and related organelles. Philos Trans R Soc Lond B Biol Sci.
2015;370(1678):20140326.
36.   Tovar J, Leon-Avila G, Sanchez LB, Sutak R, Tachezy J, van der Giezen M, et al.
Mitochondrial remnant organelles of Giardia function in iron-sulphur protein maturation. Nature.
2003;426(6963):172-6.
37.   Shiflett AM, Johnson PJ. Mitochondrion-related organelles in eukaryotic protists. Annu
Rev Microbiol. 2010;64:409-29.
38.   Tzagoloff A, Myers AM. Genetics of mitochondrial biogenesis. Annu Rev Biochem.
1986;55:249-85.
39.   King MP, Attardi G. Human cells lacking mtDNA: repopulation with exogenous
mitochondria by complementation. Science. 1989;246(4929):500-3.
40.   Schnaufer A, Clark-Walker GD, Steinberg AG, Stuart K. The F1-ATP synthase complex
in bloodstream stage trypanosomes has an unusual and essential function. EMBO J.
2005;24(23):4029-40.
41.   Lill R, Kispal G. Maturation of cellular Fe-S proteins: an essential function of
mitochondria. Trends Biochem Sci. 2000;25(8):352-6.
42.   Stehling O, Elsasser HP, Bruckel B, Muhlenhoff U, Lill R. Iron-sulfur protein maturation
in human cells: evidence for a function of frataxin. Hum Mol Genet. 2004;13(23):3007-15.
43.   Schnaufer A, Domingo GJ, Stuart K. Natural and induced dyskinetoplastic
trypanosomatids: how to live without mitochondrial DNA. Int J Parasitol. 2002;32(9):1071-84.
44.   Sherman IW. Biochemistry of Plasmodium (malarial parasites). Microbiol Rev.
1979;43(4):453-95.
45.   Vaidya AB. Mitochondrial physiology as a target for atovaquone and other antimalarials.
In: Sherman IW, editor. Malaria: Parasite biology, pathogenesis, and protection. Washington, DC:
ASM Press; 1998. p. 355-68.
46.   Espino-Sanchez Tanya J, Wienkers H, Marvin Rebecca G, Nalder S-a, García-Guerrero
Aldo E, VanNatta Peter E, et al. Direct tests of cytochrome c and c_1 functions in the electron
transport chain of malaria parasites. Proceedings of the National Academy of Sciences.
2023;120(19):e2301047120.

Acknowledgements

We thank Dr. Dale Chaput at University of South Florida Advanced Research Core for Mass

Spectrometry for assistance with proteomic analysis. This work was supported by Grant number

R01 AI028398 from National Institutes of Health.

S1: **Maps of plasmids used for transfections.** Maps of the two gRNA plasmids (A) and (B), and
the map for double cross over modification of MPPα locus (C) are shown.

S2: **MPPα disruption in a second parasite line not expressing yDHOD.**(A) Assessment of growth

of MPPα + and MPPα - parasites in Dd2attb parasites. Growth assay was conducted in triplicate

by counting 500 parasites via Giemsa stain. (B) Western blot analysis using anti-HA and anti-

Aldolase antibodies.

S3: **Susceptibility of MPPα - parasites to antimalarials remains unchanged.** Parasite growth

inhibition assessed by ^{3}H-hypoxanthine incorporation following DSM1, atovaquone, PA21A092

treatment. For MPPα - parasites, aTc was removed for 48 h followed by 3H-hypoxanthine

addition for 72 hr.

A

Donor Plasmid
5'HR | 3xHA | 10x apt | TetRDozi-BSD | HSP86 UTR | 3'HR

Endogenous Locus
5'UTR | MPPα CDS | 5'HR | 3'HR | 3'UTR
sgRNA
Homologous Recombination

Intact locus: PCR1 : 1598 bp

Integrated
5'UTR | MPPα CDS+5'HR | 3xHA | 10x apt | TetRDozi-BSD | HSP86 UTR | 3'UTR+HR

Integrated construct: PCR2 : 1003 bp

B

WT PCR1 WT PCR2

C

Brightfield | Anti-HA | Anti-HSP60 | DAPI | HA/HSP60 | Merge

A

	Merge	MitoTracker	Hoechst	Brightfield
Day 2 MPPα (+)				
Day 2 MPPα (−)				
Day 4 MPPα (+)				
Day 4 MPPα (−)				
Day 6 MPPα (+)				
Day 6 MPPα (−)				

B

%Mitochondrial Morphology

Diffused
Tubular

Day 2 MPPα (+)
Day 2 MPPα (−)
Day 4 MPPα (+)
Day 4 MPPα (−)
Day 6 MPPα (+)
Day 6 MPPα (−)

A

B

Compound	EC50 MPPα (+)	EC50 MPPα (-)
DSM1	> 10,000	> 10,000
Atovaquone	> 10,000	> 10,000
PA21A092	18.9±0.8	21.8±0.6
Proguanil	1585±2.1	84±5.1
Proguanil w/ Decylubiquinone	2785±3.3	107.1±3.6

C