1

**RiboTRAP-seq identifies spatially distinct functions for the anterior and posterior intestine in immune and metabolic regulation in *C. elegans***

Chung-Chih Liu[1,3], Nicolas Seban[1,2,3], Supriya Srinivasan[1]*

[1]Department of Neuroscience and Dorris Neuroscience Center, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA. [2]Current affiliation: UC Davis Center for Neuroscience, Davis, CA 95618, USA. [3]These authors contributed equally.

*Corresponding Author

## **Abstract**

The intestine integrates food-derived cues to coordinate organismal physiology, yet the

molecular specialization of discrete regions along the intestinal epithelium remains unclear.

Here, we generate cell type-specific translatomes of the *Caenorhabditis elegans* intestine during

fasting and refeeding using discrete promoters for the anterior quartet of intestinal cells (INT1)

and the remaining 8 pairs of intestinal cells (INT2-9). We found that the anterior-most INT1 cells

are a translationally distinct intestinal sub-compartment that is particularly enriched for immune-

and stress-response genes. Functional assays using novel INT1-specific genes emerging from

this study reveal that these specialized cells play a previously unappreciated role in pathogen

avoidance and organismal survival. A second critical function of INT1 cells is their role in

sensing and responding to the contents of the gut lumen. We show that luminal pyruvate is the

key signal linking bacterial nutrients in the gut, to secretion of the gut insulin antagonist INS-7.

These findings establish INT1 as sentinel enteroendocrine cells that integrate metabolic and

immune cues to couple food status with immune and endocrine responses. Our studies also

provide a rich resource for dissecting segment-specific intestinal biology, an overlooked and

fertile area for future research.

## Introduction

The intestine is a critical organ for food digestion and nutrient absorption in animals. *Caenorhabditis elegans*, as a bacterivore, constantly ingests a variety of bacteria with an intestinal transit time as short as two minutes in adults [1]. This rapid process implies the existence of food-sensing mechanisms within the intestine that can continuously monitor and respond to the acute fluctuation in food availability. When food is absent, a transcriptional metabolic response is evoked via the induction of the fasting response genes including lipases, peroxisomal and mitochondrial β-oxidation genes and some fatty acid desaturases in the intestine to maintain energy homeostasis [2-4]. However, it has been unclear whether discrete segments within the intestine have specific food-sensing capabilities or exhibit different responses to fasting in comparison to the organ as a whole. To our knowledge, no studies to date have addressed this question with high spatial and temporal resolution. The *C. elegans* intestine is comprised of 20 cells designated INT1-INT9 [5]. The anterior ring comprises 4 cells called INT1, which are the most proximal intestinal cells adjacent to the pharynx and represent the initial location of contact with ingested food, which for *C. elegans* is digested bacterial components. The remaining 16 cells are arranged in pairs to form the latter 8 posterior rings (INT2-9) arranged in a stereotypical structure. A number of years ago *Sulston et al.* had observed that the microvilli of INT1 cells are shorter than those in the rest of the intestine [5], which implies that INT1 cells may not have the same food absorption capacity that INT2-9 cells have.

In addition to the morphological differences, we recently reported that INT1 functions as an enteroendocrine cell, and secretes the gut peptide INS-7 shortly after food withdrawal. INS-7 secretion levels increase over the duration of fasting, which are reset by the re-introduction of food. Gut INS-7 acts in the nervous system to suppress the fat loss-stimulating peptide FLP-7, functional ortholog of the *C. elegans* tachykinin peptide [6]. This mechanism allows the stimulation of fat loss to be regulated, in part, by incoming food signals from the gut. Furthermore, *Quintin et al.* showed that worms treated with $H_2O_2$ have an induction of the antioxidant enzyme PRDX-2 specifically in INT1 cells, suggesting that INT1 cells may have additional oxidative stress-sensing functions [7]. Although these lines of evidence have suggested that INT1 cells may be a group of specialized intestinal cells that sense the quality of the ingested food and have distinct fasting responses, many questions remain. First, the molecular features of INT1 cells and the molecular distinctions between INT1 and the rest of the intestine (INT2-9) have not been defined. Second, the role of INT1 cells in modulating

physiological responses to changes in food availability remains undetermined. Finally, the

differential effects of metabolic challenge via fasting and re-feeding in the subsets of intestinal

cells are unclear.

In this study, we aim to address these questions using spatial translatomics. In short, the

approach is as follows: actively translating mRNAs are bound to clusters of ribosomes, referred

to as polysomes [8, 9]. Translating ribosome affinity purification (TRAP) was developed to

isolate cell type-specific polysome-mRNA complexes [10, 11], allowing quantitative analysis of

translational regulation [12]. Here, we performed TRAP-Seq using a modified protocol based on

*McLachlan et al.* [13], in which RPL-22 was tagged with HA to enable affinity purification of

polysome-mRNA complexes, which is followed by RNA sequencing. This approach provided a

comprehensive spatially distinct translatomic profile of actively translating mRNAs in INT1 and

INT2-9 cells. To discern novel temporal features in response to metabolic challenge, we

examined the translational responses of these intestinal subsets to acute and chronic fasting

and refeeding paradigms. Our datasets reveal that INT1 cells exhibit a distinct molecular identity,

characterized by the enrichment of immune- and stress-responsive genes. Furthermore, fasting

and refeeding elicit divergent regulatory programs in INT1 and INT2-9 cells. To our knowledge,

this represents the first study to determine the molecular identities of discrete intestinal cell

subsets in response to fasting and refeeding in *C. elegans*.

## Results

### Isolation and sequencing of translating RNA from INT1 and INT2-9 cells

To identify the molecular differences between INT1 and INT2-9 cells in fed, fasted and refed

states, we employed translating ribosome affinity purification (TRAP) [10, 11] to isolate

polysomal RNA from INT1 and INT2-9 cells followed by RNA sequencing. This approach

captures actively translating mRNAs, thereby providing a cell type-specific snapshot of the

translational landscape under fasting and refeeding conditions. To selectively target INT1 and

INT2-9 cells, we expressed the HA-tagged ribosomal protein (RPL-22-3xHA) under the control

of the *Pges-1ΔB* [6, 14] and *Ppho-1* [15] promoters, respectively. As published [6], we noted that

*Pges-1ΔB* is strictly expressed in INT1 cells, whereas *Ppho-1* is expressed in INT2-9 cells with

detectable, albeit lower expression in INT2 cells (Fig 1A).

We operationally defined the 'acute' conditions at 30 min fasting followed by 30 min refeeding,

because under these conditions, intestinal fat content remains unchanged (Fig 1B and 1C).

However, INS-7 secretion was induced within the 30 min acute fasting period and restored to

baseline levels within 30 min upon refeeding [6] , indicating that intestinal responses to food

withdrawal can occur before detectable changes in lipid stores. Chronic conditions were

established as 3 hr fasting followed by 90 min refeeding. In this paradigm, animals lost

approximately half of intestinal lipid stores after chronic fasting, which were subsequently

replenished upon refeeding for 90 min (Fig 1D and 1E). Thus, chronic conditions represent a

more pronounced metabolic challenge. For TRAP-Seq sample preparation, we cultured 100,000

*INT1::TRAP* or *INT2-9::TRAP* animals to the Day 1 adult stage, subjected them to acute or

chronic conditions (Fig 1F), and performed RNA isolation following a modified TRAP protocol

[13]. These experiments were conducted 3 times.

## Stress response genes distinguish INT1 from INT2-9 translatomes across acute and chronic conditions

Principal component analysis (PCA) of all TRAP-Seq datasets revealed clearly distinct

clustering by promoter identity, indicating that INT1 and INT2-9 cells maintain fundamentally

different molecular identities independent of any and all acute or chronic conditions (Fig 2A). To

systematically define these differences, we performed weighted correlation network analysis

(WGCNA) [16], which unbiasedly identified 12 gene modules associated with condition-specific

responses across the INT1 and INT2-9 datasets (Fig 2B and 2C, S1A Fig). Notably, the

expression of genes in module red was positively correlated to all INT1 conditions but negatively

correlated to all INT2-9 conditions, suggesting a transcriptional program preferentially translated

in INT1 cells (Fig 2C and S1A Fig). Examination of normalized counts confirmed uniformly

higher expression of module red genes across all INT1 conditions compared with INT2-9 (Fig

2D). To identify the categories and functions of module red genes, we used WormCat [17, 18],

an annotation and category enrichment tool with near-complete annotation of the *C. elegans*

genome. Among the 100 genes from module red, we found genes involved in "Stress Response"

including C-type lectin, pathogen and heat categories (Fig 2E). In addition, Gene Ontology (GO)

term enrichment analysis of module red genes further highlighted the enrichment for the terms

"Immune system process" and "Response to biotic stimulus" (S1B Fig).

Recently, *Ghaddar et al.* presented a gene expression atlas of adult *C. elegans* with
single-cell RNA sequencing [19]. They categorized the intestinal cells into three subtypes:
intestine anterior, intestine middle and intestine posterior. To place our findings in the context of
previously defined intestinal cell subtypes, we compared our datasets to theirs. Genes
annotated as 'intestine anterior' were upregulated in INT1 cells, whereas 'intestine posterior'
genes were downregulated in INT1 relative to INT2-9, supporting our conclusion that INT1
represents a translationally distinct intestinal subset (S1C-E Fig).

Next, we decided to comprehensively characterize the translatomic differences between
INT1 and INT2-9 cells under each acute and chronic condition. We began with the differentially
expressed genes in INT1 cells relative to INT2-9 cells in the fed state. There were 153
upregulated and 58 downregulated genes (Fig 2F), and we found *ins-7* among the upregulated
genes, consistent with our prior report of its restricted expression in INT1 [6]. To identify the
categories of the differentially expressed genes in INT1 cells under the fed condition, we
performed WormCat category enrichment analysis and also used ShinyGO [20] to conduct
Gene Ontology (GO) term enrichment analysis with a hierarchical clustering tree, allowing us to
visualize the overlapping relationships among enriched GO terms. In the upregulated genes
with WormCat analysis, we found genes involved in "Stress Response" including pathogen, C-
type lectin, and heat-shock categories (Fig 2G). Further, the results from ShinyGO
demonstrated a cluster of GO terms related to "Immune Response" and "Unfolded Protein
Response" (Fig 2H). In contrast, no functional enrichment was detected among the INT1-
downregulated genes. Together, these findings show that INT1 cells are characterized by a
stress- and immune-responsive translational program in the fed state, marking a strong and
clear distinction from the rest of the intestine.

**C-type lectins and ShKT domain-containing proteins are enriched in INT1 cells**

We noticed that many INT1-differentially expressed genes were not assigned to a category by
WormCat or annotated by ShinyGO. To further explore their molecular functions, we used the
InterPro database to annotate the protein domains for each gene and assign the domains into
several main categories, such as "Enzyme", "Membrane Protein", and "Transcription Factor".
Genes encoding domains that did not fall into these categories, as well as those lacking
identifiable domains, were grouped as "Genes with Unknown Functions" (S2A and S2B Fig).
Among the 149 upregulated genes in INT1, we found 11 encoding C-type lectins (S2A Fig). C-

type-lectin-domain-containing proteins  are notable for pathogen recognition and the initiation of

immune responses in vertebrates [21]. In *C. elegans,* the expression of many distinct C-type

lectins has previously been found to be induced in a pathogen-specific exposure [22, 23].

In addition, we discovered a subset of INT1 upregulated genes encoding ShKT domains

within the genes with unknown functions (S2A Fig). The ShK toxin (ShKT) is a potassium ($K^+$)

channel blocker originally isolated from the sea anemone *Stichodactyla helianthus* [24]. ShKT

domain-containing proteins have been reported in parasitic nematode secretomes [25] and

shown to modulate the activity of human effector memory T cells by blocking voltage-gated

potassium channels [26]. Their biological functions in *C. elegans*, however, remain unclear. Our

findings that specific ShKT domain-containing proteins are specifically expressed in INT1 cells

are intriguing. Notably, several INT1-upregulated ShKT proteins, including F01D5.1, F01D5.5

and SYSM-1, were previously found to be upregulated by PMK-1 (p38 MAPK) [27], a key

regulator of pathogen-induced immune responses in *C. elegans* [27, 28], suggesting that these

ShKT proteins may contribute to intestinal immune functions.

To validate our TRAP-Seq findings and define the spatial expression of INT1-enriched

genes, we chose key candidate genes with high fold change and low adjusted p-value

(therefore, the most significantly altered genes, Fig 2I and S1F-J Fig) to generate transcriptional

reporters using their endogenous promoters and 3'UTRs. We started with *B0024.4*, a

membrane raft protein-coding gene with the lowest adjusted p-value in the INT1-enriched genes,

which revealed  tight expression restricted to INT1 cells. We next decided to examine reporters

for two C-type lectin genes, *clec-86* and *clec-160*, and two ShKT protein-coding genes, *F01D5.1*

and *F01D5.5.* Reporters for *clec-86, clec-160 and F01D5.5* also exhibited highly specific

expression in INT1, whereas *F01D5.1* was expressed in both INT1 and INT2. Out of curiosity,

we also generated a reporter line for the most highly downregulated gene *npr-28* in INT1, and

interestingly, found that it was restricted to the posterior intestinal cells INT7-9, with no

expression in the anterior intestine (Fig 2J). Collectively, these analyses corroborate the spatial

restriction of our TRAP-Seq method  and demonstrate that INT1-upregulated and INT1-

downregulated genes exhibit distinct spatial expression patterns along the intestine.

**Comparisons of enriched categories under acute and chronic conditions identify**

**ShKT domain-containing proteins as stable INT1-specific markers**

We next wished to evaluate whether there are specific differences between INT1 and INT2-9

cells under acute fasting and refeeding. In the acute fasted state, INT1 displayed 90

upregulated and 12 downregulated genes (Fig 3A). WormCat analysis of upregulated genes

revealed those involved in "Stress Response" including C-type lectin and pathogen categories

(Fig 3B), while ShinyGO identified a cluster of GO terms related to "Defense Response" (Fig

3C). In the acute refed state, we observed 52 INT1-upregulated and 35 INT1-downregulated

genes (Fig 3D). Enriched categories of the upregulated genes again included "Stress

Response", particularly pathogen and C-type lectins by WormCat (Fig 3E), with GO terms

related to "Defense Response" and "Immune Response" by ShinyGO (Fig 3F). InterPro

annotations of INT1-upregulated genes revealed consistent representation of "Enzyme", "C-type

Lectin" and "Membrane Protein" categories across fed, acute fasted, and acute refed states

(S3A and S3C Fig). Although enriched functional categories were similar between the 2

intestinal sub-compartments, a Venn diagram analysis of the "Stress Response" genes

demonstrated only partial overlap across states (Fig. 4A), suggesting that translational

differences between INT1 and INT2-9 vary depending on physiological states.

To further test this possibility, we compared gene lists from 4 major InterPro categories

("Enzyme", "Membrane Protein", "C-type Lectin" and "ShKT Protein") across fed, acute fasted

and acute refed states (Fig 4B-H). While "Enzyme", "Membrane Protein", and "C-type Lectin"

categories exhibited only partial overlap, the "ShKT Protein" category showed near-complete

overlap across conditions (Fig 4E), suggesting that ShKT proteins represent stable INT1-

specific markers under acute conditions.

We next extended our analysis to chronic fasting and refeeding, which elicit more

pronounced metabolic shifts in the intestine. In the chronic fasted state, INT1 exhibited 111

upregulated and 77 downregulated genes (Fig 3G). WormCat analysis of the upregulated genes

again identified enrichment for "Stress Response" categories, including C-type lectins and

pathogen-related genes (Fig 3H), while ShinyGO revealed a cluster of GO terms related to

"Immune Response" (Fig 3I). Similar enrichments were observed in the chronic refed state (Fig

3K and 3L). InterPro annotations of INT1 upregulated genes revealed categories consistent with

acute conditions, including "Enzyme", "Membrane Protein" and "C-type Lectin" (S4A and S4C

Fig). Comparison of gene lists demonstrated partial overlap within these categories (Fig 4J-L,

4N-P). Strikingly, however, upregulated ShKT genes showed near-complete overlap across

chronic conditions (Fig 4M).

Finally, we compared the INT1-upregulated ShKT genes across both acute and chronic

conditions. Venn diagram analysis revealed complete overlap, with the same set of 11 ShKT

genes consistently upregulated in all conditions (Fig 4R). These findings demonstrate that ShKT

domain-containing proteins represent robust INT1-specific markers, stable across both acute

and chronic states. Thus, pathogen-sensing, detoxification and mounting an organismal

response to incoming information from the gut lumen are the principal functions of INT1.

**INT1 and INT2-9 cells exhibit distinct fasting and refeeding responses under**

**acute and chronic conditions**

Given the fundamentally distinct translatomes of the INT1 and INT2-9 cells, we next

investigated whether these differences extend to their responses to fasting and refeeding.

Specifically, we sought to determine if the translatomic profiles of INT1 and INT2-9 cells exhibit

distinct regulatory patterns in response to metabolic shifts induced by fasting and subsequent

refeeding. Looking at the concordance of differentially expressed genes during chronic and

acute fasting between INT1 and INT2-9 cells, we identified fasting-response genes with

opposing regulation between the two intestinal subsets (Fig 5A and 5B). For instance, 270

genes were upregulated in INT1 following acute fasting but downregulated in INT2-9 under the

same condition (Fig 5A), indicating divergent translational responses to metabolic shifts.

We next listed all differentially expressed genes (DEGs) in INT1 and INT2-9 under acute

and chronic conditions. Overall, INT1 exhibited fewer DEGs than INT2-9 (Fig 5C). To rule out

the possibility that INT1 DEGs represented a subset of INT2-9 DEGs, we used Venn diagrams

to compare all the INT1 and INT2-9 DEGs from both acute and chronic conditions (Fig 5D). We

found two major features of these fasting and refeeding response genes. Notably, INT1 DEGs

were not a subset of INT2-9 DEGs regardless of acute or chronic conditions, but rather

represented distinct gene sets (Fig 5D). Another key distinction is that for both INT1 and INT2-9,

the acute response genes did not overlap substantially with chronic response genes, indicating

that distinct translational programs are engaged during acute versus chronic conditions (Fig 5D).

To identify the enriched categories of these INT1 and INT2-9 DEGs under each

condition, we performed WormCat analysis and found that INT1 datasets generally had fewer

enriched categories than INT2-9 datasets under each condition. (S5A and S5B Fig, S6A and

S6B Fig). In the acute fasted state, upregulated genes in both INT1 and INT2-9 were enriched

for "Metabolism", whereas categories including "Stress Response", "Transcription Factor", and

"Lysosome" were specific to INT2-9 (S5A Fig). Among the downregulated genes in both cell

types in the acute fasted state, we found genes involved in "Ribosome", including ribosomal

subunit genes (S5B Fig). Interestingly, in the acute refed state, genes involved in "Ribosome"

were upregulated in INT2-9 (S5C Fig), suggesting the sensitivity of ribosomal gene expression

to acute fasting and refeeding. Additional categories, such as "Transcription Factor" and

"Proteolysis Proteasome", were enriched exclusively among INT2-9 downregulated genes in the

acute refed state (S5D Fig).

Further, we observed that chronic conditions elicited broader and more diverse

enrichment patterns than acute states in both INT1 and INT2-9 (S5, S6 and S7 Figs). Although

this is an expected result, the category term "Stress Response" was surprisingly enriched only

among INT1-upregulated genes in the chronic fasted state (Fig 5E), whereas "Pathogen" was

uniquely enriched among INT1-upregulated genes in the chronic refed state (Fig 5F),

underscoring an important potential specialization of INT1 in stress and immune responses.

Conversely, both INT1 and INT2-9 datasets shared enrichment for "Ribosome" and "Metabolism"

among downregulated genes in the chronic fasted state and upregulated genes in the chronic

refed state (S6B and S7A Figs), suggesting that protein translation and metabolic regulation are

highly responsive to chronic fasting and refeeding.

Taken together, these results demonstrate that INT1 and INT2-9 cells engage distinct

translational programs during fasting and refeeding. The divergence is particularly evident under

chronic conditions, where INT1 uniquely activates stress- and pathogen-related programs,

suggesting specialized roles for INT1 in maintaining homeostasis under chronic conditions.

However, because a large portion of the INT1 DEGs remain unassigned by WormCat, the full

extent of INT1-specific regulatory programs may not be fully captured by gene set enrichment

analyses.

**INT1 cells modulate pathogen avoidance, survival, and peptide secretion**

To begin to annotate and assign functions to the genes emerging from our datasets, we

examined the pathogen response and peptide secretion functions of the INT1 cells, and the

roles of the most highly regulated genes therein. To study pathogen response, we noted that

previous studies had implicated INT1-enriched genes in immune responses. For example,

expression of *B0024.4* is induced upon exposure to pathogenic *S. maltophilia* JV3 strain [29],

and *clec-42,* one of our INT1-upregulated genes, is exclusively expressed in INT1 at all larval

stages and mediates resistance to pathogen Bt247 [23]. To test whether INT1-upregulated

genes contribute to immune responses, we performed the PA14 avoidance- and PA14 slow-

killing assays [30] following INT1-specific RNAi knockdown of *clec-86* and *clec-160* RNAi.

Interestingly, animals with INT1-specific knockdown of either gene exhibited impaired PA14

avoidance (Fig 6A and 6B) and significantly increased survival on PA14 (Fig 6C). These

findings suggest that INT1 cells execute specialized, gene-dependent immune functions during

pathogen exposure.

From the WGCNA, we noted that some modules (brown, magenta, turquoise) contained

genes with an undulating change in expression levels in response to fasting and refeeding

under both acute and chronic conditions (Fig 6D-F). These genes could be classified as "fasted

up/refed down" or "fasted down/refed up". WormCat analysis of brown module genes identified

enrichment for "Ribosome" and "Metabolism", consistent with categories previously linked to

ribosomal subunit genes (S6B, S7A, and S8A Figs). Notably, modules magenta and turquoise

contained "fasted up/refed down" genes, but the induction in INT2-9 cells after acute fasting was

weaker in the turquoise module, suggesting these undulating genes were more INT1-specific

(Fig 6E and 6F).

Given our prior finding that the gut peptide INS-7 is secreted from INT1 cells in response

to acute fasting [6], we next investigated whether INT1-specific undulating genes could

modulate this response. Using a strain enabling INT1-specific RNAi (*sid-1;Pges-1ΔB::sid-1*)

crossed into an INS-7 secretion reporter, we confirmed that vector RNAi-treated animals

displayed the expected undulating INS-7 secretion levels upon acute fasting and refeeding (Fig

6G), as previously described REF. We next targeted two INT1-specific undulating genes, *aakg-*

*4* and *fmo-2*, which showed the largest absolute fold changes across acute fasting and

refeeding for INT1-specific RNAi (Fig 6H and 6J). *aakg-4* is the gene encoding one of the five $\gamma$

subunits of the AMP-activated protein kinase (AMPK) in *C. elegans* [31]. However, the

physiological role of *aakg-4* in the intestine is unknown. Remarkably, INT1-specific *aakg-4*

knockdown prevented INS-7 secretion from returning to basal levels after refeeding, indicating a

requirement for *aakg-4* in resetting INS-7 secretion and suggesting that INS-7 release from

INT1 cells is subject to energy-dependent regulation, consistent with its modulation by fasting

and refeeding (Fig 6I). The second gene *fmo-2*, encoding flavin-containing monooxygenase-2,

is known to promote longevity under hypoxia and dietary restriction [32]. FMOs contain the

prosthetic group that binds $FAD^+$ and utilize the cofactor NADPH together with $O_2$ to catalyze

substrate oxygenation, generating $NADP^+$ as a by-product [33]. Animals subjected to INT1-

specific *fmo-2* knockdown displayed a normal fasting-induced INS-7 secretion, but exhibited

reduced basal INS-7 secretion and impaired refeeding response (Fig 6K). Given that

NADPH/NADP$^+$ ratio is a central determinant of cellular redox state and energy metabolism [34],

this result further supports the notion that regulation of INS-7 secretion in INT1 cells is coupled

to energy-sensing metabolic pathways.

Together, these results demonstrate that INT1 cells integrate immune and metabolic

cues. By regulating pathogen avoidance and survival through lectin genes and modulating INS-

7 secretion via energy-responsive genes such as *aakg-4* and *fmo-2,* INT1 cells function as a

specialized intestinal subset that senses energy states and coordinates gut peptide secretion to

maintain organismal homeostasis.

## Pyruvate is crucial for regulating INS-7 peptide secretion from INT1 cells

We were intrigued by the findings with *aakg-4* and *fmo-2*, which suggested that cellular energy

status may govern the refeeding response of INS-7 secretion. Notably, the effect does not

appear to be mediated by changes in intestinal fat content, as lipid levels remain unaltered

during acute fasting and refeeding (Fig 1B and 1C). These observations led us to hypothesize

that INT1 cells may integrate specific sensory cues to reset INS-7 secretion in response to food

availability. To explore this possibility, we first evaluated the contribution of mechanosensory

inputs. Fasted worms refed with 0.5 $\mu$m latex beads, comparable in size to *E. coli* [35, 36], failed

to restore INS-7 secretion to basal levels, whereas refeeding with OP50 fully normalized INS-7

secretion (Fig 7A), suggesting that mechano-stimulus alone was not sufficient to induce the

refeeding response of INS-7 secretion. Because *C. elegans* can detect secreted molecules from

bacteria and trigger a stress response [37], we next exposed fasted animals to cell-free OP50

culture supernatant [37, 38]. Supernatant alone likewise failed to reset INS-7 secretion (Fig 7A).

A combination of beads and supernatant was similarly ineffective (Fig 7A). In contrast, UV-killed

OP50 completely reinstated basal INS-7 secretion (Fig 7A), indicating that chemicals or

nutrients from bacteria, rather than mechano-stimulus or secreted molecules, mediate the

refeeding response.

Next, we set out to pinpoint the bacterial nutrient or chemical most critical for regulating

INS-7 secretion during refeeding. Based on the requirement of *aakg-4* and *fmo-2* for INS-7

secretion dynamics, we tested whether restoring cellular energy states after fasting could return

INS-7 secretion to baseline. Glucose serves as a primary cellular energy source, undergoing

glycolysis to generate pyruvate, which in turn fuels downstream metabolic pathways to produce

energy. However, refeeding fasted worms with 2% glucose, a concentration considered a high-

glucose diet for *C. elegans* [47], failed to normalize INS-7 secretion, which remained elevated

(Fig 7B). In contrast, remarkably, refeeding with 100 mM pyruvate, a dose effective in prior

studies [47], fully recapitulated the refeeding response observed with OP50 (Fig 7B). This effect

was dose-dependent (S9A Fig). Thus, pyruvate, a key molecule derived from glycolysis and a

precursor metabolite for numerous biochemical processes, is sufficient to substitute for bacterial

food in restoring INS-7 secretion to baseline after fasting.

After entering the cell, pyruvate can be converted to lactate or alanine in the cytosol or

transported into the mitochondrial matrix via the mitochondrial pyruvate carrier (MPC) [39, 40] to

enter the citric acid cycle, which generates energy [41]. INT1-specific RNAi of the two *C.*

*elegans* MPC genes (*mpc-1* and *mpc-2*) abolished the pyruvate-induced restoration of INS-7

secretion (Fig 7C and 7D), demonstrating that mitochondrial import of pyruvate is required.

Within the matrix, pyruvate is converted to acetyl-CoA by pyruvate dehydrogenase (*pdha-1* and

*pdhb-1* in *C. elegans*) or to oxaloacetate by pyruvate carboxylase (*pyc-1* in *C. elegans*). INT1-

specific knockdown of either pathway suppressed fasting-induced INS-7 secretion and

abolished secretion dynamics (Fig 7E and 7F). Since chronic *pdha-1* and *pdhb-1* knockdown

lowered basal INS-7 secretion level, we acutely inhibited pyruvate dehydrogenase during

refeeding with the specific inhibitor 3-fluropyruvate (3-FP); this intervention prevented

normalization of INS-7 secretion (S9B Fig), suggesting that the conversion of pyruvate to acetyl-

CoA is required for the refeeding response of INS-7 secretion.

If intracellular pyruvate is the crucial determinant of INS-7 secretion, we predicted that

maintaining pyruvate levels in the INT1 cells during fasting should suppress the fasting-induced

INS-7 secretion. Indeed, worms fasted on NGM agar supplemented with 100 mM pyruvate

showed no induction of INS-7 secretion (Fig 7G), suggesting pyruvate alone is sufficient to

suppress fasting-induced INS-7 secretion. Collectively, these findings identify mitochondrial

pyruvate metabolism as the key signal sensed by INT1 enteroendocrine cells to couple

nutritional availability with the fluctuation in INS-7 peptide secretion during fasting and refeeding.

## Discussion

In the human intestine, both epithelial and immune cell compositions shift dramatically along the

anterior-posterior axis, with region-specific cellular neighborhoods for epithelial, stromal, and

immune cells contributing to distinct physiological functions [42]. In contrast, the *C. elegans*
intestine comprises only 20 epithelial cells and lacks resident immune cells, raising the question
of whether spatially defined molecular and functional heterogeneity also exists in this simplified
but powerful model system. Our findings, together with prior observations, support the view that
*C. elegans* intestinal subsets are molecularly and functionally distinct.

In the present study, we utilized INT1- and INT2-9-specific promoters to perform TRAP-
Seq analyses under fasting and refeeding conditions. While comparisons between INT1 and
INT2-9 cells revealed marked differences, this approach likely underrepresents heterogeneity
among the INT2-9 cells themselves. Prior studies demonstrated that posterior intestinal cells
initiate the rhythmic defecation motor program [43, 44], and our dataset identified *npr-28*, which
is specifically expressed in the posterior intestinal cells INT7-9 (Fig 2J), further underscoring
molecular diversity across intestinal segments. Future identification of promoters with more
restricted expression within the INT2-9 subset will be crucial for functionally resolving the spatial
map of the *C. elegans* intestine.

We previously identified INT1 as enteroendocrine cells that secrete the gut peptide INS-
7 preferentially during acute food withdrawal to suppress neuronally stimulated fat loss [6].
Additional lines of evidence also point to the specialization of the anterior-most INT1 cells. First,
acid phosphatase staining for intestinal brush border could not be detected in INT1 cells [15, 45].
Second, Methyl red staining showed strong red staining of acidic vacuoles in INT1 cells. From
INT2 to INT9 cells, vacuoles were stained gradually from orange-yellow (neutral-alkaline) to
yellow (alkaline) with methyl red [45], suggesting the pH of intracellular vacuoles is lower in
INT1 cells. Third, ultrastructural analysis highlighted enrichment of rough ER, Golgi, and
mitochondria in INT1, consistent with a secretory function, whereas INT2-9 cells preferentially
accumulate yolk and lipid vacuoles for storage [46]. Fourth, components of a cullin-RING
ubiquitin ligase complex critical for thermotolerance and the intracellular pathogen response
(IPR), including CUL-6 and SKR-5, were found to be specifically expressed in INT1 cells [47],
suggesting that INT1 cells may play an integral role in proteostasis responses to intracellular
pathogen infection. Collectively, these findings establish INT1 as a specialized cell-type that
integrates endocrine, stress, and immune functions within the *C. elegans* intestine.

We uncovered pyruvate as a crucial metabolite that functions as a true surrogate for
bacterial food to regulate the secretion of the endogenous insulin antagonist INS-7 [6].
Pyruvate must enter the TCA cycle to elicit this effect, suggesting that either actual metabolism
of pyruvate is required, or that downstream metabolites may also act as mediators of this

process. In other systems, TCA cycle intermediates such as acetyl-CoA, itaconate, and L-2-

hydroxyglutarate (L-2-HG) modulate immune responses [48], while succinate and fumarate

regulate hypoxia sensing upon cytosolic release [49]. A similar mechanism may underpin the

communication between mitochondria and the nucleus in *C. elegans*, exemplified by the

translocation of ATFS-1 during the activation of the mitochondrial unfolded protein response [50].

Thus, pyruvate metabolism may provide a metabolic checkpoint that couples nutrient availability

to enteroendocrine signaling.

Many bacterial species including *Y. enterocolitica*, *C. freundii*, *E. aerogenes* and *E. coli*

excrete pyruvate into the extracellular environment during nutrient-replete conditions [51, 52]

and reabsorb extracellular pyruvate during nutrient insufficiency [53, 54]. Thus, the presence of

extracellular pyruvate may serve as an indicator of the nutrient abundance in bacterial

environments, a sensing mechanism that *C. elegans* may have co-evolved by leveraging the

INT1 cells to couple environmental conditions with host physiology. It was reported that *C.*

*elegans* exhibits chemotaxis to volatile pyruvic acid via AWC neurons [55]. However, the

molecular mechanisms underlying both INT1 and neuronal sensing of pyruvate remain

uncharacterized. Thus both intracellular and extracellular bacterial pyruvate appear to regulate

the metabolism or behavior of *C. elegans*, indicating that pyruvate is a crucial signaling

molecule.

In summary, we expect our TRAP-Seq datasets presented here to provide the scientific

community with valuable resources for investigating the dynamic translatomes of specific

subsets of intestinal cells in response to fasting and refeeding. More broadly, the discovery of

cell-type heterogeneity within the *C. elegans* intestine highlights the evolutionary conservation of

spatially organized intestinal biology and lays a foundation for future mechanistic studies of

metabolic and immune regulation in the intestine.

**Limitations of the TRAP-Seq in the current study**

Not all previous studies have demonstrated a strong correlation between mRNA expression

levels and protein abundance [56-58]. TRAP-Seq provides a more precise assessment of gene

expression by detecting actively translating mRNAs. Specifically, TRAP-Seq better reflects the

translational responses under various conditions, such as temperature fluctuations or fasting.

Moreover, by employing tagged ribosomal proteins, TRAP-Seq enables the selective purification

of polysomes without the need for cell dissociation or sorting, thereby reducing RNA

contamination from neighboring cells or tissues.

To obtain sufficient ribosome-mRNA complexes for RNA sequencing, we generated

strains with an integrated extrachromosomal array expressing HA-tagged RPL22 in either INT1

or INT2-9 cells. While this strategy enables robust affinity purification, the elevated expression

of HA-tagged RPL22 in this system could potentially introduce unanticipated effects on

transcript utilization or translational bias. Nevertheless, because all analyses rely on internal

referencing, the differential gene expression signatures remain valid. In addition, the random

integration of multi-copy transgenes into unknown genomic loci may alter the expression of

adjacent genes, leading to additional variability. Importantly, we did not observe any changes in

growth rate, fertility or fecundity in the TRAP-seq transgenic lines, suggesting that potential

confounding effects are minimal.

TRAP has traditionally tagged proteins of the 60S large ribosomal subunit, including

RPL10A [10, 11, 59-62] and RPL22 [13, 63-65], to enable tissue- or cell type-specific capture of

polysome-mRNA complexes followed by RNA sequencing. A key distinction among previous

studies lies in the choice of ribosomal protein for tagging. Both RPL10A and RPL22 are

evolutionarily conserved ribosomal proteins. In *C. elegans*, RPL10A [62, 66, 67] and RPL22 [13,

65, 68] have been successfully used for tagging in TRAP-Seq experiments, yielding robust

results. However, recent findings in mammalian systems indicate that RPL10A is significantly

sub-stoichiometric in polysomes of mouse embryonic stem cells [69], suggesting it is not

associated with all actively translating mRNAs. Similarly, in *C. elegans*, RPL10A shows limited

expression in the germline and weak association with ribosomes, impairing the co-purification of

translating mRNAs from the germline [68]. These observations suggest that tagging RPL10A in

TRAP experiments may not capture the complete translatome in specific cell types or tissues.

Further, *Nousch et al.* recently demonstrated that RPL-4 and RPL-9 are more effective than

RPL10A and RPL22 for purifying polysome-mRNA complexes from germ cells in *C. elegans*

[68], suggesting that the choice of ribosomal proteins for tagging in TRAP-Seq experiments may

vary depending on the tissue of interest. Specifically, certain ribosomal proteins may offer

improved specificity and efficiency for isolating tissue-specific mRNAs. Importantly, the RPL22-

based TRAP in our experiments achieved robust immunoprecipitation efficiency (about 40% of

input) for intestinal polysomal mRNAs (S4 table), thereby validating our approach. Future

studies may further investigate whether alternative ribosomal proteins could provide enhanced

isolation of translating mRNAs in different tissues including the intestine.

## Materials and methods

### *C. elegans* maintenance and strains

Worms were cultured on nematode growth medium (NGM) agar plates with *Escherichia coli* strain OP50 at 20°C as described [70]. Bristol strain N2 was obtained from the *Caenorhabditis* Genetic Center (CGC) and used as the wild-type strain. All transgenic strains used in this study are listed in S1 Table. Worms were synchronized for Oil Red O staining by hypochlorite treatment, after which hatched L1 larvae were seeded on plates with the appropriate bacteria; worms were synchronized for INS-7 peptide secretion assay, PA14 avoidance and slow killing assays by letting gravid adult worms lay eggs on plates with the appropriate bacteria for 1.5 hr. All experiments were performed on Day 1 adults.

### Oil Red O staining

Oil Red O staining was performed as described [71]. Worms were harvested with phosphate-buffered saline (PBS) and incubated on ice for 10 minutes. Worms were fixed as described [71]. Following fixation, worms were stained in filtered Oil Red O (Thermo Scientific) working solution (60% Oil Red O in isopropanol: 70% water) overnight. Approximately 2000 worms were fixed and stained for all conditions within one experiment. We visually examined about 100 worms on slides and then blindly chose 20 worms for imaging. All experiments were repeated at least three times.

### Image acquisition and quantitation

Oil Red O-stained worms were imaged using 10X objective on a Zeiss Axio Imager microscope. Images were acquired with the Software AxioVision (Zeiss). Lipid droplet staining in the first four pairs of intestinal cells was quantified using ImageJ software (NIH). All reported results were consistent across biological replicates. Fluorescent images of mNeonGreen reporters were acquired with software AxioVision (Zeiss) using a 10X objective on a Zeiss Axio Imager microscope. Fluorescent images of reporters for INS-7 secretion were acquired with software AxioVision (Zeiss) using a 20X objective on a Zeiss Axio Imager microscope. The first pair of coelomocytes was imaged. mCherry fluorescence intensity in one of the two imaged

507  coelomocytes was quantified and normalized to the surface area of the coelomocyte (*Punc-*

508  *122::GFP*) as previously described and validated [72]. Within each experiment, at least 15

509  worms from each condition were quantified using ImageJ software (NIH).

## 511  **Cloning and transgenic strain construction**

For generating *Pges-1ΔB::rpl-22-3xHA* and *Ppho-1::rpl-22-3xHA* constructs, promoters and *rpl-*

*22-3xHA* were amplified with standard PCR techniques and subcloned into donor vectors using

Gateway Cloning Technology$^{TM}$ (Life Technologies). All plasmids for NeonGreen reporters were

generated by Gibson Assembly$^{TM}$. Primers used for cloning in this study are listed in S3 Table.

Transgenic strains were generated by injecting plasmids into the germline of wild-type worms

followed by visual selection of co-injection markers (*Pmyo-2::mCherry*, *Pges-1ΔB::GFP* or

*Ppho-1::mCherry*) under the fluorescence microscope. The INS-7 secretion line was previously

developed and validated [73]. For *INT1::TRAP* strain, N2 worms were injected with 10 ng/µL of

the *Pges-1ΔB::rpl-22-3xHA* plasmid, 20 ng/µL of the *Pges-1ΔB::GFP* plasmid, and 70 ng/µL of

an empty vector to bring the final concentration of injection mix to 100 ng/µL. For *INT2-9::TRAP*

strain, N2 worms were injected with 25 ng/µL of the *Ppho-1::rpl-22-3xHA* plasmid, 50 ng/µL of

the *Ppho-1::mCherry* plasmid, and 25 ng/µL of an empty vector to bring the final concentration

of injection mix to 100 ng/µL. A transgenic line with high transmission rate and consistent

expression was integrated using the Stratalinker UV Crosslinker 2400 (Stratagene) and

backcrossed four times before experimentation. For other microinjections, we injected the

animals with 30 ng/µL of the desired plasmid, 10 ng/µL of the *Pmyo-2::mCherry* and 60 ng/µL

empty vector to maintain a final injection mix concentration of 100 ng/µL. Two lines were

selected for experimentation based on the transmission rate and consistency of expression.

## 531  **RNAi**

532  RNAi experiments were performed as described [74, 75]. Carbenicillin-IPTG plates were seeded

533  with HT115 bacteria containing the empty or the relevant RNAi clone four days before seeding

534  larvae.

## 536  **PA14 avoidance assay**

The PA14 avoidance assay was performed as previously described [76]. *Pseudomonas*

*aeruginosa* strain PA14 was cultured by inoculating a single colony into 2 mL of LB broth and

incubating at 37 °C for 6 hr. 20 μL of the culture was then seeded onto 3.5 cm slow-killing (SK)

plates and incubated at 37 °C for 15 hr. For each assay, 30 synchronized Day 1 gravid adults

(containing 6-10 eggs) grown on RNAi bacteria were cleaned and transferred to the agar

surface outside the PA14 lawn. The number of animals on and off the PA14 lawns was scored

after 8 h. Three plates were used per RNAi treatment in each experiment.

**PA14 slow killing assay**

PA14 slow killing assay was performed as described [30]. Briefly, overnight cultures of

*Pseudomonas aeruginosa* strain PA14 were seeded onto slow-killing (SK) plates and incubated

at 37°C for 24 hr, followed by 25°C for an additional 24 hr. Synchronized animals at the closed

vulva stage subjected to RNAi treatment were transferred to 3.5 cm SK plates seeded with

PA14 (30 animals per plate), with three plates per RNAi treatment. Plates were maintained at

25°C, and worm survival was monitored until all animals had died. Survival curves were

analyzed using GraphPad Prism, and statistical significance was determined by the log-rank

test.

**Quantitative RT-PCR**

Total RNA was extracted from Day1 adults using TRIzol reagent (Invitrogen). Genomic

DNA was removed using an RNase-free DNase set (QIAGEN). cDNA was prepared

using iScript Reverse Transcription Supermix for RT-qPCR kit (Bio-Rad) according to

the manufacturer's instructions. Quantitative RT-PCR was performed using the

SsoAdvanced Universal SYBR Green Supermix (Bio-Rad) following the manufacturer's

instructions. Data were normalized to pan-actin (*actin-1, -3, -4*) mRNA. Primer

sequences used in this study are listed in S3 Table.

**Translating ribosome affinity purification**

The protocol was adapted from reference [13] and described in detail in S1 Protocol. Briefly,

transgenic worms containing the Ribo-TRAP constructs were made to drive expression of *rpl-*

*22-3xHA* in INT1 and INT2-9. Age-matched transgenic worms for each experimental condition

were washed, flash-frozen, and lysed to release ribosomes and cellular components in the

presence of cycloheximide to pause translation as well as RNAase and protease inhibitors.

Immunoprecipitation of ribosomes from our desired tissue was performed with a monoclonal

anti-HA antibody and protein G-beads. Nascent RNA was eluted from ribosomes with beta

mercaptoethanol (BMP). Percent IP RNA compared to input, on average less then 1%, was

used to determine if the IP was successful.

## RNA purification

576 RNA from the IP, flowthrough, and input was purified with Agilent Absolutely RNA Nanoprep

577 Kit$^{TM}$ according to the manufacturer's instructions including the on-column DNase treatment.

578 Final RNA isolates were eluted in 15 μl and used to make diluted aliquots for sequencing,

579 validation, and QC.

## RNA quality assessment

We tested for RNA degradation and quality by running the input, flowthrough, and IP samples

on an Agilent Bioanalyzer chip (nano or pico, depending on concentration). Because IP samples

had a small concentration it was difficult to accurately measure the RNA integrity number (RIN)

in all conditions ,but the quality of RNA for these unknown samples could be extrapolated from

the flowthrough and input samples. In all conditions the average RIN across sample types was

greater than 7.5 indicating a RNA integrity sufficient for sequencing, and in the majority of

conditions it was greater than 9 indicative of high-quality samples.

## RNAseq library prep, sequencing, and alignment

Total RNA samples were prepared into RNAseq libraries using the NEBNext® Ultra™

Directional RNA Library Prep Kit for Illumina® following the manufacturer's

recommended protocol. Briefly, for each sample 100ng total RNA was polyA-selected,

converted to double stranded cDNA followed by fragmentation and ligation of sequencing

adapters. The libraries were then PCR amplified 12 cycles using barcoded PCR primers,

purified and size selected using AMPure XP Beads before loading onto an Illumina NextSeq500

for 75 base single read sequencing. Reads were aligned to *C. elegans* genome (WBcel235/r67)

using STAR (version 2.6.1d) [77], and gene counts were produced using featureCounts (version

1.6.4) [78]. On average, 10.6 million unique single-end (SE) reads aligned per INT1 sample and

19.5 million unique SE reads per INT2-9 sample, differences were due to the larger RNA input

for INT2-9 samples. Data quality was assessed using FastQC (version 0.11.8) [79]. Raw RNA-

seq FASTQ files and a gene count matrix are available on GEO (GSE306302). Bioinformatic

analysis was performed using R programming language (version 4.4.2) (R Development Core

Team, 2015) and RStudio integrated development environment (version 2024.12.1) (Team R,

2018). Plots were generated using ggplot2 R package (version 3.5.2).

**Differential expression analysis**

Differential expression analysis was performed using the DESeq2 R package (version 1.46.0)

[80] which models the gene expression counts with a negative binomial distribution while

adjusting for library size and dispersion. Differentially expressed genes were identified as having

adjusted p-value less then 0.05 using a Benjamini-Hochberg false discovery rate correction.

DESeq2 variance stabilized counts were used to generate normalized count matrices that were

used for plotting expression data of individual genes and those averaged across WGCNA

modules. PCA was performed using the top 500 variable genes from variance stabilized counts

output in DESeq2.

**WGCNA**

WGCNA was performed with the WGCNA R package (version 1.73) [16] to identify modules of

co-expressed genes and their associations with external traits in the analysis. Prior to analysis,

samples with low expression values were removed to reduce variation associated with batch, a

30 min fasting and 30 min refeeding sample were removed from the *Pges-1ΔB* experiment

based on this criteria. DESeq2 variance stabilized counts were used as the input and only

genes in the $95^{th}$ percentile of variance were kept to reduce noise (n = 2,338). A soft-

thresholding power of 10 was selected using the scale-free topology criterion to approximate

scale-free network topology. A signed adjacency matrix was constructed and transformed into a

topological overlap matrix (TOM), followed by hierarchical clustering to detect modules of highly

correlated genes. Modules were identified using dynamic tree cutting with a minimum module

size of 30 genes and subsequently merged based on eigengene similarity (cut height = 0.25).

Module eigengenes (MEs) were correlated with external traits (e.g., line and fasting condition),

and Pearson correlation coefficients along with associated p-values were computed. Modules

showing significant trait associations were further investigated for biological relevance and

functional enrichment. Module GO enrichment analysis was performed using the ClusterProfiler

R package version (version 4.14.6)[81]. For GO analysis, we reported terms with FDR < 0.1.

The test set of module genes was compared against the background set of genes in the

WGNCA analysis passing the variance criteria.

**Statistics**

Wild-type animals were included as controls for every experiment. Error bars represent standard

error of the mean (SEM). Student's *t*-test and one-way ANOVA were used as indicated in the

Fig legends. All statistical analyses were performed in GraphPad Prism 9.3 (GraphPad

Software). Appropriate multiple comparison corrections were used for ANOVAs.

**Acknowledgements**

This work was supported by research grants to S.S. from the NIH/NIDDK (R01 DK124706) and

NIH/NIA (R01 AG056648). This work was supported by the resources and staff at the Scripps

Research Genomics Core Facility. C.L. was supported by a Dorris Scholar Award from the

Dorris Neuroscience Center, The Scripps Research Institute. Elements of Fig 1F were created

with BioRender.com.

653 **FIGURES and FIGURE LEGENDS**

**Figure 1**

Fig 1.

Conditions

**and workflow for INT1- and INT2-9-specific translatomics.**

(A) Representative images of wild-type animals bearing integrated *INT1::GFP* and *INT2-9::mCherry* transgenes. Upper panel, GFP expression in INT1 cells and mCherry expression in INT2-9 cells; lower panel, merge with DIC. Scale bar, 100 μm. (B) Representative images of wild-type animals in fed, 30 min fasted, and 30 min refed states fixed and stained with Oil Red O. (C) Fat content was quantified for each condition and expressed as a percentage of wild-type animals in fed state ± SEM (n=20). [ns]p>0.05 by one-way ANOVA (Tukey). (D) Representative images of wild-type animals in fed, 180 min fasted, 30 min refed, 90 min refed, and 180 min refed states fixed and stained with Oil Red O. (E) Fat content was quantified for each condition

681 and expressed as a percentage of wild-type animals in fed state $\pm$ SEM (n=18-20). [***]p<0.001,

682 [ns]p>0.05 by one-way ANOVA (Sidak). (F) 100000 *C. elegans* expressing *INT1::RPL-22::3xHA*

683 or *INT2-9::RPL-22::3xHA* were grown to day 1 adults. Worms were fasted and refed for the time

684 indicated and harvested for TRAP-Seq.

**Figure 2**

**Fig 2. Identification of the fundamental differences between INT1 and INT2-9 translatomes.**

(A) Principal component analysis showing translatomic variability across the first two principle components of TRAP-Seq data from INT1 and INT2-9 promoters under both acute and chronic conditions (fed, fasted, and refed). (B) Cluster dendrogram produced by average linkage hierarchical clustering in WGCNA. The color row under the dendrogram indicates the module assignment. (C) Pearsons correlations of TRAP-seq conditions with WGCNA modules. Each

row represents a WGCNA module and its means correlation to the indicated trait across

replicates.The color of each cell indicates the mean pearsons correlation coefficient between

the module and the condition; red represents a positive correlation and blue represents a

negative correlation. (D) Average expression of red modules genes in the TRAP-seq conditions.

The y-axis shows mean variance stabilized expression across all genes and replicates per

condition with standard deviation indicated by the error bars. (E) WormCat visualization of

categories enriched in genes of module red. Categories 1 are all bold uppercase; Categories 2

are capitalized; Categories 3 are gray italics. The size of the bubbles indicates the gene counts

in the category and the color of the bubble represents the adjusted p-value. (F) Volcano plot of

differentially expressed genes for INT1 fed versus INT2-9 fed. High confidence differentially

expressed genes are labeled. The horizontal dashed line indicates an adjusted p-value cutoff of

0.05. (G) WormCat visualization of categories enriched in genes upregulated in INT1 cells

versus INT2-9 cells in the fed state. Categories 1 are all bold uppercase; Categories 2 are

capitalized; Categories 3 are gray italics. The size of the bubbles indicates the gene counts in

the category and the color of the bubble represents the adjusted p-value. (H) A hierarchical

clustering tree of significantly enriched GO terms (Biological Process, BP) in the upregulated

genes for INT1 fed versus INT2-9 fed by ShinyGO. GO terms with many shared genes are

clustered together. Larger dots indicate more significant adjusted p-values. (I) Fluorescent

images of transgenic animals bearing *PB0024.4::mNeonGreen, Pclec-86::mNeonGreen*, *Pclec-*

*160::mNeonGreen*, *PF01D5.5::mNeonGreen or PF01D5.1::mNeonGreen* transgenes. The white

arrowhead indicates *mNeonGreen* expression in INT1 cells. Scale bar, 100 μm. (J) Fluorescent

image of a transgenic animal bearing *Pnpr-28::mNeonGreen* transgene. Scale bar, 100 μm.

**Figure 3**

**Fig 3. Identification of the differences between INT1 and INT2-9 translatomes under acute and chronic conditions.**

(A, D, G, J) Volcano plot of differentially expressed genes for INT1 versus INT2-9 in acute fasted, acute refed, chronic fasted and chronic refed states. High confidence differentially expressed genes. The horizontal dashed line indicates an adjusted p-value cutoff of 0.05. (B, E, H, K) WormCat visualization of categories enriched in genes upregulated in INT1 cells versus

748 INT2-9 cells in acute fasted, acute refed, chronic fasted and chronic refed states. Categories 1

749 are all bold uppercase; Categories 2 are capitalized; Categories 3 are gray italics. The size of

750 the bubbles indicates the gene counts in the category and the color of the bubble represents the

751 adjusted p-value. (C, F, I, L) A hierarchical clustering tree of significantly enriched GO terms

752 (Biological Process, BP) in the upregulated genes for INT1 versus INT2-9 in acute fasted, acute

753 refed, chronic fasted and chronic refed states by ShinyGO. GO terms with many shared genes

754 are clustered together. Larger dots indicate more significant adjusted p-values.

**Figure 4**

**A**    INT1 vs. INT2-9 Upregulated Stress Response Genes

**B**    INT1 vs. INT2-9 Upregulated Enzymes

**C**    INT1 vs. INT2-9 Upregulated Membrane Proteins

**D**    INT1 vs. INT2-9 Upregulated C-type Lectins

**E**    INT1 vs. INT2-9 Upregulated ShKT proteins

**F**    INT1 vs. INT2-9 Downregulated Enzymes

**G**    INT1 vs. INT2-9 Downregulated Membrane Proteins

**H**    INT1 vs. INT2-9 Downregulated C-type Lectins

**I**    INT1 vs. INT2-9 Upregulated Stress Response Genes

**J**    INT1 vs. INT2-9 Upregulated Enzymes

**K**    INT1 vs. INT2-9 Upregulated Membrane Proteins

**L**    INT1 vs. INT2-9 Upregulated C-type Lectins

**M**    INT1 vs. INT2-9 Upregulated ShKT proteins

**N**    INT1 vs. INT2-9 Downregulated Enzymes

**O**    INT1 vs. INT2-9 Downregulated Membrane Proteins

**P**    INT1 vs. INT2-9 Downregulated C-type Lectins

**Q**    INT1 vs. INT2-9 Downregulated ShKT proteins

**R**    INT1 vs. INT2-9 Overlapped Upregulated ShKT proteins

**Fig 4. Comparisons of InterPro categories for INT1 versus INT2-9 under acute and**
**chronic conditions.**

(A-H) Venn diagrams showing the number of upregulated or downregulated genes in specific
categories for INT1 versus INT2-9 under the three acute conditions (fed, fasted, and refed). (I-Q)
Venn diagrams showing the number of genes in specific categories for INT1 versus INT2-9
under the three chronic conditions (fed, fasted, and refed). (R) Venn diagram showing the
overlap between overlapped upregulated ShKT proteins for INT1 versus INT2-9 under acute
and chronic conditions.

**Figure 5**

**A** Acute fasted vs. fed

n=139      n=656
n=612      n=270

INT2-9 [log$_2$ fold change] vs INT1 [log$_2$ fold change]

**B** Chronic fasted vs. fed

n=22      n=767
n=710      n=46

INT2-9 [log$_2$ fold change] vs INT1 [log$_2$ fold change]

**C**

| Expression | Differentially Expressed Genes | 30min Fasted | 30min Refed | 180min Fasted | 90min Refed |
|---|---|---|---|---|---|
| *INT1::TRAP* | Upregulated | 41 | 1 | 510 | 362 |
| *INT1::TRAP* | Downregulated | 17 | 19 | 457 | 427 |
| *INT2-9::TRAP* | Upregulated | 134 | 141 | 1519 | 921 |
| *INT2-9::TRAP* | Downregulated | 72 | 130 | 1195 | 759 |

**D**

Fasted Up: 45, 264, 1, 7, 195, 18, 0, 29, 1233, 1, 10, 41, 4, 1, 6

Fasted Down: 22, 65, 3, 3, 357, 3, 0, 23, 791, 0, 8, 13, 1, 0, 2

Refed Up: 46, 94, 0, 2, 230, 1, 0, 36, 598, 0, 0, 57, 0, 0, 0

Refed Down: 56, 165, 0, 10, 211, 10, 0, 35, 478, 1, 2, 27, 3, 0, 3

INT1::TRAP Acute (pink)   INT1::TRAP Chronic (green)
INT2-9::TRAP Acute (blue)   INT2-9::TRAP Chronic (yellow)

**E** Chronic Fasted Up

INT1   INT2-9

**METABOLISM**
Lipid
*CoA*
**STRESS RESPONSE**
**SIGNALING**
Small GTPase
**TRANSCRIPTION FACTOR**
NHR
*NHR*

**F** Chronic Refed Up

INT1   INT2-9

**STRESS RESPONSE**
Pathogen
Detoxification
*GST*
**RIBOSOME**
Subunit
*subunit*

Count: 100, 75, 50, 25, 10, 5, 2

*P*-value: 10$^{-40}$, 10$^{-10}$, 10$^{-5}$, 0.001, NS

**Fig 5. Comparison of fasting and refeeding responses between INT1 and INT2-9 cells under acute and chronic conditions.**

(A, B) Scatterplots comparing the acute and chronic fasting response genes in INT1 vs INT2-9 differential expression. The top 1000 DEGs by significance were selected for each line and fasting response to be compared, with a regression line fitted to compared INT1 vs INT2-9 response. (C) Table of differentially expressed genes (FDR < 0.05) under acute and chronic conditions for INT1 and INT2-9 datasets. (D) Venn diagrams showing the number of differentially expressed genes (FDR < 0.05) in fasting and refeeding conditions for INT1 and INT2-9 datasets. (E, F) WormCat visualization of categories enriched in upregulated genes in INT1 cells and INT2-9 cells under chronic fasting or refeeding conditions. Categories 1 are all bold uppercase; Categories 2 are capitalized; Categories 3 are gray italics. The size of the bubbles indicates the gene counts in the category and the color of the bubble represents the adjusted p-value.

**Fig 6. INT1-specific genes modulate behavior and metabolism.**

(A) The vector, *clec-86*, and *clec-160* RNAi treated *sid-1;INT1::sid-1* animals after 8 hr exposure

to PA14 lawns are shown. (B) The distribution of indicated RNAi-treated *sid-1;INT1::sid-1*

animals at hour 8 after being transferred to PA14 plates. At hour 0, a total of 30 animals were

placed near the PA14 lawn. Data are expressed as a percentage of the animals staying inside

PA14 lawn at hour 8 ± SEM (n=3, 3 plates per RNAi treatment, 30 animals on each plate).

**p<0.01 by one-way ANOVA (Dunnett). (C) The lifespan of each indicated RNAi-treated animal

after being transferred to PA14 lawn was determined by counting the number of live and dead

animals every 6hr until all animals had died. Data are plotted as the percentage of animals that

survived on any given hour relative to the number of live animals at hour 0. The number of

animals for each RNAi treatment is indicated in the figure panels. ***p<0.001 by Log-rank Test.

(D-F) Average expression of brown, magenta, and turquoise modules genes in the TRAP-seq

conditions. The y-axis shows mean variance stabilized expression across all genes and

replicates per condition and module with standard deviation indicated by the error bars. (G, I, K)

*sid-1;INT1::sid-1* animals bearing integrated *INT1::INS-7mCherry* and *CLM::GFP* transgenes

were treated with vector, *aakg-4* or *fmo-2* RNAi. INS-7 secretion levels were measured after 30

min fasting and after 30 min refeeding. The intensity of INS-7mCherry fluorescence within a

single coelomocyte was quantified and normalized to the area of CLM::GFP expression for each

RNAi treatment. Data are expressed as a percentage of the normalized INS-7mCherry

fluorescence intensity of fed animals treated with indicated RNAi $\pm$ SEM (n=21-27). *p<0.05,

**p<0.01, ***p<0.001 by one-way ANOVA (Dunnett T3). (H, J) Normalized expression plots of

*aakg-4* and *fmo-2* under all conditions in INT1 and INT2-9 cells by replicate.

Figure 7

**888  Fig. 7. Pyruvate is critical for maintaining INS-7 secretion dynamics in INT1 cells.**

(A) INS-7 secretion dynamics during fasting and re-feeding with OP50, latex beads, Day 1

OP50 supernatant, Day 1 OP50 supernatant with latex beads, and UV-killed OP50 were

determined at the indicated time points. The intensity of INS-7mCherry fluorescence within a

single coelomocyte was quantified and normalized to the area of CLM::GFP expression for each

time point. Data are expressed as a percentage of the normalized INS-7mCherry fluorescence

intensity of wild-type fed animals $\pm$ SEM (n=20-26). **p<0.01, ***p<0.001, [ns]p>0.05 by one-way

ANOVA (Dunnett T3). (B) INS-7 secretion dynamics during fasting and re-feeding with OP50,

2% glucose and 100 mM pyruvate were determined at the indicated time points. The intensity of

INS-7mCherry fluorescence within a single coelomocyte was quantified and normalized to the

area of CLM::GFP expression for each time point. Data are expressed as a percentage of the

normalized INS-7mCherry fluorescence intensity of wild-type fed animals $\pm$ SEM (n=14-26).

**p<0.01, ***p<0.001, [ns]p>0.05 by one-way ANOVA (Dunnett T3). (C-F) INS-7 secretion

dynamics in INT1-specific vector RNAi, *mpc-1* and *mpc-2* RNAi, *pyc-1* RNAi, *pdha-1* and *pdhb-*

*1* RNAi-treated animals during fasting and re-feeding with 100 mM pyruvate were determined at

the indicated time points. The intensity of INS-7mCherry fluorescence within a single

coelomocyte was quantified and normalized to the area of CLM::GFP expression for each time

point. Data are expressed as a percentage of the normalized INS-7mCherry fluorescence

intensity of vector RNAi fed animals ± SEM (n=13-24). $*p<0.05$, $**p<0.01$, $***p<0.001$, $^{ns}p>0.05$

by one-way ANOVA (Dunnett T3). (G) INS-7 secretion dynamics during fasting and fasting with

the presence of 100 mM pyruvate were determined at the indicated time points. The intensity of

INS-7mCherry fluorescence within a single coelomocyte was quantified and normalized to the

area of CLM::GFP expression for each time point. Data are expressed as a percentage of the

normalized INS-7mCherry fluorescence intensity of fed animals ± SEM (n=16-19). $***p<0.001$,

$^{ns}p>0.05$ by one-way ANOVA (Dunnett T3).

**S1 Fig. The genes differentially expressed in INT1 cells.**

(A) Module-trait relationships for selected conditions in WGCNA. Rows indicate the WGCNA
module and the columns comparative traits. The color indicates the strength of the pearsons
correlation and bracketed number the significance. The color red represents a positive
correlation; the color green represents a negative correlation. (B) GO enrichment of selected
modules in the WGCNA analysis. Size indicates the strength of the significance and color the
significance. (C-E) Plots showing the fold change of the intestinal anterior, middle and posterior
genes from Ghaddar et al. in our INT1 vs. INT2-9 Fed datasets. (F-J) Expression profiles of
*B0024.4, clec-86, clec-160, F01D5.5* and *F01D5.1* under all conditions in INT1 and INT2-9 cells.

**S2 Fig**

**A** INT1 vs. INT2-9 (Fed-Upregulated Genes) InterPro

**B** INT1 vs. INT2-9 (Fed-Downregulated Genes) InterPro

 **S2 Fig. InterPro annotations of differentially expressed genes for INT1 Fed versus INT2-9**

940 **Fed.**

941 (A, B) Bar charts of the upregulated or downregulated genes categorized by the protein

942 domains predicted by InterPro for INT1 fed versus INT2-9 fed.

**S3 Fig**

**A** INT1 vs. INT2-9 (Acute Fasted Up) InterPro

**B** INT1 vs. INT2-9 (Acute Fasted Down) InterPro

**C** INT1 vs. INT2-9 (Acute Refed Up) InterPro

**D** INT1 vs. INT2-9 (Acute Refed Down) InterPro

**S3 Fig. InterPro annotations of differentially expressed genes for INT1 versus INT2-9 under acute conditions.**

972    (A, B) Bar charts of the upregulated or downregulated genes categorized by the protein

973    domains predicted by InterPro for INT1 30 min fasted versus INT2-9 30 min fasted. (C, D) Bar

974    charts of the upregulated or downregulated genes categorized by the protein domains predicted

975    by InterPro for INT1 30 min refed versus INT2-9 30 min refed.

**S4 Fig**

**A** INT1 vs. INT2-9 (Chronic Fasted Up)
InterPro

**B** INT1 vs. INT2-9 (Chronic Fasted Down)
InterPro

**C** INT1 vs. INT2-9 (Chronic Refed Up)
InterPro

**D** INT1 vs. INT2-9 (Chronic Refed Down)
InterPro

**S4 Fig. InterPro annotations of differentially expressed genes for INT1 versus INT2-9**

**under chronic conditions.**

(A, B) Bar charts of the upregulated or downregulated genes categorized by the protein

domains predicted by InterPro for INT1 180 min fasted versus INT2-9 180 min fasted. (C, D) Bar

charts of the upregulated or downregulated genes categorized by the protein domains predicted

by InterPro for INT1 90 min refed versus INT2-9 90 min refed.

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

**S5 Fig. WormCat annotations of differentially expressed genes under acute fasting and refeeding conditions for INT1 and INT2-9 cells.**

(A, B) WormCat visualization of categories enriched in differentially expressed genes in INT1 cells and INT2-9 cells under acute fasting condition. (C, D) WormCat visualization of categories enriched in differentially expressed genes in INT1 cells and INT2-9 cells under acute refeeding condition. Categories 1 are all bold uppercase; Categories 2 are capitalized; Categories 3 are gray italics. The size of the bubbles indicates the gene counts in the category and the color of the bubble represents the adjusted p-value.

**S6 Fig**

**S6 Fig. WormCat annotations of differentially expressed genes under chronic fasting condition for INT1 and INT2-9 cells.**

1040  (A, B) WormCat visualization of categories enriched in differentially expressed genes in INT1

1041  cells and INT2-9 cells under chronic fasting condition. Categories 1 are all bold uppercase;

1042  Categories 2 are capitalized; Categories 3 are gray italics. The size of the bubbles indicates the

1043  gene counts in the category and the color of the bubble represents the adjusted p-value.

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

**S7 Fig**

**A**

Chronic Refed Up

**B**

Chronic Refed Down

**S7 Fig. WormCat annotations of differentially expressed genes under chronic refeeding**

**condition for INT1 and INT2-9 cells.**

(A, B) WormCat visualization of categories enriched in differentially expressed genes in INT1

cells and INT2-9 cells under chronic refeeding condition. Categories 1 are all bold uppercase;

Categories 2 are capitalized; Categories 3 are gray italics. The size of the bubbles indicates the

gene counts in the category and the color of the bubble represents the adjusted p-value.

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

**S8 Fig**

A   Module Brown (166 genes)   B   Module Turquoise (1188 genes)

1129   **S8 Fig. WormCat annotations of undulating genes in INT1 and INT2-9 cells.**

1130   (A, B) WormCat visualization of categories enriched in genes of module brown and turquoise.

1131   Categories 1 are all bold uppercase; Categories 2 are capitalized; Categories 3 are gray italics.

1132   The size of the bubbles indicates the gene counts in the category and the color of the bubble

1133   represents the adjusted p-value.

1134

1135

1136

1137

**S9 Fig**

1138

**S9 Fig. The role of pyruvate in refeeding responses of INS-7 secretion from INT1 cells.**

(A) INS-7 secretion dynamics during fasting and re-feeding with different concentrations of pyruvate (0.01 mM - 100 mM) were determined at the indicated time points. The intensity of INS-7mCherry fluorescence within a single coelomocyte was quantified and normalized to the area of CLM::GFP expression for each time point. Data are expressed as a percentage of the normalized INS-7mCherry fluorescence intensity of wild-type fed animals $\pm$ SEM (n=15-23). \*\*\*p<0.001, $^{ns}$p>0.05 by one-way ANOVA (Dunnett T3). (B) INS-7 secretion dynamics during fasting and re-feeding with 10 mM pyruvate, 10 mM 3-FP, and 10 mM pyruvate and 3-FP were determined at the indicated time points. The intensity of INS-7mCherry fluorescence within a single coelomocyte was quantified and normalized to the area of CLM::GFP expression for each time point. Data are expressed as a percentage of the normalized INS-7mCherry fluorescence intensity of wild-type fed animals $\pm$ SEM (n=17-20). \*p<0.05, \*\*\*p<0.001, $^{ns}$p>0.05 by one-way ANOVA (Dunnett T3).

1152

1153

1154

1155

1156

1157

1158

## References

1160     1.      Ghafouri S, McGhee JD. Bacterial residence time in the intestine of Caenorhabditis elegans. Nematology. 2007;9:87–91.

2.      Van Gilst MR, Hadjivassiliou H, Yamamoto KR. A Caenorhabditis elegans nutrient response system partially dependent on nuclear receptor NHR-49. Proc Natl Acad Sci U S A. 2005;102(38):13496–501.

3.      O'Rourke EJ, Ruvkun G. MXL-3 and HLH-30 transcriptionally link lipolysis and autophagy to nutrient availability. Nat Cell Biol. 2013;15(6):668–76.

4.      Jo H, Shim J, Lee JH, Lee J, Kim JB. IRE-1 and HSP-4 contribute to energy homeostasis via fasting-induced lipases in C. elegans. Cell Metab. 2009;9(5):440–8.

5.      Sulston JE, Schierenberg E, White JG, Thomson JN. The embryonic cell lineage of the nematode Caenorhabditis elegans. Dev Biol. 1983;100(1):64–119.

6.      Liu CC, Khan A, Seban N, Littlejohn N, Shah A, Srinivasan S. A homeostatic gut-to-brain insulin antagonist restrains neuronally stimulated fat loss. Nat Commun. 2024;15(1):6869.

7.      Quintin S, Aspert T, Ye T, Charvin G. Distinct mechanisms underlie H2O2 sensing in C. elegans head and tail. PLoS One. 2022;17(9):e0274226.

8.      Warner JR, Knopf PM. The discovery of polyribosomes. Trends Biochem Sci. 2002;27(7):376–80.

9.      Warner JR, Knopf PM, Rich A. A multiple ribosomal structure in protein synthesis. Proc Natl Acad Sci U S A. 1963;49(1):122–9.

10.     Heiman M, Kulicke R, Fenster RJ, Greengard P, Heintz N. Cell type-specific mRNA purification by translating ribosome affinity purification (TRAP). Nat Protoc. 2014;9(6):1282–91.

11.     Heiman M, Schaefer A, Gong S, Peterson JD, Day M, Ramsey KE, et al. A translational profiling approach for the molecular characterization of CNS cell types. Cell. 2008;135(4):738–48.

12.     Zhao J, Qin B, Nikolay R, Spahn CMT, Zhang G. Translatomics: The Global View of Translation. Int J Mol Sci. 2019;20(1).

13.     McLachlan IG, Flavell SW. Cell Type-specific mRNA Purification in Caenorhabditis elegans via Translating Ribosome Affinity Purification. Bio Protoc. 2019;9(15):e3328.

14.     Schroeder DF, McGhee JD. Anterior-posterior patterning within the Caenorhabditis elegans endoderm. Development. 1998;125(24):4877–87.

15.    Beh CT, Ferrari DC, Chung MA, McGhee JD. An acid phosphatase as a biochemical

marker for intestinal development in the nematode Caenorhabditis elegans. Dev Biol.

1991;147(1):133–43.

16.    Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network

analysis. BMC Bioinformatics. 2008;9:559.

17.    Higgins DP, Weisman CM, Lui DS, D'Agostino FA, Walker AK. Defining characteristics

and conservation of poorly annotated genes in Caenorhabditis elegans using WormCat 2.0.

Genetics. 2022;221(4).

18.    Holdorf AD, Higgins DP, Hart AC, Boag PR, Pazour GJ, Walhout AJM, et al. WormCat:

An Online Tool for Annotation and Visualization of Caenorhabditis elegans Genome-Scale Data.

Genetics. 2020;214(2):279–94.

19.    Ghaddar A, Armingol E, Huynh C, Gevirtzman L, Lewis NE, Waterston R, et al. Whole-

body gene expression atlas of an adult metazoan. Sci Adv. 2023;9(25):eadg0506.

20.    Ge SX, Jung D, Yao R. ShinyGO: a graphical gene-set enrichment tool for animals and

plants. Bioinformatics. 2020;36(8):2628–9.

21.    Mayer S, Raulf MK, Lepenies B. C-type lectins: their network and roles in pathogen

recognition and immunity. Histochem Cell Biol. 2017;147(2):223–37.

22.    Schulenburg H, Hoeppner MP, Weiner J, 3rd, Bornberg-Bauer E. Specificity of the

innate immune system and diversity of C-type lectin domain (CTLD) proteins in the nematode

Caenorhabditis elegans. Immunobiology. 2008;213(3-4):237–50.

23.    Pees B, Yang W, Kloock A, Petersen C, Peters L, Fan L, et al. Effector and regulator:

Diverse functions of C. elegans C-type lectin-like domain proteins. PLoS Pathog.

2021;17(4):e1009454.

24.    Castaneda O, Sotolongo V, Amor AM, Stocklin R, Anderson AJ, Harvey AL, et al.

Characterization of a potassium channel toxin from the Caribbean Sea anemone Stichodactyla

helianthus. Toxicon. 1995;33(5):603–13.

25.    Loukas A, Hintz M, Linder D, Mullin NP, Parkinson J, Tetteh KK, et al. A family of

secreted mucins from the parasitic nematode Toxocara canis bears diverse mucin domains but

shares similar flanking six-cysteine repeat motifs. J Biol Chem. 2000;275(50):39600–7.

26.    Chhabra S, Chang SC, Nguyen HM, Huq R, Tanner MR, Londono LM, et al. Kv1.3

channel-blocking immunomodulatory peptides from parasitic worms: implications for

autoimmune diseases. FASEB J. 2014;28(9):3952–64.

27.    Troemel ER, Chu SW, Reinke V, Lee SS, Ausubel FM, Kim DH. p38 MAPK regulates

expression of immune response genes and contributes to longevity in C. elegans. PLoS Genet.

2006;2(11):e183.

28.    Shivers RP, Youngman MJ, Kim DH. Transcriptional responses to pathogens in

Caenorhabditis elegans. Curr Opin Microbiol. 2008;11(3):251–6.

29.    Radeke LJ, Herman MA. Identification and characterization of differentially expressed

genes in Caenorhabditis elegans in response to pathogenic and nonpathogenic

Stenotrophomonas maltophilia. BMC Microbiol. 2020;20(1):170.

30.    Tan MW, Mahajan-Miklos S, Ausubel FM. Killing of Caenorhabditis elegans by

Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. Proc Natl Acad

Sci U S A. 1999;96(2):715–20.

31.    Tullet JM, Araiz C, Sanders MJ, Au C, Benedetto A, Papatheodorou I, et al. DAF-

16/FoxO directly regulates an atypical AMP-activated protein kinase gamma isoform to mediate

the effects of insulin/IGF-1 signaling on aging in Caenorhabditis elegans. PLoS Genet.

2014;10(2):e1004109.

32.    Leiser SF, Miller H, Rossner R, Fletcher M, Leonard A, Primitivo M, et al. Cell

nonautonomous activation of flavin-containing monooxygenase promotes longevity and health

span. Science. 2015;350(6266):1375–8.

33.    Ziegler DM. An overview of the mechanism, substrate specificities, and structure of

FMOs. Drug Metab Rev. 2002;34(3):503–11.

34.    Xiao W, Wang RS, Handy DE, Loscalzo J. NAD(H) and NADP(H) Redox Couples and

Cellular Energy Metabolism. Antioxid Redox Signal. 2018;28(3):251–72.

35.    Raizen D, Song BM, Trojanowski N, You YJ. Methods for measuring pharyngeal

behaviors. WormBook. 2012:1–13.

36.    Fukushige T, Hawkins MG, McGhee JD. The GATA-factor elt-2 is essential for formation

of the Caenorhabditis elegans intestine. Dev Biol. 1998;198(2):286–302.

37.    Samuel BS, Rowedder H, Braendle C, Felix MA, Ruvkun G. Caenorhabditis elegans

responses to bacteria from its natural habitats. Proc Natl Acad Sci U S A. 2016;113(27):E3941–

9.

38.    Zaslaver A, Liani I, Shtangel O, Ginzburg S, Yee L, Sternberg PW. Hierarchical sparse

coding in the sensory system of Caenorhabditis elegans. Proc Natl Acad Sci U S A.

2015;112(4):1185–9.

39.  Bricker DK, Taylor EB, Schell JC, Orsak T, Boutron A, Chen YC, et al. A mitochondrial

pyruvate carrier required for pyruvate uptake in yeast, Drosophila, and humans. Science.

2012;337(6090):96–100.

40.  Herzig S, Raemy E, Montessuit S, Veuthey JL, Zamboni N, Westermann B, et al.

Identification and functional expression of the mitochondrial pyruvate carrier. Science.

2012;337(6090):93–6.

41.  Gray LR, Tompkins SC, Taylor EB. Regulation of pyruvate metabolism and human

disease. Cell Mol Life Sci. 2014;71(14):2577–604.

42.  Hickey JW, Becker WR, Nevins SA, Horning A, Perez AE, Zhu C, et al. Organization of

the human intestine at single-cell resolution. Nature. 2023;619(7970):572–84.

43.  Mahoney TR, Luo S, Round EK, Brauner M, Gottschalk A, Thomas JH, et al. Intestinal

signaling to GABAergic neurons regulates a rhythmic behavior in Caenorhabditis elegans. Proc

Natl Acad Sci U S A. 2008;105(42):16350–5.

44.  Beg AA, Ernstrom GG, Nix P, Davis MW, Jorgensen EM. Protons act as a transmitter for

muscle contraction in C. elegans. Cell. 2008;132(1):149–60.

45.  Borgonie Gt, Claeys M, Waele DD, Coomans A. In-vivo and in-vitro characterization of

the intestine of fifteen bacteriophagous nematodes (Nematoda: Rhabditida). Fundamental and

applied nematology. 1995;18:115–22.

46.  Borgonie G, Claeys M, Dewaele D, Coomans A. Ultrastructure of the Intestine of the

Bacteriophagous Nematodes Caenorhabditis-Elegans, Panagrolaimus Superbus and

Acrobeloides Maximus (Nematoda, Rhabditida). Fundamental and Applied Nematology.

1995;18(2):123–33.

47.  Panek J, Gang SS, Reddy KC, Luallen RJ, Fulzele A, Bennett EJ, et al. A cullin-RING

ubiquitin ligase promotes thermotolerance as part of the intracellular pathogen response in

Caenorhabditis elegans. Proc Natl Acad Sci U S A. 2020;117(14):7950–60.

48.  Martinez-Reyes I, Chandel NS. Mitochondrial TCA cycle metabolites control physiology

and disease. Nat Commun. 2020;11(1):102.

49.  Picard M, Shirihai OS. Mitochondrial signal transduction. Cell Metab. 2022;34(11):1620–

53.

50.  Nargund AM, Fiorese CJ, Pellegrino MW, Deng P, Haynes CM. Mitochondrial and

nuclear accumulation of the transcription factor ATFS-1 promotes OXPHOS recovery during the

UPR(mt). Mol Cell. 2015;58(1):123–33.

51.      Behr S, Brameyer S, Witting M, Schmitt-Kopplin P, Jung K. Comparative analysis of
LytS/LytTR-type histidine kinase/response regulator systems in gamma-proteobacteria. PLoS
One. 2017;12(8):e0182993.

52.      Fried L, Behr S, Jung K. Identification of a target gene and activating stimulus for the
YpdA/YpdB histidine kinase/response regulator system in Escherichia coli. J Bacteriol.
2013;195(4):807–15.

53.      Paczia N, Nilgen A, Lehmann T, Gatgens J, Wiechert W, Noack S. Extensive
exometabolome analysis reveals extended overflow metabolism in various microorganisms.
Microb Cell Fact. 2012;11:122.

54.      Yasid NA, Rolfe MD, Green J, Williamson MP. Homeostasis of metabolites in
Escherichia coli on transition from anaerobic to aerobic conditions and the transient secretion of
pyruvate. R Soc Open Sci. 2016;3(8):160187.

55.      Zhang Y, Chou JH, Bradley J, Bargmann CI, Zinn K. The Caenorhabditis elegans seven-
transmembrane protein ODR-10 functions as an odorant receptor in mammalian cells. Proc Natl
Acad Sci U S A. 1997;94(22):12162–7.

56.      Harvald EB, Sprenger RR, Dall KB, Ejsing CS, Nielsen R, Mandrup S, et al. Multi-omics
Analyses of Starvation Responses Reveal a Central Role for Lipoprotein Metabolism in Acute
Starvation Survival in C. elegans. Cell Syst. 2017;5(1):38–52 e4.

57.      Maier T, Guell M, Serrano L. Correlation of mRNA and protein in complex biological
samples. FEBS Lett. 2009;583(24):3966–73.

58.      Vogel C, Marcotte EM. Insights into the regulation of protein abundance from proteomic
and transcriptomic analyses. Nat Rev Genet. 2012;13(4):227–32.

59.      Thomas A, Lee PJ, Dalton JE, Nomie KJ, Stoica L, Costa-Mattioli M, et al. A versatile
method for cell-specific profiling of translated mRNAs in Drosophila. PLoS One.
2012;7(7):e40276.

60.      Huang Y, Ainsley JA, Reijmers LG, Jackson FR. Translational profiling of clock cells
reveals circadianly synchronized protein synthesis. PLoS Biol. 2013;11(11):e1001703.

61.      Zhang KX, Tan L, Pellegrini M, Zipursky SL, McEwen JM. Rapid Changes in the
Translatome during the Conversion of Growth Cones to Synaptic Terminals. Cell Rep.
2016;14(5):1258–71.

62.      Gracida X, Calarco JA. Cell type-specific transcriptome profiling in C. elegans using the
Translating Ribosome Affinity Purification technique. Methods. 2017;126:130–7.

63.    Sanz E, Yang L, Su T, Morris DR, McKnight GS, Amieux PS. Cell-type-specific isolation

of ribosome-associated mRNA from complex tissues. Proc Natl Acad Sci U S A.

2009;106(33):13939–44.

64.    Wester LE, Lanjuin A, Bruckisch EHW, Perez-Matos MC, Stine PG, Heintz C, et al. A

single-copy knockin translating ribosome immunoprecipitation toolkit for tissue-specific profiling

of actively translated mRNAs in C. elegans. Cell Rep Methods. 2023;3(3):100433.

65.    Rhoades JL, Nelson JC, Nwabudike I, Yu SK, McLachlan IG, Madan GK, et al. ASICs

Mediate Food Responses in an Enteric Serotonergic Neuron that Controls Foraging Behaviors.

Cell. 2019;176(1-2):85–97 e14.

66.    Gracida X, Dion MF, Harris G, Zhang Y, Calarco JA. An Elongin-Cullin-SOCS Box

Complex Regulates Stress-Induced Serotonergic Neuromodulation. Cell Rep.

2017;21(11):3089–101.

67.    Koterniak B, Pilaka PP, Gracida X, Schneider LM, Pritisanac I, Zhang Y, et al. Global

regulatory features of alternative splicing across tissues and within the nervous system of C.

elegans. Genome Res. 2020;30(12):1766–80.

68.    Nousch M. RPL-4 and RPL-9 Mediated Ribosome Purifications Facilitate the Efficient

Analysis of Gene Expression in Caenorhabditis elegans Germ Cells. G3 (Bethesda).

2020;10(11):4063–9.

69.    Shi Z, Fujii K, Kovary KM, Genuth NR, Rost HL, Teruel MN, et al. Heterogeneous

Ribosomes Preferentially Translate Distinct Subpools of mRNAs Genome-wide. Mol Cell.

2017;67(1):71–83 e7.

70.    Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77(1):71–94.

71.    Noble T, Stieglitz J, Srinivasan S. An integrated serotonin and octopamine neuronal

circuit directs the release of an endocrine signal to control C. elegans body fat. Cell Metab.

2013;18(5):672–84.

72.    Palamiuc L, Noble T, Witham E, Ratanpal H, Vaughan M, Srinivasan S. A tachykinin-like

neuroendocrine signalling axis couples central serotonin action and nutrient sensing with

peripheral lipid metabolism. Nat Commun. 2017;8:14237.

73.    Liu CC, Khan A, Seban N, Littlejohn N, Srinivasan S. A homeostatic gut-to-brain insulin

antagonist restrains neuronally stimulated fat loss. bioRxiv. 2023.

74.    Kamath RS, Ahringer J. Genome-wide RNAi screening in Caenorhabditis elegans.

Methods. 2003;30(4):313–21.

75.    Srinivasan S, Sadegh L, Elle IC, Christensen AG, Faergeman NJ, Ashrafi K. Serotonin
regulates C. elegans fat and feeding through independent molecular mechanisms. Cell Metab.
2008;7(6):533–44.
76.    Filipowicz A, Lalsiamthara J, Aballay A. Dissection of a sensorimotor circuit underlying
pathogen aversion in C. elegans. BMC Biol. 2022;20(1):229.
77.    Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15–21.
78.    Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923–30.
79.    Andrews S. FastQC:  A Quality Control Tool for High Throughput Sequence Data
[Online]. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. 2010.
80.    Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.
81.    Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal
enrichment tool for interpreting omics data. Innovation (Camb). 2021;2(3):100141.

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

**1377**
**1378** **Supplementary Table 1:** *C. elegans* **strains used in this study.**

| Strains used in this study | Source | Identifier | Additional Information |
|---|---|---|---|
| *N2; ssrIs1248[Pges-1△B::rpl-22-3xHA]; ssrIs1191[Pges-1△B::GFP]* | This paper | SSR1578 | Integrated. Backcrossed 4X |
| *N2; ssrIs1191[Pges-1△B::GFP]; ssrIs1210[Ppho-1::mCherry]* | This paper | SSR1582 | Integrated. Backcrossed 4X. |
| *N2; ssrIs1251[Ppho-1::rpl-22-3xHA]; ssrIs1210[Ppho-1::mCherry]* | This paper | SSR1605 | Integrated. Backcrossed 4X. |
| *sid-1(qt9); ssrIs1220[Pges-1△B::sid-1::GFP]; ssrIs1240[Pges-1△B::ins-7mCherry]; ssrIs615[Punc-122::GFP]* | PMID: 37961386 | SSR1634 | |
| *N2; ssrEx1317[Pclec-160::mNeonGreen]; ssrEx1031[Pmyo-2::mCherry]* | This paper | SSR1701 | |
| *N2; ssrEx1319[Pclec-86::mNeonGreen]; ssrEx1031[Pmyo-2::mCherry]* | This paper | SSR1702 | |
| *N2; ssrEx1318[PB0024.4::mNeonGreen]; ssrEx1031[Pmyo-2::mCherry]* | This paper | SSR1703 | |
| *N2; ssrEx1323[PF01D5.5::mNeonGreen]; ssrEx1031[Pmyo-2::mCherry]* | This paper | SSR1705 | |
| *N2; ssrEx1324[Pnpr-28::mNeonGreen]; ssrEx1031[Pmyo-2::mCherry]* | This paper | SSR1706 | |
| *N2; ssrEx1325[PF01D5.1::mNeonGreen]; ssrEx1031[Pmyo-2::mCherry]* | This paper | SSR1713 | |

**1379**
**1380** **Supplementary Table 2: Plasmids used in this study.**
**1381**

| Plasmids used in this study | Source | Identifier |
|---|---|---|
| *Pges-1△B::GFP* | This paper | pSS1191 |
| *Ppho-1::mCherry* | This paper | pSS1210 |
| *Pges-1△B::rpl-22-3xHA* | This paper | pSS1248 |
| *Ppho-1::rpl-22-3xHA* | This paper | pSS1251 |
| *Pclec-160::mNG::clec-160 3'UTR* | This paper | pSS1317 |
| *PB0024.4::mNG::B0024.4 3'UTR* | This paper | pSS1318 |
| *Pclec-86::mNG::clec-86 3'UTR* | This paper | pSS1319 |
| *PF01D5.5::mNG::F01D5.5 3'UTR* | This paper | pSS1323 |
| *Pnpr-28::mNG::npr-28 3'UTR* | This paper | pSS1324 |
| *PF01D5.1::mNG::F01D5.1 3'UTR* | This paper | pSS1325 |

**1382**
**1383** **Supplementary Table 3: Cloning primers used in this study.**
**1384**

| Gibson Assembly | Primer | Source | Sequence |
|---|---|---|---|
| *Pclec-160::mNG::clec-160 3'UTR* | Forward primer for cloning the *Pclec-160* sequence | IDT | CATGCCTGCAAGGAAAAAATTGGAAGGGAAAGGAAAACT |
| | Reverse primer for cloning the *Pclec-160* sequence | IDT | TTGGAGACCATCTGAATAATTCTTGGGTATTAAAAAAAAGATTTTTGTACCTATTGG |
| | Forward primer for cloning the mNG sequence | IDT | AATTATTCAGATGGTCTCCAAGGGAGAGGAG |
| | Reverse primer for cloning the mNG sequence | IDT | GAATGACATCACTACTTGTAGAGCTCGTCCATTCC |
| | Forward primer for cloning the *clec-160 3'UTR* sequence | IDT | CTACAAGTAGTGATGTCATTCTCAATTTGAATTTAAACAGAATAAACTATTTTTAAAAGT |
| | Reverse primer for cloning the *clec-160 3'UTR* sequence | IDT | AATTCGAGCTCTCGCCTCAATTCTGTGATTCATAAAAAGTTCA |
| | Forward primer for cloning the pUC19 sequence | IDT | ATTGAGGCGAGAGCTCGAATTCACTGGCCG |
| | Reverse primer for cloning the pUC19 sequence | IDT | AATTTTTTCCTTGCAGGCATGCAAGCTTG |
| *Pclec-86::mNG::clec-86 3'UTR* | Forward primer for cloning the *Pclec-86* sequence | IDT | AGCTTGCATGCCTACCCGACCGGCCG |
| | Reverse primer for cloning the *Pclec-86* sequence | IDT | CCTTGGAGACCATTTTCAGTGTCCAGTAAAATAATAAATTTTATCTGCTTATCACAG |
| | Forward primer for cloning the mNG sequence | IDT | CTGGACACTGAAAATGGTCTCCAAGGGAGAGGAG |
| | Reverse primer for cloning the mNG sequence | IDT | TTCGAATCAAAAGCTACTTGTAGAGCTCGTCCATTCCC |
| | Forward primer for cloning the *clec-86 3'UTR* sequence | IDT | GCTCTACAAGTAGCTTTTGATTCGAAAGAAAATATTTATATATTATGAGCTAAATTGGAT |
| | Reverse primer for cloning the *clec-86 3'UTR* sequence | IDT | GTGAATTCGAGCTACCTGAGCATGCCGTTTGG |

| | | | |
|---|---|---|---|
| | Forward primer for cloning the pUC19 sequence | IDT | GGCATGCTCAGGTAGCTCGAATTC ACTGGCCGTCGTTTTACA |
| | Reverse primer for cloning the pUC19 sequence | IDT | CGGCCGGTCGGGTAGGCATGCAA GCTTGGCGT |
| *PB0024.4::mNG::B002 4.4 3'UTR* | Forward primer for cloning the *PB0024.4* sequence | IDT | CTTGCATGCCTGTTTAATTTTAAATG AGAAAATAATGATTTTTTGTCATTAA ATTTAACT |
| | Reverse primer for cloning the *PB0024.4* sequence | IDT | CTTGGAGACCATTCTACTAGAAATA GAAAGTTCGTTTGGAATACCG |
| | Forward primer for cloning the mNG sequence | IDT | TTTCTAGTAGAATGGTCTCCAAGGG AGAGGAG |
| | Reverse primer for cloning the mNG sequence | IDT | ACAAATATCAACCTACTTGTAGAGC TCGTCCATTCCC |
| | Forward primer for cloning the *B0024.4 3'UTR* sequence | IDT | TCTACAAGTAGGTTGATATTTGTTTG TAATTACGGACTTATGGTGC |
| | Reverse primer for cloning the *B0024.4 3'UTR* sequence | IDT | AGTGAATTCGAGATTATTTGTGGCC ATCCCCAACTATAAAAAT |
| | Forward primer for cloning the pUC19 sequence | IDT | CCACAAATAATCTCGAATTCACTGG CCGTCG |
| | Reverse primer for cloning the pUC19 sequence | IDT | TTAAAATTAAACAGGCATGCAAGCT TGGCGT |
| *PF01D5.5::mNG::F01D 5.5 3'UTR* | Forward primer for cloning the *PF01D5.5* sequence | IDT | AGCTTGCATGCCCTGGAAATTGTTC GGGGATTTTTAATTTTTTTTTTAAGC |
| | Reverse primer for cloning the *PF01D5.5* sequence | IDT | CCTTGGAGACCATCGCGAGCTTGA AGTTTTTCTTTGTT |
| | Forward primer for cloning the mNG sequence | IDT | TTCAAGCTCGCGATGGTCTCCAAG GGAGAGGAG |
| | Reverse primer for cloning the mNG sequence | IDT | ATTTTTGTAATTACTACTTGTAGAGC TCGTCCATTCC |
| | Forward primer for cloning the *F01D5.5 3'UTR* sequence | IDT | CTCTACAAGTAGTAATTACAAAAATA TCTTTGGAATTGTAAATATTGTTATTT GAATAAA |
| | Reverse primer for cloning the *F01D5.5 3'UTR* sequence | IDT | ATTCGAGCTCGGTTGCCGTTCCCC TATCCTACCGTAC |
| | Forward primer for cloning the pUC19 sequence | IDT | AGGGGAACGGCAACCGAGCTCGAA TTCACTGGCC |
| | Reverse primer for cloning the pUC19 sequence | IDT | GAACAATTTCCAGGGCATGCAAGC TTGGCGT |
| *PF01D5.1::mNG::F01D 5.1 3'UTR* | Forward primer for cloning the *PF01D5.1* sequence | IDT | CAAGCTTGCATGCCTGGATAAAGAA AGCATGAATTTGAATTTTATATGTAC ACTATTCT |
| | Reverse primer for cloning the *PF01D5.1* sequence | IDT | TCCCTTGGAGACCATTTGTTTTCTG CAAAACTTAGTTTTTAGATAAAATGA CTTTAAGAC |
| | Forward primer for cloning the mNG sequence | IDT | TTTGCAGAAAACAAATGGTCTCCAA GGGAGAGGAG |
| | Reverse primer for cloning the mNG sequence | IDT | AATAAAAAGATTTATCTACTTGTAGA GCTCGTCCATTCCC |
| | Forward primer for cloning the *F01D5.1 3'UTR* sequence | IDT | AGCTCTACAAGTAGATAAATCTTTTT ATTTTAATTTTCATCGTAACTCGGTA GTAGAACA |
| | Reverse primer for cloning the *F01D5.1 3'UTR* sequence | IDT | CAGTGAATTCGAGCTCAGACACACT ACACCTAACTCGTTCAG |
| | Forward primer for cloning the pUC19 sequence | IDT | GTGTAGTGTGTCTGAGCTCGAATTC ACTGGCCG |
| | Reverse primer for cloning the pUC19 sequence | IDT | TGCTTTCTTTATCCAGGCATGCAAG CTTGGCGT |
| *Pnpr-28::mNG::npr-28 3'UTR* | Forward primer for cloning the *Pnpr-28* sequence | IDT | TTGCATGCCTGAGCTAATGTCCTAT TCGCCCC |
| | Reverse primer for cloning the *Pnpr-28* sequence | IDT | TTGGAGACCATAGTAGAATAACTGA CAAGTCGTGTAAAGCAAA |
| | Forward primer for cloning the mNG sequence | IDT | GTTATTCTACTATGGTCTCCAAGGG AGAGGAG |
| | Reverse primer for cloning the mNG sequence | IDT | AGTATTGTTATCTACTTGTAGAGCTC GTCCATTCCC |
| | Forward primer for cloning the *npr-28 3'UTR* sequence | IDT | TCTACAAGTAGATAACAATACTTTGA ATATGTACCTTTTTTCAACTTGTATC ATT |

| | Reverse primer for cloning the *npr-28 3'UTR* sequence | IDT | TGAATTCGAGCTATTTCGTAATATTT ATTTGTTTATCTTTTTATTTACTTATTT CACTCA |
|---|---|---|---|
| | Forward primer for cloning the pUC19 sequence | IDT | ATTACGAAATAGCTCGAATTCACTG GCCG |
| | Reverse primer for cloning the pUC19 sequence | IDT | GGACATTAGCTCAGGCATGCAAGC TTGGCGT |

1411