Characterisation of cold-selective lamina I spinal projection neurons

Aimi N. Razlan[1,4], Wenhui Ma[1,4], Allen C. Dickie[1,4], Erika Polgár[1,4], Anna McFarlane[1],
Mansi Yadav[1], Andrew H. Cooper[1], Douglas Strathdee[2], Masahiko Watanabe[3], Andrew M.
Bell[1], Andrew J. Todd[1]*, Junichi Hachisuka[1]*

[1]School of Psychology and Neuroscience, College of Medical, Veterinary and Life
Sciences, University of Glasgow, Glasgow G12 8QQ, UK

[2]School of Cancer Sciences, College of Medical, Veterinary and Life Sciences, University
of Glasgow, Bearsden G61 1BD UK

[3]Department of Anatomy, Hokkaido University School of Medicine, Sapporo 060-8638,
Japan

[4]Equal contribution

*Corresponding Authors: Andrew.Todd@glasgow.ac.uk and
Junichi.Hachisuka@Glasgow.ac.uk

Click here to enter text.

ABSTRACT

Skin cooling is detected by primary afferents that express the Trpm8 channel, but how this information is conveyed to the brain remains poorly understood. We have previously identified a population of lamina I projection neurons belonging to the anterolateral system (ALS) that receive numerous contacts from Trpm8-expressing primary afferents. Here, using a semi-intact somatosensory preparation, we provide evidence that these cells correspond to the cold-selective ALS neurons identified in previous physiological studies. We also confirm the presence of synapses from Trpm8 afferents onto these cells at the ultrastructural level and with optogenetics. Based on our previous transcriptomic findings, we identify calbindin as a molecular marker, and show that this can be used to target the cold-selective ALS neurons for anterograde tracing studies. We provide evidence that they project to brain regions that have been implicated in thermosensation: the rostralmost part of the lateral parabrachial area, the caudal part of the periaqueductal grey matter, and the posterior triangular and ventral posterolateral nuclei of the thalamus. Our findings provide important insights into the organisation of neuronal circuits that underlie thermoregulation and the perception of cold stimuli applied to the skin.

Click here to enter text.

INTRODUCTION

The anterolateral system (ALS) consists of spinal cord neurons that project to various brain regions, including the caudal ventrolateral medulla (CVLM), the nucleus of the solitary tract (NTS), the lateral parabrachial area (LPB), the periaqueductal grey matter (PAG), the superior colliculus and the thalamus. ALS neurons are required for the perception of pain, itch and skin temperature, as shown by the loss of these sensations on the side contralateral to an anterolateral cordotomy[1,2]. Although axons belonging to this pathway ascend in the anterolateral quadrant in humans, they are located in the dorsolateral white matter in rodents[3-6]. ALS neurons account for only a small proportion (probably less than 1%) of spinal cord neurons[7,8] and are unevenly distributed, with a relatively high density in lamina I and the lateral spinal nucleus (LSN), and scattered cells in deeper laminae (III-VII) and around the central canal[9-12]. ALS neurons are functionally heterogeneous, and we recently identified 5 distinct transcriptomic populations, which we named ALS1-5[13]. This was based on single-nucleus RNA sequencing of a major subset of these cells, those that transiently express the transcription factor Phox2a during development[14].

Perception of skin cooling is dependent on primary afferent neurons that express the Trpm8 channel[15-19], and these terminate mainly in lamina I of the spinal cord[20-22]. Consistent with this, it has been shown that many lamina I projection neurons respond to cool or cold stimuli applied to the skin, with a distinctive subset of "cold-selective" neurons responding predominantly or exclusively to skin cooling[23-33].

How information is transmitted from Trpm8-expressing primary afferent neurons to these cold-selective lamina I ALS cells is currently not known, although a recent report has suggested that this occurs via a dedicated population of excitatory interneurons[34]. However, we had previously identified a population of lamina I ALS neurons that received dense synaptic input from Trpm8-expressing primary afferents[13]. These neurons had dendrites that were intimately associated with bundles of Trpm8-positive afferents, while their cell bodies generally also received numerous contacts, and were sometimes surrounded by these afferents. We therefore proposed that these cells correspond to the cold-selective neurons described above, and also provided evidence that they belonged to the ALS3 population[13].

The aims of this study were to characterise and validate the Trpm8[Flp] mouse line that we had used to reveal this input, to confirm the presence of synaptic input, and to test the prediction that the lamina I projection neurons with dense synaptic input from Trpm8-

Click here to enter text.

expressing primary afferents do indeed correspond to the cold-selective population

described in physiological studies[23-33]. In addition, since Calb1 (which encodes the

calcium-binding protein calbindin) is one of the top differentially expressed genes in the

ALS3 population, we used mice in which Cre recombinase is targeted to the Calb1 locus

(Calb1[Cre]) to investigate the projections of the ALS3 cells to the brain.

RESULTS

Validation of Trpm8[Flp] mouse line and characterisation of Flp-expressing cells.

Trpm8[Flp] mice were crossed with the Flp reporter line RCE:FRT to generate

Trpm8[Flp];RCE:FRT mice, in which cells that express Flp at any stage during development

are labelled with GFP. Tissue was obtained from mice aged between 5-9 weeks for this

part of the study. We used a stereological technique to quantify neurons in an L4 dorsal

root ganglion (DRG) from 3 of these mice and found that 5.7% (range 5.4-6.7%) of all

DRG neurons contained GFP (Figure 1A). We then performed fluorescence *in situ*

hybridisation (FISH) to examine the extent of overlap between GFP and *Trpm8* mRNA

(Figure 1B). In DRGs from 3 mice, we found that 92% (88%-98%) of GFP-containing cells

had *Trpm8* message, while 83% (80%-86%) of cells with *Trpm8* were GFP-positive. These

findings show a high degree of fidelity in the Trpm8[Flp] line, and are consistent with the

reported expression of *Trpm8* mRNA in 5-10% of mouse DRG cells[15]. We also used this

tissue to determine the soma size of the GFP-positive cells (Figure 1C), and found that

these were small, with a mean diameter of 14.6 µm (± 3.4 µm, SD), showing a very similar

size range to that reported for cells captured with a different Trpm8[Flp] mouse line[35]. Trpm8-

positive cells that lacked GFP, as well as the few GFP-labelled cells that lacked *Trpm8*

mRNA, showed a similar size distribution.

Transcriptomic studies have demonstrated that Trpm8-expressing primary afferents can

co-express certain genes that serve as markers for primary afferent populations, such as

Slc17a8 (which encodes Vglut3), Trpv1 and Tac1 (the gene coding for substance P), but

lack other neuropeptides such as calcitonin gene-related peptide (CGRP) and

somatostatin[36,37]. We therefore performed immunohistochemistry to look for expression of

the corresponding proteins/peptides in dorsal root ganglia from 3 mice in each case. We

found that 11.9% (9.8-13.8%) of GFP-containing cells were immunoreactive for Vglut3 and

26.4% (22.4-30.9%) for Trpv1 (Figure 1–figure supplement 1A-D). Among the

neuropeptides, we found minimal co-localisation with CGRP (mean 2.5%, range 1.4-4.5%)

and none with somatostatin. However, 56% (range 55.8-56.3%) of the GFP cells contained

Click here to enter text.

substance P, although this was expressed at relatively low levels (Figure 1–figure

supplement 1G-M). These results are therefore consistent with the transcriptomic findings

described above[36,37].

Central projections of Trpm8-expressing primary afferents have been shown to

terminate mainly in lamina I[20-22], and we found a similar pattern in the spinal cords of

Trpm8[Flp];RCE:FRT mice, with the great majority of GFP-labelled axons remaining in

lamina I, and very few passing ventrally (Figure 2). A similar pattern of labelling was

observed in all spinal segments examined. We had previously noted that when lamina I

was viewed in horizontal sections, Trpm8 afferents were arranged in interweaving bundles

that frequently followed the dendrites of a distinctive subset of projection neurons[13].

Consistent with this, we found that Trpm8 afferents labelled with GFP showed a

discontinuous arrangement within lamina I when viewed in transverse sections and did not

occupy the whole medio-lateral extent of the lamina (Figure 2). The GFP-labelled afferents

did not show any obvious restriction in the medio-lateral axis, and could be seen in medial,

central and lateral parts of lamina I. Since the great majority of Trpm8-expressing afferents

contain GFP in this genetic cross, it is unlikely that the large spaces between bundles of

GFP-labelled afferents in lamina I are filled by Trpm8-expressing afferents that lack GFP.

This "incomplete" innervation of lamina I by Trpm8 afferents is therefore likely to be

genuine.

It has been reported that some lamina I projection neurons were associated with

boutons that were immunoreactive for Vglut3[38] and since 12% of Trpm8 afferent cell

bodies contained Vglut3, we looked for expression of the transporter in their central

terminals. In transverse sections, Vglut3-immunoreactive axons formed a dense band in

inner lamina II that corresponds to central terminals of C-low threshold mechanoreceptors

(C-LTMRs)[39,40]. However, there were a few Vglut3-immunoreactive boutons in lamina I

that were also GFP-positive, indicating that central terminals of some Trpm8-expressing

primary afferents had detectable levels of the transporter (Figure 2–figure supplement 1A-

F). We generated a mouse genetic cross (Phox2a::Cre;Ai9;Trpm8[Flp];RCE:FRT) in which a

subset of ALS cells express tdTomato and Trpm8 afferents express GFP. In horizontal

sections from these mice we identified lamina I projection neurons that were associated

with numerous GFP-labelled (Trpm8-expressing) boutons and found that some of these

boutons contained Vglut3 (Figure 2–figure supplement 1G-J). The proportion of GFP-

labelled boutons that contained Vglut3 varied considerably between cells. Since C-LTMRs

do not appear to arborise in lamina I[35], it is likely that the Vglut3-containing boutons that

have been found to contact projection neurons in this lamina[38] belong to Trpm8 afferents.

Even though many Trpm8-positive cell bodies showed Trpv1 and substance P

immunoreactivities, we were not able to detect either of these in their central terminals

(Figure 2–figure supplement 1K-M).

Although we did not investigate the distribution of GFP in the brains of

Trpm8[Flp];RCE:FRT mice in detail, we noted that there was dense axonal labelling in the

spinal trigeminal nucleus, while scattered GFP-labelled cell bodies were present in various

brain regions that are known to contain Trpm8-expressing neurons, including the preoptic

area and the reticular nucleus of the thalamus[41] (data not shown).

Synaptic input from Trpm8-expressing afferents to lamina I projection neurons.

We had previously shown that ~20% of lamina I projection neurons retrogradely

labelled from the LPB received numerous contacts from GFP-labelled afferents in

Trpm8[Flp];RCE:FRT mice, and that these contacts were associated with punctate staining

for Homer1[13] (a marker for glutamatergic synapses[42]). To reveal synapses formed

between Trpm8 afferents and these projection neurons at the ultrastructural level, we used

a combined confocal/electron microscopic method[43] on spinal cord tissue from

Phox2a::Cre;Ai9;Trpm8[Flp];RCE:FRT mice. This genetic cross was chosen because we had

previously shown that the Phox2a-lineage includes many lamina I ALS neurons with dense

input from Trpm8 afferents[13]. Horizontal sections from lumbar spinal cord segments of 2 of

these mice were immunostained to reveal tdTomato and GFP, and 3 tdTomato-labelled

cells with numerous contacts from GFP-labelled axons were scanned with a confocal

microscope (Figure 3, Figure 3–figure supplements 1 and 2). The tissue was subsequently

processed with an immunoperoxidase method to label the GFP-positive axons with

diaminobenzidine (DAB), and examined with an electron microscope (Figure 3, Figure 3–

figure supplement 2). Although the cell bodies and dendrites of the projection neurons did

not contain an electron-dense label, they could be readily identified by their location within

the ultrathin sections, by their association with numerous DAB-labelled axonal boutons

and by the morphology of their somata and proximal dendrites (Figure 3A,B). Because of

the very high density of GFP-labelled boutons that were seen to contact the cells in the

confocal images, it was not always possible to relate individual boutons from confocal

images to those seen with the electron microscope. However, we were able to identify

numerous axosomatic and axodendritic synapses (a minimum of 10 synapses per cell)

between the GFP-immunoreactive boutons and the lamina I projection neurons. These

Click here to enter text.

could be recognised by the presence of darkening of the membranes and in some cases a

faint postsynaptic density, together with clustering of synaptic vesicles on the presynaptic

side (Figure 3B-D, Figure 3–figure supplement 2C-J). Interestingly, the synapses between

the GFP-containing boutons and the projection neurons generally had relatively faint

postsynaptic densities and only small regions where the vesicles were clustered on the

presynaptic side. These findings, together with our previous observation of Homer1-

immunoreactive puncta located between GFP-expressing afferent boutons and the cell

bodies and dendrites of retrogradely labelled lamina I projection neurons in

Trpm8[Flp];RCE:FRT mice[13], indicate that ALS cells belonging to this population receive

numerous synapses from Trpm8-expressing primary afferents.

To provide electrophysiological evidence for synapses between Trpm8 afferents and

this population of lamina I ALS cells, we carried out optogenetic experiments. Two

Trpm8[Flp];RCE:FRT mice received intraperitoneal injections of

AAV.PHP.S.Flp[ON].ChR2_YFP as neonates to label Trpm8-expressing primary afferents

with channelrhodopsin. They were then injected with AAV11.tdTomato into the CVLM or

LPB ~5 weeks later (Figure 3E). Whole-cell patch-clamp recordings were subsequently

obtained from tdTomato-labelled lamina I neurons in a whole-cord preparation. We used

the Trpm8[Flp];RCE:FRT cross for these experiments in order to allow un-biased

identification of ALS neurons that were surrounded by Trpm8 afferents. Importantly,

although the viral delivery approach that we used will have specifically targeted Trpm8

afferents, it will not have resulted in adequate expression of channelrhodopsin in all of

these cells.

We identified 8 tdTomato cells that were densely surrounded by GFP-labelled (Trpm8)

afferents and found that 1 ms blue light pulses induced EPSCs in 5 of these cells (Figure

3E,F). We measured the latency of the first EPSC after blue light pulses (0.2 Hz) and

found that the latency jitter was less than 1 ms, indicating that these neurons receive

monosynaptic inputs[44] (Figure 3F; n = 5, Average latency: 5.8 ± 0.86 ms, latency jitter:

0.68 ± 0.14 ms). Note that the relatively long latency is likely to reflect the time taken for

action potential initiation in the Trpm8 boutons[31]. Two of the 5 neurons showed multiple

peaks with a larger latency jitter, indicating that there may also have been polysynaptic

inputs (Figure 3–figure supplement 3). It is likely that the remaining 3 cells received input

from Trpm8 afferents that did not express sufficiently high levels of channelrhodopsin.

Together, these findings demonstrate that Trpm8 afferents directly synapse onto lamina I

ALS cells that are surrounded by these afferents.

Click here to enter text.

Lamina I ALS neurons that are surrounded by Trpm8 afferents are cold-selective

We next tested whether the lamina I projection neurons that are densely innervated by Trpm8 afferents correspond to the cold-selective neurons identified in physiological studies[23-33]. To do this, we carried out whole-cell patch-clamp recordings in preparations in which ALS neurons were retrogradely labelled with either mCherry or tdTomato, and Trpm8 axons expressed GFP. Retrograde labelling was achieved by injecting AAV9.mCherry into the CVLM or LPB of Trpm8[Flp];RCE:FRT mice, resulting in mCherry labelling, or else by injecting AAV9.Cre_GFP into the CVLM of Trpm8[Flp];RCE:FRT;Ai9 mice leading to tdTomato labelling (Figure 4A). We used the semi-intact *ex vivo* somatosensory preparation, in which the spinal cord, L2 and L3 roots and ganglia, saphenous and femoral cutaneous nerves, and hindlimb skin are dissected in continuity (Fig 4B)[31,32].

Visualisation of the spinal cord with filter sets for GFP or mCherry/tdTomato revealed lamina I projection neurons (labelled with mCherry or tdTomato) as well as a plexus of GFP-labelled (Trpm8-expressing) primary afferents (Figure 4). We recorded from 6 mCherry/tdTomato-labeled projection neurons that had dendrites (and in some cases also cell bodies) that were clearly associated with numerous GFP-labelled afferents, as well as from 5 projection neurons for which there was no clear association with these afferents. We assessed the responses of these cells to a 4 g von Frey filament, hot saline (50°C), and cold saline (15°C) applied to the skin[31]. All 6 of the projection neurons that had numerous contacts from GFP-labelled axons responded exclusively to application of cold saline. In 4 cases recordings were made in current clamp and the cells fired action potentials (n = 3) or generated EPSPs (subthreshold, n = 1) following application of cold saline, but showed no response to mechanical or heat stimuli (Figure 4C-E). The time course of the response was consistent with that seen for cold-selective projection neurons that we previously reported (Figure 4–figure supplement 1)[32]. The other two cells with numerous GFP contacts were recorded in voltage clamp, and these showed EPSCs in response to cold, but not mechanical or heat, stimuli. All six of these neurons were therefore classified as cold-selective. The five projection neurons that were not preferentially associated with GFP-labelled afferents were all recorded in current clamp mode, and all of these cells exhibited action potential firing in response to mechanical stimulation. Three of them also responded to heat, and one additionally to cold stimuli. The

five neurons that were not associated with numerous GFP-labelled afferents were

therefore classified as either mechano-selective or polymodal (Figure 4F-H).

We previously reported that cold-selective projection neurons are distinct from other

lamina I projection neurons[32], and a defining feature of these cells was their low frequency

of spontaneous EPSCs (sEPSCs). We therefore quantified sEPSC frequency and

amplitude in lamina I projection neurons with or without dense Trpm8 input. As expected,

neurons with dense Trpm8 input had significantly lower sEPSC frequencies compared to

those that lacked dense Trpm8 input (Figure 4–figure supplement 1). Interestingly, the

amplitude of sEPSCs in the projection neurons with dense Trpm8 input was smaller than

in those that lacked this input, which had not been observed in our previous study[32].

Together, these results strongly suggest that lamina I projection neurons receiving dense

Trpm8 afferent innervation constitute a distinct neuronal population specialised for cold

sensation, and correspond to the cold-selective ALS neurons identified in previous

studies[23-33].

Our anatomical studies indicate that the cells densely coated with Trpm8-expressing

primary afferents receive many synapses from these afferents (Figure 3), and we had

shown previously that these cells had numerous Homer1-immunoreactive puncta, with

~60% being associated with a Trpm8-expressing primary afferent bouton[13]. The low

frequency of sEPSCs is therefore not likely to result from a low density of synapses, but

presumably reflects a very low release probability at these synapses in these experimental

conditions. The lower amplitude of sEPSCs may be related to our EM observation that

contacts between Trpm8 afferents and densely coated lamina I projection neurons

typically have small synaptic specialisations (Figure 3, Figure 3–figure supplement 2).

Targeting of cold-selective lamina I ALS neurons with the Calb1^{Cre} mouse

We had previously demonstrated that lamina I projection neurons that are densely

innervated by Trpm8-expressing afferents are included among those of the Phox2a-

lineage, and contain mRNA for *Hs3st1*, one of the top differentially expressed genes for

ALS3. We had therefore assigned the Trpm8-innervated cells to the ALS3 cluster[13]. Calb1,

which encodes the calcium-binding protein calbindin, is also among the top 20 differentially

expressed genes for ALS3 (Figure 5–figure supplement 1A, see also Figure 1D of

reference[13]). However, calbindin is not restricted to the ALS3 cluster, as it is also

expressed by some cells in the ALS2, ALS4 and ALS5 clusters[13] (Figure 5–figure

supplement 1A), and by many ALS neurons in the lateral spinal nucleus (LSN)[45], most of

which are not derived from the Phox2a-lineage[14,46].

To assess the relationship between Calb1-expressing ALS neurons and those with

dense Trpm8 innervation, we injected an AAV coding for Cre-dependent tdTomato

(AAV11.Cre^{ON}.tdTomato[47]) unilaterally into the LPB of 4 Calb1^{Cre};Trpm8^{Flp};RCE:FRT mice

(Figure 5–figure supplement 1B). As expected from the known distribution of calbindin-

expressing ALS cells in the rat[45], we observed tdTomato-positive cells in lamina I (mainly

on the contralateral side), as well as bilaterally in the LSN and in deeper laminae (Figure

5A, Figure 5–figure supplement 1C-F). When we examined horizontal sections through

lamina I, we found that many of the tdTomato-labelled cells, were densely innervated by

Trpm8 afferents (Figure 5C-H), and these were located throughout the mediolateral extent

of the dorsal horn. The proportion of tdTomato cells with dense Trpm8 input varied

between 38.7-59.6% across the L2-L4 segments (Figure 5B, RM ANOVA, p = 0.048).

Interestingly, the proportion was significantly higher in L2 (59.6%) than in L4 (38.7%), (RM

ANOVA, Holm-Šídák's multiple comparisons test, p = 0.0056, n = 4), and this may reflect a

difference in the proportion of lamina I cells with dense Trpm8 innervation between

segments. When results across the 3 segments were pooled, the total numbers of

retrogradely labelled cells ranged from 133 to 170 (mean 151), while the proportions with

dense Trpm8 input ranged from 46.5%-54.1% (mean 50.6%). To confirm expression of

calbindin in lamina I projection neurons with dense Trpm8 input, we carried out

immunofluorescence labelling on horizontal sections from 3 Trpm8^{Flp};RCE:FRT mice that

had received injections of AAV9.mCherry into the LPB. We identified a total of 32

mCherry-positive lamina I neurons that were associated with dense GFP labelling (6-14

per mouse), and found that all but four of these (27/31, 87%) were calbindin-

immunoreactive (Figure 5–figure supplement 1G).

We also used the semi-intact preparation to record from tdTomato-labelled lamina I

neurons in spinal cords from 4 Calb1^{Cre} mice that had received injections of

AAV11.Cre^{ON}.tdTomato into the CVLM (Figure 5–figure supplement 1B, Figure 5–figure

supplement 2). We recorded from 6 retrogradely labelled neurons and found that 5 were

cold-selective, while one was polymodal, responding to brushing of the skin, as well as to

application of both hot and cold stimuli (Figure 5I,J). These cold-selective neurons

showed a similar time course in response to cold stimulation to those described above

(Figure 5–figure supplement 2C). Taken together, these findings indicate that while Calb1

Click here to enter text.

is not restricted to cold-selective cells, many of the Calb1-expressing projection neurons in

lamina I belong to the cold-selective population.

Brain projections of putative cold-selective lamina I neurons

Because Calb1 is expressed by cold-selective ALS cells in lamina I, we carried out

anterograde tracing by injecting AAV coding for Cre-dependent expression of tdTomato

(AAV1.Cre[ON].tdTom) into the lumbar spinal cords of Calb1[Cre] mice. Our aim was to test for

projections to two potential brainstem targets that receive input from the ALS[11,12] and also

contain many neurons that express the immediate early gene Fos in mice exposed to low

ambient temperatures[48-51]: the caudal part of the PAG (cPAG), and the rostral part of the

LPB, in particular a region that has been named PBrel (rostral to external lateral)[49]. Two

cortical areas are activated by skin cooling, the primary somatosensory and posterior

insular cortices[52]. We therefore looked for evidence of an input to the ventral posterolateral

(VPL) and posterior triangular (PoT) thalamic nuclei, which are thought to provide the

thermosensory inputs to these cortical areas in rodents[53]. Initially, we examined labelling

resulting from injections of AAV1.Cre[ON].tdTomato that were targeted on the central part of

the dorsal horn, either in the L3 segment, or in the L3, L4 and L5 segments, using a similar

approach to that which we had taken with other genotypes (Figure 6–figure supplement

1A,B; Table 1). Labelling patterns were consistent among the mice used in this part of the

study, but labelling was denser in all regions in those that had received injections into L3,

L4 and L5. This approach revealed that axons ascended in the dorsolateral white matter of

the spinal cord and gave branches to several regions on the contralateral side of the

brainstem, including LPB and the cPAG (Figure 6–figure supplement 1C,D). Within the

LPB, axonal labelling was very dense in the rostral-most part, which corresponds to

PBrel[49], and moderately dense in the centrolateral (PBcl) nucleus. In addition, there was

some labelling in other nuclei, including external lateral (PBel), dorsolateral (PBdl) and

internal lateral (PBil). There was dense labelling in the cPAG and in addition, labelling was

seen within the superior colliculus (Figure 6–figure supplement 1C). In the thalamus, there

was sparse labelling in the VPL, posterior (Po) and PoT nuclei contralateral to the injection

site, and many labelled profiles were present in the medial thalamus on both sides (Figure

6–figure supplement 1E). These findings were therefore consistent with the expected

projections of cold-selective ALS cells to PBrel, cPAG and the VPL nucleus of thalamus.

However, this strategy will also have labelled Calb1-expressing projection neurons in

other regions, including the LSN and the lateral reticulated part of lamina V (Figure 6–

Click here to enter text.

figure supplement 1B). Consistent with this interpretation, we observed labelling in PBil

and the medial thalamus, both of which are known to receive input from ALS cells in the

deep dorsal horn[54-56], including cells belonging to the ALS4 population in lateral lamina V[5].

We therefore altered the injection technique by reducing the volume and injecting at a

more medial location (250 μm rather than 400 μm lateral to the midline). As with the

previous experiments, these injections were either made into the L3 segment, or into each

of L3, L4 and L5 (Table 1). Again, the pattern of labelling was consistent among these

mice, but labelling was denser in those that had received injections into L3, L4 and L5.

This strategy resulted in injection sites that were restricted to the medial part of the dorsal

horn and excluded the LSN (Fig 6A-C). Although very weak tdTomato labelling was seen

in the LSN in these experiments, this is likely to have arisen from transport by axons of

local calbindin-expressing excitatory interneurons, since many excitatory interneurons in

laminae I-II have axons that enter the LSN[57-59]. Importantly, no tdTomato-labelled cell

bodies were seen in the LSN or in the lateral reticulated part of lamina V in these cases.

Since these injections were likely to have included a much higher proportion of lamina I

neurons with dense Trpm8 innervation, we analysed the anterograde labelling in more

detail.

Axons ascended in the dorsal quadrant of the spinal cord, mainly on the contralateral

side (Figure 6–figure supplement 2A). Although a few collaterals arose at all segmental

levels examined, these were far less numerous than those originating from the

somatostatin-expressing ALS cells that occupy the lateral part of lamina V[5]. Ascending

axons lay near the ventrolateral surface of the medulla, and sent collateral branches to the

NTS (Figure 6D), but unlike Gpr83- and Tacr1-expressing ALS cells[5,60], these Calb1-

positive cells did not innervate the inferior olive. Labelling in the LPB was much more

restricted following medial spinal injections, compared to the central injections (Figure 6E,

Figure 6–figure supplement 1C,D, Figure 6–figure supplement 2B). In particular, labelling

in PBcl and PBil was greatly reduced, while the dense input to the PBrel was retained. The

input to PBrel was confirmed by immunostaining for the transcription factor Foxp2, which is

expressed in PBrel neurons (Fig 6G,H)[49]. There was also strong labelling in the lateral part

of the cPAG, (Figure 6E), but only very sparse labelling in the rostral part of the PAG

(Figure 6–figure supplement 2C,E). Input to both LPB and PAG was bilateral, but denser

on the side contralateral to the spinal injections. Labelling in the superior colliculus was

almost completely lost following medial spinal injections (Figure 6E). Within the thalamus,

the sparse labelling in VPL and Po was still present, but that in the medial thalamus was

Click here to enter text.

far weaker (Figure 6F,I,J). In addition, medial spinal injections resulted in labelling in the PoT nucleus of the thalamus (Figure 6–figure supplement 2C,D), which is a known target for lamina I ALS neurons[56,61] and is thought to convey non-noxious thermal information to the posterior insular cortex[53]. Very few, if any, tdTomato-labelled axons were seen in any other parts of the diencephalon[62]. High magnification scans showed that in the regions targeted by tdTomato labelled axons (NTS, LPB, PAG, thalamus) these axons possessed numerous varicosities, which presumably correspond to axonal boutons (Figure 6–figure supplement 2F). Immunostaining for the postsynaptic density protein Homer1 revealed that varicosities were often apposed to Homer1 puncta (Figure 6–figure supplement 2G,H), confirming that they constituted the presynaptic elements at glutamatergic synapses.

We have previously demonstrated that although spinothalamic lamina I neurons are relatively common in the rodent cervical enlargement, they are far less numerous at lumbar levels, accounting for <5% of lamina I ALS neurons[46,63]. We therefore injected AAV1.Cre^{ON}.tdTom into the cervical enlargement of 3 Calb1^{Cre} mice (Table 1, Figure 6–figure supplement 3A,B). This resulted in far denser axonal labelling within the VPL and PoT nuclei on the side contralateral to the spinal injection (Figure 6–figure supplement 3C-E).

The differences between projection patterns of medial and central injections into the lumbar enlargement provide additional information about potential targets of Calb1 cells that were not located in lamina I. As noted above, the input to PBil and medial thalamus seen following the central injections is likely to result at least in part from capture of cells belonging to the ALS4 population in lateral lamina V[13], since we have shown that these cells (targeted in a Sst^{Cre} mouse line) project to these two sites[5]. However, we found only very sparse input from the Sst-expressing (ALS4) cells to the superior colliculus[5], raising the possibility that input to the colliculus originates mainly from cells in the LSN, many of which express Calb1. Consistent with this suggestion, Choi et al[60] observed a projection to the superior colliculus from Tacr1-expressing ALS cells, and many ALS neurons in the LSN express the neurokinin 1 receptor (NK1r), which is encoded by Tacr1[64-66].

To confirm the input to PAG from cold-selective lamina I cells, we targeted injections of AAV11.Cre^{ON}.tdTomato to the caudal PAG in 3 Calb1^{Cre};Trpm8^{Flp};RCE:FRT mice. In these experiments, the total numbers of retrogradely labelled (tdTomato-positive) lamina I cells in the L2-L4 segments were lower than was the case for the LPB injection experiments (26, 59 and 99 in the three cases), but the proportion of these cells that had dense Trpm8

Click here to enter text.

input was significantly higher: 84.6%, 81.4% and 77.8% (p < 0.0001, unpaired t-test with

Welch's correction; Figure 6–figure supplement 4). This indicates that, at least for

calbindin-expressing lamina I ALS neurons, cold-selective cells are highly enriched among

those that project to the caudal part of the PAG.

Finally, we looked for further evidence of a projection from cold-selective lamina I cells

to the lateral part of the thalamus by making small injections of cholera toxin B subunit

(CTB) in 3 Trpm8^{Flp};RCE:FRT mice. CTB was used as a retrograde tracer for these

experiments, because it can result in relatively small and well-defined injection sites. In all

3 cases, the injection was centred on the ventral posterior nucleus of thalamus, and

included both lateral (VPL) and medial (VPM) parts (Figure 6–figure supplement 5A-C).

There was some spread into Po and the ventrolateral (VL) nucleus, but none into the

medial thalamus and none of the injections extended as far caudally as the PoT. As

expected, from the low number of lamina I spinothalamic neurons at lumbar levels[46,63], few

(if any) retrogradely labelled cells were seen in the lumbar spinal cord in these

experiments (5 cells in the L4 segment of one mouse, but no cells in this segment in the

other two), and for this reason we focused our search on the cervical enlargement. In

horizontal sections through the C7 segment, we identified a large number of retrogradely

labelled lamina I cells in each case. Although the dendritic filling was less complete than

that seen with retrograde viral tracing, we were able to identify many CTB-positive cells

that received numerous contacts on cell bodies and dendrites from Trpm8-expressing

primary afferents (Figure 6–figure supplement 5D-I). In the one case in which retrogradely

labelled cells were detected in the L4 segment, 3 of the 5 cells were also densely coated

with GFP-labelled afferents (Figure 6–figure supplement 5J-L). These findings

demonstrate that cold-selective lamina I ALS neurons project to the lateral thalamus.

DISCUSSION

The main findings of this study are that: (1) lamina I projection neurons that are densely

innervated by Trpm8-expressing primary afferents correspond to the cold-selective cells

identified in physiological studies and receive monosynaptic input from Trpm8 afferents;

(2) these are captured when AAVs encoding Cre-dependent constructs are injected into

the brains of Calb1^{Cre} mice; and (3) axons of Calb1-positive neurons in the medial part of

the dorsal horn project to brain regions known to be involved in processing information

related to cold input, PBrel, caudal PAG, as well as the PoT and VPL nuclei of the

thalamus. This suggests that these sites are directly innervated by cold-selective ALS

neurons in lamina I.

Lamina I ALS neurons with dense Trpm8 input are cold-selective

We had previously observed that Trpm8-expressing primary afferents are closely

associated with the cell bodies and dendrites of a subset of ALS cells in lamina I[13]. Based

on their association with Homer1 puncta, we were also able to estimate that GFP-labelled

(Trpm8-expressing) afferents accounted for ~60% of the excitatory synapses on these

ALS cells. Here we directly demonstrate the presence of synapses with both electron

microscopy and optogenetics, and confirm our prediction[13] that these densely Trpm8-

innervated ALS cells correspond to the cold-selective neurons identified in previous

physiological studies[23-33].

Cold-selective lamina I projection neurons have been reported in several species,

including monkey[27], cat[23-26], rat[28,30] and mouse[29,31-33], and a recent *in vivo* calcium-imaging

study reported that they accounted for ~15% of lamina I spinoparabrachial neurons in the

mouse[33]. Although cold-selective lamina I projection neurons respond predominantly to

skin cooling, some show a weak "paradoxical" response when noxious heat is applied to

the skin[24,26,27,29]. We did not detect any response from cold-selective cells when hot saline

was applied to the skin in the experiments reported here, or in our previous study of cold-

selective neurons[32], and this may be because although the temperature of the hot saline

was 50°C, the small volume applied to the skin meant that skin temperature did not reach

a sufficiently high level to result in activation. It is known that some Trpm8-expressing

primary afferents have low levels of Trpv1[20,35,37], and some Trpm8 afferents have been

shown to respond weakly to noxious heat *in vivo*[35]. It has been proposed that there are two

different populations of Trpm8-expressing afferents: one consisting of cells that lack Trpv1

and respond to innocuous cooling, and the other of cells that express Trpv1, and may

function as "cold nociceptors"[67]. Since we found Trpv1-immunoreactivity in around a

quarter of GFP-labelled dorsal root ganglion cells, it is likely that both populations are

included amongst the GFP-expressing cells in the Trpm8[Flp];RCE:FRT mice. However, we

were unable to detect Trpv1-immunoreactivity in the central terminals of any GFP-positive

axons (presumably reflecting the low level of Trpv1 expressed by these cells), and we

therefore do not know whether the Trpm8-positive axons synapsing on cold-selective

lamina I ALS neurons include those that express Trpv1. However, if this is the case, it is

Click here to enter text.

likely to explain the weak response to noxious heat reported for some cold-selective

projection neurons[24,26,27,29].

Direct and indirect input from Trpm8 afferents to cold-selective ALS neurons

Our findings clearly demonstrate a major direct (monosynaptic) input from Trpm8-

expressing primary afferents to cold-selective lamina I projection neurons. We had

previously estimated that a mean of 62% of the Homer1 puncta (which correspond to

excitatory synapses) on cell bodies and dendrites of these cells were associated with

GFP-positive boutons in Trpm8[Flp]:RCE:FRT mice[13]. However, we show here that this line

only labels ~83% of Trpm8-expressing primary afferent neurons with GFP. If this

proportion applies equally to Trpm8 afferents that synapse on cold-selective ALS cells,

then Trpm8 afferents would give rise to ~75% of the excitatory synapses on these cells.

Lee et al[34] recently proposed that there is a dedicated population of superficial dorsal

horn excitatory interneurons that express the thyrotropin releasing hormone receptor (Trhr)

and form a disynaptic link between Trpm8 afferents and cold-selective lamina I projection

neurons. Interestingly, in their initial experiments, they ablated or silenced calbindin-

expressing dorsal horn neurons and found that this reduced behavioural responses to skin

cooling. Although they interpreted this as resulting from loss of function of calbindin-

expressing excitatory interneurons, our findings indicate that they are also likely to have

ablated or silenced calbindin-expressing cold-selective lamina I ALS neurons. This could

therefore have made a major contribution to the reduced sensitivity to skin cooling. Having

obtained evidence that cold-responsive Calb1+ neurons in the dorsal horn often co-

expressed Trhr, Lee et al generated a Trhr[Cre] mouse line and used spinal injection of AAVs

coding for Cre-dependent constructs, or a genetic intersectional approach, to target Trhr-

expressing spinal neurons[34]. They ablated these cells and found that this selectively

reduced responses to skin cooling. They also reported that when AAV.Cre[ON].YFP was

injected intraspinally in Trhr[Cre] mice, there was no axonal labelling in the brain, suggesting

that Trhr-expressing dorsal horn neurons were exclusively interneurons. However, this

observation is not consistent with our transcriptomic dataset[13], which reveals Trhr

expression in some Phox2a-derived ALS neurons, including some of those in the ALS3

cluster (Figure 7-figure supplement 1). It is therefore not clear why a brain projection from

Trhr cells was not detected by Lee et al, although it is possible that the survival time

allowed for anterograde transport of YFP (reported as 2-3 weeks) was not sufficient. Lee

et al[34] also reported that the cold-selective ALS neurons were marked by expression of the

calcitonin receptor-like receptor (encoded by the *Calcrl* gene). However, this is unlikely to

be a specific marker, as our data show expression of Calcrl among several populations of

Phox2a-derived ALS neurons[13] (Figure 7-figure supplement 1), and it has been reported

that Calcrl is also expressed by spinoparabrachial neurons that are required for

mechanical itch[68].

While our findings do not exclude the possibility that there is a specific population of

interneurons that link Trpm8-expressing primary afferents to cold-selective ALS neurons in

lamina I, they clearly demonstrate a substantial direct (monosynaptic) input from Trpm8

afferents to cold-selective lamina I projection neurons, and indicate the need for caution in

relying on the specificity of individual genetic markers to define neurons and circuits in the

dorsal horn.

Brain targets of cold-selective ALS cells

At present there is no known molecular marker that could be used to target cold-

selective lamina I ALS cells exclusively. We chose calbindin because *Calb1* is among the

top differentially expressed genes in the ALS3 cluster[13], and there is a well-established

knock-in Cre line available[69]. Importantly, our transcriptomic analysis[13] was restricted to

projection neurons derived from the Phox2a lineage[14], which excludes most ALS cells in

the LSN and ~40% of those in lamina I[14,46]. Menetrey et al[45] had demonstrated that

calbindin-immunoreactive spinoparabrachial neurons in rat were located in lamina I, the

LSN and lateral lamina V, with scattered cells around the central canal, and this closely

matches the distribution of cells labelled when AAV11.Cre^{ON}.tdTomato was injected into

the LPB of Calb1^{Cre} mice. For anterograde tracing, we were able to exclude LSN and

lateral lamina V ALS cells by injecting smaller volumes into the medial part of the dorsal

horn, and this was confirmed by the lack of axonal labelling in regions known to receive

input from these cells (e.g., PBil and medial thalamus)[5,54,56]. However, although our medial

injections preferentially captured cold-selective lamina I ALS neurons they will also have

labelled cells that are not cold-selective, including some lamina I cells in the ALS2 cluster[13]

and potentially some of those that are not derived from the Phox2a lineage[46]. We therefore

focused our analysis on brain regions known to be involved in cold sensing, and tested

whether these received input from the Calb1-positive ALS cells.

Among the ALS projection targets that contain cells responding to cold, we were able to

show a dense input to the rostral part of the LPB and to the cPAG. Two studies have

mapped neurons in LPB that express Fos in mice exposed to low ambient

Click here to enter text.

temperatures[49,50], and both found that Fos-positive cells were concentrated in the rostral

part of the LPB, corresponding to a level ~4.9 mm caudal to bregma (or 1.1 mm caudal to

the interaural line)[62]. Yang et al[50] identified the region containing Fos+ cells as part of the

external lateral (PBel) nucleus, while Geerling et al[49] defined it as the PBreI, and showed

that this area could be recognised by the expression of Foxp2. Yang et al[50] reported that

cold-activated cells in this region projected to the preoptic area (POA), which is known to

form part of a cold-defence circuit[70], as well as to the dorsomedial hypothalamus (DMH).

They also showed that parabrachial neurons targeting DMH promoted activation of brown

adipose tissue, suggesting a role for both of these hypothalamic areas in maintenance of

body temperature in cold conditions. The cold-selective lamina I ALS cells are therefore

likely to provide an important source of input to POA and DMH via their projection to PBreI,

thus forming part of the afferent limb of a cold defence pathway (Figure 7).

Neurons in the cPAG show increased Fos expression in rats exposed to low ambient

temperatures[48], and this region receives an input from cold-activated cells in the

hypothalamus[51]. Our results demonstrate that Calb1-positive neurons in the dorsal horn

also provide a dense input to the cPAG. Although we cannot be certain that this originates

from the cold-selective ALS neurons, there are two lines of evidence to support this

suggestion. First, Li et al[71] demonstrated innervation of mouse lamina I spino-PAG

neurons by Trpm8-expressing primary afferents, and second, we found that ~80% of

Calb1 neurons retrogradely labelled from cPAG received dense Trpm8 input. Consistent

with this interpretation, we had previously reported that rat lamina I neurons retrogradely

labelled from the PAG seldom showed strong expression of the NK1r, unlike those labelled

from the LPB or CVLM[72], and cold-selective lamina I neurons have been shown to respond

weakly, if at all, to substance P, the main ligand for this receptor[32]. These findings suggest

that the cold-selective lamina I neurons provide a direct input to cPAG, which presumably

contributes to the responsiveness of neurons in this region to cooling[48]. Activation of

neurons in the cPAG results in brown adipose tissue thermogenesis[73], suggesting a role in

maintenance of body temperature in cold conditions (Figure 7).

Our CTB injections into the lateral thalamus were centred on the ventral posterior

nucleus, with a varying degree of spread into Po, which was also found to receive input

from Calb1-positive ALS neurons. It is not possible to determine precisely the region(s)

from which retrograde transport would occur, but since we identified lamina I cells in the

C7 segment (and in one case also in the L4 segment) with relatively strong CTB labelling

that were associated with GFP-labelled axons, these are likely to have projected to the

core of the injection site, i.e. the ventral posterior nucleus. As noted above, spinothalamic

lamina I neurons are infrequent in the rodent lumbar enlargement[46,63], and this presumably

accounts for the very sparse labelling in VPL that we saw following injections of

AAV1.Cre^ON.tdTomato into the lumbar spinal cord, compared to that seen after cervical

injections. However, our findings from the retrograde tracing experiments suggest that

cold-selective cells project to VPL, and this is consistent with the demonstration of input

from cervical lamina I neurons to VPL in the rat[56]. In addtion, we identified an input to the

PoT nucleus of thalamus, which projects to the posterior insular cortex[53]. Together, the

input from cold-selective lamina I ALS cells to PoT and VPL is likely to contribute to the

perception of cold stimuli applied to the skin through the projections of these thalamic

nuclei to the posterior insular and somatosensory cortices[53] (Figure 7).

MATERIALS AND METHODS

Key resources table

Reagent or resource	Source	Identifier
Mouse lines		
Trpm8^Flp	Dr Mark Hoon	N/A
B6.Cg-Gt(ROSA)26Sor^tm9(CAG-tdTomato)Hze/J (Ai9)	Prof. Hongkui Zheng; available from The Jackson Laboratory, ME, USA	Cat#: 007909 RRID: IMSR_JAX:007909
Phox2a::Cre	Prof Artur Kania	N/A
STOCK Gt(ROSA)26Sor^tm1.2(CAG-EGFP)Fsh/Mmjax (RCE:FRT)	The Jackson Laboratory, ME, USA	Cat#: 032038 RRID:MMRRC_032038-JAX
B6;129S-Calb1^tm2.1(cre)Hze/J (Calb1^Cre)	The Jackson Laboratory, ME, USA	Cat#: 028532 RRID:IMSR_JAX:028532
AAV constructs		
pENN.AAV.CB7.CI.mCherry.WPRE.RBG (AAV9.mCherry)	Penn Vector Core, PA, USA; available from Addgene, MA, USA	Cat#: 105544-AAV9 RRID:Addgene_105544
pssAAV-2-CAG-EGFP_Cre-WPRE-SV40p(A) (AAV9.EGFP_Cre)	Viral Vector Facility, University of Zurich, Switzerland	Cat#: v25-9
pssAAV-2-CAG-dlox-tdTomato(rev)-dlox-WPRE-bGHp(A) (AAV1.Cre^ON.tdTomato)	Viral Vector Facility, University of Zurich, Switzerland	Cat#: v167-1
AAV11.CAG.Cre^ON.tdTomato (AAV11.Cre^ON.tdTomato)	BrainCase Biotechnology Co Ltd, Wuhan, China	Cat#: BC-0870
AAV.PHP.S.Flp^ON.ChR2_YFP	Viral Vector Facility, University of Zurich, Switzerland	Cat#: v237-PHP.S
AAV11.CAG.tdTomato (AAV11.tdTomato)	BrainCase Biotechnology Co Ltd, Wuhan, China	Cat#: BC-0868

Click here to enter text.

Antibodies		
Chicken anti-GFP (1:1000)	Abcam plc	Cat#: ab13970 RRID: AB_300798
Rabbit anti-GFP (1:1000)	M Watanabe	RRID:AB_2571573
Rat anti-mCherry (1:1000)	Thermo Fisher Scientific	Cat#: M11217 RRID: AB_2536611
Guinea pig anti-NeuN (1:1000)	Synaptic Systems	Cat#: 266 004 RRID:AB_2619988
Rabbit anti-NeuN-Alexa Fluor® 647 (1:1000)	Abcam plc	Cat#: EPR12763 RRID:AB_2532109
Rat anti-substance P (1:200)	Oxford biotechnology	Cat#: OBT06435
Rabbit anti-substance P (1:1000)	Peninsula	Cat#: T-4107 RRID:AB_518630
Rabbit anti-TRPV1 (1:1000)	Synaptic Systems	Cat#: 444 013 RRID:AB_2864792
Sheep anti-CGRP (1:2000)	Enzo	Cat#: BML-CA1137 RRID:AB_2243859
Rabbit anti-somatostatin (1:500)	Peninsula	Cat#: T-4103 RRID:AB_518614
Guinea pig anti-VGLUT3 (1:100)	M Watanabe	RRID:AB_2571856
Sheep anti-Foxp2 (1:500)	Biotechne	Cat#: AF5647 RRID:AB_2107133
Goat anti-cholera toxin B subunit (1:2000 for immunofluorescence; 1:100,000 for immunoperoxidase)	List Biological	Cat#: 703 RRID:AB_10013220
Goat anti-Homer1 (1:500)	M Watanabe	RRID:AB_2631104
Rabbit anti-calbindin (1:1000)	Swant	Cat#: CB38 RRID:AB_10000340
Alexa Fluor™ Plus 405 Donkey anti-Rabbit IgG (1:500)	Thermo-Fisher Scientific	Cat#: A48258 RRID: AB_2890547
Alexa Fluor® 488 Donkey Anti-Chicken IgY (1:500)	Jackson ImmunoResearch	Cat#: 703-545-155 RRID: AB_2340375
Alexa Fluor® 488 Donkey anti-Goat IgG (1:500)	Jackson ImmunoResearch	Cat#: 705-545-147 RRID: AB_2336933
Alexa Fluor® 488 Donkey Anti-Rabbit IgG (1:500)	Jackson ImmunoResearch	Cat#: 711-545-152 RRID: AB_2313584
Alexa Fluor™ Plus 488 Donkey anti-Rabbit IgG (1:500)	Thermo-Fisher Scientific	Cat#: A32790 RRID: AB_2762833
Rhodamine Red™-X Donkey Anti-Guinea pig IgG (1:100)	Jackson ImmunoResearch	Cat#: 706-295-148 RRID: AB_2340468
Rhodamine Red™-X Donkey Anti-Rabbit IgG (1:100)	Jackson ImmunoResearch	Cat#: 711-295-152 RRID: AB_2340613
Rhodamine Red™-X Donkey Anti-Rat IgG (1:100)	Jackson ImmunoResearch	Cat#: 712-295-153 RRID: AB_2340676
Alexa Fluor™ Plus 555 Donkey anti-Rat IgG (1:500)	Thermo-Fisher Scientific	Cat#: A48270 RRID:AB_2896336
Alexa Fluor™ Plus 555 Donkey anti-Goat IgG (1:500)	Thermo-Fisher Scientific	Cat#: A32816 RRID:AB_2762839

Click here to enter text.

Alexa Fluor® 647 Donkey Anti-Rabbit IgG (1:500)	Jackson ImmunoResearch	Cat#: 711-605-152 RRID:AB_2492288
Alexa Fluor® 647 Donkey Anti-Goat IgG (1:500)	Jackson ImmunoResearch	Cat#: 705-605-147 RRID: AB_2340437
Alexa Fluor® 647 Donkey Anti-Guinea pig IgG (1:500)	Jackson ImmunoResearch	Cat#: 706-605-148 RRID: AB_2340476
Biotin-SP Donkey Anti-Goat IgG (1:500)	Jackson ImmunoResearch	Cat#: 705-065-147 RRID: AB_2340397
Biotin-SP Donkey Anti-Guinea pig IgG (1:500)	Jackson ImmunoResearch	Cat#: 706-065-148 RRID: AB_2340451
Biotin-SP Donkey Anti-Rabbit IgG (1:500)	Jackson ImmunoResearch	Cat#: 711-065-152 RRID: AB_2340593
Avidin conjugates		
Extravidin-Peroxidase	Sigma-Aldrich	Cat#: E2886 RRID:AB_2620165
Streptavidin-Pacific Blue	Life Technologies	Cat#: S11222
HCR probe		
Mm-Trpm8 v3.0	Molecular Instruments	
Chemicals		
Cholera toxin B subunit	Sigma-Aldrich	Cat#: C9972
DAPI	Sigma-Aldrich	Cat#: D9542
Sodium borohydride	Sigma-Aldrich	Cat#: 452882
3,3'-Diaminobenzidine	Revvity	Cat#: NEL938001EA
Hydrogen peroxide	Sigma-Aldrich	Cat#: H1009
Osmium tetroxide	Agar Scientific	Cat#: R1016
Uranyl acetate	Agar Scientific	Cat#: R1260A
Lead citrate	Agar Scientific	Cat#: R1210
Software		
Neurolucida	MBF Bioscience	https://www.mbfbioscience.com/neurolucida RRID: SCR_001775
Neurolucida Explorer	MBF Bioscience	https://www.mbfbioscience.com/neurolucida-explorer RRID: SCR_017348
pClamp	Molecular Devices	https://www.moleculardevices.com/products/axon-patch-clamp-system/acquisition-and-analysis-software/pclamp-software-suite#gref RRID: SCR_011323
Easy Electrophysiology	Easy Electrophysiology LTD	https://www.easyelectrophysiology.com
Zen Black	Carl Zeiss	https://www.zeiss.com/microscopy/int/products/microscope-software/zen.html RRID: SCR_018163
Prism	GraphPad Software	https://www.graphpad.com/scientific-software/prism/

Click here to enter text.

		RRID: SCR_002798
Photoshop CS6	Adobe Systems Incorporated	https://www.adobe.com/
Xara Xtreme v2.0	Xara Group Ltd	https://www.xara.com/
Inkscape	Software Freedom Conservancy	https://inkscape.org/

Experimental Model and Subject Details

All experiments were approved by the Ethical Review Process Applications Panel of the University of Glasgow, and were carried out in accordance with the UK Animals (Scientific Procedures) Act 1986 and ARRIVE guidelines. The following transgenic mouse lines were used in this study: the BAC transgenic Phox2a::Cre line, which expresses Cre recombinase under control of the Phox2a promoter[14]; the Ai9 Cre reporter line, in which a loxP-flanked STOP cassette prevents CAG promoter-driven transcription of tdTomato[69]; the Trpm8[Flp] line, in which 2A-linked FlpO recombinase is fused with the last codon (and replaces the stop codon) of the *Trpm8* gene[13]; the RCE:FRT line, which carries the R26R CAG-boosted EGFP reporter allele with a FRT-flanked STOP cassette upstream of the EGFP gene[74]; and the Calb1[Cre] line which has an IRES2 sequence and a Cre recombinase gene inserted downstream of the translational STOP codon of the calbindin 1 gene[75]. Further details of these lines can be found in the Key Resources table. The following crosses were generated by cross-breeding these lines: Trpm8[Flp];RCE:FRT mice, which were used for anatomical analysis of Trpm8-expressing DRG cells, for electrophysiological experiments, and for retrograde tracing of spinothalamic neurons; Trpm8[Flp];RCE:FRT;Ai9 mice, which were used for some electrophysiological experiments; Trpm8[Flp];RCE:FRT;Phox2a::Cre;Ai9 mice, which were used for the combined confocal and electron microscopic analysis; and Calb1[Cre];Trpm8[Flp];RCE:FRT mice, which were used to determine the proportion of retrogradely labelled Calb1-expressing cells that received dense Trpm8 input.

Perfusion fixation was performed while mice were deeply anaesthetised with pentobarbitone (20 mg i.p.). A brief rinse with Ringer's solution was followed by perfusion with ~250 ml of fixative. In all cases apart from those used for combined confocal/electron microscopy the fixative consisted of 4% freshly depolymerised formaldehyde in phosphate buffer (PB). Spinal cords and brains were dissected and stored in the same fixative for 2 hours, before being transferred to 30% sucrose in PB.

Dorsal root ganglia for FISH were obtained from Trpm8[Flp];RCE:FRT mice, which had been killed with an overdose of pentobarbital (40 mg i.p.) and perfused with cold phosphate buffered saline (PBS). The ganglia were fixed in 4% formaldehyde for 2 hours.

Click here to enter text.

Mice of both sexes, aged 4-14 weeks and weighing between 13 and 25 g at the start of
any surgical procedures were used in the study.

Fluorescent *in situ* hybridisation, immunohistochemistry and imaging

FISH was performed on intact lumbar dorsal root ganglia from Trpm8[Flp];RCE:FRT mice
using HCR RNA-FISH v3.0 reagents. Tissues were dehydrated in graded ethanol
solutions, followed by rinsing in PBS. Hybridisation and amplification were performed with
a probe against mouse *Trpm8* mRNA and hairpins conjugated to Alexa Fluor 546 (HCR
v3.0, Molecular Instruments, Inc.) according to the manufacturer's instructions. Tissues
were mounted using antifade mounting medium and stored at -20°C.

Sections of brain, spinal cord and dorsal root ganglia for immunohistochemical
processing were cut on either a vibrating blade microtome (Leica VT1000 or VT1200) or a
cryostat (Leica CM1950). Section thickness was 50 μm for brain and 60 μm for spinal cord
and dorsal root ganglia. Multiple-label immunofluorescence microscopy was carried out as
described previously[5,59,76]. Reactions were performed on free-floating sections, and the
sources and concentrations of antibodies are listed in the Key Resources Table. Sections
were incubated for 1-3 days at 4°C or room temperature in primary antibodies diluted in
PBS that contained 0.3M NaCl, 0.3% Triton X-100 (except for tissue processed by the
confocal/EM method) and 5% normal donkey serum. After rinsing they were incubated for
2-18 hours in appropriate species-specific secondary antibodies that were raised in
donkey and conjugated to Alexa Fluor Plus 405, Alexa Fluor 488, Alexa Fluor Plus 488,
Alexa Fluor Plus 555, Alexa Fluor 647 or Rhodamine Red, in the same diluent. Following
the reaction, sections were mounted with anti-fade medium and stored at -20°C.

Tissue sections and dorsal root ganglia that had undergone immunofluorescence
labelling or FISH were scanned with either a Zeiss LSM710 confocal microscope with
Argon multi-line, 405 nm diode, 561 nm solid state and 633 nm HeNe lasers, or with a
Zeiss LSM900 Airyscan confocal microscope with 405, 488, 561 and 640 nm diode lasers.
Confocal image stacks were obtained through a 20x dry lens (numerical aperture, NA,
0.8), or through 40x (NA 1.3) or 63x (NA 1.4) oil immersion lenses with the confocal
aperture set to 1 Airy unit or less.

The method for combining confocal and electron microscopy was the same as that
described previously[77]. Two Trpm8[Flp];RCE:FRT;Phox2a::Cre;Ai9 mice were perfused with
fixative that contained 0.5% glutaraldehyde and 4% formaldehyde. Horizontal sections of
spinal cord were obtained from lumbar segments, and these were treated with 50%

ethanol for 30 mins, to enhance antibody penetration, and then with 1% sodium borohydride for 30 mins to quench unbound aldehyde sites. After rinsing, they were incubated for 3 days in primary antibodies against GFP (raised in rabbit) and mCherry (raised in rat), and then overnight in a mixture of secondary antibodies: donkey anti-rabbit conjugated to Alexa Fluor 488, donkey anti-rabbit conjugated to biotin and donkey anti-rat conjugated to Rhodamine Red. Note that the anti-mCherry antibody recognises tdTomato. The antibody diluent consisted of PBS that contained 0.3M NaCl and 5% normal donkey serum. They were then incubated in avidin conjugated to horseradish peroxidase (HRP) and mounted on slides with anti-fade medium. Sections were viewed with a confocal microscope, and regions that contained tdTomato-positive cells that were densely coated with GFP-labelled axons were selected. These were scanned with the confocal microscope to generate z-series through the cell bodies and dendritic trees of the neurons. The sections were then removed from the slides, rinsed in PBS and then PB, processed with 3,3'-diaminobenzidine (DAB) and H_2O_2 to reveal peroxidase-labelled (Trpm8-positive) axons, osmicated, stained with uranyl acetate, dehydrated and flat-embedded in Durcupan resin between acetate sheets[77]. Series of ultrathin sections (silver interference colour) were cut with a diamond knife and collected on Formvar-coated slot grids. These were further stained with lead citrate and viewed with either a Philips CM100 or Jeol JEM-1400Flash electron microscope.

Intracranial and intraspinal injections

All surgical procedures involving intraspinal and/or intracranial injections were performed while the mice were anaesthetised with isoflurane (1-2%), and all animals received perioperative analgesia (buprenorphine 0.1 mg/kg and carprofen 10 mg/kg, s.c.). For both types of injection, the mice were placed in a stereotaxic frame.

For intracranial injections, lidocaine (10 mg/kg, s.c.) was injected into the skin over the skull, which was then incised. Burr holes were drilled to allow targeting at the following co-ordinates (CVLM: 1.1-1.3 mm lateral to midline, 3.5 mm caudal to lambdoid suture, 3.3-3.7 mm below brain surface; LPB: 1.1 mm lateral to the midline, 0.35 mm caudal to lambdoid suture or 4.5 mm caudal to bregma, 3.0 mm below brain surface, PAG: 0.4 mm lateral to midline, 0.4 mm caudal to the interaural line, 3.1 mm below brain surface; VPL: 1.6-1.7 lateral to the midline, 0.79-1.11 caudal to bregma, 3.4 mm below brain surface). Injections were made through glass micropipettes (outer tip diameter ~60 μm) attached to a Harvard Pump 11 Elite (for AAV injections) or a PV 800 Pneumatic Picopump (WPI; for CTB

injections). For optogenetic experiments, Trpm8^{Flp}RCE:FRT mice that had received

intraperitoneal injections of AAV.PHP.S.Flp^{ON}.ChR2_YFP (see below) were injected with

AAV11.tdTomato (2.5×10^9 GC in 500 nl) into the LPB or CVLM. Brain injections for the

electrophysiological experiments in which the semi-intact preparation was used to record

from cells with or without dense Trpm8 input consisted of AAV9.mCherry (7.5×10^8-

1.2×10^{10} GC in 500 nl) injected into the LPB or CVLM of Trpm8^{Flp};RCE:FRT mice, or

AAV9.Cre_GFP (3.5×10^9 GC in 500 nl) injected into the CVLM of a Trpm8^{Flp};RCE:FRT;Ai9

mouse. For anatomical experiments involving Calb1^{Cre};Trpm8^{Flp};RCE:FRT mice

AAV11.Cre^{ON}.tdTomato was injected into LPB (2×10^9 GC in 400 nl) or PAG (1×10^9 GC in

200 nl). For the immunohistochemical analysis of calbindin expression in Trpm8-

innervated neurons, Trpm8^{Flp};RCE:FRT mice received injections of AAV9.mCherry

(7.5×10^8-1.2×10^{10} GC in 500 nl) into the LPB. For electrophysiological recording of Calb1-

expressing neurons in the semi-intact preparation, 4 Calb1^{Cre} mice were injected with

AAV11.Cre^{ON}.tdTomato into the CVLM (2×10^9 GC in 400 nl). Injections into the lateral

thalamus consisted of 100 nl of 1% CTB. Micropipettes were left in place for ~5 mins after

the injection to limit leakage up the track. For electrophysiological experiments mice were

allowed to recover for between 9 and 22 days (see below for further details). For

anatomical experiments, survival times were 3 days for CTB injections and 12-16 days for

AAV injections and these were followed by perfusion fixation, as described above.

For intraspinal injections in lumbar spinal cord the T12 and L1 vertebrae were exposed

and clamped. Injections were made through glass micropipettes (outer tip diameter ~60

µm) attached to a 10 µL Hamilton syringe, on the right side of the spinal cord into either

the L3 segment, or the L3, L4 and L5 segments. Injections into L3 and L5 were made

through the intervertebral spaces on either side of the T13 vertebra, while those into L4

were made through a small hole drilled in the lamina of the T13 vertebra. In initial

experiments we used a similar strategy to that described in previous papers[5,64], and these

are referred to as "central" injections. We subsequently modified the technique in order to

restrict injection sites to the medial part of the dorsal horn and exclude the LSN, and we

refer to these as "medial" injections. For cervical injections, the head was held in place

with ear-bars and the T2 spine was clamped. Injections were made (as described above)

in the intervertebral space between the C6 and C7 vertebrae. The mice survived between

26 and 57 days and were then deeply anaesthetised and perfused with fixative as

described above (for further details see Table 1).

Click here to enter text.

Image analysis

Lamina I ALS neurons that received numerous contacts from GFP-positive (Trpm8-expressing) primary afferents were identified in several parts of this study, either following retrograde labelling or through expression of tdTomato (in crosses that involved Phox2a::Cre and Ai9). The cells could be readily identified by the close association of GFP-labelled axons with their dendrites, which received multiple contacts from GFP-positive boutons. In most cases the cell bodies were also associated with numerous GFP-labelled boutons, and these sometimes completely surrounded the cell body[13].

All quantitative image analyses were performed with Neurolucida for Confocal software (MBF Bioscience, Williston, VT, USA).

To determine the proportion of DRG neurons that were labelled with GFP in Trpm8[Flp];RCE:FRT mice we scanned 3 or 4 sections immunostained to reveal GFP and NeuN, and counterstained with DAPI from the L4 dorsal root ganglia from each of 3 mice (1 male, 2 female) with a confocal microscope. Z-stacks that included the complete cross-sectional area of the ganglion were analysed using a modified disector method[78]. Reference and look-up sections were set 10 μm apart, and all intervening sections were examined to reveal any neurons that might have been located between these two levels. Initially we viewed the NeuN and DAPI channels, and in this way, we identified all neurons (between 904 and 1383 per mouse) for which the top surface of the nucleus was located between reference and look-up section. We then revealed the channel for GFP and counted any of the sampled neurons that were GFP-immunoreactive.

For the comparison of *Trpm8* mRNA and GFP dorsal root ganglia from 3 Trpm8[Flp];RCE:FRT mice (1 male, 2 female) that had been reacted as described above were scanned with a confocal microscope to generate z-series through the entire volume of the ganglion. Initially, both fluorescent channels (GFP and Alexa Fluor 546) were combined such that any cell with either GFP or *Trpm8* mRNA was visible. For all of these cells (between 155-209 per mouse), the cell size was determined by drawing an outline of the cell at its maximum point and converting cross-sectional area to diameter by assuming a circular shape. The two fluorescent channels were then separated, and for each cell the presence of GFP and/or *Trpm8* mRNA was determined.

Because we found a relatively limited range of sizes for GFP/*Trpm8*-positive cells (see Figure 1C) we did not use a stereological method for analysing co-localisation with other neurochemical markers. We scanned sections of dorsal root ganglia immunoreacted to reveal GFP together with: (1) VGLUT3, (2) Trpv1, (3) CGRP and substance P, or (4)

Click here to enter text.

somatostatin. In each case tissue from 3 mice (either 2 male and 1 female, or 1 male and 2 female) was used, and z-stacks covering the entire cross-sectional area of the ganglion were analysed. Initially the GFP channel was viewed and the locations of all labelled cells were marked (between 126 and 350 per mouse for each reaction). The other channel(s) was/were then examined and the presence or absence of the other marker(s) was noted.

To determine the proportion of retrogradely labelled lamina I neurons that had dense input from Trpm8-expressing afferents in the Calb1^{Cre};Trpm8^{Flp};RCE:FRT mice injected with AAV11.Cre^{ON}.tdTomato into either LPB (4 mice, 2 male and 2 female) or PAG (3 mice, 1 male, 2 female), we examined horizontal sections cut through the L2, L3 and L4 segments. These had been immunostained to reveal GFP and tdTomato, and were scanned with a confocal microscope. The resulting z-stacks were viewed with Neurolucida software, initially with only the tdTomato channel visible, and the locations of retrogradely labelled lamina I neurons were recorded (between 32 and 81 cells per segment in each animal for the experiments involving LPB injections). For the experiments involving PAG injections, numbers of retrogradely labelled neurons were much lower, and results from the 3 segments were pooled for each animal (between 26 and 99 cells per mouse). The GFP channel was then viewed, and cells that received dense Trpm8 input were identified. The L5 segments of the 4 Calb1^{Cre};Trpm8^{Flp};RCE:FRT mice that had received injections into LPB mice were cut transversely and the locations of tdTomato-labelled cell bodies on between 8 and 10 individual sections were identified. These were then morphed onto a standard outline of the L5 segment in Inkscape, resulting in single scalable vector graphics files for each mouse.

To test whether cold-selective cells were included among those projecting to the lateral thalamus, we examined horizontal sections from the L4 and C7 segments of 3 Trpm8^{Flp};RCE:FRT mice (2 male, 1 female) that had received CTB injections into the lateral thalamus. Sections were immunoreacted to reveal CTB and GFP and scanned with a confocal microscope. We then searched for CTB-labelled cells that were associated with numerous GFP-positive axons.

For anterograde tracing following intraspinal injections of AAV1.Cre^{ON}.tdTomato in Calb1^{Cre} mice, sections were first viewed with a fluorescence microscope (Zeiss Axioscope 5) and selected sections were scanned with a confocal microscope. Identification of brain regions that contained tdTomato-labelled axons was generally based on the atlas of Franklin and Paxinos[62], apart from the LPB. For this, we used the same terminology as in

Click here to enter text.

788 our previous paper[64], and identified the PBrel nucleus based on the findings of Geerling et
789 al[49].
790
Whole spinal cord preparation

The whole spinal cord preparation was made as described previously with minor
modifications[44]. Animals were anaesthetised with isoflurane, followed by a lethal dose of
pentobarbital, and then quickly perfused with oxygenated sucrose-based artificial
cerebrospinal fluid (ACSF; in mM; 3.0 KCl, 1.2 NaH2PO4, 0.5 CaCl2, 7.0 MgCl2, 26.0
NaHCO3, 15.0 glucose, 251.6 sucrose, 1.0 Na ascorbate, and 1.0 Na pyruvate).
Immediately after perfusion, the back skin was incised and the spinal cord was quickly
excised and placed into a sucrose-based ACSF. Dura and pia-arachnoid membranes were
removed. The spinal cord was pinned into a chamber wall made from Sylgard.The spinal
cord was perfused with oxygenated normal ACSF (in mM; 125.8 NaCl, 3.0 KCl, 1.2
NaH_2PO_4, 2.4 $CaCl_2$, 1.3 $MgCl_2$, 26.0 $NaHCO_3$, and 15.0 glucose) at 25˚C.
802
Semi-intact somatosensory preparation

The semi-intact somatosensory preparation was made as described previously[31,32] with
minor modifications. Mice were anaesthetised with isoflurane, followed by a lethal dose of
pentobarbital, and then quickly perfused with oxygenated sucrose-based artificial
cerebrospinal fluid (ACSF; in mM; 3.0 KCl, 1.2 NaH_2PO_4, 0.5 $CaCl_2$, 7.0 $MgCl_2$, 26.0
$NaHCO_3$, 15.0 glucose, 251.6 sucrose, 1.0 Na ascorbate, and 1.0 Na pyruvate). The back
skin was incised, and the spinal cord was exposed by dorsal laminectomy. The right
hindlimb and the right side of the trunk with the spinal cord were excised and transferred to
the dissection/recording chamber, and tissues were submerged in the oxygenated
sucrose-based ACSF. The right hind paw skin, saphenous nerve, and femoral cutaneous
nerve were carefully isolated from the surrounding tissues. Only the L2 and L3 ganglia on
the right side were left attached to the spine. All dorsal roots except L2 and L3, dural and
pial membranes were carefully removed. The spinal cord was pinned onto a Sylgard
chamber with the right dorsal horn facing upward. The tissued was perfused with
oxygenated normal ACSF (in mM; 125.8 NaCl, 3.0 KCl, 1.2 NaH_2PO_4, 2.4 $CaCl_2$, 1.3
$MgCl_2$, 26.0 $NaHCO_3$, and 15.0 glucose) at 25˚C. Recordings were performed for up to 4
hours after dissection. The hindpaw skin was also at a temperature of 25˚C. At this
temperature Trpm8 afferents will have been active, but are likely to have adapted over the

course of the experiment. In addition, the conduction velocity of their axons will have been

lower than in the *in vivo* state.

Patch-clamp recording from lamina I projection neurons

Neurons were visualised using a fixed stage upright microscope (SliceScope;

Scientifica, Uckfield, UK) equipped with a 40x water immersion objective and a CMOS

Camera (Prime BSI Express, Teledyne Vision Solutions, Ontario, Canada). A narrow-

beam infrared LED (Opto SFH 4550; Osram, Munich, Germany, emission peak, 860 nm)

was positioned outside the solution meniscus. Fluorescent cells and axons were visualised

using LED illumination (pE-300 ultra, CoolLED, Andover, UK). Whole-cell patch clamp

recordings were made with a pipette constructed from thin-walled single-filamented

borosilicate glass using a microelectrode puller (P-1000, Sutter Instrument, CA, USA).

Pipette resistances ranged from 6 to 10 MΩ. Electrodes were filled with an intracellular

solution (in mM; 130.0 K gluconate, 10.0 KCl, 2.0 $MgCl_2$, 10.0 HEPES, 0.5 EGTA, 2.0

ATP-Na, 0.5 GTP-Na, and 0.2% Neurobiotin, pH adjusted to 7.3 with 1.0 M KOH). Data

were recorded and acquired with an amplifier (Axopatch 200B; Molecular Devices,

Wokingham, UK). The data were low-pass filtered at 2 kHz and digitised at 10 kHz with an

A/D converter (Digidata 1550B; Molecular Devices, Wokingham, UK) and stored using a

data acquisition program (Clampex 11; Molecular Devices, Wokingham, UK).

Optogenetic activation

Two Trpm8[Flp];RCE:FRT mice were given intraperitoneal injections of

AAV.PHP.S.Flp[ON].ChR2_YFP ($1.3×10^{11}$ GC in 10 µl) on postnatal day 2 and 5 weeks later

they received brain injections of AAV11.tdTomato, as described above. A blue light pulse

(pE-300 ultra, CoolLED, Andover, UK) was applied through the objective (40x) of the

microscope for 1 ms. The shutter was controlled by a TTL pulse from the A/D converter

(Digidata 1550B; Molecular Devices, Wokingham, UK). To determine whether the

recorded neuron received monosynaptic or polysynaptic input from Trpm8 afferents, we

applied 0.2 Hz photostimulation (1 ms pulse width) based on a previous study[44] with minor

modifications. Input was considered monosynaptic if there was no failure and the latency

jitter was smaller than 1ms.

Natural stimulation to the skin

To test whether cells with dense Trpm8 input were cold-selective, we made patch-clamp

recordings in mice in which Trpm8 axons were revealed with GFP and projection neurons

were labelled with either mCherry or tdTomato using semi-intact somatosensory

preparation (Fig 4A). Successful recordings were made in 6 out of 16 experiments: 5

Trpm8^{Flp};RCE:FRT mice (2 male, 3 female) that had received injections of AAV9.mCherry

into the LPB or CVLM, and one male Trpm8^{Flp};RCE:FRT;Ai9 mouse injected in the CVLM

with AAV9.EGFP_Cre. To test whether cells with dense Trpm8 input were cold-selective,

we made patch-clamp recordings in 4 Trpm8^{Flp};RCE:FRT mice (1 male, 3 female) that had

received injections of AAV9.mCherry into the LPB or CVLM, and one male

Trpm8^{Flp};RCE:FRT;Ai9 mouse injected in the CVLM with AAV9.EGFP_Cre. For the

investigation of Calb1-Cre-positive lamina I neurons, recordings were made in 4 female

Calb1^{Cre} mice that had received injections of AAV11.Cre^{ON}.tdTomato into the CVLM.

Recordings were initially made in voltage clamp mode (VH = -70mV) to reveal

spontaneous EPSCs. To find the receptive field of the recorded neurons, mechanical, hot,

and cold stimuli were gently applied to the skin. Mechanical stimulation was applied using

von Frey filaments (4-g or 10-g), and in some cases the skin was brushed with a

paintbrush. Hot and cold stimulation were administered by applying a small amount of

15˚C saline or 50˚C saline to the skin using an eye dropper. The temperature at the skin

surface was 20˚C and 38˚C, respectively. Once the receptive field was identified, each

stimulus was reapplied directly to the receptive field for 1 second. The evoked EPSCs

were recorded in voltage clamp mode (VH = -70mV), and the evoked action potentials

were recorded in current clamp mode (IH = 0 pA) or both. The events were detected with

Easy Electrophysiology software (Easy Electrophysiology, London, UK).

Quantification and statistical analysis

Data are reported as mean ± SD, unless stated otherwise. Statistical analyses were

performed using Prism software (v10, GraphPad Software, CA, USA) or SciPy library

(https://docs.scipy.org/doc/) in Python. The statistical tests used for each experiment,

including tests for multiple comparisons, are given in the appropriate figure legends. A p

value of <0.05 was considered significant, and significance markers are denoted within

figures as follows: *p < 0.05, **p < 0.01, ***p < 0.001.

Data and materials availability

Upon acceptance of this manuscript, all data will be made accessible from the Enlighten: Research Data open repository hosted at the University of Glasgow. This study did not generate any new materials, reagents or code.

ACKNOWLEDGMENTS

This research was funded in whole, or in part, by the Wellcome Trust (Grant numbers 219433/Z/19/Z and 304005/Z/23/Z) and the Medical Research Council (Grant number MR/V033638/1). For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. We are grateful to Robert Kerr, Iain Plenderleith and Erin Dunn for expert technical assistance, and to Mark Hoon and Artur Kania for the gifts of Trpm8^{Flp} and Phox2a::Cre mice.

AUTHOR CONTRIBUTIONS

A.N.R., W.M., A.C.D., E.P, A.M.B., A.J.T and J.H. conceived the project and designed experiments. A.J.T., A.M.B. and J.H. obtained funding. A.N.R., W.M., A.C.D., E.P., A.M., M.Y., A.H.C., D.S., A.M.B. and J.H. performed experiments. A.N.R., W.M., A.M., A.J.T. and J.H. analysed data. M.W. provided reagents. A.N.R., W.M., A.M.B., A.J.T. and J.H. wrote the paper. All of the authors provided feedback and contributed to editing of the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Javed, S., Viswanathan, A., and Abdi, S. (2020). Cordotomy for intractable cancer pain: a narrative review. Pain Physician *23*, 283-292.
2. Lahuerta, J., Bowsher, D., Campbell, J., and Lipton, S. (1990). Clinical and instrumental evaluation of sensory function before and after percutaneous anterolateral cordotomy at cervical level in man. Pain *42*, 23-30. 10.1016/0304-3959(90)91087-Y.
3. Chen, H., Bleimeister, I.H., Nguyen, E.K., Li, J., Cui, A.Y., Stratton, H.J., Smith, K.M., Baccei, M.L., and Ross, S.E. (2024). The functional and anatomical characterization of three spinal output pathways of the anterolateral tract. Cell reports *43*, 113829. 10.1016/j.celrep.2024.113829.
4. Ma, W., Dickie, A.C., Polgar, E., Yadav, M., Quillet, R., Gutierrez-Mecinas, M., Bell, A.M., and Todd, A.J. (2025). Expression of Tacr1 and Gpr83 by spinal projection neurons. Molecular pain *21*, 17448069251342409. 10.1177/17448069251342409.
5. Ma, W., Polgar, E., Dickie, A.C., Hajer, M.A., Quillet, R., Gutierrez-Mecinas, M., Yadav, M., Hachisuka, J., Todd, A.J., and Bell, A.M. (2025). Anatomical characterisation of somatostatin-expressing neurons belonging to the anterolateral system. Scientific reports *15*, 9549. 10.1038/s41598-025-93816-6.
6. McMahon, S.B., and Wall, P.D. (1983). A system of rat spinal cord lamina 1 cells projecting through the contralateral dorsolateral funiculus. The Journal of comparative neurology *214*, 217-223. 10.1002/cne.902140209.
7. Chung, K., Kevetter, G.A., Willis, W.D., and Coggeshall, R.E. (1984). An estimate of the ratio of propriospinal to long tract neurons in the sacral spinal cord of the rat. Neuroscience letters *44*, 173-177. 10.1016/0304-3940(84)90077-6.
8. Abraira, V.E., and Ginty, D.D. (2013). The sensory neurons of touch. Neuron *79*, 618-639. 10.1016/j.neuron.2013.07.051.
9. Ma, Q. (2022). A functional subdivision within the somatosensory system and its implications for pain research. Neuron *110*, 749-769. 10.1016/j.neuron.2021.12.015.
10. Todd, A.J. (2010). Neuronal circuitry for pain processing in the dorsal horn. Nature reviews. Neuroscience *11*, 823-836. nrn2947 [pii] 10.1038/nrn2947.
11. Wang, L.H., Ding, W.Q., and Sun, Y.G. (2022). Spinal ascending pathways for somatosensory information processing. Trends Neurosci *45*, 594-607. 10.1016/j.tins.2022.05.005.
12. Wercberger, R., and Basbaum, A.I. (2019). Spinal cord projection neurons: a superficial, and also deep, analysis. Curr Opin Physiol *11*, 109-115. 10.1016/j.cophys.2019.10.002.
13. Bell, A.M., Utting, C., Dickie, A.C., Kucharczyk, M.W., Quillet, R., Gutierrez-Mecinas, M., Razlan, A.N.B., Cooper, A.H., Lan, Y., Hachisuka, J., et al. (2024). Deep sequencing of Phox2a nuclei reveals five classes of anterolateral system neurons. Proceedings of the National Academy of Sciences of the United States of America *121*, e2314213121. 10.1073/pnas.2314213121.
14. Roome, R.B., Bourojeni, F.B., Mona, B., Rastegar-Pouyani, S., Blain, R., Dumouchel, A., Salesse, C., Thompson, W.S., Brookbank, M., Gitton, Y., et al. (2020). Phox2a defines a developmental origin of the anterolateral system in mice and humans. Cell reports *33*, 108425. 10.1016/j.celrep.2020.108425.
15. Peier, A.M., Moqrich, A., Hergarden, A.C., Reeve, A.J., Andersson, D.A., Story, G.M., Earley, T.J., Dragoni, I., McIntyre, P., Bevan, S., and Patapoutian, A. (2002). A TRP channel that senses cold stimuli and menthol. Cell *108*, 705-715. 10.1016/s0092-8674(02)00652-9.
16. McKemy, D.D., Neuhausser, W.M., and Julius, D. (2002). Identification of a cold receptor reveals a general role for TRP channels in thermosensation. Nature *416*, 52-58. 10.1038/nature719.
17. Lewis, C.M., and Griffith, T.N. (2022). The mechanisms of cold encoding. Current opinion in neurobiology *75*, 102571. 10.1016/j.conb.2022.102571.
18. Knowlton, W.M., Palkar, R., Lippoldt, E.K., McCoy, D.D., Baluch, F., Chen, J., and McKemy, D.D. (2013). A sensory-labeled line for cold: TRPM8-expressing sensory neurons define the cellular basis

for cold, cold pain, and cooling-mediated analgesia. The Journal of neuroscience : the official

journal of the Society for Neuroscience *33*, 2837-2848. 10.1523/JNEUROSCI.1943-12.2013.

19. Pogorzala, L.A., Mishra, S.K., and Hoon, M.A. (2013). The cellular code for mammalian

thermosensation. The Journal of neuroscience : the official journal of the Society for Neuroscience

33, 5533-5541. 10.1523/JNEUROSCI.5788-12.2013.

20. Takashima, Y., Daniels, R.L., Knowlton, W., Teng, J., Liman, E.R., and McKemy, D.D. (2007). Diversity

in the neural circuitry of cold sensing revealed by genetic axonal labeling of transient receptor

potential melastatin 8 neurons. The Journal of neuroscience : the official journal of the Society for

Neuroscience *27*, 14147-14157. 10.1523/JNEUROSCI.4578-07.2007.

21. Kim, Y.S., Park, J.H., Choi, S.J., Bae, J.Y., Ahn, D.K., McKemy, D.D., and Bae, Y.C. (2014). Central

connectivity of transient receptor potential melastatin 8-expressing axons in the brain stem and

spinal dorsal horn. PloS one *9*, e94080. 10.1371/journal.pone.0094080.

22. Dhaka, A., Earley, T.J., Watson, J., and Patapoutian, A. (2008). Visualizing cold spots: TRPM8-

expressing sensory neurons and their projections. The Journal of neuroscience : the official journal

of the Society for Neuroscience *28*, 566-575. 28/3/566 [pii] 10.1523/JNEUROSCI.3976-07.2008.

23. Craig, A.D., and Kniffki, K.D. (1985). Spinothalamic lumbosacral lamina I cells responsive to skin and

muscle stimulation in the cat. The Journal of physiology *365*, 197-221.

10.1113/jphysiol.1985.sp015767.

24. Craig, A.D., Krout, K., and Andrew, D. (2001). Quantitative response characteristics of

thermoreceptive and nociceptive lamina I spinothalamic neurons in the cat. Journal of

neurophysiology *86*, 1459-1480. 10.1152/jn.2001.86.3.1459.

25. Craig, A.D., and Dostrovsky, J.O. (2001). Differential projections of thermoreceptive and nociceptive

lamina I trigeminothalamic and spinothalamic neurons in the cat. Journal of neurophysiology *86*,

856-870. 10.1152/jn.2001.86.2.856.

26. Craig, A.D., and Hunsley, S.J. (1991). Morphine enhances the activity of thermoreceptive cold-

specific lamina I spinothalamic neurons in the cat. Brain research *558*, 93-97. 10.1016/0006-

8993(91)90719-c.

27. Dostrovsky, J.O., and Craig, A.D. (1996). Cooling-specific spinothalamic neurons in the monkey.

Journal of neurophysiology *76*, 3656-3665. 10.1152/jn.1996.76.6.3656.

28. Zhang, X., Davidson, S., and Giesler, G.J., Jr. (2006). Thermally identified subgroups of marginal zone

neurons project to distinct regions of the ventral posterior lateral nucleus in rats. The Journal of

neuroscience : the official journal of the Society for Neuroscience *26*, 5215-5223.

10.1523/JNEUROSCI.0701-06.2006.

29. Allard, J. (2019). Physiological properties of the lamina I spinoparabrachial neurons in the mouse.

The Journal of physiology *597*, 2097-2113. 10.1113/JP277447.

30. Andrew, D. (2009). Sensitization of lamina I spinoparabrachial neurons parallels heat hyperalgesia

in the chronic constriction injury model of neuropathic pain. The Journal of physiology *587*, 2005-

2017.

31. Hachisuka, J., Baumbauer, K.M., Omori, Y., Snyder, L.M., Koerber, H.R., and Ross, S.E. (2016). Semi-

intact ex vivo approach to investigate spinal somatosensory circuits. Elife *5*, e22866.

10.7554/eLife.22866.

32. Hachisuka, J., Koerber, H.R., and Ross, S.E. (2020). Selective-cold output through a distinct subset of

lamina I spinoparabrachial neurons. Pain *161*, 185-194. 10.1097/j.pain.0000000000001710.

33. Chisholm, K.I., Lo Re, L., Polgar, E., Gutierrez-Mecinas, M., Todd, A.J., and McMahon, S.B. (2021).

The encoding of cutaneous stimuli by lamina I projection neurons. Pain *162*, 2405-2417.

10.1097/j.pain.0000000000002226.

34. Lee, H., Hor, C.C., Horwitz, L.R., Xiong, A., Su, X.Y., Soden, D.R., Yang, S., Cai, W., Zhang, W., Li, C., et

al. (2025). A dedicated skin-to-brain circuit for cool sensation in mice. Nature communications *16*,

6731. 10.1038/s41467-025-61562-y.

35. Qi, L., Iskols, M., Shi, D., Reddy, P., Walker, C., Lezgiyeva, K., Voisin, T., Pawlak, M., Kuchroo, V.K.,

Chiu, I.M., et al. (2024). A mouse DRG genetic toolkit reveals morphological and physiological

Click here to enter text.

1012 diversity of somatosensory neuron subtypes. Cell *187*, 1508-1526 e1516.
1013 10.1016/j.cell.2024.02.006.

1014 36. Sharma, N., Flaherty, K., Lezgiyeva, K., Wagner, D.E., Klein, A.M., and Ginty, D.D. (2020). The
1015 emergence of transcriptional identity in somatosensory neurons. Nature *577*, 392-398.
1016 10.1038/s41586-019-1900-1.

1017 37. Zeisel, A., Hochgerner, H., Lonnerberg, P., Johnsson, A., Memic, F., van der Zwan, J., Haring, M.,
1018 Braun, E., Borm, L.E., La Manno, G., et al. (2018). Molecular Architecture of the Mouse Nervous
1019 System. Cell *174*, 999-1014 e1022. 10.1016/j.cell.2018.06.021.

1020 38. Li, J., Kritzer, E., Craig, P.E., and Baccei, M.L. (2015). Aberrant synaptic integration in adult lamina I
1021 projection neurons following neonatal tissue damage. The Journal of neuroscience : the official
1022 journal of the Society for Neuroscience *35*, 2438-2451. 10.1523/JNEUROSCI.3585-14.2015.

1023 39. Seal, R.P., Wang, X., Guan, Y., Raja, S.N., Woodbury, C.J., Basbaum, A.I., and Edwards, R.H. (2009).
1024 Injury-induced mechanical hypersensitivity requires C-low threshold mechanoreceptors. Nature
1025 *462*, 651-655.

1026 40. Dickie, A.C., Bell, A.M., Iwagaki, N., Polgar, E., Gutierrez-Mecinas, M., Kelly, R., Lyon, H., Turnbull, K.,
1027 West, S.J., Etlin, A., et al. (2019). Morphological and functional properties distinguish the substance
1028 P and gastrin-releasing peptide subsets of excitatory interneuron in the spinal cord dorsal horn.
1029 Pain *160*, 442-462. 10.1097/j.pain.0000000000001406.

1030 41. Ordas, P., Hernandez-Ortego, P., Vara, H., Fernandez-Pena, C., Reimundez, A., Morenilla-Palao, C.,
1031 Guadano-Ferraz, A., Gomis, A., Hoon, M., Viana, F., and Senaris, R. (2021). Expression of the cold
1032 thermoreceptor TRPM8 in rodent brain thermoregulatory circuits. The Journal of comparative
1033 neurology *529*, 234-256. 10.1002/cne.24694.

1034 42. Gutierrez-Mecinas, M., Kuehn, E.D., Abraira, V.E., Polgár, E., Watanabe, M., and Todd, A.J. (2016).
1035 Immunostaining for Homer reveals the majority of excitatory synapses in laminae I-III of the mouse
1036 spinal dorsal horn. Neuroscience *329*, 171-181. 10.1016/j.neuroscience.2016.05.009.

1037 43. Todd, A.J. (1997). A method for combining confocal and electron microscopic examination of
1038 sections processed for double- or triple-labelling immunocytochemistry. Journal of neuroscience
1039 methods *73*, 149-157. S0165-0270(97)02222-X [pii].

1040 44. Hachisuka, J., Omori, Y., Chiang, M.C., Gold, M.S., Koerber, H.R., and Ross, S.E. (2018). Wind-up in
1041 lamina I spinoparabrachial neurons: a role for reverberatory circuits. Pain *159*, 1484-1493.
1042 10.1097/j.pain.0000000000001229.

1043 45. Menetrey, D., De Pommery, J., Thomasset, M., and Baimbridge, K.G. (1992). Calbindin-D28K
1044 (CaBP28k)-like Immunoreactivity in Ascending Projections. The European journal of neuroscience *4*,
1045 70-76. 10.1111/j.1460-9568.1992.tb00110.x.

1046 46. Alsulaiman, W.A.A., Quillet, R., Bell, A.M., Dickie, A.C., Polgar, E., Boyle, K.A., Watanabe, M.,
1047 Roome, R.B., Kania, A., Todd, A.J., and Gutierrez-Mecinas, M. (2021). Characterisation of lamina I
1048 anterolateral system neurons that express Cre in a Phox2a-Cre mouse line. Scientific reports *11*,
1049 17912. 10.1038/s41598-021-97105-w.

1050 47. Han, Z., Luo, N., Ma, W., Liu, X., Cai, Y., Kou, J., Wang, J., Li, L., Peng, S., Xu, Z., et al. (2023). AAV11
1051 enables efficient retrograde targeting of projection neurons and enhances astrocyte-directed
1052 transduction. Nature communications *14*, 3792. 10.1038/s41467-023-39554-7.

1053 48. Cano, G., Passerin, A.M., Schiltz, J.C., Card, J.P., Morrison, S.F., and Sved, A.F. (2003). Anatomical
1054 substrates for the central control of sympathetic outflow to interscapular adipose tissue during cold
1055 exposure. The Journal of comparative neurology *460*, 303-326. 10.1002/cne.10643.

1056 49. Geerling, J.C., Kim, M., Mahoney, C.E., Abbott, S.B., Agostinelli, L.J., Garfield, A.S., Krashes, M.J.,
1057 Lowell, B.B., and Scammell, T.E. (2016). Genetic identity of thermosensory relay neurons in the
1058 lateral parabrachial nucleus. Am J Physiol Regul Integr Comp Physiol *310*, R41-54.
1059 10.1152/ajpregu.00094.2015.

1060 50. Yang, W.Z., Xie, H., Du, X., Zhou, Q., Xiao, Y., Zhao, Z., Jia, X., Xu, J., Zhang, W., Cai, S., et al. (2023). A
1061 parabrachial-hypothalamic parallel circuit governs cold defense in mice. Nature communications
1062 *14*, 4924. 10.1038/s41467-023-40504-6.

Click here to enter text.

51.   Yoshida, K., Konishi, M., Nagashima, K., Saper, C.B., and Kanosue, K. (2005). Fos activation in
hypothalamic neurons during cold or warm exposure: projections to periaqueductal gray matter.
Neuroscience *133*, 1039-1046. 10.1016/j.neuroscience.2005.03.044.
52.   Vestergaard, M., Carta, M., Guney, G., and Poulet, J.F.A. (2023). The cellular coding of temperature
in the mammalian cortex. Nature *614*, 725-731. 10.1038/s41586-023-05705-5.
53.   Leva, T.M., and Whitmire, C.J. (2023). Thermosensory thalamus: parallel processing across model
organisms. Front Neurosci *17*, 1210949. 10.3389/fnins.2023.1210949.
54.   Bernard, J.F., Dallel, R., Raboisson, P., Villanueva, L., and Le Bars, D. (1995). Organization of the
efferent projections from the spinal cervical enlargement to the parabrachial area and
periaqueductal gray: a PHA-L study in the rat. The Journal of comparative neurology *353*, 480-505.
10.1002/cne.903530403.
55.   Feil, K., and Herbert, H. (1995). Topographic organization of spinal and trigeminal somatosensory
pathways to the rat parabrachial and Kolliker-Fuse nuclei. The Journal of comparative neurology
353, 506-528. 10.1002/cne.903530404.
56.   Gauriau, C., and Bernard, J.F. (2004). A comparative reappraisal of projections from the superficial
laminae of the dorsal horn in the rat: the forebrain. The Journal of comparative neurology *468*, 24-
56. 10.1002/cne.10873.
57.   Gutierrez-Mecinas, M., Polgar, E., Bell, A.M., Herau, M., and Todd, A.J. (2018). Substance P-
expressing excitatory interneurons in the mouse superficial dorsal horn provide a propriospinal
input to the lateral spinal nucleus. Brain structure & function *223*, 2377-2392. 10.1007/s00429-018-
1629-x.
58.   Gutierrez-Mecinas, M., Bell, A., Polgar, E., Watanabe, M., and Todd, A.J. (2019). Expression of
neuropeptide FF defines a population of excitatory interneurons in the superficial dorsal horn of
the mouse spinal cord that respond to noxious and pruritic stimuli. Neuroscience *416*, 281-293.
10.1016/j.neuroscience.2019.08.013.
59.   Polgár, E., Dickie, A.C., Gutierrez-Mecinas, M., Bell, A.M., Boyle, K.A., Quillet, R., Ab Rashid, E., Clark,
R.A., German, M.T., Watanabe, M., et al. (2023). Grpr expression defines a population of superficial
dorsal horn vertical cells that have a role in both itch and pain. Pain *164*, 149-170.
10.1097/j.pain.0000000000002677.
60.   Choi, S., Hachisuka, J., Brett, M.A., Magee, A.R., Omori, Y., Iqbal, N.U., Zhang, D., DeLisle, M.M.,
Wolfson, R.L., Bai, L., et al. (2020). Parallel ascending spinal pathways for affective touch and pain.
Nature *587*, 258-263. 10.1038/s41586-020-2860-1.
61.   Al-Khater, K.M., Kerr, R., and Todd, A.J. (2008). A quantitative study of spinothalamic neurons in
laminae I, III, and IV in lumbar and cervical segments of the rat spinal cord. The Journal of
comparative neurology *511*, 1-18.
62.   Franklin, K.B.J., and Paxinos, G. (2007). The Mouse Brain in Stereotaxic Coordinates. Third Edition
(Elsevier).
63.   Al-Khater, K.M., and Todd, A.J. (2009). Collateral projections of neurons in laminae I, III, and IV of
rat spinal cord to thalamus, periaqueductal gray matter, and lateral parabrachial area. The Journal
of comparative neurology *515*, 629-646.
64.   Ma, W., Dickie, A., Polgar, E., Yadav, M., Quillet, R., Gutierrez-Mecinas, M., Bell, A.M., and Todd, A.
(2025). Expression of Tacr1 and Gpr83 by spinal projection neurons. Molecular pain,
17448069251342409. 10.1177/17448069251342409.
65.   Ding, Y.Q., Takada, M., Shigemoto, R., and Mizumo, N. (1995). Spinoparabrachial tract neurons
showing substance P receptor-like immunoreactivity in the lumbar spinal cord of the rat. Brain
research *674*, 336-340. 000689939500022I [pii].
66.   Marshall, G.E., Shehab, S.A., Spike, R.C., and Todd, A.J. (1996). Neurokinin-1 receptors on lumbar
spinothalamic neurons in the rat. Neuroscience *72*, 255-263. 0306452295005587 [pii].
67.   McKemy, D.D. (2013). The molecular and cellular basis of cold sensation. ACS Chem Neurosci *4*,
238-247. 10.1021/cn300193h.

Click here to enter text.

68. Ren, X., Liu, S., Virlogeux, A., Kang, S.J., Brusch, J., Liu, Y., Dymecki, S.M., Han, S., Goulding, M., and
Acton, D. (2023). Identification of an essential spinoparabrachial pathway for mechanical itch.
Neuron *111*, 1812-1829 e1816. 10.1016/j.neuron.2023.03.013.
69. Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D.,
Hawrylycz, M.J., Jones, A.R., et al. (2010). A robust and high-throughput Cre reporting and
characterization system for the whole mouse brain. Nature neuroscience *13*, 133-140. nn.2467 [pii]
10.1038/nn.2467.
70. Morrison, S.F. (2018). Efferent neural pathways for the control of brown adipose tissue
thermogenesis and shivering. Handb Clin Neurol *156*, 281-303. 10.1016/B978-0-444-63912-
7.00017-5.
71. Li, J., Serafin, E.K., and Baccei, M.L. (2023). Intrinsic and synaptic properties of adult mouse
spinoperiaqueductal gray neurons and the influence of neonatal tissue damage. Pain *164*, 905-917.
10.1097/j.pain.0000000000002787.
72. Spike, R.C., Puskar, Z., Andrew, D., and Todd, A.J. (2003). A quantitative and morphological study of
projection neurons in lamina I of the rat lumbar spinal cord. The European journal of neuroscience
18, 2433-2448.
73. Chen, X.M., Nishi, M., Taniguchi, A., Nagashima, K., Shibata, M., and Kanosue, K. (2002). The caudal
periaqueductal gray participates in the activation of brown adipose tissue in rats. Neuroscience
letters *331*, 17-20. 10.1016/s0304-3940(02)00757-7.
74. Sousa, V.H., Miyoshi, G., Hjerling-Leffler, J., Karayannis, T., and Fishell, G. (2009). Characterization
of Nkx6-2-derived neocortical interneuron lineages. Cerebral cortex *19 Suppl 1*, i1-10.
10.1093/cercor/bhp038.
75. Daigle, T.L., Madisen, L., Hage, T.A., Valley, M.T., Knoblich, U., Larsen, R.S., Takeno, M.M., Huang, L.,
Gu, H., Larsen, R., et al. (2018). A suite of transgenic driver and reporter mouse lines with enhanced
brain-cell-type targeting and functionality. Cell *174*, 465-480 e422. 10.1016/j.cell.2018.06.035.
76. Quillet, R., Gutierrez-Mecinas, M., Polgar, E., Dickie, A.C., Boyle, K.A., Watanabe, M., and Todd, A.J.
(2023). Synaptic circuits involving gastrin-releasing peptide receptor-expressing neurons in the
dorsal horn of the mouse spinal cord. Front Mol Neurosci *16*, 1294994.
10.3389/fnmol.2023.1294994.
77. Bell, A.M., Gutierrez-Mecinas, M., Stevenson, A., Casas-Benito, A., Wildner, H., West, S.J.,
Watanabe, M., and Todd, A.J. (2020). Expression of green fluorescent protein defines a specific
population of lamina II excitatory interneurons in the GRP::eGFP mouse. Scientific reports *10*,
13176. 10.1038/s41598-020-69711-7.
78. Polgár, E., Gray, S., Riddell, J.S., and Todd, A.J. (2004). Lack of evidence for significant neuronal loss
in laminae I-III of the spinal dorsal horn of the rat in the chronic constriction injury model. Pain *111*,
144-150. 10.1016/j.pain.2004.06.011 S0304395904002994 [pii].

1149

1150

Click here to enter text.

MAIN FIGURES

Figure 1. The Trpm8^{Flp} mouse line captures Trpm8-expressing primary afferent neurons. (**A**) Immunohistochemical staining for GFP (yellow) and NeuN (blue) in a dorsal root ganglion from a Trpm8^{Flp};RCE:FRT mouse. GFP is present in some of the small neurons. (**B**) Fluorescence *in situ* hybridisation with a probe for *Trpm8* mRNA (magenta) superimposed on GFP native fluorescence (green). The great majority of GFP cells contain *Trpm8* message, and *vice versa*. Some of the double-labelled cells are indicated with arrows, while the arrowhead points to a *Trpm8*-positive cell that lacks GFP. (**C**) Frequency distribution of soma diameter for cells that were GFP-/*Trpm8*+ (red bars), GFP+/*Trpm8*+ (blue bars) or GFP+/*Trpm8*- (green bars). Scale bars: (**A**) = 25 μm, (**B**) = 50 μm.

1164

Figure 2. Distribution of Trpm8 afferents in the spinal dorsal horn.

Immunohistochemical staining for GFP at different segmental levels through the spinal

cord in Trpm8^{Flp};RCE:FRT mice. GFP-labelled axons are largely restricted to lamina I,

although occasional fibres penetrate into deeper parts of the dorsal horn (arrow). Within

lamina I the distribution of GFP axons is patchy, and does not occupy the entire

mediolateral extent. Scale bar = 200 µm.

1171

1172

Click here to enter text.

Click here to enter text.

Figure 3. Evidence for monosynaptic input from Trpm8 afferents to lamina I ALS neurons that were surrounded by these afferents. (**A**-**D**) Combined confocal and electron microscopic examination of a lamina I projection neuron with numerous contacts from Trpm8-expressing afferents. (**A**) A confocal image (single optical section) from a horizontal section through lamina I, showing the cell body and part of a proximal dendrite of one of the tdTomato-expressing neurons that were analysed. The tissue was obtained from a Phox2a::Cre;Ai9;Trpm8^{Flp};RCE:FRT mouse. TdTomato expressed by the projection neuron is shown in magenta and GFP immunoreactivity in Trpm8 afferents in green. Several Trpm8 boutons contact the cell, and five of these are indicated with arrowheads. RC, rostro-caudal; ML, mediolateral. (**B**) A low magnification electron micrograph through the cell at approximately the same z-level as that shown in **A**. Although the cell does not contain an electron dense label, it can be recognised by its shape, and because it is surrounded by GFP-expressing axonal profiles that were labelled with an immunoperoxidase method. The locations of those corresponding to the boutons shown in **A** are again marked with arrowheads. The cell has been pseudocoloured magenta to show its location. (**C,D**) Progressively higher magnification EM images showing the five GFP-labelled boutons marked with arrowheads in **A** and **B**. In **D** the boutons are marked with asterisks and the locations of membrane darkening and vesicle clusters that presumably correspond to synapses on the cell body and dendrite of the projection neuron are indicated with arrowheads. (**E**) The experimental approach used for optogenetic testing of Trpm8 input to ALS neurons. Stimulation (1 ms, 0.2 Hz) evoked EPSCs in 5 of the ALS neurons that were densely coated by Trpm8 afferents. (**F**) shows an example of the response in one cell (upper trace), the latencies of EPSCs in in response to 10 consecutive stimuli (n = 5, lower graph), and on the right, the latency jitter for each of these cells (n = 5). Scale bars: (**A**,**B**) = 10 µm, (**C**) = 1 µm (**D**) = 500 nm.

Figure 4. Electrophysiological characterisation of retrogradely labelled lamina I projection neurons recorded in Trpm8^{Flp};RCE:FRT mice (**A**) Cells were identified by

Click here to enter text.

expression of mCherry following injection of AAV9.mCherry into the CVLM or LPB of

Trpm8^{Flp};RCE:FRT mice, or by expression of tdTomato following injection of

AAV9.Cre_GFP into the CVLM of Trpm8^{Flp};RCE:FRT;Ai9 mice. Note that the level of GFP

expression resulting from injection of AAV9.Cre_GFP was extremely low, and would have

been restricted to the nucleus of labelled neurons, due to the presence of a nuclear

localisation signal. In both cases, Trpm8-expressing afferents were labelled with GFP. (**B**)

The semi-intact somatosensory preparation retained skin attached to lumbar spinal cord

through intact peripheral nerves and was used for recording responses to natural skin

stimuli: Hot saline (50°C), Cold saline (15°C) or von Frey filaments (4 or 10 g) (**C**)

Fluorescence microscopy image of a retrogradely labelled (mCherry-positive) lamina I ALS

neuron (soma marked with asterisk) that is densely coated by Trpm8-expressing (GFP-

positive) axons. Note that bundles of GFP axons are closely associated with the dendrites

of the retrogradely labelled neuron, which are marked with arrowheads. (**D**) Whole-cell

recording from this cell reveals that only cold stimulation evoked action potentials (APs).

(**E**) Pie chart showing the response properties of the 6 lamina I ALS neurons that were

densely coated with Trpm8 afferents to mechanical, cold and heat stimulation of the skin.

All of these cells responded only to cold stimulation: 3 cells showed APs, 1 cell showed

EPSPs but no APs (subthreshold), while 2 cells were recorded in voltage clamp and

showed EPSCs. (**F**) Fluorescence microscopy image of a lamina I ALS neuron that was

not densely coated with Trpm8 axons. Again, the soma is marked with an asterisk and

dendrites with arrowheads. Note that although the dendrites of this cell pass through

bundles of GFP axons these bundles are not aligned with the dendrites. (**G**) Whole-cell

recording from the cell shown in (**F**). AP firing was evoked by mechanical stimulation (von

Frey filament 4.0g), but not by application of hot or cold saline to the skin. (**H**) Pie chart of

response properties of 5 lamina I ALS neurons that lacked dense Trpm8 input to

mechanical, cold and heat stimulation to the skin. Two cells were mechano-selective, while

three were polymodal (2 cells: mechano, heat, 1 cell: mechano, heat and cold). Scale bar

(**C**,**F**) = 50 μm. Scale bars in **D** apply to **G**.

1231

Click here to enter text.

1232

Figure 5 Characterisation of Calb1-expressing projection neurons. (A) The
distribution of retrogradely labelled cells in 10 transverse sections from the L5 segment of
one of the Calb1^Cre mice that had received an injection of AAV11.Cre^ON.tdTomato into the
LPB. The right side of the plot is the side contralateral to the brain injection. (B) Bar chart

showing the proportion of retrogradely labelled cells in the L2, L3 and L4 segments of the

Calb1^{Cre} mice injected with AAV11.Cre^{ON}.tdTomato into the LPB that received dense

Trpm8 input. Repeated measures 1-way ANOVA showed a significant difference (p =

0.048), and post-hoc Holm-Šídák's multiple comparisons test revealed a difference only

between L2 and L4 (p = 0.0056). (**C**) Part of a horizontal section through the L2 segment

of one of the mice, showing 4 lamina I neurons (numbered 1-4) that were retrogradely

labelled with tdTomato (tdTom, magenta). RC, rostrocaudal; ML, mediolateral (applies to

C-H). (**D**) The same field scanned to reveal both tdTomato and GFP (green). (**E-H**)

detailed views of the 4 cells shown in **C**,**D**. Three of these cells (1-3) have numerous

contacts from GFP-labelled axons on their dendrites (marked with arrowheads) and cell

bodies. The fourth cell (4) has very few such contacts. (**I**) Recordings from one of the cold-

selective cells (upper traces) and the polymodal cell (lower traces) in experiments carried

out in Calb1^{Cre} mice that had received injections of AAV11.Cre^{ON}.tdTomato into the CVLM.

The cell in the upper traces shows action potential firing in response to application of cold

saline to the skin, but no response to application of a 10g von Frey (VF) hair or hot saline.

The other cell responds to brushing of the skin with a paint brush, as well as to application

of hot and cold saline. (**J**) Pie chart showing response characteristics of the 6 cells

recorded in these experiments. Scale bars: (**C**,**D**) = 50 μm, (**E-H**) = 25 μm. **P<0.01.

1256

Click here to enter text.

Figure 6. Anterograde tracing following injections of AAV1.Cre^{ON}.tdTomato into the medial dorsal horn of Calb1^{Cre} mice (**A**-**C**) Injection sites in the medial parts of the L3, L4 and L5 segments of one of the mice used in this part of the study. Immunostaining for tdTomato is shown in yellow and for NeuN in blue. Labelling of cell bodies is largely restricted to the medial part of the dorsal horn. There is some labelling in the lateral spinal nucleus (LSN), marked by asterisks, but this is likely to result from transport in axons of local interneurons. (**D**-**F**) Sections through selected regions of the brain immunostained to reveal tdTomato (yellow), with the corresponding dark-field images shown in blue. (**D**) At the level of the caudal medulla, the main ascending fibre bundle (arrowhead) lies in the ventrolateral region on the side contralateral to the spinal injections. Collateral axons innervate the nucleus of the solitary tract (NTS). (**E**) There is a dense projection of axons

Click here to enter text.

to the rostralmost part of the lateral parabrachial area (PBrel) and the caudal part of the

PAG (cPAG) on the side contralateral to the injections. (**F**) Within the diencephalon there

is sparse labelling in the contralateral ventral posterolateral nucleus (VPL) and posterior

nucleus (Po) as well as within the medial thalamus. (**G**,**H**) The region indicated in the box

in **E** is shown at higher magnification with immunostaining for Foxp2 shown in magenta.

The overlap between arborisation of tdTomato and Foxp2-positive nuclei indicates that this

region is indeed PBrel. The box in (**G**) corresponds to the region shown at higher

magnification in Figure 6–figure supplement 2F. (**I**, **J**) show higher magnification views of

tdTomato-positive axons in the VPL and Po, respectively, corresponding to the boxes in

(**F**). Numbers in **D-F** show approximate rostrocaudal locations in relation to the interaural

line. Images in **A-E**, **G-H** are from animal #8, and those in **F**, **I**, **J** are from animal #7 in

Table 1. Scale bars: (**A-F**) = 1 mm, (**G-J**) = 200 µm.

1281

Figure 7. Proposed circuits involving cold-selective lamina I ALS neurons Our results suggest that cold-selective lamina I projection neurons, which belong to the ALS3 cluster[13] project, among other sites, to PBrel, the caudal part of the PAG (cPAG) and the ventral posterolateral nucleus of the thalamus (VPL). Cold-responsive cells in PBrel innervate the preoptic area and dorsomedial hypothalamus, which are integral components of circuits that underlie cold defence. The cPAG can indirectly activate brown adipose tissue (BAT) metabolism, also contributing to cold defence. The projection to PoT and VPL, through their connections to the posterior insular and primary somatosensory (S1) cortices, presumably underlies conscious perception of cold stimuli applied to the skin. Although we show a single ALS3 neuron projecting to different brain regions, it is possible that branches to different brain nuclei originate from specific subsets of ALS3 neurons. Note

Click here to enter text.

1295   that in addition to the direct (monosynaptic) input from Trpm8-expressing primary afferents

1296   to ALS3 neurons, there may also be indirect (e.g. polysynaptic) inputs arriving via

1297   excitatory interneurons.

1298

FIGURE SUPPLEMENTS

Figure 1–figure supplement 1. Immunohistochemical characterisation of GFP-expressing somata in dorsal root ganglia of Trpm8^{Flp};RCE:FRT mice. In each set of images GFP immunoreactivity is shown in green and other types of immunoreactivity in

magenta, red or blue. (**A-C**) GFP and Vglut3 are present in largely separate populations,

although some cells contain both proteins (one cell in this field, marked with an arrow).

Two cells containing only GFP are indicated with arrowheads, and two cells containing

only Vglut3 with double arrowheads. (**D-F**) There is a greater degree of overlap between

GFP and Trpv1, and three double-labelled cells are shown with arrows. Some cells with

only GFP or Trpv1 are indicated with single and double arrowheads, respectively. (**G-I**)

GFP and somatostatin (Som) are found in separate populations, and some labelled cells

are marked with single and double arrowheads, respectively. Note that somatostatin is

seen as a partial ring around the nucleus in cells that express the peptide. (**J-M**) GFP cells

often contain substance P (SP) but lack calcitonin gene-related peptide (CGRP), and 2 of

these are marked with arrows. A single and double arrowhead mark cells with CGRP only

and with both SP and CGRP, respectively. Scale bar: (**A-M**) = 100 μm.

1317

Click here to enter text.

1318

Figure 2–figure supplement 1. Expression of Vglut3 and lack of detectable Trpv1 in
1319

the central terminals of Trpm8 afferents. (**A**-**C**) A low magnification confocal image from
1320

a transverse section of the L2 segment in a Trpm8^{Flp};RCE:FRT mouse immunostained for
1321

GFP (green) and Vglut3 (magenta). GFP-labelled axons are restricted to lamina I, while a
1322

dense band of Vglut3-immunoreactive terminals in the inner part of lamina II (marked with
1323

arrowheads) corresponds to the central terminals of C low-threshold mechanoreceptors.
1324

The dashed line in C shows the outline of the dorsal horn and the box indicates the area
1325

shown in (**D-F**). (**D-F**) At higher magnification a few Vglut3 profiles are also seen in lamina
1326

I (arrows) and these co-express GFP. (**G-J**) A horizontal section through lamina I from a

Phox2a::Cre;Ai9;Trpm8^{Flp};RCE:FRT mouse immunostained to reveal tdTomato (tdTom,

red), GFP (green) and Vglut3 (blue). This field includes a tdTomato labelled projection

neuron that receives numerous contacts from GFP-labelled axons. Some of these are also

immunoreactive for Vglut3 and these are seen more clearly (some marked with arrows)

when only tdTomato and Vglut3 are revealed (**J**). The insets in J show two of these Vglut3

profiles (those marked with arrows numbered 1 and 2) contacting dendrites of the

projection neuron in limited z-projections (2 optical sections at 0.5 μm z-spacing). In these

images the coexpression of Vglut3 (blue) and GFP (green) is clearly visible. (**K-M**) A

transverse section through lamina I from a Trpm8^{Flp};RCE:FRT mouse immunostained to

reveal GFP (green) and Trpv1 (magenta). Although numerous Trpv1-immunoreactive

profiles are visible, labelling for Trpv1 is generally not detected in the Trpm8 afferents,

which express GFP. Scale bars: (**A-C**) = 100 μm, (**D-F**) = 50 μm, (**G-J**) = 50 μm, (**K-M**) =

20 μm, (insets in **J**) = 2 μm.

1341

1342

Click here to enter text.

1343

Figure 3–figure supplement 1. The relationship between tdTomato and GFP labelling

for the cell illustrated in Figure 3. The top pane shows immunostaining for tdTomato

(tdTom, magenta) in a horizontal section through lamina I from a

Phox2a::Cre;Ai9;Trpm8^{Flp};RCE:FRT mouse. The tdTomato-labelled projection neuron

illustrated in Figure 3 is visible. The soma is marked with an asterisk and parts of the

dendritic tree with arrowheads. The cell body and dendrites are associated with dense

bundles of GFP-labelled axons. Scale bar = 50 µm.

1351

1352

1353

1354

Click here to enter text.

1355

Figure 3–figure supplement 2. Confocal and electron microscope images of two Trpm8-innervated cells examined with the combined method in spinal cord sections from a Phox2a::Cre;Ai9;Trpm8^{Flp};RCE:FRT mouse. (**A**) Projected confocal z-stack showing parts of the two cells (numbered 1, 2) labelled with tdTomato (magenta) in a horizontal section. (**B**) a single optical section from the same confocal z-stack showing the association between axons labelled with GFP (green) and the cell body of cell 1 and dendrites of both cells. (**C-J**) Examples of synapses between the GFP-containing boutons, which are labelled with an immunoperoxidase method, and (unlabelled) dendrites (D) or soma (S) of the tdTomato-positive projection neurons. **C-F** show synapses on cell 1, and **G-J** those on cell 2. Presumed synaptic active zones (marked by arrowheads) can be recognised by the presence of vesicles on the presynaptic side and darkening of the

Click here to enter text.

1367   membrane. However, these synapses do not show prominent postsynaptic densities.

1368   Scale bars: (**A**,**B**) = 50 μm, (**C-J**) = 500 nm.

1369

Click here to enter text.

1370

Figure 3–figure supplement 3. Optogenetic evidence for possible polysynaptic input from Trpm8 afferents to a lamina I ALS cell with dense Trpm8 input An example trace of an ALS cell with monosynaptic, and possibly also polysynaptic, inputs. Note that the delayed EPSCs exhibit latency jitter.

Click here to enter text.

1375

Figure 4–Figure supplement 1. Characteristics of spontaneous EPSCs in cold-

selective and other lamina I ALS cells. (**A**) Representative traces from an ALS neuron

with dense Trpm8 input (left) and an ALS neuron without dense Trpm8 input (right). (**B**,**C**)

Quantification of sEPSC frequency (**B**) and amplitude (**C**) of ALS neurons. Box plots are

median and interquartile range with symbols representing data points from individual ALS

neurons (with dense Trpm8 input: n=6, without dense Trpm8 input: n=5; * indicates p <

0.05, Mann-Whitney U test). (**D**) Overlay of the firing time course of 3 cold-selective cells in

response to cold, heat and mechanical stimulation. Note that action potentials frequency

was determined from counts in 1 sec bins. Individual traces shown in grey and averages in

black.

Figure 5–figure supplement 1. **Further characterisation of Calb1-expressing**

projection neurons. (**A**) UMAP plot showing the distribution of cells in the ALS1-5

clusters, together with a plot for expression of Calb1. Note the presence of many Calb1-

positive cells in the ALS3 cluster. (**B**) The experimental approach used to generate data

presented in this figure and in Figure 5. (**C**) An example of a section through the L5

segment from a Calb1^{Cre} mouse injected with AAV11.Cre^{ON}.tdTomato in the LPB.

Expression of tdTomato (red) is superimposed on a dark field image. (**D-F**) The distribution

of retrogradely labelled cells in 9 (**D,F**) or 8 (**E**) transverse sections from the L5 segments

of the remaining 3 Calb1^{Cre} mice injected with AAV11.Cre^{ON}.tdTomato in the LPB. The

labelling for the other mouse is shown in Figure 5A. (**G**) Horizontal section through lamina I

from the L4 segment of a Trpm8^{Flp};RCE:FRT mouse that had been injected with

AAV9.mCherry into the LPB. The section has been immunoreacted to reveal mCherry

1400 (red), GFP (green) and calbindin (blue). A mCherry-positive lamina I neuron that is

1401 surrounded by GFP axons shows strong calbindin-immunoreactivity. Scale bar: (**C**) = 500

1402 µm, (**G**) = 20 µm.

1403

1404

Figure 5–figure supplement 2. **Electrophysiological characterisation of Calb1-**

expressing projection neurons. (**A, B**) sEPSC frequencies and amplitudes for the 5

cold-selective cells and the 1 polymodal cell recorded in Calb1[Cre] mice injected with

AAV11.Cre[ON].tdTomato in the CVLM. Note that although 2 of the cold-selective cells in this

sample had relatively high sEPSC frequencies, when data from these cells were pooled

with those reported for other sample of cold-selective cells (shown in Figure 4–Figure

supplement 1) and compared with the polymodal cells shown in that Figure supplement,

there was still a significant difference for both sEPSC frequency (p = 0.037) and amplitude

(p = 0.015) (Mann-Whitney U test). (**C**) Overlay of the firing time course of 5 cold-selective

cells in response to cold, heat and mechanical stimulation. Note that action potentials

frequency was determined from counts in 1 sec bins. Individual traces shown in grey and

averages in black.

Click here to enter text.

Figure 6–figure supplement 1. Anterograde labelling following injections of AAV1.Cre^{ON}.tdTomato into the central part of the spinal dorsal horn of Calb1^{Cre} mice. (**A**) The experimental approach used to generate data for Figure 6 and for Figure 6–figure supplements 1-3. Intraspinal injections were made into either the L3 or the L3, L4 and L5 segments, and were targeted on either the central or medial part of the dorsal horn. (**B**) Example of an injection into the central part of the L4 segment of a Calb1^{Cre} mouse. Immunostaining for tdTomato is shown in yellow and for NeuN in blue. There is extensive

tdTomato labelling throughout most of the dorsal horn, and this extends into the lateral spinal nucleus (arrowhead), which contains tdTomato-labelled cell bodies. (**C-E**) Sections through selected regions of the brain have been immunostained to reveal tdTomato (yellow), and these are shown on corresponding dark-field images (blue). (**C**) There is strong labelling in the rostralmost part of the contralateral LPB, corresponding to the PBrel, as well as in the caudal PAG (cPAG). Unlike the pattern seen with injections restricted to the medial dorsal horn, there is also strong labelling in the contralateral superior colliculus (SC). (**D**) At a slightly more caudal level there is axonal labelling in several subnuclei of the contralateral LPB, including external lateral (el), dorsolateral (dl), central lateral (cl) and internal lateral (il), but not ventrolateral (vl). The location of the superior cerebellar peduncle (SCP) is indicated. (**E**) Within the diencephalon there is sparse axonal labelling in contralateral VPL and posterior (Po) thalamic nuclei, as seen following medial injections. However, there is also dense labelling in the medial thalamus, which was not seen with medial injections. Numbers in **C-E** show approximate rostrocaudal locations in relation to the interaural line. All images are from animal #2 in Table 1. Scale bars: (**B,C,E**) = 1 mm, (**D**) = 500 μm.

Click here to enter text.

1443

Figure 6–figure supplement 2. Anterograde labelling following injections of

AAV.Cre^{ON}.tdTomato into the medial dorsal horn of Calb1^{Cre} mice. (A) A section

through the C7 segment of a mouse that received injections into L3, L4 and L5 segments.

Immunostaining for tdTomato is shown in yellow and for NeuN in blue. There is a bundle of

axons (arrow) in the contralateral dorsolateral white matter, with a much more weakly

stained bundle in the same location on the opposite side. A few collateral branches are

present in the deep dorsal horn and the area around the central canal. (**B**,**C**) Sections

through selected regions of the brain have been immunostained to reveal tdTomato

(yellow), and this is shown on corresponding dark-field images (blue). (**B**) At this level of

the contralateral LPB, there is axonal labelling in the external lateral (el) and dorsolateral

(dl) subnuclei, with very little labelling in ventrolateral (vl), central lateral (cl) or internal

lateral (il). The location of the superior cerebellar peduncle (SCP) is indicated. (**C**) At a

level corresponding to ~ 0.4 mm rostral to the interaural line there is very sparse axonal

labelling in the posterior triangular nucleus of the thalamus (PoT) and in the rostral PAG

(rPAG) on the side contralateral to the spinal injection. This is shown at higher

magnification in (**D**) and (**E**), which correspond to the boxes in (**C**). (**F**) shows a maximum

intensity projection of a z-series of 39 optical sections (0.5 µm z-separation) corresponding

to part of the image shown in Figure 6G. Several varicosities, which probably correspond

to synaptic boutons, are visible, and some are indicated with arrows. (**G**, **H**) High

magnification views of anterogradely labelled axon terminals in the PBrel of a mouse that

received injections into the L3, L4 and L5 segments. Immunostaining for tdTomato is

shown in magenta and for Homer1 in green. The image in (**G**) is a maximum intensity

projection from 23 optical sections at 0.5 µm z-separation, showing labelled axons with

varicosities. (H) shows a projection of 3 optical sections (0.5 µm z-separation). Three of

the varicosities in this field (arrows) are associated with Homer1 puncta and these are

shown at higher magnification in insets. Note that the colour used to represent tdTomato

has been altered to display the relation of Homer1 puncta to tdTomato varicosities.

Numbers in **B** and **C** show approximate rostrocaudal locations in relation to the interaural

line. Images in **A** and **C**-**E** are from animal #7, those in **B** and **F** are from animal # 8, and

those in **G** and **H** are from animal #10 in Table 1. Scale bars: (**A**,**C**) = 1 mm, (**B**,**D**,**E**) = 500

µm, (**F**-**H**) = 20 µm.

1475

1476

Click here to enter text.

1477

Figure 6–figure supplement 3. Anterograde labelling following injections of

AAV.Cre^ON.tdTomato into the cervical dorsal horn of Calb1^Cre mice (**A**) The

experimental approach used to generate data for this figure. Injections were made into the

cervical enlargement. (**B**) Injection site in the C8 segment. Immunostaining for tdTomato is

shown in yellow and for NeuN in blue. (**C**) A section through the diencephalon scanned to

reveal tdTomato (yellow) and dark-field illumination (blue) shows dense tdTomato labelling

Click here to enter text.

1484   in the VPL and Po thalamic nuclei. The region in the box is shown at higher magnification

1485   in **D**. (**D**) Dense tdTomato labelling can be seen in the VPL. (**E**) At a more caudal level,

1486   there is also dense tdTomato labelling in the PoT nucleus of thalamus. The number in C

1487   shows the approximate location in relation to the interaural line. All images are from animal

1488   #14 in Table 1. Scale bars: (**B**,**C**) = 1 mm, (**D**,**E**) = 250 μm.

Click here to enter text.

Figure 6–figure supplement 4. Retrograde labelling of Calb1-expressing ALS neurons from PAG. (**A**) TdTomato labelling at the site of injection of AAV11.Cre[ON].tdTomato in one of the 3 Calb1[Cre];Trpm8[Flp];RCE:FRT mice used in this part of the study. Note that this will not reveal the precise location of the injection, because only nearby Calb1-expressing cells will be labelled. (**B**-**D**) Labelling for tdTomato (tdTom) and GFP in a horizontal section through lamina I from the case illustrated in (**A**). Several tdTom-positive neurons are visible in this field, and all of them are densely coated with GFP-labelled (Trpm8-positive) axons. (**E**) Histogram showing the percentages of retrogradely labelled neurons in the L2-L4 segments that were densely coated with Trpm8-positive axons in this set of experiments (PAG), compared with the corresponding values from the 4 cases in which AAV11.Cre[ON].tdTomato was injected into the LPB. There was a highly significant difference between the percentages between the two injection sites (p < 0.0001, unpaired t-test with Welch's correction). Scale bars: (**A**) = 1 mm, (**B**-**D**) = 100 µm.

Click here to enter text.

Figure 6–figure supplement 5. Identification of spinothalamic lamina I neurons with dense Trpm8 input. (**A-C**) CTB injection sites in the 3 experiments. In each case there is dense labelling in the ventral posterolateral (VPL) and ventral posteromedial (VPM) nuclei, together with some labelling in the posterior (Po) nucleus, and in two cases also in the ventrolateral (VL) nucleus. Note the lack of labelling in medial thalamic nuclei (centrolateral, CL; centromedial, CM; mediodorsal, MD). Numbers show approximate rostrocaudal locations in relation to the interaural line. (**D-I**) Examples of retrogradely labelled (CTB-positive) lamina I neurons (marked with asterisks) that are associated with numerous GFP-labelled (Trpm8-expressing) axons in horizontal sections through the C7

Click here to enter text.

segments in the experiments illustrated in **A** and **B**. (**J-L**) In the experiment illustrated in **C**,

we also identified 5 CTB-labelled cells in the L4 segment, and one of these, which is

associated with numerous GFP-labelled boutons, is shown here (asterisk). Note that we

also found CTB-labelled cells associated with numerous GFP-positive boutons in the C7

segment in this experiment. In each case, the images are shown below the injection site

from the corresponding experiment. Scale bars: (**A**-**C**) = 1 mm, (**D-L**) = 50 µm.

1521

Click here to enter text.

1522

Figure 7–Figure supplement 1 UMAP plot showing the distribution of cells in the ALS1-5
clusters, together with plots for expression of Trhr and Calcrl. Note the presence of both
Trhr and Calcrl in several of the ALS clusters, including ALS3.

Click here to enter text.

1526 Table 1 Experimental details for Calb1^{Cre} mice used in anterograde tracing experiments.

1527

Animal	Sex	Injected segments	Injection location	Injection volume (nl)	Viral load (GC)	Survival (days)
1	M	L3, L4, L5	central	300	9.48×10^7	26
2	M	L3, L4, L5	central	300	9.48×10^7	55
3	F	L3, L4, L5	central	300	9.48×10^7	57
4	M	L3	central	300	9.48×10^7	42
5	F	L3	central	300	9.48×10^7	40
6	F	L3	central	300	9.48×10^7	40
7	M	L3, L4, L5	medial	150	4.74×10^7	55
8	M	L3, L4, L5	medial	150	4.74×10^7	54
9	F	L3, L4, L5	medial	150	4.74×10^7	41
10	F	L3, L4, L5	medial	150	4.74×10^7	55
11	F	L3	medial	150	4.74×10^7	41
12	F	L3	medial	150	4.74×10^7	41
13	F	L3	medial	150	4.74×10^7	53
14	M	C8	central	500	1.58×10^7	35
15	M	C8	central	500	1.58×10^7	44
16	M	C8	central	500	1.58×10^7	43

1528

1529 Mice received injections of AAV1.Cre^{ON}.tdTomato into the L3 segment only, the L3, L4 and

1530 L5 segments, or the C8 segments on the right side. Injections in lumbar cord were either

1531 targeted centrally (400 µm lateral to the midline) or medially (250 µm lateral to the midline)

1532 within the dorsal horn. Those in cervical cord were located centrally (430-450 µm lateral to

Click here to enter text.

1533   the midline). All injections were made at 300 µm below the surface of the spinal cord.

1534   Injection volume and viral load refer to each injection in animals 1-3 and 7-9.

Click here to enter text.