Postnatal maternal care impacts hypothalamic *Esrrg* gene expression, co-expression profiles, and the DNA methylome in prenatal bisphenol-exposed rats

Short title: Postnatal maternal care impacts in prenatal bisphenol-exposed rats

Samantha C. Lauby[a,b] (0000-0001-5676-1333), Dennis C. Wylie[c], Hannah E. Lapp[a] (0000-0002-3656-1753), Melissa Salazar[a], Amy E. Margolis[d] (0000-0002-3334-6783) & Frances A. Champagne[a,b*] (0000-0002-3922-084X)

[a]Department of Psychology, University of Texas at Austin, Austin, Texas, USA
[b]Center for Molecular Carcinogenesis and Toxicology, University of Texas at Austin, Austin, Texas, USA
[c]Center for Biomedical Research Support, University of Texas at Austin, Austin, Texas, USA
[d]Department of Psychiatry, The Ohio State University, Columbus, Ohio, USA

*Corresponding Author
Corresponding Author: Frances A. Champagne
Email: franceschampagne@utexas.edu
Postal Address: 108 E. Dean Keeton St., Austin, TX, USA 78712

23 **Abstract**
24

Environmental exposures co-occurring during early life have a profound influence on
neurodevelopment. Our previous work in rats suggests that postnatal maternal care
modulates the effects of prenatal exposure to bisphenols, an estrogenic endocrine
disrupting chemical, on offspring neurodevelopment. Elevated postnatal maternal
licking/grooming and prenatal bisphenol exposure have known opposing effects on
estrogen receptor alpha (*Esr1*) expression in the medial preoptic area (MPOA) of the
hypothalamus, which could impact expression of estrogen-responsive genes. Based on
this previous work, we hypothesized that postnatal maternal licking/grooming would
mitigate the effects of prenatal bisphenol exposure on *Esr1* expression and estrogen-
responsive genes in the developing MPOA. In addition, we hypothesized that there would
be interactive effects of prenatal bisphenol exposure and postnatal maternal
licking/grooming on DNA methylation, particularly nearby estrogen responsive elements.
Our results suggest that maternal postnatal licking/grooming normalized prenatal
bisphenol-induced upregulation of estrogen-related receptor gamma (*Esrrg*) expression in
female pups. These mitigating impacts were also evident in co-expression gene profiles in
female pups; the majority of which were enriched for estrogen-responsive genes. Finally,
DNA methylation analyses indicated that adding postnatal maternal licking/grooming as a
covariate influenced the number of differentially methylated regions for prenatal
bisphenol-exposed male and female pups. These differentially methylated regions were
enriched for binding sites for transcription factors that are known to interact with estrogen
receptors, suggesting some secondary effects on postnatal gene regulation. These results
suggest a novel biological mechanism in which postnatal maternal care can mitigate the
negative neurodevelopmental impacts of prenatal bisphenol exposure. These results also
suggest that postnatal tactile stimulation might be a potential intervention strategy to
mitigate the neurodevelopmental risks from prenatal endocrine disrupting chemical
exposure.

51
52 **Keywords:** Bisphenols, maternal care, estrogen receptor signaling, transcriptome, DNA
53 methylation, estrogen-related receptor gamma
54
55

Author Summary

Neurodevelopment can be shaped by both aversive and positive experiences early in life, in part due to 'epigenetic' mechanisms such as DNA methylation. Here, we follow up on our previous studies that suggest high levels of postnatal maternal care could mitigate the negative impacts of prenatal bisphenol exposure, an estrogenic endocrine disrupting chemical. We focused on gene expression and DNA methylation changes in the developing medial preoptic area, a brain area that is enriched in estrogen receptors and important for sex-specific social behaviors. We found that maternal postnatal licking/grooming normalized prenatal bisphenol-induced upregulation of estrogen-related receptor gamma (*Esrrg*) expression in female pups only. We find similar patterns in multiple co-expressed gene networks that are enriched in estrogen-responsive genes. Finally, postnatal maternal licking/grooming influenced DNA methylation patterns for prenatal bisphenol-exposed male and female pups. These results suggest a novel biological mechanism in which postnatal maternal care can mitigate the negative neurodevelopmental impacts of prenatal bisphenol exposure. This is important because it suggests that postnatal tactile stimulation could be an effective intervention against the negative neurodevelopmental impacts from prenatal endocrine disrupting chemical exposure.

1. Introduction

Early-life environmental exposures have a profound influence on neurodevelopmental trajectories and can lead to stable changes in behavior throughout life. Outside of controlled laboratory conditions, individuals are exposed to a mixture of risk and protective factors that shape neurodevelopmental trajectories via distinct and common biological pathways. In particular, interactions between early-life environmental and social experiences appear to be prevalent in predicting risk and resilience to neurodevelopmental and mental health conditions (1,2). Some examples include early-life foster care for institutionalized children (3,4), perinatal nurse-family visitation programs for families in low socioeconomic conditions (5,6), and postnatal kangaroo care/skin-to-skin contact for preterm infants (7,8). The biological mechanisms underlying these interactions are largely unknown but are critical for developing effective intervention programs to mitigate the impacts of early-life adversity.

Prenatal exposure to endocrine disrupting chemicals has been shown in human and nonhuman animal studies to alter neurodevelopmental trajectories and increase the risk of neurodevelopmental and mental health conditions (9–11). Bisphenols (BP) are commonly used plasticizers that are present in many plastics, such as food storage containers and reusable water bottles, as well as the lining of food cans. Bisphenols such as BPA and other "BPA-free" structural analogs, including BPF and BPS, have also been shown to act as endocrine disruptors that primarily affect estrogen receptor signaling (12,13). Previous studies have demonstrated that bisphenols can bind to both estrogen receptor alpha (ESR1) and estrogen receptor beta (ESR2) to exert effects on gene transcription (12,14,15). Prenatal exposure to bisphenols has been associated with negative neurodevelopmental and behavior outcomes for children in human epidemiological studies (16–22) as well as in rodent models of prenatal bisphenol exposure (23–28).

Bisphenol exposure during pregnancy has also been linked to impairments in postnatal maternal care (25,29–32), though previous work in our lab has also suggested that the impacts of prenatal bisphenol exposure on offspring neurodevelopment and some aspects of postnatal maternal care might be dissociable (33,34). Regardless of the impact of prenatal bisphenols on postnatal maternal care, variations in caregiving experienced during postnatal development can have a profound impact on molecular, physiological and behavioral trajectories (35–38). In our previous work, we found that postnatal maternal care (licking/grooming and nest attendance) interacts with prenatal bisphenol exposure to predict offspring phenotype, including neurodevelopmental milestones (eye opening timing), adult behavior (attentional set shifting performance and anxiety-like behavior in the open field), transcriptional profiles in the developing brain, and DNA methylation in the adult brain (34,39). These effects were observed to be sex-specific and dose-dependent. Interestingly, we also found that higher postnatal maternal care can mitigate the impacts of prenatal bisphenol exposure on a subset of offspring phenotypes (34), suggesting that higher postnatal maternal care received and prenatal bisphenol exposure impact similar biological pathways in opposing directions. Understanding these biological pathways are

119 important to identify and propose more targeted and effective interventions to mitigate the
120 deleterious neurodevelopmental impacts of prenatal bisphenol exposure.
121
Prenatal bisphenol exposure (25,27,40,41) and variations in postnatal maternal care (42–
45) are associated with changes in DNA methylation levels across the genome which can
lead to stable changes in gene expression. However, the underlying mechanisms linking
prenatal bisphenol exposure as well as postnatal maternal care with DNA methylation
modifications at specific loci are not well-understood. There is accumulating evidence that
steroid hormone receptor signaling links environmental exposures to DNA methylation
modifications proximal to their respective transcription factor binding sites (46–50). This
includes estrogen receptor signaling, where estrogen receptor binding at estrogen
responsive elements (EREs) has been associated with downstream changes in DNA
methylation levels proximal to EREs (51–54). Rats with perinatal BPA exposure have
decreased transcript abundance of *Esr1* in the medial preoptic area (MPOA) of the
hypothalamus later in life (55,56), particularly in females. Prenatal bisphenol exposure has
also been associated with altered expression of estrogen-responsive genes with
corresponding changes in DNA methylation levels within these genes (40,57). Therefore,
the effects of bisphenols on *Esr1* expression and estrogen receptor signaling at EREs may
confer changes in DNA methylation levels at proximal CpG sites. Concurrently, high levels
of postnatal maternal care received (licking/grooming or licking-like stimulation) has been
associated with increased gene expression and decreased DNA methylation of *Esr1*
throughout life in the MPOA (44,45,58), an effect also occurring predominantly in females.
Therefore, higher maternal licking/grooming received might mitigate the effects of prenatal
bisphenol exposure on DNA methylation proximal to EREs later in life by normalizing *Esr1*
transcript abundance and ESR1 binding to DNA in the developing MPOA, particularly in
female offspring. Consistent with this hypothesis, our prior transcriptome analyses of the
developing prefrontal cortex and amygdala found significant interactive effects of prenatal
bisphenol exposure and postnatal maternal care on co-expressed gene modules that are
related to estrogen receptor signaling (34), further suggesting that the interactions between
prenatal bisphenol exposure and postnatal maternal care converge on estrogen receptor
signaling changes. However, we have also found that the impacts of prenatal bisphenol
exposure and postnatal maternal care on DNA methylation in adult rats can be highly brain
region-specific (39), so it is important to examine the MPOA directly to confirm that these
interactive effects also extend to this brain region.
153
The primary objective of the current study was to examine the interactive effects of
prenatal bisphenol exposure and postnatal maternal licking/grooming on the transcriptome
and DNA methylation modifications at specific transcription factor binding sites, notably
EREs, in the developing MPOA. The MPOA may be especially vulnerable to prenatal
bisphenol exposure because there is a high concentration of estrogen receptors in this
brain region. The MPOA is also highly sexually dimorphic and might contribute to the sex-
specific effects observed in prenatal bisphenol exposure studies in humans and rodents.
We chose to examine PND10 brains because this age shortly follows the sensitive period
for both when exogenous estrogens can permanently alter brain sexual maturation in

females (59,60) and when variations in maternal care impact MPOA ESR1 levels (61). In
addition, we adapted a correlative method to identify DNA motifs and potential
transcription factor binding sites at differentially methylated regions. Suffix Array Kernel
Smoothing (SArKS) has been previously implemented in RNA-seq datasets and allows the
user to correlate a continuous range of gene expression changes to proximal transcription
factors binding sites (62). We hypothesized that postnatal maternal licking/grooming would
mitigate the effects of prenatal bisphenol exposure on *Esr1* expression and estrogen-
responsive genes. In addition, we hypothesized that there would be interactions between
prenatal bisphenol exposure and postnatal maternal licking/grooming on DNA methylation
modifications, particularly nearby EREs and other transcription factor binding sites. Finally,
we hypothesized that there would be sex-specific effects, specifically that there would be
larger impacts in female offspring compared to male offspring. Overall, we found that
postnatal maternal licking/grooming mitigated the impacts of prenatal bisphenol exposure
on *Esrrg* expression rather than *Esr1* in female pups only. This corresponded to significant
interactions between postnatal maternal licking/grooming and prenatal bisphenol
exposure in genome-wide gene expression and DNA methylation changes, with some
enrichment in estrogen-responsive genes and even higher enrichment in genes that are
known to interact with estrogen receptors.
181
182

183
184 **Fig 1.** Preprocessing and analysis pipelines for gene expression, DNA methylation, and
185 multi-omic analyses.
186

2. Results

187 **2. Results**
188 The preprocessing and analysis pipelines for 3' tag sequencing (tag-seq), oxidative reduced
189 representation bisulfite sequencing (oxRRBS), and multi-omic factor analyses (MOFA) are
190 summarized in **Fig 1.**
191

2.1 Interactive effects of prenatal bisphenol exposure and postnatal maternal licking/grooming on Esrrg expression in the MPOA of female pups

2.1 Interactive effects of prenatal bisphenol exposure and postnatal maternal
licking/grooming on Esrrg expression in the MPOA of female pups
We found a significant interaction between the 150 μg/kg Mixed BP exposure group and
maternal licking/grooming on estrogen-related receptor gamma expression (*Esrrg*; p =
0.0004) in the female pups (**Fig 2A**) but not the male pups (**Fig 2B**). There were no other
significant interactions with the other estrogen and estrogen-related receptors or in the
male pups. There were also no significant interactions found between prenatal bisphenol
exposure and nest attendance.
200

201
Fig 2. Sex-specific interaction between prenatal bisphenol exposure and postnatal
maternal licking/grooming on *Esrrg* expression in the medial preoptic area. A significant
interaction between the 150 μg/kg Mixed BP exposure group and maternal licking/grooming
was found in female pups (**A**) but not male pups (**B**). In females from the 150 μg/kg Mixed
BP group, maternal licking/grooming was negatively correlated with Esrrg expression.
Scatterplots are displayed with linear regression lines for each prenatal treatment group. *
p < 0.05 interaction between prenatal treatment and postnatal maternal care; # p < 0.10
interaction between prenatal treatment and postnatal maternal care
210

2.2 Limited interactive effects of prenatal bisphenol exposure and postnatal maternal licking/grooming on differential expressed genes (DEGs)

2.2 Limited interactive effects of prenatal bisphenol exposure and postnatal maternal
licking/grooming on differential expressed genes (DEGs)
We examined the effects of prenatal bisphenol exposure and its interactions with postnatal
maternal care on DEGs. **Supplementary Table S1** summarizes the DESeq2 output.
Overall, there were very few DEGs found in all comparisons. The only exception was that
there were 63 DEGs (FDR ≤ 0.10) found when examining the interactions between the 150
μg/kg Mixed BP exposure group and maternal licking/grooming in female pups (including
Esrrg); however, none of these DEGs passed FDR correction of ≤ 0.05.
219

2.3 WGCNA gene expression modules associated with interactive effects of prenatal
bisphenol exposure and postnatal maternal licking/grooming
For the WGCNA analyses, a total of 7 modules for male pups and 13 modules for female
pups were analyzed. For the male pups, there was only a significant main effect of prenatal
treatment group found for one module, where the eigenvalues for the 50 µg/kg Mixed BP
exposure group were lower than the Corn Oil group. For the female pups, there were
significant or marginal interactions between the 150 µg/kg Mixed BP exposure group and
maternal licking/grooming on five modules (Blue, Brown, Magenta, Turquoise, Yellow). More
specifically, there were significant interactions for the Brown module (t = -2.544, p =
0.0292; 2,481 genes; **Fig 3A**), the Turquoise module (t = 2.291, p = 0.0449; 5,229 genes; **Fig**
3B), and the Yellow module (t = 2.061, p = 0.0264;1,445 genes; **Fig 3C**). There were also
marginal interactions for the Blue module (t = -2.179, p = 0.0544; 4,022 genes;
Supplemental Fig S1A) and the Magenta module (t = -2.205, p = 0.0520; 980 genes;
Supplemental Fig S1B), Top GO terms for each of the modules with significant interactions
in female offspring are presented in **Fig 3D;** the top GO terms for the modules with marginal
interactions are presented in **Supplemental Fig S1C**. There were no significant interactions
found between prenatal bisphenol exposure and nest attendance; however, including nest
attendance and the interaction between prenatal bisphenol exposure and nest attendance
in the linear model was required to reveal the interactions between the 150 µg/kg Mixed BP
exposure group and maternal licking/grooming.
240

Fig 3. Interactions between prenatal bisphenol exposure and postnatal maternal
licking/grooming on co-expressed gene modules in the medial preoptic area of female
pups. Significant interactions between the 150 μg/kg Mixed BP exposure group and
maternal licking/grooming were found in the Brown (**A**), Turquoise (**B**), and Yellow (**C**)
eigengene modules. In females from the 150 μg/kg Mixed BP group, maternal
licking/grooming was negatively correlated with the Brown eigengene values and positively
correlated with the Turquoise and Yellow eigengene values. Top GO terms (**D**) from these
modules were related to estrogen receptor signaling, neurodevelopment,
neurotransmission, and cellular metabolism. Scatterplots are displayed with linear
regression lines for each prenatal treatment group. * p < 0.05 interaction between prenatal
treatment and postnatal maternal care

2.4 Gene expression module enrichment for estrogen-responsive genes and genes
that interact with estrogen receptors
Among the modules in which there were significant interactions between prenatal BP and
postnatal maternal care in female offspring (**Fig 3**), we examined enrichment for estrogen-
related and -responsive genes. In the Brown module, there was no enrichment of ESR1
found in the list of ENCODE and ChEA Consensus TFs from ChIP-X or Transcription Factor
Protein-Protein Interactions. Notably, this module contains *Esr1*, *Esr2*, and *Esrrg*. In the
Turquoise module, there was a marginal enrichment of ESR1 found in the list of ENCODE
and ChEA Consensus TFs from ChIP-X (FDR = 0.054) and a significant enrichment of ESR1
found in the list of Transcription Factor Protein-Protein Interactions (FDR = 0.003). In the
Yellow module, there was a significant enrichment of ESR1 found in the list of ENCODE and
ChEA Consensus TFs from ChIP-X (FDR = 0.038) and Transcription Factor Protein-Protein
Interactions (FDR < 0.001).

2.5 Connectivity between Esrrg and all gene expression modules with interactions
between prenatal bisphenol exposure and postnatal maternal licking/grooming
We examined the connectivity of each of the estrogen and estrogen-related receptors for
each module in which there were significant interactions between prenatal BP and
postnatal maternal care in female offspring. **Table 1** summarizes these results. Most
notably, *Esrrg* had moderate-high positive connectivity with the Brown module (73%) and
moderate-high negative connectivity with the Turquoise and Yellow modules (-69% and -
54%, respectively).

Table 1. Connectivity (kME) of estrogen receptors with genes within the co-expressed gene
modules generated by the WGCNA analysis. Values are represented as percentages of
genes within each module that have significant correlations. Negative values indicate
inverse correlations.
281

	kME Brown	kME Turquoise	kME Yellow
Esr1	32.93	-13.40	-1.53
Esr2	41.06	-34.39	-48.27

Esrra	18.76	-20.91	-14.89
Esrrb	25.80	-54.04	-17.94
Esrrg	73.13	-68.89	-53.58

2.6 Impact of postnatal maternal licking/grooming on differentially methylated regions in prenatal bisphenol-exposed pups

We examined the influence of postnatal maternal licking/grooming on differentially methylated regions (DMRs) in the developing MPOA among the prenatal treatment groups. To do so, we assessed the degree of overlap in DMRs when we include or exclude maternal licking/grooming as a covariate while comparing DNA methylation levels between each prenatal bisphenol treatment group to the Corn Oil group. Overall, while many DMRs were unchanged when including maternal licking/grooming as a covariate, there were some discrepancies found for all comparisons between prenatal bisphenol treatment groups in male and female pups. Notably, in most comparisons there were more DMRs that were added than lost when including maternal licking/grooming as a covariate.

2.6.1 DMRs influenced by maternal licking/grooming in male pups For male pups with prenatal 50 µg/kg BPA exposure, there were 43 DMRs (corresponding to 32 genes) that were unchanged, 49 DMRs (corresponding to 31 genes) that were lost, and 54 DMRs (corresponding to 38 genes) that were added when including maternal licking/grooming as a covariate (**Fig 4A**; **Supplementary Table S2**). In the prenatal 50 µg/kg Mixed BP group, there were 31 DMRs (corresponding to 16 genes) that were unchanged, 7 DMRs (corresponding to one gene) that were lost, and 47 DMRs (corresponding to 35 genes) that were added when including maternal licking/grooming as a covariate (**Fig 4B**; **Supplementary Table S3**). In the prenatal 150 µg/kg Mixed BP group, there were 44 DMRs (corresponding to 27 genes) that were unchanged, 6 DMRs (corresponding to four genes) that were lost, and 38 DMRs (corresponding to 21 genes) that were added when including maternal licking/grooming as a covariate (**Fig 4C**; **Supplementary Table S4**).

2.6.2 DMRs influenced by maternal licking/grooming in female pups For female pups with prenatal 50 µg/kg BPA exposure, there were 20 DMRs (corresponding to nine genes) that were unchanged, 23 DMRs (corresponding to 14 genes) that were lost, and 28 DMRs (corresponding to 15 genes) that were added when including maternal licking/grooming as a covariate (**Fig 4D; Supplementary Table S5**). In the prenatal 50 µg/kg Mixed BP group, there were 46 DMRs (corresponding to 27 genes) that were unchanged, 25 DMRs (corresponding to 19 genes) that were lost, and three DMRs (corresponding to two genes) that were added when including maternal licking/grooming as a covariate (**Fig 4E; Supplementary Table S6**). In the prenatal 150 µg/kg Mixed BP group, there were 60 DMRs (corresponding to 48 genes) that were unchanged, 10 DMRs (corresponding to eight genes) that were lost, and 25 DMRs (corresponding to 16 genes) that were added when including maternal licking/grooming as a covariate (**Fig 4F; Supplementary Table S7**).

Fig 4. Influence of postnatal maternal licking/grooming on differentially methylated regions (DMRs) of prenatal bisphenol-exposed male and female pups. Including maternal licking/grooming as a covariate altered DMRs in all prenatal treatment groups in both male (**A-C**) and female (**D-F**) pups. For all comparisons except for the females in the 50 μg/kg Mixed BP group, more DMRs were added than lost when adding licking/grooming as a covariate. Venn diagrams are displayed with the number of DMRs and annotated genes when excluding (green) and including (orange) maternal licking/grooming as a covariate.

329   DMRs that do not overlap are considered to be influenced by postnatal maternal
330   licking/grooming.
331
While the gene lists for each comparison were too small to run any functional enrichment
analyses (e.g., GO), there were genes related to neurotransmission (e.g., Grik1, Kcnip1,
Hcn2), neurodevelopment (e.g., Epha6, Nlgn3, Sash1), gene regulation (e.g., Arx, Zfp521,
Mir135b), and mitochondrial function (e.g., Mtfr2, Gpam, Mthfd2) throughout all of the
comparisons between prenatal bisphenol treatment and Corn Oil groups, similar to the
gene modules of interest in the WGCNA analysis. There were also several notable
neuroendocrine-related genes (Ddc, Maoa, Kiss1, Glpr1, Igf2bp2, Drd5, Gper1, Hsd17b2)
found throughout all comparisons between prenatal bisphenol treatment and Corn Oil
groups.
341

2.7 Impact of postnatal maternal licking/grooming on differentially methylated
transcription factor binding sites in prenatal bisphenol-exposed pups
We next examined the influence of postnatal maternal licking/grooming on differential
methylation of proximal transcription factor binding sites and EREs. To assess this
outcome, we performed a motif analysis using two different methods (SArKS and MEME
Suite) to identify overrepresented sequence motifs and JASPAR to match the motifs to
known transcription factor binding sites.
349

2.7.1 Differentially methylated transcription factor binding sites in male pups When
underlined{excluding} licking/grooming as a covariate in male pups, SArKS identified four significant
motifs associated with four unique transcription factors in hypomethylated regions with
prenatal 50 μg/kg BPA exposure, 23 significant motifs associated with 17 unique
transcription factors in hypermethylated regions with prenatal 50 μg/kg Mixed BP exposure,
and six significant motifs associated with five unique transcription factors in
hypomethylated regions with prenatal 150 μg/kg Mixed BP exposure (**Supplementary Table
S8**). When including licking/grooming as a covariate, SArKS identified 18 significant motifs
associated with seven unique transcription factors in hypomethylated regions with
prenatal 50 μg/kg BPA exposure, 16 significant motifs associated with 13 unique
transcription factors in hypermethylated regions with prenatal 50 μg/kg Mixed BP exposure,
and four significant motifs associated with four unique transcription factors in
hypomethylated regions with prenatal 150 μg/kg Mixed BP exposure (**Supplementary Table
S9**). Overall, when including maternal licking/grooming as a covariate, three new
transcription factor binding sites (EBF2, GLI3, TFAP2C) were identified in hypomethylated
regions with prenatal 50 μg/kg BPA exposure, three new transcription factor binding sites
(NRF1, E2F3, EGR3) were identified in hypermethylated regions with prenatal 50 μg/kg
Mixed BP exposure, and two new transcription factor binding sites (EGR3, E2F7) were
identified in hypomethylated regions with prenatal 150 μg/kg Mixed BP exposure (**Fig 5A**).
369

2.7.2 Differentially methylated transcription factor binding sites in female pups When
underlined{excluding} licking/grooming as a covariate in female pups, SArKS identified 34 significant
motifs associated with 10 unique transcription factors in hypomethylated regions and two

significant motifs associated with two transcription factors in hypermethylated regions with
prenatal 50 µg/kg Mixed BP exposure only (**Supplementary Table S10**). When <u>including</u>
licking/grooming as a covariate, SArKS identified three significant motifs associated with
two unique transcription factors in hypermethylated regions with prenatal 50 µg/kg BPA
exposure as well as two significant motifs associated with two unique transcription factors
in hypomethylated regions and four significant motifs associated with four transcription
factors in hypermethylated regions with prenatal 50 µg/kg Mixed BP exposure
(**Supplementary Table S11**). Overall, when including maternal licking/grooming as a
covariate, two new transcription factor binding sites (SP5, ZNF610) were identified with
prenatal 50 µg/kg BPA exposure and two new transcription factor binding sites (HAND2,
IKZF2) were identified with prenatal 50 µg/kg Mixed BP exposure, all within
hypermethylated regions (**Fig 5B**).
385

386
Fig 5. Influence of postnatal maternal licking/grooming on differentially methylated
transcription factor binding sites of prenatal bisphenol-exposed male and female pups

using the SArKS method. Including maternal licking/grooming as a covariate altered
overrepresented sequence motifs and predicted transcription factor binding sites in all
prenatal treatment groups in male (**A**) and female (**B**) pups (except for females from the 150
μg/kg Mixed BP group). Venn diagrams are displayed with the number of motifs and unique
transcription factors when excluding (blue) and including (purple) maternal
licking/grooming as a covariate. Motifs that do not overlap are considered to be influenced
by postnatal maternal licking/grooming.

There were several predicted transcription factor binding sites that reappeared using the
MEME Suite method in male offspring (SP5, HAND2, RBPJ) and female offspring (ELK1,
IKZF2, SP5, RBPJ, HAND2). There were also changes in motifs and predicted transcription
factor binding sites when adding postnatal maternal licking/grooming as a covariate with
MEME Suite (**Supplementary Fig S2**). In many comparisons, there were a larger number of
motifs and predicted transcription factor binding sites that were altered with postnatal
maternal licking/grooming and less overlap using the MEME Suite method than the SArKS
method.

While no estrogen receptors were found in the list of predicted transcription factor binding
sites, many of the predicted transcription factors have known interactions with estrogen
receptors. This includes transcription factors that are known to dimerize with ESR1, block
estrogen receptors from binding to DNA, facilitate transcription of *Esr1*, or are estrogen-
responsive genes themselves (**Table 2**).

Table 2. List of unique transcription factors associated with prenatal bisphenol exposure
and/or postnatal maternal licking/grooming in male and female pups and their known
interactions with estrogen receptors.

Analysis	Methylation comparison	Transcription factor name	Relationship to estrogen receptors
Males			
Unchanged with Licking/Grooming			
50 μg/kg BPA vs. Corn Oil	Hypomethylated	**ZEB1**	(63,64)
50 μg/kg BPA vs. Corn Oil	Hypomethylated	**RBPJ**	(65)
50 μg/kg BPA vs. Corn Oil	Hypomethylated	SP5	
50 μg/kg BPA vs. Corn Oil	Hypomethylated	**HAND2**	(66,67)
50 μg/kg Mixed BP vs. Corn Oil	Hypermethylated	ZNF610	
50 μg/kg Mixed BP vs. Corn Oil	Hypermethylated	E2F7	
50 μg/kg Mixed BP vs. Corn Oil	Hypermethylated	**NHLH2**	(68)
50 μg/kg Mixed BP vs. Corn Oil	Hypermethylated	ZNF417	
50 μg/kg Mixed BP vs. Corn Oil	Hypermethylated	ZFP57	
50 μg/kg Mixed BP vs. Corn Oil	Hypermethylated	**SP3**	(69)
50 μg/kg Mixed BP vs. Corn Oil	Hypermethylated	**KLF15**	(70–72)

50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	**SP1**	(69,73,74)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	YY2	
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	ZBED1	
150 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	ZNF417	
150 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	SP5	
Lost with Licking/Grooming			
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	GCM2	
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	SP5	
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	**GLI3**	(75)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	**KLF13**	(76)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	**EBF1**	(77,78)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	EBF4	
150 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	EBF2	
150 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	**HAND2**	(66,67)
150 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	**RBPJ**	(65)
Added with Licking/Grooming			
50 µg/kg BPA vs. Corn Oil	Hypomethylated	EBF2	
50 µg/kg BPA vs. Corn Oil	Hypomethylated	**GLI3**	(75)
50 µg/kg BPA vs. Corn Oil	Hypomethylated	**TFAP2C**	(79,80)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	**NRF1**	(81–83)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	**EGR3**	(84)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	E2F3	
150 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	**EGR3**	(84)
150 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	E2F7	
Females			
Unchanged with Licking/Grooming			
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	LHX1	
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	BHLHE22	
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	**RBPJ**	(65)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	SP5	
Lost with Licking/Grooming			
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	**KLF15**	(70–72)
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	ERF::NHLH1	
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	**ELK1**	(85)
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	LHX5	
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	ARGFX	
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	POU6F1	
50 µg/kg Mixed BP vs. Corn Oil	Hypomethylated	E2F7	
Added with Licking/Grooming			
50 µg/kg BPA vs. Corn Oil	Hypermethylated	ZNF610	

50 µg/kg BPA vs. Corn Oil	Hypermethylated	SP5	
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	**HAND2**	(66,67)
50 µg/kg Mixed BP vs. Corn Oil	Hypermethylated	IKZF2	

2.8 Interactive effects of prenatal bisphenol exposure and postnatal maternal licking/grooming on the multi-omic epigenome in the MPOA of female pups

We examined the influence of postnatal maternal licking/grooming on the combined DNA methylome and transcriptome in the developing MPOA using MOFA. For male pups, there were two factors that showed significant interactions of prenatal BP exposure and maternal licking/grooming; however, upon further inspection these interactions appear to be solely driven by one outlier in both factors (data not shown). For female pups, there was one factor (Factor 4) that showed significant interactions between the 150 µg/kg Mixed BP exposure group and maternal licking/grooming (t = 4.456, p = 0.0043; **Fig 6A**) as well as nest attendance (t = 3.182, p = 0.0190; **Fig 6B**). Similar to the WGCNA analysis, including nest attendance and the interaction between prenatal bisphenol exposure and nest attendance in the linear model was required to reveal the significant interaction between the 150 µg/kg Mixed BP exposure group and maternal licking/grooming.

Fig 6. Interactions between prenatal bisphenol exposure and postnatal maternal licking/grooming on Factor 4 generated from the multi-omic factor analysis (MOFA) in female pups. Significant interactions between the 150 µg/kg Mixed BP exposure group and maternal licking/grooming (**A**) and nest attendance (**B**) were found when combining the transcriptome and DNA methylome datasets. In females from the 150 µg/kg Mixed BP group, Factor 4 values were positively correlated with maternal licking/grooming and negatively correlated with nest attendance. Scatterplots are displayed with linear regression lines for each prenatal treatment group. * p < 0.05 interaction between prenatal treatment and postnatal maternal care

2.8.1 Factor module enrichment for estrogen-responsive genes and genes that interact with estrogen receptors
For Factor 4, there were only 14 genes that overlapped between the top genes in the tag-seq and oxRRBS datasets. For the top genes in the tag-seq dataset (467 genes analyzed) there was a significant enrichment of ESR1 found in the list of

446 Transcription Factor Protein-Protein Interactions (FDR < 0.001). For the top genes in the
447 oxRRBS dataset (423 genes analyzed) there was a significant enrichment of ESR1 found in
448 the list of ENCODE and ChEA Consensus TFs from ChIP-X (FDR = 0.031). There was also a
449 nominal enrichment of ESR1 found in the list of Transcription Factor Protein-Protein
450 Interactions (p = 0.0178), but it did not pass FDR correction (FDR > 0.10).
451
452 **3. Discussion**
453
In this study, we examined the interacting and possibly mitigating impacts of postnatal
maternal licking/grooming in prenatal bisphenol-exposed pups on the transcriptome and
the DNA methylome in the developing MPOA. We also investigated the role of estrogen
receptor signaling changes as a potential biological mechanism underlying these
interactions. Overall, we found that postnatal maternal licking/grooming has a substantial
influence on gene expression and DNA methylation changes, particularly in female pups.
This might be driven by the mitigating effect of maternal licking/grooming on *Esrrg*
expression rather than *Esr1*. Given that ESRRG has a high affinity for bisphenols (86,87) and
can facilitate gene expression by binding to estrogen responsive elements or its own
specific binding motif, these results suggest a novel biological mechanism in which
postnatal maternal care can mitigate the negative neurodevelopmental impacts of prenatal
bisphenol exposure. Moreover, these results suggest that postnatal licking-like tactile
stimulation may be an effective intervention against the adverse neurodevelopmental
consequences of prenatal bisphenol exposure.
468
We hypothesized that the interactions between prenatal bisphenol exposure and postnatal
maternal licking/grooming would primarily involve changes in *Esr1* expression in the MPOA
and that the effects would be stronger in female pups than male pups. However, we found
sex-specific interactions in *Esrrg* expression that appear to be associated with multiple co-
expressed gene modules in the WGCNA analysis. While it is known that BPA exposure can
stimulate expression of *Esrrg* in a sex-specific manner (25,88) and has a high binding
affinity to ESRRG (86,87), less is known about the effects of postnatal maternal care on
Esrrg expression or signaling changes. A prior study examining prenatal BPA exposure and
postnatal maternal care conducted in our lab found brain region and sex-specific effects of
maternal care on *Esrrg* expression, where there was an increase in expression in the
prefrontal cortex in female offspring only and in the hypothalamus in male and female
offspring (25). There were no interactions between prenatal BPA and postnatal maternal
care on *Esrrg* expression noted in the study. In our current study, we found postnatal
maternal licking/grooming decreased and normalized *Esrrg* expression in the female pups
with prenatal 150 µg/kg Mixed BP exposure. The discrepancies may be primarily due to the
different ages assessed (weanlings vs. PND10 pups), the bisphenol doses used, and the
bisphenol mixture used in our study may have a different effect than BPA alone. While the
sex-specific interactions in gene expression were consistent in our hypothesis, we also
found that postnatal maternal licking/grooming has a substantial influence on DNA
methylation changes in prenatal bisphenol-exposed male offspring. We also found in prior
studies that postnatal maternal care can interact and mitigate the impacts of prenatal

bisphenol exposure on later-life phenotypes and gene expression in the prefrontal cortex
and amygdala at PND10 in male offspring (34). Therefore, it is possible that postnatal
maternal licking/grooming can mitigate the impacts of prenatal bisphenol exposure in male
offspring, but the underlying biological mechanisms are currently unclear and/or are not
strongly related to estrogen receptor signaling. As an example, previous work using another
estrogenic endocrine disrupting chemical (Aroclor 1221) found estradiol levels predicted
behavioral phenotypes in prenatal Aroclor 1221-exposed females while dopamine-related
gene expression predicted behavioral phenotypes in prenatal Aroclor 1221-exposed male
offspring (89). These sex-specific impacts are worth exploring in future studies.
499
Our WGCNA analysis indicated that including postnatal nest attendance in our statistical
models was required to unveil the significant interactions between the 150 µg/kg Mixed BP
exposure group and maternal licking/grooming but was not significantly associated with
these modules by itself. Postnatal nest attendance includes licking/grooming and other
forms of pup contact, including thermotactile contact. This suggests that postnatal
licking/grooming is the primary driver of these interactions and overall pup contact plays a
minor but possibly important role (47) in gene expression changes for prenatal bisphenol-
exposed female pups. Directly manipulating licking-like tactile stimulation while keeping
other forms of pup contact consistent would confirm these findings and is an ongoing area
of investigation in our lab.
510
Our DMR analysis indicates postnatal maternal licking/grooming has a substantial
influence on DNA methylation changes in prenatal bisphenol-exposed pups. The finding
that more DMRs were added than lost when including postnatal maternal licking/grooming
as a covariate suggests that postnatal maternal licking/grooming plays more of a
moderating role in shaping the DNA methylome in prenatal bisphenol-exposed pups,
consistent with our prior findings (34,39). However, it was apparent that the differential
methylation findings did not map consistently onto differential gene expression. While it is
assumed that changes in DNA methylation and gene expression would be strongly linked,
there are several reasons why there may be a discordance. First, we did not implement any
substantial environmental challenges to the pups immediately prior to brain collection.
While DNA methylation changes are relatively stable, gene expression changes are
dependent on both the availability of chromatin to allow transcription and transcriptional
activity itself. Therefore, the associations between DNA methylation and gene expression
changes may be stimulus-dependent. The second reason is that there are inherent
differences in the approach in which these changes are assessed. While we assessed the
whole transcriptome, we only probed a small portion of CpG sites in the genome. In
addition, many of these CpG sites are intergenic and we have an incomplete understanding
of the role of DNA methylation in distal CpG sites on gene expression. Finally, in any
particular cell, DNA methylation is mainly binary (methylated or unmethylated with rare
instances of hemi-methylation) while gene expression changes can have a large range. This
could complicate analyses when using bulk tissue and would need to be resolved at the
single-cell level. Nevertheless, when we integrated the DNA methylation and gene
expression data using MOFA, we generally replicate the gene expression findings. This

confirms that the significant interactions between the 150 μg/kg Mixed BP exposure group
and maternal licking/grooming in female pups apply to both gene expression and DNA
methylation.
537
We explored potential transcription factor binding sites associated with DNA methylation
changes with prenatal bisphenol exposure and postnatal maternal licking/grooming. To do
so, we adapted the SArKS method to RRBS data so we can correlate DNA methylation
changes in prenatal bisphenol-exposed pups with and without postnatal maternal
licking/grooming as a covariate to motifs and possible transcription factor binding sites. As
a comparison, we also used MEME Suite to identify motifs based on the changes in DMRs
with and without postnatal maternal licking/grooming as a covariate. The main difference
with using a correlative-based method is that it accounts for the continuous range of DNA
methylation changes (0-100%) instead of a binary representation (i.e., it is differentially
methylated or not differentially methylated). A correlative-based method can be
particularly advantageous with more complex study designs where multiple independent
variables are assessed. When a binary representation is applied to DMR data with and
without covariates, there is an assumption that all DMRs that change will also have the
same effect size changes and would be equally important for motif discovery. In contrast,
accounting for the continuous range of differential methylation and effect size changes
with covariates can provide a more accurate representation of the potential transcription
factors involved with differential methylation at specific loci. With the SArKS method, we
found numerous motifs and transcription factor binding site predictions that change when
including postnatal maternal licking/grooming as a covariate, as we would have expected
based on the DMR analysis. This was also the case with the MEME Suite method and the
changes were sometimes even more pronounced, possibly due to the binary
representation of the input sequences as described above.
560
A consistent theme throughout our analyses was that the genes and transcription factor
binding site predictions associated with prenatal bisphenol exposure and postnatal
maternal licking/grooming as well as their interactions were either related to estrogen
receptor signaling or were estrogen-responsive genes. In prior transcriptome analyses in
the PND10 prefrontal cortex and amygdala, we found consistently strong and significant
enrichment of ESR1 in the Transcription Factor Protein-Protein Interactions module in
Enrichr (34). This was also the case in the PND10 MPOA, but we also found significant
(though weaker) enrichment of ESR1 in the ENCODE and ChEA Consensus TFs from ChIP-X
module. These findings are consistent with our hypothesis that the interactions between
prenatal bisphenol exposure and postnatal maternal licking/grooming converge on
estrogen receptor signaling, with some brain region-specific differences, and that we are
primarily finding secondary effects of estrogen receptor signaling changes by PND10.
573
It is worth noting that the mitigating effects of maternal licking/grooming on ESRRG
signaling are speculative since they are derived from RNA-seq data. Future work will need
to test these impacts more directly, such as with chromatin immunoprecipitation or
luciferase reporter assays for ESRRG. One other notable limitation of this study was while

we found consistent interactions between postnatal licking/grooming and the 150 μg/kg
Mixed BP dose, this dosage is not considered environmentally relevant and is generally
higher than human exposure levels. However, bisphenol exposure follows a non-monotonic
dose response curve (90,91), meaning that very high levels and very low levels will exert
biological effects. In our study, the 50 μg/kg Mixed BP and BPA-only dose did not facilitate
any increases of *Esrrg* expression in female pups; however, *Esrrg* expression may change
with doses lower than 50 μg/kg, which would be environmentally relevant. Consistent with
this logic, our lab's previous work on prenatal BPA exposure found that the 2 μg/kg and 20
μg/kg doses alter *Esrrg* expression in the prefrontal cortex and hypothalamus (25). While
the dosages used in the current study were informative for discovering potential biological
mechanisms, additional studies using lower doses are needed to ensure the translational
potential of our current findings. Finally, we do not have any data on how maternal
licking/grooming might mitigate the impacts of prenatal bisphenol exposure on MPOA-
dependent behaviors later in life, such as social behavior and maternal care provisioning to
the next generation. However, given that there is already evidence that both prenatal
bisphenol exposure and postnatal maternal care can impact these later-life phenotypes
(25,92–98), it is likely that interactions between these two early-life environmental
exposures exist.
596
Overall, we found that postnatal maternal licking/grooming can mitigate the impacts of
prenatal bisphenol exposure on *Esrrg* expression and co-expressed gene modules in the
developing MPOA in a dose-dependent and sex-specific manner. This appears to
correspond to changes in DNA methylation, which can persist beyond both transient
environmental exposures and impact neurodevelopmental trajectories and behavioral
phenotype. Whether licking-like tactile stimulation can be implemented as an effective
intervention against the adverse neurodevelopmental impacts of prenatal bisphenol
exposure in female offspring remains an open question but will be important to examine in
the future. Exploring this issue will allow us to infer causality and confirm that prenatal
bisphenol exposure and postnatal maternal licking/grooming are acting on the same
biological pathways in opposing directions. This approach will be essential to identifying
potential avenues, such as postnatal touch stimulation, to mitigate the effects of prenatal
bisphenol exposure and improve health and well-being in human populations.
610
611   **4. Materials and Methods**
612
613   **Supplementary Table S12** displays all the code files and data files used for the
614   bioinformatic analyses, as well as the information from all samples used for analyses.
615
4.1 Husbandry and breeding
All animal procedures were approved by the Institutional Animal Care and Use Committee
at the University of Texas at Austin and conformed to the guidelines of the American
Association for Laboratory Animal Science. The rat pups used for the current study were
part of a larger cohort that has been described in previous publications (34,39). Briefly,
breeder female and male Long-Evans rats (Charles River) were housed in same-sex pairs

on a 12:12 hour inverse light-dark cycle with ad libitum access to standard chow diet
(#5LL2, Lab Diet) and water. To limit external exposure to bisphenols and other
xenoestrogens, all rats were provided glass water bottles, polysulfone cages, and aspen
wood shavings for bedding material.
626

4.2 Gestational bisphenol administration
Male and female rats were screened for breeding receptivity by pairing one male with one
or two females and observing mounting behavior for the male and lordosis for the
female(s). From gestational day 8 until parturition, cage-pairs of dams were randomly
assigned to receive control Corn Oil, 50 µg/kg BPA, 50 µg/kg Mixed Bisphenols (BP), or 150
µg/kg Mixed BP. This time period includes the start of brain sexual differentiation and when
the expression of estrogen receptors is apparent in the fetal brain (99). Stock solutions of
bisphenol A (BPA; #B0494, TCI, ≥ 99 %), bisphenol S (BPS; #A17342, Alpha Aesar, ≥ 99 %),
or bisphenol F (BPF; #A11471, Alfa Aesar, ≥ 98 %) were dissolved in corn oil (#405435000,
Acros Organics). Equal parts BPA, BPS, and BPF were used for both Mixed BP treatment
groups. Experimenters involved with the administration of treatments were blinded to the
treatment groups. After parturition (postnatal day 0; PND0), offspring were sex-determined
using relative anogenital distance and culled to six male and six female pups per litter.
Following the maternal care video recordings at PND10, the brains from one male pup and
one female pup per litter were collected and flash-frozen in hexanes on dry ice. All brains
were stored at -80°C until cryosectioning.
643

4.3 Postnatal maternal care observations and quantification
Home cage maternal behavior was recorded for one hour per day starting one hour after
lights-off from PND1-10 using Raspberry Pi 3B+ minicomputers. Maternal behavior was
scored by either manual coding or through the AMBER pipeline (100). All postnatal
maternal care measures were normalized to seconds of observed behavior per day and
then scaled and centered where the average is 0 and standard deviation is 1 for all
downstream bioinformatic analyses. Based on prior analyses (34,39) we used
licking/grooming and nest attendance from PND1-5 as independent variables or covariates
for the current study. Maternal behavior measures used in the current study did not vary
between prenatal bisphenol exposure groups (34).
654

4.4 RNA extraction and 3' tag sequencing (Tag-seq)
Power analyses were done using the R package PROPER for RNA-seq data with the Gilad
dataset (medium biological variation) and 25 data simulations to calculate effect size.
Analyzing 6 biological samples per group allowed 72% power to correctly identify
differentially expressed genes with an FDR value < 0.05 and log fold change of at least 0.5.
PND10 brains (n = 6 per sex per prenatal treatment group) were cryosectioned in 50 µm
slices using a ThermoFisher Scientific CryoStar NX50 cryostat or a Leica CM3050S
cryostat. The medial preoptic area (MPOA; 0.20 to -0.60 mm Bregma) was microdissected
using an atlas for the developing rat brain (101) and a supplementary atlas for the PND10
rat brain (102). RNA was extracted using the MagMAX FFPE DNA/RNA Ultra Kit (#A31881,
ThermoFisher Scientific). For unfixed frozen tissue, 210 µl of protease solution was

pipetted onto each MPOA sample and allowed to incubate at 55°C at 900 rpm for at least
one hour. The homogenate was immediately processed for RNA and DNA extraction using
the Kingfisher Flex System following the manufacturer's instructions. RNA quantity was
assessed with the Quant-iT RNA Assay kit (#Q33140, ThermoFisher Scientific) and RNA
quality was assessed using the RNA 6000 Pico Assay kit (#5067-1513, Agilent
Technologies). All RNA samples were diluted to 20 ng/µl before submission for library
preparation and tag-seq (103,104) at the Genome Sequence and Analysis Facility at the
University of Texas at Austin. Reads were sequenced on the NovaSeq S1 (100 bp single-end
reads).
675
4.5 DNA extraction and oxidative reduced representation bisulfite sequencing
(oxRRBS)
DNA was extracted using the MagMAX FFPE DNA/RNA Ultra Kit (#A31881, ThermoFisher
Scientific). A subset of samples (two 150 µg/kg Mixed BP females, one 150 µg/kg Mixed BP
male) did not contain sufficient DNA for library preparation for oxRRBS. Another subset of
samples (one 50 µg/kg BPA male, one 150 µg/kg Mixed BP female, one Corn Oil female)
likely had substantial anterior pituitary gland contamination based on the tag-seq results
(described in the tag-seq preprocessing section) and were not processed for oxRRBS. To
maintain a sample size of 6 per sex per prenatal treatment group, DNA was extracted from
additional MPOA samples using the MagMAX DNA Multi-Sample Ultra 2.0 Kit following the
manufacturer's instructions for tissue samples (#A45721, ThermoFisher Scientific). DNA
quantity was assessed with the Quantifluor dsDNA system (#E2670, Promega).
688
Library preparation for oxRRBS was done using the Ovation RRBS Methyl-Seq System with
the TrueMethyl® oxBS module (Tecan) using 55 ng of input DNA. Between 12-20 samples
were processed per batch of library preparation with a total of 5 batches, balanced by
prenatal treatment group and sex. The oxidation of DNA prior to bisulfite conversion
converts 5hmC to 5fC and allows for discrimination between 5mC and 5hmC at single
base-pair resolution. For this study, oxidative bisulfite conversion was used to remove
5hmC and detect only 5mC in the samples. Final libraries were quality-checked using the
Agilent Technologies High Sensitivity DNA Kit. Five libraries (one 50 µg/kg BPA female, one
50 µg/kg Mixed BP male, one 50 µg/kg Mixed BP female, and two 150 µg/kg Mixed BP
females) were removed due to insufficient final yield. The remaining libraries (n = 4-6 per
sex per prenatal treatment group) were pooled with 8-9 libraries per pool (balanced by
prenatal treatment group and sex) and submitted for an additional Ampure XP bead clean-
up and sequencing to the Genome Sequence and Analysis Facility at the University of Texas
at Austin. Reads were sequenced on the NovaSeq SP (150 bp paired-end reads).
703
4.6 Pipeline and Analyses: Tag-seq
4.6.1 Preprocessing The tag-seq pipeline overview can be found in **Fig 1** and all data and
code for the differential gene expression and weighted gene co-expression network
analysis can be found at https://github.com/SLauby/bisphenols_maternalcare_epigenome.
The analysis pipeline followed previously published tag-seq analyses (34). Adaptor
sequences, poly-A tails, and low-quality reads were trimmed (https://github.com/z0on/tag-

based_RNAseq) followed by cutadapt (version 1.18). Trimmed reads were mapped to the
rat genome (mRatBN7.2; RefSeq) using STAR (version 2.7.3a) with the default settings and
sorted using samtools (version 1.10). About 90% of reads were uniquely mapped to the rat
genome for all samples. Counts for each gene were calculated for each sample using the
bedtools (version 2.27.1) multicov function. Three samples (one 50 µg/kg BPA male, one
150 µg/kg Mixed BP female, one Corn Oil female) had unusually high counts (over 100x of
the other samples) for anterior pituitary gland-specific genes, including *Pomc, Gnrhr,* and
Prl. These samples were removed from the dataset prior to all downstream analyses.
718
4.6.2 Differential gene expression analysis For all downstream tag-seq analyses,
comparisons between prenatal treatment groups and interactions with prenatal treatment
group and postnatal maternal care were always done relative to the Corn Oil group. To
examine differential expression of individual genes, including all estrogen and estrogen-
related receptors, gene count data was normalized and analyzed for differential expression
using the DESeq2 R package (version 1.34.0), separated by sex. Genes with more than 5
counts for more than 6 samples were analyzed. Prenatal treatment and the postnatal
maternal care measures (licking/grooming, nest attendance) from PND1-5 were used as
independent variables. The estrogen and estrogen-related receptors (*Esr1, Esr2, Esrra,*
Esrrb, Esrrg) were specifically examined for significant interactions between each prenatal
treatment group and licking/grooming. Other potential differentially expressed genes
(DEGs) were identified with a false discovery rate (FDR) alpha $\leq$ 0.10 and $\leq$ 0.05 with
Benjamani-Hochburg correction. All effects were reported as statistically significant if p $\leq$
0.05 and marginally significant if p $\leq$ 0.10.
733
4.6.3 Weighted gene co-expression network analysis (WGCNA) To examine differential
expression of co-expressing gene profiles, gene networks were created with the WGCNA R
package (version 1.72-1) using normalized and log-transformed gene count data, split by
sex. Genes with very low counts and variation were filtered out using the
goodSamplesGenes function in R (verbosity = 3). WGCNA calculates Pearson correlations
of count data between every gene for network construction and clusters highly correlated
genes into eigengene modules (at least 30 genes per module). A soft power threshold of 8
was used for network construction. We excluded modules with eigengene values that were
characterized by one extreme outlier (> 0.90) and low variation following outlier removal
(standard deviation < 0.10) from further analyses. The eigengene values from the remaining
modules were analyzed using the lm function in R. Prenatal treatment and the postnatal
maternal care measures (licking/grooming, nest attendance) from PND1-5 were used as
independent variables. For any eigengene module that showed statistically significant
interactions, the gene sets were extracted and analyzed using Enrichr (105,106) using the
full gene list used for network construction as the background
(https://maayanlab.cloud/Enrichr/). To examine if the gene sets were enriched in estrogen-
responsive genes, ESR1 was examined in the ENCODE and ChEA Consensus TFs from
ChIP-X module. To examine if the gene sets were enriched in genes that are known to
interact with estrogen receptors, ESR1 was examined in the Transcription Factor Protein-
Protein Interaction module. To examine the degree of connectivity with the co-expressed

gene modules with specific estrogen receptors, the kME was calculated for all estrogen
and estrogen-related receptors in the WGCNA package. The kME determines how closely a
gene of interest is related to a particular module and is represented as a positive or
negative percentage of all genes that have highly correlated expression to the gene of
interest. Negative values indicate inverse relationships. All effects were reported as
statistically significant if $p \leq 0.05$ and marginally significant if $p \leq 0.10$.
760
4.7 Pipeline and Analyses: oxRRBS
<u>**4.7.1 Preprocessing**</u> The oxRRBS pipeline overview can be found in **Fig 1** and all code for
the differentially methylated region analysis can be found at
https://github.com/SLauby/bisphenols_maternalcare_epigenome. The bismark-processed
DNA methylation dataset can be accessed in Zenodo
(https://doi.org/10.5281/zenodo.17273659). Adaptor sequences and low-quality reads (q <
30) were trimmed using trim galore (version 0.6.10) and cutadapt (version 4.6). Diversity
sequences and reads without the MspI site (CGG) were removed using a custom python
script provided by the manufacturer of the oxRRBS kit
(https://github.com/nugentechnologies/NuMetRRBS). Paired-end reads were mapped to
the rat genome (mRatBN7.2; RefSeq) using bismark (version 0.22.3) and bowtie2 (version
2.5.2) with the default settings. About 75% of reads were uniquely mapped to the rat
genome for all samples. PCR duplicates were removed based on the genomic coordinates
of the reads and a 6-bp unique molecular identifier found in the read index using a custom
python script provided by the manufacturer of the oxRRBS kit
(https://github.com/tecangenomics/nudup). An average of about 8.5 million uniquely
aligned, de-duplicated reads per sample were analyzed for DNA methylation levels using
the bismark methylation extractor. Bisulfite conversion rate was > 99% for all samples.
779
<u>**4.7.2 Differentially methylated region (DMR) analysis**</u> For all downstream oxRRBS
analyses, pairwise comparisons between prenatal treatment groups were always done
relative to the Corn Oil group. Differentially methylated regions (500 bp) between prenatal
treatment groups were examined using Methylkit (version 1.26.0), separated by sex.
Regions were included in the analysis if there were over 10x reads for at least 4 samples per
prenatal treatment group. Batch of library preparation was used as a covariate in all
analyses. DMRs were identified with an FDR q-value ≤ 0.05 using the SLIM method and had
> 5% overall methylation difference between groups. DMR analyses are currently limited to
pairwise comparisons, so an alternative approach was undertaken to examine potential
interactions between prenatal bisphenol exposure and postnatal maternal
licking/grooming. Each comparison between the prenatal treatments groups and Corn Oil
group was done including and excluding postnatal maternal licking/grooming as a covariate
and the degree of overlap in DMRs was identified. DMRs that were lost or added when
including postnatal maternal licking/grooming as a covariate in the analysis were
considered to be influenced by postnatal maternal licking/grooming. More specifically,
DMRs that were added were assumed to be due to interactions between prenatal
bisphenol exposure and postnatal maternal licking/grooming.
797

4.7.3 Differentially methylated motif and transcription factor binding site analysis:
Suffix Array Kernel Smoothing (SArKS) To adapt the SArKS method to RRBS data, genomic
regions with methylation data were created as input sequences for the SArKS algorithm to
identify overrepresented sequence motifs. The code for creating those input sequences
can be found at https://github.com/denniscwylie/prenatal_bp_x_lg_epigenome_analysis.
For each individual sample, observed methylation sites were partitioned into disjointed
genomic ranges separated by at least 51 base pairs. The bedtools merge command was
then applied (using the bedtoolsr::bt.merge function in a custom R script) to combine the
resulting partitioned range sets from each individual sample into a single set of partitioned
genomic ranges in which every individual sample range is contained within one of the
merged range intervals. The merged set of genomic ranges were then trimmed using the
bedtools coverage command together with a custom R script, only retaining positions
which were covered by one of the individual sample-partitioned ranges for at least 25 (out
of 43 total) of the individual samples analyzed. For each sample, a local methylation rate
was assigned to each genomic position within the merged set of (trimmed) genomic ranges
defined above. Each individual base position within each of these partitioned regions is
assigned a local methylation rate equal to the estimated methylation percentage for the
nearest site observed methylation site (within 50 bp). If the nearest site observed
methylation site is > 50 bp, the local methylation rate is considered undefined for the
sample. Finally, a single average methylation level for each sample was then computed for
each of the merged (and trimmed) genomic ranges as the mean of the assigned individual
base methylation rates over all positions within the range (omitting any positions with
undefined local methylation rates).
821
For each genomic range R in the merged (and trimmed) set, a linear model was
incorporated to model the arcsin-transformed range-averaged methylation rates as the
dependent variable and the library preparation batch covariate and prenatal treatment
group as independent predictor variables. Each comparison was done including and
excluding postnatal maternal licking/grooming as a covariate. In this way, an estimated
model coefficient was obtained for each combination of genomic range R and prenatal
bisphenol treatment group to the Corn Oil group with and without postnatal maternal
licking/grooming as a covariate. For each prenatal bisphenol treatment group comparison,
SArKS (https://github.com/denniscwylie/sarks; (62)) was used to identify overrepresented
sequence motifs found within the various merged genomic ranges whose presence within
genomic range R correlates with more extreme (either higher or lower, depending on which
direction the analysis was run) estimated model coefficients for each prenatal treatment
group comparison associated with the region R. Step-by-step instructions and pseudocode
for SArKS can be found at
https://bioconductor.posit.co/packages/3.23/bioc/vignettes/sarks/inst/doc/sarks-
vignette.pdf.
838
4.7.4 Differentially methylated motif and transcription factor binding site analysis:
MEME Suite DMRs that were found using Methylkit were used to identify overrepresented
sequence motifs with MEME Suite (Version 5.5.7; (107)) using previously published

parameters (108) with some modifications. To make the analysis more comparable to
SArKS, we included regions that had differential methylation with a nominal p-value ≤ 0.05
and had > 0% overall methylation difference between groups. In addition, MEME was used
in discriminative mode using a set of sequences that were not differentially methylated
between the prenatal treatment groups and Corn Oil group.

For both SArKS and MEME Suite methods, overrepresented sequence motifs were
examined for potential transcription factor binding sites using the JASPAR 2024 CORE (non-
redundant) database for vertebrates (109). E-values ≤ 0.05 were considered statistically
significant.

4.8 Multi-omic factor analysis (MOFA) of tag-seq and oxRRBS datasets
Directly comparing gene expression differences and DNA methylation modifications can be
challenging when the analysis pipeline for each measure is not integrated. To examine the
interactions between prenatal BP exposure and maternal care on the epigenome in a more
holistic manner, a MOFA was performed using the MOFA2 package in R (version 1.10.0),
separated by sex. Multi-omic analyses can only be performed if the tag-seq and oxRRBS
datasets were derived from the same MPOA samples; therefore, the sample sizes were
smaller than either the tag-seq or oxRRBS analyses (n = 3-6 per sex per prenatal treatment
group). Gene count data were normalized with DESeq2. DNA methylome data were
restricted to one 500 bp region per 2000 bp window to reduce the influence of the high
covariance structure of DNA methylation in proximal regions. Proportion of DNA
methylation (p) was then log-transformed, similar to M-values in microarray datasets:

$$y = \text{Log2}\left(\left(\frac{p}{1.025} - p\right) + 0.025\right)$$

The gene count dataset was filtered to only include the top 5,000 most variable genes and
the DNA methylome dataset was filtered to only include the top 10,000 most variable
regions. MOFA2 was performed using the default options, except that the scale views
option was set to TRUE, convergence mode was set to slow, and the number of factors was
reduced to 5. The weights from each sample and factor were extracted and analyzed using
the lm function in R. Prenatal treatment and the postnatal maternal care measures
(licking/grooming, nest attendance) from PND1-5 were used as independent variables. For
any factor that showed statistically significant main effects or interactions, the top genes
and regions by weight (250 positive weights and 250 negative weights) were extracted. The
most proximal gene was determined from each region (if within the promoter or gene body)
from the oxRRBS dataset and examined for any overlap with the top genes from the tag-seq
dataset. The top genes (removing the uncharacterized genes) from the tag-seq and oxRRBS
datasets were separately analyzed with Enrichr. Enrichment for ESR1 was examined in the
list of ENCODE and ChEA Consensus TFs from ChIP-X and Transcription Factor Protein-
Protein Interaction modules. All effects were reported as statistically significant if p ≤ 0.05
and marginally significant if p ≤ 0.10.

5. Acknowledgements

We thank Isha Agarwal, Taylor Hite, and Madeline Severson for their assistance in the
prenatal bisphenol administration.

6. References

1. Appleton AA, Holdsworth EA, Kubzansky LD. A Systematic Review of the Interplay
Between Social Determinants and Environmental Exposures for Early-Life Outcomes.
Curr Environ Health Rep. 2016 Sep;3(3):287–301. doi:10.1007/s40572-016-0099-7

2. Tamiz AP, Koroshetz WJ, Dhruv NT, Jett DA. A focus on the neural exposome. Neuron.
2022 Apr;110(8):1286–9. doi:10.1016/j.neuron.2022.03.019

3. King LS, Guyon-Harris KL, Valadez EA, Radulescu A, Fox NA, Nelson CA, et al. A
Comprehensive Multilevel Analysis of the Bucharest Early Intervention Project: Causal
Effects on Recovery From Early Severe Deprivation. Am J Psychiatry. 2023 Aug
1;180(8):573–83. doi:10.1176/appi.ajp.20220672

4. Zeanah CH, Humphreys KL, Fox NA, Nelson CA. Alternatives for abandoned children:
insights from the Bucharest Early Intervention Project. Curr Opin Psychol. 2017
Jun;15:182–8. doi:10.1016/j.copsyc.2017.02.024

5. Catherine NLA, MacMillan H, Cullen A, Zheng Y, Xie H, Boyle M, et al. Effectiveness of
nurse-home visiting in improving child and maternal outcomes prenatally to age two
years: a randomised controlled trial (British Columbia Healthy Connections Project). J
Child Psychol Psychiatry. 2024 May;65(5):644–55. doi:10.1111/jcpp.13846

6. Olds DL, Robinson J, O'Brien R, Luckey DW, Pettitt LM, Henderson CR, et al. Home
Visiting by Paraprofessionals and by Nurses: A Randomized, Controlled Trial.
Pediatrics. 2002 Sep 1;110(3):486–96. doi:10.1542/peds.110.3.486

7. Akbari E, Binnoon-Erez N, Rodrigues M, Ricci A, Schneider J, Madigan S, et al. Kangaroo
mother care and infant biopsychosocial outcomes in the first year: A meta-analysis.
Early Hum Dev. 2018 Jul;122:22–31. doi:10.1016/j.earlhumdev.2018.05.004

8. Lode-Kolz K, Jonas W, Hetland HB, Hovland Instebø KH, Tokvam H, Pike H, et al.
Immediate Skin-to-Skin Contact at Very Preterm Birth and Neurodevelopment the
First Two Years: Secondary Outcomes from a Randomised Clinical Trial. Children.
2025 Jul 27;12(8):986. doi:10.3390/children12080986

9. Ghassabian A, Trasande L. Disruption in Thyroid Signaling Pathway: A Mechanism for
the Effect of Endocrine-Disrupting Chemicals on Child Neurodevelopment. Front
Endocrinol. 2018 Apr 30;9:204. doi:10.3389/fendo.2018.00204

10. Schug TT, Blawas AM, Gray K, Heindel JJ, Lawler CP. Elucidating the Links Between
Endocrine Disruptors and Neurodevelopment. Endocrinology. 2015 Jun
1;156(6):1941–51. doi:10.1210/en.2014-1734

11. Nesan D, Kurrasch DM. Gestational Exposure to Common Endocrine Disrupting
Chemicals and Their Impact on Neurodevelopment and Behavior. Annu Rev Physiol.
2020 Feb 10;82(1):177–202. doi:10.1146/annurev-physiol-021119-034555

12. Wetherill YB, Akingbemi BT, Kanno J, McLachlan JA, Nadal A, Sonnenschein C, et al. In
vitro molecular mechanisms of bisphenol A action. Reprod Toxicol. 2007;24(2):178–
98. doi:10.1016/j.reprotox.2007.05.010 PubMed PMID: 17628395.

13. Eladak S, Grisin T, Moison D, Guerquin MJ, N'Tumba-Byn T, Pozzi-Gaudin S, et al. A new
chapter in the bisphenol a story: Bisphenol S and bisphenol F are not safe alternatives
to this compound. Fertil Steril. 2015 Jan 1;103(1):11–21.
doi:10.1016/j.fertnstert.2014.11.005 PubMed PMID: 25475787.

14. Gertz J, Reddy TE, Varley KE, Garabedian MJ, Myers RM. Genistein and bisphenol A
exposure cause estrogen receptor 1 to bind thousands of sites in a cell type-specific
manner. Genome Res. 2012;22(11):2153–62. doi:10.1101/gr.135681.111 PubMed
PMID: 23019147.

15. Matthews JB, Twomey K, Zacharewski TR. In Vitro and in Vivo Interactions of Bisphenol
A and Its Metabolite, Bisphenol A Glucuronide, with Estrogen Receptors α and β.
Chem Res Toxicol. 2001 Feb 1;14(2):149–57. doi:10.1021/tx0001833

16. Rochester JR, Bolden AL, Kwiatkowski CF. Prenatal exposure to bisphenol A and
hyperactivity in children: a systematic review and meta-analysis. Environ Int.
2018;114(December 2017):343–56. doi:10.1016/j.envint.2017.12.028 PubMed PMID:
29525285.

17. Braun JM, Yolton K, Dietrich KN, Hornung R, Ye X, Calafat AM, et al. Prenatal bisphenol
A exposure and early childhood behavior. Environ Health Perspect.
2009;117(12):1945–52. doi:10.1289/ehp.0900979 PubMed PMID: 20049216.

18. Roen EL, Wang Y, Calafat AM, Wang S, Margolis A, Herbstman J, et al. Bisphenol A
exposure and behavioral problems among inner city children at 7-9 years of age.
Environ Res. 2015;142:739–45. doi:10.1016/j.envres.2015.01.014 PubMed PMID:
25724466.

19. Perera F, Vishnevetsky J, Herbstman JB, Calafat AM, Xiong W, Rauh V, et al. Prenatal
Bisphenol A Exposure and Child Behavior in an Inner-City Cohort. Environ Health
Perspect. 2012 Aug;120(8):1190–4. doi:10.1289/ehp.1104492

20. Perera F, Nolte ELR, Wang Y, Margolis AE, Calafat AM, Wang S, et al. Bisphenol A
exposure and symptoms of anxiety and depression among inner city children at 10–12

years of age. Environ Res. 2016;151:195–202. doi:10.1016/j.envres.2016.07.028
PubMed PMID: 27497082.

21. Braun JM, Muckle G, Arbuckle T, Bouchard MF, Fraser WD, Ouellet E, et al. Associations
of Prenatal Urinary Bisphenol A Concentrations with Child Behaviors and Cognitive
Abilities. Environ Health Perspect. 2017 Jun 23;125(6):067008. doi:10.1289/EHP984

22. Braun JM, Kalkbrenner AE, Calafat AM, Yolton K, Ye X, Dietrich KN, et al. Impact of Early-
Life Bisphenol A Exposure on Behavior and Executive Function in Children. Pediatrics.
2011 Nov 1;128(5):873–82. doi:10.1542/peds.2011-1335

23. Castro B, Sánchez P, Torres JM, Ortega E. Bisphenol A, bisphenol F and bisphenol S
affect differently 5α-reductase expression and dopamine-serotonin systems in the
prefrontal cortex of juvenile female rats. Environ Res. 2015 Oct 1;142:281–7.
doi:10.1016/j.envres.2015.07.001 PubMed PMID: 26186136.

24. Fan AM, Chou WC, Lin P. Bisphenol A—toxicity and risk assessment update with
academic and regulatory perspectives and physiologically based pharmacokinetic
modeling. In: Reproductive and Developmental Toxicology [Internet]. Elsevier; 2022. p.
779–801. Available from: http://dx.doi.org/10.1016/B978-0-323-89773-0.00039-4
doi:10.1016/B978-0-323-89773-0.00039-4

25. Kundakovic M, Gudsnuk K, Franks B, Madrid J, Miller RL, Perera FP, et al. Sex-specific
epigenetic disruption and behavioral changes following low-dose in utero bisphenol A
exposure. Proc Natl Acad Sci. 2013 Jun 11;110(24):9956–61.
doi:10.1073/pnas.1214056110

26. Arambula SE, Jima D, Patisaul HB. Prenatal bisphenol A (BPA) exposure alters the
transcriptome of the neonate rat amygdala in a sex-specific manner: a CLARITY-BPA
consortium study. NeuroToxicology. 2018 Mar 1;65:207–20.
doi:10.1016/j.neuro.2017.10.005 PubMed PMID: 29097150.

27. Cheong A, Johnson SA, Howald EC, Ellersieck MR, Camacho L, Lewis SM, et al. Gene
expression and DNA methylation changes in the hypothalamus and hippocampus of
adult rats developmentally exposed to bisphenol A or ethinyl estradiol: a CLARITY-BPA
consortium study. Epigenetics. 2018;13(7):704–20.
doi:10.1080/15592294.2018.1497388 PubMed PMID: 30001178.

28. Witchey SK, Fuchs J, Patisaul HB. Perinatal bisphenol A (BPA) exposure alters brain
oxytocin receptor (OTR) expression in a sex- and region- specific manner: A CLARITY-
BPA consortium follow-up study. NeuroToxicology. 2019 Sep;74:139–48.
doi:10.1016/j.neuro.2019.06.007

29. Catanese MC, Vandenberg LN. Bisphenol S (BPS) alters maternal behavior and brain in
mice exposed during pregnancy/lactation and their daughters. Endocrinology.
2017;158(3):516–30. doi:10.1210/en.2016-1723 PubMed PMID: 28005399.

30. LaPlante CD, Catanese MC, Bansal R, Vandenberg LN. Bisphenol S Alters the Lactating
Mammary Gland and Nursing Behaviors in Mice Exposed During Pregnancy and
Lactation. Endocrinology. 2017 Oct 1;158(10):3448–61. doi:10.1210/en.2017-00437

31. Seta DD, Minder I, Dessì-Fulgheri F, Farabollini F. Bisphenol-A exposure during
pregnancy and lactation affects maternal behavior in rats. Brain Res Bull.
2005;65(3):255–60. doi:10.1016/j.brainresbull.2004.11.017 PubMed PMID: 15811589.

32. Bonaldo B, Gioiosa L, Panzica G, Marraudino M. Exposure to either Bisphenol A or S
Represents a Risk for Crucial Behaviors for Pup Survival, Such as Spontaneous
Maternal Behavior in Mice. Neuroendocrinology. 2023;113(12):1283–97.
doi:10.1159/000526074

33. Lapp HE, Margolis AE, Champagne FA. Impact of a bisphenol A, F, and S mixture and
maternal care on the brain transcriptome of rat dams and pups. NeuroToxicology.
2022 Dec;93:22–36. doi:10.1016/j.neuro.2022.08.014

34. Lauby SC, Lapp HE, Salazar M, Semyrenko S, Chauhan D, Margolis AE, et al. Postnatal
maternal care moderates the effects of prenatal bisphenol exposure on offspring
neurodevelopmental, behavioral, and transcriptomic outcomes. Torrens C, editor.
PLOS ONE. 2024 Jun 11;19(6):e0305256. doi:10.1371/journal.pone.0305256

35. Hane AA, Henderson HA, Reeb-Sutherland BC, Fox NA. Ordinary variations in human
maternal caregiving in infancy and biobehavioral development in early childhood: A
follow-up study. Dev Psychobiol. 2010 Sep;52(6):558–67. doi:10.1002/dev.20461

36. Narvaez D, Gleason T, Wang L, Brooks J, Lefever JB, Cheng Y. The evolved development
niche: Longitudinal effects of caregiving practices on early childhood psychosocial
development. Early Child Res Q. 2013 Oct;28(4):759–73.
doi:10.1016/j.ecresq.2013.07.003

37. Champagne FA. Epigenetic mechanisms and the transgenerational effects of maternal
care. Front Neuroendocrinol. 2008. doi:10.1016/j.yfrne.2008.03.003

38. Curley JP, Champagne FA. Influence of maternal care on the developing brain:
Mechanisms, temporal dynamics and sensitive periods. Front Neuroendocrinol. 2016.
doi:10.1016/j.yfrne.2015.11.001

39. Lauby SC, Agarwal I, Lapp HE, Salazar M, Semyrenko S, Chauhan D, et al. Interplay
between prenatal bisphenol exposure, postnatal maternal care, and offspring sex in
predicting DNA methylation relevant to anxiety-like behavior in rats. Horm Behav. 2025
Jun;172:105745. doi:10.1016/j.yhbeh.2025.105745

40. Jorgensen EM, Alderman MH, Taylor HS. Preferential epigenetic programming of
estrogen response after *in utero* xenoestrogen (bisphenol-A) exposure. FASEB J. 2016
Sep 16;30(9):3194–201. doi:10.1096/fj.201500089R

41. Yaoi T, Itoh K, Nakamura K, Ogi H, Fujiwara Y, Fushiki S. Genome-wide analysis of
epigenomic alterations in fetal mouse forebrain after exposure to low doses of
bisphenol A. Biochem Biophys Res Commun. 2008;376(3):563–7.
doi:10.1016/j.bbrc.2008.09.028 PubMed PMID: 18804091.

42. McGowan PO, Suderman M, Sasaki A, Huang TCT, Hallett M, Meaney MJ, et al. Broad
epigenetic signature of maternal care in the brain of adult rats. PLoS ONE. 2011.
doi:10.1371/journal.pone.0014739

43. Weaver ICG, Cervoni N, Champagne F a, D'Alessio AC, Sharma S, Seckl JR, et al.
Epigenetic programming by maternal behavior. Nat Neurosci. 2004;7(8):847–54.
doi:10.1038/nn1276 PubMed PMID: 15220929.

44. Pena CJ, Neugut YD, Champagne FA. Developmental timing of the effects of maternal
care on gene expression and epigenetic regulation of hormone receptor levels in
female rats. Endocrinology. 2013;154(11):4340–51. doi:10.1210/en.2013-1595
PubMed PMID: 24002038.

45. Champagne FA, Weaver ICG, Diorio J, Dymov S, Szyf M, Meaney MJ. Maternal care
associated with methylation of the estrogen receptor-α1b promoter and estrogen
receptor-α expression in the medial preoptic area of female offspring. Endocrinology.
2006. doi:10.1210/en.2005-1119

46. Seebacher F, Little AG. Thyroid hormone links environmental signals to DNA
methylation. Philos Trans R Soc B Biol Sci. 2024 Mar 25;379(1898):20220506.
doi:10.1098/rstb.2022.0506

47. Lauby SC, McGowan PO. Early life variations in temperature exposure affect the
epigenetic regulation of the paraventricular nucleus in female rat pups. Proc R Soc B
Biol Sci. 2020 Oct 28;287(1937):20201991. doi:10.1098/rspb.2020.1991 PubMed
PMID: 33109014.

48. Gray JD, Kogan JF, Marrocco J, McEwen BS. Genomic and epigenomic mechanisms of
glucocorticoids in the brain. Nat Rev Endocrinol. 2017 Nov;13(11):661–73.
doi:10.1038/nrendo.2017.97

49. Sugrue VJ, Prescott M, Glendining KA, Bond DM, Horvath S, Anderson GM, et al. The
androgen clock is an epigenetic predictor of long-term male hormone exposure. Proc
Natl Acad Sci. 2025 Jan 21;122(3):e2420087121. doi:10.1073/pnas.2420087121

50. Crudo A, Petropoulos S, Suderman M, Moisiadis VG, Kostaki A, Hallett M, et al. Effects
of antenatal synthetic glucocorticoid on glucocorticoid receptor binding, DNA

methylation, and genome-wide mRNA levels in the fetal male hippocampus.
Endocrinology. 2013. doi:10.1210/en.2013-1484 PubMed PMID: 24029241.

51. Ariazi EA, Taylor JC, Black MA, Nicolas E, Slifker MJ, Azzam DJ, et al. A new role for ERα:
Silencing via DNA methylation of basal, stem cell, and EMT genes. Mol Cancer Res.
2017;15(2):152–64. doi:10.1158/1541-7786.MCR-16-0283 PubMed PMID: 28108626.

52. Kovács T, Szabó-Meleg E, Ábrahám IM. Estradiol-induced epigenetically mediated
mechanisms and regulation of gene expression. Int J Mol Sci. 2020;21(9).
doi:10.3390/ijms21093177 PubMed PMID: 32365920.

53. Ung M, Ma X, Johnson KC, Christensen BC, Cheng C. Effect of estrogen receptor α
binding on functional DNA methylation in breast cancer. Epigenetics. 2014;9(4):523–
32. doi:10.4161/epi.27688 PubMed PMID: 24434785.

54. Dumasia K, Kumar A, Deshpande S, Balasinor NH. Estrogen signaling, through estrogen
receptor β, regulates DNA methylation and its machinery in male germ line in adult
rats. Epigenetics. 2017 Jun 3;12(6):476–83. doi:10.1080/15592294.2017.1309489

55. Monje L, Varayoud J, Luque EH, Ramos JG. Neonatal exposure to bisphenol A modifies
the abundance of estrogen receptor α transcripts with alternative 5'-untranslated
regions in the female rat preoptic area. J Endocrinol. 2007;194(1):201–12.
doi:10.1677/JOE-07-0014 PubMed PMID: 17592034.

56. Cao J, Mickens JA, McCaffrey KA, Leyrer SM, Patisaul HB. Neonatal Bisphenol A
exposure alters sexually dimorphic gene expression in the postnatal rat
hypothalamus. NeuroToxicology. 2012;33(1):23–36. doi:10.1016/j.neuro.2011.11.002
PubMed PMID: 22101008.

57. Melzer D, Harrie L, Cipelli R, Henley W, Money C, Mccormack P, et al. Bisphenol a
exposure is associated with in vivo estrogenic gene expression in adults. Environ
Health Perspect. 2011;119(12):1788–93. doi:10.1289/ehp.1103809 PubMed PMID:
21831745.

58. Kurian JR, Olesen KM, Auger AP. Sex differences in epigenetic regulation of the estrogen
receptor-α promoter within the developing preoptic area. Endocrinology.
2010;151(5):2297–305. doi:10.1210/en.2009-0649 PubMed PMID: 20237133.

59. McCarthy MM, Herold K, Stockman SL. Fast, furious and enduring: Sensitive versus
critical periods in sexual differentiation of the brain. Physiol Behav. 2018 Apr;187:13–
9. doi:10.1016/j.physbeh.2017.10.030

60. MacLusky NJ, Chaptal C, McEwen BS. The development of estrogen receptor systems
in the rat brain and pituitary: Postnatal development. Brain Res. 1979 Dec;178(1):143–
60. doi:10.1016/0006-8993(79)90094-5

61. Peña CJ, Neugut YD, Champagne FA. Developmental Timing of the Effects of Maternal
Care on Gene Expression and Epigenetic Regulation of Hormone Receptor Levels in
Female Rats. Endocrinology. 2013 Nov 1;154(11):4340–51. doi:10.1210/en.2013-1595

62. Wylie DC, Hofmann HA, Zemelman BV. SArKS: *de novo* discovery of gene expression
regulatory motif sites and domains by suffix array kernel smoothing. Birol I, editor.
Bioinformatics. 2019 Oct 15;35(20):3944–52. doi:10.1093/bioinformatics/btz198

63. Mohammadi Ghahhari N, Sznurkowska MK, Hulo N, Bernasconi L, Aceto N, Picard D.
Cooperative interaction between ERα and the EMT-inducer ZEB1 reprograms breast
cancer cells for bone metastasis. Nat Commun. 2022 Apr 19;13(1):2104.
doi:10.1038/s41467-022-29723-5

64. Zhang J, Zhou C, Jiang H, Liang L, Shi W, Zhang Q, et al. ZEB1 induces ER-α promoter
hypermethylation and confers antiestrogen resistance in breast cancer. Cell Death
Dis. 2017 Apr 6;8(4):e2732–e2732. doi:10.1038/cddis.2017.154

65. Zhang S, Kong S, Wang B, Cheng X, Chen Y, Wu W, et al. Uterine Rbpj is required for
embryonic-uterine orientation and decidual remodeling via Notch pathway-
independent and -dependent mechanisms. Cell Res. 2014 Aug;24(8):925–42.
doi:10.1038/cr.2014.82

66. Fukuda T, Shirane A, Wada-Hiraike O, Oda K, Tanikawa M, Sakuabashi A, et al. HAND2-
mediated proteolysis negatively regulates the function of estrogen receptor α. Mol
Med Rep. 2015 Oct;12(4):5538–44. doi:10.3892/mmr.2015.4070

67. Mestre-Citrinovitz AC, Kleff V, Vallejo G, Winterhager E, Saragüeta P. A Suppressive
Antagonism Evidences Progesterone and Estrogen Receptor Pathway Interaction with
Concomitant Regulation of Hand2, Bmp2 and ERK during Early Decidualization. Jeong
JW, editor. PLOS ONE. 2015 Apr 21;10(4):e0124756.
doi:10.1371/journal.pone.0124756

68. Johnson SA, Marín-Bivens CL, Miele M, Coyle CA, Fissore R, Good DJ. The Nhlh2
transcription factor is required for female sexual behavior and reproductive longevity.
Horm Behav. 2004 Nov;46(4):420–7. doi:10.1016/j.yhbeh.2004.03.006

69. Li D, Mitchell D, Luo J, Yi Z, Cho SG, Guo J, et al. Estrogen Regulates KiSS1 Gene
Expression through Estrogen Receptor α and SP Protein Complexes. Endocrinology.
2007 Oct 1;148(10):4821–8. doi:10.1210/en.2007-0154

70. Hoa N, Ge L, Korach KS, Levin ER. Estrogen receptor beta maintains expression of
KLF15 to prevent cardiac myocyte hypertrophy in female rodents. Mol Cell Endocrinol.
2018 Jul;470:240–50. doi:10.1016/j.mce.2017.11.004

71. Ray S, Pollard JW. KLF15 negatively regulates estrogen-induced epithelial cell
proliferation by inhibition of DNA replication licensing. Proc Natl Acad Sci. 2012 May
22;109(21). doi:10.1073/pnas.1118515109

72. Zhou J, Xiong Z, Long X, Yang L, Jin W, Han X. Estradiol inhibits endometrial injury by
promoting the stability of the KLF15 protein and the recovery of mitochondrial
function. Exp Cell Res. 2025 Jul;450(2):114651. doi:10.1016/j.yexcr.2025.114651

73. deGraffenried LA, Hilsenbeck SG, Fuqua SAW. Sp1 is essential for estrogen receptor α
gene transcription. J Steroid Biochem Mol Biol. 2002 Sep;82(1):7–18.
doi:10.1016/S0960-0760(02)00151-6

74. Schultz JR, Petz LN, Nardulli AM. Estrogen receptor α and Sp1 regulate progesterone
receptor gene expression. Mol Cell Endocrinol. 2003 Mar;201(1–2):165–75.
doi:10.1016/S0303-7207(02)00415-X

75. Massah S, Foo J, Li N, Truong S, Nouri M, Xie L, et al. Gli activation by the estrogen
receptor in breast cancer cells: Regulation of cancer cell growth by Gli3. Mol Cell
Endocrinol. 2021 Feb;522:111136. doi:10.1016/j.mce.2020.111136

76. Heard ME, Velarde MC, Giudice LC, Simmen FA, Simmen RCM. Krüppel-Like Factor 13
Deficiency in Uterine Endometrial Cells Contributes to Defective Steroid Hormone
Receptor Signaling but Not Lesion Establishment in a Mouse Model of
Endometriosis1. Biol Reprod. 2015 Jun 1;92(6). doi:10.1095/biolreprod.115.130260

77. Le TP, Sun M, Luo X, Kraus WL, Greene GL. Mapping ERβ Genomic Binding Sites Reveals
Unique Genomic Features and Identifies EBF1 as an ERβ Interactor. Dahlman-Wright
K, editor. PLoS ONE. 2013 Aug 8;8(8):e71355. doi:10.1371/journal.pone.0071355

78. Lotesto MJ, Raimondi SL, Department of Biology, Elmhurst University, Elmhurst, IL,
USA. EBF1 Exhibits Crosstalk Regulation with ERα and ERβ in Some Hormone-Based
Cancers. OBM Genet. 2020 Oct 6;4(4). doi:10.21926/obm.genet.2004117

79. Woodfield GW, Hitchler MJ, Chen Y, Domann FE, Weigel RJ. Interaction of TFAP2C with
the Estrogen Receptor-α Promoter Is Controlled by Chromatin Structure. Clin Cancer
Res. 2009 Jun 1;15(11):3672–9. doi:10.1158/1078-0432.CCR-08-2343

80. Woodfield GW, Horan AD, Chen Y, Weigel RJ. TFAP2C Controls Hormone Response in
Breast Cancer Cells through Multiple Pathways of Estrogen Signaling. Cancer Res.
2007 Sep 15;67(18):8439–43. doi:10.1158/0008-5472.CAN-07-2293

81. Ivanova MM, Radde BN, Son J, Mehta FF, Chung SH, Klinge CM. Estradiol and tamoxifen
regulate NRF-1 and mitochondrial function in mouse mammary gland and uterus. J
Mol Endocrinol. 2013 Oct;51(2):233–46. doi:10.1530/JME-13-0051

82. Zhao W, Hou Y, Zhang Q, Yu H, Meng M, Zhang H, et al. Estrogen receptor β exerts
neuroprotective effects by fine-tuning mitochondrial homeostasis through
NRF1/PGC-1α. Neurochem Int. 2023 Dec;171:105636.
doi:10.1016/j.neuint.2023.105636

83. Mattingly KA, Ivanova MM, Riggs KA, Wickramasinghe NS, Barch MJ, Klinge CM.
Estradiol Stimulates Transcription of Nuclear Respiratory Factor-1 and Increases
Mitochondrial Biogenesis. Mol Endocrinol. 2008 Mar 1;22(3):609–22.
doi:10.1210/me.2007-0029

84. Inoue A, Omoto Y, Yamaguchi Y, Kiyama R, Hayashi S. Transcription factor EGR3 is
involved in the estrogen-signaling pathway in breast cancer cells. J Mol Endocrinol.
2004 Jun 1;32(3):649–61. doi:10.1677/jme.0.0320649

85. Cavalcanti FN, Lucas TFG, Lazari MFM, Porto CS. Estrogen receptor ESR1 mediates
activation of ERK1/2, CREB, and ELK1 in the corpus of the epididymis. J Mol
Endocrinol. 2015 Jun;54(3):339–49. doi:10.1530/JME-15-0086

86. Takayanagi S, Tokunaga T, Liu X, Okada H, Matsushima A, Shimohigashi Y. Endocrine
disruptor bisphenol A strongly binds to human estrogen-related receptor γ (ERRγ) with
high constitutive activity. Toxicol Lett. 2006 Dec;167(2):95–105.
doi:10.1016/j.toxlet.2006.08.012

87. Okada H, Tokunaga T, Liu X, Takayanagi S, Matsushima A, Shimohigashi Y. Direct
Evidence Revealing Structural Elements Essential for the High Binding Ability of
Bisphenol A to Human Estrogen-Related Receptor-γ. Environ Health Perspect. 2008
Jan;116(1):32–8. doi:10.1289/ehp.10587

88. Zou Z, Harris LK, Forbes K, Heazell AEP. Sex-specific effects of bisphenol A on the
signaling pathway of ESRRG in the human placenta. Biol Reprod. 2022 Jun
13;106(6):1278–91. doi:10.1093/biolre/ioac044

89. Hilz EN, Schnurer C, Bhamidipati S, Deka J, Thompson LM, Gore AC. Cognitive effects
of early life exposure to PCBs in rats: Sex-specific behavioral, hormonal and
neuromolecular mechanisms involving the brain dopamine system. Horm Behav.
2025 Mar;169:105697. doi:10.1016/j.yhbeh.2025.105697

90. Vandenberg LN. Non-Monotonic Dose Responses in Studies of Endocrine Disrupting
Chemicals: Bisphenol a as a Case Study. Dose-Response. 2014 Apr 1;12(2):dose-
response.1. doi:10.2203/dose-response.13-020.Vandenberg

91. Vandenberg LN, Ehrlich S, Belcher SM, Ben-Jonathan N, Dolinoy DC, Hugo ER, et al.
Low dose effects of bisphenol A: An integrated review of in vitro, laboratory animal,
and epidemiology studies. Endocr Disruptors. 2013 Oct;1(1):e26490.
doi:10.4161/endo.26490

92. Wolstenholme JT, Taylor JA, Shetty SRJ, Edwards M, Connelly JJ, Rissman EF.
Gestational Exposure to Low Dose Bisphenol A Alters Social Behavior in Juvenile Mice.
Ferrari PF, editor. PLoS ONE. 2011 Sep 28;6(9):e25448.
doi:10.1371/journal.pone.0025448

93. van Hasselt FNN, Tieskens JMM, Trezza V, Krugers HJJ, Vanderschuren LJMJJMJ, Joëls M.
Within-litter variation in maternal care received by individual pups correlates with
adolescent social play behavior in male rats. Physiol Behav. 2012;106(5):701–6.
doi:10.1016/j.physbeh.2011.12.007 PubMed PMID: 22210522.

94. Melo AI, Lovic V, Gonzalez A, Madden M, Sinopoli K, Fleming AS. Maternal and
littermate deprivation disrupts maternal behavior and social-learning of food
preference in adulthood: Tactile stimulation, nest odor, and social rearing prevent
these effects. Dev Psychobiol. 2006. doi:10.1002/dev.20130

95. Gonzalez A, Lovic V, Ward GR, Wainwright PE, Fleming AS. Intergenerational effects of
complete maternal deprivation and replacement stimulation on maternal behavior
and emotionality in female rats. Dev Psychobiol. 2001;38(1):11–32. doi:10.1002/1098-
2302(2001)38:1%3C11::AID-DEV2%3E3.0.CO;2-B PubMed PMID: 11150058.

96. Moon HJ, Shin HS, Lee SH, Hong EJ, Ahn C, Yoo YM, et al. Effects of prenatal bisphenol
S and bisphenol F exposure on behavior of offspring mice. Anim Cells Syst. 2023 Dec
11;27(1):260–71. doi:10.1080/19768354.2023.2264905

97. Starr-Phillips EJ, Beery AK. Natural variation in maternal care shapes adult social
behavior in rats. Dev Psychobiol. 2014 Jul;56(5):1017–26. doi:10.1002/dev.21182

98. Franks B, Champagne FA, Curley JP. Postnatal maternal care predicts divergent
weaning strategies and the development of social behavior. Dev Psychobiol. 2015
Nov;57(7):809–17. doi:10.1002/dev.21326

99. Wallen K, Baum MJ. Masculinization and Defeminization in Altricial and Precocial
Mammals: Comparative Aspects of Steroid Hormone Action. In: Hormones, Brain, and
Behavior. Academic Press; 2002. p. 385–423.

100. Lapp HE, Salazar MG, Champagne FA. Automated maternal behavior during early life
in rodents (AMBER) pipeline. Sci Rep. 2023 Oct 25;13(1):18277. doi:10.1038/s41598-
023-45495-4

101. Paxinos G, Tork I, L.H. T, Valentino K. Atlas of the Developing Rat Brain. Academic
Press; 1990.

102. Khazipov R, Zaynutdinova D, Ogievetsky E, Valeeva G, Mitrukhina O, Manent JB, et al.
Atlas of the Postnatal Rat Brain in Stereotaxic Coordinates. Front Neuroanat. 2015;9.
doi:10.3389/fnana.2015.00161 PubMed PMID: 26778970.

103. Lohman BK, Weber JN, Bolnick DI. Evaluation of TagSeq, a reliable low-cost alternative
for RNAseq. Mol Ecol Resour. 2016;16(6):1315–21. doi:https://doi.org/10.1111/1755-
0998.12529

104. MEYER E, AGLYAMOVA GV, MATZ MV. Profiling gene expression responses of coral
larvae (Acropora millepora) to elevated temperature and settlement inducers using a
novel RNA-Seq procedure. Mol Ecol. 2011;20(17):3599–616.
doi:https://doi.org/10.1111/j.1365-294X.2011.05205.x

105. Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, et al. Enrichr: interactive and
collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics. 2013
Dec;14(1):128. doi:10.1186/1471-2105-14-128

106. Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr: a
comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids
Res. 2016 Jul 8;44(W1):W90–7. doi:10.1093/nar/gkw377

107. Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. Nucleic Acids Res. 2015
Jul 1;43(W1):W39–49. doi:10.1093/nar/gkv416

108. Sasaki A, Eng ME, Lee AH, Kostaki A, Matthews SG. DNA methylome signatures of
prenatal exposure to synthetic glucocorticoids in hippocampus and peripheral whole
blood of female guinea pigs in early life. Transl Psychiatry. 2021;11(1).
doi:10.1038/s41398-020-01186-6 PubMed PMID: 33462183.

109. Rauluseviciute I, Riudavets-Puig R, Blanc-Mathieu R, Castro-Mondragon JA, Ferenc K,
Kumar V, et al. JASPAR 2024: 20th anniversary of the open-access database of
transcription factor binding profiles. Nucleic Acids Res. 2024 Jan 5;52(D1):D174–82.
doi:10.1093/nar/gkad1059

110. Lauby S, Champagne FA. DNA methylation data for "Postnatal maternal care impacts
hypothalamic Esrrg gene expression, co-expression profiles, and the DNA methylome
in prenatal bisphenol-exposed rats" [Internet]. Zenodo; 2025 [cited 2025 Oct 6].
Available from: https://zenodo.org/doi/10.5281/zenodo.17273659
doi:10.5281/ZENODO.17273659

1262