**Environmental DNA Metabarcoding Effectively Detects Invasive Species, Pests, and
Community Changes in Taiwan's Rice Fields**

Pritam Banerjee[1,2,*], Gobinda Dey[2], Kathryn A. Stewart[3], Matthew A. Barnes[4], Md. Taharia[2],
Mathew Seymour[5], Chin-Wen Wang[2], Raju Kumar Sharma[2], Jyoti Prakash Maity[6], Chien-Yen
Chen[2, 8,*]

[1]*Department of Environmental Science, Policy, and Management, University of California,
University Avenue and Oxford St, Berkeley, CA, 94720, USA*

[2]*Department of Earth and Environmental Sciences, National Chung Cheng University, 168
University Road, Min-Hsiung, Chiayi County 62102, Taiwan.*

[3]*Institute of Environmental Sciences, Leiden University, Leiden, 2333 CC, The Netherlands*

[4]*Department of Natural Resources Management, Texas Tech University, Lubbock, Texas, USA*

[5]*School of Biological Sciences, The University of Hong Kong SAR, China*

[6]*Department of Chemistry, School of Applied Sciences, KIIT Deemed to be University,
Bhubaneswar, Odisha 751024, India.*

[8]*Center for Nano Bio-Detection, Center for Innovative Research on Aging Society, AIM-HI,
National Chung Cheng University, Chiayi 62102, Taiwan*

Corresponding Authors*

Chien-Yen Chen[2,8*] and Pritam Banerjee[1,2*]

*Address for correspondence: Department of Earth and Environmental Sciences, National Chung
Cheng University, 168 University Road, Min-Hsiung, Chiayi County 62102, Taiwan. And
Department of Environmental Science, Policy, and Management, University of California,

23 University Avenue and Oxford St, Berkeley, CA 94720, USA Email: chien-

24 yen.chen@oriel.oxon.org; yen@eq.ccu.edu.tw (C.Y. Chen); pritam8683@gmail.com (P.

25 Banerjee); Tel.: +886-5-2720411 (ext. 66220); Fax: +886-5-2720807.

Abstract

Rice fields represent man-made semi-aquatic wetlands primed for invasive pests. Monitoring rice

field biodiversity using conventional methods, however, is time-consuming and laborious.

Environmental DNA (eDNA) methods can provide a fast and effective means to monitor rice

field communities and inform management decisions. Our study provides proof-of-concept of

rice field eDNA biodiversity assessments, with a focus on native and non-native pests across

cultivation phases. We collected eDNA samples from locations in southern Taiwan during

planting and harvesting, employing eDNA metabarcoding (COI) to detect diverse taxonomic

groups. We assigned 77 ASVs across all sites to animal taxa, 34 of which were identified to

species. Overall, 18 species were designated as native or non-native (83.3% and 16.6%,

respectively), including three major rice pests, *Chilo suppressalis* (native), *Coptotermes*

formosanus (native), and *Pomacea canaliculata* (non-native). Cultivation status affected overall

diversity, with higher species richness during planting compared to harvesting. No significant

differences were observed between native and non-native taxa between cultivation phases.

Altogether, we detected a complex environment across trophic levels comprised of both native

and non-native agricultural pests using limited sampling effort, demonstrating eDNA analysis as

an efficient biomonitoring approach in rice agroecosystems with direct applications for pest,

invasive   species,   and   vector   surveillance   within   Taiwan.

Introduction

Rice fields represent man-made, wetland agroecosystems that support a wide range of biodiversity, from microorganisms to macroinvertebrates and amphibians (Bambaradeniya & Amarasinghe, 2004). Moreover, their alternating hydrologic dynamics, from flooded during the wet seasons to dry conditions in between, mean rice fields house a unique array of biodiversity (Watanabe et al., 2013). Additional ecological characteristics, such as nutrient-rich environments, human influences, and the effects of climate change, collectively make these ephemeral wetlands particularly vulnerable to invasive species (Flanagan et al., 2015). Major threats to the ecological stability and biodiversity in the rice fields include anthropogenic pressures (e.g., intensive monoculture, agrochemical use) and the spread of invasive species (Tilman et al., 2002; Horgan, 2023).

In Taiwan, rice is a major staple crop, deeply embedded in the economy and culture, with almost half of the arable land consisting of rice fields (Hsing, 2014). However, rice fields are one of the most susceptible places for a myriad of invasive pests (Valls et al., 2014). Taiwan's subtropical climate and extensive irrigation networks further exacerbate invasions by non-native species, posing serious harm to crop productivity and native biodiversity (Barker, 2002; Huang et al., 2010; Wu et al., 2010). Several invasive pest species severely impact rice cultivation in Taiwan; for example, the introduction and establishment of *Pomacea* spp. have become serious threats to rice agriculture (Barker, 2002; Wu et al., 2010; Banerjee et al., 2022), and are currently responsible for large economic losses throughout Asia (Jiang et al., 2022). Migratory rice planthoppers (*Nilaparvata lugens*) (Otuka et al., 2012) and the recently introduced fall armyworm (*Spodoptera frugiperda*) (Tsai et al., 2000) also represent major economic concerns. Compounding impacts, native pests, such as rice stem borers (*Chilo suppressalis*, *Sesamia*

inferens, and *Scirpophaga incertulas*), planthoppers (e.g., *Sogatella furcifera*) (Huang et al.,

2010), leafhoppers (e.g., *Nephotettix* spp.), and the rice black bug (*Scotinophara coarctata*),

additionally contribute to considerable crop damage (Cheng et al., 2010, 2022; Triapitsyn et al.,

2021).

Conventional methods of biomonitoring, such as dipnet and funnel traps, are well

established and effective for understanding the co-occurrence of organisms associated with rice

cultivation (Song & Kuo, 2022), particularly invertebrates. However, conventional methods are

also time-consuming and require taxonomic expertise, limiting the scale and scope needed for

establishing large-scale temporal monitoring needed to enable adaptive resource management

(Baird & Hajibabaei, 2012). Environmental DNA (eDNA)-based metabarcoding has recently

revolutionized biodiversity monitoring by enabling the non-destructive detection of targeted or

multiple species directly from environmental samples (Taberlet et al., 2012; Valentini et al.,

2016; Didaskalou et al., 2022). The adoption of eDNA metabarcoding has proven valuable in

agricultural ecosystems, where ecological interactions, including both beneficial and pest

organisms, can be monitored for proper management decisions (Kestel et al., 2022; Llanos et al.,

2025). However, broad-scale implementation of eDNA-based monitoring in agricultural

practices remains limited (Kestel et al., 2022) to a few studies assessing general community

observations in rice fields (Ushio et al., 2023; Katayama et al., 2024; Vastano et al., 2024).

Previous studies have established species-specific approaches for an early detection method for

cryptic invasive gastropods (Banerjee et al., 2022) and fall armyworm (Tsai et al., 2020) in

Taiwan. However, more robust and community-level monitoring is necessary for early

management action that goes beyond species-specific targets.

Recent advancements in high-throughput sequencing technologies have enhanced the

resolution of eDNA metabarcoding approaches, enabling detailed taxonomic and functional

assessments of communities present within agricultural environments (Valentini et al., 2016;

Seymour et al., 2021; Ushio et al., 2023; Katayama et al., 2024; Vastano et al., 2024). Our study

aimed to elucidate a comprehensive understanding of co-occurring species associated with rice

field cultivation across two phases of rice cultivation (i.e., planting and harvesting) in Taiwan.

To do so, we implemented an eDNA-based metabarcoding approach targeting the mitochondrial

cytochrome c oxidase I (COI) gene to investigate the community dynamics of native, non-native,

and pest species associated with various rice fields. As environmental conditions (e.g., pH,

salinity, temperature, and other abiotic factors) putatively influence the shaping of community

composition and function within agroecosystems (Hartmann & Six, 2022) and potentially also

modulate eDNA detectability and persistence (Barnes & Turner, 2016; Stewart, 2019), we co-

measured water and soil parameters alongside our eDNA sampling.

We hypothesized that (i) eDNA metabarcoding would be an effective method for

detecting the vast array of interactive animal species from rice fields via water samples; and (ii)

during the planting season, species diversity would be high, but community composition would

demonstrate a higher proportion of colonizing non-natives and pests compared to harvesting time

due to the drier environment.

Materials and Methods

Sample collection and environmental parameter analysis

We collected eDNA samples from eight rice fields in Chiayi County, Taiwan (Figure 1, Table 1).

To understand the changes in the community during rice growth, including native and non-native

species, all fields were visited two times: (i) planting phase (denoted as samples "PF1-PF8") which consist of the period of time at the beginning of the growing season where crops are seeded and vegetative growth initiates, and (ii) harvesting phase ("HF1-HF8"), characterized by the ripening phase, with mature rice seedling and dry environment. In our study, planting phase sampling occurred on 22 February 2022, and the harvesting phase sampling occurred on 31 May 2022. We collected five 500 mL surface water samples from each field (N = 40 samples per season). Water samples were kept on ice and transported directly to the Department of Earth and Environmental Sciences, National Chung Cheng University, for analysis.

After collection of eDNA, we measured water chemistry at each site including pH, dissolved oxygen (DO) (mg/L), temperature (°C), salinity (PSU), oxidation-reduction potential (ORP) (mV), conductivity (mS/cm), total dissolved solids (TDS) (mg/L), and resistivity (Ω-cm) using a Hanna multiparameter waterproof meter (HANNA instruments, model HI98194, USA), replicating each measurement ten times per site. Subsequently, the water sample was acidified with 5 M HNO☐ to achieve pH < 2, followed by filtration using 0.22-µm nylon syringe filters (13☐mm; PureTech™, Taiwan) into 50-ml centrifuge tubes for further heavy metal analysis (see the analytical methods below). Additionally, 3 x 300 g soil samples were collected from each site to measure the physicochemical properties (such as pH and percentage of sand, silt, and clay with texture) and heavy metal concentration, following protocols outlined in Dey et al. (2022). Soil pH was measured using a 1:2.5 soil-to-deionized water suspension with a pH meter. Soil particle size distribution was analyzed by the hydrometer method and classified into textural classes following USDA-NRCS guidelines (Bouyoucos, 1962).

Concentrations of elements Bi, Cd, Mn, Cu, Ni, Co, Cr, Sr, Zn, Pb, and Fe in filtered acid-digested water samples were analyzed using Inductively Coupled Plasma Optical Emission

Spectrometry (ICP-OES; Agilent Technologies 5100 ICP-OES, CA, USA). To ensure accuracy

and precision, a negative control ultrapure water (PP1-185UV Type I, UNISS, Taiwan) and a

positive reference standard were included (ICP multi-element standard solution IV). The

recovery rates for the analyzed metals ranged between 94% and 115%, with precision maintained

under a 10% relative standard deviation (RSD) for all elements. The soil samples were dried at

70°C for 48 hours, followed by grinding in a mortar and pestle, and then were prepared for heavy

metal concentrations analysis using ICP-OES (Agilent Technologies 5100 ICP-OES, USA),

based on the method described by Bakshi et al. (2018). Briefly, 0.4 g dried soil was digested with

8 ml aqua regia (a 3:1 mixture of HCl and HNO3) and 2 ml 40% hydrofluoric acid (HF). After

cooling, the solution was diluted with 30 mL of deionized water and then filtered with 0.22 µm

nylon syringe filters (13 mm; PureTech™, Taiwan) before the ICP-OES analysis.

eDNA Filtration, Extraction, and Amplification

In the laboratory, all water samples were passed through a GN-6 Metricel membrane mixed

cellulose ester filter paper (GN-6 Metricel ®, Pall Corporation, USA) with a diameter of 47 mm

and a pore size of 0.45 µm. As rice field water has high TDS, any clogging issues were mitigated

by using multiple filter papers as sub-samples (e.g., Hunter et al. 2019). Once our filtration was

complete, half of each filter paper was processed for immediate eDNA extraction, and the latter

half was preserved at -20 °C in Longmire's buffer for future use (Longmire et al., 1997). For

samples that required multiple sub-filters to mitigate clogging, all filter papers corresponding to

the same sample were placed in a single tube for DNA extraction. During the filtration process,

three filtration blanks (distilled water) were included to determine contamination and false

positive detection during laboratory analysis. Sterilization processes were maintained throughout

the workflow, like (i) frequent decontamination with 70% ethanol and 10% bleach, (ii)

unidirectional workflow with each step (i.e., filtration, DNA extraction, PCR, and post-PCR)

performed in a separate space, and (iii) PCRs were performed in a well-maintained hood.

We extracted eDNA from filters using DNeasy Blood & Tissue kits (Qiagen, Germany)

according to the manufacturer's protocol, but with some modifications as follows: (i) the filter

papers were incubated at 60°C overnight in a mixture of 180 µL of ATL lysis buffer and 20 µL

Proteinase K; and (ii) the elution step was performed twice, where the spin columns were

incubated with the same 50 µL elution buffer at 37°C for 10 minutes (Spens et al., 2017). The

quality and quantity of extracted DNA samples were assessed using a Nanodrop

spectrophotometer (Implen, Germany) and an Invitrogen Qubit 3.0 fluorometer (Thermo Fisher

Scientific, USA). For budgetary reasons, a pooling approach was used for sequencing (Melcher

et al., 2024). After extractions, all subsamples (5 samples from each field) were pooled together

based on their quantity, thus making 8 pooled samples for each planting (n = 8) and harvesting (n

= 8). Finally, these 16 pooled samples were used for further downstream analysis.

A portion of mitochondrial cytochrome c oxidase subunit I (COI) gene was amplified

with the following primers: mlCOIintF- 5′- GGWACWGGWTGAACWGTWTAYCCYCC- 3′

and jgHCO2198 - 5′-TAIACYTCIGGRTGICCRAARAAYCA- 3′, specially designed for

metazoans (Leray et al., 2013). The reaction mixture of 25 µL for conventional PCR was

prepared by mixing 5 µL 5x Fast-Run^{TM} Taq Master Mix with Dye (Protech Technology

Enterprise, Taiwan), 0.5 µL each 10 pmol forward and reverse primers (Genomics, Taiwan), 3

µL eDNA sample, and adding high-quality sterile distilled water (Thermo Fisher Scientific,

USA) to make up the final volume. For PCR, we tried to add 3 µL eDNA instead of 1µL to allow

the potential to amplify all pooled samples. Three PCR replicates for each sample (including

blank samples) were carried out. The PCR reactions were performed with the following

conditions: an initial denaturation at 94°C for 5 min, followed by a continuous amplification

process of 36 cycles, with denaturation at 94°C for 30 sec, annealing at 55°C for 30 sec, and

extension at 72°C for 1 min. The final extension was at 72°C for 5 min after the completion of

36 cycles. The PCR products were visualized in a 1.5% agarose gel electrophoresis, and all

subsamples/replicates were mixed in equal density ratios (based on band intensity) using the Gel

Doc XR system with the QUANTITY ONE™ software (Bio-Rad, USA). No visible

amplification was observed in any blank sample (filtration or extraction blanks); these were

therefore excluded from further analysis. The PCR products were then purified using 1.5× SPRI

beads (AMPure XP; Beckman Coulter, USA). Purified amplicons were quantified using an

Invitrogen Qubit 3.0 fluorometer and submitted for library preparation and sequencing at

Genomics Bioscience and Technology Co., Ltd., Taiwan.

Library preparation and next-generation sequencing

Libraries were prepared using the TruSeq Nano DNA Library Prep Kit (Illumina, USA), and

quality was assessed using an Agilent Bioanalyzer 2100 system (Agilent Technologies, USA).

The target inserts (~313 bp of the mitochondrial COI gene fragment) were derived from the PCR

amplification described earlier. The final library size, including Illumina adapters, was

approximately 450-500 bp, as measured by Bioanalyzer. Libraries were individually indexed

(barcoded) during preparation and pooled (multiplexed) in equimolar concentrations before

sequencing. Library quantification was performed using qPCR to ensure accurate equimolar

pooling. All qualified libraries were sequenced using 300 bp paired-end reads on an Illumina

MiSeq platform (Illumina, USA), operated by Genomics Bioscience and Technology Co., Ltd.,

Taiwan. Sequencing was carried out on a multiplexed standard v3 MiSeq flow cell.

Data processing

Raw sequencing data were demultiplexed using an in-house script developed by Genomics

Bioscience and Technology Co., Ltd., Taiwan, to assign reads to their corresponding samples

based on barcode sequences. Trimmomatic (v0.39) was used to remove the adapter sequences,

and low-quality bases (QV < 3) using with the following parameters:

ILLUMINACLIP:adapter:2:30:10:1:true LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15.

This configuration removes Illumina adapter sequences (allowing up to two seed mismatches),

trims low-quality bases from both ends of the reads, and applies sliding window trimming when

the average Phred quality score within the window falls below 15 (Bolger et al., 2014).

Sequences corresponding to both ends of the mlCOIintF and jgHCO2198 primers were trimmed

using Cutadapt (v3.4) (Martin, 2011). Primer trimming was performed with a minimum read

length cutoff of 150 bp to remove short, low-information reads, and reads lacking primer

sequences were discarded to ensure that only primer-containing sequences were retained. The

trimming allowed a minimum overlap of 3 bp and an error rate of 0.1 for primer matching.

Denoising, chimera removal, and assignment to Amplicon Sequence Variants (ASVs) were

performed using DADA2 (Callahan et al., 2016), following the approach of (Sigsgaard et al.,

2021). After error removal, all remaining sequences were then compared against the GenBank

nucleotide database (Sayers et al., 2022) (as of September 2022) using BLASTN, and taxa were

selected based on the Lowest Common Ancestor (LCA) method

(https://github.com/timkahlke/BASTA). The criteria for sequence alignment included a

maximum of 500 aligned sequences per query, with a ≥90% query cover and 70-100% sequence

similarity. For taxonomic assignments, we applied similarity thresholds as follows: 80-84% for

class-level, 85-89% for order-level, 90-94% for family-level, 95-97% for genus-level, and >98%

for species-level (Sigsgaard et al., 2021). In cases where multiple taxonomic matches occurred at

a given level, the ASV was assigned to the next higher taxonomic rank. The species detected via

eDNA metabarcoding were classified as native or non-native based on previous literature and the

Catalog of Life in Taiwan (https://taicol.tw/). Furthermore, the ecological roles were documented

from previously published literature accessed through Google Scholar, PubMed, and Web of

Science.

Statistical analysis

All statistical analyses were performed using R version 4.3.3 (R Core Team, 2025). Alpha

diversity was estimated as observed ASV richness (number of unique ASVs per sample) and

visualized across field types using *ggplot2*. Differences in richness between cultivational phases

were assessed using Wilcoxon rank-sum tests. Beta-diversity was assessed using non-metric

multidimensional scaling (NMDS) based on Bray–Curtis dissimilarities with the metaMDS()

function in the vegan package (Oksanen et al., 2001). Group differences in community

composition between cultivational stages were tested using PERMANOVA (*adonis2*). To

visualize the associations between species and cultivation phases (harvesting and planting), a

sankey diagram was constructed using the *networkD3* package (Allaire et al., 2014). All

visualization of taxonomic distributions and sample overlaps were performed using the *ggplot2*

and *eulerr* packages (Wickham, 2016; Yan, 2021). For the environmental parameter, mean

values ± standard error (SE) were calculated, and differences between harvesting and planting

periods were evaluated using Welch's two-sample t-test (α = 0.05).

Results

Community-level dynamics of rice fields

Sequencing reads and taxonomic composition

After filtering, 1,014,225 sequencing reads were retained, resulting in 67,615 ± 10,984 reads

(mean ± standard error; n=15) per sample. Overall, 196 amplicon sequence variants (ASVs) were

initially identified across taxa, including Metazoa, Protozoa, Algae, Fungi, and Bacteria. We

discarded one sample (HF2) due to low sequencing reads (< 3000). Of the total sequence reads,

54.92% originated from Animalia (80 ASVs), 12.16% from Protozoa (25 ASVs), 4.24% from

Bacteria (13 ASVs), 3.29% from Algae (38 ASVs), and 1.40% from Fungi (38 ASVs) (Table 2).

Additionally, 23.95% of reads could not be taxonomically assigned with high confidence, as they

fell below the 80% similarity threshold (2 ASVs) (Table 2). After removing the unassigned taxa

(2 human ASVs were also removed) and targeting animal taxa only (one ASV that could only

identify up to Animalia was removed), a total of 77 ASVs were retained for downstream

analysis. The majority of reads (93.12%) were from Arthropoda (43 ASVs), followed by

Annelida (11 ASVs), Rotifera (8 ASVs), Mollusca (5 ASVs), Chordata (4 ASVs), Gastrotricha

(2 ASVs), Cnidaria (2 ASVs), and Nematoda (2 ASVs) (Table 3). In total, 34 species were

identified from various groups, and among them, 18 species could be designated as either native

or non-native based on published literature, involving 83.3% native, 16.6% non-native (Table 4).

Environmental parameters

Soil pH was slightly alkaline (7.74 ± 0.05 at harvesting, 7.54 ± 0.07 at planting) across sites

(Table 5), with textures ranging from sandy loam to loamy sand. Particle size composition

remained consistent across both cultivation times, dominated by sand (approximately 72%).

Heavy metal concentrations in soil were found to exhibit temporal variation, with Cd, Co, Cr,

Cu, Ni, Zn, and Ag generally higher during harvesting, though differences were not statistically

significant (Table 5). In contrast, both pH and Lead (Pb) were significantly elevated in soil at

harvest ($p < 0.05$; Table 5).

Water parameters also reflected alkaline pH (7.52 ± 0.05 at harvesting, 7.54 ± 0.05 at planting).

ORP was significantly higher during the planting phase (188.99 ± 3.77) compared to the

harvesting phase (152.37 ± 11.67; $p = 0.0270$). Dissolved oxygen (DO) was also higher during

planting (5.27 ± 0.26 mg/L) than harvesting (4.45 ± 0.02 mg/L; $p = 0.0197$). Similarly,

conductivity increased during planting (0.55 ± 0.05 mS/cm) relative to harvesting (0.40 ± 0.04

mS/cm; $p = 0.0363$), and total dissolved solids (TDS) were greater in planting (259.80 ± 22.12

mg/L) than harvesting (187.08 ± 20.39 mg/L; $p = 0.0424$). In contrast, temperature was

significantly higher during the harvesting phase (29.32 ± 0.88 °C) compared to the planting

phase (26.08 ± 0.76 °C; $p = 0.0231$). Among trace metals, only lithium (Li) showed a statistically

significant difference, with slightly higher values during harvesting (0.27 ± 0.00 mg/L) than

planting (0.26 ± 0.00 mg/L; $p = 0.0082$). Other parameters, including pH, mV pH, resistivity,

PSU, atmospheric pressure, Fe, Mn, and Sr, did not differ significantly between phases ($p >$

0.05).

Planting vs harvesting phase

The relative abundance of the top 10 taxa at different taxonomic levels (order, family, genus, and

species) varied markedly between harvesting and planting phases (Figure 2). At the order level,

Diplostraca was the dominant order in both field types, accompanied primarily by Diptera in

harvesting fields and by Crassiclitellata and Cyclopoida in planting fields. At the family level,

Moinidae dominated both harvesting and planting fields, co-occurring with Culicidae and

Daphniidae in harvesting fields and with Megascolecidae and Cyclopidae in planting fields. At

the genus level, *Moina* sp., dominated both harvesting and planting fields, co-occurring with

Culex sp., and *Ceriodaphnia* sp., in harvesting fields and with *Metaphire* sp., and *Thermocyclops*

sp., in planting fields. At the species level, *Moina macrocopa* was dominant in both fields,

accompanied by *Culex tritaeniorhynchus* and *Ceriodaphnia cornuta* in harvesting fields and by

Metaphire posthuma in planting fields.

Of note, species overlap between planting and harvesting stages was substantial (44.1%;

Figure 3A) but also demonstrated some exclusive taxa between both cultivation phases, with

more unique species during the planting (35.7%) than the harvesting phase (14.3%) (Figures 3A,

B). This was corroborated by a Sankey diagram showing the association between the cultivation

phases and the relative abundance (based on sequencing reads) of taxa (Figure 3B). Moreover,

species richness was higher during the planting phase, with 12 native, 3 non-native, and 14

undetermined (status) taxa, compared to 9 native, 2 non-native, and 9 undetermined (status) taxa

in the harvesting phase (Figure 4; Table 4). Still, alpha-diversity did not differ significantly

between harvesting and planting phases (p-value = 0.52) (Figure 5A). The median species

richness was slightly higher during harvesting compared to planting phases, although the

variation was larger during planting, with a few high diversity outliers (Figure 5A).

Despite no significant differences in alpha-diversity across cultivation times, the non-

metric multidimensional scaling (NMDS) showed a pattern of differing community composition

between cultivation times (Figure 5B) (stress = 0.164). The PERMANOVA test confirmed that

the difference between the two growth phases was statistically significant ($p = 0.003$).

Additionally, no significant differences were observed between native and non-native taxa across

the two cultivation phases (Wilcoxon test, $p > 0.99$).

Discussion

In this study, we aimed to establish eDNA metabarcoding as an effective method for detecting

the vast array of interactive animal species from rice fields. Overall, the eDNA-based method

successfully detected a wide range of taxonomic groups during both planting and harvesting

phases, even with minimal sampling time and effort (Figure 3B, Table 4), consistent with the

first hypothesis (H1). We successfully detected rice pests, including *Chilo suppressalis* (native),

Coptotermes formosanus (native), and *Pomacea canaliculata* (non-native), among others (Table

4). Our results indicate that changes in rice cultivation phases correspond to significant changes

in community composition (Figure 3A, B), though the species richness between the phases

remains relatively static (Figure 5A). We didn't observe any significant difference between

native and non-native taxa across the two cultivation phases (Figure 4), counter to our second

hypothesis (H2). However, this may also be a consequence of the relatively small sample size

and limited study area. Future studies incorporating a larger number of samples and broader

spatial coverage would improve the robustness of the analysis and may reveal patterns that were

not detectable in the present study.

Functional role of the detected organism

Our analysis of the rice field ecosystem revealed a diverse community spanning multiple trophic levels, including primary consumers, detritivores, predators, omnivores, and native and non-native agricultural pests (Table 4). Primary producers included rice crops and various phytoplankton. Detritivores such as *Metaphire posthuma* (Tsai et al., 2020), aquatic oligochaetes (*Allonais*, *Dero*, *Pristina*), soil invertebrates (e.g., *Homidia socia*, *Coptotermes formosanus*) (Cheng et al., 2022; Blumenfeld et al., 2021), and gastropods (*Melanoides tuberculata*) have an important role in nutrient cycling (Chang, 1991). Phytoplankton grazers included cladocerans (e.g., *Moina macrocopa*, *Ceriodaphnia cornuta*), rotifers (*Brachionus*, *Keratella*), and gastropods (*Pomacea canaliculata*, *Austropeplea ollula*) (Table 4). Omnivores such as ants (*Monomorium pharaonis*, *Pheidole* sp.) (Wetterer, 2010), and birds (*Passer montanus*, *Acridotheres tristis*) were also detected (Ding et al., 2020). In this context, birds may play a potential role as insect predators (Sottomayor et al., 2024). Secondary consumers included *Thermocyclops decipiens* and *T. taihokuensis* (Reid, 1989), while predators such as frogs (*Microhyla fissipes*, *Fejervarya* sp.) (Jang-Liaw & Chou, 2015) and aquatic insects (*Diplonychus esakii*, *Microvelia douglasi*) further structured the food web (Rieger, 1995; Ye et al., 2014).

Two taxa of particular concern, *Chilo suppressalis* and *Pomacea canaliculata*, are major rice pests responsible for economic damage (Chu, 1971); however, the detection of *Coptotermes formosanus* is also a concern, as they may damage the rice plants (Li et al., 2009). We also recorded high abundances of *Culex tritaeniorhynchus* (Figure 2), a vector of Japanese encephalitis virus, which may raise concerns about the management of stagnant water in rice fields (Table 4).

Planting vs harvesting phase

The NMDS results demonstrate that arthropod community structure differs significantly between

harvesting and planting phases. The tighter clustering of harvesting compared to the planting

phase suggests more homogeneous community assemblages under harvesting conditions,

possibly due to low availability of water or selective pressures associated with crop removal

(Flannagan & Marshall, 2012; Saksongmuang et al., 2024). In contrast, the broader spread of

identified ASVs during planting phases indicates greater community heterogeneity, which may

reflect colonization dynamics, variability in early vegetation structure, or differences in resource

availability (Mukherjee & Khan, 2017; Saksongmuang et al., 2024). This pattern is also

supported by the measured physicochemical parameters, with the planting phase showing higher

DO, ORP, conductivity, and TDS compared to the harvesting phase, possibly due to longer water

residence time and more biological activity, whereas the harvesting stage was characterized by

relatively drier conditions and reduced water availability (Table 6). However, in this study, the

relationship between environmental parameters and indicator taxa could be better resolved with

more robust seasonal sampling.

The significant PERMANOVA results support our proposition that these biodiversity

compositional differences are not random but instead driven by the phase of rice cultivation. This

may highlight that agricultural practices, even across short time scales, can influence community

composition, possibly through habitat modification, disturbance regimes, and resource

distribution (Mukherjee & Khan, 2017). The harvesting phase encompasses warmer conditions

compared to the planting phase, driven by warmer season and slight increases in soil pH and Pb,

probably due to cumulative agrochemical deposition.

Native and non-native status

Two frog species were identified as likely native to Taiwan. Among these, *Microhyla fissipes* is recognized as native and widely distributed (Jang-Liaw & Chou, 2015). The other taxon, *Fejervarya* sp., was identified only to the genus level, which includes both a native species (*Fejervarya limnocharis*) and a non-native species (*Fejervarya cancrivora*) present in Taiwan (Jang-Liaw & Chou, 2015; Lee et al., 2019). One native (*Passer montanus*) and an introduced bird (*Acridotheres tristis*) with widespread distribution in Taiwan was also detected from the rice field water (Ding et al., 2020). Among arthropods, *Coptotermes formosanus* is reported as a native pest species (Li et al., 2009; Blumenfeld et al., 2021); however, this species is considered among the top 100 invasive species in the world and its introduced population has been established in Japan, Hawaii, and the southeastern United States (Simberloff & Rejmanek, 2019; Blumenfeld et al., 2021). The native arthropod taxa detected included *Homidia socia*, *Culex tritaeniorhynchus*, *Sciara humeralis*, *Diplonychus esakii*, *Microvelia douglasi*, *Monomorium pharaonis*, and *Chilo suppressalis*. The non-native earthworm *Metaphire posthuma* was detected (Tsai et al., 2000); this species is now naturalized and does not appear to negatively impact the local environment. Additionally, one invasive snail (*Pomacea canaliculata*) which is responsible for various crop damage, and two native snails (*Austropeplea ollula* and *Melanoides tuberculata*), were detected together in the rice fields (Chang, 1991).

Although eDNA analysis of rice field water provided a comprehensive assessment of species presence, the array of biodiversity detection was still less than other previous conventional surveys conducted in Taiwan (Song & Kuo, 2022; Chen et al., 2023). In our results, we detected 77 animal ASVs, of which 34 were assigned to species level, which is lower than reported in previous conventional studies (Song & Kuo, 2022; Chen et al., 2023). The disparity

with other studies may be due to lower sampling effort (16 pooled samples) in this study, where

repetitive monitoring and an increase in sampling events can increase the detection (Yates et al.,

2023). We did not explore the effects of eDNA persistence dynamics under hot climatic

conditions and potential PCR inhibition (McCartin et al., 2022), but their influence cannot be

neglected. Furthermore, universal concerns related to primer bias and limitations of reference

databases require further investigation (Korbel et al., 2024). Taiwan has a good reference library

(Hu et al., 2024), a direct comparison between the conventional detection and eDNA analysis

can collect more evidence of disparity. Thus, future studies should implement a comparative

analysis of conventional and eDNA methods to understand the potential community shift during

cultivation and the robustness of eDNA biomonitoring compared to conventional approaches.

Fewer detections with eDNA sampling compared to conventional approaches may also

suggest that the water sampling volume used here might not be sufficient to capture broad

diversity (Cantera et al., 2019). Frequent clogging of filter papers from eDNA in turbid habitats,

such as wetlands, is a key limitation (Hinlo et al., 2017). Future studies may look to test for pre-

filtration (e.g., Turner et al. 2014), pooling of extractions from multiple smaller volume samples,

or implement passive filtration methods (Stevens et al., 2024). Moreover, during the harvesting

stage when less water is available, eDNA approaches may also create bias due to the increased

inhibitors within the water samples, putatively impacting amplification and thus biodiversity

recovery. Thus, additional soil and swabbing, or washing of rice plants for eDNA, could possibly

maximize detection. Still, our eDNA-based approach detected a broad range of taxonomic

groups with only limited sampling effort and without requiring extensive taxonomic expertise.

Thus, our method provides a fast and effective tool for the regular monitoring of native and non-

native pests in rice fields, and with developing technology, it will be easy to use for farmers.

Furthermore, incorporating multiple sample types (e.g., water, soil, leaf swabs), using group-

specific primers, and increasing the number of samples could further maximize the effectiveness

of this monitoring approach.

Conclusions

Environmental DNA metabarcoding is an effective tool in revealing the diversity and

composition of communities in rice fields across Taiwan. Through this study, a wide range of

taxa were detected in a limited number of sampling events, providing valuable insights into the

pest community. Three detected pests (*Chilo suppressalis*, *Coptotermes formosanus*, and

Pomacea canaliculata) are highly concerning. Notably, the presence of non-native *Pomacea*

canaliculata dominates over native snails, raising concerns about the impact on native

ecosystems and agricultural productivity. Furthermore, the detection of disease vectors like the

Japanese Encephalitis virus from the stagnant rice field water is a rising concern for human

health. These findings are particularly important for sensitive agricultural systems where

ecological and biological processes are becoming increasingly recognized as essential for

maintaining their ecological services. Overall, this study highlights the transformative potential

of eDNA metabarcoding for rapidly evaluating dynamic biological and environmental systems.

Caption of the Figures

Figure 1. Locations of eight eDNA sampling sites near Chiayi County, Taiwan (February and

May 2022)

Figure 2. Relative abundance of the top metazoan 10 taxa by field type. Stacked bar plots

showing the relative abundance (%) of the top 10 taxa detected in *harvesting* and *planting* field

types, displayed at four taxonomic levels: (A) orders, (B) families, (C) genera, and (D) species.

Figure 3. (A) Venn diagram of species detected in planting and harvesting fields. Circle and

overlap size denote proportional species diversity representation across field types. (B) Sankey

diagram showing the association between rice field types (planting and harvesting) and

detected species.

Figure 4. (A) Comparison of alpha diversity (observed ASVs) between field types (harvesting,

orange; planting, green) across sites (n=8). (B) Non-metric multidimensional scaling (NMDS)

plot based on Bray–Curtis dissimilarity showing beta diversity of communities between planting

(green) and harvesting (orange) rice fields detected from eDNA metabarcoding. Each point

represents a site, with ellipses indicating 95% confidence intervals for each field type.

Figure 5. Functional roles and species status across rice field types. (Left) Relative abundance

of species categorized by functional roles during harvesting and planting. (Right) Species

richness grouped by published literature and the Catalog of Life in Taiwan (https://taicol.tw)

(Table 4; native, non-native, and undermined/NA).

Caption of Tables

Table 1. Metadata of rice field samples collected at two crop stages. Planting-stage samples

(PF1–PF8) were collected on 10 February 2022, and harvesting-stage samples (HF1–HF8)

were collected on 31 May 2022, from four rice fields located in Chiayi County, Taiwan.

Coordinates for each sampling site are provided.

Table 2. Taxonomic distribution of amplicon sequence variants (ASVs) across major groups.

The table summarizes the total number of reads, ASV richness, and percentage of reads

assigned to each group. Unclassified reads with <80% similarity are denoted as "Below 80".

Table 3. Distribution of Animalia amplicon sequence variants (ASVs) across phyla. The table

shows the total number of reads, ASV richness, and relative percentage of reads assigned to

each phylum.

Table 4. Taxonomic composition, ecological roles, and native/non-native status of organisms

detected in Taiwan rice field ecosystems. Species are categorized by phylum, with ecological

roles spanning primary consumers, detritivores, predators, and omnivores; status and species

type (native, non-native, or not available - N/A) assigned based on references.

Table 5. The physiochemical properties and heavy metal concentration in agricultural soil during

harvesting and planting. p-value are from Welch's two-sample t-test ($\alpha=0.05$). a and b subscripts

represent significant differences.

Table 6. The physicochemical properties and heavy metal concentration of water at agricultural

fields during harvesting and planting. p-value are from Welch's two-sample t-test ($\alpha=0.05$). a

and b subscripts represent significant differences.

ACKNOWLEDGEMENTS

P.B. has been supported by Overseas Research Scholarships (ORS) from National Chung Cheng

University as well as the Ministry of Education (MOE)- Industry- Academia project (Taiwan).

The authors would also like to thank the Chiayi Christian Hospital (CYCH-CCU-2023-01) and

National Science and Technology Council (NSTC:112-2116-M-194- 010) for financial support.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Data sharing supporting the study is openly available in Dryad

(Dataset

http://datadryad.org/share/LINK_NOT_FOR_PUBLICATION/v8gjEC9xdWbgRxdx0GQ1BvlO

bvD6UjM8EyZWVZOFdvg)

References

Allaire, J. J., C. Gandrud, K. Russell, & C. Yetman, 2014. networkD3: D3 JavaScript Network

Graphs from R. , 0.4.1.

Baird, D. J., & M. Hajibabaei, 2012. Biomonitoring 2.0: a new paradigm in ecosystem

assessment made possible by next☐generation DNA sequencing. Molecular Ecology 21: 2039–

2044. https://doi.org/10.1111/j.1365-294X.2012.05519.x.

Bambaradeniya, C. N. B., & F. P. Amarasinghe, 2004. Biodiversity associated with the rice field

agroecosystem in Asian countries: a brief review. International Water Management Institute,

Colombo, Sri Lanka.

Banerjee, P., K. A. Stewart, G. Dey, R. K. Sharma, J. P. Maity, M. W. Y. Chan, K. P. Chang, T.-

H. Chen, C.-T. Hsu, & C.-Y. Chen, 2022. When conventional methods fall short: identification

of invasive cryptic Golden Apple Snails (Pomacea canaliculata; P. maculata) using

environmental DNA. Hydrobiologia 849: 4241–4257. https://doi.org/10.1007/s10750-022-

04979-6.

Barker, G. M., 2002. Molluscs as crop pests. CABI publ, Oxon.

Barnes, M. A., & C. R. Turner, 2016. The ecology of environmental DNA and implications for

conservation genetics. Conservation Genetics 17: 1–17. https://doi.org/10.1007/s10592-015-

0775-4.

Blumenfeld, A. J., P.-A. Eyer, C. Husseneder, J. Mo, L. N. L. Johnson, C. Wang, J. Kenneth

Grace, T. Chouvenc, S. Wang, & E. L. Vargo, 2021. Bridgehead effect and multiple

introductions shape the global invasion history of a termite. Communications Biology 4: 196.

https://doi.org/10.1038/s42003-021-01725-x.

Bolger, A. M., M. Lohse, & B. Usadel, 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30: 2114–2120. https://doi.org/10.1093/bioinformatics/btu170.

Bouyoucos, G. J., 1962. Hydrometer Method Improved for Making Particle Size Analyses of

Soils[1]. Agronomy Journal 54: 464–465.

https://doi.org/10.2134/agronj1962.00021962005400050028x.

Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, & S. P. Holmes,

2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods

13: 581–583. https://doi.org/10.1038/nmeth.3869.

Cantera, I., K. Cilleros, A. Valentini, A. Cerdan, T. Dejean, A. Iribar, P. Taberlet, R. Vigouroux,

& S. Brosse, 2019. Optimizing environmental DNA sampling effort for fish inventories in

tropical streams and rivers. Scientific Reports 9: 3085. https://doi.org/10.1038/s41598-019-

39399-5.

Chang, K. M., 1991. Catalogue of Fresh Water Shells of Taiwan.

Chen, J.H., Liao, Y.C., Chiu, M.C. and Yang, M.M., 2023. Arthropod communities of rice

agroecosystems can be shaped both by local agricultural practices and the surrounding

landscape. Frontiers in Environmental Science, 11, p.1221247.

Cheng, H.-J., F. Janssens, & C.-H. Chang, 2022. An updated checklist of Collembola in Taiwan,

with DNA barcoding of Papirioidesjacobsoni Folsom, 1924 (Symphypleona, Dicyrtomidae).

ZooKeys 1123: 123–146. https://doi.org/10.3897/zookeys.1123.90202.

Cheng, X., C. Chang, & S. Dai, 2010. Responses of striped stem borer, *Chilo suppressalis*

(Lepidoptera: Pyralidae), from Taiwan to a range of insecticides. Pest Management Science 66:

762–766. https://doi.org/10.1002/ps.1939.

Chu, Y. I., 1971. In Proceedings of the Symposium on Rice Insects. In Proceedings of the

Symposium on Rice Insects, Tokyo, Japan. pp. 19-24.

Dey, G., P. Banerjee, J. P. Maity, R. K. Sharma, G. Gnanachandrasamy, Y.-H. Huang, H.-B.

Huang, & C.-Y. Chen, 2022. Heavy metals distribution and ecological risk assessment including

arsenic resistant PGPR in tidal mangrove ecosystem. Marine Pollution Bulletin 181: 113905.

https://doi.org/10.1016/j.marpolbul.2022.113905.

Didaskalou, E. A., K. B. Trimbos, & K. A. Stewart, 2022. Environmental DNA. Current Biology

32: R1250–R1252. https://doi.org/10.1016/j.cub.2022.09.052.

Ding, T.-S., C.-S. Juan, R.-S. Lin, Y.-J. T. Tsai, J.-L. Wu, J. Wu, & Y.-H. Yang, 2020. The 2020

TWBF Checklist of the Birds of Taiwan. Bird Record Committee, Taiwan Wild Bird Federation,

Fl. 2, No.3, Ln. 50, Tacheng St., Datong Dist., Taipei City 103, Taiwan.

Flanagan, N. E., C. J. Richardson, & M. Ho, 2015. Connecting differential responses of native

and invasive riparian plants to climate change and environmental alteration. Ecological

Applications 25: 753–767. https://doi.org/10.1890/14-0767.1.

Flannagan, J. F., & K. E. Marshall, 2012. Advances in Ephemeroptera biology. Springer Science

& Business Media.

Hartmann, M., & J. Six, 2022. Soil structure and microbiome functions in agroecosystems.

Nature Reviews Earth & Environment 4: 4–18. https://doi.org/10.1038/s43017-022-00366-w.

Hinlo, R., D. Gleeson, M. Lintermans, & E. Furlan, 2017. Methods to maximise recovery of

environmental DNA from water samples. PLOS ONE 12: e0179251.

https://doi.org/10.1371/journal.pone.0179251.

Horgan, F. G., 2023. Integrated pest management in cereal cultivation. Burleigh Dodds Science

Publishing, Cambridge.

Hsing, Y. I. C., 2014. Rice in Taiwan. In Selin, H. (ed) Encyclopaedia of the History of Science,

Technology, and Medicine in Non-Western Cultures. Springer Netherlands, Dordrecht. pp. 1–3.

Hu, F.S., Arriaga-Varela, E., Biffi, G., Bocák, L., Bulirsch, P., Damaška, A.F., Frisch, J., Hájek,

J., Hlaváč, P., Ho, B.H. and Ho, Y.H., 2024. Forest leaf litter beetles of Taiwan: first DNA

barcodes and first insight into the fauna. Deutsche Entomologische Zeitschrift, 71(1), pp.17-47.

Huang, S.-H., C.-H. Cheng, C.-N. Chen, W.-J. Wu, & A. Otuka, 2010. Estimating the

immigration source of rice planthoppers, Nilaparvata lugens (Stål) and Sogatella furcifera

(Horváth) (Homoptera: Delphacidae), in Taiwan. Applied Entomology and Zoology 45: 521–

531. https://doi.org/10.1303/aez.2010.521.

Jang-Liaw, N.-H., & W.-H. Chou, 2015. Anuran Fauna of Taiwan and Adjacent Islands Based

on Valid Specimen Records.

Jiang, X., P. Zheng, I. Soto, P. J. Haubrock, J. Chen, & L. Ji, 2022. Global economic costs and

knowledge gaps of invasive gastropods. Ecological Indicators 145: 109614.

https://doi.org/10.1016/j.ecolind.2022.109614.

Katayama, N., S. Yamamoto, Y. G. Baba, K. Ito, & J. Yamasako, 2024. Complementary role of

environmental DNA for line-transect bird surveys: A field test in a Japanese rice landscape.

Ecological Indicators 166: 112442. https://doi.org/10.1016/j.ecolind.2024.112442.

Kestel, J. H., D. L. Field, P. W. Bateman, N. E. White, M. E. Allentoft, A. J. M. Hopkins, M.

Gibberd, & P. Nevill, 2022. Applications of environmental DNA (eDNA) in agricultural

systems: Current uses, limitations and future prospects. Science of The Total Environment 847:

157556. https://doi.org/10.1016/j.scitotenv.2022.157556.

Korbel, K. L., G. C. Hose, C. Karwautz, P. Greenfield, H. Wang, A. A. Chariton, & C. Griebler,

2024. Detection, movement and persistence of invertebrate eDNA in groundwater. Scientific

Reports 14: 17151. https://doi.org/10.1038/s41598-024-67349-3.

Lee, K.-H., T.-H. Chen, G. Shang, S. Clulow, Y.-J. Yang, & S.-M. Lin, 2019. A check list and

population trends of invasive amphibians and reptiles in Taiwan. ZooKeys 829: 85–130.

https://doi.org/10.3897/zookeys.829.27535.

Li, H.-F., W. Ye, N.-Y. Su, & N. Kanzaki, 2009. Phylogeography of *Coptotermes gestroi* and

Coptotermes formosanus (Isoptera: Rhinotermitidae) in Taiwan. Annals of the Entomological

Society of America 102: 684–693. https://doi.org/10.1603/008.102.0413.

Llanos, J., H. Hipperson, G. Horsburgh, M. G. Lappage, K. H. Maher, T. Burke, J. R. Leake, &

P. J. Watt, 2025. Environmental DNA is more effective than hand sorting in evaluating

earthworm biodiversity recovery under regenerative agriculture. Science of The Total

Environment 968: 178793. https://doi.org/10.1016/j.scitotenv.2025.178793.

Longmire, J. L., M. Maltbie, R. J. Baker, & Texas Tech University., 1997. Use of "Lysis Buffer"

in DNA isolation and its implication for museum collections. Museum of Texas Tech University,

Lubbock, TX.

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 17: 10. https://doi.org/10.14806/ej.17.1.200.

McCartin, L.J., Vohsen, S.A., Ambrose, S.W., Layden, M., McFadden, C.S., Cordes, E.E.,

McDermott, J.M. and Herrera, S., 2022. Temperature controls eDNA persistence across

physicochemical conditions in seawater. Environmental Science & Technology, 56(12),

pp.8629-8639.

Melcher, A. C., S. Weber, K. Birkhofer, D. Harms, & H. Krehenwinkel, 2024. To pool or not to

pool: Pooled metabarcoding does not affect estimates of prey diversity in spider gut content

analysis. Ecological Entomology 49: 768–778. https://doi.org/10.1111/een.13382.

Mukherjee, P., & M. Khan, 2017. Abundance of arthropod insect pests and natural enemies in

rice field as influenced by rice growth stages and neighboring crops. Bangladesh Journal of

Agricultural Research 42: 309–319. https://doi.org/10.3329/bjar.v42i2.32817.

Oksanen, J., G. L. Simpson, F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara,

P. Solymos, M. H. H. Stevens, E. Szoecs, H. Wagner, M. Barbour, M. Bedward, B. Bolker, D.

Borcard, T. Borman, G. Carvalho, M. Chirico, M. De Caceres, S. Durand, H. B. A. Evangelista,

R. FitzJohn, M. Friendly, B. Furneaux, G. Hannigan, M. O. Hill, L. Lahti, C. Martino, D.

McGlinn, M.-H. Ouellette, E. Ribeiro Cunha, T. Smith, A. Stier, C. J. F. Ter Braak, & J.

Weedon, 2001. vegan: Community Ecology Package. , 2.7-1.

Otuka, A., S.-H. Huang, S. Sanada-Morimura, & M. Matsumura, 2012. Migration analysis of

Nilaparvata lugens (Hemiptera: Delphacidae) from the Philippines to Taiwan under typhoon-

induced   windy   conditions.   Applied   Entomology   and   Zoology   47:   263–271.

https://doi.org/10.1007/s13355-012-0118-6.

R Core Team, 2025. R: A Language and Environment for Statistical Computing.

Saksongmuang, V., R. Michalko, B. Petcharad, & S. Bumrungsri, 2024. Changes in community

composition and prey capture of web-building spiders during rice field development. Basic and

Applied Ecology 79: 29–37. https://doi.org/10.1016/j.baae.2024.05.003.

Sayers, E. W., M. Cavanaugh, K. Clark, K. D. Pruitt, C. L. Schoch, S. T. Sherry, & I. Karsch-

Mizrachi,   2022.   GenBank.   Nucleic   Acids   Research   50:   D161–D164.

https://doi.org/10.1093/nar/gkab1135.

Schenekar, T., M. Schletterer, L. A. Lecaudey, & S. J. Weiss, 2020. Reference databases, primer

choice, and assay sensitivity for environmental metabarcoding: Lessons learnt from a

re□evaluation of an eDNA fish assessment in the Volga headwaters. River Research and

Applications 36: 1004–1013. https://doi.org/10.1002/rra.3610.

Seymour, M., F. K. Edwards, B. J. Cosby, I. Bista, P. M. Scarlett, F. L. Brailsford, H. C.

Glanville, M. De Bruyn, G. R. Carvalho, & S. Creer, 2021. Environmental DNA provides higher

resolution assessment of riverine biodiversity and ecosystem function via spatio-temporal

nestedness   and   turnover   partitioning.   Communications   Biology   4:   512.

https://doi.org/10.1038/s42003-021-02031-2.

Sigsgaard, E. E., K. Olsen, M. D. D. Hansen, O. L. P. Hansen, T. T. Høye, J. Svenning, & P. F.

Thomsen, 2021. Environmental DNA metabarcoding of cow dung reveals taxonomic and

functional diversity of invertebrate assemblages. Molecular Ecology 30: 3374–3389.

https://doi.org/10.1111/mec.15734.

Simberloff, D., & M. Rejmanek (eds), 2019. 100 of the World's Worst Invasive Alien Species: A

Selection From The Global Invasive Species Database. In Encyclopedia of Biological Invasions.

University of California Press. pp. 715–716.

Song, J., & C. Kuo, 2022. Farming practice affects rice field animal biota during cultivation but

not fallow periods in Taiwan. Ecosphere 13: e4069. https://doi.org/10.1002/ecs2.4069.

Sottomayor, M., Palmeirim, A.F., Meyer, C.F., De Lima, R.F., Rocha, R. and Rainho, A., 2024.

Nature-based solutions to increase rice yield: An experimental assessment of the role of birds

and bats as agricultural pest suppressors in West Africa. Agriculture, Ecosystems &

Environment, 370, p.109067.

Spens, J., A. R. Evans, D. Halfmaerten, S. W. Knudsen, M. E. Sengupta, S. S. T. Mak, E. E.

Sigsgaard, & M. Hellström, 2017. Comparison of capture and storage methods for aqueous

macrobial EDNA using an optimized extraction protocol: advantage of enclosed filter. Methods

in Ecology and Evolution 8: 635–645. https://doi.org/10.1111/2041-210X.12683.

Stevens, E. R., J. Hyde, L. S. Beesley, D. C. Gwinn, S. Thompson, L. Morris, P. R. Wilson, & D.

B. Gleeson, 2024. Fishy Business—Assessing the Efficacy of Active and Passive EDNA to

Describe the Fish Assemblage of a River in Southwestern Western Australia to Support Effective

Monitoring. Environmental DNA 6: e70040. https://doi.org/10.1002/edn3.70040.

Stewart, K. A., 2019. Understanding the effects of biotic and abiotic factors on sources of aquatic

environmental DNA. Biodiversity and Conservation 28: 983–1001.

https://doi.org/10.1007/s10531-019-01709-8.

Taberlet, P., E. Coissac, M. Hajibabaei, & L. H. Rieseberg, 2012. Environmental DNA.

Molecular Ecology 21: 1789–1793. https://doi.org/10.1111/j.1365-294X.2012.05542.x.

Tilman, D., K. G. Cassman, P. A. Matson, R. Naylor, & S. Polasky, 2002. Agricultural

sustainability and intensive production practices. Nature 418: 671–677.

https://doi.org/10.1038/nature01014.

Triapitsyn, S. V., H.-T. Shih, S.-H. Huang, & M.-J. Tseng, 2021. Identification of egg parasitoids

of rice leafhoppers and planthoppers (Hemiptera: Cicadellidae and Delphacidae) of economic

importance in Taiwan, part 1: Mymaridae (Hymenoptera). Journal of Asia-Pacific Entomology 24: 77–90. https://doi.org/10.1016/j.aspen.2020.11.002.

Tsai, C.-F., H.-P. Shen, & S.-C. Tsai, 2000. Native and Exotic Species of Terrestrial Earthworms (Oligochaeta) in Taiwan with Reference to Northeast Asia 4: 285–294.

Tsai, C.-L., I.-H. Chu, M.-H. Chou, T. Chareonviriyaphap, M.-Y. Chiang, P.-A. Lin, K.-H. Lu, & W.-B. Yeh, 2020. Rapid identification of the invasive fall armyworm Spodoptera frugiperda (Lepidoptera, Noctuidae) using species-specific primers in multiplex PCR. Scientific Reports 10: 16508. https://doi.org/10.1038/s41598-020-73786-7.

Turner, C.R., Barnes, M.A., Xu, C.C.Y., Jones, S.E., Jerde, C.L., & Lodge, D.M. 2014. Particle size distribution and optimal capture of aqueous macrobial eDNA. Methods in Ecology and Evolution 5: 676–684.

Ushio, M., H. Saito, M. Tojo, & A. J. Nagano, 2023. An ecological network approach for detecting and validating influential organisms for rice growth. eLife 12: RP87202. https://doi.org/10.7554/eLife.87202.3.

Valentini, A., P. Taberlet, C. Miaud, R. Civade, J. Herder, P. F. Thomsen, E. Bellemain, A. Besnard, E. Coissac, F. Boyer, C. Gaboriaud, P. Jean, N. Poulet, N. Roset, G. H. Copp, P. Geniez, D. Pont, C. Argillier, J. Baudoin, T. Peroux, A. J. Crivelli, A. Olivier, M. Acqueberge, M. Le Brun, P. R. Møller, E. Willerslev, & T. Dejean, 2016. Next□generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. Molecular Ecology 25: 929–942. https://doi.org/10.1111/mec.13428.

Valls, L., J. Rueda, & F. Mesquita-Joanes, 2014. Rice fields as facilitators of freshwater

invasions in protected wetlands: the case of Ostracoda (Crustacea) in the Albufera Natural Park

(E Spain). Zoological Studies 53: 68. https://doi.org/10.1186/s40555-014-0068-5.

Vastano, A. R., M. C. Allen, C. J. Penca, O. C. Stringham, M. J. Domingue, & J. L. Lockwood,

2024. Can environmental DNA be used within pest insect agricultural biosecurity? Detecting

khapra beetle within stored rice. Environmental DNA 6: e585. https://doi.org/10.1002/edn3.585.

Watanabe, K., S. Koji, K. Hidaka, & K. Nakamura, 2013. Abundance, Diversity, and Seasonal

Population Dynamics of Aquatic Coleoptera and Heteroptera in Rice Fields: Effects of Direct

Seeding Management. Environmental Entomology 42: 841–850.

https://doi.org/10.1603/EN13109.

Wickham, H., 2016. Data Analysis. In ggplot2. Springer International Publishing, Cham. pp.

189–201.

Wu, J.-Y., P.-J. Meng, M.-Y. Liu, Y.-W. Chiu, & L.-L. Liu, 2010. A High Incidence of Imposex

in Pomacea Apple Snails in Taiwan: A Decade after Triphenyltin Was Banned. Zoological

Studies 49(1): 85–93.

Yan, M. L., 2021. ggvenn: Draw Venn diagram by'ggplot2'.

Yates, M.C., Gaudet☐Boulay, M., Garcia Machado, E., Côté, G., Gilbert, A. and Bernatchez, L.,

2023. How much is enough? Examining the sampling effort necessary to estimate mean eDNA

concentrations in lentic systems. Environmental DNA, 5(6), pp.1527-1540.

Table 1. Metadata of rice field samples collected at two crop stages. Planting-stage samples (PF1–PF8)
were collected on 10 February 2022 and harvesting-stage samples (HF1–HF8) were collected on 31 May
2022, from four rice fields located in Chiayi County, Taiwan. Coordinates for each sampling site are
provided.

Sample name	Field type	Location	Date
PF1	planting	23.567737, 120.454625	10-Feb-22
PF2			
PF3		23.540607, 120.380341	
PF4			
PF5		23.567776, 120.454035	
PF6			
PF7		23.543891, 120.384877	
PF8			
HF1	harvesting	23.567737, 120.454625	31-May-22
HF3		23.540607, 120.380341	
HF4			
HF5		23.567776, 120.454035	
HF6			
HF7		23.543891, 120.384877	
HF8			

725
726

727 **Table 2.** Taxonomic distribution of amplicon sequence variants (ASVs) across major groups. The table
728 summarizes the total number of sequencing reads, ASV richness, and percentage of reads assigned to
729 each group. The majority of reads were assigned to Animalia, followed by unclassified reads with <80%
730 similarity ("Below 80"), Protozoa, Bacteria, Algae, and Fungi.

Phylum	Total reads	ASVs	Percentage of reads
Animalia	557061	80	54.92
Below 80	242931	2	23.95
Protozoa	123431	25	12.16
Bacteria	43090	13	4.24
Algae	33448	38	3.29
Fungi	14264	38	1.40

731
732

Table 3. Distribution of Animalia amplicon sequence variants (ASVs) across phyla. The table shows the total number of sequencing reads, ASV richness, and relative percentage of reads assigned to each phylum. Arthropoda was the dominant phylum, followed by Annelida and Rotifera, while other phyla were represented at much lower abundances.

Phylum	Total reads	ASVs	Percentage of reads
Arthropoda	518717	43	93.12
Annelida	31958	11	5.73
Rotifera	3257	8	0.58
Mollusca	1614	5	0.28
Gastrotricha	779	2	0.13
Cnidaria	355	2	0.06
Nematoda	108	2	0.01
Chordata	53	4	~0.01

Table 4. Taxonomic composition, ecological roles, and native/non-native status of organisms detected in rice field ecosystems. Species are categorized by phylum, with ecological roles spanning primary consumers, detritivores, predators, and omnivores. Status is assigned based on literature references, indicating whether species are native, non-native, or not available (N/A).

Phylum	Species	Ecological role	Status
Annelida	*Metaphire posthuma*	Detritivores	Non-native (Tsai et al., 2000)
	Allonais sp.		N/A
	Bothrioneurum vejdovskyanum		N/A
	Dero sp.		N/A
	Pristina sp.		N/A
Arthropoda	*Ceriodaphnia cornuta*	Primary consumer	N/A
	Daphnia sp.	Primary consumer	N/A
	Moina macrocopa	Primary consumer	N/A
	Homidia socia	Detritivore	Native (Cheng et al., 2022)
	Thermocyclops	Secondary consumer	Native (Reid,

	decipiens		1989)
	Thermocyclops taihokuensis	Secondary consumer	Native (Reid, 1989)
	Coptotermes formosanus	Detritivores	Native (Blumenfeld et al., 2021)
	Tanypus kraatzi	Detritivores, and Omnivorous	NA
	Tanytarsus formosanus	Detritivores, and Omnivorous	Native (Gao et al., 2024)
	Culex tritaeniorhynchus	detritivore/filter-feeder (larva) and hematophagous (adult)	Native (Chung et al., 2024)
	Psilopa sp.	Detritivores	N/A
	Sciara humeralis	Detritivores (Larva), feed on nectar or fungi (adult)	Native
	Diplonychus esakii	Predator	Native (Rieger, 1995)
	Microvelia douglasi	Predator	Native (Ye et al., 2014)
	Monomorium pharaonis	Omnivorous	Native (Wetterer, 2010)
	Pheidole sp.	Omnivorous	N/A
	Chilo suppressalis	Primary consumer (Rice crop pest)	Native (Chu, 1971)
Chordata	*Microhyla fissipes*	Predator	Native (Jang-Liaw & Chou, 2015)
	Fejervarya sp.	Predator	N/A
	Passer montanus	Omnivorous	Native (Ding et al., 2020)
	Acridotheres tristis	Omnivorous	Non-native (Ding et al., 2020)
Gastrotricha	*Chaetonotus* sp.	Detritivores, microphagus	N/A
Mollusca	*Pomacea canaliculata*	Detritivorous herbivores. Primary consumer, Rice crop pest	Non-native (Banerjee et al., 2024)
	Austropeplea ollula	Detritivorous herbivores. Primary consumer	Native (Chang, 1991)

	Melanoides tuberculata	Detritivorous herbivores. Primary consumer	Native (Chang, 1991)
Rotifera	*Brachionus* sp.	Detritivores	N/A
	Keratella sp.		
	Lecane sp.		
	Polyarthra sp.		
	Rotaria sp.		

Table 5. The physiochemical properties and heavy metal concentration in agricultural soil during harvesting and planting. p-value are from Welch's two-sample t-test (α=0.05). a and b subscripts represent significant differences.

Parameter	Harvesting (Mean ± SE)	Planting (Mean ± SE)	p (Welch t-test)
pH	7.74 ± 0.05 [a]	7.54 ± 0.07 [b]	**0.0394**
Silt (%)	14.53 ± 1.69 [a]	15.15 ± 1.59 [a]	0.8075
Clay (%)	13.10 ± 1.65 [a]	12.80 ± 1.47 [a]	0.9016
Sand (%)	72.37 ± 1.63 [a]	72.05 ± 1.46 [a]	0.8937
Cd (mg/kg)	0.44 ± 0.07 [a]	0.30 ± 0.06 [a]	0.1694
Co (mg/kg)	2.41 ± 0.57 [a]	2.02 ± 0.35 [a]	0.6029
Cr (mg/kg)	70.94 ± 6.48 [a]	64.66 ± 7.12 [a]	0.5536
Cu (mg/kg)	19.40 ± 1.77 [a]	16.46 ± 2.10 [a]	0.3379
Fe (mg/kg)	29781.25 ± 2729.60 [a]	24943.06 ± 2848.57 [a]	0.2743
Li (mg/kg)	37.34 ± 2.49 [a]	37.73 ± 1.77 [a]	0.9074
Mn (mg/kg)	256.67 ± 32.80 [a]	243.13 ± 36.91 [a]	0.8026
Ni (mg/kg)	35.11 ± 2.85 [a]	32.98 ± 3.93 [a]	0.6899
Pb (mg/kg)	35.81 ± 1.39 [a]	29.04 ± 2.49 [b]	**0.0492**
Sr (mg/kg)	269.93 ± 34.18 [a]	306.87 ± 27.48a	0.4489
Zn (mg/kg)	92.91 ± 9.04 [a]	86.12 ± 7.95a	0.6092
Ag (mg/kg)	1.26 ± 0.15 [a]	1.14 ± 0.21a	0.6702

Table 6. The physicochemical properties and heavy metal concentration of water at agricultural fields during harvesting and planting. p-value are from Welch's two-sample t-test (α=0.05). a and b subscripts represent significant differences.

Parameter	Harvesting (Mean ± SE)	Planting (Mean ± SE)	p (Welch t-test)
pH	7.52 ± 0.06 [a]	7.54 ± 0.05 [a]	0.7818
mV pH	-21.48 ± 2.63 [a]	-26.70 ± 2.52 [a]	0.2059
ORP	152.37 ± 11.67 [b]	188.99 ± 3.77 [a]	**0.0270**
DO (mg/L)	4.45 ± 0.02 [b]	5.27 ± 0.26 [a]	**0.0197**
Conductivity (mS/cm)	0.40 ± 0.04 [b]	0.55 ± 0.05 [a]	**0.0363**
Resistivity (ohm·cm)	3070.52 ± 382.50 [a]	2111.05 ± 220.36 [a]	0.0723
TDS (mg/L)	187.08 ± 20.39 [b]	259.80 ± 22.12 [a]	**0.0424**
PSU	0.26 ± 0.02 [a]	0.25 ± 0.02 [a]	0.7992
Temperature (°C)	29.32 ± 0.88 [a]	26.08 ± 0.76 [b]	**0.0231**
Atm. Pressure (PSI)	14.57 ± 0.10 [a]	14.76 ± 0.02 [a]	0.1412
Fe (mg/L)	0.15 ± 0.07 [a]	0.03 ± 0.01 [ab]	0.1676
Li (mg/L)	0.27 ± 0.00 [a]	0.26 ± 0.00 [b]	**0.0082**
Mn (mg/L)	0.31 ± 0.10 [a]	0.24 ± 0.11 [a]	0.6609
Sr (mg/L)	3.90 ± 0.87 [a]	6.61 ± 1.09 [a]	0.0938

757
758