Title: Colony-level pollen collection reflects visitation of managed bumble bees (*Bombus impatiens*) in strawberry fields and surrounding landscapes without reducing pollen limitation.

Authors & Affiliations

Leeah I Richardson[1], Olivia Miller[2], David Sossa[2], Aaron Iverson[3], Scott McArt[2], Katja Poveda[2], and Heather Grab[4]

[1] University of Texas at Austin, Department of Integrative Biology

[2] Cornell University, Department of Entomology

[3] St. Lawrence University, Department of Environmental Studies

[4] Department of Entomology, The Pennsylvania State University, University Park, PA 16802

Corresponding author: Leeah I Richardson; Email: LeeahRichardson@utexas.edu

ORCiD:

Leeah I Richardson: https://orcid.org/0000-0002-2118-8765

Olivia Miler: https://orcid.org/0009-0007-7217-5095

David Sossa: https://orcid.org/0009-0000-9069-9107

Aaron Iverson: https://orcid.org/0000-0001-6343-4226

Scott McArt: https://orcid.org/0000-0001-7157-9011

Katja Poveda: https://orcid.org/0000-0002-5222-6947

Heather Grab: https://orcid.org/0000-0002-1073-8805

Acknowledgements:

We thank the growers that allowed us access to their fields, as well as to Kass Urban-Mead and Diana Obregon for pollen staining advice, and to Nate Flicker for help with data collection. This work was supported by the USDA NIFA Hatch Grant NYC-139303, USDA NIFA 2018-08603 and Hatch Grant NYC-139440.

Data Sharing & Availability:

Data used in this manuscript will be made available from the Dryad Digital Repository upon publication.

Conflict of Interest statement:

We have no conflict of interest to declare.

Abstract
Managed bees are frequently used to supplement pollination of a range of crop plants. Yet, their
effectiveness varies and much of their foraging can be outside of the focal crop. Bee foraging
behavior is highly sensitive to the availability of alternative floral resources which depend on the
composition of the surrounding landscape. To understand how foraging and crop pollination
dynamics of the managed common eastern bumble bee (*Bombus impatiens*) are influenced by
landscape composition, we assessed their visitation and pollination in cultivated strawberry
fields along a landscape gradient. We also used pollen extracted from bumble bee (*Bombus*
impatiens) colony wax to understand whole-colony foraging over the blooming period. Bumble
bee visits to strawberry crop flowers were very low and pollen limitation was not reduced at sites
with greater visitation; however, colonies in fields with higher visitation rates had more pollen
from the Rosaceae family (including strawberry). We found that landscape composition affected
foraging effort on strawberry at the colony level, as a lower proportion of rosaceous pollen was
collected in landscapes with more surrounding pasture, while a greater proportion of rosaceous
pollen was collected at sites with larger strawberry fields. Overall, our results suggest that
managed bumble bees do forage on strawberry crop flowers, but they predominantly forage
elsewhere, especially in surrounding pastures when strawberry fields are small. Therefore, the
addition of bumble bee colonies may not be economically beneficial for strawberry growers at
least in diverse agricultural landscapes similar to that of our study region.
53

Introduction
Pollinators serve an important role in food production, with about 30% of agricultural food
production relying to some extent on animal mediated pollination (Klein *et al.*, 2007; Khalifa *et*
al., 2021). Yet, agriculture itself is a major driver of pollinator decline (Cane & Tepedino, 2001)
due to factors like habitat loss and exposure to pesticides (Goulson *et al.*, 2015) leading to
reductions in the services provided by wild bees frequently observed in landscapes that are
more agriculturally dominated (Ricketts *et al.*, 2018; Grab *et al.*, 2019). To achieve sufficient
pollination, growers can bring in managed bees to supplement natural pollination (Rucker *et al.*,
2012). However, supplemental managed bees may not always improve yields (Petersen *et al.*,
2013) and factors that moderate their efficacy are unclear.
64
Supplementing crop pollination with managed pollinators does not always increase pollination
rates in the focal crop (Petersen *et al.*, 2013; Trillo *et al.*, 2018) for two potential reasons. First,
in landscapes with high amounts of natural habitat, wild pollinators are often more abundant and

more likely to meet crop pollination demands on their own (Garibaldi *et al.*, 2011; Kennedy *et*
al., 2013; Mallinger & Gratton, 2015). Second, in environments with abundant natural habitats,
managed pollinators may preferentially forage in these areas and overlook the focal crop
(Guzman *et al.*, 2019). For example, in cranberry agroecosystems honey bee stocking density
was found to have a strong positive relationship with yield in fields that have low forest cover but
no relationship to yield in landscapes with high forest cover (Gaines-Day & Gratton, 2016),
where around only 15% of pollen from hives was from cranberry (Guzman *et al.*, 2019). These
low rates of focal crop pollen collection by managed pollinators have been reported from a
range of other crop systems (8.7% apple (McArt *et al.*, 2017a), 0.4-2% pumpkin (Brochu *et al.*,
2020), 19-26% strawberry (Bänsch *et al.*, 2020), 0.5-26% blueberry (Graham *et al.*, 2023)), and
suggest that managed bees are frequently using resources outside of the focal crop. For these
reasons, we expect managed bees to be more effective in simple, agriculturally dominated
landscapes than in landscapes containing more natural and semi-natural habitats.
81
The honey bee (*Apis mellifera*) is not the only managed bee that can be used to supplement pollination
services (Osterman *et al.*, 2021). The common eastern bumble bee (*Bombus impatiens*) is widely used
across the eastern United States as a commercially available alternative to honey bees for supplemental
pollination services, as they are generalist foragers that are capable of pollinating many different crops
(Evans, 2017). While these colonies are primarily used for pollination within greenhouses (Evans, 2017),
they are also sold to supplement wild pollinators for crops grown in fields (Stanghellini et al., 1998;
Desjardins & De Oliveira, 2006; Artz & Nault, 2011). Though bumble bee colonies have fewer
individuals than honey bees, they can be more effective at pollinating certain crops (ex. blueberry,
tomatoes, etc) due to traits like hairiness (pilosity) (Stavert et al., 2016), tolerance to foraging in colder
temperatures (Corbet *et al.,* 1993), and the ability to buzz pollinate (De Luca & Vallejo-Marín, 2013).
They can also have a higher per-visit efficiency and colonies (which only live one season) can be
purchased for prices comparable to or even less expensive than renting commercial honey bees for one
season of pollination (Drummond, 2012).
95
Strawberries are a crop that has been shown to benefit from bumble bee pollination in
greenhouses and other enclosed environments (Dimou *et al.*, 2008; Martin *et al.*, 2019).
However, the majority of strawberry production occurs in open field conditions (*USDA NASS,*
2022). Insect pollination generally benefits strawberry crops (Gudowska *et al.*, 2024; Menzel,
2023). However, adding managed honey bee colonies to strawberry plantings has been found
to negatively affect yield (Angelella et al. 2021), while increased wild bee abundance is

associated with improved yield (Connelly *et al.,* 2015). In our study region, the most common
pollinators to strawberry during the early spring are bees in the genus *Andrena, Lasioglossum,*
and *Augochlorella;* followed by honey bees, which were found to make up less than 7% of the
pollinator community (Connelly *et al.,* 2015). This is similar to the composition detected in
another study region, also in eastern North America, although it was less dominated by honey
bees (MacInnis & Forrest, 2019). Bumble bees are considered effective pollinators generally
based on their size and loosely packed pollen (Nye & Anderson, 1974), while managed bumble
bees (*B. terrestris*) have been shown to dedicate much of their foraging effort to strawberry
plantings in Spain (Trillo *et al.*, 2020).

In the northeast United States, wild bumble bee foundress queens begin to start colonies during
strawberry bloom from May to early June, but are not established to produce foraging workers
until after the bloom of spring strawberry has completed (Lanterman *et al.*, 2019). Indeed,
previous work suggests that bumble bees do not visit spring strawberry plantings in New York at
sites where they are not actively managed (Connelly *et al.*, 2015). This suggests the potential
for commercially available bumble bee colonies to provide supplemental services, particularly in
agriculturally dominated landscapes where pollination services from wild bees may be lower
(Connelly *et al.*, 2015).

Understanding which floral resources bumble bees are foraging on by identifying pollen grains
offers an important insight into their foraging patterns (Guzman *et al.*, 2019). It is typically
difficult to get this colony level information because floral usage in bumble bees varies between
individuals (Saifuddin & Jha, 2014), across the season (Wood *et al.*, 2018), and even over the
course of a single day (Inouye, 1978). This means that short sampling periods are unlikely to
capture the true breadth of floral resource use, and it is typically impractical to look at the pollen
loads of every returning forager to a colony. However, techniques can be adapted for extracting
pollen from beeswax (Furnessa, 1994) to look at the pollen brought back at the colony level,
which allows integration over time and across individuals with a single sample from a colony.
These methods have recently been adapted to look at the pollen found in medieval wax seals
(Kasso *et al.*, 2023), and to understand historic honey bee foraging patterns from museum
comb specimens (Kasso *et al.*, 2024); their use in investigating variation in contemporary
bumble bee foraging offers a promising new direction to understanding season-long colony level
foraging efforts.

In this study, we assessed the degree to which supplemental managed colonies of *B. impatiens* visit strawberry plantings in open field settings during peak bloom. Specifically, we focus our analyses on four questions: (1) Does managed bumble bee visitation rate to strawberry depend on surrounding land cover? (2) Do fields with greater visitation by managed bumble bees have reduced pollen limitation? (3) Is visitation rate by managed bumble bees reflected in colony Rosaceae pollen collection? and (4) Is pollen collection by managed bumble bees is dependent on landscape context?

Methods

Landscape composition and colony placements:

In the spring of 2018, we surveyed 10 strawberry fields across upstate New York (Fig. 1A) that were a minimum of 500 m apart, placing commercially available bumble bee (Bombus impatiens) colonies from Koppert Biological Systems (MI, USA) in or along the edge of each field. Colonies were placed in the fields between May 14–15th (all within a 24 hour period), just prior to strawberry bloom, and remained in the fields until fruit had set in mid June. Fields ranged in size from 0.069 hectares to 3.035 hectares. The number of colonies placed in each field ranged from 1–16 depending on field size, such that stocking densities were maintained at approximately two colonies per acre (0.4 ha) as recommended by Koppert Biological Systems, for a total of 43 colonies across all sites.

We determined the landscape composition around each field using the 2018 Cropland Data Layer (https://nassgeodata.gmu.edu/CropScape/). The proportion of each land cover category was calculated at 500, 750 and 1000 m through ArcGIS 10.3 (ESRI 2018) for three different land cover types: agricultural (cultivated crops; ranging from 5.9%-86.1% at 1000 m), semi-natural (forests, shrublands, wetlands; ranging from 2.4%-24.4% at 1000 m), and pasture/grass/hay (alfalfa, meadows, clover, wildflowers, grasses; ranging from 5.8% to 30.0% at 1000 m), hereafter referred to as 'pasture' (Figure 1).

Bumble bee visitation:

We conducted visual surveys to determine the visitation rate of bumble bees to strawberry during the blooming period. Four times during strawberry bloom (May 16th - June 7th) we walked a 10 min transect in each field at a consistent slow pace and counted each bee that visited a flower. Visits were defined as contact with the flower petals or stigmas (Grab *et al.*, 2018), and we visually identified each bee as either a bumble bee or any other type of bee (clade: Anthophila), which were primarily bees in the genera *Andrena, Lasioglossum,*

Augochlorella, and *Apis.* Because wild bumble bee workers were not yet foraging in this region

during this time period (Lanterman *et al.*, 2019) and previous studies in the region reported no

bumble bee visitors to strawberry at fields without managed colonies (Connelly *et al.*, 2015), we

assumed that all bumble bee visits during our sampling periods were from our managed *B.*

impatiens colonies.

Sentinel plants for assessing pollination:

We started bare-root strawberry plants of the cultivar 'Seascape' (Indiana Berry, IN, USA) in 10

cm pots in a greenhouse approximately 5 weeks before the onset of spring field strawberry

bloom to use as sentinel plants flowering simultaneously with field-grown plants. Most of our 10

sites received 24 plants, although one site only received 18. Strawberry flowers tend to bifurcate

from the stem internodes such that one large "primary" flower head is on the same stem as two

smaller paired "secondary" flowers that typically open on the same day and will lead to fruits of

similar sizes (Petran & Hoover, 2018). We removed the primary flowers from our plants, leaving

the paired secondary flower buds, and dried the anthers from these primary flowers until they

dehisced. On the day of secondary flower anthesis we used pollen collected from primary

flowers to hand-pollinate one of the secondary flowers by gently brushing pollen across all

stigmatic surfaces with a small paint brush (hand-pollinated) while we left the other secondary

flower to be pollinated in the field (open-pollinated). These were marked with a colored piece of yarn tied

loosely around the base of each flower (Fig. 1C).

Plants were brought out to the fields on the same day that they were hand-pollinated and placed in trays

that held ~3 cm of water in groups of 6 within the field rows when the field was flowering. Sentinel plants

were only being open-pollinated when the field strawberries were also blooming and were present

throughout the blooming period as we brought out trays on four dates throughout the season (first trays of

plants to fields from May 14th - May 25th and the final plants were brought to fields May 31st - June

7th). We watered the plants during all field visits, and brought them back to be cared for in the

greenhouse when the flowers had senesced (June 6th - June 22nd). Once the fruits had fully developed

(June 7th - July 5th) we weighed each as a measure of pollination quality (Gudowska et al.,

2024), and calculated the degree of pollen limitation in the field as the weight of the open-

pollinated fruit subtracted from the hand-pollinated fruit then divided by the weight of the hand-

pollinated fruit for each plant (Larson & Barrett, 1999; Baskin & Baskin, 2018). This allowed us

to understand how close the open-pollinated fruits were to being fully pollinated (represented by

the hand-pollinated treatment) to quantify pollen limitation in this system. If we detected no

difference between these treatments it would indicate that pollination was sufficient, due to animal-

mediated pollination and/or self pollination.

We were able to retrieve paired fruits from 154 plants for an average (+/- standard deviation) of

15.40 +/- 4.64 plants per field. Plants or their fruits that were lost were due primarily to herbivory

or disease, which was distributed fairly evenly across fields (75 total plants were excluded from

analyses: average 7.5 plants +/- 4.67 SD per site).

Wax and pollen extraction:

At the end of strawberry bloom, we removed approximately 10 g of wax from a single randomly

selected representative colony from each field to quantify pollen composition such that each site

was represented by one colony. Because each colony grows over the course of the season, old

wax is located in the center of the colony (Sakagami *et al.*, 1967) and is much darker in color

(Michener, 1974) so it is possible to only remove newer wax that had built up along the entire

periphery after the colony was put into the strawberry fields. We expected the pollen present in

the wax to be representative of pollen collection over time by random contamination into the

wax during larval feeding and wax manipulation for construction (Hines *et al.*, 2007). We

pulverized the wax samples from these representative colonies in liquid nitrogen with a mortar

and pestle for homogenization and to maximize the available surface area for pollen extraction.

We used a small sample (~0.5 g) of this pulverized wax to undergo an acetolysis protocol to

extract and identify the pollen that was present. We separated the pollen grains from the wax by

adding 1 mL of glacial acetic acid to the wax sample in a 2 mL centrifuge tube and allowing it to

soak for 24 hours so that pollen grains entered solution (Furnessa, 1994). This solution was

then drained from the wax residue and centrifuged for 6 min at 13,000 rpm to form a pollen

pellet.

A standard acetolysis protocol was then carried out as follows (*sensu* Jones, 2014): we drained

the remaining acetic acid and then added 1 mL of a 9:1 solution of acetic anhydride and sulfuric

acid and vortexed each tube. These samples were immediately placed into a 5 min hot water

bath at 90°C and were then centrifuged for 5 min at 13,000 rpm. We removed the acetic

anhydride and sulfuric acid solution from the pellet and added a 50% ethanol solution to rinse

away any remaining solution. Samples were vortexed again and then underwent a final 5 min

centrifuge at 13,000 rpm. The pellet was again rinsed with the 50% ethanol solution and then

gently mixed with a small volume of glycerol to store the pollen grains for later identification.

Pollen counting and identification:

After a minimum of 24 hours we added 15 uL of this glycerol solution and a small volume of

fuschin stain to slides, and sealed the cover slip to the slide with clear nail polish. For

consistency, we counted pollen at 400x magnification along horizontal transects, starting at the

midline of the slide and establishing further transects bisecting the midline and slide edge until

the first 300 pollen grains were counted to systematically quantify the composition of pollen

collected by a colony (Jones & Bryant, 1998). A pollen grain was only counted if it was entirely

visible in the field of view, and all pollen grains visible in the field of view once a count of 300

was achieved were counted (for a final count of 300-306 per slide). Pollen was identified as

being within the family Rosaceae (Kapp *et al.*, 2000), or was categorized as non-rosaceous.

Statistical analyses:

All analyses were done with R version 4.4.2 (R Core Team, 2024), and checked for violations of

model assumptions (package: *DHARMa*; Hartig 2024). To test if bumble bee visitation rates

differed from visitation rates by all other bees to strawberry flowers we used a linear mixed

effect model (package: *lme4*; Bates *et al.*, 2015) to assess if the average number of bees

recorded was predicted by type of bee (bumble bee vs other bees) with field site as a random

factor.

1. <u>Does managed bumble bee visitation rate to strawberry depend on surrounding land cover?</u>

We evaluated if landscape composition surrounding each field influenced bumble bee visitation

to strawberry flowers by running three individual linear models, one for each land cover category

(agriculture, pasture, and semi-natural) as a predictor with the transformed (log + 1) average

number of bumble bee visits as our response variable. We also evaluated if field size influenced

bumble bee visitation with linear models. The average number of bumble bee visits was log + 1

transformed in all analyses to aid in model fit. We estimated predictor significance with ANOVA

likelihood ratio tests. We use the landscape 1000 m surrounding the fields because this scale

has previously been shown to be most relevant to bees foraging on strawberry in this region

(Connelly et al. 2015) and relevant to bumble bee foraging distances (mean: 842 m (McArt *et*

al., 2017b)), but also evaluated the 750 m and 500 m scales (Table 1).

2. <u>Do fields with greater visitation by managed bumble bees have reduced pollen limitation?</u>

To then assess if fields with greater observed bee visitation had reduced pollen limitation we ran

a linear mixed effects model (package: *lme4*; Bates *et al.*, 2015) with the difference between the

open and hand-pollinated fruit divided by the hand-pollinated fruit as the response and the

transformed (log + 1) average number of bumble bee visits as the predictor, we also ran these

models with average number of other bee visits (log + 1) and with average total number of bee

visits (log + 1) as predictors to understand the role of increased visitation by all bees. Last, we

ran a model with field size as a predictor to understand the role that the amount of cultivated

strawberry around the colonies had on pollen limitation. We included site as a random factor to

account for the multiple plants from each field. We estimated predictor significance with ANOVA

likelihood ratio tests.

3. <u>Is visitation rate by managed bumble bees reflected in colony Rosaceae pollen collection?</u>

To understand the relationship between observed bumble bee visitation at our focal crop and

the proportion of rosaceous pollen we recovered from the colonies we ran a beta regression

model (package: *betareg*; Cribari-Neto & Zeileis, 2012) with the transformed (log + 1) average

number of bumble bee visits as the predictor and the proportion of rosaceous pollen as the

response. We estimated predictor significance for the beta regression with a likelihood ratio test

(package: *lmtest;* Zeileis & Hothorn, 2002).

4. <u>Is pollen collection by managed bumble bees dependent on landscape context?</u>

We then evaluated if landscape composition surrounding each field influenced the proportion of

rosaceous pollen from the colony by running three models (package: *betareg*; Cribari-Neto &

Zeileis, 2012), one for each land cover category (agriculture, pasture, and semi-natural) as a

predictor with the proportion rosaceous pollen as the response variable. We also ran a model

with field size as a predictor to understand the role that the amount of cultivated strawberry

around the colonies had on rosaecous pollen collection. As above, we estimated predictor

significance with likelihood ratio tests (package: *lmtest;* Zeileis & Hothorn, 2002).

Results

Across the 4 visitation surveys to the 10 sites we observed a total of 648 bees visiting

strawberry flowers: 72 bumble bees, and 576 other bees. Bumble bees provided substantially

fewer visits to strawberry flowers compared to other bees (Figure 1B; X^2 = 12.786, df = 1, p <

0.001). Average bumble bee visits in the 10 min observation period across the 4 rounds per site

ranged from 0 to 6.75 bees (mean = 1.78) while visits from other bees ranged from 1 to 26.80

bees (mean = 11.08).

1. <u>Does managed bumble bee visitation rate to strawberry depend on surrounding land cover?</u>

The average number of bumble bee visits was not affected by surrounding pasture ($F_{1,8}$ = 1.043,

p = 0.337), semi-natural area ($F_{1,8}$ = 0.689, p = 0.431), or agriculture ($F_{1,8}$ = 0.002, p = 0.968) at

any scale (Table 1). Strawberry field size also did not affect bumble bee visitation ($F_{1,8}$ = 0.983,

p = 0.350).

2. <u>Do fields with greater visitation by managed bumble bees have improved pollination?</u>

There was no evidence that managed bumble bees improved pollination to strawberry as sites

with greater bumble bee visitation did not have reduced pollen limitation (Figure 2B; X^2 = 0.099,

df = 1, p = 0.754); increased visitation by other bees also did not have an effect (X^2 < 0.001, df =

1, p = 0.979), nor did increased total visitation (X^2 = 0.194, df = 1, p = 0.659). We also did not

find that surrounding landscape (1000 m surrounding the site) or field size affected pollen

limitation (pasture: X^2 = 1.266, df = 1, p = 0.260; agriculture: X^2 = 2.996, df = 1, p = 0.083; semi-

natural area: X^2 = 2.372, df = 1, p = 0.123; field size: X^2 = 0.877, df = 1, p = 0.348) at any spatial

scale (Table 1).

3. <u>Is visitation rate by managed bumble bees reflected in colony Rosaceae pollen collection?</u>

In line with the low visitation rates by bumble bees to strawberry, we recovered correspondingly

low proportions of rosaceous pollen from the colony wax, ranging from 0 to 0.20 of total pollen

grains screened (mean proportion ± sd= 0.04 ± 0.059). However, fields with more observed

bumble bee visits to strawberry flowers had a greater proportion of rosaceous pollen (Figure 2A;

z = 3.272, p = 0.001).

4. <u>Is pollen collection by managed bumble bees dependent on landscape context?</u>

Colonies at sites with larger strawberry fields collected more rosaceous pollen (Figure 3A; z =

3.878, p < 0.001). Bumble bee colonies collected less rosaceous pollen when there was more

pasture in the surrounding landscape at a 1000 m radius (Figure 3B; z = -2.092, p = 0.036). The

proportion of rosaceous pollen was not influenced by surrounding semi-natural area (z= -1.540,

p = 0.123) or by surrounding agriculture (z = 0.846, p = 0.397) at any scale (Table 1).

Discussion

Landscape composition is known to influence crop visitation by wild bees (Kennedy *et al.*, 2013) but the role of landscape in mediating the efficacy of managed bees is less well understood. Here, we address this gap using a novel approach that infers whole-colony foraging by extracting pollen from nest wax. We find that managed bumble bees (*B. impatiens*) provided a relatively low number of visits to our focal strawberry crop (family: Rosaceae) compared to other bees, collected a low proportion of rosaceous pollen, and increased visitation by bumble bees did not reduce pollen limitation. These findings improve out knowledge of how managed bumble bees can be effectively used in agroecosystems and suggest that supplementing strawberry pollination with managed bumble bee colonies is not likely to improve pollination, particularly in landscapes where other resources found in pasture habitats may be more attractive to bumble bees, although future work could directly compare pollen limitation between sites with and without managed bumble bees.

1. <u>Does managed bumble bee visitation rate to strawberry depend on surrounding land cover?</u>
While observed bumble bee visitation was not influenced by landscape context, their collection of rosaceous pollen at the colony level was both positively correlated with focal crop visitation and field size, and lower at fields with more pasture in the surrounding landscape. This indicates that although there may not be fewer bumblebees (*B. impatiens)* visiting strawberry flowers as pasture increases, the colony overall is dedicating less of its foraging efforts toward the focal strawberry crop when these other resources are present. Similarly, while we did not observe more bumble bees on strawberry flowers when fields were larger, the colony seems to have been dedicating relatively more of its foraging effort toward strawberry, which could be because a greater proportion of their foraging radius is occupied by strawberry as they are central place foragers (Bell 1990). Future work could use a similar design but identify all pollen grains to species instead of just categorizing as either rosaceous or not to determine which floral resources compete the most with strawberry for bumble bee visits since bumble bees are known to have strong floral preferences (Lanterman Novotny *et al.*, 2023).

2. <u>Do fields with greater visitation by managed bumble bees have improved pollination?</u>
Managed bumble bees in this system appeared to be primarily visiting other plants across our 10 sites since an average of only 4.3% of the pollen that these colonies collected was from plants within the family Rosaceae. This is counter to their intended purpose as supplemental pollinators to strawberry placed in the fields, but corresponds with work in other systems. For instance, managed honey bees collect less focal cranberry pollen in landscapes with high woodland cover which presumably contains

preferred floral resources (Guzman *et al.*, 2019). Bumble bees are known to have strong foraging

preferences based both on floral traits (Harder, 1988), resource availability (Aleixo *et al.*, 2017), and

nutritional needs of the colony (Vaudo *et al.*, 2016). When managed bumble bee colonies are added to

squash fields less than five percent of the pollen collected is from squash (Brochu *et al.*, 2020), and

managed colonies are not effective at providing supplemental pollination services for squash, showing no

increase in the amount of fruit produced per plant (Petersen *et al.*, 2013). Similarly, we also find that

greater visitation by bumble bees to strawberry did not reduce pollen limitation. Our hand-pollinated

fruits tended to be larger than those that were open-pollinated, which suggests that our sites were

generally pollen limited and so there may not have been enough bee visitation at any site to see improved

pollination which corresponds with our finding that total bee visitation was not related to improved fruit

set. As our sites appeared to be generally pollen-limited (our open-pollinated fruits tended to be

smaller than the hand-pollinated fruits), this suggests that animal-mediated pollination and/or

self fertilization did not fully maximize pollination. Increased stocking density of managed bumble

bees may improve pollination, as stocking rates twice our stocking density have been observed to improve

pollination in blueberry (Drummond et al. 2012). This is suggested by our finding that colonies in larger

fields collected more rosaceous pollen despite similar colony densities per acre, inconsistent with a

scenario in which strawberry flowers were saturated by managed bees and thus limited in pollen

availability.

3. <u>Is visitation rate by managed bumble bees reflected in colony Rosaceae pollen collection?</u>

In this study we adapted a method of extracting pollen from museum wax sculptures by soaking wax

acetic acid prior to acetolysis (Furnessa, 1994) and studying historic honey bee foraging (Kasso *et al.*,

2023, 2024) for use in a contemporary ecological context to understand bumble bee foraging efforts at the

colony scale. It has recently been suggested that when senesced bumble bee colonies are

incidentally found in the wild data should be collected on colony demographics, size,

architecture and pollen stored in pots to gain a better understanding of these understudied

aspects of bumble bee colonies across species (Smith *et al.*, 2025), and we suggest that

extracting pollen from the wax broadly could also lend useful insights to the floral resources

used. Due to the habit of building outward from a central point as the bumble bee colony grows

(Sakagami *et al.*, 1967) it may also be possible to look at colony-level pollen collection over

time. This could allow for broader studies of colony foraging dynamics and resource usage;

including from species that are not commercially managed.

4. <u>Is pollen collection by managed bumble bees dependent on landscape context?</u>

Bee diet composition can be influenced by the surrounding landscape (Inês da Silva *et al.*,
2024). For instance, landscapes with more semi-natural habitats and lower agricultural cover
have been associated with greater diet diversity and lower collection of Rosaceae pollen in the
solitary managed bee, *Osmia cornifrons* (Centrella *et al.*, 2020). While we do not see an effect
of semi-natural or agricultural landcover, we do find our colonies collecting less rosaceous
pollen in landscapes with more surrounding pasture, which similarly suggests that crop pollen
collection is influenced by what other resources are available in the surrounding landscape. For
example, clover (*Trifolium* spp.) is a preferred floral resource for *Bombus impatiens* (Lanterman
Novotny *et al.*, 2023) and is included in our pasture areas, while our semi-natural areas were
largely forests with primarily wind pollinated species blooming during our study period (Regal,
1982). Other rosaceous crops (apple and raspberry) and wild rosaceous plants (ex: *Crataegus*
spp*., Rubus* spp*.,* and *Pyrus* spp.) were present in low abundance around some fields, and as
pollen was not identified to species it is possible that visitation to these plants contributed to the
patterns that we find. For example, co-blooming apple is known to reduce visitation to
strawberry during peak bloom and then increase visitation later in the season (Grab *et al.*, 2017)
so if other rosaceous plants in the landscape were co-blooming with strawberry there could be
increased pollen collection from these plants while simultaneously influencing strawberry
visitation. However, the link between strawberry visitation and field size to Rosaceae pollen
suggests the rosaceous pollen is likely strawberry.
425
Conclusions:
There are potential costs associated with bringing in managed bees to supplement pollination.
There is the financial cost of the bees themselves (Baylis *et al.*, 2021), but also they can have
negative effects on wild pollinators (Paini, 2004) that may already be providing these services
(Garibaldi *et al.*, 2014). Managed bees can outcompete wild bees (Mallinger *et al.*, 2017),
spread disease (Graystock *et al.*, 2013, 2016; Martin *et al.*, 2021), and can even function as
ecological traps that directly lead to wild bumble bee queen mortality (Miller *et al.*, 2023). Only
assessing these risks to the pollinator community at a focal crop will poorly capture the true
impact if they are largely foraging elsewhere.
435
To accurately assess the potential benefits of managed bees we need to understand whole colony foraging
dynamics. Managed bumble bees are effective pollinators for many pollinator-dependent greenhouse
crops as they are not accessible to wild pollinators (Velthuis & Doorn, 2006) and some field crops
(Stubbs & Drummond, 2001; Drummond, 2012). We find that bumble bee colony-level foraging effort

dedicated to our focal crop was generally very low and landscape-context dependent. Furthermore, we

found that pollen limitation did not vary based on the observed bumble bee visitation rates; however,

future work could expand to include sites without bumble bees present across a larger number of sites to

better understand the role that bumble bees may play in strawberry pollination. By applying a novel

method to extract pollen from colony wax, we quantify whole-colony bumble bee foraging and assess

whether colonies preferentially forage on a focal crop and which wild pollinator communities they may

impact.

Figure 1: A) Site map depicting 2018 landscape categories by color (brown indicating

agricultural land cover, yellow indicating pasture land cover, and green indicating semi-natural

land cover). **B)** Charts showing the proportion and number of observed bee visits to strawberry

flowers by bumble bees vs all other bees at each of the 10 fields. The average number of visits

to strawberry flowers by bumble bees (shown in yellow) and by other bees (shown in purple) are

also displayed for each site. **C)** Method of collecting pollen from removed primary flowers to

hand-pollinate one of the secondary flowers while leaving the other to be open-pollinated in the

fields to assess the influence of bee visitation on pollination limitation.

Figure 2: Increased bumble bee visitation led to more rosaceous pollen in the colony wax but did not affect pollen limitation. **A:** Relationship between the transformed average number of bumble bee visits to strawberry flowers observed across the 4x 10 min visitation surveys and

the proportion of rosaceous pollen recovered from the colony wax with beta-regression 95%

confidence intervals. **B:** Pollen limitation data with each point representing the value calculated

from the masses of a pair of fruit from a single plant ((Fruit$_{hand}$ - Fruit$_{open}$) / Fruit$_{hand}$). The mean

+/- SE for each site is depicted in black. The grey dotted line at zero represents if there was no

difference between the hand-pollinated and open-pollinated fruit, all values above zero show

pairs where the hand-pollinated fruit was larger than the open-pollinated fruit.

Figure 3: More rosaceous pollen was recovered from colony wax when strawberry fields were

larger, and less rosaceous pollen was recovered when there was more pasture in the

surrounding landscape. **A:** Relationship between strawberry field size (hectares) and the

proportion of rosaceous pollen recovered from the colony wax with beta-regression 95%

confidence interval. **B:** Relationship between the proportion of pasture surrounding 1000m

around fields and the proportion of rosaceous pollen recovered from the colony wax with beta-

regression 95% confidence interval.

Table 1: Results from models across three spatial scales (1000 m, 750 m, and 500 m) for

landcover categories. Significant effects are bolded. See visualizations of effects at the 1000 m

scale in supplementary information (Fig. S1).

Response	Landcover Category	Surrounding Scale	Test Statistic	P-Value
Bumble bee visitation (log + 1 transformed)	Pasture	1000 m	F = 1.043	p = 0.337
		750 m	F = 0.476	p = 0.537
		500 m	F = 0.414	p = 0.537
	Agriculture	1000 m	F = 0.002	p = 0.968
		750 m	F = 0.001	p = 0.972
		500 m	F = 0.011	p = 0.918
	Semi-Natural	1000 m	F = 0.689	p = 0.431
		750 m	F = 0.692	p = 0.429
		500 m	F = 0.077	p = 0.788
Proportion rosaceous pollen	Pasture	**1000 m**	**z = -2.092**	**p = 0.036**
		750 m	z = -1.634	p = 0.102
		500 m	z = -1.599	p = 0.109
	Agriculture	1000 m	z = 0.846	p = 0.397
		750 m	z = 0.827	p = 0.408
		500 m	z = 1.060	p = 0.289
	Semi-Natural	1000 m	z = -1.540	p = 0.123
		750 m	z = -1.433	p = 0.151
		500 m	z = -0.651	p = 0.514

Pollen limitation: (hand - open) / hand	Pasture	1000 m	$X^2 = 1.266$	$p = 0.260$
		500 m	$X^2 = 0.776$	$p = 0.378$
		750 m	$X^2 = 0.566$	$p = 0.451$
	Agriculture	1000 m	$X^2 = 2.996$	$p = 0.083$
		750 m	$X^2 = 2.412$	$p = 0.120$
		500 m	$X^2 = 1.745$	$p = 0.186$
	Semi-Natural	1000 m	$X^2 = 2.372$	$p = 0.123$
		750 m	$X^2 = 2.427$	$p = 0.119$
		500 m	$X^2 = 1.754$	$p = 0.185$

References

References

Aleixo, K.P., Menezes, C., Imperatriz Fonseca, V.L. & Silva, C.I. da. (2017) Seasonal availability
of floral resources and ambient temperature shape stingless bee foraging
behavior (Scaptotrigona aff. depilis). *Apidologie*, **48**, 117–127.

Artz, D.R. & Nault, B.A. (2011) Performance of Apis mellifera, Bombus impatiens, and
Peponapis pruinosa (Hymenoptera: Apidae) as Pollinators of Pumpkin. *Journal of*
Economic Entomology, **104**, 1153–1161.

Bänsch, S., Tscharntke, T., Wünschiers, R., Netter, L., Brenig, B., Gabriel, D., *et al.* (2020)
Using ITS2 metabarcoding and microscopy to analyse shifts in pollen diets of honey
bees and bumble bees along a mass-flowering crop gradient. *Molecular Ecology*, **29**,
5003–5018.

Baskin, J.M. & Baskin, C.C. (2018) Pollen limitation and its effect on seed germination. *Seed*
Science Research, **28**, 253–260.

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software*, **67**, 1–48.

Baylis, K., Lichtenberg, E.M. & Lichtenberg, E. (2021) Economics of Pollination. *Annual Review*
of Resource Economics, **13**, 335–354.

Bell, W.J. (1990) Central place foraging. *Search Behavior: Chapman and Hall Animal Behaviour*
Series. Springer, Dordrecht.

Brochu, K.K., Dyke, M.T. van, Milano, N.J., Petersen, J.D., McArt, S.H., Nault, B.A., *et al.* (2020)
Pollen defenses negatively impact foraging and fitness in a generalist bee (Bombus
impatiens: Apidae). *Scientific Reports*, **10**, 3112.

Cane, J.H. & Tepedino, V.J. (2001) Causes and Extent of Declines among Native North
American Invertebrate Pollinators: Detection, Evidence, and Consequences.
Conservation Ecology, **5**, art1.

Centrella, M., Russo, L., Moreno Ramírez, N., Eitzer, B., Dyke, M. van, Danforth, B., *et al.*
(2020) Diet diversity and pesticide risk mediate the negative effects of land use change
on solitary bee offspring production. *Journal of Applied Ecology*, **57**, 1031-1042.

Corbet, S.A., Fussell, M., Ake, R., Fraser, A., Gunson, C., Savage, A., *et al.* (1993)
Temperature and the pollinating activity of social bees. *Ecological Entomology*, **18**, 17–
30.

Connelly, H., Poveda, K. & Loeb, G. (2015) Landscape simplification decreases wild bee
pollination services to strawberry. *Agriculture, Ecosystems & Environment*, **211**, 51–56.

Cribari-Neto, F. & Zeileis, A. (2012) Beta Regression in R. *Journal of Statistical Software*, **34**, 1–
25.

Cropland Data Layer. (2022). USDA NASS.

De Luca, P.A. & Vallejo-Marín, M. (2013) What's the 'buzz' about? The ecology and

evolutionary significance of buzz-pollination. *Current Opinion in Plant Biology*, **16**, 429–435.

Desjardins, È.-C. & De Oliveira, D. (2006) Commercial Bumble Bee Bombus impatiens (Hymenoptera: Apidae) as a Pollinator in Lowbush Blueberry (Ericale: Ericaceae) Fields. *Journal of Economic Entomology*, **99**, 443–449.

Dimou, M., Taraza, S., Thrasyvoulou, A. & Vasilakakis, M. (2008) Effect of bumble bee pollination on greenhouse strawberry production. *Journal of Apicultural Research*, **47**, 99–101.

Drummond, F. (2012) Commercial Bumble Bee Pollination of Lowbush Blueberry. *International Journal of Fruit Science*, **12**, 54–64.

Evans, E. (2017) From Humble Bee to Greenhouse Pollination Workhorse: Can We Mitigate Risks for Bumble Bees? *Bee World*, **94**, 34–41.

Furnessa, C.A. (1994) The extraction and identification of pollen from a beeswax statue. *Grana*, **33**, 49–52.

Gaines-Day, H.R. & Gratton, C. (2016) Crop yield is correlated with honey bee hive density but not in high-woodland landscapes. *Agriculture, Ecosystems & Environment*, **218**, 53–57.

Garibaldi, L.A., Carvalheiro, L.G., Leonhardt, S.D., Aizen, M.A., Blaauw, B.R., Isaacs, R., *et al.* (2014) From research to action: enhancing crop yield through wild pollinators. *Frontiers in Ecology and the Environment*, **12**, 439–447.

Garibaldi, L.A., Steffan-Dewenter, I., Kremen, C., Morales, J.M., Bommarco, R., Cunningham, S.A., *et al.* (2011) Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecology Letters*, **14**, 1062–1072.

Goulson, D., Nicholls, E., Botías, C. & Rotheray, E.L. (2015) Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, **347**, 1255957.

Grab, H., Blitzer, E.J., Danforth, B., Loeb, G. & Poveda, K. (2017) Temporally dependent pollinator competition and facilitation with mass flowering crops affects yield in co-blooming crops. *Scientific Reports*, **7**, 45296.

Grab, H., Branstetter, M.G., Amon, N., Urban-Mead, K.R., Park, M.G., Gibbs, J., *et al.* (2019) Agriculturally dominated landscapes reduce bee phylogenetic diversity and pollination services. *Science*, **363**, 282–284.

Grab, H., Poveda, K., Danforth, B. & Loeb, G. (2018) Landscape context shifts the balance of costs and benefits from wildflower borders on multiple ecosystem services. *Proceedings of the Royal Society B: Biological Sciences*, **285**, 20181102.

Graham, K.K., Milbrath, M.O., Killewald, M., Soehnlen, A., Zhang, Y. & Isaacs, R. (2023) Identity and diversity of pollens collected by two managed bee species while in blueberry fields for pollination. *Environmental Entomology*, **52**, 907–917.

Graystock, P., Blane, E.J., McFrederick, Q.S., Goulson, D. & Hughes, W.O.H. (2016) Do

managed bees drive parasite spread and emergence in wild bees? *International Journal*
for Parasitology: Parasites and Wildlife, Including articles from the 25th WAAVP
Conference, Liverpool, August 2015, **5**, 64–75.

Graystock, P., Yates, K., Evison, S.E.F., Darvill, B., Goulson, D. & Hughes, W.O.H. (2013) The
Trojan hives: pollinator pathogens, imported and distributed in bumblebee colonies.
Journal of Applied Ecology, n/a-n/a.

Gudowska, A., Cwajna, A., Marjańska, E. & Moroń, D. (2024) Pollinators enhance the
production of a superior strawberry – A global review and meta-analysis. *Agriculture,*
Ecosystems & Environment, **362**, 108815.

Guzman, A., Gaines-Day, H.R., Lois, A.N., Steffan, S.A., Brunet, J., Zalapa, J., *et al.* (2019)
Surrounding landscape and spatial arrangement of honey bee hives affect pollen
foraging and yield in cranberry. *Agriculture, Ecosystems & Environment*, **286**, 106624.

Hines, H.M., Cameron, S.A., & Deans, A.R. (2007) Nest architecture and foraging behavior in
Bombus pullatus (Hymenoptera: Apidae), while comparisons to other tropical bumble
bees. *Journal of the Kansas Entomological Society*, **80(1)**, 1-15.

Harder, L.D. (1988) Choice of individual flowers by bumble bees: interaction of morphology,
time and energy. *Behaviour*, **104**, 60–77.

Hertig, F. (2024) DHARMa: Residual diagnostics for hierarchial (Multi-Level / Mixed) regression
models. 10.32614/CRAN.package.DHARMa

Inês da Silva, C., Victorino Nicolosi Arena, M., Cristina Mathias da Silva, E., Roberto Martines,
M., Malaspina, O., Chiovatto, G., *et al.* (2024) Landscape and land use affect
composition and nutritional values of bees' food. *Journal of Environmental Management*,
352, 120031.

Inouye, D.W. (1978) Resource Partitioning in Bumblebees: Experimental Studies of Foraging
Behavior. *Ecology*, **59**, 672–678.

Jones, G.D. (2014) Pollen analyses for pollination researh, acetolysis, **13**, 15.

Jones, G.D. & Bryant, V.M. (1998) Are all counts created equal? *Contributions Series-American*
Association. of Stratigraphic Palynologists Foundation, **33**, 115–120.

Kapp, R.O., Davis, O.K. & King, J.E. (2000) *Pollen and spores*. 2nd edn. American Association
of Stratigraphic Palynologists Foundation, Dallas, TX.

Kasso, T.M., Bruun Jensen, A., Johns, S., Roffet-Salque, M., Collins, M.J. & Enevold, R. (2024)
Beescapes – extracting pollen from historical Danish beeswax to explore honeybee
foraging. *Palynology*, **48**, 2288220.

Kasso, T.M., Enevold, R., Johns, S., Rangel-Piñeros, G., Taurozzi, A.J., Sutherland, A., *et al.*
(2023) ArcHives—combined palynological, genomic and lipid analysis of medieval wax
seals. *Heritage Science*, **11**.

Kennedy, C.M., Lonsdorf, E., Neel, M.C., Williams, N.M., Ricketts, T.H., Winfree, R., *et al.*

(2013) A global quantitative synthesis of local and landscape effects on wild bee
pollinators in agroecosystems. *Ecology Letters*, **16**, 584–599.

Khalifa, S.A.M., Elshafiey, E.H., Shetaia, A.A., El-Wahed, A.A.A., Algethami, A.F., Musharraf,
S.G., *et al.* (2021) Overview of Bee Pollination and Its Economic Value for Crop
Production. *Insects*, **12**, 688.

Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C.,
et al. (2007) Importance of pollinators in changing landscapes for world crops.
Proceedings of the Royal Society B: Biological Sciences, **274**, 303–313.

Lanterman, J., Reeher, P., Mitchell, R.J. & Goodell, K. (2019) Habitat Preference and
Phenology of Nest Seeking and Foraging Spring Bumble Bee Queens in Northeastern
North America (Hymenoptera: Apidae: Bombus). *The American Midland Naturalist*, **182**,
131.

Lanterman Novotny, J., Lybbert, A., Reeher, P., Mitchell, R.J. & Goodell, K. (2023) Bumble bee
banquet: Genus- and species-level floral selection by Midwestern Bombus. *Ecosphere*,
14, e4425.

Larson, B.M.H. & Barrett, S.C.H. (1999) The ecology of pollen limitation in buzz☐pollinated
Rhexia virginica (Melastomataceae). *Journal of Ecology*, **87**, 371–381.

Mallinger, R.E., Gaines-Day, H.R. & Gratton, C. (2017) Do managed bees have negative effects
on wild bees?: A systematic review of the literature. *PLoS ONE*, **12**.

Mallinger, R.E. & Gratton, C. (2015) Species richness of wild bees, but not the use of managed
honeybees, increases fruit set of a pollinator-dependent crop. *Journal of Applied
Ecology*, **52**, 323–330.

Martin, C.D., Fountain, M.T. & Brown, M.J.F. (2019) Varietal and seasonal differences in the
effects of commercial bumblebees on fruit quality in strawberry crops. *Agriculture,
Ecosystems & Environment*, **281**, 124–133.

McArt, S.H., Fersch, A.A., Milano, N.J., Truitt, L.L. & Böröczky, K. (2017a) High pesticide risk to
honey bees despite low focal crop pollen collection during pollination of a mass blooming
crop. *Scientific Reports*, **7**, 1–10.

McArt, S.H., Urbanowicz, C., McCoshum, S., Irwin, R.E. & Adler, L.S. (2017b) Landscape
predictors of pathogen prevalence and range contractions in US bumblebees.
Proceedings of the Royal Society B: Biological Sciences, **284**, 20172181.

Michener, C.D. (1974) *The Social Behavior of the Bees: A Comparative Study*. Harvard
University Press.

Miller, O., Hale, C., Richardson, L., Sossa, D., Iverson, A., McArt, S., *et al.* (2023) Commercial
Bombus impatiens colonies function as ecological traps for wild queens. *Journal of
Applied Ecology*, **60**, 592–600.

Osterman, J., Aizen, M.A., Biesmeijer, J.C., Bosch, J., Howlett, B.G., Inouye, D.W., *et al.* (2021)
Global trends in the number and diversity of managed pollinator species. *Agriculture,*

Ecosystems & Environment, **322**, 107653.

Paini, D.R. (2004) Impact of the introduced honey bee (Apis mellifera) (Hymenoptera: Apidae)
on native bees: A review. *Austral Ecology*, **29**, 399–407.

Petersen, J.D., Reiners, S. & Nault, B.A. (2013) Pollination Services Provided by Bees in
Pumpkin Fields Supplemented with Either Apis mellifera or Bombus impatiens or Not
Supplemented. *PLoS ONE*, **8**, e69819.

Petran, A. & Hoover, E. (2018) The Flowers of Fragaria × ananassa. In *Horticultural Reviews*.
John Wiley & Sons, Ltd, pp. 1–31.

R Core Team. (2024) R: A language and environment for statistical computing.

Regal, P.J. (1982) Pollination by Wind and Animals: Ecology of Geographic Patterns. *Annual*
Review of Ecology and Systematics, **13**, 497–524.

Ricketts, T.H., Regetz, J., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C., Bogdanski, A.,
et al. (2018) Landscape effects on crop pollination services: are there general patterns?
Ecology Letters, 499–515.

Rucker, R.R., Thurman, W.N. & Burgett, M. (2012) Honey Bee Pollination Markets and the
Internalization of Reciprocal Benefits. *American Journal of Agricultural Economics*, **94**,
956–977.

Saifuddin, M. & Jha, S. (2014) Colony-Level Variation in Pollen Collection and Foraging
Preferences Among Wild-Caught Bumble Bees (Hymenoptera: Apidae). *Environmental*
Entomology, **43**, 393–401.

Sakagami, S.F., Akahira, Y. & Zucchi, R. (1967) Nest architecture and brood development in a
neotropical bumblebee,Bombus atratus. *Insectes Sociaux*, **14**, 389–413.

Smith, T.A., Boone, M., Choy, S., Evans, E., Everett, J.G., Palmer, J.E., *et al.* (2025) Answering
key bumble bee conservation questions by studying discovered wild nests: A *Bombus*
affinis case study. *Insect Conservation and Diversity*, **18**, 303–318.

Stanghellini, M.S., Ambrose, J.T. & Schultheis, J.R. (1998) Using Commercial Bumble Bee
Colonies as Backup Pollinators for Honey Bees to Produce Cucumbers and
Watermelons. *HortTechnology*, **8**, 590–594.

Stavert, J.R., Liñán-Cembrano, G., Beggs, J.R., Howlett, B.G., Pattemore, D.E. & Bartomeus, I.
(2016) Hairiness: the missing link between pollinators and pollination. *PeerJ*, **4**, e2779.

Stubbs, C.S. & Drummond, F.A. (2001) *Bombus impatiens* (Hymenoptera: Apidae): An
Alternative to *Apis mellifera* (Hymenoptera: Apidae) for Lowbush Blueberry Pollination.
Journal of Economic Entomology, **94**, 609–616.

Trillo, A., Herrera, J.M. & Vilà, M. (2018) Managed bumble bees increase flower visitation but
not fruit weight in polytunnel strawberry crops. *Basic and Applied Ecology*, **30**, 32–40.

Vaudo, A.D., Patch, H.M., Mortensen, D.A., Tooker, J.F. & Grozinger, C.M. (2016)

Macronutrient ratios in pollen shape bumble bee (Bombus impatiens) foraging strategies
and floral preferences. *Proceedings of the National Academy of Sciences*, **113**, E4035–
E4042.

Velthuis, H.H.W. & Doorn, A. van. (2006) A century of advances in bumblebee domestication
and the economic and environmental aspects of its commercialization for pollination.
Apidologie, **37**, 421–451.

Wood, T.J., Kaplan, I. & Szendrei, Z. (2018) Wild Bee Pollen Diets Reveal Patterns of Seasonal
Foraging Resources for Honey Bees. *Frontiers in Ecology and Evolution*, **6**, 210.

Zeileis, A. & Hothorn, T. (2002) Diagnostic Checking in Regresssion Relationships. *R News*, **2**,
7–10.