1 **Title:** Colony-level pollen collection reflects visitation of managed bumble bees (*Bombus*
2 *impatiens*) in strawberry fields and surrounding landscapes without improving crop yield.
3
Abstract

Managed bees are frequently used to supplement pollination of a range of crop plants. Yet, their
effectiveness varies and much of their foraging can be outside of the focal crop. Bee foraging
behavior is highly sensitive to the availability of alternative floral resources which depend on the
composition of the surrounding landscape. To understand how foraging and crop pollination
dynamics of the managed common eastern bumble bee (*Bombus impatiens*) are influenced by
landscape composition, we assessed their visitation and pollination in cultivated strawberry
fields along a landscape gradient. We also used pollen extracted from bumble bee (*Bombus*
impatiens) colony wax to understand whole-colony foraging over the blooming period. Bumble
bee visits to strawberry crop flowers were very low and did not improve pollination; however,
colonies in fields with higher visitation rates had more pollen from the Rosaceae family
(including strawberry). We found that landscape composition affected foraging effort on
strawberry at the colony level, as a lower proportion of rosaceous pollen was collected in
landscapes with more surrounding pasture, while a greater proportion of rosaceous pollen was
collected at sites with larger strawberry fields. Overall, our results suggest that managed bumble
bees do forage on strawberry crop flowers, but they predominantly forage elsewhere, especially
in surrounding pastures when strawberry fields are small. Therefore, addition of bumble bee
colonies may not be economically beneficial for strawberry growers at least in diverse
agricultural landscapes similar to that of our study region.

Introduction
Pollinators serve an important role in food production, with about 30% of agricultural food
production relying to some extent on animal mediated pollination (Klein *et al.*, 2007; Khalifa *et*
al., 2021). Yet, agriculture itself is a major driver of pollinator decline (Cane & Tepedino, 2001)
due to factors like habitat loss and exposure to pesticides (Goulson *et al.*, 2015) leading to
reductions in the services provided by wild bees frequently observed in landscapes that are
more agriculturally dominated (Ricketts *et al.*, 2018; Grab *et al.*, 2019). To achieve sufficient
pollination, growers will sometimes bring in managed bees to supplement natural pollination
(Rucker *et al.*, 2012). However, supplemental managed bees may not always improve yields
(Petersen *et al.*, 2013) and factors that moderate their efficacy are unclear.
34

Supplementing crop pollination with managed pollinators does not always increase pollination

rates in the focal crop (Petersen *et al.*, 2013; Trillo *et al.*, 2018) for two potential reasons. First,

in landscapes with high amounts of natural habitat, wild pollinators are often more abundant and

more likely to meet crop pollination demands on their own (Garibaldi *et al.*, 2011; Kennedy *et

al.*, 2013; Mallinger & Gratton, 2015). Second, in environments with abundant natural habitats,

managed pollinators may preferentially forage in these areas and overlook the focal crop

(Guzman *et al.*, 2019). For example, in cranberry agroecosystems honey bee stocking density

was found to have a strong positive relationship with yield in fields that have low forest cover but

no relationship to yield in landscapes with high forest cover (Gaines-Day & Gratton, 2016),

where around only 15% of pollen from hives was from cranberry (Guzman *et al.*, 2019). These

low rates of focal crop pollen collection by managed pollinators have been reported from a

range of other crop systems (8.7% apple (McArt *et al.*, 2017a), 0.4-2% pumpkin (Brochu *et al.*,

2020), 19-26% strawberry (Bänsch *et al.*, 2020), 0.5-26% blueberry (Graham *et al.*, 2023)), and

suggest that managed bees are frequently using resources outside of the focal crop. For these

reasons, we expect managed bees to be more effective in simple, agriculturally dominated

landscapes than in landscapes containing more natural and semi-natural habitats.

The honey bee (*Apis mellifera*) is not the only managed bee that can be used to supplement

pollination services (Osterman *et al.*, 2021). The common eastern bumble bee (*Bombus

impatiens*) is widely used across the eastern United States as a commercially available

alternative to honey bees for supplemental pollination services (Evans, 2017). While these

colonies are primarily used for pollination within greenhouses (Evans, 2017), they are also sold

to supplement wild pollinators for crops grown in fields (Stanghellini *et al.*, 1998; Desjardins &

De Oliveira, 2006; Artz & Nault, 2011). Bumble bees can be very effective at pollinating certain

crops (ex. blueberry, tomatoes, etc) due to traits like hairiness (pilosity) (Stavert *et al.*, 2016)

and buzz pollination (De Luca & Vallejo-Marín, 2013). They can also have a higher per-visit

efficiency and can be obtained for prices that can be comparable or even less expensive than

renting commercial honey bees for pollination (Drummond, 2012).

Strawberries are a crop which have been shown to benefit from bumble bee pollination in

greenhouses and other enclosed environments (Dimou *et al.*, 2008; Martin *et al.*, 2019).

However, the majority of strawberry production occurs in open field conditions (*USDA NASS*,

2022). In the northeast United States, wild bumble bee foundress queens begin to start colonies

during strawberry bloom from May to early June, but are not established to produce foraging

workers until after the bloom of spring strawberry has completed (Lanterman *et al.*, 2019).

Indeed, previous work suggests that bumble bees do not visit spring strawberry plantings in

New York at sites where they are not actively managed (Connelly *et al.*, 2015). This suggests

the potential for commercially available bumble bee colonies to provide supplemental services,

particularly in agriculturally dominated landscapes where pollination services from wild bees

may be lower (Connelly *et al.*, 2015).

Understanding which floral resources bumble bees are foraging on by identifying pollen grains

offers an important insight into their foraging patterns (Guzman *et al.*, 2019). It is typically

difficult to get this colony level information because floral usage in bumble bees varies between

individuals (Saifuddin & Jha, 2014), across the season (Wood *et al.*, 2018), and even over the

course of a single day (Inouye, 1978). This means that short sampling periods are unlikely to

capture the true breadth of floral resource use, and it is typically impractical to look at the pollen

loads of every returning forager to a colony. However, techniques can be adapted for extracting

pollen from beeswax (Furnessa, 1994) to look at the pollen brought back at the colony level,

which allows integration over time and across individuals with a single sample from a colony.

These methods have recently been adapted to look at the pollen found in medieval wax seals

(Kasso *et al.*, 2023), and to understand historic honey bee foraging patterns from museum

comb specimens (Kasso *et al.*, 2024); their use in investigating variation in contemporary

bumble bee foraging offers a promising new direction to understanding season-long colony level

foraging efforts.

In this study, we assessed the degree to which supplemental managed colonies of *B. impatiens*

visit strawberry plantings in open field settings. Specifically, we focus our analyses on four

questions: (1) Does managed bumble bee visitation rate to strawberry depend on surrounding

land cover? (2) Do fields with greater visitation by managed bumble bees have improved

pollination? (3) Is visitation rate by managed bumble bees reflected in colony Rosaceae pollen

collection? and (4) Is pollen collection by managed bumble bees dependent on landscape

context?

Methods

Landscape composition and colony placements:

In the spring of 2018, 43 commercially available bumble bee (*Bombus impatiens)* colonies from

Koppert Biological Systems (MI, USA) were placed in or along the edge of 10 strawberry fields

across upstate New York (Fig. 1A) that were a minimum of 500 m apart. Colonies were placed
in the fields between May 14-15th, just prior to strawberry bloom, and remained in the fields until
fruit had set in mid June. The number of colonies placed in each field ranged from 1-16
depending on the field size so that stocking densities were maintained at approximately two
colonies per acre (0.4 ha) as recommended by Koppert Biological Systems. Fields ranged in
size from 0.069 hectares to 3.035 hectares.
109
We determined the landscape composition around each field using the 2018 Cropland Data
Layer (https://nassgeodata.gmu.edu/CropScape/). The proportion of each land cover category
was calculated at 500, 750 and 1000 m through ArcGIS 10.3 (ESRI 2018) for three different
land cover types: agricultural (cultivated crops; ranging from 5.9%-86.1% at 1000 m), semi-
natural (forests, shrublands, wetlands; ranging from 2.4%-24.4% at 1000 m), and
pasture/grass/hay (alfalfa, meadows, clover, wildflowers, grasses; ranging from 5.8% to 30.0%
at 1000 m), hereafter referred to as 'pasture' (Figure 1).
117

Bumble bee visitation:
We conducted visual surveys to determine the visitation rate of bumble bees to strawberry
during the blooming period. Four times during strawberry bloom (May 16th - June 7th) we
walked a 10 min transect in each field at a consistent slow pace and counted each bee that
visited a flower. Visits were defined as contact with the flower petals or stigmas (Grab *et al.*,
2018), and we visually identified each bee as either a bumble bee or any other type of bee.
Because wild bumble bee workers were not yet foraging in this region during this time period
(Lanterman *et al.*, 2019) and previous studies in the region reported no bumble bee visitors to
strawberry at fields without managed colonies (Connelly *et al.*, 2015), we assumed that all
bumble bee visits during our sampling periods were from our managed *B. impatiens* colonies.
128

Sentinel plants for assessing pollination:
We started bare-root strawberry plants of the cultivar 'Seascape' (Indiana Berry, IN, USA) in 10
cm pots in a greenhouse approximately 5 weeks before the onset of spring field strawberry
bloom to use as sentinel plants flowering simultaneously with field-grown plants. Most sites
received 24 plants, although one site only received 18. Strawberry flowers tend to bifurcate from
the stem internodes such that one large "primary" flower head is on the same stem as two
smaller paired "secondary" flowers that typically open on the same day and will lead to fruits of
similar sizes (Petran & Hoover, 2018). We removed the primary flowers from our plants, leaving

the paired secondary flower buds, and dried the anthers from these primary flowers until they

dehisced. On the day of secondary flower anthesis we used pollen collected from primary

flowers to hand pollinate one of the secondary flowers by gently brushing pollen across all

stigmatic surfaces with a small paint brush (hand-pollinated) while we left the other secondary

flower to be pollinated in the field (open-pollinated). These were marked with a colored piece of

yarn tied loosely around the base of each flower (Fig. 1C).

Plants were brought out to the fields on the same day that they were hand-pollinated and placed

in trays that held ~3 cm of water in groups of 6 within the field rows when the field was

flowering. Sentinel plants were only being open pollinated when the field strawberries were also

blooming and were present throughout the blooming period as we brought out trays on four

dates throughout the season (first trays of plants to fields from May 14th - May 25th and the final

plants were brought to fields May 31st - June 7th). We watered the plants during all field visits,

and brought them back to be cared for in the greenhouse when the flowers had senesced (June

6th - June 22nd). Once the fruits had fully developed (June 7th - July 5th) we weighed each as a

measure of pollination quality (Gudowska *et al.*, 2024), and calculated the degree of pollen

limitation in the field as the weight of the open-pollinated fruit subtracted from the hand-

pollinated fruit then divided by the weight of the hand-pollinated fruit for each plant (Larson &

Barrett, 1999; Baskin & Baskin, 2018). We were able to retrieve paired fruits from 154 plants for

an average (+/- standard deviation) of 15.40 +/- 4.64 plants per field. Plants or their fruits that

were lost were due primarily to herbivory or disease, which was distributed fairly evenly across

fields (75 total plants were excluded from analyses: average 7.5 plants +/- 4.67 SD per site).

Wax and pollen extraction:

At the end of strawberry bloom, we removed approximately 10 g of wax from a single randomly

selected representative colony from each field to quantify pollen composition. Because each

colony grows over the course of the season, old wax is located in the center of the colony

(Sakagami *et al.*, 1967) and is much darker in color (Michener, 1974) so it is possible to only

remove newer wax that had built up along the entire periphery after the colony was put into the

strawberry fields. We expected the pollen present in the wax to be representative of pollen

collection over time by random contamination into the wax during larval feeding and wax

manipulation for construction (Hines *et al.*, 2007). We pulverized the wax samples from these

representative colonies in liquid nitrogen with a mortar and pestle for homogenization and to

maximize the available surface area for pollen extraction.

We used a small sample (~0.5 g) of this pulverized wax to undergo an acetolysis protocol to

extract and identify the pollen that was present. We separated the pollen grains from the wax by

adding 1 mL of glacial acetic acid to the wax sample in a 2 mL centrifuge tube and allowing it to

soak for 24 hours so that pollen grains entered solution (Furnessa, 1994). This solution was

then drained from the wax residue and centrifuged for 6 min at 13,000 rpm to form a pollen

pellet.

A standard acetolysis protocol was then carried out as follows (*sensu* Jones, 2014): we drained

the remaining acetic acid and then added 1 mL of a 9:1 solution of acetic anhydride and sulfuric

acid and vortexed each tube. These samples were immediately placed into a 5 min hot water

bath at 90°C and were then centrifuged for 5 min at 13,000 rpm. We removed the acetic

anhydride and sulfuric acid solution from the pellet and added a 50% ethanol solution to rinse

away any remaining solution. Samples were vortexed again and then underwent a final 5 min

centrifuge at 13,000 rpm. The pellet was again rinsed with the 50% ethanol solution and then

gently mixed with a small volume of glycerol to store the pollen grains for later identification.

Pollen counting and identification:

After a minimum of 24 hours we added 15 uL of this glycerol solution and a small volume of

fuschin stain to slides, and sealed the cover slip to the slide with clear nail polish. For

consistency, we counted pollen at 400x magnification along horizontal transects, starting at the

midline of the slide and establishing further transects bisecting the midline and slide edge until

the first 300 pollen grains were counted to systematically quantify the composition of pollen

collected by a colony (Jones & Bryant, 1998). A pollen grain was only counted if it was entirely

visible in the field of view, and all pollen grains visible in the field of view once a count of 300

was achieved were counted (for a final count of 300-306 per slide). Pollen was identified as

being within the family Rosaceae (Kapp *et al.*, 2000), or was categorized as non-rosaceous.

Statistical analyses:

We evaluated if landscape composition surrounding each field influenced bumble bee visitation

to strawberry flowers by running three individual linear models, one for each land cover category

(agriculture, pasture, and semi-natural) as a predictor with the transformed (log + 1) average

number of bumble bee visits as our response variable. We also evaluated if field size influenced

bumble bee visitation with linear models. The average number of bumble bee visits was log + 1

transformed in all analyses to aid in model fit. We estimated predictor significance with ANOVA

likelihood ratio tests. We use the landscape 1000 m surrounding the fields because this scale

has previously been shown to be most relevant to bees foraging on strawberry in this region

(Connelly et al. 2015) and relevant to bumble bee foraging distances (mean: 842 m (McArt *et*

al., 2017b)), but also evaluated the 750 m and 500 m scales (Table 1).

To test if bumble bee visitation rates differed from visitation rates by all other bees to strawberry

flowers we used a linear mixed effect model ((Bates *et al.*, 2015) to assess if the average

number of bees recorded was predicted by type of bee (bumble bee vs other bees) with field

site as a random factor. To then assess if fields with greater observed bee visitation had

reduced pollen limitation we ran a linear mixed effects model ((Bates *et al.*, 2015) with the

difference between the open and hand-pollinated fruit divided by the hand-pollinated fruit as the

response and the transformed (log + 1) average number of bumble bee visits as the predictor,

we also ran these models with average number of other bee visits (log + 1) and with average

total number of bee visits (log + 1) as predictors. Last, we ran a model with field size as a

predictor to understand the role that the amount of cultivated strawberry around the colonies

had on pollen limitation. We included site as a random factor to account for the multiple plants

from each field. We estimated predictor significance with ANOVA likelihood ratio tests.

To understand the relationship between observed bumble bee visitation at our focal crop and

the proportion of rosaceous pollen we recovered from the colonies we ran a beta regression

model (Cribari-Neto & Zeileis, 2012) with the transformed (log + 1) average number of bumble

bee visits as the predictor and the proportion of rosaceous pollen as the response. We

estimated predictor significance for the beta regression with a likelihood ratio test (Zeileis &

Hothorn, 2002).

We then evaluated if landscape composition surrounding each field influenced the proportion of

rosaceous pollen from the colony by running three models (Cribari-Neto & Zeileis, 2012), one

for each land cover category (agriculture, pasture, and semi-natural) as a predictor with the

proportion rosaceous pollen as the response variable. We also ran a model with field size as a

predictor to understand the role that the amount of cultivated strawberry around the colonies

had on rosaecous pollen collection. As above, we estimated predictor significance with

likelihood ratio tests (Zeileis & Hothorn, 2002).

All analyses were done with R version 4.4.2 (R Core Team, 2024).

Results

Bumble bees provided substantially fewer visits to strawberry flowers compared to other bees

(Figure 1B; X^2 = 12.786, df = 1, p < 0.001). Average bumble bee visits in the 10 min observation

period across the 4 rounds per site ranged from 0 to 6.75 (mean = 1.78) while visits from other

bees ranged from 1 to 26.80 (mean = 11.08). The average number of bumble bee visits was not

affected by surrounding pasture ($F_{1,8}$ = 1.043, p = 0.337), semi-natural area ($F_{1,8}$ = 0.689, p =

0.431), or agriculture ($F_{1,8}$ = 0.002, p = 0.968) at any scale (Table 1). Strawberry field size also

did not affect bumble bee visitation ($F_{1,8}$ = 0.983, p = 0.350). There was no evidence that

managed bumble bees improved pollination to strawberry as fields with greater bumble bee

visitation did not have reduced pollen limitation (Figure 2B; X^2 = 0.099, df = 1, p = 0.754);

increased visitation by other bees also did not have an effect (X^2 < 0.001, df = 1, p = 0.979), nor

did increased total visitation (X^2 = 0.194, df = 1, p = 0.659). We also did not find that

surrounding landscape (1000 m surrounding the site) or field size affected pollen limitation

(pasture: X^2 = 1.266, df = 1, p = 0.260; agriculture: X^2 = 2.996, df = 1, p = 0.083; semi-natural

area: X^2 = 2.372, df = 1, p = 0.123; field size: X^2 = 0.877, df = 1, p = 0.348) at any spatial scale

(Table 1).

In line with the low visitation rates by bumble bees to strawberry, we recovered correspondingly

low proportions of rosaceous pollen from the colony wax, ranging from 0 to 0.20 of total pollen

grains screened (mean proportion ± sd= 0.04 ± 0.059). However, fields with more observed

bumble bee visits to strawberry flowers had a greater proportion of rosaceous pollen (Figure 2A;

z = 3.272, p = 0.001), and colonies at sites with larger strawberry fields collected more

rosaceous pollen (Figure 3A; z = 3.878, p < 0.001). Bumble bee colonies collected less

rosaceous pollen when there was more pasture in the surrounding landscape at a 1000 m

radius (Figure 3B; z = -2.092, p = 0.036). The proportion of rosaceous pollen was not influenced

by surrounding semi-natural area (z= -1.540, p = 0.123) or by surrounding agriculture (z =

0.846, p = 0.397) at any scale (Table 1).

Discussion

Landscape composition is known to influence crop visitation by wild bees (Kennedy *et al.*, 2013)

but the role of landscape in mediating the efficacy of managed bees is less well understood.

Here we find that managed bumble bees (*B. impatiens*) provided a relatively low number of

visits to our focal strawberry crop (family: Rosaceae) compared to other bees, collected a low

proportion of rosaceous pollen, and did not improve pollination. These findings suggest that

supplementing strawberry pollination with managed bumble bee colonies is not likely to improve

pollination, particularly in landscapes where other resources found in pasture habitats may be

more attractive to bumble bees.

While observed bumble bee visitation was not influenced by landscape context, their collection

of rosaceous pollen at the colony level was both positively correlated with focal crop visitation

and field size, and lower at fields with more pasture in the surrounding landscape. Bee diet

composition can be influenced by the surrounding landscape (Inês da Silva *et al.*, 2024). For

instance, landscapes with more semi-natural habitats and lower agricultural cover have been

associated with greater diet diversity and lower collection of Rosaceae pollen in the solitary

managed bee, *Osmia cornifrons* (Centrella *et al.*, 2020). While we do not see an effect of semi-

natural or agricultural landcover, we do find our colonies collecting less rosaceous pollen in

landscapes with more surrounding pasture, which similarly suggests that crop pollen collection

is influenced by what other resources are available in the surrounding landscape. For example,

clover (*Trifolium* spp.) is a preferred floral resource for *Bombus impatiens* (Lanterman Novotny

et al., 2023) and is included in our pasture areas, while our semi-natural areas were largely

forests with primarily wind pollinated species blooming during our study period (Regal, 1982).

Other rosaceous crops (apple and raspberry) and wild rosaceous plants (ex: *Crataegus* spp.*,*

Rubus spp.*,* and *Pyrus* spp.) were present in low abundance around some fields, and as pollen

was not identified to species it is possible that visitation to these plants contributed to the

patterns that we find. For example, co-blooming apple is known to reduce visitation to

strawberry during peak bloom and then increase visitation later in the season (Grab *et al.*, 2017)

so if other rosaceous plants in the landscape were co-blooming with strawberry there could be

increased pollen collection from these plants while simultaneously influencing strawberry

visitation. However, the link between strawberry visits and field size to Rosaceae pollen

suggests the rosaceous pollen is likely strawberry. This indicates that although there may not be

fewer bumblebees (*B. impatiens)* visiting strawberry flowers as pasture increases, the colony

overall is dedicating less of its foraging efforts toward the focal strawberry crop when these

other resources are present. Similarly, while we did not observe more bumble bees on

strawberry flowers when fields were larger, the colony seems to have been dedicating relatively

more of its foraging effort toward strawberry. Future work could use a similar design but identify

all pollen grains to species to determine which floral resources compete the most with

strawberry for bumble bee visits.

Managed bumble bees in this system appeared to be primarily visiting other plants since an

average of only 4.3% of the pollen that these colonies collected was from plants within the

family Rosaceae. This is counter to their intended purpose as supplemental pollinators to

strawberry placed in the fields, but corresponds with work in other systems. For instance,

managed honey bees collect less focal cranberry pollen in landscapes with high woodland cover

which presumably contains preferred floral resources (Guzman *et al.*, 2019). Bumble bees are

known to have strong foraging preferences based both on floral traits (Harder, 1988), resource

availability (Aleixo *et al.*, 2017), and nutritional needs of the colony (Vaudo *et al.*, 2016). When

managed bumble bee colonies are added to squash fields less than five percent of the pollen

collected is from squash (Brochu *et al.*, 2020), and managed colonies are not effective at

providing supplemental pollination services for squash, showing no increase in the amount of

fruit produced per plant (Petersen *et al.*, 2013). Similarly, we also find that greater visitation by

bumble bees to strawberry did not reduce pollen limitation. Our hand pollinated fruits tended to

be larger than those that were open pollinated, which suggests that our sites were generally

pollen limited and so there may not have been enough bee visitation at any site to see improved

pollination which corresponds with our finding that total bee visitation was not related to

improved fruit set. Increased stocking density of managed bumble bees may improve

pollination, as stocking rates twice our stocking density have been observed to improve

pollination in blueberry (Drummond et al. 2012). This could be suggested in our system by our

finding that colonies in larger fields collected more rosaceous pollen despite similar densities

per acre, a pattern unlikely if strawberry flowers were saturated with our managed bees.

In this study we adapted a method of extracting pollen from museum wax sculptures (Furnessa,

1994) and studying historic honey bee foraging (Kasso *et al.*, 2023, 2024) for use in a

contemporary ecological context to understand bumble bee foraging efforts at the colony scale.

It has recently been suggested that when senesced bumble bee colonies are incidentally found

in the wild data should be collected on colony demographics, size, architecture and pollen

stored in pots to gain a better understanding of these understudied aspects of bumble bee

colonies across species (Smith *et al.*, 2025), and we suggest that extracting pollen from the wax

broadly could also lend useful insights to the floral resources used. Due to the habit of building

outward from a central point as the bumble bee colony grows (Sakagami *et al.*, 1967) it may

340 also be possible to look at colony-level pollen collection over time. This could allow for broader

341 studies of colony foraging dynamics and resource usage; including from species that are not

342 commercially managed.

344 There are potential costs associated with bringing in managed bees to supplement pollination.

345 There is the financial cost of the bees themselves (Baylis *et al.*, 2021), but also they can have

346 negative effects on wild pollinators (Paini, 2004) that may already be providing these services

347 (Garibaldi *et al.*, 2014). Managed bees can outcompete wild bees (Mallinger *et al.*, 2017),

348 spread disease (Graystock *et al.*, 2013, 2016), and can even function as ecological traps that

349 directly lead to wild bumble bee queen mortality (Miller *et al.*, 2023). Only assessing these risks

350 to the pollinator community at a focal crop will poorly capture the true impact if they are largely

351 foraging elsewhere.

353 To accurately assess the potential benefits of managed bees we need to understand whole

354 colony foraging dynamics. Managed bumble bees are effective pollinators for many pollinator

355 dependent greenhouse crops as they are not accessible to wild pollinators (Velthuis & Doorn,

356 2006) and some field crops (Stubbs & Drummond, 2001; Drummond, 2012). We find that the

357 colony-level foraging effort dedicated to our focal crop was generally very low and landscape

358 dependent, with no improvement to yield. We also cannot accurately assess their costs unless

359 we know which plant-pollinator communities they are interacting with, as their interactions with

360 the strawberry pollinator community appear to be limited. By studying whole colony foraging, we

361 can better understand if they preferentially forage on a focal crop and also which wild pollinator

362 communities they may be impacting.

Figure 1: A) Site map depicting 2018 landscape categories by color (brown indicating
agricultural land cover, yellow indicating pasture land cover, and green indicating semi-natural

366   land cover). **B)** Pie charts showing the proportion of observed bee visits to strawberry flowers by

367   bumble bees vs all other bees at each of the 10 fields. **C)** Method of collecting pollen from

368   removed primary flowers to hand pollinate one of the secondary flowers while leaving the other

369   to be open pollinated in the fields to assess the influence of bee visitation on pollination

370   limitation.

Figure 2: Increased bumble bee visitation led to more rosaceous pollen in the colony wax but did not affect pollen limitation. **A:** Relationship between the transformed average number of bumble bee visits to strawberry flowers observed across the 4x 10 min visitation surveys and

the proportion of rosaceous pollen recovered from the colony wax with beta-regression 95%

confidence intervals. **B:** Pollen limitation data with each point representing the value calculated

from the masses of a pair of fruit from a single plant (($Fruit_{hand}$ - $Fruit_{open}$) / $Fruit_{hand}$). The mean

+/- SE for each site is depicted in black. The grey dotted line at zero represents if there was no

difference between the hand pollinated and open pollinated fruit, all values above zero show

pairs where the hand pollinated fruit was larger than the open pollinated fruit.

Figure 3: More rosaceous pollen was recovered from colony wax when strawberry fields were

larger, and less rosaceous pollen was recovered when there was more pasture in the

surrounding landscape. **A:** Relationship between strawberry field size (hectares) and the

proportion of rosaceous pollen recovered from the colony wax with beta-regression 95%

confidence interval. **B:** Relationship between the proportion of pasture surrounding 1000m

around fields and the proportion of rosaceous pollen recovered from the colony wax with beta-

regression 95% confidence interval.

Table 1: Results from models across three spatial scales (1000 m, 750 m, and 500 m) for
landcover categories. Significant effects are bolded.

Response	Landcover Category	Surrounding Scale	Test Statistic	P-Value
Bumble bee visitation (log + 1 transformed)	Pasture	1000 m	F = 1.043	p = 0.337
		750 m	F = 0.476	p = 0.537
		500 m	F = 0.414	p = 0.537
	Agriculture	1000 m	F = 0.002	p = 0.968
		750 m	F = 0.001	p = 0.972
		500 m	F = 0.011	p = 0.918
	Semi-Natural	1000 m	F = 0.689	p = 0.431
		750 m	F = 0.692	p = 0.429
		500 m	F = 0.077	p = 0.788
Proportion rosaceous pollen	Pasture	**1000 m**	**z = -2.092**	**p = 0.036**
		750 m	z = -1.634	p = 0.102
		500 m	z = -1.599	p = 0.109
	Agriculture	1000 m	z = 0.846	p = 0.397
		750 m	z = 0.827	p = 0.408
		500 m	z = 1.060	p = 0.289
	Semi-Natural	1000 m	z = -1.540	p = 0.123

		750 m	z = -1.433	p = 0.151
		500 m	z = -0.651	p = 0.514
Pollen limitation: (hand - open) / hand	Pasture	1000 m	X^2 = 1.266	p = 0.260
		500 m	X^2 = 0.776	p = 0.378
		750 m	X^2 = 0.566	p = 0.451
	Agriculture	1000 m	X^2 = 2.996	p = 0.083
		500 m	X^2 = 2.412	p = 0.120
		750 m	X^2 = 1.745	p = 0.186
	Semi-Natural	1000 m	X^2 = 2.372	p = 0.123
		500 m	X^2 = 2.427	p = 0.119
		750 m	X^2 = 1.754	p = 0.185

References
Aleixo, K.P., Menezes, C., Imperatriz Fonseca, V.L. & Silva, C.I. da. (2017) Seasonal availability
of floral resources and ambient temperature shape stingless bee foraging
behavior (Scaptotrigona aff. depilis). *Apidologie*, **48**, 117–127.

Artz, D.R. & Nault, B.A. (2011) Performance of Apis mellifera, Bombus impatiens, and
Peponapis pruinosa (Hymenoptera: Apidae) as Pollinators of Pumpkin. *Journal of
Economic Entomology*, **104**, 1153–1161.

Bänsch, S., Tscharntke, T., Wünschiers, R., Netter, L., Brenig, B., Gabriel, D., *et al.* (2020)
Using ITS2 metabarcoding and microscopy to analyse shifts in pollen diets of honey
bees and bumble bees along a mass-flowering crop gradient. *Molecular Ecology*, **29**,
5003–5018.

Baskin, J.M. & Baskin, C.C. (2018) Pollen limitation and its effect on seed germination. *Seed
Science Research*, **28**, 253–260.

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software*, **67**, 1–48.

Baylis, K., Lichtenberg, E.M. & Lichtenberg, E. (2021) Economics of Pollination. *Annual Review
of Resource Economics*, **13**, 335–354.

Brochu, K.K., Dyke, M.T. van, Milano, N.J., Petersen, J.D., McArt, S.H., Nault, B.A., *et al.* (2020)
Pollen defenses negatively impact foraging and fitness in a generalist bee (Bombus
impatiens: Apidae). *Scientific Reports*, **10**, 3112.

Cane, J.H. & Tepedino, V.J. (2001) Causes and Extent of Declines among Native North
American Invertebrate Pollinators: Detection, Evidence, and Consequences.
Conservation Ecology, **5**, art1.

Centrella, M., Russo, L., Moreno Ramírez, N., Eitzer, B., Dyke, M. van, Danforth, B., *et al.*
(2020) Diet diversity and pesticide risk mediate the negative effects of land use change
on solitary bee offspring production. *Journal of Applied Ecology*, **57**, 1031-1042.

Connelly, H., Poveda, K. & Loeb, G. (2015) Landscape simplification decreases wild bee
pollination services to strawberry. *Agriculture, Ecosystems & Environment*, **211**, 51–56.

Cribari-Neto, F. & Zeileis, A. (2012) Beta Regression in R. *Journal of Statistical Software*, **34**, 1–
25.

Cropland Data Layer. (2022). USDA NASS.

De Luca, P.A. & Vallejo-Marín, M. (2013) What's the 'buzz' about? The ecology and
evolutionary significance of buzz-pollination. *Current Opinion in Plant Biology*, **16**, 429–
435.

Desjardins, È.-C. & De Oliveira, D. (2006) Commercial Bumble Bee Bombus impatiens
(Hymenoptera: Apidae) as a Pollinator in Lowbush Blueberry (Ericale: Ericaceae) Fields.
Journal of Economic Entomology, **99**, 443–449.

Dimou, M., Taraza, S., Thrasyvoulou, A. & Vasilakakis, M. (2008) Effect of bumble bee pollination on greenhouse strawberry production. *Journal of Apicultural Research*, **47**, 99–101.

Drummond, F. (2012) Commercial Bumble Bee Pollination of Lowbush Blueberry. *International Journal of Fruit Science*, **12**, 54–64.

Evans, E. (2017) From Humble Bee to Greenhouse Pollination Workhorse: Can We Mitigate Risks for Bumble Bees? *Bee World*, **94**, 34–41.

Furnessa, C.A. (1994) The extraction and identification of pollen from a beeswax statue. *Grana*, **33**, 49–52.

Gaines-Day, H.R. & Gratton, C. (2016) Crop yield is correlated with honey bee hive density but not in high-woodland landscapes. *Agriculture, Ecosystems & Environment*, **218**, 53–57.

Garibaldi, L.A., Carvalheiro, L.G., Leonhardt, S.D., Aizen, M.A., Blaauw, B.R., Isaacs, R., *et al.* (2014) From research to action: enhancing crop yield through wild pollinators. *Frontiers in Ecology and the Environment*, **12**, 439–447.

Garibaldi, L.A., Steffan-Dewenter, I., Kremen, C., Morales, J.M., Bommarco, R., Cunningham, S.A., *et al.* (2011) Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecology Letters*, **14**, 1062–1072.

Goulson, D., Nicholls, E., Botías, C. & Rotheray, E.L. (2015) Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, **347**, 1255957.

Grab, H., Blitzer, E.J., Danforth, B., Loeb, G. & Poveda, K. (2017) Temporally dependent pollinator competition and facilitation with mass flowering crops affects yield in co-blooming crops. *Scientific Reports*, **7**, 45296.

Grab, H., Branstetter, M.G., Amon, N., Urban-Mead, K.R., Park, M.G., Gibbs, J., *et al.* (2019) Agriculturally dominated landscapes reduce bee phylogenetic diversity and pollination services. *Science*, **363**, 282–284.

Grab, H., Poveda, K., Danforth, B. & Loeb, G. (2018) Landscape context shifts the balance of costs and benefits from wildflower borders on multiple ecosystem services. *Proceedings of the Royal Society B: Biological Sciences*, **285**, 20181102.

Graham, K.K., Milbrath, M.O., Killewald, M., Soehnlen, A., Zhang, Y. & Isaacs, R. (2023) Identity and diversity of pollens collected by two managed bee species while in blueberry fields for pollination. *Environmental Entomology*, **52**, 907–917.

Graystock, P., Blane, E.J., McFrederick, Q.S., Goulson, D. & Hughes, W.O.H. (2016) Do managed bees drive parasite spread and emergence in wild bees? *International Journal for Parasitology: Parasites and Wildlife*, Including articles from the 25th WAAVP Conference, Liverpool, August 2015, **5**, 64–75.

Graystock, P., Yates, K., Evison, S.E.F., Darvill, B., Goulson, D. & Hughes, W.O.H. (2013) The Trojan hives: pollinator pathogens, imported and distributed in bumblebee colonies. *Journal of Applied Ecology*, n/a-n/a.

Gudowska, A., Cwajna, A., Marjańska, E. & Moroń, D. (2024) Pollinators enhance the production of a superior strawberry – A global review and meta-analysis. *Agriculture, Ecosystems & Environment*, **362**, 108815.

Guzman, A., Gaines-Day, H.R., Lois, A.N., Steffan, S.A., Brunet, J., Zalapa, J., *et al.* (2019) Surrounding landscape and spatial arrangement of honey bee hives affect pollen foraging and yield in cranberry. *Agriculture, Ecosystems & Environment*, **286**, 106624.

Hines, H.M., Cameron, S.A., & Deans, A.R. (2007) Nest architecture and foraging behavior in *Bombus pullatus* (Hymenoptera: Apidae), while comparisons to other tropical bumble bees. *Journal of the Kansas Entomological Society*, **80(1)**, 1-15.

Harder, L.D. (1988) Choice of individual flowers by bumble bees: interaction of morphology, time and energy. *Behaviour*, **104**, 60–77.

Inês da Silva, C., Victorino Nicolosi Arena, M., Cristina Mathias da Silva, E., Roberto Martines, M., Malaspina, O., Chiovatto, G., *et al.* (2024) Landscape and land use affect composition and nutritional values of bees' food. *Journal of Environmental Management*, **352**, 120031.

Inouye, D.W. (1978) Resource Partitioning in Bumblebees: Experimental Studies of Foraging Behavior. *Ecology*, **59**, 672–678.

Jones, G.D. (2014) Pollen analyses for pollination researh, acetolysis, **13**, 15.

Jones, G.D. & Bryant, V.M. (1998) Are all counts created equal? *Contributions Series-American Association. of Stratigraphic Palynologists Foundation*, **33**, 115–120.

Kapp, R.O., Davis, O.K. & King, J.E. (2000) *Pollen and spores*. 2nd edn. American Association of Stratigraphic Palynologists Foundation, Dallas, TX.

Kasso, T.M., Bruun Jensen, A., Johns, S., Roffet-Salque, M., Collins, M.J. & Enevold, R. (2024) Beescapes – extracting pollen from historical Danish beeswax to explore honeybee foraging. *Palynology*, **48**, 2288220.

Kasso, T.M., Enevold, R., Johns, S., Rangel-Piñeros, G., Taurozzi, A.J., Sutherland, A., *et al.* (2023) ArcHives—combined palynological, genomic and lipid analysis of medieval wax seals. *Heritage Science*, **11**.

Kennedy, C.M., Lonsdorf, E., Neel, M.C., Williams, N.M., Ricketts, T.H., Winfree, R., *et al.* (2013) A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecology Letters*, **16**, 584–599.

Khalifa, S.A.M., Elshafiey, E.H., Shetaia, A.A., El-Wahed, A.A.A., Algethami, A.F., Musharraf, S.G., *et al.* (2021) Overview of Bee Pollination and Its Economic Value for Crop Production. *Insects*, **12**, 688.

Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C., *et al.* (2007) Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences*, **274**, 303–313.

Lanterman, J., Reeher, P., Mitchell, R.J. & Goodell, K. (2019) Habitat Preference and
Phenology of Nest Seeking and Foraging Spring Bumble Bee Queens in Northeastern
North America (Hymenoptera: Apidae: Bombus). *The American Midland Naturalist*, **182**,
131.

Lanterman Novotny, J., Lybbert, A., Reeher, P., Mitchell, R.J. & Goodell, K. (2023) Bumble bee
banquet: Genus- and species-level floral selection by Midwestern Bombus. *Ecosphere*,
14, e4425.

Larson, B.M.H. & Barrett, S.C.H. (1999) The ecology of pollen limitation in buzz☐pollinated
Rhexia virginica (Melastomataceae). *Journal of Ecology*, **87**, 371–381.

Mallinger, R.E., Gaines-Day, H.R. & Gratton, C. (2017) Do managed bees have negative effects
on wild bees?: A systematic review of the literature. *PLoS ONE*, **12**.

Mallinger, R.E. & Gratton, C. (2015) Species richness of wild bees, but not the use of managed
honeybees, increases fruit set of a pollinator-dependent crop. *Journal of Applied
Ecology*, **52**, 323–330.

Martin, C.D., Fountain, M.T. & Brown, M.J.F. (2019) Varietal and seasonal differences in the
effects of commercial bumblebees on fruit quality in strawberry crops. *Agriculture,
Ecosystems & Environment*, **281**, 124–133.

McArt, S.H., Fersch, A.A., Milano, N.J., Truitt, L.L. & Böröczky, K. (2017a) High pesticide risk to
honey bees despite low focal crop pollen collection during pollination of a mass blooming
crop. *Scientific Reports*, **7**, 1–10.

McArt, S.H., Urbanowicz, C., McCoshum, S., Irwin, R.E. & Adler, L.S. (2017b) Landscape
predictors of pathogen prevalence and range contractions in US bumblebees.
Proceedings of the Royal Society B: Biological Sciences, **284**, 20172181.

Michener, C.D. (1974) *The Social Behavior of the Bees: A Comparative Study*. Harvard
University Press.

Miller, O., Hale, C., Richardson, L., Sossa, D., Iverson, A., McArt, S., *et al.* (2023) Commercial
Bombus impatiens colonies function as ecological traps for wild queens. *Journal of
Applied Ecology*, **60**, 592–600.

Osterman, J., Aizen, M.A., Biesmeijer, J.C., Bosch, J., Howlett, B.G., Inouye, D.W., *et al.* (2021)
Global trends in the number and diversity of managed pollinator species. *Agriculture,
Ecosystems & Environment*, **322**, 107653.

Paini, D.R. (2004) Impact of the introduced honey bee (Apis mellifera) (Hymenoptera: Apidae)
on native bees: A review. *Austral Ecology*, **29**, 399–407.

Petersen, J.D., Reiners, S. & Nault, B.A. (2013) Pollination Services Provided by Bees in
Pumpkin Fields Supplemented with Either Apis mellifera or Bombus impatiens or Not
Supplemented. *PLoS ONE*, **8**, e69819.

Petran, A. & Hoover, E. (2018) The Flowers of Fragaria ✕ ananassa. In *Horticultural Reviews*.
John Wiley & Sons, Ltd, pp. 1–31.

R Core Team. (2024) R: A language and environment for statistical computing.

Regal, P.J. (1982) Pollination by Wind and Animals: Ecology of Geographic Patterns. *Annual*
Review of Ecology and Systematics, **13**, 497–524.

Ricketts, T.H., Regetz, J., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C., Bogdanski, A.,
et al. (2018) Landscape effects on crop pollination services: are there general patterns?
Ecology Letters, 499–515.

Rucker, R.R., Thurman, W.N. & Burgett, M. (2012) Honey Bee Pollination Markets and the
Internalization of Reciprocal Benefits. *American Journal of Agricultural Economics*, **94**,
956–977.

Saifuddin, M. & Jha, S. (2014) Colony-Level Variation in Pollen Collection and Foraging
Preferences Among Wild-Caught Bumble Bees (Hymenoptera: Apidae). *Environmental*
Entomology, **43**, 393–401.

Sakagami, S.F., Akahira, Y. & Zucchi, R. (1967) Nest architecture and brood development in a
neotropical bumblebee,Bombus atratus. *Insectes Sociaux*, **14**, 389–413.

Smith, T.A., Boone, M., Choy, S., Evans, E., Everett, J.G., Palmer, J.E., *et al.* (2025) Answering
key bumble bee conservation questions by studying discovered wild nests: A *Bombus*
affinis case study. *Insect Conservation and Diversity*, **18**, 303–318.

Stanghellini, M.S., Ambrose, J.T. & Schultheis, J.R. (1998) Using Commercial Bumble Bee
Colonies as Backup Pollinators for Honey Bees to Produce Cucumbers and
Watermelons. *HortTechnology*, **8**, 590–594.

Stavert, J.R., Liñán-Cembrano, G., Beggs, J.R., Howlett, B.G., Pattemore, D.E. & Bartomeus, I.
(2016) Hairiness: the missing link between pollinators and pollination. *PeerJ*, **4**, e2779.

Stubbs, C.S. & Drummond, F.A. (2001) *Bombus impatiens* (Hymenoptera: Apidae): An
Alternative to *Apis mellifera* (Hymenoptera: Apidae) for Lowbush Blueberry Pollination.
Journal of Economic Entomology, **94**, 609–616.

Trillo, A., Herrera, J.M. & Vilà, M. (2018) Managed bumble bees increase flower visitation but
not fruit weight in polytunnel strawberry crops. *Basic and Applied Ecology*, **30**, 32–40.

Vaudo, A.D., Patch, H.M., Mortensen, D.A., Tooker, J.F. & Grozinger, C.M. (2016)
Macronutrient ratios in pollen shape bumble bee (Bombus impatiens) foraging strategies
and floral preferences. *Proceedings of the National Academy of Sciences*, **113**, E4035–
E4042.

Velthuis, H.H.W. & Doorn, A. van. (2006) A century of advances in bumblebee domestication
and the economic and environmental aspects of its commercialization for pollination.
Apidologie, **37**, 421–451.

Wood, T.J., Kaplan, I. & Szendrei, Z. (2018) Wild Bee Pollen Diets Reveal Patterns of Seasonal
Foraging Resources for Honey Bees. *Frontiers in Ecology and Evolution*, **6**, 210.

Zeileis, A. & Hothorn, T. (2002) Diagnostic Checking in Regresssion Relationships. *R News*, **2**,