Incentive Salience, not Psychomotor Sensitization or Tolerance, Drives Escalation of Cocaine Self-Administration in Heterogeneous Stock Rats

Jarryd Ramborger[1,2], Joseph Mosquera[1], Molly Brennan[1], Benjamin Sichel[1], Dyar Othman[1], Sonja Plasil, PhD[1], Elizabeth Sneddon, PhD[1], Selene Bonnet-Zahedi[1,3], Alex Morgan[1], Supakorn Chonwattanagul[1], Kathleen Bai[1], Lindsey China[1], Tu La[1], Lisa Maturin[1], Lieselot L. G. Carrette, PhD[1], Olivier George, PhD[1]

[1] Department of Psychiatry, University of California, San Diego, La Jolla, CA 92093, USA

[2] Department of Psychology, Arizona State University, Tempe, AZ 85287, USA

[3]Institut de Neurosciences de la Timone, Aix-Marseille Université, Marseille, France

Corresponding Author:

Name: Olivier George, PhD

Email: olgeorge@health.ucsd.edu

Present Address:

Department of Psychiatry, University of California, San Diego, La Jolla, CA 92093, USA

Abstract:

Sensitization and tolerance are two phenomena often studied independently

despite overlapping neurobiological substrates. Each has extensive research showing

their influence on the development and maintenance of addiction, but the degree to

which they drive escalation in cocaine self-administration is poorly understood. Using

self-administration, noncontingent methods, and pose-estimation, we demonstrate that

incentive salience, not psychomotor sensitization or tolerance, drives the escalation of

cocaine self-administration in heterogenous stock rats. Individual differences in

psychomotor sensitization or tolerance were found to have no effect on cocaine intake.

Incentive salience as measured by locomotion and active lever entrances per meter

occurring before the self-administration session began (pre-lever) during Short Access

was found to predict intake during Long Access. Both pre-lever locomotion and active

lever entrances per meter were found to increase during Long Access and after two-to-

three days of abstinence. Critically, rats with low pre-lever activity during Short Access

escalated both their intake and pre-lever measures by the end of Long Access to levels

comparable with high pre-lever activity rats who maintained their elevated responding.

These findings support the notion that incentive salience during Short Access is a

catalyst to escalated use and an early marker of addiction vulnerability. Moreover, they

suggest that individuals initially resistant to incentive salience can, with sufficient

exposure, become sensitized and escalate cocaine use to the same level as more

susceptible individuals. Analysis of pre-lever activity offers a novel longitudinal

behavioral marker to predict vulnerability and provides a framework for understanding

individual trajectories of addiction.

1. Introduction

Addiction is a chronic, relapsing disorder characterized by cycles of binge/intoxication, withdrawal/negative affect, and preoccupation/anticipation, leading to long-lasting neuroadaptations in stress, executive function, and reward processing [1,2]. Among the most studied adaptations to repeated drug exposure are tolerance, a reduced response to the drug, and sensitization, an enhanced response. Both processes have been demonstrated in preclinical models of addiction, but the expression of either phenomenon varies widely depending on experimental parameters such as drug of choice, method of exposure (e.g., noncontingent injections versus self-administration), schedule of access (e.g. Short Access, Long Access, Intermittent, etc.), and outcome variables (e.g. locomotion, physiological measures, and operant behavior) [3-13]. Despite evidence that tolerance and sensitization share overlapping neurobiological substrates, they are often investigated in isolation [14-17], leading to two competing models in the transition to addiction.

According to the hedonic homeostatic dysregulation theory, repeated drug use leads to tolerance to the positive reinforcing effects of drugs and the allostatic recruitment of stress circuitry, thereby shifting the hedonic set point and motivating continued use to restore homeostasis [1,18]. In contrast, the incentive-sensitization theory proposes that repeated use hypersensitizes neural circuits leading to *incentive salience*—drug-related cues become increasingly "wanting" even as the "liking" diminishes—and to *behavioral sensitization*—a heightened psychomotor response observed primarily in psychostimulants like amphetamines and cocaine [19]. There is no doubt that both phenomena contribute to the addiction cycle, however, the degree to

which sensitization and tolerance drive escalation of cocaine self-administration is

poorly understood. While a number of studies have demonstrated the expression of

tolerance or sensitization to cocaine with noncontingent methods of continuous,

repeated, or single infusions [7,9,20-23], or the use of different lengths of self-

administration access [3-6,24-26], none have in a longitudinal paradigm employing

noncontingent challenges, Short Access, and Long Access.

To address this gap, using an operant cocaine self-administration paradigm in

heterogeneous stock rats in conjunction with machine vision pose-estimation, we

assessed multiple behavioral indices to differentiate individual development of tolerance

or sensitization across time and conditions and the subsequent influence on cocaine

self-administration. These included: (1) locomotion following a single noncontingent

intravenous infusion of cocaine at three time points: before Short Access, after Short

Access 09, and after Long Access 14; (2) locomotion for 15 minutes before the self-

administration session begins (pre-lever); (3) drug-seeking during pre-lever and

Noncontingent sessions, measured as nose entrances into the active lever area per

meter; (4) cocaine intake within the first 15 minutes and (5) across the full session; and

(6) breakpoint on a progressive ratio schedule.

2. Materials and Methods

A more detailed account of sections 2.1-2.3 below has been published elsewhere

[27-28].

2.1.   Subjects

Heterogeneous stock rats (N=37; n=21 males; n=16 females) were obtained from the from the University of California, San Diego colony [29] and housed in pairs on a 12-hour light/dark cycle. Rats were anesthetized with 1–5% isoflurane for jugular vein catheterization using Micro-Renathane tubing (Braintree Scientific) connected to a guide cannula that exits through the back and secured with mesh and dental acrylic. Post-surgery, rats received flunixin subcutaneously (2.5 mg/kg) and cefazolin intramuscularly (330 mg/kg). Catheters were flushed daily with heparinized saline (10 U/ml, American Pharmaceutical Partners) mixed with 0.9% bacteriostatic sodium chloride (Hospira) and cefazolin (52.4 mg/0.2 ml). Catheter patency was tested intravenously with brevital sodium (1.5 mg, American Pharmaceutical Partners) at experiment start and end, with failing rats removed from the experiment. Rats analyzed demonstrated catheter patency. All procedures were conducted in strict adherence to the National Institutes of Health Guide for the Care and Use of Laboratory Animals and were approved by the Institutional Animal Care and Use Committees of UC San Diego.

2.2. Drugs

Cocaine HCl (National Institute on Drug Abuse, Bethesda, MD) is dissolved in 0.9% sterile saline. During self-administration sessions, a press of the active lever resulted in an intravenous infusion of 0.1ml at 0.5 mg/kg over six seconds by pump, followed by a 20-second timeout where the active lever is deactivated. During Noncontingent sessions, a single intravenous infusion of 0.3ml at 1.5 mg/kg (i.e. three times received in self-administration) was delivered in under five seconds. Intravenous injections were chosen to address the differences in absorption, clearance, and blood

concentrations between intravenous and intraperitoneal methods and the influence of

intravenous infusion rate on the development of sensitization [30-36].

2.3.   Intravenous Self-Administration

Over 8 weeks (Monday-Friday), rats participated in 30 self-administration

sessions with various lengths and ratios. Under fixed ratio one, rats participated in 10

two-hour Short Access sessions,15 six-hour Long Access sessions, and one-hour

PRESHOCK and SHOCK sessions. During SHOCK, rats had a 30% chance to have an

infusion paired with a footshock (0.3 mA for 0.5 sec) through the grid floor. Under

progressive ratio, rats participated in three six-hour sessions in which infusions followed

the equation $[5e(\text{injection number} \times 0.2)] - 5$. The breakpoint was defined as the last

ratio before a 60-minute period in which a ratio was not completed, in which case the

session ended. A complete experimental timeline is shown in **Fig. 1A**.

2.4.   Noncontingent Sessions

Seven days after surgeries, rats underwent the first noncontingent session

(Noncontingent 01). Rats were placed in the operant chamber and video recorded for

30 minutes to establish baseline measurements (Baseline 01). Rats were then given an

intravenous saline injection, placed back in the chamber, and recorded for 30 minutes

(Saline 01), followed by a noncontingent intravenous injection of cocaine (1.5 mg/kg)

and recorded for 30 minutes (Drug 01). The same procedures were replicated for the

subsequent noncontingent sessions following Short Access 09 (Noncontingent 02) and

Long Access 14 (Noncontingent 03).

2.5.   Video Recording and Processing

2.5.1. *Raspberry Pi Video Recording*

Video recordings of sessions were conducted as described previously [37] with a

few alterations. A static FFmpeg build (ffmpeg-6.0-armhf-static;

https://johnvansickle.com/ffmpeg/) was installed onto each Raspberry Pi for video

processing to reduce video size threefold without sacrifice to video quality. All sessions

were recorded for 30 minutes. During self-administration sessions, levers were

extended in the chamber after 15 minutes of recording. Each csv file was processed in

RStudio (R version 4.3.1 (2023-06-16 ucrt)) to ensure consistent length and alignment

to the lever extension, resulting in ~25 minutes per csv, or ~12.5 minutes for both pre-

and post-lever data. Out of a possible 1,110 csv files (37 rats over 30 sessions), 1,084

are present in the study due to various issues such as catheter or infrared light

disconnections. For eight rats during Long Access 04 the Med Associates program was

initiated too early, resulting in no pre-lever data. For Noncontingent sessions, R

processing resulted in ~25 minutes per csv. Out of a possible 333 csv files, 327 are

present in the data, as six failed to track during SLEAP processing.

2.5.2. Social LEAP Estimates Animal Poses (SLEAP) Modeling

A pose-estimation model was trained using SLEAP [38] on ~5200 labeled frames

over 89 videos from past cohorts utilizing the top-down configuration and a unet

backbone. Thirteen nodes were tracked: nose, both ears, neck, four paws, base (where

tail met the body), active and inactive levers, active lever cue light, and the catheter

node, which was chosen as the model anchor point. A second model was trained to

track the first Noncontingent session, where the rats were mistakenly not plugged into

their catheter lines. This model focused on only six nodes to accelerate training: nose,

both ears, neck, base, and the catheter as the anchor node, utilizing ~2200 labeled

frames from 40 videos.

2.5.3. *High Throughput Computing on Open Science Grid (OSG)*

Video recordings were processed on the OSG Consortium supercomputer

system [39-42]. A python code with SLEAP command line interface functions processed

each video to identify one instance (track) using an "instance" identification method and

Hungarian approach between frames. Results of each node for each frame per second

inference registered as (x, y) coordinates were followed by an interpolation method to fill

missing data points (https://sleap.ai/notebooks/Analysis_examples.html).

2.6.   Dependent variable calculation

Calculations were conducted in RStudio version 4.3.1. Euclidian distances were

used to calculate movement of the catheter (locomotion) and the nose (nose motion)

nodes frame by frame and converted from pixels to centimeters using Image2 [43] and

the known measurements of the ENV-005 grid. Given the barrel distortion of a fisheye

lens, the conversion factor was computed by measuring the grid floor at five separate

segments and averaging the two length measurements (left and right), averaging three

width measurements (top, middle, and bottom), and taking the average of the two,

resulting in ~0.111cm/px. Locomotion variables are the sum of the distances traveled

and then smoothed by a window of 10 (10 FPS, i.e., 0.33 seconds), then converted to

meters. Locomotion (m) pre-lever corresponds to the locomotion measured for the 15

minutes before the levers are extended into the chamber. Active and inactive lever

entrances were calculated by producing a bounding box from the position of the (x,y)

nodes, 30 +/- x and 18 +/- y, and counting each entrance of the nose node into the lever

areas. Active and inactive lever entrances per meter are the total lever entrances

normalized by locomotion. Cocaine infusions are the total infusions delivered during the

session, and cocaine infusions first 15 minutes are those that occurred during the video

recordings. Session number corresponds to an individual session (e.g. Short Access

01). Session type refers to experimental phase (e.g. Long Access). Deprivation refers to

the time between the end of one session and the beginning of the next, where 18 Hours

translates to consecutive days, 66 Hours as a two-day weekend, and 90 Hours as a

three-day weekend. Pre-lever variables were examined by deprivation to account for

abstinence effects that could facilitate sensitization [6,14-17,20,33-35].

2.7. Statistical Analyses

Statistical comparisons were made by fitting a linear mixed effects model utilizing

the lmerTest package [44-45] or a generalized linear mixed effects model utilizing the

glmmTMB package [46] in R. Given that linear and generalized linear mixed-effects

models are inherently robust to moderate deviations from normality and extreme values,

outliers for all variables were winsorized at the 95th percentile to minimize their

influence while retaining potentially meaningful behavioral variability. Inferential tests of

fixed effects for linear mixed models were conducted using Type III ANOVA with

Satterthwaite degrees of freedom approximations (Type III ANOVA test) which have

been shown to control for Type I errors in hierarchical models [47], while for generalized

linear models a Type III Wald chi-square analysis of deviance was conducted (Type III

Wald test). Tests for model assumptions and fit were analyzed with the

simulateResdiduals() function from the DHARMa package [48]. The model fitting

process and chosen models of each analysis can be found in the Supplementary

Material with best adherence to suggestions previously published [49] but prioritizing the

results of DHARMa tests. Transformations of response variables where necessary were

achieved using Box-Cox or Yeo-Johnson after identifying the most effective lambda

using powerTransform() from the car package and reported at two decimal places [50].

Post-hoc analyses of all significance results were conducted using contrasts(), while

direct comparisons to baseline used test(), both within the emmeans package and

calculate the pairwise comparisons of the estimated marginal means of mixed effects

models and are therefore model-derived ratios or mean differences [51]. All p-values

were adjusted by the Benjamini-Yekutieli (BY) method due to its consideration of

dependence and correlations in nested data such as mixed models [52]. From the

rstatix package, Student's t-tests were conducted with t_test() with Bonferroni

adjustments and correlations with cor_test() by either Pearson or Spearman Rank as

determined by a normality test with shapiro_test() [53]. Plots represent the raw or

percent difference from baseline.

3. Results

3.1. Cocaine Infusions

3.1.1. Total cocaine infusions escalate during both Short and Long Access

To determine whether cocaine intake changed during Short Access, a Type III

Wald test was performed. A significant main effect of session number was observed

($\chi^2(9)=100.94$, $p<.001$). Post-hoc analyses revealed all sessions to have significantly

more cocaine infusions in comparison to Short Access 01. These results confirm that

cocaine intake increased as early as Short Access 02 (**Fig. 1B**).

To determine whether cocaine intake changed during Long Access, a Type III

Wald test was performed. A significant main effect of session number was identified

($\chi^2(13)=173.01$, $p<.001$). Post-hoc comparisons indicated that Long Access 09 through

Long Access 14 had significantly more infusions relative to Long Access 01. These

results show that cocaine intake continued to escalate into the third week of Long

Access (**Fig. 1B**).

3.1.2. Cocaine infusions during the first 15 minutes escalate across sessions

To evaluate whether cocaine infusions during the initial 15 minutes of each

session escalated, a Type III Wald test was conducted. A significant main effect of

session number was identified (χ^2(23)=304.87, $p<.001$). Post-hoc analyses revealed

that cocaine infusions were significantly greater from Short Access 07 through Long

Access 14 compared to Short Access 01. Additionally, when compared to Long Access

01, only sessions Long Access 09, Long Access 13, and Long Access 14 showed

significantly elevated infusions (**Fig. 1C**). These results indicate that escalation of

cocaine intake during the first 15 minutes emerged during the later stages of the Short

Access period and continued throughout Long Access, particularly after a period of

deprivation.

Figure 1. Experimental timeline and cocaine infusions during Short (ShA) and

Long Access (LgA). (A.) Experimental timeline with 66 hours (pink) and 90 hours

(blue) indicating instances of deprivation. **(B.)** Cocaine infusions during the entire

session. **(C.)** Cocaine infusions during the first 15 minutes of the session. Plots

represent means and SEM. *$p<0.05$, **$p<0.01$, ***$p<.001$ vs. Short Access 01; ##$p<.01$,

##$p<.001$ vs. Long Access 01.

3.2.   Locomotion During Noncontingent Sessions

3.2.1. Locomotion increases after noncontingent cocaine administration with no

evidence of psychomotor sensitization or tolerance after Short or Long Access to

cocaine self-administration.

To determine if raw locomotion differs between Noncontingent sessions following

a noncontingent intravenous infusion of cocaine, a Type III ANOVA test was performed.

A significant main effect of session number was found ($F_{(8, 290.88)}=17.38$, $p<.001$).

Post-hoc analyses revealed significant differences only within Noncontingent sessions,

as each Drug session had significantly greater locomotion than their respective Baseline

and Saline sessions (**Fig. 2A**).

To assess the change in the locomotor response to noncontingent cocaine

administration, data was normalized to each rat's baseline and analyzed as percent

difference from the respective Baseline sessions. A Type III ANOVA revealed a

significant main effect of session number ($F_{(5, 176.97)}=16.36$, $p<.001$). Post-hoc

analyses revealed similar significant differences, where Drug sessions were significantly

greater than their respective Saline and Baseline sessions, but no significant differences

between Drug sessions (**Fig. 2B**). These results show that locomotion did not increase

in successive Noncontingent sessions following a noncontingent intravenous infusion of

cocaine at the population level.

3.2.2. Rats differentially develop psychomotor sensitization or tolerance after a history

of cocaine self-administration

To assess each individual rat's response to successive noncontingent infusions,

rats were grouped by their locomotion difference from Drug 01 to Drug 03. Rats who

increased were labeled "Sensitization" and rats who *decreased* were labeled as

"Tolerance". The locomotion in each Drug session was then analyzed by a percentage

change from Baseline (Drug 01). Following a 0.13 lambda Box-Cox power

transformation, a Type III Wald test found no significant main effect of behavioral

expression ($\chi^2(1)$=4.8, p=.059), but a significant effect of session number ($\chi^2(1)$=6.47,

p=.031) and their interaction ($\chi^2(1)$=15.2, p<.001). Post-hoc analyses revealed rats in

the Tolerance group to have significantly decreased in their locomotion as early as

Noncontingent 02, with a further decrease in Noncontingent 03. Similarly, although not

significantly increased from baseline, rats in the Sensitization group showed greater

locomotion in Noncontingent 02 than the Tolerance group that progressed further in

Noncontingent 03. These results demonstrate markedly divergent profiles in the

psychomotor response to cocaine that begin as early as the first nine Short Access

sessions and continue through Long Access (**Fig. 2C**).

3.2.3. *Psychomotor sensitization and tolerance show no effect on cocaine self-administration*

To test whether the development of psychomotor sensitization or tolerance had an effect on cocaine self-administration over the entire session, a Type III Wald test was performed for both Short and Long Access. During Short Access, a main effect of session number was found ($x^2(9)=64.36$, $p<.001$), but not of behavioral expression ($x^2(1)=0.98$, $p=.967$), or their interaction ($x^2(9)=8.72$, $p=.967$. Similarly, in Long Access, a main effect of session number was found ($x^2(13)=107.44$, $p<.001$), but not of behavioral expression ($x^2(1)=0.47$, $p=1$), or their interaction ($x^2(13)=11.54$, $p=1$). These results suggest that the development of psychomotor sensitization or tolerance to the effect of cocaine has no effect on cocaine intake during Short or Long Access (**Fig. 2D**).

To determine if the development of psychomotor sensitization or tolerance has an effect on cocaine infusions in the first 15 minutes, a Type III Wald test was performed. A main effect of session number was found ($x^2(23)=154.13$, $p<.001$), but not of behavioral expression ($x^2(1)=0.003$, $p=1$), or their interaction ($x^2(23)=30.78$, $p=.356$). These results confirm that the development of psychomotor sensitization or tolerance to the effect of cocaine has no effect on the loading behavior during cocaine self-administration (**Fig. 2E**).

Figure 2. Psychomotor sensitization and tolerance show no effect on cocaine self-administration. (A.) Raw locomotion (m) across Noncontingent sessions. *###p<.001 vs Baseline; ***p<.001 vs Saline.* **(B.)** Locomotion (m) percent difference from Baseline. *#p<.05, ##p<.01, ###p<.001 vs Baseline; *p<.05, ***p<.001 vs Saline.* **(C.)** Locomotion (m) percentage difference from Baseline (Drug 01) grouped by behavioral expression. Drug 01 < Drug 03, Sensitization (n = 14); Drug 01 ≥ Drug 03, Tolerance (n = 22). *#p<.05, ###p<.001 vs Baseline; Within: $p<.05 vs Drug 02;*

Between: ***$p<.001$ vs Tolerance Drug 03. **(D-E.)** Cocaine infusions over the entire

session (D.) and within the first 15 minutes (E.) grouped by behavioral expression. Plots

represent means and SEM.

3.3. **Pre-Lever Activity**

3.3.1. Locomotion pre-lever increases from Short to Long Access and predicts cocaine

intake during the first 15 minutes of self-administration

To determine if locomotion pre-lever increased over the experiment, a Type III

Wald test was performed. A significant main effect of session number was found

following a 0.29 lambda Box-Cox power transformation ($\chi^2(23)=204.7$, $p<.001$). Post-

hoc analyses revealed significant increases at Short Access 07, Short Access 08, Long

Access 09, and Long Access 13 in comparison with Short Access 01 (**Fig. 3A**). These

results indicate that locomotion pre-lever increased from Short to Long Access.

To assess the influence of session type and deprivation on locomotion pre-lever,

a separate analysis was conducted following a 0.28 lambda Box-Cox transformation. A

Type III Wald test found significant main effects of session type ($\chi^2(1)=11.93$, $p=.002$),

deprivation ($\chi^2(2)=14$, $p=.002$), and their interaction ($\chi^2(2)=27.57$, $p<.001$). Within Short

Access, post-hoc analyses revealed locomotion pre-lever at 90 hours was significantly

greater than 18 hours. Within Long Access, 66 and 90 hours were found to be

significantly greater than 18 hours. Interactively, at 18 hours there was a significant

decrease in locomotion pre-lever from Short to Long Access. At 66 hours, there was a

significant increase from Short to Long Access. Overall, locomotion pre-lever was

significantly greater in Long Access than in Short Access across deprivation (**Fig. 3B**).

These results show that locomotion pre-lever increases with drug history and hours of

deprivation.

To assess locomotion pre-lever predictability of cocaine intake in the first 15

minutes of the session and break point, correlation analyses were performed. Between

locomotion pre-lever and cocaine infusions during the first 15 minutes, a positive

correlation was found during Short Access ($r_s(4222)=.5$, $p=.002$) (Figure 3.C.), and Long

Access ($r(4.29)=.59$, $p<.001$, 95% CI [0.33, 0.77]) (Figure 3.E.), but not during SHOCK

($r_s(6952)=.18$, $p=.298$) (Figure 3.G.). Between locomotion pre-lever and breaking point,

no correlations were found in both Progressive Ratio 01 ($r_s(7497)=.04$, $p=.84$) or

Progressive Ratio 02 ($r_s(6689)=.21$, $p=.219$) (**Fig. 3D & 3F**). These results indicate that

locomotion pre-lever is predictive of cocaine intake during the first 15 minutes in Short

and Long Access, but is not predictive of the motivation to seek cocaine as measured

by breaking point.

3.3.2. Active lever entrances per meter pre-lever escalate from Short to Long Access

and predict cocaine intake during the first 15 minutes of self-administration

Initial analyses indicated that nose entrances into the active lever area may be

predictive of intake, however, there were large individual variability in locomotion. To

account for animals with higher general activity having a greater probability of entering

the active lever area by chance, we normalized the number of active lever area

entrances to the overall locomotion in the chamber (active lever entrances per meter).

To determine if active lever entrances per meter increased over the experiment, a Type

III Wald test was performed. A significant main effect of session number was found

($\chi^2(23)$=234.44, p<.001). Post-hoc comparisons with Short Access 01 revealed

significant increases in all sessions from Short Access 07 to Long Access 14, excluding

Long Access 01, 03, and 07 (**Fig. 3H**). These results confirm that active lever entrances

per meter increased over the experiment.

To assess the effects of session type and deprivation on active lever entrances

per meter, a separate Type III Wald test was performed. No effect of deprivation was

found ($\chi^2(2)$=1.7, p=.89), but a significant main effect of session type ($\chi^2(1)$=13.66,

p<.001), and their interaction ($\chi^2(2)$=17.47, p<.001). Post-hoc analyses revealed no

significant differences between deprivation levels within Short Access. Within Long

Access, significant increases were found at 66 and 90 hours in comparison to 18 hours.

Interactively, at every level of deprivation there was found to be a significant increase

from Short to Long Access (**Fig. 3I**). These results confirm that active lever entrances

per meter increase with drug experience, with the strongest increases occurring after

longer periods of abstinence.

To assess active lever entrances per meter predictability of cocaine intake in the

first 15 minutes of the session and break point, correlation analyses were performed.

Between active lever entrances per meter and cocaine infusions during the first 15

minutes, a significant positive correlation was found during Short Access (r_s(3002)=.64,

p<.001) (**Fig. 3J**), Long Access (r(4.65)=.62, p<.001, 95% CI [0.37, 0.79]) (**Fig. 3L**), and

SHOCK (r_s(5199)=.38, p=.019) (**Fig. 3N**). Between active lever entrances per meter and

break point, a significant positive correlation was found during Progressive Ratio 01

$(r_s(5173)=.33, p=.046)$ (**Fig. 3K**), but not Progressive Ratio 02 $(r_s(7985)=.05, p=.753)$

(**Fig. 3M**). These results show that active lever entrances per meter pre-lever are

predictive of both cocaine intake within the first 15 minutes and motivation measured as

break point following Short Access, but not after Long Access.

Locomotion (m) Pre-Lever

Active Lever Entrances per Meter Pre-Lever

Figure 3. Pre-lever activity increases with Long Access, hours of deprivation, and is predictive of cocaine infusions within the first 15 minutes of self-administration. (A.) Locomotion pre-lever across experiment. *$p<.05$, **$p<.01$ vs Short Access 01. **(B.)** Comparison of deprivation levels within and between session types. Within session type: ***$p<.001$ vs 18 hours; between session type and deprivation: ##$p<.01$ vs 18 hours Long Access, ###$p<.001$ vs 66 hours Short Access; between session type: $$$$$$p<.001$ vs Short Access. **(C.)** Mean locomotion pre-lever during Short Access correlates with mean cocaine infusions during first 15 minutes of Short Access. **(D.)** Locomotion pre-lever during Progressive Ratio 01 does not correlate with break point. **(E**.**)** Mean locomotion pre-lever during Long Access correlates with mean cocaine infusions during first 15 minutes of Long Access. **(F.)** Locomotion pre-lever during Progressive Ratio 02 does not correlate with break point. **(G.)** Locomotion pre-lever during SHOCK does not correlate with cocaine infusions during first 15 minutes of SHOCK. **(H.)** Active lever entrances per meter pre-lever across experiment. *$p<.05$, **$p<.01$, ***$p<.001$ vs ShA01. **(I.)** Comparison of deprivation levels within and between session types. Within session type: ***$p<.001$ vs 18 hours; between session type and deprivation: ##$p<.01$ vs 18, 66, and 90 hours Short Access; between session type: $$$$$$p<.001$ vs Short Access. **(J.)** Mean active lever entrances per meter pre-lever during Short Access correlates with mean cocaine infusions during first 15 minutes of Short Access. **(K.)** Active lever entrances per meter pre-lever during Progressive Ratio 01 correlates with break point. **(L.)** Mean active lever entrances per meter pre-lever during Long Access correlates with mean cocaine infusions during first 15 minutes of Long Access. **(M.)** Active lever entrances per meter pre-lever during Progressive Ratio 02

does not correlate with break point. **(N.)** Active lever entrances per meter pre-lever

during SHOCK correlates with cocaine infusions during first 15 minutes of SHOCK.

Plots A, B, H, and I represent mean and SEM. Correlation plots E and L are by Pearson,

while remaining correlations are by Spearman Rank.

3.4.   **Lever Entrances During Noncontingent Sessions**

3.4.1. Lever entrances per meter as percent difference from Noncontingent 01 show a

preference for the active lever after a history of self-administration

To determine if seeking behavior was expressed during Noncontingent sessions,

entrances per meter into the active and inactive lever areas were analyzed. Considering

the levers never extend during Noncontingent sessions they were treated as pre-lever

sessions. Examination of raw data showed low variation within Noncontingent sessions

and therefore all three sessions (e.g. Baseline, Saline, and Drug) were grouped to

analyze nose entrances into the active and inactive lever areas per meter.

Noncontingent 01 was then treated as the Baseline, and Noncontingent 02 and 03

active and inactive lever entrances per meter for each rat were normalized by the group

mean of Noncontingent 01 to account for high individual variation. Following a 0.23

lambda Yeo-Johnson power transformation, a Type III Wald test indicated a significant

main effect of session number ($\chi^2(1)=12.46$, $p=.002$), but not a significant main effect of

levers ($\chi^2(1)=3.12$, $p=.162$) or their interaction ($\chi^2(1)=3.49$, $p=.162$). Post-hoc analyses

revealed Noncontingent 03 to be significantly greater than Noncontingent 02, prompting

a paired Student's t-test between the active and inactive levers in Noncontingent 03

where active was found to be greater than inactive. Direct comparisons to Noncontingent 01 found that the inactive lever entrances per meter in Noncontingent 02 were significantly less than Noncontingent 01, while the active lever entrances per meter in Noncontingent 03 were significantly greater than baseline. (**Fig. 4A**). These results indicate that by Noncontingent 03 rats demonstrate preferential seeking behavior toward the drug-associated active lever area.

To visualize the development of interest in the active lever, density heat maps of each rat's nose node coordinates (x, y) for all sessions within each Noncontingent session were overlain. Bounding boxes corresponding to the active and inactive lever areas and the rounded mean lever entrances into each lever area were also plotted (Noncontingent 01 active: M=22.2, SD=13.9; inactive: M=23.4, SD=12.3; Noncontingent 02 active: M=21.8, SD=16.1; inactive: M=18.4, SD=16.5; Noncontingent 03 active: M=43.5, SD=42.1; inactive: M=24.7, SD=22.1). Student's t-tests between active and inactive lever entrances within Noncontingent sessions revealed only Noncontingent 03 following a 0.23 lambda Yeo-Johnson power transformation achieved significance, with active greater than inactive (**Fig. 4B**). These results highlight the development of interest in the active lever after a history or self-administration as the high activity regions migrate from the inactive side of the chamber in Noncontingent 01 to the active side in Noncontingent 03.

3.4.2. Pre-lever activity during Short Access sessions 07-09 predicts active lever entrances per meter during Noncontingent sessions.

To determine if the interest in the active lever was developed during self-

administration, rats were grouped as "High" (n=19) or "Low" (n=18) by a median-split z-

score of their pre-lever activity measures (locomotion and active lever entrances per

meter) during Short Access 07-09. These sessions were chosen as they were the first to

show significance compared to Short Access 01 and are the last Short Access sessions

before Noncontingent 02. Following a 0.2 lambda Yeo-Johnson power transformation, a

Type III Wald test indicated no main effect of session number ($\chi^2(1)=4.81$, $p=.059$), but a

significant effect of pre-lever activity score ($\chi^2(1)=29.53$, $p<.001$) and their interaction

($\chi^2(1)=9.39$, $p=.006$). Post-hoc analyses revealed that both Noncontingent 02 and 03 in

the High pre-lever activity group had significantly greater active lever entrances per

meter than baseline, and versus the Low pre-lever activity group in Noncontingent 02.

For the Low pre-lever activity group, Noncontingent 02 was significantly less than both

baseline and Noncontingent 03. There was no difference between groups in

Noncontingent 03 (**Fig. 4C**). These results confirm that rats with high pre-lever activity

during Short Access 07-09 led to significant seeking in Noncontingent 02 that persisted

in Noncontingent 03. Furthermore, the Low pre-lever activity group developed an

interest during Long Access before Noncontingent 03.

3.4.3. Pre-lever activity during Short Access sessions 07-09 predicts cocaine self-

administration in Short and Long Access

To determine if pre-lever activity during Short Access sessions 07-09 predicts

cocaine intake over the entire session in Short Access, a Type III Wald test was

performed. A significant main effect was found in session number ($\chi^2(9)=125.52$,

$p<.001$), but not a significant effect of pre-lever activity score ($\chi^2(1)=3.51$, $p=.169$) or

their interaction ($\chi^2(9)=16.9$, $p=.169$) (**Fig. 4D**). These results show that cocaine intake

did not differ between groups in Short Access.

To determine if pre-lever activity during Short Access sessions 07-09 predicts

cocaine intake over the entire session in Long Access, a Type III Wald test was

performed. Significant main effects were found in session number ($\chi^2(13)=62.03$,

$p<.001$), pre-lever activity score ($\chi^2(1)=13.49$, $p<.001$) and their interaction

($\chi^2(13)=28.79$, $p=.015$). Post-hoc analyses revealed the High pre-lever activity group to

have significantly greater cocaine infusions over the entire session in all Long Access

sessions except Long Access 11, 13, and 14 (**Fig. 4D**). These results confirm that pre-

lever activity during Short Access 07-09 was initially predictive of cocaine intake during

Long Access, but that groups eventually converged.

To determine if pre-lever activity during Short Access sessions 07-09 predicts

cocaine intake within the first 15 minutes of the session, a Type III Wald test was

performed. A significant main effect was found in session number ($\chi^2(23)=208.9$,

$p<.001$), but not in pre-lever activity score ($\chi^2(1)=0.85$, $p=.742$) or their interaction

($\chi^2(23)=35.32$, $p=.134$) (**Fig 4E**). These results suggest that pre-lever activity during

Short Access 07-09 is not predictive of loading behavior between groups.

Considering the incongruency of the significant association between pre-lever

variables and cocaine intake variables (**Fig. 3**), and the previous nonsignificant results

between pre-lever activity groups and cocaine infusions in the first 15 minutes (**Fig. 4D**),

correlations were performed between cocaine intake variables and pre-lever activity z-

scores in Short Access and three separate phases of Long Access. For cocaine

infusions during the entire session and pre-lever activity, significant positive correlations

were found in Short Access ($r_s(2960)=.65$, $p<.001$; **Fig. 4F**), Long Access 01-05

($r_s(3284)=.61$, $p<.001$; **Fig. 4G**), Long Access 06-10 ($r_s(3422)=.59$, $p<.001$; **Fig. 4H**), but

not during Long Access 11-14 ($r_s(7013)=.17$, $p=.318$; **Fig. 4I**). For cocaine infusions

during the first 15 minutes and pre-lever activity, significant positive correlations were

similarly found in Short Access ($r_s(3094)=.63$, $p<.001$; **Fig. 4J**), Long Access 01-05

($r_s(3301)=.61$, $p<.001$; **Fig. 4K**), Long Access 06-10 ($r_s(4023)=.52$, $p<.001$; **Fig. 4L**), but

not during Long Access 11-14 ($r_s(6735)=.2$, $p=.231$; **Fig. 4M**). These results confirm that

rats who showed markedly increased pre-lever activity during Short Access sessions

07-09 were more likely to consume more cocaine during the first 15 minutes and over

the entire session in both Short and Long Access, but converged by the end of Long

Access. Furthermore, these results support those shown in **Fig. 4C,** where no

significant difference was found in active lever entrances per meter in Noncontingent

03, which occurs following Long Access 14.

Figure 4. Pre-lever activity during Short Access 07-09 predicts cocaine intake in Short and Long Access and seeking behavior during Noncontingent sessions.

(A.) Active (green) and inactive (red) lever entrances per meter as percentage difference from Baseline (Noncontingent 01). ##p<.01, ###p<.001 vs Baseline; *p<.05 vs inactive lever. **(B.)** Nose (*x, y)* pixel coordinate heatmaps of all rats within each Noncontingent session in relation to active (green) and inactive (red) Levers. Each plot represents ~109 sessions overlain revealing high and low activity zones. Numbers within lever bounding boxes are the mean sum of active and inactive lever entrances for all rats within each Noncontingent session. *p<.05 vs inactive lever. **(C.)** Active lever entrances per meter percent difference from Baseline (Noncontingent 01) grouped by pre-lever activity z-score during Short Access 07-09. ###p<.001 vs Baseline; ***p<.001 vs Noncontingent 02 Low group. **(D-E.)** Cocaine infusions over the entire session (D.) and within the first 15 minutes (E.) grouped by Short Access 07-09 pre-lever activity z-score. *p<.05, **p<.01, ***p<.001, ###p<.001 vs Low. **(F-I.)** Cocaine infusions over the entire session correlated using Spearman Rank with Short Access 07-09 pre-lever activity z-score in Short Access (F.), Long Access 01-05 (G.), Long Access 06-10 (H.), but not during Long Access 11-14 (I.). **(J-M.)** Cocaine infusions within the first 15 minutes correlated using Spearman Rank with Short Access 07-09 pre-lever activity z-score in Short Access (J.), Long Access 01-05 (K.), Long Access 06-10 (L.), but not during Long Access 11-14 (M.). Plots represent means and SEM.

4. Discussion

Here we demonstrate that cocaine infusions over the full session and within the first 15 minutes escalated during Short and Long Access in heterogeneous stock rats. Locomotion increased after a noncontingent cocaine infusion, and group-level analysis showed no signs of psychomotor sensitization or tolerance following Short Access (Noncontingent 02) or Long Access (Noncontingent 03). Individual analyses revealed strong divergence with some rats developing psychomotor sensitization and others showing tolerance, though neither influenced the level of cocaine self-administration. Pre-lever locomotion and active lever entrances per meter escalated from Short to Long Access, increased after instances of deprivation, and predicted early cocaine intake. Rats with high pre-lever activity during Short Access sessions 07-09 escalated lever entries across Noncontingent sessions and earned more infusions during Short Access and first two weeks of Long Access than low pre-lever activity rats, but these differences disappeared by Long Access 13 as low-activity rats escalated to similar levels. Together, the data dissociate psychomotor sensitization and tolerance from cocaine intake and identify pre-lever activity as a robust predictor of cocaine intake and escalation.

Rats increased total cocaine intake as early as the second Short Access session and within the first 15 minutes as early as Short Access 07, while exhibiting significant escalation of intake after two weeks of Long Access. These findings align with our previous reports that heterogenous stock rats show increased cocaine intake during Short and Long Access [28], and the larger literature that Long Access induces escalation of intake [3,4,24,54].

Rats showed individual differences in the propensity to develop psychomotor

sensitization or tolerance to noncontingent cocaine administration. One alternative

interpretation is that analyzing only ambulatory activity may misclassify some rats

displaying stereotypic behavior, including head bobbing and weaving, sniffing or

gnawing at high cocaine doses [8,55]. However, this is unlikely for several reasons.

First, the dose used (1.5 mg/kg) is known to produce minimal stereotypic behavior and

robust sensitization [6,33-35]. Second, the nose motion (a proxy for head bobbing,

weaving, sniffing or gnawing) was highly correlated with locomotion (r_s(606)=.93,

p<.001), suggesting very few stereotypic behaviors. Third, visual inspection of the

videos show that the most tolerant rat (M2677) did not exhibit stereotypic behavior while

the most sensitized rat (M2668) shows pronounced psychomotor sensitization. A

possible explanation for the difference between sensitized vs. tolerant rats is that rats

classified as sensitized consumed less cocaine, while tolerant rats consumed more.

However, intake analyses do not support this hypothesis: the two groups showed no

differences in cocaine intake across the Short or Long Access session or within the first

15 minutes. These findings challenge the idea that Short Access produces only

sensitization and Long Access only tolerance to the effects of cocaine [1,3-6,18,24], and

they show that neither tolerance nor sensitization reliably predict future cocaine intake.

Pre-lever variables locomotion and active lever entrances per meter increased

from Short to Long Access and intensified following two-to-three days of abstinence.

Similarly, rats showed a significant increase in the active lever area entrances per meter

in Noncontingent 03 in comparison to Noncontingent 01, suggesting sensitization of

drug-seeking behavior. In contrast to measures of psychomotor sensitization or

tolerance expression, both locomotion and active lever entrances per meter were highly

predictive of intake. These findings align with the incentive salience literature, showing

that repeated drug use leads to salience attribution to cues and environments that

particularly manifests following a period of abstinence [5,6,15-16,19-20,33-35,57-64].

To investigate the individual propensity to attribute salience to drug-associated

cues, a median z-score of pre-lever activity was calculated. Short Access sessions 07–

09 were selected as they occurred within the same week (i.e., no deprivation periods)

and were the first sessions in which both pre-lever variables (**Fig. 3A & 3H**) and

infusions within the first 15 minutes (**Fig. 1C**) reached significance. Comparisons

between High and Low pre-lever activity groups revealed that High pre-lever activity

promoted an early escalation of cocaine intake during Long Access, but ultimately both

High and Low pre-lever activity groups reached the same level of escalation after two

weeks. One possible explanation is that the Low pre-lever activity group in Short Access

was slower in developing incentive salience, and once expressed, their intake escalated

to levels comparable to the High pre-lever activity group. Indeed, the significant

difference in active lever entrances per meter between activity groups in Noncontingent

02 and its absence in Noncontingent 03 supports this interpretation. These findings

align with previous reports that individuals differ in their susceptibility to incentive

salience attribution [55,58,60-61], and that incentive salience may act as a catalyst of

escalation and an early predictor of vulnerability [26]. Another interpretation is that

negative affect or allostatic load may be driving the incentive salience and escalation in

Long Access [1,3,54,56].

The phenotype we observed (rats with high pre-lever activity measured as

elevated active lever entrances per meter and locomotion) shares conceptual parallels

with the sign-tracker/goal-tracker framework [58]. Sign-trackers preferentially interact

with a discrete cue (the lever CS), while goal-trackers orient to the site of reward

delivery. Sign-trackers have been shown to initially self-administer more cocaine [62]

and choose cocaine over food [65]. However, important differences exist: sign-tracking

is defined in a purely Pavlovian context, with lever pressing having no instrumental

consequence, whereas our measure reflects operant-conditioned lever-directed

exploration. Notably, recent work has shown that sign-/goal-tracker status does not

reliably predict overall cocaine self-administration or escalation, although sign-trackers

may show greater resistance to punishment [66]. Thus, our results extend the sign-

/goal-tracker framework into an operant cocaine paradigm, highlighting that cue-bias

and responsivity can shape early intake and drive escalation of use even in resistant

populations with prolonged access.

The present study has a few limitations. One is the lack of a control group without

periods of abstinence (no weekends). A separate group participating in all sessions in

succession without instances of deprivation could further confirm the influence of

deprivation on incentive salience incubation. However, noncontingent sessions 02 and

03 did not follow a period of deprivation, and expressions of incentive salience and both

psychomotor sensitization and tolerance were still observed. Another limitation is the

use of pose-estimation over manual scoring of behavior could result in inaccuracies.

However, pose-estimation with SLEAP is remarkably accurate [38,67], and the instance

rendered sample videos display precise tracking of the catheter and nose nodes (see

supplementary videos).

In summary, this report demonstrates that individual differences in psychomotor

sensitization or tolerance had no influence on cocaine intake. In contrast, sensitization

of incentive salience during Short Access acts as a catalyst or early predictor of

escalation in cocaine self-administration during the first two weeks of Long Access, but

does not predict established escalation of cocaine intake after three weeks of Long

Access. Moreover, individuals initially resistant to incentive salience can, with sufficient

exposure, become sensitized and escalate cocaine use to the same level as more

susceptible individuals. Expression of incentive salience sensitization as reflected in

pre-lever activity measures of active lever entrances per meter and locomotion and their

subsequent influence on cocaine intake supports the idea that incentive salience and

psychomotor sensitization are mediated by distinct mechanisms [6-7,21,34]. The

development of the incentive salience profiles is accelerated by periods of abstinence

(weekend), which allows for incubation of craving. The measure of approach toward the

active lever area using video recording before the start of the self-administration session

may represent a novel longitudinal behavioral marker of addiction vulnerability that can

precisely track the motivation for the drug with very high accuracy without using any

operant measures, the drug itself, or any other training or conditioning that could

interfere with the drug self-administration. Such a measure may also be useful for

medication development to allow dissociation between drug-dependent and drug-

independent effect of experimental medication on drug seeking and taking.

Data Availability Statement

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

This research was done using services provided by the OSG Consortium [39-42] which is supported by the National Science Foundation awards #2030508 and #1836650.

Author Contributions

Jarryd Ramborger: Manuscript drafting and revising, data collection, experiment execution, experimental design, data analysis and interpretation

Joseph Mosquera: Data collection, experiment execution

Molly Brenna: Data collection, experiment execution

Benjamin Sichel: Data collection, experiment execution

Dyar Othman: Data collection, experiment execution

Sonja Plasil, PhD: Data collection, experiment execution

Elizabeth Sneddon, PhD: Data collection, experiment execution

Selene Bonnet-Zahedi: Data collection, experiment execution

Alex Morgan: Data collection, experiment execution

Supakorn Chonwattanagul: Data collection, experiment execution

Kathleen Bai: Data collection, experiment execution

Lindsey China: Data collection, experiment execution

Tu La: Data collection, experiment execution

Lisa Maturin: Data collection

Lieselot L. G. Carrette, PhD: Experimental design, manuscript drafting and revising

Olivier George, PhD: Manuscript drafting and revising, experimental design, data
analysis and interpretation, funding and resources, final approval

Funding

Funding was supported by DA043799 and the Preclinical Addiction Research
Consortium.

Competing Interests

689   Authors declare no competing interests.

References

1. Koob, G. F., & Le Moal, M. (1997). Drug abuse: Hedonic homeostatic dysregulation. *Science, 278*(5335), 52–58. https://doi.org/10.1126/science.278.5335.52

2. Koob, G. F., & Volkow, N. D. (2016). Neurobiology of addiction: a neurocircuitry analysis. *The lancet. Psychiatry*, *3*(8), 760–773. https://doi.org/10.1016/S2215-0366(16)00104-8.

3. Ahmed, S. H., & Koob, G. F. (1998). Transition from moderate to excessive drug intake: Change in hedonic set point. *Science, 282*(5387), 298-300. https://doi.org/10.1126/science.282.5387.298.

4. Ben-Shahar, O., Ahmed, S. H., Koob, G. F., & Ettenberg, A. (2004). The transition from controlled to compulsive drug use is associated with a loss of sensitization. *Brain Research, 995*(1), 46–54. https://doi.org/10.1016/j.brainres.2003.09.053

5. Calipari, E. S., Ferris, M. J., Zimmer, B. A., Roberts, D. C. S., & Jones, S. R. (2013). Temporal pattern of cocaine intake determines tolerance vs sensitization of cocaine effects at the dopamine transporter. *Neuropsychopharmacology, 38*(12), 2385–2392. https://doi.org/10.1038/npp.2013.136.

6. Carr, C. C., Ferrario, C. R., & Robinson, T. E. (2020). Intermittent access cocaine self-administration produces psychomotor sensitization: Effects of withdrawal, sex, and cross-sensitization. *Psychopharmacology, 237*(6), 1795–1812. https://doi.org/10.1007/s00213-020-05500-4.

7. Dalia AD, Norman MK, Tabet MR, Schlueter KT, Tsibulsky VL, Norman AB (1998) Transient amelioration of the sensitization of cocaine-induced behaviors in rats

by the induction of tolerance. Brain Res 797:29–34.

https://doi.org/10.1016/S0006-8993(98)00323-0.

8.   Flagel, S. B., & Robinson, T. E. (2007). Quantifying the psychomotor activating

effects of cocaine in the rat. *Behavioural Pharmacology, 18*(4), 297–302.

https://doi.org/10.1097/FBP.0b013e3281f522a4.

9.   Lett, B. T. (1989). Repeated exposures intensify rather than diminish the

rewarding effects of amphetamine, morphine, and cocaine.

Psychopharmacology, 98(3), 357-362. https://doi.org/10.1007/BF00451688.

10.   Porrino, L. J. (1993). Functional consequences of acute cocaine treatment

depend on route of administration. *Psychopharmacology, 112*(3), 343–351.

https://doi.org/10.1007/BF02244948.

11.   Swain, Y., Muelken, P., LeSage, M. G., Gewirtz, J. C., & Harris, A. C. (2018).

Locomotor activity does not predict individual differences in morphine self-

administration in rats. *Pharmacology Biochemistry and Behavior*, 166, 48-56.

https://doi.org/10.1016/j.pbb.2018.01.008.

12.   Vanderschuren, L. J. M. J., Schmidt, E. D., De Vries, T. J., Van Moorsel, C. A. P.,

Tilders, F. J. H., & Schoffelmeer, A. N. M. (1999). A single exposure to

amphetamine is sufficient to induce long-term behavioral, neuroendocrine, and

neurochemical sensitization in rats. *Journal of Neuroscience, 19*(21), 9579–9586.

https://doi.org/10.1523/JNEUROSCI.19-21-09579.1999

13.   Vanderschuren, L. J. M. J., De Vries, T. J., Wardeh, G., Hogenboom, F. A. C. M.,

& Schoffelmeer, A. N. M. (2001). A single exposure to morphine induces long-

lasting behavioural and neurochemical sensitization in rats. *European Journal of*

Neuroscience, 14(8), 1533–1538. https://doi.org/10.1046/j.0953-816x.2001.01771.x

14. Hammer, R. P. Jr., Egilmez, Y., & Emmett-Oglesby, M. W. (1997). Neural mechanisms of tolerance to the effects of cocaine. *Behavioural Brain Research*, *84*(1-2), 225-239. https://doi.org/10.1016/s0166-4328(97)83332-3.

15. Kalivas PW, Sorg BA, Hooks MS (1993). The pharmacology and neural circuitry of sensitization to psychostimulants. Behav Pharmacol. 1993;4(4):315-334. PMID: 11224200.

16. Vanderschuren, L. J. M. J., & Kalivas, P. W. (2000). Alterations in dopaminergic and glutamatergic transmission in the induction and expression of behavioral sensitization: A critical review of preclinical studies. *Psychopharmacology, 151*(2), 99–120. https://doi.org/10.1007/s002130000493.

17. Vezina, P. (2004). Sensitization of midbrain dopamine neuron reactivity and the self-administration of psychomotor stimulant drugs. *Neuroscience & Biobehavioral Reviews, 27*(8), 827–839. https://doi.org/10.1016/j.neubiorev.2003.11.001.

18. Solomon, R. L. (1980). The opponent-process theory of acquired motivation: The costs of pleasure and the benefits of pain. *American Psychologist*, 35(8), 691-712. https://doi.org/10.1037/0003-066X.35.8.691

19. Robinson, T. E. & Berridge, K. C.(1993). The neural basis of drug craving: An incentive-sensitization theory of addiction. *Brain Research Reviews*, 18(3), 247-291. https://doi.org/10.1016/0165-0173(93)90013-P.

759   20.   Ciccocioppo, R., Martin-Fardon, R., & Weiss, F. (2004). Stimuli associated with a

760   single cocaine experience elicit long-lasting cocaine-seeking. *Nature*

761   *Neuroscience, 7*(5), 495-496. https://doi.org/10.1038/nn1239.

762   21.   Izenwasser, S., & French, D. (2002). Tolerance and sensitization to the

763   locomotor-activating effects of cocaine are mediated via independent

764   mechanisms. *Pharmacology Biochemistry and Behavior, 73*(4), 877–882.

765   https://doi.org/10.1016/s0091-3057(02)00942-5.

766   22.   Kalivas, P. W., & Alesdatter, J. E. (1993). Involvement of N-methyl-D-aspartate

767   receptor stimulation in the ventral tegmental area and amygdala in behavioral

768   sensitization to cocaine. *Journal of Pharmacology and Experimental*

769   *Therapeutics, 267*(1), 486-495. Retrieved from

770   http://jpet.aspetjournals.org/content/267/1/486.abstract.

771   23.   Ungless, M., Whistler, J., Malenka, R. *et al.* Single cocaine exposure *in vivo*

772   induces long-term potentiation in dopamine neurons. *Nature* 411, 583–587

773   (2001). https://doi.org/10.1038/35079077

774   24.   Ahmed, S. H., & Cador, M. (2006). Dissociation of psychomotor sensitization

775   from compulsive cocaine consumption. Neuropsychopharmacology, 31(3), 563–

776   571. https://doi.org/10.1038/sj.npp.1300834.

777   25.   Kawa AB, Bentzley BS, Robinson TE. Less is more: prolonged intermittent

778   access cocaine self-administration produces incentive-sensitization and

779   addiction-like behavior. Psychopharmacology (Berl). 2016 Oct;233(19-20):3587-

780   602. doi: 10.1007/s00213-016-4393-8. Epub 2016 Aug 2. PMID: 27481050;

781   PMCID: PMC5023484.

782   26.   Kawa AB, Valenta AC, Kennedy RT, Robinson TE. Incentive and dopamine

783   sensitization produced by intermittent but not long access cocaine self-

784   administration. Eur J Neurosci. 2019 Aug;50(4):2663-2682. doi:

785   10.1111/ejn.14418. Epub 2019 May 2. PMID: 30968487; PMCID: PMC6742545.

786   27.   Carrette, L. L. G., de Guglielmo, G., Kallupi, M., Maturin, L., Brennan, M.,

787   Boomhower, B., et al.. (2021). The Cocaine and Oxycodone Biobanks, two

788   repositories from genetically diverse and behaviorally characterized rats for the

789   study of addiction. *eNeuro*, *8*(3), ENEURO.0033-21.2021.

790   https://doi.org/10.1523/ENEURO.0033-21.2021.

791   28.   Giordano de Guglielmo, Carrette, L. L. G., Kallupi, M., Brennan, M., Boomhower,

792   B., Maturin, L., et al.. (2024). Large-scale characterization of cocaine addiction-

793   like behaviors reveals that escalation of intake, aversion-resistant responding,

794   and breaking-points are highly correlated measures of the same construct. *eLife*,

795   *12*, RP90422. https://doi.org/10.7554/eLife.90422

796   29.   Solberg Woods LC, Palmer AA (2019) Using heterogeneous stocks for fine-

797   mapping genetically complex traits. *Methods Mol Biol* 2018:233–247.

798   doi: 10.1007/978-1-4939-9581-3_11.

799   30.   Argilli, E., Sibley, D. R., Malenka, R. C., England, P. M., & Bonci, A. (2008).

800   Mechanism and time course of cocaine-induced long-term potentiation in the

801   ventral tegmental area. *The Journal of Neuroscience, 28*(37), 9092–9100.

802   https://doi.org/10.1523/JNEUROSCI.1001-08.2008.

803   31.   Booze, R. M., Lehner, A. F., Wallace, D. R., Welch, M. A., & Mactutus, C. F.

804   (1997). Dose-response cocaine pharmacokinetics and metabolite profile

following intravenous administration and arterial sampling in unanesthetized,

freely moving male rats. *Neurotoxicology and Teratology*, 19(1), 7–15.

https://doi.org/10.1016/S0892-0362(96)00180-8.

32. Pan, H.-T., Menacherry, S., & Justice, J. B., Jr. (1991). Differences in the

pharmacokinetics of cocaine in naive and cocaine-experienced rats. *Journal of*

Neurochemistry, 56(4), 1299–1306. https://doi.org/10.1111/j.1471-

4159.1991.tb11429.x

33. Samaha, A. N., Li, Y., & Robinson, T. E. (2002). The rate of intravenous cocaine

administration determines susceptibility to sensitization. *The Journal of*

neuroscience : the official journal of the Society for Neuroscience, 22(8), 3244–

3250. https://doi.org/10.1523/JNEUROSCI.22-08-03244.2002.

34. Samaha, A.-N., Mallet, N., Ferguson, S. M., Gonon, F., & Robinson, T. E. (2004).

The rate of cocaine administration alters gene regulation and behavioral

plasticity: Implications for addiction. *The Journal of Neuroscience*, *24*(28), 6362–

6370. https://doi.org/10.1523/JNEUROSCI.1424-04.2004.

35. Samaha, A.-N., & Robinson, T. E. (2005). Why does the rapid delivery of drugs to

the brain promote addiction? *Trends in Pharmacological Sciences, 26*(2), 82-87.

https://doi.org/10.1016/j.tips.2004.12.007.

36. Woolverton, W. L., & Wang, Z. (2004). Relationship between injection duration,

transporter occupancy, and reinforcing strength of cocaine. *European Journal of*

Pharmacology, *486*(3), 251–257. https://doi.org/10.1016/j.ejphar.2004.01.003.

37. Ramborger, J., Kalra, S., Mosquera, J., Smith, A. C. W., & George, O. (2024).

High quality, high throughput, and low-cost simultaneous video recording of 60

828  animals in operant chambers using PiRATeMC. *Journal of Neuroscience*

829  *Methods, 411*, 110270. https://doi.org/10.1016/j.jneumeth.2024.110270.

830 38. T.D. Pereira, N. Tabris, A. Matsliah, D. M. Turner, J. Li, S. Ravindranath, et al..

831  Sleap: A deep learning system for multi-animal pose tracking. *Nature Methods*,

832  19(4), 2022. https://doi.org/10.1038/s41592-022-01426-1.

833 39. OSG. (2006). OSPool. OSG. https://doi.org/10.21231/906P-4D78

834 40. Pordes, R., Petravick, D., Kramer, B., Olson, D., Livny, M., Roy, A., et al.. (2007).

835  The open science grid. J. Phys. Conf. Ser., 78, 012057.

836  https://doi.org/10.1088/1742-6596/78/1/012057.

837 41. Sfiligoi, I., Bradley, D. C., Holzman, B., Mhashilkar, P., Padhi, S., & Wurthwein, F.

838  (2009). The pilot way to grid resources using glideinWMS. 2009 WRI World

839  Congress on Computer Science and Information Engineering, 2, 428–432.

840  https://doi.org/10.1109/CSIE.2009.950.

841 42. OSG. (2015). Open Science Data Federation. OSG.

842  https://doi.org/10.21231/0KVZ-VE57

843 43. Rueden, C.T., Schindelin, J., Hiner, M.C. *et al.* ImageJ2: ImageJ for the next

844  generation of scientific image data. *BMC Bioinformatics* **18**, 529 (2017).

845  https://doi.org/10.1186/s12859-017-1934-z.

846 44. Bates D, Mächler M, Bolker B, Walker S (2015). "Fitting Linear Mixed-Effects

847  Models Using lme4." *Journal of Statistical Software*, **67**(1), 1–

848  48. doi:10.18637/jss.v067.i01.

849   45.   Kuznetsova A, Brockhoff PB, Christensen RHB (2017). "lmerTest Package: Tests

850   in Linear Mixed Effects Models." *Journal of Statistical Software*, **82**(13), 1–

851   26. doi:10.18637/jss.v082.i13.

852   46.   Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A,

853   et al. (2017). "glmmTMB Balances Speed and Flexibility Among Packages for

854   Zero-inflated Generalized Linear Mixed Modeling." *The R Journal*, 9(2), 378–

855   400. doi:10.32614/RJ-2017-066.

856   47.   Luke, S.G. Evaluating significance in linear mixed-effects models in R. *Behav*

857   *Res* **49**, 1494–1502 (2017). https://doi.org/10.3758/s13428-016-0809-y.

858   48.   Hartig, F. (2024). *DHARMa: Residual diagnostics for hierarchical (multi-level /*

859   *mixed) regression models* (Version 0.4.7) [R package]. CRAN.

860   https://doi.org/10.32614/CRAN.package.DHARMa.

861   49.   Meteyard, L., & Davies, R. A. I. (2020). Best practice guidance for linear mixed-

862   effects models in psychological science. *Journal of Memory and Language, 112*,

863   104092. https://doi.org/10.1016/j.jml.2020.104092.

864   50.   Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (3rd ed.).

865   Sage. https://www.john-fox.ca/Companion/.

866   51.   Lenth, R. V. (2025). *emmeans: Estimated marginal means, aka least-squares*

867   *means* (Version 1.10.7) [R package]. CRAN.

868   https://doi.org/10.32614/CRAN.package.emmeans.

869   52.   Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in

870   multiple testing under dependency. *The Annals of Statistics, 29*(4), 1165–1188.

871   https://doi.org/10.1214/aos/1013699998.

872   53.   Kassambara, A. (2023). rstatix: Pipe-friendly framework for basic statistical tests

873   (Version 0.7.2) [Computer software].

874   https://rpkgs.datanovia.com/rstatix/[](https://rpkgs.datanovia.com/rstatix/authors.h

875   tml).

876   54.   Ahmed, S. H., & Koob, G. F. (1999). Longlasting increase in the set point for

877   cocaine selfadministration after escalation in rats. *Psychopharmacology, 146*(3),

878   303–312. https://doi.org/10.1007/s002130051121.

879   55.   Flagel, S. B., Watson, S. J., Akil, H., & Robinson, T. E. (2008). Individual

880   differences in the attribution of incentive salience to a reward-related cue:

881   Influence on cocaine sensitization. *Behavioural Brain Research, 186*(1), 48–56.

882   https://doi.org/10.1016/j.bbr.2007.07.022

883   56.   Ahmed, S. H., Kenny, P. J., Koob, G. F., & Markou, A. (2002). Neurobiological

884   evidence for hedonic allostasis associated with escalating cocaine use. *Nature*

885   *Neuroscience, 5*(7), 625–626. https://doi.org/10.1038/nn872.

886   57.   Blanchard, M. M., Mendelsohn, D., & Stamp, J. A. (2009). The HR/LR model:

887   Further evidence as an animal model of sensation seeking. *Neuroscience &*

888   *Biobehavioral Reviews, 33*(7), 1145-1154.

889   https://doi.org/10.1016/j.neubiorev.2009.05.009.

890   58.   Flagel, S.B., Watson, S.J., Robinson, T.E. *et al.* Individual differences in the

891   propensity to approach signals vs goals promote different adaptations in the

892   dopamine system of rats. *Psychopharmacology* **191**, 599–607 (2007).

893   https://doi.org/10.1007/s00213-006-0535-8.

894   59.   Piazza, P. V., Deroche-Gamonent, V., Rouge-Pont, F., & Le Moal, M. (2000).

895   Vertical Shifts in Self-Administration Dose–Response Functions Predict a Drug-

896   Vulnerable Phenotype Predisposed to Addiction. *The Journal of Neuroscience*,

897   20(11), 4226-4232. https://doi.org/10.1523/JNEUROSCI.20-11-04226.2000.

898   60.   Post, R. M., Lockfeld, A., Squillace, K. M., & Contel, N. R. (1981). Drug-

899   environment interaction: Context dependency of cocaine-induced behavioral

900   sensitization. *Life Sciences*, *28*, 755–760. https://doi.org/10.1016/0024-

901   3205(81)90157-0.

902   61.   Saunders, B. T., & Robinson, T. E. (2010). A cocaine cue acts as an incentive

903   stimulus in some but not others: Implications for addiction. Biological Psychiatry,

904   67(8), 730–736. https://doi.org/10.1016/j.biopsych.2009.11.015

905   62.   Saunders, B. T., & Robinson, T. E. (2011). Individual variation in the motivational

906   properties of cocaine. *Neuropsychopharmacology, 36*(8), 1668–1676.

907   https://doi.org/10.1038/npp.2011.48.

908   63.   Stewart, J., & Badiani, A. (1993). Tolerance and sensitization to the behavioral

909   effects of drugs. *Behavioural Pharmacology, 4*(4), 289-312.

910   64.   Weiss, S. R. B., Post, R. M., Pert, A., Woodward, R., & Murman, D. (1989).

911   Context-dependent cocaine sensitization: Differential effect of haloperidol on

912   development versus expression. *Pharmacology, Biochemistry, and Behavior,*

913   *34*(3), 655-661. https://doi.org/10.1016/S0091-3057(89)80001-2.

914   65.   Tunstall, B. J., & Kearns, D. N. (2015). Sign-tracking predicts increased choice of

915   cocaine over food in rats. Behavioural Brain Research, 281, 222–228.

916   https://doi.org/10.1016/j.bbr.2014.12.034.

66.   Pohořalá, V., Enkel, T., Bartsch, D. *et al.* Sign- and goal-tracking score does not

correlate with addiction-like behavior following prolonged cocaine self-

administration. *Psychopharmacology* **238**, 2335–2346 (2021).

https://doi.org/10.1007/s00213-021-05858-z

67.   Pereira, T. D., Shaevitz, J. W., & Murthy, M. (2020). Quantifying behavior to

understand the brain. *Nature Neuroscience, 23*(12), 1537–1549.

https://doi.org/10.1038/s41593-020-00734-z.