Incentive Salience, not Psychomotor Sensitization or Tolerance, Drives Escalation

of Cocaine Self-Administration in Heterogeneous Stock Rats

Jarryd Ramborger[1,2], Joseph Mosquera[1], Molly Brennan[1], Benjamin Sichel[1], Dyar

Othman[1], Sonja Plasil, PhD[1], Elizabeth Sneddon, PhD[1], Selene Zahedi[1,3], Alex

Morgan[1], Supakorn Chonwattanagul[1], Kathleen Bai[1], Lindsey China[1], Tu La[1], Lisa

Maturin[1], Lieselot L. G. Carrette, PhD[1], Olivier George, PhD[1]

[1] Department of Psychiatry, University of California, San Diego, La Jolla, CA 92093,

USA

[2] Department of Psychology, Arizona State University, Tempe, AZ 85287, USA

[3]Institut de Neurosciences de la Timone, Aix-Marseille Université, Marseille, France

Corresponding Author:

Name: Olivier George, PhD

Email: olgeorge@health.ucsd.edu

Present Address:

Department of Psychiatry, University of California, San Diego, La Jolla, CA 92093, USA

Abstract:

Sensitization and tolerance are two phenomena often studied independently

despite overlapping neurobiological substrates. Each has extensive research showing

their influence on the development and maintenance of addiction, but the degree to

which they drive escalation in cocaine self-administration is poorly understood. Using

self-administration, intravenous noncontingent infusions, and pose-estimation machine

vision, we find that incentive salience, not psychomotor sensitization or tolerance, drives

the escalation of cocaine self-administration in heterogenous stock rats. Individual

differences in psychomotor sensitization or tolerance were found to have no effect on

cocaine intake. Incentive salience as measured by locomotion and active lever

entrances per meter traveled occurring before the self-administration session began

(pre-lever activity measures) during Short Access (2-hours) was found to predict intake

during Long Access (6-hours). Both pre-lever locomotion and active lever entrances per

meter were found to increase during Long Access and after two-to-three days of

abstinence. Critically, rats with low pre-lever activity during Short Access escalated both

their intake and pre-lever measures by the end of Long Access to levels comparable

with high pre-lever activity rats who maintained their elevated responding. These

findings support the notion that incentive salience during Short Access is a catalyst to

escalated use and an early marker of addiction vulnerability. Moreover, they suggest

that individuals initially resistant to incentive salience can, with sufficient exposure,

become sensitized and escalate cocaine use to the same level as more susceptible

individuals. Analysis of pre-lever activity offers a novel longitudinal behavioral marker to

predict vulnerability and provides a framework for understanding individual trajectories

of addiction.

1. Introduction

Addiction is a chronic, relapsing disorder characterized by cycles of binge/intoxication, withdrawal/negative affect, and preoccupation/anticipation, leading to long-lasting neuroadaptations in stress, executive function, and reward processing [1,2]. Within this framework, two of the most studied adaptations to repeated drug exposure are tolerance, a reduced response to the drug, and sensitization, an enhanced response. Both processes have been demonstrated in preclinical models of addiction, but the expression of either phenomenon varies widely depending on experimental parameters such as drug of choice, method of exposure (e.g., noncontingent injections versus self-administration), schedule of access (e.g. Short Access, Long Access, Intermittent, etc.), and outcome variables (e.g. locomotion, physiological measures, and operant behavior) [3-13]. Despite evidence that tolerance and sensitization share overlapping neurobiological substrates, they are often investigated in isolation [14-17], leading to two competing models in the transition to addiction.

According to the hedonic homeostatic dysregulation theory, 1) repeated drug use leads to tolerance to the positive reinforcing effects of drugs, and 2) the allostatic recruitment of stress circuitry, thereby shifting the hedonic set point and motivating continued use to restore homeostasis [1,18]. In contrast, the incentive-sensitization theory proposes that repeated use hypersensitizes neural circuits leading to 1) *incentive salience*—drug-related cues become increasingly "wanting" even as "liking" diminishes—and 2) *behavioral sensitization*—a heightened psychomotor response observed primarily in psychostimulants [19]. Both phenomena can contribute to the addiction cycle; however, the degree to which sensitization and tolerance drive

escalation of cocaine self-administration is poorly understood. While several studies

have demonstrated the expression of tolerance or sensitization to cocaine with

noncontingent methods of continuous, repeated, or single infusions [7,9,20-23], or the

use of different lengths of self-administration access [3-6,24-26], none have in a

longitudinal paradigm employing noncontingent challenges, Short Access, and Long

Access.

To address this gap, using an operant cocaine self-administration paradigm in

conjunction with machine vision pose-estimation, we assessed multiple behavioral

indices to differentiate individual development of tolerance or sensitization and the

subsequent influence on cocaine self-administration. These included: 1) locomotion

following a single noncontingent intravenous infusion of cocaine at three time points

(see **Fig. 1A** for timeline); 2) locomotion for 15 minutes before the self-administration

session begins (pre-lever); 3) drug-seeking during pre-lever and Noncontingent

sessions, measured as nose entrances into the active lever area per meter traveled; 4)

cocaine intake within the first 15 minutes and 5) full session; and 6) breakpoint on a

progressive ratio schedule. Psychomotor tolerance and sensitization will be assessed

by locomotion during Noncontingent sessions. Incentive salience will be assessed by

both locomotion and nose entrances into the active lever area per meter during the pre-

lever phase of self-administration.

2. Materials and Methods

Procedures in sections 2.1-2.3 below have been published elsewhere in more

detail [27-28]. ChatGPT4 was utilized for coding assistance and authors take full

responsibility for output.

2.1.   Subjects

Heterogeneous stock rats (N=37; n=21 males; n=16 females) obtained from the

UC San Diego colony [29] were used to mimic human genetic diversity and housed in

pairs on a 12-hour light/dark cycle. Rats were anesthetized with 1–5% isoflurane for

jugular vein catheterization using Micro-Renathane tubing (Braintree Scientific)

connected to a guide cannula that exits through the back and secured with mesh and

dental acrylic. Post-surgery, rats received flunixin subcutaneously (2.5 mg/kg) and

cefazolin intramuscularly (330 mg/kg). Catheters were flushed daily with heparinized

saline (10 U/ml, American Pharmaceutical Partners) mixed with 0.9% bacteriostatic

sodium chloride (Hospira) and cefazolin (52.4 mg/0.2 ml). Catheter patency was tested

intravenously with brevital sodium (1.5 mg, American Pharmaceutical Partners) at

experiment start and end, with failing rats removed from the experiment. All procedures

were conducted in strict adherence to the National Institutes of Health Guide for the

Care and Use of Laboratory Animals and were approved by the Institutional Animal

Care and Use Committees of UC San Diego.

2.2.   Drugs

Cocaine HCl (National Institute on Drug Abuse, Bethesda, MD) was dissolved in

0.9% sterile saline. During self-administration sessions, a press of the active lever

resulted in an intravenous infusion of 0.1ml at 0.5 mg/kg over six seconds by pump,

followed by a 20-second timeout where the active lever had no programmed effect.

During Noncontingent sessions, a single intravenous infusion of 0.3ml at 1.5 mg/kg (i.e.

three times the dose received in self-administration) was delivered in under five

seconds in the same operant chamber as self-administration to address influences of

environment and intravenous infusion rate on the development of sensitization [30-33].

2.3. Self-Administration

Over 8 weeks (Monday-Friday), rats participated in 30 self-administration

sessions of various lengths and ratios. Under fixed ratio one, rats participated in 10 two-

hour Short Access sessions,15 six-hour Long Access sessions, one one-hour Preshock,

and one one-hour Shock session. During Shock, each infusion had a 30% chance to be

paired with a footshock (0.3 mA for 0.5 sec) through the grid floor. Under progressive

ratio, rats participated in three six-hour sessions in which the response requirements for

an infusion followed the equation $[5e(\text{injection number} \times 0.2)] - 5$. The breakpoint was

defined as the last ratio before a 60-minute period in which a ratio was not completed, in

which case the session ended. A complete experimental timeline is shown in **Fig. 1A**.

2.4. Noncontingent Sessions

Seven days after surgeries, rats underwent the first noncontingent session

(Noncontingent 01) which occurred before Short Access 01. Rats were first video

recorded for 30 minutes to establish baseline measurements (Baseline 01), then given

an intravenous saline injection and recorded for 30 minutes (Saline 01), followed by a

noncontingent intravenous injection of cocaine (1.5 mg/kg) and recorded for 30 minutes

(Drug 01). The same procedures were replicated for the subsequent noncontingent

sessions.

2.5.   Video Recording and Processing

2.5.1. Video Recording

Video recordings were conducted as described previously [34] with a few

alterations. A static FFmpeg build (ffmpeg-6.0-armhf-static;

https://johnvansickle.com/ffmpeg/) was installed onto each Raspberry Pi for video

processing to reduce video size threefold without sacrifice to video quality. All sessions

were recorded for 30 minutes. During self-administration sessions, levers were

extended into the chamber after 15 minutes of recording. Each comma-separated value

(csv) file was processed in R version 4.3.1 using RStudio [35-36] to ensure consistent

length and alignment to the lever extension, resulting in ~25 minutes per csv, or ~12.5

minutes for both pre- and post-lever data. Out of 1,110 csv files (37 rats over 30

sessions), 26 were excluded for various issues such as catheter or infrared light

disconnections. For eight rats during Long Access 04 the Med Associates program was

initiated too early, resulting in no pre-lever data. For Noncontingent sessions, post-

processing resulted in ~25 minutes per csv. Out of 333 csv files, six files failed to track

during Social LEAP Estimates Animal Poses (SLEAP) processing.

2.5.2. SLEAP Modeling

A pose-estimation model was trained using SLEAP on ~5200 labeled frames

over 89 videos utilizing the top-down configuration and a unet backbone [37]. Thirteen

nodes were tracked: nose, ears, neck, four paws, base (where tail met the body), active

and inactive levers, active lever cue light, and the catheter node, which was chosen as

the model anchor point. A second model was trained to track Noncontingent 01 as rats

were mistakenly not plugged into their catheter lines. This model focused on six nodes

to accelerate training: nose, ears, neck, base, and catheter as the anchor node, utilizing

~2200 labeled frames from 40 videos.

2.5.3. High Throughput Computing on Open Science Grid

Video recordings were processed on the OSG Consortium supercomputer

system [38-41]. A python script with SLEAP command line functions processed each

video to identify one instance (track) using an "instance" identification method and

Hungarian approach between frames. Results of each node for each frame inference

registered as (x, y) coordinates were interpolated to fill missing data points

(https://sleap.ai/notebooks/Analysis_examples.html).

2.6.   Dependent and Independent Variables

Statistical calculations were conducted using RStudio. Euclidian distances

calculated movement of the catheter (locomotion) and the nose (nose motion) nodes

frame-by-frame and converted from pixels to centimeters using Image2 [42] and the

known measurements of the ENV-005 grid. Given the barrel distortion of a fisheye lens,

the conversion factor was computed by averaging two length measurements (left and

right) and three width measurements (top, middle, and bottom), resulting in

~0.111cm/px. Locomotion variables are the total distances traveled smoothed by a

window of 10 (10 FPS, i.e., 0.33 seconds) and converted to meters. Locomotion (m)

pre-lever is the distance traveled during the 15 minutes before the levers are extended

into the chamber. Active and inactive lever entrances were calculated by producing a

bounding box around the lever nodes (x,y) at 30 +/- x and 18 +/- y, and counting each

entrance of the nose node into the lever areas. Active and inactive lever entrances per

meter are the total entrances normalized by locomotion. Cocaine infusions are the total

infusions delivered during the session, and cocaine infusions first 15 minutes are those

that occurred during the video recordings. Session corresponds to an individual session

(e.g. Short Access 01). Session type refers to experimental phase (e.g. Long Access).

Abstinence refers to the time between the end of one session and the beginning of the

next, where 18 Hours translates to consecutive days, 66 Hours to a two-day weekend,

and 90 Hours to a three-day weekend. Pre-lever variables were examined by

abstinence to test for effects that could facilitate sensitization [6,14-17,20,30-34,43-44].

Analyses with sex as a factor were performed and reported in the Supplementary

Material as most effects were not sex dependent.

2.7.   Statistical Analyses

Statistical comparisons were made by fitting a linear mixed-effects model utilizing

the lmerTest package [45-46] or a generalized linear mixed-effects model utilizing the

glmmTMB package [47] in R. Given that linear and generalized linear models are

inherently robust to moderate deviations from normality and extreme values, outliers for

all dependent variables were winsorized at the 95th percentile to minimize their

influence while retaining potentially meaningful behavioral variability. Inferential tests of

fixed effects for linear models were conducted using Type III ANOVA with Satterthwaite

degrees of freedom approximations (Type III ANOVA test) which have been shown to

control for Type I errors in hierarchical models [48], while generalized linear models a

Type III Wald chi-square analysis of deviance was conducted (Type III Wald test). Model

assumption tests and fit were analyzed with the simulateResdiduals() function from the

DHARMa package [49]. The model fitting process and chosen models of each analysis

can be found in the Supplementary Material with best adherence to suggestions

previously published [50] but prioritizing the results of DHARMa tests. Transformations

of response variables where necessary (see Supplemental Material) were achieved

using Box-Cox or Yeo-Johnson after identifying the most effective lambda using

powerTransform() from the car package [51]. Post hoc analyses were conducted using

contrasts(), while direct baseline comparisons used test(), both within the emmeans

package and calculate the pairwise comparisons of the estimated marginal means and

are therefore model-derived ratios or mean differences [52]. All *p*-values were adjusted

by the Benjamini-Yekutieli (BY) method due to its consideration of dependence and

correlations in nested data such as mixed models [53]. From the rstatix package,

Student's t-tests were conducted with t_test() with Bonferroni adjustments, and

correlations with cor_test() by either Pearson or Spearman Rank as determined by a

normality test with shapiro_test() [54]. Plots represent the raw average values or

percent difference from baseline.

224 3. Results

225 3.1. Cocaine Infusions

3.1.1. Cocaine infusions escalate during both Short and Long Access

To determine whether cocaine intake changed during Short Access, a Type III

Wald test was performed. A significant main effect of session was observed

($\chi^2(9)=100.94$, *p*<.001). Post hoc analyses revealed all sessions to have significantly

more cocaine infusions in comparison to Short Access 01. These results confirm that

cocaine intake increased as early as Short Access 02 (**Fig. 1B**).

To determine whether cocaine intake changed during Long Access, a Type III

Wald test was performed. A significant main effect of session was identified

($\chi^2(13)=173.01$, *p*<.001). Post hoc comparisons indicated that Long Access 09 through

Long Access 14 had significantly more infusions relative to Long Access 01. These

results show that cocaine intake continued to escalate into the third week of Long

Access (**Fig. 1B**).

3.1.2. Cocaine infusions during the first 15 minutes escalate across sessions

To evaluate whether cocaine infusions during the initial 15 minutes of each

session escalated, a Type III Wald test was conducted. A significant main effect of

session was identified (χ^2(23)=304.87, p<.001). Post hoc analyses revealed that

cocaine infusions were significantly greater from Short Access 07 through Long Access

14 compared to Short Access 01. Additionally, when compared to Long Access 01, only

sessions Long Access 09, 13, and 14 showed significantly elevated infusions (**Fig. 1C**).

These results indicate that escalation of cocaine intake during the first 15 minutes

emerged at the end of Short Access and continued throughout Long Access.

Figure 1. Experimental timeline and cocaine infusions during Short (ShA) and Long Access (LgA). (A.) Experimental timeline with 66 hours (pink) and 90 hours (blue) indicating instances of abstinence. **(B.)** Cocaine infusions during the entire session. **(C.)** Cocaine infusions during the first 15 minutes of the session. Plots represent means and SEM. *$p<.05$, **$p<.01$, ***$p<.001$ vs. Short Access 01; ##$p<.01$, ##$p<.001$ vs. Long Access 01.

3.2. Locomotion During Noncontingent Sessions

3.2.1. Noncontingent cocaine administration increases locomotion with no evidence of psychomotor sensitization or tolerance following Short or Long Access cocaine self-administration.

To determine if raw locomotion differs between Noncontingent sessions following noncontingent administration of cocaine, a Type III ANOVA test was performed. A significant main effect of session was found ($F(8,290.88)=17.38$, $p<.001$). Post hoc analyses only found significant differences within each respective Noncontingent session. Specifically, Drug sessions had significantly greater locomotion than their respective Baseline and Saline sessions (**Fig. 2A**).

To assess the change in the locomotor response to noncontingent cocaine administration, data was analyzed as percent difference from each rat's respective Baseline session. A Type III ANOVA revealed a significant main effect of session ($F(5,176.97)=16.36$, $p<.001$). Post hoc analyses revealed that Drug sessions were significantly greater than their respective Saline and Baseline sessions, but no significant differences between Drug sessions (**Fig. 2B**). These results show that locomotion did not differ between Noncontingent sessions following noncontingent administration of cocaine at the population level.

3.2.2. Rats differentially develop psychomotor sensitization or tolerance after a history of cocaine self-administration

To assess each individual rat's response to noncontingent infusions, rats were

grouped by their locomotion difference from Drug 01 to Drug 03. Rats who *increased*

were labeled "Sensitization" and rats who *decreased* were labeled as "Tolerance".

Locomotion between Drug sessions was then analyzed by a percentage change from

baseline (Drug 01). Following a 0.13 lambda Box-Cox power transformation, a Type III

Wald test found no significant main effect of behavioral expression ($\chi^2(1)=4.8$, $p=.059$),

but a significant effect of session ($\chi^2(1)=6.47$, $p=.031$) and their interaction ($\chi^2(1)=15.2$,

$p<.001$). Post hoc analyses revealed that Tolerant rats significantly decreased in

locomotion in Drug 02, with a further decrease in Drug 03. Conversely, Sensitized rats

increased in Drug 03 and were greater than Tolerant rats in both Drug 02 and 03. These

results demonstrate divergent profiles in the psychomotor response to cocaine that

begins after just nine Short Access sessions and continues through Long Access (**Fig.**

2C). The effect observed was not due to stereotypies in Tolerant rats (**Fig. S1**).

3.2.3. Psychomotor sensitization and tolerance show no effect on cocaine self-

administration

To test whether the development of psychomotor sensitization or tolerance

influenced cocaine self-administration, a Type III Wald test was performed for both Short

and Long Access. During Short Access, a main effect of session was found

($\chi^2(9)=64.36$, $p<.001$), but not behavioral expression ($\chi^2(1)=0.98$, $p=.967$), or their

interaction ($\chi^2(9)=8.72$, $p=.967$. Similarly, in Long Access, a main effect of session was

found ($\chi^2(13)=107.44$, $p<.001$), but not behavioral expression ($\chi^2(1)=0.47$, $p=1$), or their

interaction ($\chi^2(13)=11.54$, $p=1$). These results suggest that the development of

psychomotor sensitization or tolerance to the effect of cocaine has no effect on cocaine

intake during Short or Long Access (**Fig. 2D**).

To determine if the development of psychomotor sensitization or tolerance

influences cocaine infusions in the first 15 minutes, a Type III Wald test was performed.

A main effect of session was found ($\chi^2(23)=154.13$, $p<.001$), but not behavioral

expression ($\chi^2(1)=0.003$, $p=1$), or their interaction ($\chi^2(23)=30.78$, $p=.356$). These results

confirm that the development of psychomotor sensitization or tolerance to the effect of

cocaine has no effect on loading behavior during cocaine self-administration (**Fig. 2E**).

Figure 2. Psychomotor sensitization and tolerance show no effect on cocaine

self-administration. (A.) Raw locomotion (m) across Noncontingent sessions.

###p<.001 vs Baseline*; ***p*<.001 vs Saline. **(B.)** Locomotion (m) percent difference

from Baseline. *#p*<.05, *###p*<.001 vs Baseline*; *p*<.05, ****p*<.001 vs Saline. **(C.)**

Locomotion (m) percentage difference from Baseline (Drug 01) grouped by behavioral

expression. Drug 01 < Drug 03, Sensitization (n = 14); Drug 01 > Drug 03, Tolerance (n

= 22). *#p*<.05, *###p*<.001 vs Baseline*;* Within behavioral expression groups: *$p*<.05,

316    $\$\$p<.01$ vs Drug 02; Between behavioral expression groups: $*p<.05$ vs Tolerance Drug

317   02, $***p<.001$ vs Tolerance Drug 03. **(D-E.)** Cocaine infusions over the entire session

318   (D.) and within the first 15 minutes (E.) grouped by behavioral expression. Plots

319   represent means and SEM.

3.3. Pre-Lever Activity

3.3.1. Locomotion pre-lever increases from Short to Long Access and predicts cocaine

        intake during the first 15 minutes of self-administration

324   To determine if locomotion pre-lever increased over the experiment, a Type III

325   Wald test was performed. A significant main effect of session was found following a 0.29

326   lambda Box-Cox power transformation ($\chi^2(23)=204.7$, $p<.001$). Post hoc analyses

327   revealed significant increases at Short Access 07, Short Access 08, Long Access 09,

328   and Long Access 13 in comparison with Short Access 01 (**Fig. 3A**). These results

329   indicate that locomotion pre-lever increased from Short to Long Access.

330   To assess the influence of session type and abstinence on locomotion pre-lever,

a Type III Wald test was conducted following a 0.28 lambda Box-Cox transformation.

Significant main effects of session type ($\chi^2(1)=11.93$, $p=.002$), abstinence ($\chi^2(2)=14$,

$p=.002$), and their interaction ($\chi^2(2)=27.57$, $p<.001$) were found. Within Short Access,

post hoc analyses revealed locomotion pre-lever at 90 hours was significantly greater

than 18 hours. Within Long Access, 66 and 90 hours was significantly greater than 18

hours. Interactively, locomotion pre-lever significantly decreased at 18 hours from Short

to Long Access. At 66 hours, there was a significant increase from Short to Long

Access. Overall, locomotion pre-lever was significantly greater in Long Access than in

Short Access (**Fig. 3B**). These results show that locomotion pre-lever increases with

drug history and hours of abstinence.

To assess locomotion pre-lever predictability of cocaine intake in the first 15

minutes of the session and break point, correlation analyses were performed. Between

locomotion pre-lever and cocaine infusions during the first 15 minutes, a positive

correlation was found during Short Access (r_s(4222)=.5, p=.002) (**Fig. 3C**), and Long

Access (r(4.29)=.59, p<.001, 95% CI [0.33, 0.77]) (**Fig. 3E**), but not during Shock (r-

$_s$(6952)=.18, p=.298) (**Fig. 3G**). Between locomotion pre-lever and breaking point, no

correlations were found in both Progressive Ratio 01 (r_s(7497)=.04, p=.84) or

Progressive Ratio 02 (r_s(6689)=.21, p=.219) (**Fig. 3D & 3F**). These results indicate that

locomotion pre-lever is predictive of cocaine intake during the first 15 minutes in Short

and Long Access, but is not predictive of breaking point.

3.3.2. Active lever entrances per meter pre-lever escalate from Short to Long Access

and predict cocaine intake during the first 15 minutes of self-administration

Initial analyses indicated that nose entrances into the active lever area may be

predictive of intake. However, there was large individual variability in locomotion. To

account for animals with higher general activity having a greater probability of entering

the active lever area by chance, active lever area entrances were normalized by

locomotion. To determine if active lever entrances per meter increased over the

experiment, a Type III Wald test was performed. A significant main effect of session was

found (χ^2(23)=234.44, p<.001). Post hoc comparisons with Short Access 01 revealed

significant increases in all sessions from Short Access 07 to Long Access 14, excluding

Long Access 01, 03, and 07 (**Fig. 3H**). These results confirm that active lever entrances

per meter increased over the experiment.

To assess the effects of session type and abstinence on active lever entrances

per meter, a Type III Wald test was performed. No effect of abstinence was found

(χ^2(2)=1.7, p=.89), but a significant main effect of session type (χ^2(1)=13.66, p<.001),

and their interaction (χ^2(2)=17.47, p<.001). Post hoc analyses revealed no significant

differences between abstinence levels within Short Access. Within Long Access,

significant increases were found at 66 and 90 hours in comparison to 18 hours.

Interactively, at every level of abstinence there was found to be a significant increase

from Short to Long Access (**Fig. 3I**). These results confirm that active lever entrances

per meter increase with drug experience, particularly after longer periods of abstinence.

To assess active lever entrances per meter predictability of cocaine intake in the

first 15 minutes of the session and break point, correlation analyses were performed.

Between active lever entrances per meter and cocaine infusions during the first 15

minutes, a significant positive correlation was found during Short Access (r_s(3002)=.64,

p<.001) (**Fig. 3J**), Long Access (r(4.65)=.62, p<.001, 95% CI [0.37, 0.79]) (**Fig. 3L**), and

Shock (r_s(5199)=.38, p=.019) (**Fig. 3N**). Between active lever entrances per meter and

break point, a significant positive correlation was found during Progressive Ratio 01 (r_s

(5173)=.33, p=.046) (**Fig. 3K**), but not Progressive Ratio 02 (r_s(7985)=.05, p=.753)

(**Fig. 3M**). These results show that active lever entrances per meter pre-lever are

predictive of both cocaine intake within the first 15 minutes and motivation measured as

break point following Short Access, but not after Long Access.

Locomotion (m) Pre-Lever

Active Lever Entrances per Meter Pre-Lever

Figure 3. Pre-lever activity increases with Long Access, hours of abstinence, and is predictive of cocaine infusions within the first 15 minutes of self-administration. (A.) Locomotion pre-lever across experiment. *$p<.05$, **$p<.01$ vs Short Access 01. **(B.)** Comparison of abstinence levels within and between session types. Within session type: **$p<.01$, ***$p<.001$ vs 18 hours; between session type and abstinence: ###$p<.001$ vs 18 hours Long Access, ###$p<.001$ vs 66 hours Short Access; between session type: $$$p<.01$ vs Short Access. **(C.)** Mean locomotion pre-lever during Short Access correlates with mean cocaine infusions during first 15 minutes of Short Access. **(D.)** Locomotion pre-lever during Progressive Ratio 01 does not correlate with break point. **(E.)** Mean locomotion pre-lever during Long Access correlates with mean cocaine infusions during first 15 minutes of Long Access. **(F.)** Locomotion pre-lever during Progressive Ratio 02 does not correlate with break point. **(G.)** Locomotion pre-lever during Shock does not correlate with cocaine infusions during first 15 minutes of Shock. **(H.)** Active lever entrances per meter pre-lever across experiment. *$p<.05$, **$p<.01$, ***$p<.001$ vs ShA01. **(I.)** Comparison of abstinence levels within and between session types. Within session type: ***$p<.001$ vs 18 hours; between session type and abstinence: ###$p<.001$ vs 18, 66, and 90 hours Short Access; between session type: $$$$p<.001$ vs Short Access. **(J.)** Mean active lever entrances per meter pre-lever during Short Access correlates with mean cocaine infusions during first 15 minutes of Short Access. **(K.)** Active lever entrances per meter pre-lever during Progressive Ratio 01 correlates with break point. **(L.)** Mean active lever entrances per meter pre-lever during Long Access correlates with mean cocaine infusions during first 15 minutes of Long Access. **(M.)** Active lever entrances per meter pre-lever during Progressive Ratio 02

408 does not correlate with break point. **(N.)** Active lever entrances per meter pre-lever

409 during Shock correlates with cocaine infusions during first 15 minutes of Shock. Plots A,

410 B, H, and I represent mean and SEM. Correlation plots E and L are by Pearson, while

411 remaining correlations are by Spearman Rank.

3.4. Lever Entrances During Noncontingent Sessions

3.4.1. Lever entrances per meter show a preference for the active lever after a history of self-administration

To determine if seeking behavior was expressed during Noncontingent sessions, entrances per meter into the active and inactive lever areas were analyzed. Considering the levers never extend during Noncontingent sessions they were treated as pre-lever sessions. Examination of raw data showed low variation within Noncontingent sessions and therefore all three sessions (e.g. Baseline, Saline, and Drug) were grouped for analyses. Noncontingent 01 was then treated as the Baseline, and Noncontingent 02 and 03 active and inactive lever entrances per meter were normalized by the group mean of Noncontingent 01 to account for high individual variation. Following a 0.23 lambda Yeo-Johnson power transformation, a Type III Wald test indicated a significant main effect of session ($\chi^2(1)=12.46$, $p=.002$), but not a significant main effect of levers ($\chi^2(1)=3.12$, $p=.162$) or their interaction ($\chi^2(1)=3.49$, $p=.162$). Post hoc analyses revealed Noncontingent 03 to be significantly greater than Noncontingent 02, prompting a paired Student's t-test between the active and inactive levers in Noncontingent 03 where active was found to be greater than inactive. Direct comparisons to

Noncontingent 01 found that the inactive lever entrances per meter in Noncontingent 02

were significantly less than baseline, while the active lever entrances per meter in

Noncontingent 03 were significantly greater than baseline. (**Fig. 4A**). These results

indicate that by Noncontingent 03 rats demonstrate preferential seeking behavior

toward the drug-associated active lever area.

To visualize the development of interest in the active lever, density heat maps of

each rat's nose node coordinates (*x, y*) for all sessions within each Noncontingent

session were overlain. Bounding boxes corresponding to the active and inactive lever

areas were plotted with the rounded mean lever entrances. Student's t-tests between

active and inactive lever entrances within Noncontingent sessions revealed only

Noncontingent 03 following a 0.23 lambda Yeo-Johnson power transformation achieved

significance, with active greater than inactive (**Fig. 4B**). These results highlight the

development of interest in the active lever after a history or self-administration as the

high activity regions migrate to the active lever side by Noncontingent 03.

3.4.2. Pre-lever activity during Short Access sessions 07-09 predicts active lever

entrances per meter during Noncontingent sessions.

To determine if the interest in the active lever was developed during self-

administration, rats were grouped as "High" (n=19) or "Low" (n=18) by a median-split z-

score of their pre-lever activity measures (locomotion and active lever entrances per

meter) during Short Access 07-09. These sessions were chosen as they were the first to

show significance compared to Short Access 01 and are the last Short Access sessions

before Noncontingent 02. First, the average of each variable was computed across the

three sessions before being standardized within the respective variables. The two z-

scores were then averaged, where higher scores equate to greater pre-session activity.

Following a median z-score split, those greater than or equal to the median were

deemed "High", and those less than were deemed "Low".

Following a 0.2 lambda Yeo-Johnson power transformation, a Type III Wald test

indicated no main effect of session ($\chi^2(1)=4.81$, $p=.059$), but a significant effect of pre-

lever activity ($\chi^2(1)=29.53$, $p<.001$) and their interaction ($\chi^2(1)=9.39$, $p=.006$). Post hoc

analyses revealed the High pre-lever activity group had significantly greater active lever

entrances per meter in both Noncontingent 02 and 03 than baseline and versus the Low

pre-lever activity group within Noncontingent 02. For the Low pre-lever activity group,

Noncontingent 02 was significantly less than both baseline and Noncontingent 03.

There was no difference between groups in Noncontingent 03 (**Fig. 4C**). These results

confirm that rats with high pre-lever activity during Short Access 07-09 led to significant

seeking in Noncontingent 02 that persisted in Noncontingent 03. Furthermore, the Low

pre-lever activity group developed an interest during Long Access before Noncontingent

03.

3.4.3. Pre-lever activity during Short Access sessions 07-09 predicts cocaine self-

administration in Short and Long Access

To determine if pre-lever activity during Short Access sessions 07-09 predicts

cocaine intake over the entire session in Short Access, a Type III Wald test was

performed. A significant main effect was found in session ($\chi^2(9)=125.52$, $p<.001$), but

not a significant effect of pre-lever activity ($\chi^2(1)$=3.51, p=.169) or their interaction

($\chi^2(9)$=16.9, p=.169) (**Fig. 4D**). These results show that cocaine intake did not differ

between groups in Short Access.

To determine if pre-lever activity during Short Access sessions 07-09 predicts

cocaine intake over the entire session in Long Access, a Type III Wald test was

performed. Significant main effects were found in session ($\chi^2(13)$=62.03, p<.001), pre-

lever activity ($\chi^2(1)$=13.49, p<.001) and their interaction ($\chi^2(13)$=28.79, p=.015). Post

hoc analyses revealed the High pre-lever activity group to have significantly greater

cocaine infusions over the entire session in all Long Access sessions except Long

Access 11, 13, and 14 (**Fig. 4D**). These results confirm that pre-lever activity during

Short Access 07-09 was initially predictive of cocaine intake during Long Access, but

that groups eventually converged.

To determine if pre-lever activity during Short Access sessions 07-09 predicts

cocaine intake within the first 15 minutes of the session, a Type III Wald test was

performed. A significant main effect was found in session ($\chi^2(23)$=208.9, p<.001), but

not in pre-lever activity ($\chi^2(1)$=0.85, p=.742) or their interaction ($\chi^2(23)$=35.32, p=.134)

(**Fig 4E**). These results suggest that pre-lever activity during Short Access 07-09 is not

predictive of loading behavior between groups.

Considering the incongruency of the significant association between pre-lever

variables and cocaine intake variables (**Fig. 3**), and the previous nonsignificant results

between pre-lever activity groups and cocaine infusions in the first 15 minutes (**Fig. 4D**),

correlations were performed between cocaine intake variables and pre-lever activity z-

scores in Short Access and three separate phases of Long Access. For cocaine

infusions during the entire session and pre-lever activity, significant positive correlations were found in Short Access (r_s(2960)=.65, p<.001; **Fig. 4F**), Long Access 01-05 (r_s(3284)=.61, p<.001; **Fig. 4G**), Long Access 06-10 (r_s(3422)=.59, p<.001; **Fig. 4H**), but not during Long Access 11-14 (r_s(7013)=.17, p=.318; **Fig. 4I**). For cocaine infusions during the first 15 minutes and pre-lever activity, significant positive correlations were similarly found in Short Access (r_s(3094)=.63, p<.001; **Fig. 4J**), Long Access 01-05 (r_s(3301)=.61, p<.001; **Fig. 4K**), Long Access 06-10 (r_s(4023)=.52, p<.001; **Fig. 4L**), but not during Long Access 11-14 (r_s(6735)=.2, p=.231; **Fig. 4M**). These results confirm that rats who showed markedly increased pre-lever activity during Short Access sessions 07-09 were more likely to consume more cocaine during the first 15 minutes and over the entire session in both Short and Long Access, but converged by the end of Long Access. Furthermore, these results support those shown in **Fig. 4C,** where no significant difference was found in active lever entrances per meter in Noncontingent 03, which occurs after Long Access 14.

Figure 4. Pre-lever activity during Short Access 07-09 predicts cocaine intake in Short and Long Access and seeking behavior during Noncontingent sessions.

(A.) Active (green) and inactive (red) lever entrances per meter as percentage difference from Baseline (Noncontingent 01). #p<.05, ##p<.01 vs Baseline; *p<.05 vs inactive lever. **(B.)** Nose (x, y) pixel coordinate heatmaps of all rats within each Noncontingent session in relation to active (green) and inactive (red) Levers. Each plot represents ~109 sessions overlain revealing high and low activity zones. Numbers within lever bounding boxes are the mean sum of active and inactive lever entrances for all rats within each Noncontingent session (Noncontingent 01 active: M=22.2, SD=13.9; inactive: M=23.4, SD=12.3; Noncontingent 02 active: M=21.8, SD=16.1; inactive: M=18.4, SD=16.5; Noncontingent 03 active: M=43.5, SD=42.1; inactive: M=24.7, SD=22.1). *p<.05 vs inactive lever. **(C.)** Active lever entrances per meter percent difference from Baseline (Noncontingent 01) grouped by pre-lever activity z-score during Short Access 07-09. ###p<.001 vs Baseline; Within pre-lever activity group: $\$p$<.05, $\$\$\$p$<.001 vs Noncontingent 02; Between pre-lever activity groups: ***p<.001 vs Noncontingent 02 Low group. **(D-E.)** Cocaine infusions over the entire session (D.) and within the first 15 minutes (E.) grouped by Short Access 07-09 pre-lever activity z-score. *p<.05, **p<.01, ***p<.001, ###p<.001 vs Low. **(F-I.)** Cocaine infusions over the entire session correlated using Spearman Rank with Short Access 07-09 pre-lever activity z-score in Short Access (F.), Long Access 01-05 (G.), Long Access 06-10 (H.), but not during Long Access 11-14 (I.). **(J-M.)** Cocaine infusions within the first 15 minutes correlated using Spearman Rank with Short Access 07-09 pre-lever activity z-

535   score in Short Access (J.), Long Access 01-05 (K.), Long Access 06-10 (L.), but not

536   during Long Access 11-14 (M.). Plots represent means and SEM.

4. Discussion

Here, we demonstrate that cocaine infusions over the full session and within the

first 15 minutes escalated during Short and Long Access in heterogeneous stock rats.

Locomotion increased after a noncontingent cocaine infusion, and population-level

analysis showed no signs of psychomotor sensitization or tolerance following Short

Access (Noncontingent 02) or Long Access (Noncontingent 03). Individual analyses

revealed strong divergence in psychomotor sensitization or tolerance expression,

though neither influenced cocaine self-administration. Pre-lever locomotion and active

lever entrances per meter escalated from Short to Long Access, increased after

instances of abstinence, and predicted early session intake. Rats with high pre-lever

activity during Short Access sessions 07-09 escalated active lever entrances across

Noncontingent sessions, and earned more infusions during Short Access and first two

weeks of Long Access than low pre-lever activity rats, but these differences

disappeared by Long Access 13 as low-activity rats escalated to similar levels. Together,

the data dissociate psychomotor sensitization and tolerance from cocaine intake and

identify pre-lever activity as a robust predictor of cocaine intake and escalation.

4.1.   Cocaine Intake

Rats increased total cocaine intake as early as Short Access 02 and within the first

15 minutes as early as Short Access 07, while exhibiting significant escalation of intake

after two weeks of Long Access. These findings align with our previous reports that

heterogenous stock rats show increased cocaine intake during Short and Long Access

[28], and the larger literature that Long Access induces escalation of intake [3,4,24,55].

4.2.   Psychomotor Expression

Population-level analyses revealed locomotion increased following a

noncontingent infusion of cocaine, but no differences between Drug sessions. Saline 02

was significantly greater than Baseline 02, suggesting an anticipatory conditioned

stimulus effect as the last time they were handled as such they received a

noncontingent infusion of cocaine (Drug 01). However, the effect is not seen following

Long Access, suggesting that this conditioned responding is short-lasting and unlikely to

drive escalation of cocaine self-administration. Individually, rats did show differences in

the propensity to develop psychomotor sensitization or tolerance to noncontingent

cocaine administration. One interpretation is that analyzing only ambulatory activity may

misclassify some rats displaying stereotypic behavior, such as head bobbing, weaving,

sniffing, or gnawing at high cocaine doses [8,56]. However, this is unlikely for several

reasons. First, the dose used (1.5 mg/kg) is known to produce minimal stereotypic

behavior and robust sensitization [6,30-32]. Second, the nose motion (a proxy for

stereotypic behavior) was not found to increase in Tolerant rats (**Fig. S1**). Third, visual

inspection of the videos show that the most tolerant rat (M2677) did not exhibit

stereotypic behavior while the most sensitized rat (M2668) showed pronounced

psychomotor sensitization. A possible explanation for the difference between sensitized

vs. tolerant rats is that sensitized rats consumed less cocaine, while tolerant rats

consumed more. However, intake analyses do not support this hypothesis: groups

showed no differences in cocaine intake in Short or Long Access, or within the first 15

minutes. These findings challenge the idea that Short Access produces only

sensitization and Long Access only tolerance to the effects of cocaine [1,3-6,18,24,57],

and show that neither reliably predicts cocaine intake.

4.3.   Pre-Lever Activity Measures

Pre-lever variables locomotion and active lever entrances per meter increased

from Short to Long Access and intensified following two-to-three days of abstinence.

Similarly, rats showed a significant increase in the active lever area entrances per meter

in Noncontingent 03 in comparison to Noncontingent 01, suggesting sensitization of

drug-seeking behavior. In contrast to measures of psychomotor sensitization or

tolerance, both locomotion and active lever entrances per meter were highly predictive

of intake. These findings align with the incentive salience literature, showing that

repeated drug use leads to salience attribution to cues and environments that

particularly manifests following a period of abstinence [5-6,15-16,19-20,30-32,46-47,58-

65].

4.4.   Individual Differences in Pre-Lever Activity Measures

To investigate the individual propensity to attribute salience to drug-associated

cues, a median z-score of pre-lever activity was calculated. Short Access sessions 07–

09 were selected as they occurred within the same week (no abstinence periods) and

were the first sessions in which both pre-lever variables and infusions within the first 15

minutes reached significance (**Fig. 1C, 3A, & 3H**). Comparisons between High and Low

pre-lever activity groups revealed that high-activity rats exhibited an early escalation of

cocaine intake during Long Access, but ultimately both pre-lever activity groups reached

a ceiling level of intake after two weeks when groups no longer differed.

One possible explanation is that Low pre-lever activity rats in Short Access were

slower in developing incentive salience, and once expressed, escalated their intake to

levels comparable to High pre-lever activity rats. Indeed, the significant difference in

active lever entrances per meter between pre-lever activity groups in Noncontingent 02

and its absence in Noncontingent 03 supports this interpretation. These findings align

with previous reports that individuals differ in their susceptibility to attribute incentive

salience, which may serve as an early predictor of vulnerability [56,59,62-63].

Furthermore, they suggest that incentive salience is a catalyst of escalation, contrary to

previous findings that incentive salience and escalation are dissociable [26].

Another interpretation is that escalating pre-lever activity during Long Access

reflects the combined influence of allostatic load [1,3,55,57], incentive habit formation

[66-68], and the incubation of craving [69-70]. Allostasis theory predicts that repeated

use progressively recruits stress systems and negative affect, shifting the hedonic set

point and motivating drug seeking to alleviate withdrawal-related distress. Consistent

with this view, both animal and human studies show that acute withdrawal, withdrawal-

conditioned cues, negative affect, and psychosocial stress increase craving, invigorate

drug seeking, and amplify cue-evoked neural activity [1,3,55,57,71-75]. The incentive

habit framework suggests that repeated drug-cue pairing transforms early cue reactivity

into motivationally charged habitual responses, which may lead to automatic approach

toward drug-associated cues even before drug availability [66-68]. Finally, incubation of

craving provides an additional, complementary mechanism: periods of abstinence

without obvious negative emotional states enhance the motivational impact of drug cues

over time, strengthening cue-triggered approach behavior [69-70]. Follow up studies will

be needed to dissect the role of each of these processes.

The high pre-lever activity phenotype we observed shares conceptual parallels

with the sign-tracker/goal-tracker framework [59]. Sign-trackers preferentially interact

with a discrete cue (the lever CS), while goal-trackers orient to the site of reward

delivery. Sign-trackers have been shown to initially self-administer more cocaine [63]

and choose cocaine over food [76]. However, important differences exist: sign-tracking

is defined in a purely Pavlovian context, with lever pressing having no instrumental

consequence, whereas our measure reflects operant-conditioned lever-directed

exploration. Notably, recent work has shown that sign-/goal-tracker status does not

reliably predict overall cocaine self-administration or escalation, although sign-trackers

may show greater resistance to punishment [77]. Thus, our results extend the sign-

/goal-tracker framework into an operant cocaine paradigm, highlighting that cue-bias

and responsivity can shape early intake and drive escalation of intake with prolonged

access even in initially resistant animals.

4.5.   Limitations

The present study has a few limitations. One is the lack of a control group without

periods of abstinence. A control group participating in successive sessions could further

confirm the influence of abstinence on incentive salience incubation. However,

noncontingent sessions 02 and 03 did not follow a period of abstinence, and

expressions of incentive salience and both psychomotor sensitization and tolerance

were still observed. Another limitation is the use of pose-estimation over manual scoring

of behavior could result in inaccuracies. However, pose-estimation with SLEAP is

remarkably accurate [37,78], and the instance rendered sample videos display precise

tracking of the catheter and nose nodes (see supplementary videos).

4.6.   Summary

In summary, this report demonstrates that individual differences in psychomotor

sensitization or tolerance had no influence on cocaine intake. In contrast, incentive

salience during Short Access acts as a catalyst and early predictor of escalation in

cocaine self-administration during the first two weeks of Long Access, but does not

predict established escalation of cocaine intake after three weeks of Long Access.

Moreover, individuals initially resistant to incentive salience can, with sufficient

exposure, become sensitized and escalate cocaine use to the same level as more

susceptible individuals. Expression of incentive salience sensitization as reflected in

pre-lever activity measures and their subsequent influence on cocaine intake supports

the idea that incentive salience and psychomotor sensitization are mediated by distinct

mechanisms [6-7,21,31]. The development of the incentive salience profiles is

accelerated by periods of abstinence (weekend), which allows for incubation of craving,

a progressive increase of craving during early days of abstinence that can remain

elevated following prolonged abstinence [44,69-70]. The measure of approach toward

the active lever area using video recording before the start of the self-administration

session may represent a novel behavioral marker of addiction vulnerability that can

precisely track the motivation for the drug with high accuracy without using any operant

measures, the drug itself, or any other training or conditioning that could interfere with

the drug self-administration. Such a measure may also be useful for medication

development to allow dissociation between drug-dependent and drug-independent

effects of experimental medication on drug seeking and taking.

Data Availability Statement

The datasets generated during and/or analyzed during the current study are

available from the corresponding author on reasonable request.

Acknowledgements

This research was done using services provided by the OSG Consortium [38-41]

which is supported by the National Science Foundation awards #2030508 and

#1836650.

Author Contributions

Jarryd Ramborger: Manuscript drafting and revising, data collection, experiment

execution, experimental design, data analysis and interpretation

Joseph Mosquera: Data collection, experiment execution

Molly Brenna: Data collection, experiment execution

Benjamin Sichel: Data collection, experiment execution

Dyar Othman: Data collection, experiment execution

Sonja Plasil, PhD: Data collection, experiment execution

Elizabeth Sneddon, PhD: Data collection, experiment execution

Selene Zahedi: Data collection, experiment execution

Alex Morgan: Data collection, experiment execution

Supakorn Chonwattanagul: Data collection, experiment execution

Kathleen Bai: Data collection, experiment execution

Lindsey China: Data collection, experiment execution

Tu La: Data collection, experiment execution

Lisa Maturin: Data collection

Lieselot L. G. Carrette, PhD: Experimental design, manuscript drafting and revising

Olivier George, PhD: Manuscript drafting and revising, experimental design, data
analysis and interpretation, funding and resources, final approval

Funding

Funding was supported by DA043799 and the Preclinical Addiction Research
Consortium.

Competing Interests

Authors declare no competing interests.

References

1. Koob, G. F., & Le Moal, M. (1997). Drug abuse: Hedonic homeostatic dysregulation. *Science, 278*(5335), 52–58. https://doi.org/10.1126/science.278.5335.52

2. Koob, G. F., & Volkow, N. D. (2016). Neurobiology of addiction: a neurocircuitry analysis. *The lancet. Psychiatry, 3*(8), 760–773. https://doi.org/10.1016/S2215-0366(16)00104-8.

3. Ahmed, S. H., & Koob, G. F. (1998). Transition from moderate to excessive drug intake: Change in hedonic set point. *Science, 282*(5387), 298-300. https://doi.org/10.1126/science.282.5387.298.

4. Ben-Shahar, O., Ahmed, S. H., Koob, G. F., & Ettenberg, A. (2004). The transition from controlled to compulsive drug use is associated with a loss of sensitization. *Brain Research, 995*(1), 46–54. https://doi.org/10.1016/j.brainres.2003.09.053

5. Calipari, E. S., Ferris, M. J., Zimmer, B. A., Roberts, D. C. S., & Jones, S. R. (2013). Temporal pattern of cocaine intake determines tolerance vs sensitization of cocaine effects at the dopamine transporter. *Neuropsychopharmacology, 38*(12), 2385–2392. https://doi.org/10.1038/npp.2013.136.

6. Carr, C. C., Ferrario, C. R., & Robinson, T. E. (2020). Intermittent access cocaine self-administration produces psychomotor sensitization: Effects of withdrawal, sex, and cross-sensitization. *Psychopharmacology, 237*(6), 1795–1812. https://doi.org/10.1007/s00213-020-05500-4.

7. Dalia AD, Norman MK, Tabet MR, Schlueter KT, Tsibulsky VL, Norman AB (1998) Transient amelioration of the sensitization of cocaine-induced behaviors in rats

by the induction of tolerance. Brain Res 797:29–34.

https://doi.org/10.1016/S0006-8993(98)00323-0.

8.   Flagel, S. B., & Robinson, T. E. (2007). Quantifying the psychomotor activating

effects of cocaine in the rat. *Behavioural Pharmacology, 18*(4), 297–302.

https://doi.org/10.1097/FBP.0b013e3281f522a4.

9.   Lett, B. T. (1989). Repeated exposures intensify rather than diminish the

rewarding effects of amphetamine, morphine, and cocaine.

Psychopharmacology, 98(3), 357-362. https://doi.org/10.1007/BF00451688.

10.   Porrino, L. J. (1993). Functional consequences of acute cocaine treatment

depend on route of administration. *Psychopharmacology, 112*(3), 343–351.

https://doi.org/10.1007/BF02244948.

11.   Swain, Y., Muelken, P., LeSage, M. G., Gewirtz, J. C., & Harris, A. C. (2018).

Locomotor activity does not predict individual differences in morphine self-

administration in rats. *Pharmacology Biochemistry and Behavior*, 166, 48-56.

https://doi.org/10.1016/j.pbb.2018.01.008.

12.   Vanderschuren, L. J. M. J., Schmidt, E. D., De Vries, T. J., Van Moorsel, C. A. P.,

Tilders, F. J. H., & Schoffelmeer, A. N. M. (1999). A single exposure to

amphetamine is sufficient to induce long-term behavioral, neuroendocrine, and

neurochemical sensitization in rats. *Journal of Neuroscience, 19*(21), 9579–9586.

https://doi.org/10.1523/JNEUROSCI.19-21-09579.1999

13.   Vanderschuren, L. J. M. J., De Vries, T. J., Wardeh, G., Hogenboom, F. A. C. M.,

& Schoffelmeer, A. N. M. (2001). A single exposure to morphine induces long-

lasting behavioural and neurochemical sensitization in rats. *European Journal of*

Neuroscience, 14(8), 1533–1538. https://doi.org/10.1046/j.0953-

816x.2001.01771.x

14.   Hammer, R. P. Jr., Egilmez, Y., & Emmett-Oglesby, M. W. (1997). Neural

mechanisms of tolerance to the effects of cocaine. *Behavioural Brain Research*,

84(1-2), 225-239. https://doi.org/10.1016/s0166-4328(97)83332-3.

15.   Kalivas PW, Sorg BA, Hooks MS (1993). The pharmacology and neural circuitry

of sensitization to psychostimulants. Behav Pharmacol. 1993;4(4):315-334.

PMID: 11224200.

16.   Vanderschuren, L. J. M. J., & Kalivas, P. W. (2000). Alterations in dopaminergic

and glutamatergic transmission in the induction and expression of behavioral

sensitization: A critical review of preclinical studies. *Psychopharmacology, 151*(2),

99–120. https://doi.org/10.1007/s002130000493.

17.   Vezina, P. (2004). Sensitization of midbrain dopamine neuron reactivity and the

self-administration of psychomotor stimulant drugs. *Neuroscience &*

Biobehavioral Reviews, 27(8), 827–839.

https://doi.org/10.1016/j.neubiorev.2003.11.001.

18.   Solomon, R. L. (1980). The opponent-process theory of acquired motivation: The

costs of pleasure and the benefits of pain. *American Psychologist*, 35(8), 691-

712. https://doi.org/10.1037/0003-066X.35.8.691

19.   Robinson, T. E. & Berridge, K. C.(1993). The neural basis of drug craving: An

incentive-sensitization theory of addiction. *Brain Research Reviews*, 18(3), 247-

291. https://doi.org/10.1016/0165-0173(93)90013-P.

20.   Ciccocioppo, R., Martin-Fardon, R., & Weiss, F. (2004). Stimuli associated with a

single cocaine experience elicit long-lasting cocaine-seeking. *Nature*

Neuroscience, 7(5), 495-496. https://doi.org/10.1038/nn1239.

21.   Izenwasser, S., & French, D. (2002). Tolerance and sensitization to the

locomotor-activating effects of cocaine are mediated via independent

mechanisms. *Pharmacology Biochemistry and Behavior, 73*(4), 877–882.

https://doi.org/10.1016/s0091-3057(02)00942-5.

22.   Kalivas, P. W., & Alesdatter, J. E. (1993). Involvement of N-methyl-D-aspartate

receptor stimulation in the ventral tegmental area and amygdala in behavioral

sensitization to cocaine. *Journal of Pharmacology and Experimental*

Therapeutics, 267(1), 486-495. Retrieved from

http://jpet.aspetjournals.org/content/267/1/486.abstract.

23.   Ungless, M., Whistler, J., Malenka, R. *et al.* Single cocaine exposure *in vivo*

induces long-term potentiation in dopamine neurons. *Nature* 411, 583–587

(2001). https://doi.org/10.1038/35079077

24.   Ahmed, S. H., & Cador, M. (2006). Dissociation of psychomotor sensitization

from compulsive cocaine consumption. Neuropsychopharmacology, 31(3), 563–

571. https://doi.org/10.1038/sj.npp.1300834.

25.   Kawa AB, Bentzley BS, Robinson TE. Less is more: prolonged intermittent

access cocaine self-administration produces incentive-sensitization and

addiction-like behavior. Psychopharmacology (Berl). 2016 Oct;233(19-20):3587-

602. doi: 10.1007/s00213-016-4393-8. Epub 2016 Aug 2. PMID: 27481050;

PMCID: PMC5023484.

26.   Kawa AB, Valenta AC, Kennedy RT, Robinson TE. Incentive and dopamine

sensitization produced by intermittent but not long access cocaine self-

administration. Eur J Neurosci. 2019 Aug;50(4):2663-2682. doi:

10.1111/ejn.14418. Epub 2019 May 2. PMID: 30968487; PMCID: PMC6742545.

27.   Carrette, L. L. G., de Guglielmo, G., Kallupi, M., Maturin, L., Brennan, M.,

Boomhower, B., et al.. (2021). The Cocaine and Oxycodone Biobanks, two

repositories from genetically diverse and behaviorally characterized rats for the

study of addiction. eNeuro, 8(3), ENEURO.0033-21.2021.

https://doi.org/10.1523/ENEURO.0033-21.2021.

28.   Giordano de Guglielmo, Carrette, L. L. G., Kallupi, M., Brennan, M., Boomhower,

B., Maturin, L., et al.. (2024). Large-scale characterization of cocaine addiction-

like behaviors reveals that escalation of intake, aversion-resistant responding,

and breaking-points are highly correlated measures of the same construct. eLife,

12, RP90422. https://doi.org/10.7554/eLife.90422

29.   Solberg Woods LC, Palmer AA (2019) Using heterogeneous stocks for fine-

mapping genetically complex traits. Methods Mol Biol 2018:233–247.

doi: 10.1007/978-1-4939-9581-3_11.

30.   Samaha, A. N., Li, Y., & Robinson, T. E. (2002). The rate of intravenous cocaine

administration determines susceptibility to sensitization. The Journal of

neuroscience : the official journal of the Society for Neuroscience, 22(8), 3244–

3250. https://doi.org/10.1523/JNEUROSCI.22-08-03244.2002.

31.   Samaha, A.-N., Mallet, N., Ferguson, S. M., Gonon, F., & Robinson, T. E. (2004).

The rate of cocaine administration alters gene regulation and behavioral

plasticity: Implications for addiction. *The Journal of Neuroscience*, *24*(28), 6362–

6370. https://doi.org/10.1523/JNEUROSCI.1424-04.2004.

32.   Samaha, A.-N., & Robinson, T. E. (2005). Why does the rapid delivery of drugs to

the brain promote addiction? *Trends in Pharmacological Sciences, 26*(2), 82-87.

https://doi.org/10.1016/j.tips.2004.12.007.

33.   Woolverton, W. L., & Wang, Z. (2004). Relationship between injection duration,

transporter occupancy, and reinforcing strength of cocaine. *European Journal of*

Pharmacology, *486*(3), 251–257. https://doi.org/10.1016/j.ejphar.2004.01.003.

34.   Ramborger, J., Kalra, S., Mosquera, J., Smith, A. C. W., & George, O. (2024).

High quality, high throughput, and low-cost simultaneous video recording of 60

animals in operant chambers using PiRATeMC. *Journal of Neuroscience*

Methods, 411, 110270. https://doi.org/10.1016/j.jneumeth.2024.110270.

35.   R Core Team. (2023). *R: A language and environment for statistical computing*

[Computer software]. R Foundation for Statistical Computing. https://www.R-

project.org/

36.   Posit Software, PBC. (2023). *RStudio: Integrated development environment for R*

[Computer software]. https://posit.co/

37.   T.D. Pereira, N. Tabris, A. Matsliah, D. M. Turner, J. Li, S. Ravindranath, et al..

Sleap: A deep learning system for multi-animal pose tracking. *Nature Methods*,

19(4), 2022. https://doi.org/10.1038/s41592-022-01426-1.

38.   OSG. (2006). OSPool. OSG. https://doi.org/10.21231/906P-4D78

39.   Pordes, R., Petravick, D., Kramer, B., Olson, D., Livny, M., Roy, A., et al.. (2007).

846   The open science grid. J. Phys. Conf. Ser., 78, 012057.

847   https://doi.org/10.1088/1742-6596/78/1/012057.

848   40.   Sfiligoi, I., Bradley, D. C., Holzman, B., Mhashilkar, P., Padhi, S., & Wurthwein, F.

849   (2009). The pilot way to grid resources using glideinWMS. 2009 WRI World

850   Congress on Computer Science and Information Engineering, 2, 428–432.

851   https://doi.org/10.1109/CSIE.2009.950.

852   41.   OSG. (2015). Open Science Data Federation. OSG.

853   https://doi.org/10.21231/0KVZ-VE57

854   42.   Rueden, C.T., Schindelin, J., Hiner, M.C. *et al.* ImageJ2: ImageJ for the next

855   generation of scientific image data. *BMC Bioinformatics* **18**, 529 (2017).

856   https://doi.org/10.1186/s12859-017-1934-z.

857   43.   Berridge, K. C., & Robinson, T. E. (1998). What is the role of dopamine in reward:

858   Hedonic impact, reward learning, or incentive salience? *Brain Research*

859   *Reviews, 28*(3), 309–369. https://doi.org/10.1016/S0165-0173(98)00019-8.

860   44.   Li, X., Caprioli, D., & Marchant, N. J. (2014). Recent updates on incubation of

861   drug craving: A mini-review. *Addiction Biology, 20*(5), 872–876.

862   https://doi.org/10.1111/adb.12205.

863   45.   Bates D, Mächler M, Bolker B, Walker S (2015). "Fitting Linear Mixed-Effects

864   Models Using lme4." *Journal of Statistical Software*, **67**(1), 1–

865   48. doi:10.18637/jss.v067.i01.

866   46.   Kuznetsova A, Brockhoff PB, Christensen RHB (2017). "lmerTest Package: Tests

867   in Linear Mixed Effects Models." *Journal of Statistical Software*, **82**(13), 1–

868   26. doi:10.18637/jss.v082.i13.

869   47.   Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A,

870   et al. (2017). "glmmTMB Balances Speed and Flexibility Among Packages for

871   Zero-inflated Generalized Linear Mixed Modeling." *The R Journal*, 9(2), 378–

872   400. doi:10.32614/RJ-2017-066.

873   48.   Luke, S.G. Evaluating significance in linear mixed-effects models in R. *Behav*

874   *Res* **49**, 1494–1502 (2017). https://doi.org/10.3758/s13428-016-0809-y.

875   49.   Hartig, F. (2024). *DHARMa: Residual diagnostics for hierarchical (multi-level /*

876   *mixed) regression models* (Version 0.4.7) [R package]. CRAN.

877   https://doi.org/10.32614/CRAN.package.DHARMa.

878   50.   Meteyard, L., & Davies, R. A. I. (2020). Best practice guidance for linear mixed-

879   effects models in psychological science. *Journal of Memory and Language, 112*,

880   104092. https://doi.org/10.1016/j.jml.2020.104092.

881   51.   Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (3rd ed.).

882   Sage. https://www.john-fox.ca/Companion/.

883   52.   Lenth, R. V. (2025). *emmeans: Estimated marginal means, aka least-squares*

884   *means* (Version 1.10.7) [R package]. CRAN.

885   https://doi.org/10.32614/CRAN.package.emmeans.

886   53.   Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in

887   multiple testing under dependency. *The Annals of Statistics, 29*(4), 1165–1188.

888   https://doi.org/10.1214/aos/1013699998.

54.   Kassambara, A. (2023). rstatix: Pipe-friendly framework for basic statistical tests

(Version 0.7.2) [Computer software].

https://rpkgs.datanovia.com/rstatix/[](https://rpkgs.datanovia.com/rstatix/authors.h

tml).

55.   Ahmed, S. H., & Koob, G. F. (1999). Longlasting increase in the set point for

cocaine selfadministration after escalation in rats. *Psychopharmacology, 146*(3),

303–312. https://doi.org/10.1007/s002130051121.

56.   Flagel, S. B., Watson, S. J., Akil, H., & Robinson, T. E. (2008). Individual

differences in the attribution of incentive salience to a reward-related cue:

Influence on cocaine sensitization. *Behavioural Brain Research, 186*(1), 48–56.

https://doi.org/10.1016/j.bbr.2007.07.022

57.   Ahmed, S. H., Kenny, P. J., Koob, G. F., & Markou, A. (2002). Neurobiological

evidence for hedonic allostasis associated with escalating cocaine use. *Nature

Neuroscience, 5*(7), 625–626. https://doi.org/10.1038/nn872.

58.   Blanchard, M. M., Mendelsohn, D., & Stamp, J. A. (2009). The HR/LR model:

Further evidence as an animal model of sensation seeking. *Neuroscience &

Biobehavioral Reviews, 33*(7), 1145-1154.

https://doi.org/10.1016/j.neubiorev.2009.05.009.

59.   Flagel, S.B., Watson, S.J., Robinson, T.E. *et al.* Individual differences in the

propensity to approach signals vs goals promote different adaptations in the

dopamine system of rats. *Psychopharmacology* **191**, 599–607 (2007).

https://doi.org/10.1007/s00213-006-0535-8.

60.   Piazza, P. V., Deroche-Gamonent, V., Rouge-Pont, F., & Le Moal, M. (2000).

Vertical Shifts in Self-Administration Dose–Response Functions Predict a Drug-

Vulnerable Phenotype Predisposed to Addiction. *The Journal of Neuroscience*,

20(11), 4226-4232. https://doi.org/10.1523/JNEUROSCI.20-11-04226.2000.

61.   Post, R. M., Lockfeld, A., Squillace, K. M., & Contel, N. R. (1981). Drug-

environment interaction: Context dependency of cocaine-induced behavioral

sensitization. *Life Sciences*, *28*, 755–760. https://doi.org/10.1016/0024-

3205(81)90157-0.

62.   Saunders, B. T., & Robinson, T. E. (2010). A cocaine cue acts as an incentive

stimulus in some but not others: Implications for addiction. Biological Psychiatry,

67(8), 730–736. https://doi.org/10.1016/j.biopsych.2009.11.015

63.   Saunders, B. T., & Robinson, T. E. (2011). Individual variation in the motivational

properties of cocaine. *Neuropsychopharmacology, 36*(8), 1668–1676.

https://doi.org/10.1038/npp.2011.48.

64.   Stewart, J., & Badiani, A. (1993). Tolerance and sensitization to the behavioral

effects of drugs. *Behavioural Pharmacology, 4*(4), 289-312.

65.   Weiss, S. R. B., Post, R. M., Pert, A., Woodward, R., & Murman, D. (1989).

Context-dependent cocaine sensitization: Differential effect of haloperidol on

development versus expression. *Pharmacology, Biochemistry, and Behavior,*

34(3), 655-661. https://doi.org/10.1016/S0091-3057(89)80001-2.

66.   Balleine, B. W. (2011). Sensation, incentive learning, and the motivational control

of appetitive behavior. In G. F. Koob, M. Le Moal, & R. F. Thompson (Eds.),

Neurobiology of addiction (pp. 241–259). Academic Press.

67.   Belin, D., Belin-Rauscent, A., Murray, J. E., & Everitt, B. J. (2013). Addiction:

935   Failure of control over maladaptive incentive habits. Current Opinion in

936   Neurobiology, 23(4), 564–572. https://doi.org/10.1016/j.conb.2013.01.025

937   68.   Fouyssac, M., Hynes, T., Belin-Rauscent, A., Joshi, D. D., & Belin, D. (2025).

938   Incentive cocaine-seeking habits and their compulsive manifestation emerge

939   after a downregulation of the dopamine transporter in astrocytes across

940   functional domains of the striatum. European Journal of Neuroscience, 61(5),

941   e70054. https://doi.org/10.1111/ejn.70054

942   69.   Chow, J. J., Pitts, K. M., Negishi, K., Madangopal, R., Dong, Y., Wolf, M. E., &

943   Shaham, Y. (2025). Neurobiology of the incubation of drug craving: An update.

944   Pharmacology & Therapeutics Reviews, 77(2), 100022.

945   https://doi.org/10.1016/j.pharmr.2024.100022

946   70.   Grimm, J. W., Hope, B. T., Wise, R. A., & Shaham, Y. (2001). Neuroadaptation:

947   Incubation of cocaine craving after withdrawal. Nature, 412(6843), 141–142.

948   https://doi.org/10.1038/35084134

949   71.   Carmack, S. A., Kirson, E. D., & Markou, A. (2019). Heroin addiction engages

950   negative emotional learning brain circuits in rats. *The Journal of Clinical*

951   *Investigation, 129*(6), 2480–2484. https://doi.org/10.1172/JCI125534

952   72.   Hutcheson, D. M., et al. (2001). The role of withdrawal in heroin addiction. *Nature*

953   *Neuroscience, 4*(10), 1031–1034. https://doi.org/10.1038/nn0901-943

954   73.   Kenny, P. J., Chen, S. A., Kitamura, O., Markou, A., & Koob, G. F. (2006).

955   Conditioned withdrawal drives heroin consumption and decreases reward

sensitivity. *The Journal of Neuroscience, 26*(22), 5894–5900.

https://doi.org/10.1523/JNEUROSCI.0740-06.2006

74. Pantazis, C. B., Gonzalez, L. A., Tunstall, B. J., Carmack, S. A., Koob, G. F.,

Vendruscolo, L. F., … & George, F. (2021). Cues conditioned to withdrawal and

negative reinforcement: Neglected but key motivational elements driving opioid

addiction. *Science Advances, 7*(46), eabf0364.

https://doi.org/10.1126/sciadv.abf0364

75. Sinha, R., Fuse, T., Aubin, L. R., & O'Malley, S. S. (2000). Psychological stress,

drug-related cues and cocaine craving. *Psychopharmacology (Berlin), 152*(2),

140–148. https://doi.org/10.1007/s002130000499

76. Tunstall, B. J., & Kearns, D. N. (2015). Sign-tracking predicts increased choice of

cocaine over food in rats. Behavioural Brain Research, 281, 222–228.

https://doi.org/10.1016/j.bbr.2014.12.034.

77. Pohořalá, V., Enkel, T., Bartsch, D. *et al.* Sign- and goal-tracking score does not

correlate with addiction-like behavior following prolonged cocaine self-

administration. *Psychopharmacology* **238**, 2335–2346 (2021).

https://doi.org/10.1007/s00213-021-05858-z

78. Pereira, T. D., Shaevitz, J. W., & Murthy, M. (2020). Quantifying behavior to

understand the brain. *Nature Neuroscience, 23*(12), 1537–1549.

https://doi.org/10.1038/s41593-020-00734-z.

976