Incentive Salience, not Psychomotor Sensitization or Tolerance, Drives Escalation of Cocaine Self-Administration in Heterogeneous Stock Rats

Jarryd Ramborger[1,2], Joseph Mosquera[1], Molly Brennan[1], Benjamin Sichel[1], Dyar Othman[1], Sonja Plasil, PhD[1], Elizabeth Sneddon, PhD[1], Selene Zahedi[1,3]**,** Alex Morgan[1], Supakorn Chonwattanagul[1], Kathleen Bai[1], Lindsey China[1], Tu La[1], Lisa Maturin[1], Lieselot L. G. Carrette, PhD[1], Olivier George, PhD[1]

[1] Department of Psychiatry, University of California, San Diego, La Jolla, CA 92093, USA

[2] Department of Psychology, Arizona State University, Tempe, AZ 85287, USA

[3]Institut de Neurosciences de la Timone, Aix-Marseille Université, Marseille, France

Corresponding Author:

Name: Olivier George, PhD

Email: olgeorge@health.ucsd.edu

Present Address:

Department of Psychiatry, University of California, San Diego, La Jolla, CA 92093, USA

Abstract:

Sensitization and tolerance are two phenomena often studied independently despite overlapping neurobiological substrates. Each has extensive research showing their influence on the development and maintenance of addiction, but the degree to which they drive escalation in cocaine self-administration is poorly understood. Using self-administration, intravenous noncontingent infusions, and pose-estimation machine vision, we find that incentive salience, not psychomotor sensitization or tolerance, drives the escalation of cocaine self-administration in heterogenous stock rats. Individual differences in psychomotor sensitization or tolerance were found to have no effect on cocaine intake. Incentive salience as measured by locomotion and active lever entrances per meter traveled occurring before the self-administration session began (pre-lever activity measures) during Short Access (2-hours) was found to predict intake during Long Access (6-hours). Both pre-lever locomotion and active lever entrances per meter were found to increase during Long Access and after two-to-three days of abstinence. Critically, rats with low pre-lever activity during Short Access escalated both their intake and pre-lever measures by the end of Long Access to levels comparable with high pre-lever activity rats who maintained their elevated responding. These findings support the notion that incentive salience during Short Access is a catalyst to escalated use and an early marker of addiction vulnerability. Moreover, they suggest that individuals initially resistant to incentive salience can, with sufficient exposure, become sensitized and escalate cocaine use to the same level as more susceptible individuals. Analysis of pre-lever activity offers a novel longitudinal behavioral marker to

42   predict vulnerability and provides a framework for understanding individual trajectories

43   of addiction.

1. Introduction

Addiction is a chronic, relapsing disorder characterized by cycles of binge/intoxication, withdrawal/negative affect, and preoccupation/anticipation, leading to long-lasting neuroadaptations in stress, executive function, and reward processing [1,2]. Within this framework, two of the most studied adaptations to repeated drug exposure are tolerance, a reduced response to the drug, and sensitization, an enhanced response. Both processes have been demonstrated in preclinical models of addiction, but the expression of either phenomenon varies widely depending on experimental parameters such as drug of choice, method of exposure (e.g., noncontingent injections versus self-administration), schedule of access (e.g. Short Access, Long Access, Intermittent, etc.), and outcome variables (e.g. locomotion, physiological measures, and operant behavior) [3-13]. Despite evidence that tolerance and sensitization share overlapping neurobiological substrates, they are often investigated in isolation [14-17], leading to two competing models in the transition to addiction.

According to the hedonic homeostatic dysregulation theory, 1) repeated drug use leads to tolerance to the positive reinforcing effects of drugs, and 2) the allostatic recruitment of stress circuitry, thereby shifting the hedonic set point and motivating continued use to restore homeostasis [1,18]. In contrast, the incentive-sensitization theory proposes that 1) repeated use hypersensitizes neural circuits leading to *incentive salience*—drug-related cues become increasingly "wanting" even as the "liking" diminishes—and 2) *behavioral sensitization*—a heightened psychomotor response observed primarily in psychostimulants like amphetamines and cocaine [19]. Many agree that both phenomena contribute to the addiction cycle; however, the degree to

which sensitization and tolerance drive escalation of cocaine self-administration is

poorly understood. While a number of studies have demonstrated the expression of

tolerance or sensitization to cocaine with noncontingent methods of continuous,

repeated, or single infusions [7,9,20-23], or the use of different lengths of self-

administration access [3-6,24-26], none have in a longitudinal paradigm employing

noncontingent challenges, Short Access, and Long Access.

To address this gap, using an operant cocaine self-administration paradigm in

heterogeneous stock rats in conjunction with machine vision pose-estimation, we

assessed multiple behavioral indices to differentiate individual development of tolerance

or sensitization across time and conditions and the subsequent influence on cocaine

self-administration. These included: (1) locomotion following a single noncontingent

intravenous infusion of cocaine at three time points: before Short Access, after Short

Access 09, and after Long Access 14 (see **Fig. 1A** for timeline); (2) locomotion for 15

minutes before the self-administration session begins (pre-lever); (3) drug-seeking

during pre-lever and Noncontingent sessions, measured as nose entrances into the

active lever area per meter traveled; (4) cocaine intake within the first 15 minutes and

(5) across the full session; and (6) breakpoint on a progressive ratio schedule.

Tolerance and sensitization will be assessed by the measures of psychomotor

locomotion during Noncontingent sessions. Incentive salience will be assessed by both

locomotion and nose entrances into the active lever area per meter during the pre-lever

phase of self-administration.

2. Materials and Methods

A more detailed account of procedures in sections 2.1-2.3 below has been

published elsewhere [27-28]. ChatGPT4 was utilized for coding assistance in RStudio

and python, authors take full responsibility for output.

2.1.   Subjects

Heterogeneous stock rats (N=37; n=21 males; n=16 females) were used to mimic

the genetic diversity seen in the human population. Rats were obtained from the from

the University of California, San Diego colony [29] and housed in pairs on a 12-hour

light/dark cycle. Rats were anesthetized with 1–5% isoflurane for jugular vein

catheterization using Micro-Renathane tubing (Braintree Scientific) connected to a guide

cannula that exits through the back and secured with mesh and dental acrylic. Post-

surgery, rats received flunixin subcutaneously (2.5 mg/kg) and cefazolin intramuscularly

(330 mg/kg). Catheters were flushed daily with heparinized saline (10 U/ml, American

Pharmaceutical Partners) mixed with 0.9% bacteriostatic sodium chloride (Hospira) and

cefazolin (52.4 mg/0.2 ml). Catheter patency was tested intravenously with brevital

sodium (1.5 mg, American Pharmaceutical Partners) at experiment start and end, with

failing rats removed from the experiment. Rats analyzed demonstrated catheter

patency. All procedures were conducted in strict adherence to the National Institutes of

Health Guide for the Care and Use of Laboratory Animals and were approved by the

Institutional Animal Care and Use Committees of UC San Diego.

2.2.   Drugs

Cocaine HCl (National Institute on Drug Abuse, Bethesda, MD) was dissolved in 0.9% sterile saline. During self-administration sessions, a press of the active lever resulted in an intravenous infusion of 0.1ml at 0.5 mg/kg over six seconds by pump, followed by a 20-second timeout where the active lever has no programmed effect. During Noncontingent sessions, a single intravenous infusion of 0.3ml at 1.5 mg/kg (i.e. three times the dose received in self-administration) was delivered in under five seconds in the same operant chamber as self-administration. Intravenous injections were chosen to address the differences in absorption, clearance, and blood concentrations between intravenous and intraperitoneal methods [30-32]. Furthermore, administration within the same chamber and under five seconds were implemented to address influences of environment and intravenous infusion rate on the development of sensitization [33-36].

2.3.   Intravenous Self-Administration

Over 8 weeks (Monday-Friday), rats participated in 30 self-administration sessions with various lengths and ratios. Under fixed ratio one, rats participated in 10 two-hour Short Access sessions,15 six-hour Long Access sessions, and one Preshock and one Shock session, each for one-hour. During Shock, each infusion had a 30% chance to be paired with a footshock (0.3 mA for 0.5 sec) through the grid floor. Under progressive ratio, rats participated in three six-hour sessions in which the response requirements for an infusion followed the equation [5e(injection number × 0.2)] – 5. The breakpoint was defined as the last ratio before a 60-minute period in which a ratio was

not completed, in which case the session ended. A complete experimental timeline is

shown in **Fig. 1A**.

2.4. Noncontingent Sessions

Seven days after surgeries, rats underwent the first noncontingent session

(Noncontingent 01), which occurred before Short Access 01. Rats were placed in the

operant chamber and video recorded for 30 minutes to establish baseline

measurements (Baseline 01). Rats were then given an intravenous saline injection,

placed back in the chamber, and recorded for 30 minutes (Saline 01), followed by a

noncontingent intravenous injection of cocaine (1.5 mg/kg) and recorded for 30 minutes

(Drug 01). The same procedures were replicated for the subsequent noncontingent

sessions following Short Access 09 (Noncontingent 02) and Long Access 14

(Noncontingent 03).

2.5. Video Recording and Processing

2.5.1. *Raspberry Pi Video Recording*

Video recordings of sessions were conducted as described previously [37] with a

few alterations. A static FFmpeg build (ffmpeg-6.0-armhf-static;

https://johnvansickle.com/ffmpeg/) was installed onto each Raspberry Pi for video

processing to reduce video size threefold without sacrifice to video quality. All sessions

were recorded for 30 minutes. During self-administration sessions, levers were

extended in the chamber after 15 minutes of recording. Each comma-separated value

file (csv) file was processed in RStudio (R version 4.3.1 (2023-06-16 ucrt)) to ensure

consistent length and alignment to the lever extension, resulting in ~25 minutes per csv,

or ~12.5 minutes for both pre- and post-lever data. Out of a possible 1,110 csv files (37

rats over 30 sessions), 1,084 are present in the study due to exclusions for various

issues such as catheter or infrared light disconnections. For eight rats during Long

Access 04 the Med Associates program was initiated too early, resulting in no pre-lever

data. For Noncontingent sessions, R processing resulted in ~25 minutes per csv. Out of

a possible 333 csv files, 327 are present in the data, as six failed to track during Social

LEAP Estimates Animal Poses (SLEAP) processing.

2.5.2. Social LEAP Estimates Animal Poses (SLEAP) Modeling

A pose-estimation model was trained using SLEAP on ~5200 labeled frames

over 89 videos from past cohorts utilizing the top-down configuration and a unet

backbone [38]. Thirteen nodes were tracked: nose, both ears, neck, four paws, base

(where tail met the body), active and inactive levers, active lever cue light, and the

catheter node, which was chosen as the model anchor point. A second model was

trained to track the first Noncontingent session, where the rats were mistakenly not

plugged into their catheter lines. This model focused on only six nodes to accelerate

training: nose, both ears, neck, base, and the catheter as the anchor node, utilizing

~2200 labeled frames from 40 videos.

2.5.3. *High Throughput Computing on Open Science Grid (OSG)*

Video recordings were processed on the OSG Consortium supercomputer

system [39-42]. A python script with SLEAP command line interface functions processed

each video to identify one instance (track) using an "instance" identification method and

Hungarian approach between frames. Results of each node for each frame per second

inference registered as (x, y) coordinates were followed by an interpolation method to fill

missing data points (https://sleap.ai/notebooks/Analysis_examples.html).

2.6.   Dependent variable calculation

Calculations were conducted in R version 4.3.1 using RStudio [43-44]. Euclidian

distances were used to calculate movement of the catheter (locomotion) and the nose

(nose motion) nodes frame by frame and converted from pixels to centimeters using

Image2 [45] and the known measurements of the ENV-005 grid. Given the barrel

distortion of a fisheye lens, the conversion factor was computed by measuring the grid

floor at five separate segments and averaging the two length measurements (left and

right), averaging three width measurements (top, middle, and bottom), and taking the

average of the two, resulting in ~0.111cm/px. Locomotion variables are the sum of the

distances traveled and then smoothed by a window of 10 (10 FPS, i.e., 0.33 seconds),

then converted to meters. Locomotion (m) pre-lever corresponds to the locomotion

measured for the 15 minutes before the levers are extended into the chamber. Active

and inactive lever entrances were calculated by producing a bounding box from the

position of the (x,y) nodes, 30 +/- x and 18 +/- y, and counting each entrance of the

nose node into the lever areas. Active and inactive lever entrances per meter are the

total lever entrances normalized by locomotion. Cocaine infusions are the total infusions

delivered during the session, and cocaine infusions first 15 minutes are those that

occurred during the video recordings. Session number corresponds to an individual

session (e.g. Short Access 01). Session type refers to experimental phase (e.g. Long

Access). Abstinence refers to the time between the end of one session and the

beginning of the next, where 18 Hours translates to consecutive days, 66 Hours to a

two-day weekend, and 90 Hours to a three-day weekend. Pre-lever variables were

examined by abstinence to test for effects that could facilitate sensitization [6,14-

17,20,33-35,46-47]. Analyses with sex as a factor were performed, but most effects

were not sex dependent and therefore are reported in the Supplementary Material.

2.7.   Statistical Analyses

Statistical comparisons were made by fitting a linear mixed effects model utilizing

the lmerTest package [48-49] or a generalized linear mixed effects model utilizing the

glmmTMB package [50] in R. Given that linear and generalized linear mixed-effects

models are inherently robust to moderate deviations from normality and extreme values,

outliers for all variables were winsorized at the 95th percentile to minimize their

influence while retaining potentially meaningful behavioral variability. Inferential tests of

fixed effects for linear mixed models were conducted using Type III ANOVA with

Satterthwaite degrees of freedom approximations (Type III ANOVA test) which have

been shown to control for Type I errors in hierarchical models [51], while for generalized

linear models a Type III Wald chi-square analysis of deviance was conducted (Type III

Wald test). Tests for model assumptions and fit were analyzed with the

simulateResdiduals() function from the DHARMa package [52]. The model fitting

process and chosen models of each analysis can be found in the Supplementary

Material with best adherence to suggestions previously published [53] but prioritizing the

results of DHARMa tests. Transformations of response variables where necessary (see

Supplemental Material) were achieved using Box-Cox or Yeo-Johnson after identifying

the most effective lambda using powerTransform() from the car package and reported at

two decimal places [54]. Post hoc analyses of all significance results were conducted

using contrasts(), while direct comparisons to baseline used test(), both within the

emmeans package and calculate the pairwise comparisons of the estimated marginal

means of mixed effects models and are therefore model-derived ratios or mean

differences [55]. All p-values were adjusted by the Benjamini-Yekutieli (BY) method due

to its consideration of dependence and correlations in nested data such as mixed

models [56]. From the rstatix package, Student's t-tests were conducted with t_test()

with Bonferroni adjustments and correlations with cor_test() by either Pearson or

Spearman Rank as determined by a normality test with shapiro_test() [57]. Plots

represent the raw average values or percent difference from baseline.

3. Results

3.1. Cocaine Infusions

3.1.1. Total cocaine infusions escalate during both Short and Long Access

To determine whether cocaine intake changed during Short Access, a Type III

Wald test was performed. A significant main effect of session number was observed

($\chi^2(9)=100.94$, $p<.001$). Post hoc analyses revealed all sessions to have significantly

more cocaine infusions in comparison to Short Access 01. These results confirm that

cocaine intake increased as early as Short Access 02 (**Fig. 1B**).

To determine whether cocaine intake changed during Long Access, a Type III

Wald test was performed. A significant main effect of session number was identified

(χ^2(13)=173.01, *p*<.001). Post hoc comparisons indicated that Long Access 09 through

Long Access 14 had significantly more infusions relative to Long Access 01. These

results show that cocaine intake continued to escalate into the third week of Long

Access (**Fig. 1B**).

3.1.2. Cocaine infusions during the first 15 minutes escalate across sessions

To evaluate whether cocaine infusions during the initial 15 minutes of each

session escalated, a Type III Wald test was conducted. A significant main effect of

session number was identified (χ^2(23)=304.87, *p*<.001). Post hoc analyses revealed

that cocaine infusions were significantly greater from Short Access 07 through Long

Access 14 compared to Short Access 01. Additionally, when compared to Long Access

01, only sessions Long Access 09, Long Access 13, and Long Access 14 showed

significantly elevated infusions (**Fig. 1C**). These results indicate that escalation of

cocaine intake during the first 15 minutes emerged during the later stages of the Short

Access period and continued throughout Long Access, particularly after a period of

abstinence.

Figure 1. Experimental timeline and cocaine infusions during Short (ShA) and Long Access (LgA). (A.) Experimental timeline with 66 hours (pink) and 90 hours (blue) indicating instances of abstinence. **(B.)** Cocaine infusions during the entire session. **(C.)** Cocaine infusions during the first 15 minutes of the session. Plots represent means and SEM. *$p<0.05$, **$p<0.01$, ***$p<.001$ vs. Short Access 01; ##$p<.01$, ##$p<.001$ vs. Long Access 01.

3.2. Locomotion During Noncontingent Sessions

3.2.1. Locomotion increases after noncontingent cocaine administration with no evidence of psychomotor sensitization or tolerance after Short or Long Access to cocaine self-administration.

To determine if raw locomotion differs between Noncontingent sessions following a noncontingent intravenous infusion of cocaine, a Type III ANOVA test was performed. A significant main effect of session number was found ($F_{(8, 290.88)}=17.38$, $p<.001$). Post hoc analyses only found significant differences within each respective Noncontingent session. Specifically, Drug sessions had significantly greater locomotion than their respective Baseline and Saline sessions (**Fig. 2A**).

To assess the change in the locomotor response to noncontingent cocaine administration, data was normalized to each rat's baseline and analyzed as percent difference from the respective Baseline sessions. A Type III ANOVA revealed a significant main effect of session number ($F_{(5, 176.97)}=16.36$, $p<.001$). Post hoc analyses revealed similar significant differences, where Drug sessions were significantly greater than their respective Saline and Baseline sessions, but no significant differences between Drug sessions (**Fig. 2B**). However, Saline 02 was significantly greater than Baseline 02, potentially an anticipatory conditioned stimulus effect as the last time they were handled as such, they received a noncontingent infusion of cocaine (Drug 01). The fact that this effect is not seen following Long Access suggests that the conditioned stimulus effect of picking up the animals and administering saline has disappeared, perhaps because at this stage (Long Access 14+) the animals have been handled extensively (~4 weeks) and therefore the act of being picked up from the operant

chamber is not associated anymore with cocaine. These results show that locomotion

did not increase in successive Noncontingent sessions following a noncontingent

intravenous infusion of cocaine at the population level.

3.2.2. Rats differentially develop psychomotor sensitization or tolerance after a history

of cocaine self-administration

To assess each individual rat's response to successive noncontingent infusions,

rats were grouped by their locomotion difference from Drug 01 to Drug 03. Rats who

increased were labeled "Sensitization" and rats who *decreased* were labeled as

"Tolerance". The locomotion in each Drug session was then analyzed by a percentage

change from Baseline (Drug 01). Following a 0.13 lambda Box-Cox power

transformation, a Type III Wald test found no significant main effect of behavioral

expression ($\chi^2(1)=4.8$, *p*=.059), but a significant effect of session number ($\chi^2(1)=6.47$,

p=.031) and their interaction ($\chi^2(1)=15.2$, *p*<.001). Post hoc analyses revealed rats in

the Tolerance group to have significantly decreased in their locomotion as early as Drug

02, with a further decrease in Drug 03. Similarly, although not significantly increased

from baseline, rats in the Sensitization group showed greater locomotion in Drug 02

than the Tolerance group that progressed further in Drug 03. These results demonstrate

markedly divergent profiles in the psychomotor response to cocaine that begin as early

as the first nine Short Access sessions and continue through Long Access (**Fig. 2C**).

The effect observed was not due to stereotypies in Tolerant rats (**Fig. S1**).

3.2.3. *Psychomotor sensitization and tolerance show no effect on cocaine self-*

administration

To test whether the development of psychomotor sensitization or tolerance had

an effect on cocaine self-administration over the entire session, a Type III Wald test was

performed for both Short and Long Access. During Short Access, a main effect of

session number was found ($\chi^2(9)=64.36$, $p<.001$), but not of behavioral expression

($\chi^2(1)=0.98$, $p=.967$), or their interaction ($\chi^2(9)=8.72$, $p=.967$. Similarly, in Long Access,

a main effect of session number was found ($\chi^2(13)=107.44$, $p<.001$), but not of

behavioral expression ($\chi^2(1)=0.47$, $p=1$), or their interaction ($\chi^2(13)=11.54$, $p=1$). These

results suggest that the development of psychomotor sensitization or tolerance to the

effect of cocaine has no effect on cocaine intake during Short or Long Access (**Fig. 2D**).

To determine if the development of psychomotor sensitization or tolerance has an

effect on cocaine infusions in the first 15 minutes, a Type III Wald test was performed. A

main effect of session number was found ($\chi^2(23)=154.13$, $p<.001$), but not of behavioral

expression ($\chi^2(1)=0.003$, $p=1$), or their interaction ($\chi^2(23)=30.78$, $p=.356$). These results

confirm that the development of psychomotor sensitization or tolerance to the effect of

cocaine has no effect on the loading behavior during cocaine self-administration (**Fig.**

2E).

Figure 2. Psychomotor sensitization and tolerance show no effect on cocaine self-administration. (A.) Raw locomotion (m) across Noncontingent sessions. ###*p*<.001 vs Baseline; ****p*<.001 vs Saline. **(B.)** Locomotion (m) percent difference from Baseline. #*p*<.05, ##*p*<.01, ###*p*<.001 vs Baseline; **p*<.05, ****p*<.001 vs Saline. **(C.)** Locomotion (m) percentage difference from Baseline (Drug 01) grouped by behavioral expression. Drug 01 < Drug 03, Sensitization (n = 14); Drug 01 ≥ Drug 03, Tolerance (n = 22). #*p*<.05, ###*p*<.001 vs Baseline; Within: $*p*<.05 vs Drug 02;

Between: ***$p<.001$ vs Tolerance Drug 03. **(D-E.)** Cocaine infusions over the entire session (D.) and within the first 15 minutes (E.) grouped by behavioral expression. Plots represent means and SEM.

3.3. **Pre-Lever Activity**

3.3.1. *Locomotion pre-lever increases from Short to Long Access and predicts cocaine intake during the first 15 minutes of self-administration*

To determine if locomotion pre-lever increased over the experiment, a Type III Wald test was performed. A significant main effect of session number was found following a 0.29 lambda Box-Cox power transformation ($x^2(23)=204.7$, $p<.001$). Post hoc analyses revealed significant increases at Short Access 07, Short Access 08, Long Access 09, and Long Access 13 in comparison with Short Access 01 (**Fig. 3A**). These results indicate that locomotion pre-lever increased from Short to Long Access.

To assess the influence of session type and abstinence on locomotion pre-lever, a separate analysis was conducted following a 0.28 lambda Box-Cox transformation. A Type III Wald test found significant main effects of session type ($x^2(1)=11.93$, $p=.002$), abstinence ($x^2(2)=14$, $p=.002$), and their interaction ($x^2(2)=27.57$, $p<.001$). Within Short Access, post hoc analyses revealed locomotion pre-lever at 90 hours was significantly greater than 18 hours. Within Long Access, 66 and 90 hours were found to be significantly greater than 18 hours. Interactively, at 18 hours there was a significant decrease in locomotion pre-lever from Short to Long Access. At 66 hours, there was a significant increase from Short to Long Access. Overall, locomotion pre-lever was

significantly greater in Long Access than in Short Access across abstinence (**Fig. 3B**). These results show that locomotion pre-lever increases with drug history and hours of abstinence.

To assess locomotion pre-lever predictability of cocaine intake in the first 15 minutes of the session and break point, correlation analyses were performed. Between locomotion pre-lever and cocaine infusions during the first 15 minutes, a positive correlation was found during Short Access (r_s(4222)=.5, p=.002) (**Fig. 3C**), and Long Access (r(4.29)=.59, p<.001, 95% CI [0.33, 0.77]) (**Fig. 3E**), but not during Shock (r_s(6952)=.18, p=.298) (**Fig. 3G**). Between locomotion pre-lever and breaking point, no correlations were found in both Progressive Ratio 01 (r_s(7497)=.04, p=.84) or Progressive Ratio 02 (r_s(6689)=.21, p=.219) (**Fig. 3D & 3F**). These results indicate that locomotion pre-lever is predictive of cocaine intake during the first 15 minutes in Short and Long Access, but is not predictive of the motivation to seek cocaine as measured by breaking point.

3.3.2. Active lever entrances per meter pre-lever escalate from Short to Long Access
and predict cocaine intake during the first 15 minutes of self-administration

Initial analyses indicated that nose entrances into the active lever area may be predictive of intake, however, there were large individual variability in locomotion. To account for animals with higher general activity having a greater probability of entering the active lever area by chance, we normalized the number of active lever area entrances to the overall locomotion in the chamber (active lever entrances per meter).

To determine if active lever entrances per meter increased over the experiment, a Type

III Wald test was performed. A significant main effect of session number was found

($\chi^2(23)$=234.44, p<.001). Post hoc comparisons with Short Access 01 revealed

significant increases in all sessions from Short Access 07 to Long Access 14, excluding

Long Access 01, 03, and 07 (**Fig. 3H**). These results confirm that active lever entrances

per meter increased over the experiment.

To assess the effects of session type and abstinence on active lever entrances

per meter, a separate Type III Wald test was performed. No effect of abstinence was

found ($\chi^2(2)$=1.7, p=.89), but a significant main effect of session type ($\chi^2(1)$=13.66,

p<.001), and their interaction ($\chi^2(2)$=17.47, p<.001). Post hoc analyses revealed no

significant differences between abstinence levels within Short Access. Within Long

Access, significant increases were found at 66 and 90 hours in comparison to 18 hours.

Interactively, at every level of abstinence there was found to be a significant increase

from Short to Long Access (**Fig. 3I**). These results confirm that active lever entrances

per meter increase with drug experience, with the strongest increases occurring after

longer periods of abstinence.

To assess active lever entrances per meter predictability of cocaine intake in the

first 15 minutes of the session and break point, correlation analyses were performed.

Between active lever entrances per meter and cocaine infusions during the first 15

minutes, a significant positive correlation was found during Short Access (r_s(3002)=.64,

p<.001) (**Fig. 3J**), Long Access (r(4.65)=.62, p<.001, 95% CI [0.37, 0.79]) (**Fig. 3L**), and

Shock (r_s(5199)=.38, p=.019) (**Fig. 3N**). Between active lever entrances per meter and

break point, a significant positive correlation was found during Progressive Ratio 01 (r-

$_s$(5173)=.33, p=.046) (**Fig. 3K**), but not Progressive Ratio 02 (r_s(7985)=.05, p=.753)

(**Fig. 3M**). These results show that active lever entrances per meter pre-lever are

predictive of both cocaine intake within the first 15 minutes and motivation measured as

break point following Short Access, but not after Long Access.

Locomotion (m) Pre-Lever

Active Lever Entrances per Meter Pre-Lever

Figure 3. Pre-lever activity increases with Long Access, hours of abstinence, and is predictive of cocaine infusions within the first 15 minutes of self-administration. (A.) Locomotion pre-lever across experiment. *p<.05, **p<.01 vs Short Access 01. **(B.)** Comparison of abstinence levels within and between session types. Within session type: ***p<.001 vs 18 hours; between session type and abstinence: ##p<.01 vs 18 hours Long Access, ###p<.001 vs 66 hours Short Access; between session type: $$$p<.001 vs Short Access. **(C.)** Mean locomotion pre-lever during Short Access correlates with mean cocaine infusions during first 15 minutes of Short Access. **(D.)** Locomotion pre-lever during Progressive Ratio 01 does not correlate with break point. **(E.)** Mean locomotion pre-lever during Long Access correlates with mean cocaine infusions during first 15 minutes of Long Access. **(F.)** Locomotion pre-lever during Progressive Ratio 02 does not correlate with break point. **(G.)** Locomotion pre-lever during Shock does not correlate with cocaine infusions during first 15 minutes of Shock. **(H.)** Active lever entrances per meter pre-lever across experiment. *p<.05, **p<.01, ***p<.001 vs ShA01. **(I.)** Comparison of abstinence levels within and between session types. Within session type: ***p<.001 vs 18 hours; between session type and abstinence: ##p<.01 vs 18, 66, and 90 hours Short Access; between session type: $$$p<.001 vs Short Access. **(J.)** Mean active lever entrances per meter pre-lever during Short Access correlates with mean cocaine infusions during first 15 minutes of Short Access. **(K.)** Active lever entrances per meter pre-lever during Progressive Ratio 01 correlates with break point. **(L.)** Mean active lever entrances per meter pre-lever during Long Access correlates with mean cocaine infusions during first 15 minutes of Long Access. **(M.)** Active lever entrances per meter pre-lever during Progressive Ratio 02

does not correlate with break point. **(N.)** Active lever entrances per meter pre-lever

during Shock correlates with cocaine infusions during first 15 minutes of Shock. Plots A,

B, H, and I represent mean and SEM. Correlation plots E and L are by Pearson, while

remaining correlations are by Spearman Rank.

3.4.   **Lever Entrances During Noncontingent Sessions**

3.4.1. Lever entrances per meter as percent difference from Noncontingent 01 show a

preference for the active lever after a history of self-administration

To determine if seeking behavior was expressed during Noncontingent sessions,

entrances per meter into the active and inactive lever areas were analyzed. Considering

the levers never extend during Noncontingent sessions they were treated as pre-lever

sessions. Examination of raw data showed low variation within Noncontingent sessions

and therefore all three sessions (e.g. Baseline, Saline, and Drug) were grouped to

analyze nose entrances into the active and inactive lever areas per meter.

Noncontingent 01 was then treated as the Baseline, and Noncontingent 02 and 03

active and inactive lever entrances per meter for each rat were normalized by the group

mean of Noncontingent 01 to account for high individual variation. Following a 0.23

lambda Yeo-Johnson power transformation, a Type III Wald test indicated a significant

main effect of session number ($\chi^2(1)=12.46$, $p=.002$), but not a significant main effect of

levers ($\chi^2(1)=3.12$, $p=.162$) or their interaction ($\chi^2(1)=3.49$, $p=.162$). Post hoc analyses

revealed Noncontingent 03 to be significantly greater than Noncontingent 02, prompting

a paired Student's t-test between the active and inactive levers in Noncontingent 03

where active was found to be greater than inactive. Direct comparisons to

Noncontingent 01 found that the inactive lever entrances per meter in Noncontingent 02

were significantly less than Noncontingent 01, while the active lever entrances per

meter in Noncontingent 03 were significantly greater than baseline. (**Fig. 4A**). These

results indicate that by Noncontingent 03 rats demonstrate preferential seeking behavior

toward the drug-associated active lever area.

To visualize the development of interest in the active lever, density heat maps of

each rat's nose node coordinates (x, y) for all sessions within each Noncontingent

session were overlain. Bounding boxes corresponding to the active and inactive lever

areas and the rounded mean lever entrances into each lever area were also plotted

(Noncontingent 01 active: M=22.2, SD=13.9; inactive: M=23.4, SD=12.3; Noncontingent

02 active: M=21.8, SD=16.1; inactive: M=18.4, SD=16.5; Noncontingent 03 active:

M=43.5, SD=42.1; inactive: M=24.7, SD=22.1). Student's t-tests between active and

inactive lever entrances within Noncontingent sessions revealed only Noncontingent 03

following a 0.23 lambda Yeo-Johnson power transformation achieved significance, with

active greater than inactive (**Fig. 4B**). These results highlight the development of

interest in the active lever after a history or self-administration as the high activity

regions migrate from the inactive side of the chamber in Noncontingent 01 to the active

side in Noncontingent 03.

3.4.2. Pre-lever activity during Short Access sessions 07-09 predicts active lever

entrances per meter during Noncontingent sessions.

To determine if the interest in the active lever was developed during self-

administration, rats were grouped as "High" (n=19) or "Low" (n=18) by a median-split z-

score of their pre-lever activity measures (locomotion and active lever entrances per

meter) during Short Access 07-09. These sessions were chosen as they were the first to

show significance compared to Short Access 01 and are the last Short Access sessions

before Noncontingent 02. First, the average of each variable was computed across the

three sessions before being standardized within the respective variables. The two z-

scores were then summed and divided by 2, where higher scores equate to greater pre-

session activity. The median of the group was then taken, and those greater than or

equal to the median were deemed "High", and those less than were deemed "Low".

Following a 0.2 lambda Yeo-Johnson power transformation, a Type III Wald test

indicated no main effect of session number ($\chi^2(1)=4.81$, $p=.059$), but a significant effect

of pre-lever activity score ($\chi^2(1)=29.53$, $p<.001$) and their interaction ($\chi^2(1)=9.39$,

$p=.006$). Post hoc analyses revealed that both Noncontingent 02 and 03 in the High pre-

lever activity group had significantly greater active lever entrances per meter than

baseline, and versus the Low pre-lever activity group in Noncontingent 02. For the Low

pre-lever activity group, Noncontingent 02 was significantly less than both baseline and

Noncontingent 03. There was no difference between groups in Noncontingent 03 (**Fig.**

4C). These results confirm that rats with high pre-lever activity during Short Access 07-

09 led to significant seeking in Noncontingent 02 that persisted in Noncontingent 03.

Furthermore, the Low pre-lever activity group developed an interest during Long Access

before Noncontingent 03.

3.4.3. Pre-lever activity during Short Access sessions 07-09 predicts cocaine self-administration in Short and Long Access

To determine if pre-lever activity during Short Access sessions 07-09 predicts cocaine intake over the entire session in Short Access, a Type III Wald test was performed. A significant main effect was found in session number ($\chi^2(9)$=125.52, p<.001), but not a significant effect of pre-lever activity score ($\chi^2(1)$=3.51, p=.169) or their interaction ($\chi^2(9)$=16.9, p=.169) (**Fig. 4D**). These results show that cocaine intake did not differ between groups in Short Access.

To determine if pre-lever activity during Short Access sessions 07-09 predicts cocaine intake over the entire session in Long Access, a Type III Wald test was performed. Significant main effects were found in session number ($\chi^2(13)$=62.03, p<.001), pre-lever activity score ($\chi^2(1)$=13.49, p<.001) and their interaction ($\chi^2(13)$=28.79, p=.015). Post hoc analyses revealed the High pre-lever activity group to have significantly greater cocaine infusions over the entire session in all Long Access sessions except Long Access 11, 13, and 14 (**Fig. 4D**). These results confirm that pre-lever activity during Short Access 07-09 was initially predictive of cocaine intake during Long Access, but that groups eventually converged. An analysis including sex as a factor showed that among rats with low pre-lever activity, females escalated their cocaine intake more rapidly than males during Long Access, indicating that the female bias in vulnerability emerges specifically in individuals with low initial pre-lever activity measures (**Fig. S2**, see Supplementary Material for sex-specific effects).

To determine if pre-lever activity during Short Access sessions 07-09 predicts cocaine intake within the first 15 minutes of the session, a Type III Wald test was

performed. A significant main effect was found in session number ($\chi^2(23)$=208.9, $p<$.001), but not in pre-lever activity score ($\chi^2(1)$=0.85, $p=$.742) or their interaction ($\chi^2(23)$=35.32, $p=$.134) (**Fig 4E**). These results suggest that pre-lever activity during Short Access 07-09 is not predictive of loading behavior between groups.

Considering the incongruency of the significant association between pre-lever variables and cocaine intake variables (**Fig. 3**), and the previous nonsignificant results between pre-lever activity groups and cocaine infusions in the first 15 minutes (**Fig. 4D**), correlations were performed between cocaine intake variables and pre-lever activity z-scores in Short Access and three separate phases of Long Access. For cocaine infusions during the entire session and pre-lever activity, significant positive correlations were found in Short Access ($r_s(2960)$=.65, $p<$.001; **Fig. 4F**), Long Access 01-05 ($r_s(3284)$=.61, $p<$.001; **Fig. 4G**), Long Access 06-10 ($r_s(3422)$=.59, $p<$.001; **Fig. 4H**), but not during Long Access 11-14 ($r_s(7013)$=.17, $p=$.318; **Fig. 4I**). For cocaine infusions during the first 15 minutes and pre-lever activity, significant positive correlations were similarly found in Short Access ($r_s(3094)$=.63, $p<$.001; **Fig. 4J**), Long Access 01-05 ($r_s(3301)$=.61, $p<$.001; **Fig. 4K**), Long Access 06-10 ($r_s(4023)$=.52, $p<$.001; **Fig. 4L**), but not during Long Access 11-14 ($r_s(6735)$=.2, $p=$.231; **Fig. 4M**). These results confirm that rats who showed markedly increased pre-lever activity during Short Access sessions 07-09 were more likely to consume more cocaine during the first 15 minutes and over the entire session in both Short and Long Access, but converged by the end of Long Access. Furthermore, these results support those shown in **Fig. 4C,** where no significant difference was found in active lever entrances per meter in Noncontingent 03, which occurs following Long Access 14.

Figure 4. Pre-lever activity during Short Access 07-09 predicts cocaine intake in Short and Long Access and seeking behavior during Noncontingent sessions.

(A.) Active (green) and inactive (red) lever entrances per meter as percentage difference from Baseline (Noncontingent 01). ##p<.01, ###p<.001 vs Baseline; *p<.05 vs inactive lever. **(B.)** Nose (*x, y*) pixel coordinate heatmaps of all rats within each Noncontingent session in relation to active (green) and inactive (red) Levers. Each plot represents ~109 sessions overlain revealing high and low activity zones. Numbers within lever bounding boxes are the mean sum of active and inactive lever entrances for all rats within each Noncontingent session. *p<.05 vs inactive lever. **(C.)** Active lever entrances per meter percent difference from Baseline (Noncontingent 01) grouped by pre-lever activity z-score during Short Access 07-09. ###p<.001 vs Baseline; ***p<.001 vs Noncontingent 02 Low group. **(D-E.)** Cocaine infusions over the entire session (D.) and within the first 15 minutes (E.) grouped by Short Access 07-09 pre-lever activity z-score. *p<.05, **p<.01, ***p<.001, ###p<.001 vs Low. **(F-I.)** Cocaine infusions over the entire session correlated using Spearman Rank with Short Access 07-09 pre-lever activity z-score in Short Access (F.), Long Access 01-05 (G.), Long Access 06-10 (H.), but not during Long Access 11-14 (I.). **(J-M.)** Cocaine infusions within the first 15 minutes correlated using Spearman Rank with Short Access 07-09 pre-lever activity z-score in Short Access (J.), Long Access 01-05 (K.), Long Access 06-10 (L.), but not during Long Access 11-14 (M.). Plots represent means and SEM.

4. Discussion

Here we demonstrate that cocaine infusions over the full session and within the first 15 minutes escalated during Short and Long Access in heterogeneous stock rats. Locomotion increased after a noncontingent cocaine infusion, and group-level analysis showed no signs of psychomotor sensitization or tolerance following Short Access (Noncontingent 02) or Long Access (Noncontingent 03). Individual analyses revealed strong divergence with some rats developing psychomotor sensitization and others showing tolerance, though neither influenced the level of cocaine self-administration. Pre-lever locomotion and active lever entrances per meter escalated from Short to Long Access, increased after instances of abstinence, and predicted early cocaine intake. Rats with high pre-lever activity during Short Access sessions 07-09 escalated lever entries across Noncontingent sessions and earned more infusions during Short Access and first two weeks of Long Access than low pre-lever activity rats, but these differences disappeared by Long Access 13 as low-activity rats escalated to similar levels. Together, the data dissociate psychomotor sensitization and tolerance from cocaine intake and identify pre-lever activity as a robust predictor of cocaine intake and escalation.

4.1. Cocaine Intake

Rats increased total cocaine intake as early as the second Short Access session and within the first 15 minutes as early as Short Access 07, while exhibiting significant escalation of intake after two weeks of Long Access. These findings align with our previous reports that heterogenous stock rats show increased cocaine intake during Short and Long Access [28], and the larger literature that Long Access induces escalation of intake [3,4,24,58].

4.2.   Psychomotor Expression

Rats showed individual differences in the propensity to develop psychomotor sensitization or tolerance to noncontingent cocaine administration. One alternative interpretation is that analyzing only ambulatory activity may misclassify some rats displaying stereotypic behavior, including head bobbing and weaving, sniffing or gnawing at high cocaine doses [8,59]. However, this is unlikely for several reasons. First, the dose used (1.5 mg/kg) is known to produce minimal stereotypic behavior and robust sensitization [6,33-35]. Second, the nose motion (a proxy for head bobbing, weaving, sniffing or gnawing) was not found to increase in Tolerant rats (**Fig. S1**).Third, visual inspection of the videos show that the most tolerant rat (M2677) did not exhibit stereotypic behavior while the most sensitized rat (M2668) showed pronounced psychomotor sensitization. A possible explanation for the difference between sensitized vs. tolerant rats is that rats classified as sensitized consumed less cocaine, while tolerant rats consumed more. However, intake analyses do not support this hypothesis: the two groups showed no differences in cocaine intake across the Short or Long Access session or within the first 15 minutes. These findings challenge the idea that Short Access produces only sensitization and Long Access only tolerance to the effects of cocaine [1,3-6,18,24], and they show that neither tolerance nor sensitization reliably predict future cocaine intake.

4.3.   Pre-Lever Activity Measures

Pre-lever variables locomotion and active lever entrances per meter increased from Short to Long Access and intensified following two-to-three days of abstinence.

Similarly, rats showed a significant increase in the active lever area entrances per meter in Noncontingent 03 in comparison to Noncontingent 01, suggesting sensitization of drug-seeking behavior. In contrast to measures of psychomotor sensitization or tolerance expression, both locomotion and active lever entrances per meter were highly predictive of intake. These findings align with the incentive salience literature, showing that repeated drug use leads to salience attribution to cues and environments that particularly manifests following a period of abstinence [5,6,15-16,19-20,33-35,46-47,61-68].

4.4.   Individual Differences in Pre-Lever Activity Measures

To investigate the individual propensity to attribute salience to drug-associated cues, a median z-score of pre-lever activity was calculated. Short Access sessions 07–09 were selected as they occurred within the same week (i.e., no abstinence periods) and were the first sessions in which both pre-lever variables (**Fig. 3A & 3H**) and infusions within the first 15 minutes (**Fig. 1C**) reached significance. Comparisons between High and Low pre-lever activity groups revealed that High pre-lever activity promoted an early escalation of cocaine intake during Long Access, but ultimately both High and Low pre-lever activity groups reached the same level of escalation after two weeks. Although no sex differences were observed in rates of pre-lever activity, females who exhibited low pre-lever activity escalated in cocaine infusions before low pre-lever activity males in Long Access (**Fig. S2**, see Supplementary for sex-specific effects). One possible explanation is that rats in the Low pre-lever activity group in Short Access was slower in developing incentive salience, and once expressed, their intake escalated to levels comparable to the High pre-lever activity group. Indeed, the significant difference

in active lever entrances per meter between activity groups in Noncontingent 02 and its

absence in Noncontingent 03 supports this interpretation. These findings align with

previous reports that individuals differ in their susceptibility to incentive salience

attribution [59,62,64-65], and that incentive salience may act as a catalyst of escalation

and an early predictor of vulnerability [26]. Another interpretation is that negative affect

or allostatic load may be driving the incentive salience and escalation in Long Access

[1,3,58,60]. While the current report does not test this specific hypothesis, there is

compelling evidence in both animal models and humans that acute withdrawal,

withdrawal-conditioned cues, negative affect, and psychosocial stress increase drug-

induced craving, drug-induced seeking, and amplifies drug cue-evoked brain activity.

While it is unlikely that these phenomena contribute to the effects observed in this report

during Short Access (although not impossible), it is likely that they contribute during

Long Access as animals experience acute withdrawal and negative emotional state and

that abstinence produces a large increase in pre-lever activity measures before the

session only in Long Access rats [69-73].

The phenotype we observed (rats with high pre-lever activity measured as

elevated active lever entrances per meter and locomotion) shares conceptual parallels

with the sign-tracker/goal-tracker framework [62]. Sign-trackers preferentially interact

with a discrete cue (the lever CS), while goal-trackers orient to the site of reward

delivery. Sign-trackers have been shown to initially self-administer more cocaine [66]

and choose cocaine over food [74]. However, important differences exist: sign-tracking

is defined in a purely Pavlovian context, with lever pressing having no instrumental

consequence, whereas our measure reflects operant-conditioned lever-directed

exploration. Notably, recent work has shown that sign-/goal-tracker status does not

reliably predict overall cocaine self-administration or escalation, although sign-trackers

may show greater resistance to punishment [75]. Thus, our results extend the sign-

/goal-tracker framework into an operant cocaine paradigm, highlighting that cue-bias

and responsivity can shape early intake and drive escalation of use even in resistant

populations with prolonged access.

4.5.   Limitations

The present study has a few limitations. One is the lack of a control group without

periods of abstinence (no weekends). A separate group participating in all sessions in

succession without instances of abstinence could further confirm the influence of

abstinence on incentive salience incubation. However, noncontingent sessions 02 and

03 did not follow a period of abstinence, and expressions of incentive salience and both

psychomotor sensitization and tolerance were still observed. Another limitation is the

use of pose-estimation over manual scoring of behavior could result in inaccuracies.

However, pose-estimation with SLEAP is remarkably accurate [38,76], and the instance

rendered sample videos display precise tracking of the catheter and nose nodes (see

supplementary videos).

4.6.   Summary

To summarize, this report demonstrates that individual differences in

psychomotor sensitization or tolerance had no influence on cocaine intake. In contrast,

sensitization of incentive salience during Short Access acts as a catalyst or early

predictor of escalation in cocaine self-administration during the first two weeks of Long

Access, but does not predict established escalation of cocaine intake after three weeks

of Long Access. Moreover, individuals initially resistant to incentive salience can, with

sufficient exposure, become sensitized and escalate cocaine use to the same level as

more susceptible individuals. Expression of incentive salience sensitization as reflected

in pre-lever activity measures of active lever entrances per meter and locomotion and

their subsequent influence on cocaine intake supports the idea that incentive salience

and psychomotor sensitization are mediated by distinct mechanisms [6-7,21,34]. The

development of the incentive salience profiles is accelerated by periods of abstinence

(weekend), which allows for incubation of craving, a progressive increase of craving

during early days of abstinence that can remain elevated following prolonged

abstinence [47]. The measure of approach toward the active lever area using video

recording before the start of the self-administration session may represent a novel

longitudinal behavioral marker of addiction vulnerability that can precisely track the

motivation for the drug with high accuracy without using any operant measures, the

drug itself, or any other training or conditioning that could interfere with the drug self-

administration. Such a measure may also be useful for medication development to allow

dissociation between drug-dependent and drug-independent effect of experimental

medication on drug seeking and taking.

Data Availability Statement

The datasets generated during and/or analyzed during the current study are

available from the corresponding author on reasonable request.

Acknowledgements

This research was done using services provided by the OSG Consortium [39-42] which is supported by the National Science Foundation awards #2030508 and #1836650.

Author Contributions

Jarryd Ramborger: Manuscript drafting and revising, data collection, experiment execution, experimental design, data analysis and interpretation

Joseph Mosquera: Data collection, experiment execution

Molly Brenna: Data collection, experiment execution

Benjamin Sichel: Data collection, experiment execution

Dyar Othman: Data collection, experiment execution

Sonja Plasil, PhD: Data collection, experiment execution

Elizabeth Sneddon, PhD: Data collection, experiment execution

Selene Zahedi: Data collection, experiment execution

Alex Morgan: Data collection, experiment execution

Supakorn Chonwattanagul: Data collection, experiment execution

Kathleen Bai: Data collection, experiment execution

Lindsey China: Data collection, experiment execution

Tu La: Data collection, experiment execution

Lisa Maturin: Data collection

Lieselot L. G. Carrette, PhD: Experimental design, manuscript drafting and revising

Olivier George, PhD: Manuscript drafting and revising, experimental design, data

analysis and interpretation, funding and resources, final approval

Funding

Funding was supported by DA043799 and the Preclinical Addiction Research

Consortium.

Competing Interests

Authors declare no competing interests.

References

1. Koob, G. F., & Le Moal, M. (1997). Drug abuse: Hedonic homeostatic dysregulation. *Science, 278*(5335), 52–58. https://doi.org/10.1126/science.278.5335.52

2. Koob, G. F., & Volkow, N. D. (2016). Neurobiology of addiction: a neurocircuitry analysis. *The lancet. Psychiatry*, *3*(8), 760–773. https://doi.org/10.1016/S2215-0366(16)00104-8.

3. Ahmed, S. H., & Koob, G. F. (1998). Transition from moderate to excessive drug intake: Change in hedonic set point. *Science, 282*(5387), 298-300. https://doi.org/10.1126/science.282.5387.298.

4. Ben-Shahar, O., Ahmed, S. H., Koob, G. F., & Ettenberg, A. (2004). The transition from controlled to compulsive drug use is associated with a loss of sensitization. *Brain Research, 995*(1), 46–54. https://doi.org/10.1016/j.brainres.2003.09.053

5. Calipari, E. S., Ferris, M. J., Zimmer, B. A., Roberts, D. C. S., & Jones, S. R. (2013). Temporal pattern of cocaine intake determines tolerance vs sensitization of cocaine effects at the dopamine transporter. *Neuropsychopharmacology, 38*(12), 2385–2392. https://doi.org/10.1038/npp.2013.136.

6. Carr, C. C., Ferrario, C. R., & Robinson, T. E. (2020). Intermittent access cocaine self-administration produces psychomotor sensitization: Effects of withdrawal, sex, and cross-sensitization. *Psychopharmacology, 237*(6), 1795–1812. https://doi.org/10.1007/s00213-020-05500-4.

7. Dalia AD, Norman MK, Tabet MR, Schlueter KT, Tsibulsky VL, Norman AB (1998) Transient amelioration of the sensitization of cocaine-induced behaviors in rats

by the induction of tolerance. Brain Res 797:29–34.

https://doi.org/10.1016/S0006-8993(98)00323-0.

8.   Flagel, S. B., & Robinson, T. E. (2007). Quantifying the psychomotor activating

effects of cocaine in the rat. *Behavioural Pharmacology, 18*(4), 297–302.

https://doi.org/10.1097/FBP.0b013e3281f522a4.

9.   Lett, B. T. (1989). Repeated exposures intensify rather than diminish the

rewarding effects of amphetamine, morphine, and cocaine.

Psychopharmacology, 98(3), 357-362. https://doi.org/10.1007/BF00451688.

10.   Porrino, L. J. (1993). Functional consequences of acute cocaine treatment

depend on route of administration. *Psychopharmacology, 112*(3), 343–351.

https://doi.org/10.1007/BF02244948.

11.   Swain, Y., Muelken, P., LeSage, M. G., Gewirtz, J. C., & Harris, A. C. (2018).

Locomotor activity does not predict individual differences in morphine self-

administration in rats. *Pharmacology Biochemistry and Behavior*, 166, 48-56.

https://doi.org/10.1016/j.pbb.2018.01.008.

12.   Vanderschuren, L. J. M. J., Schmidt, E. D., De Vries, T. J., Van Moorsel, C. A. P.,

Tilders, F. J. H., & Schoffelmeer, A. N. M. (1999). A single exposure to

amphetamine is sufficient to induce long-term behavioral, neuroendocrine, and

neurochemical sensitization in rats. *Journal of Neuroscience, 19*(21), 9579–9586.

https://doi.org/10.1523/JNEUROSCI.19-21-09579.1999

13.   Vanderschuren, L. J. M. J., De Vries, T. J., Wardeh, G., Hogenboom, F. A. C. M.,

& Schoffelmeer, A. N. M. (2001). A single exposure to morphine induces long-

lasting behavioural and neurochemical sensitization in rats. *European Journal of*

788  *Neuroscience, 14*(8), 1533–1538. https://doi.org/10.1046/j.0953-

789  816x.2001.01771.x

790 14. Hammer, R. P. Jr., Egilmez, Y., & Emmett-Oglesby, M. W. (1997). Neural

791  mechanisms of tolerance to the effects of cocaine. *Behavioural Brain Research*,

792  *84*(1-2), 225-239. https://doi.org/10.1016/s0166-4328(97)83332-3.

793 15. Kalivas PW, Sorg BA, Hooks MS (1993). The pharmacology and neural circuitry

794  of sensitization to psychostimulants. Behav Pharmacol. 1993;4(4):315-334.

795  PMID: 11224200.

796 16. Vanderschuren, L. J. M. J., & Kalivas, P. W. (2000). Alterations in dopaminergic

797  and glutamatergic transmission in the induction and expression of behavioral

798  sensitization: A critical review of preclinical studies. *Psychopharmacology, 151*(2),

799  99–120. https://doi.org/10.1007/s002130000493.

800 17. Vezina, P. (2004). Sensitization of midbrain dopamine neuron reactivity and the

801  self-administration of psychomotor stimulant drugs. *Neuroscience &*

802  *Biobehavioral Reviews, 27*(8), 827–839.

803  https://doi.org/10.1016/j.neubiorev.2003.11.001.

804 18. Solomon, R. L. (1980). The opponent-process theory of acquired motivation: The

805  costs of pleasure and the benefits of pain. *American Psychologist*, 35(8), 691-

806  712. https://doi.org/10.1037/0003-066X.35.8.691

807 19. Robinson, T. E. & Berridge, K. C.(1993). The neural basis of drug craving: An

808  incentive-sensitization theory of addiction. *Brain Research Reviews*, 18(3), 247-

809  291. https://doi.org/10.1016/0165-0173(93)90013-P.

810   20.   Ciccocioppo, R., Martin-Fardon, R., & Weiss, F. (2004). Stimuli associated with a

811   single cocaine experience elicit long-lasting cocaine-seeking. *Nature*

812   *Neuroscience, 7*(5), 495-496. https://doi.org/10.1038/nn1239.

813   21.   Izenwasser, S., & French, D. (2002). Tolerance and sensitization to the

814   locomotor-activating effects of cocaine are mediated via independent

815   mechanisms. *Pharmacology Biochemistry and Behavior, 73*(4), 877–882.

816   https://doi.org/10.1016/s0091-3057(02)00942-5.

817   22.   Kalivas, P. W., & Alesdatter, J. E. (1993). Involvement of N-methyl-D-aspartate

818   receptor stimulation in the ventral tegmental area and amygdala in behavioral

819   sensitization to cocaine. *Journal of Pharmacology and Experimental*

820   *Therapeutics, 267*(1), 486-495. Retrieved from

821   http://jpet.aspetjournals.org/content/267/1/486.abstract.

822   23.   Ungless, M., Whistler, J., Malenka, R. *et al.* Single cocaine exposure *in vivo*

823   induces long-term potentiation in dopamine neurons. *Nature* 411, 583–587

824   (2001). https://doi.org/10.1038/35079077

825   24.   Ahmed, S. H., & Cador, M. (2006). Dissociation of psychomotor sensitization

826   from compulsive cocaine consumption. Neuropsychopharmacology, 31(3), 563–

827   571. https://doi.org/10.1038/sj.npp.1300834.

828   25.   Kawa AB, Bentzley BS, Robinson TE. Less is more: prolonged intermittent

829   access cocaine self-administration produces incentive-sensitization and

830   addiction-like behavior. Psychopharmacology (Berl). 2016 Oct;233(19-20):3587-

831   602. doi: 10.1007/s00213-016-4393-8. Epub 2016 Aug 2. PMID: 27481050;

832   PMCID: PMC5023484.

26.   Kawa AB, Valenta AC, Kennedy RT, Robinson TE. Incentive and dopamine

sensitization produced by intermittent but not long access cocaine self-

administration. Eur J Neurosci. 2019 Aug;50(4):2663-2682. doi:

10.1111/ejn.14418. Epub 2019 May 2. PMID: 30968487; PMCID: PMC6742545.

27.   Carrette, L. L. G., de Guglielmo, G., Kallupi, M., Maturin, L., Brennan, M.,

Boomhower, B., et al.. (2021). The Cocaine and Oxycodone Biobanks, two

repositories from genetically diverse and behaviorally characterized rats for the

study of addiction. *eNeuro*, *8*(3), ENEURO.0033-21.2021.

https://doi.org/10.1523/ENEURO.0033-21.2021.

28.   Giordano de Guglielmo, Carrette, L. L. G., Kallupi, M., Brennan, M., Boomhower,

B., Maturin, L., et al.. (2024). Large-scale characterization of cocaine addiction-

like behaviors reveals that escalation of intake, aversion-resistant responding,

and breaking-points are highly correlated measures of the same construct. *eLife*,

12, RP90422. https://doi.org/10.7554/eLife.90422

29.   Solberg Woods LC, Palmer AA (2019) Using heterogeneous stocks for fine-

mapping genetically complex traits. *Methods Mol Biol* 2018:233–247.

doi: 10.1007/978-1-4939-9581-3_11.

30.   Argilli, E., Sibley, D. R., Malenka, R. C., England, P. M., & Bonci, A. (2008).

Mechanism and time course of cocaine-induced long-term potentiation in the

ventral tegmental area. *The Journal of Neuroscience, 28*(37), 9092–9100.

https://doi.org/10.1523/JNEUROSCI.1001-08.2008.

31.   Booze, R. M., Lehner, A. F., Wallace, D. R., Welch, M. A., & Mactutus, C. F.

(1997). Dose-response cocaine pharmacokinetics and metabolite profile

following intravenous administration and arterial sampling in unanesthetized,

freely moving male rats. *Neurotoxicology and Teratology*, 19(1), 7–15.

https://doi.org/10.1016/S0892-0362(96)00180-8.

32. Pan, H.-T., Menacherry, S., & Justice, J. B., Jr. (1991). Differences in the

pharmacokinetics of cocaine in naive and cocaine-experienced rats. *Journal of*

Neurochemistry, 56(4), 1299–1306. https://doi.org/10.1111/j.1471-

4159.1991.tb11429.x

33. Samaha, A. N., Li, Y., & Robinson, T. E. (2002). The rate of intravenous cocaine

administration determines susceptibility to sensitization. *The Journal of*

neuroscience : the official journal of the Society for Neuroscience, 22(8), 3244–

3250. https://doi.org/10.1523/JNEUROSCI.22-08-03244.2002.

34. Samaha, A.-N., Mallet, N., Ferguson, S. M., Gonon, F., & Robinson, T. E. (2004).

The rate of cocaine administration alters gene regulation and behavioral

plasticity: Implications for addiction. *The Journal of Neuroscience*, *24*(28), 6362–

6370. https://doi.org/10.1523/JNEUROSCI.1424-04.2004.

35. Samaha, A.-N., & Robinson, T. E. (2005). Why does the rapid delivery of drugs to

the brain promote addiction? *Trends in Pharmacological Sciences, 26*(2), 82-87.

https://doi.org/10.1016/j.tips.2004.12.007.

36. Woolverton, W. L., & Wang, Z. (2004). Relationship between injection duration,

transporter occupancy, and reinforcing strength of cocaine. *European Journal of*

Pharmacology, *486*(3), 251–257. https://doi.org/10.1016/j.ejphar.2004.01.003.

37. Ramborger, J., Kalra, S., Mosquera, J., Smith, A. C. W., & George, O. (2024).

High quality, high throughput, and low-cost simultaneous video recording of 60

animals in operant chambers using PiRATeMC. *Journal of Neuroscience*

Methods, 411, 110270. https://doi.org/10.1016/j.jneumeth.2024.110270.

38.   T.D. Pereira, N. Tabris, A. Matsliah, D. M. Turner, J. Li, S. Ravindranath, et al..

Sleap: A deep learning system for multi-animal pose tracking. *Nature Methods*,

19(4), 2022. https://doi.org/10.1038/s41592-022-01426-1.

39.   OSG. (2006). OSPool. OSG. https://doi.org/10.21231/906P-4D78

40.   Pordes, R., Petravick, D., Kramer, B., Olson, D., Livny, M., Roy, A., et al.. (2007).

The open science grid. J. Phys. Conf. Ser., 78, 012057.

https://doi.org/10.1088/1742-6596/78/1/012057.

41.   Sfiligoi, I., Bradley, D. C., Holzman, B., Mhashilkar, P., Padhi, S., & Wurthwein, F.

(2009). The pilot way to grid resources using glideinWMS. 2009 WRI World

Congress on Computer Science and Information Engineering, 2, 428–432.

https://doi.org/10.1109/CSIE.2009.950.

42.   OSG. (2015). Open Science Data Federation. OSG.

https://doi.org/10.21231/0KVZ-VE57

43.   R Core Team. (2023). *R: A language and environment for statistical computing*

[Computer software]. R Foundation for Statistical Computing. https://www.R-

project.org/

44.   Posit Software, PBC. (2023). *RStudio: Integrated development environment for R*

[Computer software]. https://posit.co/

45.   Rueden, C.T., Schindelin, J., Hiner, M.C. *et al.* ImageJ2: ImageJ for the next

generation of scientific image data. *BMC Bioinformatics* **18**, 529 (2017).

https://doi.org/10.1186/s12859-017-1934-z.

46.   Berridge, K. C., & Robinson, T. E. (1998). What is the role of dopamine in reward:

903   Hedonic impact, reward learning, or incentive salience? *Brain Research*

904   *Reviews, 28*(3), 309–369. https://doi.org/10.1016/S0165-0173(98)00019-8.

905   47.   Li, X., Caprioli, D., & Marchant, N. J. (2014). Recent updates on incubation of

906   drug craving: A mini-review. *Addiction Biology, 20*(5), 872–876.

907   https://doi.org/10.1111/adb.12205.

908   48.   Bates D, Mächler M, Bolker B, Walker S (2015). "Fitting Linear Mixed-Effects

909   Models Using lme4." *Journal of Statistical Software*, **67**(1), 1–

910   48. doi:10.18637/jss.v067.i01.

911   49.   Kuznetsova A, Brockhoff PB, Christensen RHB (2017). "lmerTest Package: Tests

912   in Linear Mixed Effects Models." *Journal of Statistical Software*, **82**(13), 1–

913   26. doi:10.18637/jss.v082.i13.

914   50.   Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A,

915   et al. (2017). "glmmTMB Balances Speed and Flexibility Among Packages for

916   Zero-inflated Generalized Linear Mixed Modeling." *The R Journal*, 9(2), 378–

917   400. doi:10.32614/RJ-2017-066.

918   51.   Luke, S.G. Evaluating significance in linear mixed-effects models in R. *Behav*

919   *Res* **49**, 1494–1502 (2017). https://doi.org/10.3758/s13428-016-0809-y.

920   52.   Hartig, F. (2024). *DHARMa: Residual diagnostics for hierarchical (multi-level /*

921   *mixed) regression models* (Version 0.4.7) [R package]. CRAN.

922   https://doi.org/10.32614/CRAN.package.DHARMa.

53. Meteyard, L., & Davies, R. A. I. (2020). Best practice guidance for linear mixed-

effects models in psychological science. *Journal of Memory and Language, 112*,

104092. https://doi.org/10.1016/j.jml.2020.104092.

54. Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (3rd ed.).

Sage. https://www.john-fox.ca/Companion/.

55. Lenth, R. V. (2025). *emmeans: Estimated marginal means, aka least-squares*

means (Version 1.10.7) [R package]. CRAN.

https://doi.org/10.32614/CRAN.package.emmeans.

56. Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in

multiple testing under dependency. *The Annals of Statistics, 29*(4), 1165–1188.

https://doi.org/10.1214/aos/1013699998.

57. Kassambara, A. (2023). rstatix: Pipe-friendly framework for basic statistical tests

(Version 0.7.2) [Computer software].

https://rpkgs.datanovia.com/rstatix/[](https://rpkgs.datanovia.com/rstatix/authors.h

tml).

58. Ahmed, S. H., & Koob, G. F. (1999). Longlasting increase in the set point for

cocaine selfadministration after escalation in rats. *Psychopharmacology, 146*(3),

303–312. https://doi.org/10.1007/s002130051121.

59. Flagel, S. B., Watson, S. J., Akil, H., & Robinson, T. E. (2008). Individual

differences in the attribution of incentive salience to a reward-related cue:

Influence on cocaine sensitization. *Behavioural Brain Research, 186*(1), 48–56.

https://doi.org/10.1016/j.bbr.2007.07.022

60.   Ahmed, S. H., Kenny, P. J., Koob, G. F., & Markou, A. (2002). Neurobiological

evidence for hedonic allostasis associated with escalating cocaine use. *Nature*

Neuroscience, 5(7), 625–626. https://doi.org/10.1038/nn872.

61.   Blanchard, M. M., Mendelsohn, D., & Stamp, J. A. (2009). The HR/LR model:

Further evidence as an animal model of sensation seeking. *Neuroscience &*

Biobehavioral Reviews, 33(7), 1145-1154.

https://doi.org/10.1016/j.neubiorev.2009.05.009.

62.   Flagel, S.B., Watson, S.J., Robinson, T.E. *et al.* Individual differences in the

propensity to approach signals vs goals promote different adaptations in the

dopamine system of rats. *Psychopharmacology* **191**, 599–607 (2007).

https://doi.org/10.1007/s00213-006-0535-8.

63.   Piazza, P. V., Deroche-Gamonent, V., Rouge-Pont, F., & Le Moal, M. (2000).

Vertical Shifts in Self-Administration Dose–Response Functions Predict a Drug-

Vulnerable Phenotype Predisposed to Addiction. *The Journal of Neuroscience*,

20(11), 4226-4232. https://doi.org/10.1523/JNEUROSCI.20-11-04226.2000.

64.   Post, R. M., Lockfeld, A., Squillace, K. M., & Contel, N. R. (1981). Drug-

environment interaction: Context dependency of cocaine-induced behavioral

sensitization. *Life Sciences*, *28*, 755–760. https://doi.org/10.1016/0024-

3205(81)90157-0.

65.   Saunders, B. T., & Robinson, T. E. (2010). A cocaine cue acts as an incentive

stimulus in some but not others: Implications for addiction. Biological Psychiatry,

67(8), 730–736. https://doi.org/10.1016/j.biopsych.2009.11.015

66.   Saunders, B. T., & Robinson, T. E. (2011). Individual variation in the motivational

properties of cocaine. *Neuropsychopharmacology, 36*(8), 1668–1676.

https://doi.org/10.1038/npp.2011.48.

67.   Stewart, J., & Badiani, A. (1993). Tolerance and sensitization to the behavioral

effects of drugs. *Behavioural Pharmacology, 4*(4), 289-312.

68.   Weiss, S. R. B., Post, R. M., Pert, A., Woodward, R., & Murman, D. (1989).

Context-dependent cocaine sensitization: Differential effect of haloperidol on

development versus expression. *Pharmacology, Biochemistry, and Behavior,*

34(3), 655-661. https://doi.org/10.1016/S0091-3057(89)80001-2.

69.   Balleine, B. W. (2011). Sensation, incentive learning, and the motivational control

of appetitive behavior. In G. F. Koob, M. Le Moal, & R. F. Thompson (Eds.),

Neurobiology of addiction (pp. 241–259). Academic Press.

70.   Carmack, S. A., Kirson, E. D., & Markou, A. (2019). Heroin addiction engages

negative emotional learning brain circuits in rats. *The Journal of Clinical*

Investigation, 129(6), 2480–2484. https://doi.org/10.1172/JCI125534

71.   Hutcheson, D. M., et al. (2001). The role of withdrawal in heroin addiction. *Nature*

Neuroscience, 4(10), 1031–1034. https://doi.org/10.1038/nn0901-943

72.   Kenny, P. J., Chen, S. A., Kitamura, O., Markou, A., & Koob, G. F. (2006).

Conditioned withdrawal drives heroin consumption and decreases reward

sensitivity. *The Journal of Neuroscience, 26*(22), 5894–5900.

https://doi.org/10.1523/JNEUROSCI.0740-06.2006

73.   Pantazis, C. B., Gonzalez, L. A., Tunstall, B. J., Carmack, S. A., Koob, G. F.,

Vendruscolo, L. F., … & George, F. (2021). Cues conditioned to withdrawal and

negative reinforcement: Neglected but key motivational elements driving opioid

addiction. *Science Advances, 7*(46), eabf0364.

https://doi.org/10.1126/sciadv.abf0364

74.   Sinha, R., Fuse, T., Aubin, L. R., & O'Malley, S. S. (2000). Psychological stress,

drug-related cues and cocaine craving. *Psychopharmacology (Berlin), 152*(2),

140–148. https://doi.org/10.1007/s002130000499

75.   Tunstall, B. J., & Kearns, D. N. (2015). Sign-tracking predicts increased choice of

cocaine over food in rats. Behavioural Brain Research, 281, 222–228.

https://doi.org/10.1016/j.bbr.2014.12.034.

76.   Pohořalá, V., Enkel, T., Bartsch, D. *et al.* Sign- and goal-tracking score does not

correlate with addiction-like behavior following prolonged cocaine self-

administration. *Psychopharmacology* **238**, 2335–2346 (2021).

https://doi.org/10.1007/s00213-021-05858-z

77. Pereira, T. D., Shaevitz, J. W., & Murthy, M. (2020). Quantifying behavior to

understand the brain. *Nature Neuroscience, 23*(12), 1537–1549.

https://doi.org/10.1038/s41593-020-00734-z.

1006

Supplementary Material

Incentive Salience, not Psychomotor Sensitization or Tolerance, Drives Escalation of Cocaine Self-Administration in Heterogeneous Stock Rats

Introduction

The following document provides a detailed account of the model fitting and choice process for the linear and generalized mixed models utilized in the repeated measure ANOVA statistical analyses, a test of stereotypies to accompany **Fig. 2C**, and analyses with sex as a factor. First is the step-by-step process of testing and comparing models with different model fits. Next are the assumption test visualizations provided by the simulateResiduals() function from the RStudio DHARMa package (Hartig, 2024) for each final mixed model, including the analyses of sex and stereotypies. The function had the parameters "refit = FALSE" and "n = 500" for residual simulations. The default number of simulations is 250, and 500 was chosen for computation time and greater precision (Hartig, 2024). Each plot is identified alphabetically for how they appear in the manuscript, and each model's statistics can be found in the excel spreadsheet table. Next are the results and figure for the stereotypies analysis. And lastly are the analyses with sex included as a factor.

1. **Model Testing and Comparison of Fit Process**

To best explain the process of testing and comparing different linear and generalized linear mixed model combinations, an example is used to fit the steps in context. The chosen example is the first statistical analysis in the manuscript and

supplementary excel spreadsheet, i.e., answering the question: "Do cocaine infusions in the entire session differ between sessions in Short Access"? Therefore, the response variable is Cocaine Infusions, the fixed effect is Session Number, and given this is a repeated measure analysis, the random effect is each individual Rat. The formula, as it would appear in RStudio, would then be:

response variable ~ fixed effects + random effects

or in this example,

Cocaine Infusions ~ Session Number + (1|Rat)

This formula is then used to fit a model of the data using the linear mixed effects model function lmer() (Bates et al., 2015; Kuznetsova, Brockhoff, & Christensen, 2017) or a generalized linear mixed effects model with glmmTMB() (Brooks et al., 2017). All models were fit with maximum likelihood to allow valid statistical comparisons of fixed effects structures and to support post hoc testing as recommended (Bates et al., 2015; Luke, 2017; Meteyard & Davies, 2020). The process is as follows:

1. The first step was to determine the optimal lambda for Box-Cox of Yeo-Johnson power transformations to achieve normality for a linear regression model. Box-Cox are for response variables that do not contain zero or negatives, and Yeo-Johnson for those that do. For this example, a Yeo-Johnson was performed.

2. Two linear mixed models using lmer() were then produced: one without power transformation, and one with power transformation.

3. The two linear mixed models were then compared with a likelihood ratio test.

4. The two linear mixed models were then visualized with simulateResiduals() in the order of the likelihood ratio test results, i.e., worst to best, and their results documented. These steps elucidate the modeling done by linear methods.

5. Next, a number of generalized linear mixed models are fit using glmmTMB. Unlike linear mixed models, generalized mixed models allow for specification of both the distribution family (e.g., gaussian, logarithmic, tweedie, etc.) and the dispersion structure, enabling more flexible modeling of data with non-normal distributions or heterogeneous variance. For example, some distribution families require nonnegative and whole numbers (negative binomial distributions), while others do not have such constraints (gaussian). For this example, six separate base models were fit to the data: without transformation, with a Yeo-Johnson transformation, with a tweedie distribution, and all three available negative binomial distributions.

6. Similarly to the linear mixed models, the generalized mixed models were then compared using a likelihood ratio test.

7. The generalized mixed models were then visualized with simulateResiduals() in the order of the likelihood ratio test results from worst to best. Steps 5-7 reveals the best family distribution to be used.

8. The best-fitting model from the previous step is then used as a baseline to evaluate alternative dispersion structures within the generalized linear mixed model. Whereas linear mixed models estimate only the conditional mean of the outcome variable (e.g., $Y \sim X$), generalized linear mixed models permit simultaneous modeling of both the mean and the variance. By default, variance

is assumed to follow the form dictated by the specified distributional family. However, the inclusion of dispersion parameters enables the modeling of heteroskedasticity by allowing fixed effects (e.g., session number, session type, etc.) to account for variation in the residual variance. This step evaluates whether explicitly modeling dispersion improves overall model fit and residual behavior. In the case of this example, a few of the dispersion parameters fit were:

Cocaine Infusions ~ Session Number + (1|Rat)

Dispersion = Session Number

Dispersion = Session Number + (1|Rat)

Dispersion = Abstinence

Dispersion = Abstinence + (1|Rat)

These specifications allow the variance to vary as a function of Session, as a function of both Session and individual Rats, by each level of Abstinence, or Abstinence and each individual Rat.

9. Each generalized mixed model from step 8 was then compared using a likelihood ratio test.

10. Each model from step 8 was then visualized with simulateResiduals() in the order of the likelihood ratio test results. It is in this step that the best linear mixed model simulateResiduals() results from above in step 4 is compared with the generalized mixed model combinations.

11. The model in Step 10 that yielded the most favorable simulateResiduals() diagnostics was selected as the final model. Primary emphasis was placed on

the Kolmogorov–Smirnov test to assess residual uniformity and overall model fit to the data. However, in instances where warranted, a balanced interpretation of all diagnostic outputs—which include dispersion, outlier detection, and observed versus predicted relationships—was preferred, as the Kolmogorov–Smirnov test is known to be sensitive to large sample sizes, which may yield statistically significant deviations despite an acceptable practical fit (Hartig, 2024; Razali & Wah, 2011).

2. DHARMa simulateResiduals() Assumption Check Results

Below are the plotted outputs generated by simulateResiduals() for the final model used in each of the repeated measure statistical analyses found in the manuscript. Red text indicates a failure of an assumption test. The plots are alphabetical in the order in which they appear in the manuscript and in the accompanying supplementary excel spreadsheet. The excel spreadsheet contains a table with the final formula of the models as well as the statistical results generated by summary(). Results of models not selected are available from the corresponding author on reasonable request.

A. Cocaine Infusions in Short Access

B. Cocaine Infusions in Long Access

C. Cocaine Infusions First Fifteen Minutes

D. Locomotion Raw in Noncontingent Sessions

E. Locomotion Percent Difference From Baseline in Noncontingent Sessions

F. Locomotion Percent Difference From Drug 01 by Behavioral Expression

G. Cocaine Infusions in Short Access by Behavioral Expression

H. Cocaine Infusions in Long Access by Behavioral Expression

I. Cocaine Infusions First 15 Minutes by Behavioral Expression

J. Locomotion Pre-Lever

K. Locomotion Pre-Lever by Session Type and Abstinence

L. Active Lever Entrances per Meter Pre-Lever

M. Active Lever Entrances per Meter by Session Type and Abstinence

N. Entrances per Meter Percent Difference From Noncontingent 01

O. Active Lever Entrances per Meter Percent Difference From Noncontingent 01 by Pre-Lever Activity

P. Cocaine Infusions in Short Access by Pre-Lever Activity

Q. Cocaine Infusions in Long Access by Pre-Lever Activity

R. Cocaine Infusions First 15 Minutes by Pre-Lever Activity

S. Nose Motion Percent Difference From Drug 01 by Behavioral Expression

T. Cocaine Infusions in Short Access by Sex

U. Cocaine Infusions in Long Access by Sex

V. Locomotion Percent Difference From Baseline in Noncontingent Sessions by Sex

W. Cocaine Infusions in Short Access by Behavioral Expression and Sex

X. Cocaine Infusions in Long Access by Behavioral Expression by Sex

Y. Active Lever Entrances per Meter by Session Type, Abstinence, and Sex

Z. Entrances per Meter Percent Difference From Noncontingent 01 by Sex

AA. Cocaine Infusions in Short Access by Pre-Lever Activity and Sex

BB.   Cocaine Infusions in Long Access by Pre-Lever Activity and Sex

3. Nose Motion Analysis

3.1. Tolerant rats show no evidence of stereotypies during Drug sessions

To assess if rats in the Tolerance group were miscategorized due to stereotypies, nose motion percentage change from Baseline (Drug 01) between behavioral expression groups was analyzed. Following a 0.0002 lambda Box-Cox power transformation, a Type III Wald test found a significant main effect of behavioral expression ($\chi^2(1)$=5.77, p=.034), of session number ($\chi^2(1)$=9.87, p=.005) and their interaction ($\chi^2(1)$=18.95, p<.001). Post hoc analyses revealed rats in the Tolerance group to have significantly decreased their nose motion in Drug 03 compared to both baseline and Drug 02. Conversely, the Sensitization group showed significantly increased nose motion in both Drug 02 and Drug 03 compared to baseline, and Drug 03

greater than Drug 02. Comparatively, the Sensitization group had greater nose motion within Drug 02 and Drug 03 than the Tolerance group (**Fig. S1**). These results demonstrate that the observed effect seen in **Fig. 2C** was not due to stereotypies, and that behavioral expression groups were sufficiently categorized based on their locomotor difference from Drug 01 to Drug 03.

Supplementary Figure 1. Tolerant rats show no evidence of stereotypies during Drug sessions. Nose motion (m) percentage difference from Baseline (Drug 01) grouped by psychomotor behavioral expression. ##p<.01, ###p<.001 vs Baseline; Within group: $$p$<.05, $$$p$<.01 vs Drug 02; Between groups: *p<.05 vs Tolerance Drug 02; ***p<.001 vs Tolerance Drug 03.

4. Sex as a Factor Analyses

The following analyses were conducted with sex included as a factor. These specific tests were chosen as they are critical results to the manuscript interpretation. The main focus was the full interaction between factors and reported below.

4.1. No effect of sex on cocaine infusions in Short Access

To determine if sexes differed in cocaine infusions during Short Access, a Type III Wald test was performed. Results showed no significant interaction between sex and session number ($\chi^2(9)=14.3$, $p=.233$). These results show that cocaine intake did not differ between sexes in Short Access.

4.2. No effect of sex on cocaine infusions in Long Access

To determine if sexes differed in cocaine infusions during Long Access, a Type III Wald test was performed. Results showed no significant interaction between sex and session number ($\chi^2(13)=16.77$, $p=.583$). These results show that cocaine intake did not differ between sexes in Long Access.

4.3. No effect of sex on locomotion percent difference from baseline during Noncontingent sessions

To determine if sexes differed in their locomotion perfect difference from baseline during Noncontingent sessions, a Type III Wald test was performed. Results showed no significant interaction between sex and session number ($\chi^2(5)=10.72$, $p=.159$). These results show that locomotion percent difference from baseline did not differ between sexes in Noncontingent sessions.

4.4. No effect of sex x behavioral expression on cocaine infusions across Short Access sessions

To determine if a sex x behavioral expression x session number interaction had an effect on cocaine infusions during Short Access, a Type III Wald test was performed. Results showed the interaction was not significant ($\chi^2(9)=14.31$, $p=.607$), indicating that

cocaine intake did not differ between sexes as a function of behavioral expression across Short Access sessions.

4.5. No effect of sex x behavioral expression on cocaine infusions across Long Access sessions

To determine if a sex x behavioral expression x session number interaction had an effect on cocaine infusions during Long Access, a Type III Wald test was performed. Results showed no significant interaction effect ($\chi^2(13)=9.48$, $p=.736$), indicating that cocaine intake did not differ between sexes as a function of behavioral expression across Long Access sessions.

4.6. No effect of sex x session type x abstinence on active lever entrances per meter pre-lever

To determine if a sex x session type x abstinence interaction had an effect on active lever entrances per meter pre-lever, a Type III Wald test was performed. Results showed no significant interaction effect ($\chi^2(2)=3.96$, $p=.138$), indicating that active lever entrances per meter did not differ between sexes as a function of session type and abstinence.

4.7. No effect of sex x levers on entrances per meter percent difference from Noncontingent 01

To determine if a sex x levers interaction had an effect on entrances per meter perfect difference from Noncontingent 01, a Type III Wald test was performed. Results showed no significant interaction effect ($\chi^2(1)=0.05$, $p=1$), indicating that entrances per

meter percent difference from Noncontingent 01 did not differ between sexes towards either the active or inactive lever during Noncontingent sessions.

4.8. No effect of sex x pre-lever activity group on cocaine infusions across Short Access

To determine if a sex x pre-lever activity group interaction had an effect on cocaine infusions during Short Access, a Type III Wald test was performed. Results showed no significant interaction between sex and pre-lever activity groups ($\chi^2(9)=15.61$, $p=.438$). These results show that cocaine intake did not differ between sexes as a function of pre-lever activity group across Short Access sessions.

4.9. An effect of sex x pre-lever activity group observed on cocaine infusions across Long Access

To determine if a sex x pre-lever activity group interaction had an effect on cocaine infusions during Long Access, a Type III Wald test was performed. Results showed a significant interaction between sex x pre-lever activity group x session number ($F(13,467.25)=2.34$, $p=.023$). Post hoc analyses revealed Low Pre-Lever Activity females to have greater cocaine infusions than Low Pre-Lever Activity males from Long Access 04 to Long Access 09, excluding Long Access 07, suggesting that females are more sensitive to escalation of cocaine intake (**Fig. S2**). These results align with previous reports that females escalate in use before males (de Guglielmo et al., 2024).

Supplementary Figure 2. Low Pre-Lever Activity Female rats receive greater cocaine infusions than Low Pre-Lever Activity Male rats during Long Access.

Cocaine infusions across Short and Long Access by sex and pre-lever activity group.

Within Low Pre-Lever Activity group: *$p<.05$, **$p<.01$, ***$p<.001$ vs Females.

References

1. Hartig, F. (2024). *DHARMa: Residual diagnostics for hierarchical (multi-level / mixed) regression models* (Version 0.4.7) [R package]. CRAN. https://doi.org/10.32614/CRAN.package.DHARMa.

2. Bates D, Mächler M, Bolker B, Walker S (2015). "Fitting Linear Mixed-Effects Models Using lme4." Journal of Statistical Software, 67(1), 1–48. doi:10.18637/jss.v067.i01.

3. Kuznetsova A, Brockhoff PB, Christensen RHB (2017). "lmerTest Package: Tests in Linear Mixed Effects Models." *Journal of Statistical Software*, **82**(13), 1–26. doi:10.18637/jss.v082.i13.

4. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al. (2017). "glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling." *The R Journal*, 9(2), 378–400. doi:10.32614/RJ-2017-066.

5. Luke, S.G. Evaluating significance in linear mixed-effects models in R. Behav Res 49, 1494–1502 (2017). https://doi.org/10.3758/s13428-016-0809-y.

6. Meteyard, L., & Davies, R. A. I. (2020). Best practice guidance for linear mixed-effects models in psychological science. Journal of Memory and Language, 112, 104092. https://doi.org/10.1016/j.jml.2020.104092.

7. Razali, N. M., & Wah, Y. B. (2011). Power comparisons of Shapiro-Wilk, Kolmogorov-Smirnov, Lilliefors, and Anderson-Darling tests. *Journal of Statistical Modeling and Analytics, 2*(1), 21–33.

8. Giordano de Guglielmo, Carrette, L. L. G., Kallupi, M., Brennan, M., Boomhower, B., Maturin, L., et al.. (2024). Large-scale characterization of cocaine addiction-like

behaviors reveals that escalation of intake, aversion-resistant responding, and breaking-points are highly correlated measures of the same construct. *eLife*, *12*, RP90422.

https://doi.org/10.7554/eLife.90422