The past and potential antagonistic effects of fishing ban against nutrient loading control on eutrophication restoration in Lake Hongze

Bo Qin[1,3], Vasilis Dakos[2,3]*, Yongjiu Cai[1], Yu Zhao[1], Xiangdong Yang[1,4], Enlou Zhang[1,4], Rong Wang[1,4,5]*

[1] State Key Laboratory of Lake Science and Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, Nanjing 211135, China.

[2] ISEM, Univ Montpellier, CNRS, EPHE, IRD, 34095 Montpellier, France.

[3] Sorbonne Université, CNRS, IRD, INRAE, Université Paris Est Créteil, Université de Paris Cité, Institute of Ecology and Environmental Sciences of Paris (iEES-Paris), Paris, France.

[4] University of Chinese Academy of Sciences, Beijing 100049, P. R. China.

[5] The Fuxianhu Station of Plateau Deep Lake Field Scientific Observation and Research, Yunnan, Yuxi 653100, China.

* Correspondence:

Vasilis Dakos (vasilis.dakos@umontpellier.fr) and Rong Wang (rwang@niglas.ac.cn)

Abstract :

Integrated management strategies are often implemented to achieve comprehensive
environmental restoration; however, the potential antagonistic effects among such measures
are seldom examined. For instance, fishing bans, aimed at enhancing fish biodiversity in
eastern China, may counteract the effectiveness of nutrient control measures in mitigating
eutrophication. This study employs a dual modeling approach to evaluate this hypothesis in
Lake Hongze, the fourth largest shallow lake in China. The PCLake model was calibrated
using observational data from Lake Hongze collected between 2016 and 2020. The
calibrated model was subsequently applied in a hindcast analysis from 2020 to 2024 and a
forecast simulation from 2025 to 2030 to assess the impacts of fishing bans. Combined with
structural equation modeling (SEM) path analysis, the results support the hypothesis that
fishing bans can exhibit antagonistic effects on nutrient loading control by enhancing
top-down regulation relative to bottom-up processes. Specifically, the fishing ban led to
increased fish biomass, phytoplankton abundance, and nutrient levels over the past four
years, while reducing the abundance of submerged macrophytes, zoobenthos, and
zooplankton. Projections further indicate that such antagonistic effects would persist through
2030 if the fishing ban remains in place as planned. Additionally, model forecasts suggest
that even with adjustments to the duration and intensity of fishing bans, antagonistic effects
could be amplified under more stringent nutrient control scenarios. This study provides
valuable insights into maximizing the net benefits of combined environmental management
strategies through a systematic framework that evaluates both the effects and underlying
processes of different interventions.
41
42   **Keywords**: Lake restoration; Nutrient loading control; Fishing bans; PCLake model; Structure
43   Equation Model

44
Highlights:
1. A research framework is provided to reveal the antagonistic effects of different
managements.
2. The hypothesis that fishing ban could have antagonistic effects on nutrient loading control
by strengthening top-down effects is confirmed.
3. Such antagonism of fishing ban would be amplified under a stricter control of nutrient
loading.
52

1 Introduction

Degradations of freshwater lakes, including cultural eutrophication and biodiversity loss, have impeded global sustainable development since last century (Albert et al. 2020, Ho, Michalak and Pahlevan 2019). Major drivers of these degradations include, but are not limited to, enhanced nutrient enrichment, over-exploitation, hydrological modification, and climate change (Sinha, Michalak and Balaji 2017, Jenny et al. 2020). Achieving relevant Sustainable Development Goals (e.g.SDGs 6 and 14) requires concerted efforts from multiple stakeholders, implying that integrated management strategies targeting diverse drivers are essential (Birk et al. 2020, Reed 2008). While the synergistic effects of combined management measures have been widely studied, less attention has been given to potential antagonistic effects—which also represent critical information for policy-making.

The shallow lakes in the eastern plain region of China have experienced severe cultural eutrophication and biodiversity degradation (Qin et al. 2013, Wang et al. 2023, Chen et al. 2022). Controlling external nutrient loading is considered the primary approach for mitigating eutrophication, as it restricts phytoplankton growth and influences higher trophic levels through bottom-up resource limitation (Schindler 2012, Jeppesen et al. 2007, Tigli et al. 2024). In China, policies such as the Action Plan for Water Pollution Prevention and Control (2015) and the River and Lake Chief System (2016) are expected to facilitate the restoration of eutrophic lakes as regulations become stricter and more widespread (Yu et al. 2020, Hu et al. 2024). Concurrently, an unprecedented ten-year fishing ban has been implemented since 2020 in the Yangtze River and its connected lakes—one of the most stringent biodiversity conservation initiatives globally (Wang et al. 2022a). This year-round prohibition comprehensively restricts all harvest of natural fishery resources in designated areas. Classic biomanipulation techniques, such as removing planktivorous and benthivorous fish and introducing piscivorous species, are often applied alongside nutrient control in eutrophic lakes (Søndergaard et al. 2007, Søndergaard et al. 2008). These measures enhance zooplankton biomass by reducing grazing pressure, thereby suppressing phytoplankton proliferation. However, subtropical lakes in eastern China support more omnivorous and planktivorous-benthivorous fish and fewer piscivorous species compared to temperate systems(Jeppesen et al. 2010, Mao et al. 2020). Consequently, the fishing ban may produce effects contrary to traditional biomanipulation: by increasing planktivorous fish stocks, it could intensify predation on zooplankton and indirectly promote phytoplankton growth (Figure 1B). Although recent studies report positive effects in water quality, fish abundance, and network stability in certain lakes after fishing ban (Huang et al. 2025a, Yu et al. 2020, Zhang et al. 2024, Yao et al. 2024), the existence and magnitude of such antagonistic effects remain uncertain.

The process-based lake ecosystem model, a widely-used tool for evaluating the efficacy of restoration measures (Janssen et al. 2015), can help assess such antagonistic effects resulting from fishing bans. As a model specifically designed for shallow lakes, PCLake represents a fully mixed water body and incorporates key biological components such as phytoplankton, zooplankton, and three functional groups of fish (Janse 2005) (Figure S1). It

has been applied globally to study the causes and characteristics of eutrophication (Kuiper
2011, Janssen et al. 2021), the effectiveness of restoration measures (Qin et al. 2022, Hilt et
al. 2018),and the potential impacts of climate change (Qin et al. 2025, Rolighed et al. 2016)
across case-specific, regional, and global scales. Interactions between ecological processes
should be taken into account in order to understand the balance of feedbacks (Dakos et al.
2019). Although PCLake is built upon a rigorous theoretical framework, the complexity of
biological processes often complicates direct and concise representation of trophic cascades.
In contrast, structural equation modeling (SEM), grounded in a strong theoretical foundation,
allows for the estimation of trophic cascade strength and other causal networks through a
series of multiple regression analyses (Grace 2006, Su et al. 2019). Thus, integrating both
approaches enables a comprehensive assessment of both the outcomes and underlying
processes associated with combined management strategies.
108
This study employed two modeling approaches to evaluate the potential antagonistic effects
between nutrient control measures and fishing bans in Lake Hongze, the fourth largest
shallow lake in China (Figure 1). As a key storage reservoir for the Eastern Route of the
South-North Water Diversion Project, Lake Hongze plays a vital role in climate regulation,
flood control, and water resource management, and is currently in a critical phase of
restoration (Zhang et al. 2024). Monthly observational data from 2016 to 2020, collected prior
to the implementation of the fishing ban, were used to calibrate the PCLake model. A hindcast
simulation based on the calibrated model was then conducted to assess the ecological
impacts of the fishing ban, the trophic cascades of which were further examined using
structural equation modeling (SEM). Finally, a forecast analysis was performed to identify
strategies for maximizing management effectiveness. The results confirm the hypothesized
antagonism through a novel research framework, and the implications and limitations of these
findings for lake conservation and management are discussed.
122

2 Materials and methods

2.1 Study Area

Lake Hongze (33°06'-33°40'N, 118°10'-118°52'E), the fourth largest freshwater lake in China,
has a surface area of 1793 km^2 and a drainage basin of 160000 km^2. With a mean depth of
1.7 m and a water volume of 3.0 billion m^3 (Cai et al. 2016a), the lake exhibits a rapid
hydrological cycle and a short water retention time of approximately 35 days. The region
experiences a monsoon-influenced climate, with a mean annual precipitation of 925.5 mm, an
average wind speed of 3.7 m·s⁻¹, and mean temperatures of 2.5°C in January and 28.8°C in
July. Major inflows enter from the west and exit toward the east (Figure 1A; Table S1). The
Huaihe River, the primary inflow, accounts for about 70% of total inflow and more than half of
the external nutrient load. The main outflow occurs through the Sanhe River, which connects
Lake Hongze to the Yangtze River and conveys 60%–70% of the total discharge.
135
Since the 1980s, Lake Hongze has undergone eutrophication following marked declines in
water quality and aquatic biodiversity (Li and Pu 2003). It was listed under the Regulations of
Jiangsu Province on Protection of Lakes in 2004 to mitigate pollution and manage resource

use. Between 2017 and 2019, nine algal blooms were recorded, with affected areas
expanding from 16 km² to 36 km² (Cui et al. 2021). Spatial heterogeneity characterizes the
lake's water quality and ecological conditions (Figure 1A, S2). Zone 1 (Chengzi District) in the
north has restricted hydrologic exchange, while Zone 3 (Lihewa District) in the west functions
as a wetland reserve with shallow water. These regions support relatively abundant
populations of aquatic macrophytes, zooplankton, and macroinvertebrates (Li et al. 2019, Ni
et al. 2023, Cai et al. 2016b). In contrast, Zone 2 is an open-water area largely devoid of
submerged vegetation and exhibits higher nutrient levels, strongly influenced by inputs from
the Huaihe River.

148
Figure 1. Location of study area, hypothesis and modelling approach of this study. (A) Zones 1 and
3 exhibit relatively high vegetation coverage, whereas Zone 2 is an open water area largely devoid
of aquatic vegetation. (B) The conceptual diagram illustrates the hypothesis that fishing bans and
external nutrient loading controls may exert antagonistic effects on phytoplankton dynamics, as
they operate through distinct ecological mechanisms. (C) Model parameter calibration, hindcast
simulation, and forecast analysis will be conducted across different temporal phases. A
subsequent structural equation modeling (SEM) path analysis will be performed following the
hindcast to elucidate causal pathways.

157
2.2 Sampling and Measurement of Lake Variables

Monthly field investigations were conducted from January 2016 to December 2019 at ten
sampling sites distributed across three zones (Figure 1A). These sites are characterized by
very shallow conditions, with annual average water depths ranging between 1.9 m and 2.9 m.
At each site, Secchi depth (SD) was measured in situ, and dissolved oxygen was determined
using a YSI 6600 V2-4 Multi-Parameter Water Quality Sonde. Vertically integrated water
samples were collected using acid-cleaned 5 L plastic containers, stored under cool and
shaded conditions, and transported to the laboratory for subsequent analysis. Total nitrogen,
total phosphorus, and chlorophyll-a concentrations were measured following standard
methods described in APHA (2012). Zoobenthos samples were collected at all sites, where
the substratum predominantly consisted of muddy sediment. Sampling was performed using
a 0.05 m² modified Peterson grab, with three grabs combined to form a composite sample.
The collected material was sieved in situ through a 250 µm mesh sieve. Retained residues
were stored in a low-temperature incubator and transported to the laboratory on the same day.

There, samples were sorted on a white tray, and organisms were preserved in 7% buffered
formalin. Specimens were identified to the lowest practicable taxonomic level, enumerated,
blotted dry, and weighed for wet mass determination using an electronic balance (Sartorius
BS⬚124, precision: 0.1 mg). Taxonomic identification and classification of zoobenthos
followed the references of Liu et al. (1979) and Tang (2006).
177

2.3 Lake Ecosystem Model Configuration and Calibration

To achieve adequate model performance for Lake Hongze during 2016–2024, key model
inputs were adjusted. Fixed inputs included a mean water depth of 1.9 m, maximum depth of
4.5 m, wind fetch of 60,600 m (Janssen et al. 2014), lightly clayish soil (Janse 2005) and no
wetland zone. Time-series inputs of PCLake model, including wind speed ($m·s^{-1}$), solar
radiation ($J·m^{-2}·d^{-1}$), water temperature (°C), water inflow, evaporation ($mm·d^{-1}$), outflow
($mm·d^{-1}$), and nutrient loading ($g·m^{-2}·d^{-1}$), were derived from multiple sources: (1). Monthly
wind speed, precipitation, air temperature, and radiation from 2016 to 2024 were obtained
from National Meteorological Information Center (https://m.data.cma.cn/data) at a

meteorological station (Xuyi, 32.98°N,118.52°E, 40.8 m a.s.l) located 3 km from the lake. (2)

Water temperature from the Huaihe River (2016–2019) was used directly; an empirical
air–water temperature relationship extended these data to 2020–2024. (3). Monthly river
inflow (2016–2019) was investigated (Table S1), with the Huaihe River contributing 88.5% of
total inflow. Inflow for 2020–2024 was estimated using precipitation and historical discharge.
Daily water level (Jiangba station) ranged from 11.5 m to 13.7 m. Mean daily evaporation was
set to 2.5 mm (variance: 1.3 mm). Outflow was computed from total inflow, water level
change, and evaporation. (4). Monthly nutrient inputs (2016–2021) from the Huaihe River
averaged 4×10⬚ kg N and 2×10⬚ kg P (Meng. F. J. et al. 2023), showing a general decline,
especially after December 2018. A ratio of 0.8 was applied to estimate total nutrient loading
based on Huaihe inputs (Ji et al. 2014).
198

Two modifications were made to the PCLake model to better represent Lake Hongze's
ecosystem and address the research objectives. First, while the original model includes three
functional fish groups—zooplanktivorous juvenile whitefish, benthivorous adult whitefish, and
piscivorous fish—with benthivorous adults (sDFiAd) dominating (~80% in both clear and
turbid states), we introduced an additional group: zooplanktivorous adult whitefish. This
adjustment reflects the prevalence of zooplanktivorous and omnivorous fish in local catches.
All equations for this group were adapted from those of benthivorous adults, except for the
food limitation function, which was derived from zooplanktivorous juveniles. Second, the
original model represents fishing intesity as a first-order rate constant, which is inadequate for
simulating variable fishing pressures. To improve temporal realism, we converted fishery
intensities into time-series inputs based on annual fish yield and composition data (Mao et al.
2019), allowing dynamic simulation of fishing effort over time.
211

Default model parameters were initially derived from observations in over 40 lakes (Janse et
al. 2010). Both original and newly introduced parameters were adjusted to enhance the
performance of the modified PCLake model for Lake Hongze. Calibration was performed over

a four-year period (2016–2020) using monthly observed data for total nitrogen, total
phosphorus, dissolved oxygen, Secchi depth, chlorophyll‑a, and zoobenthos, along with
annual records of macrophyte coverage and fish harvest from published sources. Although
the vegetation presence frequency (VPF) ranges between 0.24 and 0.37 (Li et al. 2019),
submerged macrophytes remain scarce, with a peak coverage of only 11.80 km²
(approximately 0.8% of the lake area) between 2013 and 2023 (Huang et al. 2025b). Fish
catch data from 2016 to 2019 were provided by the Jiangsu Province Hongze Lake Fisheries
Management Committee, indicating that zooplanktivorous and omnivorous fish collectively
accounted for over 75% of the total catch (Lin. et al. 2013, Mao et al. 2019).
224

Simulations were conducted using a C++ compiled version of PCLake called via GRIND for
MATLAB (Mooij et al. 2014). Each simulation began from a default turbid state and was run
for a 20-year "burn-in" period under 2016 input conditions to achieve seasonal equilibrium
before calibration. Key parameters were manually calibrated based on model outputs. The
coefficient of determination (R^2) and root-mean-square error (RMSE) were employed to
evaluate model performance across whole-lake averages (Bennett et al. 2013). A higher R^2
indicates better predictive accuracy, while a lower RMSE reflects smaller deviations between
simulated and observed values. Given the absence of abrupt ecological shifts during the
calibration period, the objective was to achieve simulations where most variables exhibited
similar ranges and seasonal dynamics to the observations.
235

2.4 Hindcast and SEM analysis

Following calibration, the model simulation was extended from 2020 to 2024 to conduct a
hindcast analysis. Due to the unavailability of nutrient loading data for this period, values from
2020 were applied to the subsequent four years. Fishing intensity was set to zero to reflect
the implementation of the fishing ban (FB scenario). For comparative purposes, a
no-fishing-ban scenario (no FB) was also simulated using the average fishing intensity from
2016 to 2020. Monthly and annual averages were computed for eight model outputs.
Dissolved oxygen was excluded due to its limited influence in the food web of this large,
shallow lake, whereas zooplankton—a key group linking fish and phytoplankton
dynamics—was included. Although additional time-series observations were unavailable,
further validation was supported through comparison with previously published findings in the
discussion.
248

Structural equation modeling (SEM) was employed to analyze trophic cascades under
scenarios with and without the fishing ban. A conceptual path model was developed based on
the PCLake framework (Figure S1), incorporating seven variables; total nitrogen was
excluded due to its correlation with total phosphorus. Monthly outputs from both simulations
were used to ensure sufficient data. The model was fitted to the covariance structure of the
data, beginning with a full model that included all possible pathways, which was subsequently
refined by removing nonsignificant paths. Standardized path coefficients were used to
quantify the strength and direction of direct relationships between consecutive trophic levels.
The overall cascade strength was evaluated through the indirect effect from top to bottom of
the food web, derived from the product of sequential path coefficients. Model fit was

assessed using the chi-square/degrees of freedom ratio (CMIN/df) (Karin Schermelleh-Engel
and Helfried Moosbrugger 2003), to assess how well the proposed network structure explains
the simulated data from PCLake. All SEM analyses were performed in AMOS 21.0 (IBM
2012).

2.5 Forecast

According to the Environmental Quality Standards for Surface Water in China, the total
phosphorus level in Lake Hongze in 2023—though the lowest in recent years—still exceeded
the drinking water threshold by 46% (Zhang et al. 2024). In addition, while fish stocks
increased significantly, imbalanced fish community structure was also reported. Above facts
imply that continued measures are further needed for Lake Hongze in the near future. To
evaluate potential management outcomes, we designed eight scenarios combining different
levels of external nutrient loading and fishing intensity to simulate conditions from 2025 to
2030. Two nutrient loading scenarios were defined: the highest level over the past five years
(NP-, control) and the lowest level (NP--, approximately half of NP-). Four fishing scenarios
were established: continued full fishing ban (F0, control); a biennial ban alternating between
prohibition and permission (F0/1); full resumption of fishing at the 2020–2024 average
intensity (F1); and continuous fishing at half of F1 intensity (F0.5). Other model inputs used
2020 data as boundary conditions. As the ten-year fishing ban is scheduled to end in 2030,
we focused on the relative changes between 2030 and 2025. Differences in annual model
outputs were calculated and compared against the control scenario to standardize units
across variables.

3 Results

3.1 Lake States and PCLake Model Performance

The average nutrient concentrations in the lake water from 2016 to 2020 were 1.72 $g \cdot m^{-3}$ for
total nitrogen (TN) and 0.08 $g \cdot m^{-3}$ for total phosphorus (TP) (Figure S2). While TP exhibited a
slight increase, TN declined significantly after 2017. The mean Secchi depth was 0.35 m and
dissolved oxygen averaged 0.35 $g \cdot m^{-3}$, both remaining relatively stable throughout the period.
The chlorophyll-a concentration averaged 8 $mg \cdot m^{-3}$, with occasional peaks observed. At the
class level, Oligochaeta dominated in Zone 1, whereas Polychaeta were more abundant in
Zone 2 (Figure S3). Since the PCLake model does not explicitly include large zoobenthos,
the total dry weight of the five major classes was used to represent zoobenthos biomass.

The performance of the calibrated PCLake model for Lake Hongze was assessed using two
statistical indices based on six monthly and two annual observational variables (Figure 2).
The simulated trajectories (solid lines) generally captured both the range and seasonal
dynamics of the whole-lake observations (dots). The coefficient of determination (R^2) ranged
from 0.3 (zoobenthos) to 0.8 (dissolved oxygen), with an average of 0.53. The
root-mean-square error (RMSE) varied between 0 (fish catch) and 3.7 (chlorophyll⬚a),
averaging 0.73. Although the annual observations (macrophyte coverage and fish catch)
were limited in number, their simulations were deemed acceptable due to agreement in mean

values. Monthly water quality variables were better reproduced than biological variables
(phytoplankton and zoobenthos). The modified and calibrated model exhibited generally
higher R^2 and lower RMSE compared to the original model (Figure S4, Table S2), particularly
for biological components. Overall, the model demonstrated acceptable performance during
the 2016–2020 calibration period and was considered suitable for further analysis.

Figure 2. The result of PCLake model calibration between 2016 and 2020. Monthly observational
data from Lake Hongze comprised total nitrogen, total phosphorus, dissolved oxygen, Secchi
depth, chlorophyll-a, and zoobenthos, while macrophyte coverage and fish harvest were
annually. Data from different lake zones are presented; however, the coefficient of determination
(R²) and root mean square error (RMSE) were calculated based on whole-lake averages and
corresponding model simulations.
313

3.2 Hindcast on Fishing Ban and Trophic Cascade

The calibrated PCLake model was applied to conduct a hindcast analysis for the period
2020–2024 (Figure 3A). Under the fishing ban scenario (FB), total fish biomass and
chlorophyll-a showed an increasing trend, while macrophyte coverage declined slightly;
other variables remained largely stable. Compared to the no-fishing-ban scenario (no FB),
the FB scenario resulted in higher levels of total nitrogen, total phosphorus, chlorophyll-a,
and total fish biomass, but lower values of Secchi depth, macrophyte coverage, zoobenthos,
and zooplankton.
322

Monthly hindcast outputs were further analyzed using structural equation modeling (SEM) to
quantify trophic interactions (Figure 3B; Table S3). Both SEM models exhibited acceptable fit
(χ^2/df < 5). Under the fishing ban, the path coefficients ranged from –0.91 to 0.6 (average:
–0.38), with dominant negative relationships (red lines) across most variables except total
phosphorus and macrophyte coverage. Strong top-down effects were evident through
significant negative paths from fish to zooplankton (–0.62) and zoobenthos (–0.58). In
contrast, without the fishing ban, positive pathways (black lines) prevailed among total
phosphorus, macrophyte coverage, Secchi depth, phytoplankton, and zooplankton.
Bottom-up effects were indicated by positive sequential coefficients from total phosphorus to

phytoplankton (0.42) and phytoplankton to zooplankton (0.63).

333
Figure 3. The result of PCLake hindcast and SEM path analysis between 2020 and 2024. (A) "FB"
denotes the control scenario with a full fishing ban (fishing intensity = 0), while "no FB" represents
the scenario without fishing restrictions. Results are presented both monthly and annually. (B)
Two SEM analyses were performed using monthly data from seven variables. Symbols
representing these variables were sourced from https://ian.umces.edu/media-library/. Both SEM
models exhibited acceptable fit according to the chi-square/degrees of freedom ratio (χ^2/df), and
revealed contrasting dominant trophic cascades under fishing ban (left) and no ban (right)
conditions. Red lines indicate negative relationships, black lines positive relationships, with line
width corresponding to effect strength. Non-significant paths are shown as dashed grey lines.

343
3.3 Forecast Under Potential Managements

To evaluate potential regulatory effects from 2025 to 2030, a scenario analysis was
conducted using the Hongze Lake PCLake model. Relative changes in annual outputs by
2030 reveal distinct response patterns across scenarios compared to the control (NP- and
F0). Variables in group A show greater sensitivity to nutrient control (differences between NP-
and NP--) than to fishing intensity. In contrast, variables in group B exhibit a vertical gradient
across fishing scenarios (F0, F0/1, F0.5, F1), indicating stronger influence from fishing
pressure. Chlorophyll-a displays discernible variation along both nutrient and fishing
gradients.

353
Stricter nutrient control significantly reduced nutrient concentrations and increased oxygen
levels, whereas fishing bans had minimal impact on these variables. Chlorophyll-a also
decreased under lower external loading. Compared to continued fishing ban (F0), scenario
F1 (no ban) resulted in markedly lower total fish biomass and chlorophyll-a, but higher

zooplankton, zoobenthos, and macrophyte levels. Intermediate fishing scenarios (F0/1 and
F0.5) led to moderate effects, with F0.5 yielding slightly lower fish stock and chlorophyll-a, yet
higher macrophyte coverage than F0/1. Moreover, differences between F0 and F1 were
amplified under stricter nutrient control for chlorophyll-a and all group B variables except for
zooplankton.

Figure 4. The result of PCLake forecast between 2025 and 2030. The eight scenarios resulted
from combining two external nutrient loading conditions (controlled, NP-; strictly controlled, NP--)
with four fishing regimes (continuous ban, F0; biennial ban, F0/1; reduced effort, F0.5; no ban, F1).
The color bar indicates the relative change compared to the control scenario (NP- and F0), with
values above 1 denoting an increase relative to control. Variables are grouped into (A) and (B)
based on similar response patterns across scenarios.

4 Discussion

4.1 The Past and Potential antagonistic Effects of Nutrient Control and Fishing Ban

Despite the well-acknowledged negative effect of zooplankivorous fish stocking on water
quality, such impact remains neglected in the combination of fishing ban and nutrient control.
This neglection was mainly due to the difficulties in conducting process evaluations and the
separation of the effectiveness evaluation system. Here, we sought to fill in the gap by
combining a process-based modeling and path analysis approach in a lake with consistent
observations. We confirmed the hypotheses that fishing ban can trigger enhanced top-down
control to compromise the effectiveness of nutrient loading control on lake water quality and
such antagonism would continue until 2030 if fishing ban will be implemented as planned.
Thus, we anticipate that, in the context of comprehensive management, eutrophication
restoration would slow down due to more enhanced trophic cascade from fish.

One advantage of combining the PCLake model and SEM analysis is to disentangle both the
independent contributions and the indirect causal-effect pathways of combined
managements in lake foodweb.

First, the model revealed divergent trends in lake variables under fishing ban (FB) and
non-ban (no FB) conditions (Figure 3). From 2020 to 2024, the FB scenario was
characterized by increases in fish biomass and chlorophyll a concentration. These findings
align with reported rises in fish biodiversity and phytoplankton density by Zhang et al. (2024).
The expansion of fish stocks resulted from the cessation of fishing, particularly under
sufficient availability of zooplankton and zoobenthos as food sources. This also implies that
the likelihood of fishing bans leading to an increase in predatory fish populations and a
decrease in other fish populations is low. Although macrophytes could offer refuge for
zooplankton against predation, their limited coverage—remaining consistently low—rendered
this effect negligible. Consequently, zooplankton declined slightly, reducing grazing pressure
and contributing to phytoplankton increase. Even under no FB, a modest rise in fish biomass
was simulated, attributable to the preexisting seasonal fishing ban (February–June annually
since 2012) (Liu 2015), which had already lowered baseline fishing pressure. Continued
implementation of such partial bans could enable gradual fish recovery, as reflected in the
hindcast. Zoobenthos also increased, likely due to greater stocking of zooplanktivorous over
benthivorous fish. Notably, studies such as Ren et al. (2022) indicate that benthic fish can
promote sediment nutrient release, stimulating phytoplankton and periphyton growth, which
in turn shades and suppresses submerged macrophytes. In our hindcast, Secchi depth
remained largely stable despite correlations with other variables, suggesting that factors such
as hydrodynamics and mining activities—exacerbated by the lake's shallow depth and short

retention time (Li et al. 2019)—may exert stronger influence on light penetration and

macrophyte development.

Second, our simulations demonstrate that fishing bans can intensify top-down control,
thereby undermining the effectiveness of nutrient reduction measures in improving water
quality—an effect projected to persist until 2030 if the ban continues as scheduled (Figure 4).
Phytoplankton, positioned at the interface of bottom-up nutrient effects and top-down
predation pressure (Figure 3), exhibits complex responses under combined management
regimes (Figure 4). Lower phytoplankton levels under non-ban conditions may be attributed
to nutrient removal through fishery activity, which enhances bottom-up control. Under fishing
bans, however, phytoplankton abundance closely tracked fish biomass, as increased
predation reduced zooplankton grazing pressure. Our results indicate that stringent nutrient
control (NP--) combined with maintained fishing activity (F1) could optimize water quality by
achieving the lowest nutrient and phytoplankton levels alongside the highest transparency,
macrophyte coverage, and zoobenthos biomass (Figure 4). However, this scenario also
resulted in the lowest fish stock, illustrating a classic ecosystem service trade-off (Rodríguez
et al. 2006, Xu et al. 2017). Partial fishing restrictions (e.g., biennial F0/1 or reduced effort
F0.5) also attenuated the benefits of nutrient control, reflecting intermediate ecological

outcomes. This may result from rapid fish stock expansion during initial ban years, leading to
structural surplus (Zhang et al. 2024).
428

Generally, a 40%–75% reduction in nutrient inputs can result in a regime shift from a turbid,
algal-dominated state to a clear, macrophyte-dominated state (Søndergaard, Jensen and
Jeppesen 2005, Me et al. 2018). Although Lake Hongze is nearing this critical threshold
(Zhang et al. 2024) and both external and internal nutrient levels have declined in recent
years, no significant macrophyte recovery has been observed yet. The restoration of Lake
Hongze, which is in a critical phase of regime shift, would slow down due to more enhanced
trophic cascade from fish. With adjustments to the duration and intensity of fishing bans,
antagonistic effects could be amplified under more stringent nutrient control scenarios (Figure
4).Therefore, complementary strategies—such as reconnecting migratory pathways,
restoring habitats, and targeted species stocking (Wang et al. 2022b, Mao et al. 2020, Feng

et al. 2023)—should be integrated to simultaneously support biodiversity and eutrophication

mitigation.
441

4.2 Evaluations and Limitations of the Modeling

Based on model performance during calibration (2016–2020) and hindcasting (2020–2024),
we conclude that the boundary conditions and parameter values of the PCLake model were
appropriately configured. The calibrated simulation captured both the magnitude (RMSE) and
seasonal dynamics (R^2) of observed variables comparably to other studies (Figure 2; Table
S2). Relatively low R^2 values for zoobenthos and chlorophyll-a may be attributed to
insufficient zooplankton data, leading to uncertain estimates of grazing pressure on
phytoplankton and zoobenthos food availability. Although biological variables showed lower
fit than water quality parameters, the modified model exhibited improved performance over
the original version (Figure S4; Table S2). Hindcast simulations under the fishing ban
(2020–2024) aligned with trends reported in independent studies (Z. H. Huang et al., 2025),
further supporting model validity. Additionally, a 4.5% increase in fish species richness
(Huang et al. 2025b) and a projected 30% recovery of aquatic animal resources by 2031
(Wang et al. 2022a) corroborate the model's ecological relevance. Structural equation
modeling (SEM) was employed for trophic cascade analysis. Although a χ^2/df ratio below 3 is
ideal, values under 5 are considered acceptable (Karin Schermelleh-Engel and Helfried
Moosbrugger 2003); both models in this study (Figure 3B) fell within this range. Potential
sources of uncertainty include simplification of model structure, assumption of fixed pathways
and directions—which may vary temporally—and possible nonlinear relationships among
variables not captured in the SEM.
462

The limitations of this study primarily concern the absence of uncertainty analysis and the
generalizability of the results. First, uncertainties stemming from the PCLake model structure
and the selection of forcing functions for the control scenarios were not quantitatively
evaluated. Although originally designed for temperate lakes (Janse 2005), the model for Lake
Hongze was adapted to represent omnivorous fish indirectly by incorporating
zooplanktivorous fish and adjusting their balance with benthivorous species. Nevertheless,

the explicit absence of true omnivorous fish may reduce the reliability of simulations,
particularly for biological variables. Second, while the choice of forcing functions—such as
average nutrient loading from 2020 to 2024—likely preserves relative differences between
scenarios, it may affect the absolute values of predicted variables. Third, the mismatch in
temporal scales between lower and higher trophic levels in process-based modeling could
influence the simulated responses of certain functional groups. Additionally, although a
mutually antagonistic effect between fishing bans and nutrient reduction was identified in
Lake Hongze, synergistic outcomes might occur in lakes dominated by piscivorous fish.
477

478 4.4 Implications

Our findings hold relevance for a broad range of lakes globally. First, shallow freshwater lakes
with food web structures comparable to Lake Hongze are widespread. Many of these
systems experience severe eutrophication and water quality degradation, primarily due to
pollution and resource overexploitation. The quantitative insights provided here regarding the
contributions of different management strategies can support more informed decision-making.
Although fishing bans are not traditionally classified as biomanipulation measures, they
represent a direct policy approach to mitigating biodiversity loss by addressing key
anthropogenic pressures (Wang et al. 2022a). Such targeted interventions, aimed at
eliminating specific local stressors, are increasingly relevant in the context of achieving
Sustainable Development Goals (SDGs). While the specific antagonistic effects identified
here may be most applicable to lakes resembling Lake Hongze in physical and biological
characteristics, the proposed research framework offers a transferable methodology for
optimizing net benefits from multiple environmental policies. Importantly, the integration of
process-based modeling and path analysis enables simultaneous evaluation of management
outcomes and underlying mechanisms—an approach especially critical under future climate
change.
495

Our modeling approach reveals the presence and mechanisms of antagonistic effects
between external nutrient control and fishing bans in the restoration of eutrophic freshwater
lakes. We project that this antagonism will persist until 2030 if the fishing ban continues as
planned, and may intensify under stricter nutrient reduction. Within integrated management
frameworks, eutrophication mitigation is likely to slow due to strengthened trophic cascades
originating from fish populations. These results highlight the need for complementary
strategies that simultaneously support biodiversity recovery and water quality improvement.
Furthermore, management combinations should be implemented within systematic
frameworks—featuring quantifiable feedback mechanisms—to address inherent trade-offs
among ecosystem services. This study also demonstrates how integrating process-based
modeling with path analysis can help quantify the contributions and underlying processes of
management measures with complex interactions. Future improvements in model precision
are necessary to enhance the reliability of strategies aimed at maximizing net benefits from
multi-objective environmental policies.
510

511 **Acknowledgements**

512 This work was supported by the Joint Funds of the National Natural Science Foundation of
513 China (No. U24A20639, 42171119, 42507663), and the State Key Laboratory of Lake and
514 Watershed Science for Water Security (NKL2023-ZD02).

References

Albert, J. S., G. Destouni, S. M. Duke-Sylvester, A. E. Magurran, T. Oberdorff, R. E. Reis, K. O.

Winemiller & W. J. Ripple (2020) Scientists' warning to humanity on the freshwater

biodiversity crisis. *Ambio,* 50, 85-94.

APHA (2012) Standard Methods for the Examination of Water and Wastewater. *American Public*

Health Association, Washington, DC, 22nd edition.

Bennett, N. D., B. F. W. Croke, G. Guariso, J. H. A. Guillaume, S. H. Hamilton, A. J. Jakeman, S.

Marsili-Libelli, L. T. H. Newham, J. P. Norton, C. Perrin, S. A. Pierce, B. Robson, R. Seppelt, A.

A. Voinov, B. D. Fath & V. Andreassian (2013) Characterising performance of environmental

models. *Environmental Modelling & Software,* 40, 1-20.

Birk, S., D. Chapman, L. Carvalho, B. M. Spears, H. E. Andersen, C. Argillier, S. Auer, A.

Baattrup-Pedersen, L. Banin, M. Beklioğlu, E. Bondar-Kunze, A. Borja, P. Branco, T. Bucak, A.

D. Buijse, A. C. Cardoso, R.-M. Couture, F. Cremona, D. de Zwart, C. K. Feld, M. T. Ferreira,

H. Feuchtmayr, M. O. Gessner, A. Gieswein, L. Globevnik, D. Graeber, W. Graf, C.

Gutiérrez-Cánovas, J. Hanganu, U. Işkın, M. Järvinen, E. Jeppesen, N. Kotamäki, M. Kuijper,

J. U. Lemm, S. Lu, A. L. Solheim, U. Mischke, S. J. Moe, P. Nõges, T. Nõges, S. J. Ormerod, Y.

Panagopoulos, G. Phillips, L. Posthuma, S. Pouso, C. Prudhomme, K. Rankinen, J. J.

Rasmussen, J. Richardson, A. Sagouis, J. M. Santos, R. B. Schäfer, R. Schinegger, S.

Schmutz, S. C. Schneider, L. Schülting, P. Segurado, K. Stefanidis, B. Sures, S. J. Thackeray,

J. Turunen, M. C. Uyarra, M. Venohr, P. C. von der Ohe, N. Willby & D. Hering (2020) Impacts

of multiple stressors on freshwater biota across spatial scales and ecosystems. *Nature*

Ecology & Evolution, 4, 1060-1068.

Cai, Y., Y. Zhang, Z. Wu, Y. Chen, J. Xu & Z. Gong (2016a) Composition, diversity, and environmental

correlates of benthic macroinvertebrate communities in the five largest freshwater lakes of

China. *Hydrobiologia,* 788, 85-98.

Cai, Y. J., Y. J. Lu, J. S. Liu, X. L. Dai, H. Xu, Y. Lu & Z. J. Gong (2016b) Macrozoobenthic community

structure in a large shallow lake: Disentangling the effect of eutrophication and wind-wave

disturbance. *Limnologica,* 59, 1-9.

Chen, X., J. Liang, L. Zeng, Y. Cao & M. A. Stevenson (2022) Heterogeneity in diatom diversity

response to decadal scale eutrophication in floodplain lakes of the middle Yangtze reaches.

Journal of Environmental Management, 322.

Cui, J. Y., R. Guo, X. W. Song, Y. Zhang, C. Chen, X. Lv, Y., & Y. Dong, Y., (2021) Spatio-temporal

variations of total nitrogen and total phosphorus in lake and inflow/outflow rivers of Lake

Hongze, 2010-2019 (in Chinese). *Journal of Lake Science,* 33(6): 1727-1741.

Dakos, V., B. Matthews, A. P. Hendry, J. Levine, N. Loeuille, J. Norberg, P. Nosil, M. Scheffer & L. De

Meester (2019) Ecosystem tipping points in an evolving world. *Nat Ecol Evol,* 3, 355-362.

Feng, K., W. Deng, H. Li, Q. Guo, K. Tao, J. Yuan, J. Liu, Z. Li, S. Lek, B. Hugueny & Q. Wang (2023)

Direct and indirect effects of a fishing ban on lacustrine fish community do not result in a full

recovery. *Journal of Applied Ecology,* 60, 2210-2222.

Grace, J. (2006) Structural Equation Modeling Natural Systems. Cambridge University Press.

Hilt, S., M. M. Alirangues Nuñez, E. S. Bakker, I. Blindow, T. A. Davidson, M. Gillefalk, L.-A. Hansson,

J. H. Janse, A. B. G. Janssen, E. Jeppesen, T. Kabus, A. Kelly, J. Köhler, T. L. Lauridsen, W.

M. Mooij, R. Noordhuis, G. Phillips, J. Rücker, H.-H. Schuster, M. Søndergaard, S. Teurlincx,

K. van de Weyer, E. van Donk, A. Waterstraat, N. Willby & C. D. Sayer (2018) Response of

561    Submerged Macrophyte Communities to External and Internal Restoration Measures in North

562      Temperate Shallow Lakes. *Frontiers in Plant Science,* 9.

563  Ho, J. C., A. M. Michalak & N. Pahlevan (2019) Widespread global increase in intense lake

564      phytoplankton blooms since the 1980s. *Nature,* 574, 667-+.

565  Hu, M., R. Ma, K. Xue, Z. Cao, J. Xiong, S. A. Loiselle, M. Shen & X. Hou (2024) Eutrophication

566      evolution of lakes in China: Four decades of observations from space. *Journal of Hazardous*

567      *Materials,* 470.

568  Huang, X., Q. Xu, B. Zhang, C. Kong, L. Fang, X. Gao, L. Sun, L. Li & X. Gong (2025a) An Assessment

569      of the Population Structure and Stock Dynamics of Megalobrama skolkovii During the Early

570      Phase of the Fishing Ban in the Poyang Lake Basin. *Fishes,* 10.

571  Huang, Z. H., H. Z. Liu, Y. Q. Wu, R. H. Ma & X. M. Lu (2025b) The spatiotemporal evolution law of key

572      parameters of water environment in typical lakes in the Jianghuai Basin (in Chinese). *Jiangsu*

573      *Water Resources,* (05): 1-5.

574  IBM, C. (2012) IBM SPSS Statistics for Windows, Version 21.0. *Armonk, NY: IBM Corp.*

575  Janse, J. H. (2005) Model studies on the eutrophication of shallow lakes and ditches. *Wageningen*

576      *University* Wageningen, Ph. D. Dissertation.

577  Janse, J. H., M. Scheffer, L. Lijklema, L. Van Liere, J. S. Sloot & W. M. Mooij (2010) Estimating the

578      critical phosphorus loading of shallow lakes with the ecosystem model PCLake: Sensitivity,

579      calibration and uncertainty. *Ecological Modelling,* 221, 654-665.

580  Janssen, A. B. G., G. B. Arhonditsis, A. Beusen, K. Bolding, L. Bruce, J. Bruggeman, R. M. Couture, A.

581      S. Downing, J. A. Elliott, M. A. Frassl, G. Gal, D. J. Gerla, M. R. Hipsey, F. J. Hu, S. C. Ives, J.

582      H. Janse, E. Jeppesen, K. D. Johnk, D. Kneis, X. Z. Kong, J. J. Kuiper, M. K. Lehmann, C.

Lemmen, D. Ozkundakci, T. Petzoldt, K. Rinke, B. J. Robson, R. Sachse, S. A. Schep, M.

Schmid, H. Scholten, S. Teurlincx, D. Trolle, T. A. Troost, A. A. Van Dam, L. P. A. Van Gerven,

M. Weijerman, S. A. Wells & W. M. Mooij (2015) Exploring, exploiting and evolving diversity of

aquatic ecosystem models: a community perspective. *Aquatic Ecology,* 49, 513-548.

Janssen, A. B. G., B. Droppers, X. Kong, S. Teurlincx, Y. Tong & C. Kroeze (2021) Characterizing 19

thousand Chinese lakes, ponds and reservoirs by morphometric, climate and sediment

characteristics. *Water Research,* 202.

Janssen, A. B. G., S. Teurlincx, S. Q. An, J. H. Janse, H. W. Paerl & W. M. Mooij (2014) Alternative

stable states in large shallow lakes? *Journal of Great Lakes Research,* 40, 813-826.

Jenny, J.-P., O. Anneville, F. Arnaud, Y. Baulaz, D. Bouffard, I. Domaizon, S. A. Bocaniov, N. Chèvre,

M. Dittrich, J.-M. Dorioz, E. S. Dunlop, G. Dur, J. Guillard, T. Guinaldo, S. Jacquet, A.

Jamoneau, Z. Jawed, E. Jeppesen, G. Krantzberg, J. Lenters, B. Leoni, M. Meybeck, V. Nava,

T. Nõges, P. Nõges, M. Patelli, V. Pebbles, M.-E. Perga, S. Rasconi, C. R. Ruetz, L. Rudstam,

N. Salmaso, S. Sapna, D. Straile, O. Tammeorg, M. R. Twiss, D. G. Uzarski, A.-M. Ventelä, W.

F. Vincent, S. W. Wilhelm, S.-Å. Wängberg & G. A. Weyhenmeyer (2020) Scientists' Warning

to Humanity: Rapid degradation of the world's large lakes. *Journal of Great Lakes Research,*

46, 686-702.

Jeppesen, E., M. Meerhoff, K. Holmgren, I. Gonzalez-Bergonzoni, F. Teixeira-de Mello, S. A. J.

Declerck, L. De Meester, M. Sondergaard, T. L. Lauridsen, R. Bjerring, J. M. Conde-Porcuna,

N. Mazzeo, C. Iglesias, M. Reizenstein, H. J. Malmquist, Z. W. Liu, D. Balayla & X. Lazzaro

(2010) Impacts of climate warming on lake fish community structure and potential effects on

ecosystem function. *Hydrobiologia,* 646, 73-90.

Jeppesen, E., M. Sondergaard, M. Meerhoff, T. L. Lauridsen & J. P. Jensen (2007) Shallow lake

restoration by nutrient loading reduction - some recent findings and challenges ahead.

Hydrobiologia, 584, 239-252.

Ji, X. M., L. Wen, M. Zhang & F. Yue (2014) Flux analysis of pollutants entering Hongze Lake (in

Chinese). *Jiangsu Water Resources,* (7): 45-46, 48.

Karin Schermelleh-Engel & Helfried Moosbrugger (2003) Evaluating the Fit of Structural Equation

Models: Tests of Significance and Descriptive Goodness-of-Fit Measures. *Methods of*

Psychological Research, 8(2): 23-74.

Kuiper, J. J. (2011) Making Eco Logic and Models Work: An Integrative Approach to Lake Ecosystem

Modelling. *Wageningen University,* Wageningen, Ph. D. Dissertation.

Li, B. & P. M. Pu (2003) Study of the Evolution Tendency of Water Quality in Huai River Basin and

Hongze Lake (in Chinese). *Resources and Environment in the Yangtze Basin,* 12(1): 67-73.

Li, N., K. Shi, Y. L. Zhang, Z. J. Gong, K. Peng, Y. B. Zhang & Y. Zha (2019) Decline in Transparency

of Lake Hongze from Long-Term MODIS Observations: Possible Causes and Potential

Significance. *Remote Sensing,* 11.

Lin., M. L., T. L. Zhang., S. W. Ye., W. Li., P. Ren., Z. W. Yang., J. S. Liu. & Z. J. Li. (2013) Status of

Fish Resources, Historical Variation and Fisheries management strategies in Hongze Lake (in

Chinese). *ACTA HYDROBIOLOGICA SINICA,* 37(6): 1118-1127.

Liu, X. Z. (2015) Current Situation, Ploblems and Countermeasures of Fishery Resources in Hongze

Lake. *Nanjing Agricultural University,* Nanjing, Ph. D. Dissertation.

Liu, Y. Y., W. Z. Zhang, Y. X. Wang & E. Y. Wang (1979) Economic Fauna of China: Freshwater

Mollusca (in Chinese). *Science Press, Beijing.*

Mao, Z., X. Gu, Y. Cao, M. Zhang, Q. Zeng, H. Chen, R. Shen & E. Jeppesen (2020) The Role of

Top-Down and Bottom-Up Control for Phytoplankton in a Subtropical Shallow Eutrophic Lake:

Evidence Based on Long-Term Monitoring and Modeling. *Ecosystems,* 23, 1449-1463.

Mao, Z. G., X. H. Gu, Z. J. Gong, Q. F. Zeng, H. H. Chen, H. M. Li, S. Y. Zhang & H. Mu (2019) The

structure of fish community and changes of fishery resources in Lake Hongze (in Chinese).

Journal of Lake Sciences, 31(4): 1109-1119.

Me, W., D. P. Hamilton, C. G. McBride, J. M. Abell & B. J. Hicks (2018) Modelling hydrology and water

quality in a mixed land use catchment and eutrophic lake: Effects of nutrient load reductions

and climate change. *Environmental Modelling & Software,* 109, 114-133.

Meng. F. J., Y. X. Gao, G. D. Xu, Y. M. Zhang, T. Chen, X. W. Mao, J. Chen & Y. J. Jin (2023) Analysis

of Nitrogen and Phosphorus Pollution Flux from Huai River into Hongze Lake Based on

LOADSET Model and M-K Test. *The Administration and Technique of Environmental*

Monitoring, 35(3): 17-22.

Mooij, W. M., R. J. Brederveld, J. J. M. de Klein, D. L. DeAngelis, A. S. Downing, M. Faber, D. J. Gerla,

M. R. Hipsey, J. 't Hoen, J. H. Janse, A. B. G. Janssen, M. Jeuken, B. W. Kooi, B. Lischke, T.

Petzoldt, L. Postma, S. A. Schep, H. Scholten, S. Teurlincx, C. Thiange, D. Trolle, A. A. van

Dam, L. P. A. van Gerven, E. H. van Nes & J. J. Kuiper (2014) Serving many at once: How a

database approach can create unity in dynamical ecosystem modelling. *Environmental*

Modelling & Software, 61, 266-273.

Ni, L., H. Li, L. Zhou, J. Shi, Y. Nie, F. Zhao & S. Li (2023) Structural characteristics of zooplankton

communities in Hongze Lake driven by water environmental factors from 2016 to 2020.

Environmental Monitoring and Assessment, 195, 1503.

Qin, B., X. Kong, R. Wang, Y. Zahao & X. Yang (2022) Lake restoration time of Lake Taibai (China): a

case study based on paleolimnology and ecosystem modeling. *J Paleolimnol,* 68, 25–38.

Qin, B., M. Xu, K. Zhu, Y. Zhao, E. Zhang & R. Wang (2025) Climate warming will significantly affect

future restoration and level of ecosystem services in Lake Erhai. *Ecological Modelling,* 503,

111067.

Qin, B. Q., G. Gao, G. W. Zhu, Y. L. Zhang, Y. Z. Song, X. M. Tang, H. Xu & J. M. Deng (2013) Lake

eutrophication and its ecosystem response. *Chinese Science Bulletin,* 58, 961-970.

Reed, M. S. (2008) Stakeholder participation for environmental management: A literature review.

Biological Conservation, 141, 2417-2431.

Ren, W., Z. Wen, Y. Cao, H. Wang, C. Yuan, X. Zhang, L. Ni, P. Xie, T. Cao, K. Li & E. Jeppesen

(2022) Cascading effects of benthic fish impede reinstatement of clear water conditions in

lakes: A mesocosm study. *Journal of Environmental Management,* 301.

Rodríguez, J. P., T. D. Beard, E. M. Bennett, G. S. Cumming, S. J. Cork, J. Agard, A. P. Dobson & G.

D. Peterson (2006) Trade-offs across Space, Time, and Ecosystem Services. *Ecology and*

Society, 11.

Rolighed, J., E. Jeppesen, M. Sondergaard, R. Bjerring, J. H. Janse, W. M. Mooij & D. Trolle (2016)

Climate Change Will Make Recovery from Eutrophication More Difficult in Shallow Danish

Lake Sobygaard. *Water,* 8.

Schindler, D. W. (2012) The dilemma of controlling cultural eutrophication of lakes. *Proceedings of the*

Royal Society B-Biological Sciences, 279, 4322-4333.

Sinha, E., A. M. Michalak & V. Balaji (2017) Eutrophication will increase during the 21st century as a

result of precipitation changes. *Science,* 357, 405-408.

Søndergaard, M., J. P. Jensen & E. Jeppesen (2005) Seasonal response of nutrients to reduced

phosphorus loading in 12 Danish lakes. *Freshwater Biology,* 50, 1605-1615.

Søndergaard, M., E. Jeppesen, T. L. Lauridsen, C. Skov, E. H. Van Nes, R. Roijackers, E. Lammens &

R. Portielje (2007) Lake restoration: successes, failures and long-term effects. *Journal of*

Applied Ecology, 44, 1095-1105.

Søndergaard, M., L. Liboriussen, A. R. Pedersen & E. Jeppesen (2008) Lake Restoration by Fish

Removal: Short- and Long-Term Effects in 36 Danish Lakes. *Ecosystems,* 11, 1291-1305.

Su, H. J., J. Chen, Y. Wu, J. F. Chen, X. C. Guo, Z. B. Yan, D. Tian, J. Y. Fang & P. Xie (2019)

Morphological traits of submerged macrophytes reveal specific positive feedbacks to water

clarity in freshwater ecosystems. *Science of the Total Environment,* 684, 578-586.

Tang, H. Q. (2006) Biosystematic Study on the Chironomid Larvae in China (Diptera: Chironomidae)

(Abstract in English). *Nankai University,* TianJing, Ph. D. Dissertation.

Tigli, M., M. P. Bak, J. H. Janse, M. Strokal & A. B. G. Janssen (2024) The future of algal blooms in

lakes globally is in our hands. *Water Res,* 268, 122533.

Wang, H., P. Wang, C. Xu, Y. Sun, L. Shi, L. Zhou, E. Jeppesen, J. Chen & P. Xie (2022a) Can the

"10-year fishing ban" rescue biodiversity of the Yangtze River? *Innovation (Cambridge*

(Mass.)), 3, 100235.

Wang, R., C. P. Doncaster, W. Zheng, M. Xu, H. Yang, Y. Li, Y. Cai, Y. Zhao, E. Zhang, X. Yang & B.

Qin (2023) High phytoplankton diversity in eutrophic states impedes lake recovery. *Journal of*

Biogeography, 50, 1914-1925.

Wang, R., Y. Han, F. Fan, J. Molinos, J. Xu, K. Wang, D. Wang & Z. Mei (2022b) Need to shift in

river-lake connection scheme under the "ten-year fishing ban" in the Yangtze River, China.

693  *Ecological Indicators,* 143, 109434.

694 Xu, M., X. H. Dong, X. D. Yang, R. Wang, K. Zhang, Y. J. Zhao, T. A. Davidson & E. Jeppesen (2017)

695  Using palaeolimnological data and historical records to assess long-term dynamics of

696  ecosystem services in typical Yangtze shallow lakes (China). *Science of the Total*

697  *Environment,* 584, 791-802.

698 Yao, Y., Z. Mao, X. Gu, Q. Zeng, H. Chen, Y. Wang & E. Jeppesen (2024) Influence of fishery

699  management on trophic interactions and biomass fluxes in Lake Taihu based on a trophic

700  mass-balance model exercise on a long-term data series. *Ecological Indicators,* 158.

701 Yu, C. X., Z. Y. Li, Z. H. Xu & Z. F. Yang (2020) Lake recovery from eutrophication: Quantitative

702  response of trophic states to anthropogenic influences. *Ecological Engineering,* 143, 105697.

703 Zhang, Y. L., Y. J. Cai, K. Peng, Z. J. Gong, J. H. Luo, Y. Q. Zhou, J. B. Wiei, S. W. He, N. Li & B. Xue

704  (2024) Challenges and protection strategies of ecological environment of lakes along the

705  Eastern Route of the South to North Water Diversion Project (in Chinese). *Journal of Lake*

706  *Sciences,* 36 (5) , pp.1289-1302.

Appendix for

The past and potential antagonistic effects of fishing ban against

nutrient loading control on eutrophication restoration in Lake

Hongze

Bo Qin[1,3], Vasilis Dakos[2,3]*, Yongjiu Cai[1], Yu Zhao[1], Xiangdong Yang[1,4], Enlou Zhang[1,4], Rong Wang[1,4,5]*

[1] State Key Laboratory of Lake Science and Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, Nanjing 211135, China.

[2] ISEM, Univ Montpellier, CNRS, EPHE, IRD, 34095 Montpellier, France.

[3] Sorbonne Université, CNRS, IRD, INRAE, Université Paris Est Créteil, Université de Paris Cité, Institute of Ecology and Environmental Sciences of Paris (iEES-Paris), Paris, France.

[4] University of Chinese Academy of Sciences, Beijing 100049, P. R. China.

[5] The Fuxianhu Station of Plateau Deep Lake Field Scientific Observation and Research, Yunnan, Yuxi 653100, China.

* Correspondence:

Vasilis Dakos (vasilis.dakos@umontpellier.fr) and Rong Wang (rwang@niglas.ac.cn)

Figure S1. PCLake model structure (lake part). Cyanobacteria, diatoms and other algae are three distinguished functional groups of phytoplankton. Fish is split into a juvenile (zooplanktivorous) and an adult (benthivorous) subgroup (Janse, 2005).

Figure S2. Monthly observations of water quality. Zone 1 (Chengzi District) is in the north and Zone 3 (Lihewa District) in the west, and Zone 2 is the open water area.

Figure S3. Zoobenthos composition in biomass and total dry weight of three zones in Lake Hongze in the period of 2016-2020.

Figure S4. The simulation using original model before calibration (line) and in-lake observations of different zones in Lake Hongze between January 2016 and December 2019 (dots). The coefficient of determination (R2) and root mean squared error (RMSE) were calculated according to observations of whole-lake average and model simulations.

753
754

Table S1. Average river inflow of Lake Hongze in the period of January 2016 and December 2019.

Month	Chihe	Huaihongxinhe*	Huaihongxinhe**	Huaihe	Xinbianhe	Suihe	Laosuihe
1	1.08	0.89	26.47	610.75	0.36	11.81	0.00
2	0.33	2.01	8.73	526.75	0.20	7.46	0.00
3	0.73	1.70	19.05	618.25	0.00	6.88	0.04
4	0.00	3.70	9.15	481.75	0.08	4.23	0.00
5	24.70	0.79	18.19	830.50	0.00	2.50	0.19
6	16.75	14.74	49.66	911.00	0.24	10.02	3.75
7	24.08	37.51	85.88	1408.75	0.13	25.19	7.00
8	15.28	116.10	132.75	864.03	67.25	91.48	12.73
9	13.38	17.63	78.12	807.75	7.89	13.50	0.99
10	19.93	43.40	109.63	1380.18	4.92	41.13	3.05
11	9.68	3.83	37.95	1076.10	0.42	12.60	0.00
12	1.14	0.00	15.81	500.25	0.00	10.26	0.00
Total	127.06	242.28	591.38	10016.05	81.49	237.03	27.74
%	1.1	2.1	5.2	88.5	0.7	2.1	0.2

758 * Shuanggou station ** Xiacaowan station. The data unit for inflow is $m^3 \cdot s^{-1}$.

761 **Table S2. The performance of original and calibrated model for Lake Hongze, as well as**
762 **the model performances of other studies.**

	R2			RMSE		
	Original	Modified	Other studies*	Original	Modified	Other studies**
Total nitrogen	0.64	0.65	0.03-0.89	0.44	0.43	0.81-1.64
Total phosphorus	0.72	0.72	0-0.63	0.03	0.03	0.02-0.16
Dissolved oxygen	0.77	0.80	0-0.94	1.17	1.03	0.11-2.24
Secchi depth	0.44	0.39	0.18-0.63	0.20	0.13	0.12-0.25
Chlorophyll-a	0.34	0.42	0.05-0.94	0.47	0.23	0.35-48.1
Macrophyte coverage	0.37	0.39	0.34	19.49	3.7	4.95
Zoobenthos	-0.4	0.30	0-0.96	1.6	0.27	
Fish catch	0.61	0.60	0.18-0.83	0	0	

_ Number from other studies were collected by Qin et al. (2025)_*

Qin, B., Xu, M., Zhu, K., et al. 2025. Climate warming will significantly affect future restoration and level
of ecosystem services in Lake Erhai. ECOLOGICAL MODELLING, 503, 111067.
https://doi.org/https://doi.org/10.1016/j.ecolmodel.2025.111067

*_** Number was collected in this study from below literatures:_*

Fragoso, C. R., Marques, D. M. L. M., Ferreira, T. F., et al. 2011. Potential effects of climate change
and eutrophication on a large subtropical shallow lake. ENVIRONMENTAL MODELLING &
SOFTWARE, 26(11), 1337-1348. https://doi.org/10.1016/j.envsoft.2011.05.004

Kong, X. Z., Dong, L., He, W., et al. 2015. Estimation of the long-term nutrient budget and thresholds of
regime shift for a large shallow lake in China. ECOLOGICAL INDICATORS, 52, 231-244.
https://doi.org/10.1016/j.ecolind.2014.12.005

Kong, X., Zhan, Q., Boehrer, B., et al. 2019. High frequency data provide new insights into evaluating
and modeling nitrogen retention in reservoirs. WATER RESEARCH, 166.
https://doi.org/10.1016/j.watres.2019.115017

Andersen, T. K., Nielsen, A., Jeppesen, E., et al. 2020. Predicting ecosystem state changes in shallow
lakes using an aquatic ecosystem model: Lake Hinge, Denmark, an example. ECOLOGICAL
APPLICATIONS, 30(7). https://doi.org/10.1002/eap.2160

Qin, B., Kong, X., Wang, R., et al. 2022. Lake restoration time of Lake Taibai (China): a case study
based on paleolimnology and ecosystem modeling. J Paleolimnol, 68, 25–38.
https://doi.org/https://doi.org/10.1007/s10933-020-00165-7

Shi, T., Chen, Y., Zhang, H., et al. 2023. Numerical study on regime shifts in an urban subtropical
shallow lake: Xinglong Lake, China. ECOLOGICAL INDICATORS, 154.
https://doi.org/10.1016/j.ecolind.2023.110600

Zhang, X., Yi, Y., Cao, Y., et al. 2023. Disentangling the effects of phosphorus loading on food web
stability in a large shallow lake. JOURNAL OF ENVIRONMENTAL MANAGEMENT, 328.
https://doi.org/10.1016/j.jenvman.2022.116991

Table S3. Standardized path coefficients from SEM path analysis between lake components.

Path	Standardized Regression Weights	
	With fishing ban	Without fishing ban
Total phosphorus-Phytoplankton	0.22	0.42
Phytoplankton-Zoobenthos	-0.77	-0.28
Zoobenthos-Secchi depth	-0.91	0.00
Phytoplankton-Secchi depth	-0.73	0.56
Secchi depth-Macrophyte	0.60	0.32
Total phosphorus-Macrophyte	-0.25	-0.13
Macrophyte-Zooplankton	0.35	0.09
Phytoplankton-Zooplankton	-0.04	0.63
Zooplankton-Fish	-0.62	-0.34
Zoobenthos-Fish	-0.58	0.14