Full title: **The effects of temperature on nestling growth in a songbird depend on**

developmental and social context

Short title: **Temperature and songbird nestling growth across developmental and social**

contexts

Sage A. Madden[1]*, Rebecca J. Safran[2], Gail L. Patricelli[1], Sara R. Garcia[2], Zachary M. Laubach[2]

Affiliations:

[1] Department of Evolution and Ecology, University of California Davis, Davis, California,

United States of America

[2] Department of Ecology and Evolutionary Biology, University of Colorado Boulder, Boulder,

Colorado, United States of America

* Corresponding author

E-mail: saamadden@ucdavis.edu (SAM)

Abstract

Climate change can adversely impact animals, especially those that cannot independently thermoregulate or avoid exposure. Cold, hot, and variable temperatures may impede nestling songbird growth due to increased thermoregulatory costs and reduced food delivery by parents. At a broad scale, temperature effects on nestling growth vary across climatic zones, but how temperature effects vary with early-life developmental constraints imposed by the timing of thermoregulatory development, competition with siblings, and the amount of parental care has received less attention. We investigated whether the effect of temperature on the mass of wild barn swallow (*Hirundo rustica erythrogaster*) nestlings (n = 113 nestlings, 31 nests) in Boulder County, CO depends on timing of exposure during development, relative size within the brood, or level of parental feeding. We found that the effect of minimum temperature differed in early and late development (before versus after putative development of thermoregulatory independence) and may be less pronounced for nestlings that received more parental feeding. We did not find evidence that the smallest nestling in the brood differs from other nestlings in vulnerability to temperature. These findings indicate the existence of fine-scale heterogeneity in which the effects of temperature on nestling development are sensitive to metabolic constraints and early-life social environment, providing key insight into the factors that may ameliorate or exacerbate climate impacts on individual birds.

Introduction

Rising and increasingly variable temperatures due to climate change [1] can adversely impact animals, especially those that cannot independently thermoregulate or avoid exposure. At risk individuals include young animals lacking the mechanisms required to regulate exposure and cope with temperature extremes [2]. Exposure to excessively hot or cold temperatures is a form of developmental stress that negatively impacts not only the immediate survival of young animals in many species, but also the development of adult phenotypes [3,4]. Songbird nestlings may be particularly vulnerable to effects of extreme and variable temperatures because they are confined to a nest, ectothermic and mostly featherless early in development [5,6], and entirely dependent on parental care [7,8]. As a result, exposure to extreme or variable temperatures may impede nestling growth due to thermoregulatory costs and reduced food delivery by parent birds [9,10]. Although previous studies have revealed temperature effects on nestling growth across climate zones [9,10], we know little about how these effects may vary with localized early-life developmental constraints imposed by the timing of thermoregulatory development, competition with brood mates, and the amount of parental care. This knowledge is key for understanding heterogeneity in susceptibility of nestling birds to climate impacts and the factors that may ameliorate or exacerbate these impacts.

The effects of temperature on nestling development may differ depending on the timing of exposure during development [11–13]. Young nestlings initially lack the ability to thermoregulate independently and are mostly featherless [8,14], and body temperatures outside of a narrow range may disrupt growth [5,6,11,13]. The ability to thermoregulate develops partway through the nestling period [6,14], and extreme temperatures may require expending

energy on thermoregulation and/or entering a state of hyperthermia, leading to increased stress

and decreased growth [6,15,16]. The development of thermoregulatory independence partway

through the nestling period suggests that effects of temperature may differ across nestling

ontogeny. However, most studies have focused on the nestling period as a whole (but see [11–

13,17]), leaving unanswered whether the effects of temperature differ before versus after

development of thermoregulatory independence.

A nestling's social environment also may mitigate or exacerbate the effects of temperature on

their development [18,19]. Songbird nestlings often hatch asynchronously, resulting in size

asymmetry that is maintained across development [20]. This asymmetry may affect access to

resources—older nestlings outcompete younger nestlings for food, and younger nestlings often

grow more slowly and even die from starvation [20–22]. Heterogeneity in access to resources

within a brood may then drive differences in susceptibility to temperature. Additionally, body

size is a critical determinant of thermoregulatory ability [6,23], but most studies have

investigated the relationship between size and thermoregulation among rather than within species

[24,25]. As such, investigating whether the size of a nestling relative to its brood mates

influences responses to temperature can help elucidate how early-life social environment shapes

nestling responses to ecological conditions [21,26–28].

Parental care is another aspect of the social environment that may shape nestling responses to

temperature [18,19,29]. Nestlings are entirely dependent on food provisioned by parents to

provide energy for growth and thermoregulation. Nestlings that receive more parental care may

be less susceptible to the negative effects of temperature because they have more energetic

reserves for thermoregulation and growth. Indeed, research on burying beetles (*Nicrophorus*

vespilloides) indicates that higher levels of parental care can buffer young from effects of

temperature variation [30]. Similarly, studies of cooperatively breeding bird species suggest that

the presence of helpers at the nest may mitigate negative impacts of weather on nestlings [18]

(but see [29]). Therefore, it is plausible that parental care influences nestling vulnerability to

extreme temperatures in songbirds.

In this study, we investigated the effects of temperature at different stages across nestling

ontogeny in wild barn swallows, and we explored whether nestling size relative to brood mates

or level of parental feeding modify the effects of temperature on nestling growth. We aimed to

address the following three questions (Fig 1): 1) Does the effect of temperature on nestling mass

differ when the exposure is assessed during early versus late development? 2) Does the effect of

temperature on nestling mass depend on whether a nestling is the smallest in the brood? 3) Does

the effect of temperature on nestling mass depend on parental care? For question one, we

hypothesized that exposure to extreme and variable temperatures would impose greater

metabolic costs on nestlings before thermoregulatory independence than after, leading to

stronger negative effects of temperature on mass early in development than late. For questions

two and three, we hypothesized that advantageous social environments (e.g., larger size within

the brood hierarchy, higher levels of parental feeding) would ameliorate some of the negative

impacts of extreme and variable temperatures on nestling mass due to the nestlings' greater

access to resources. Directed acyclic graphs [31] displaying hypothesized relationships

investigated for each question are provided in S1-S3 Fig.

Fig 1. Research questions and predicted results for our hypotheses about temperature

effects on wild barn swallow nestling mass before fledging. We expected that lower minimum

temperature, higher maximum temperature, and greater temperature variability (interquartile

range) would be associated with lower nestling mass. Feeding level refers to the average nest-

level feeding rate across conditions.

Methods

Study system

We monitored nesting attempts from wild barn swallows during the summer of 2021 at

seven breeding sites in Boulder County, CO. These sites are monitored as part of a long-term

study of barn swallows and are occupied by colonies ranging in size from one to 50 breeding

pairs. Barn swallows build open mud cup nests and lay clutches of three to five eggs, which

hatch asynchronously over approximately 1-2 days [32]. Hatching asynchrony results in a size

hierarchy that persists over the course of development [33]. Hatch order is not associated with

offspring sex and there are no morphological differences between different sexes [34]. In this

socially monogamous species, both parents in a social pair provide care, including food

provisioning, until nestlings fledge (>18 days post hatch) [32].

Study design overview

We collected environmental, demographic and behavioral data at 31 first brood nests

from May through July 2021, and we measured morphometric and physiological traits on 113

nestlings across development (Fig 2). Nests were checked every three to five days to track

nestling phenology and fate. During the first check after nestlings hatched, we estimated their

ages (hatch day = day zero) based on feather emergence and other reliable developmental

characters, such as wetness (on hatch day) and ability to raise their head [35]. We took

morphological measurements of nestlings and conducted one-hour observations of parental care

behavior at three time points: days three to four, days eight to nine, and days 11-13. We

monitored temperature near the nest from hatch through last nestling measures.

Fig 2. An overview of the study design indicating the types of data collected across the

nestling period at three time points in wild barn swallows in Boulder County, CO. We

expect thermoregulatory independence to develop at ~six days post-hatch in barn swallows (see

"Near-nest temperatures").

Nestling measures

We collected standard morphometric measurements for each nestling at three time points

([36]; Fig 2). We measured mass to the nearest 100th of a gram using a digital scale and right-

wing length to the nearest half millimeter as the distance from the carpal joint to the distal end of

the second phalanx or primary wing feather, whichever was longer. On days eight to nine, we

placed a USGS metal band on the tibiotarsus of each nestling, allowing us to track individuals

through development. On days eight to nine and 11-13, we also collected two small blood

samples from each nestling as part of a separate study [36].

Using day eight to nine measures (the earliest age at which we could begin tracking

individuals), we created a categorical variable of relative nestling size, where the smallest

nestling in each nest based on right-wing length was classified as 'min' (reference group) and the

remaining nestlings were classified as 'other'. Our classification was based on right wing length,

rather than mass, because wing length measures skeletal size and may not be as strongly

influenced by short-term fluctuations as mass [37–39]. Because we were unable to track

individuals until days eight to nine, we could not determine the exact age of individual nestlings.

Therefore, we are unable to tease apart the effects of age and mass in the relationship between

relative nestling size and vulnerability to adverse temperatures.

Parental care observations

We conducted ~one-hour focal observations of each nest at three time points (see details in

[36]; Fig 2). Some nests are missing data for one or more time points due to logistical issues. The

start of each observation was preceded by collection of morphological measurements and blood

samples from nestlings (see Nestling Measures). After these measures, observers watched the

nest from a blind, and the birds were given ~15 minutes to habituate to our presence before the

observation commenced (following [36,40]). Observations began between 5:45 and 8:00 am.

During each observation, the observer logged parental care behavior in real-time on iPads or

iPhones (Apple Inc.) using "Animal Behavior Pro" [41]. Behaviors recorded included feeding

visits, where the parent delivers food to a nestling (see full ethogram in S1 Table). Given

biparental care in this species, we recorded the total number of behaviors for both social parents.

When an observer could not be present, we instead recorded nests using GoPro cameras, and

videos were scored by the same observers who conducted the in-person observations. For these

camera observations, we set up camera mounts the day before recording to allow birds to

habituate, and we waited for ~15 minutes after the cameras were started to begin scoring

behavior. Parental care behaviors are consistent when scored by different observers and between

in-person and camera observations [40].

We created an index of the level of parental care by summarizing feeding rates from both

parents across all stages of nestling development. Specifically, this index was created using a

generalized linear mixed effects model, in which total feeding count by both parents measured at

each developmental stage was the outcome and nest ID was a random intercept. We compared

model fit for several error distributions and link functions using diagnostic plots from the

package 'DHARMa,' version 0.4.6 [42], and we selected a negative binomial distribution and

log link for our final model. The model included an offset for total observation time and several

covariates that could influence parental feeding behaviors: number of days since the first nestling

hatched, number of nestlings in the nest, the median temperature near the nest during the

observation period, and the duration of time between removing nestlings from the nest and the

start of the observation. From this model, we extracted the best linear unbiased predictors

(BLUPs) and created a three-level variable of parental feeding based on nest-level tertiles of

average feeding rates summarized across development (following [36]).

Using a mixed effects model in this way allowed us to summarize parental care across

developmental stages while controlling for environmental conditions and nest-level variables that

may influence parental care. Moreover, this method is an efficient use of our data because it

allows for missing data via shrinkage of nest-level estimates toward the global sample mean.

Because parental feeding is an effect modifier in our models, we created a three-level categorical

variable for stratified analyses by classifying BLUPs as 'low' (lowest third), 'med' (middle

third), or 'high' (highest third) parental feeding. The approach of carrying BLUPs forward from

one model to another has been criticized due to its failure to propagate uncertainty [43].

However, because summarizing parental feeding using BLUPs should not substantively shift

nests from one parental feeding level to another, this approach is justified in our case given the

advantages described above.

Near-nest temperatures

We hung thermometers (Govee) in small mesh bags adjacent (10-28 cm) to each open

cup nest and logged near-nest temperature and humidity every 15 minutes throughout the data

collection period (Fig 2). At one nest, we are missing two days of temperature measurements at

the end of the nestling period. We calculated the minimum temperature, maximum temperature,

and temperature variability (defined as the interquartile range) at each nest from day zero (hatch)

to the end of day five (early development) and from day six until the last measure of nestling

mass on day 11-13 (late development). These temperature summaries correspond with before

and after the putative development of thermoregulatory independence, respectively. We chose

these time periods based on previous research in swallows and other songbirds [14,44–46],

which found that nestling swallows begin to develop the ability to thermoregulate independently

around four to five days post-hatch, and the age of the effective homeothermy (defined here as

ability to maintain relatively constant body temperature under natural conditions—in the nest

with brood mates) is likely around six days of age. Finally, we calculated the minimum

temperature, maximum temperature, and temperature variability at each nest over the entire

nestling period, from hatch day to the last measure of nestling mass.

Ethics statement

All research on wild barn swallows was approved by the Rebecca Safran's Institutional Animal

Care and Use Committee protocol (permit no. 1303.02), and all research was performed in

accordance with relevant guidelines and regulations. In addition, this work was conducted under

the Bird Banding Lab, master permit bander ID 23505. Capture using mist nests in Boulder,

Weld and Jefferson Counties, Colorado, was approved by the Colorado Parks and Wildlife

issued under permit license number 21TRb2005. Some swallow nests were located on private

property and were monitored with permission from the landowners.

Statistical analysis

We conducted three sets of formal analyses aligning with our questions. Across all

questions, we investigated: 1) minimum temperature, which captures exposure to extreme cold,

2) maximum temperature, which captures exposure to extreme heat, and 3) temperature

variability, which captures exposure to fluctuations between cold and hot [9,10]. Given our

modest sample size, we balanced our inference based on the consistency, direction, and

magnitude of effect estimates alongside of conventional significance thresholds. We focused our

reporting and interpretation of results on estimates of effect sizes and 95% confidence intervals,

providing information on the strength and likely error of our effect estimates [47], as well as p-

values interpreted using the language of evidence [48]. P-values less than 0.05 were interpreted

as providing moderate to strong evidence for an effect, p-values between 0.05 and 0.10 as

providing marginal evidence, and p-values greater than 0.10 as providing little to no evidence

[48].

Across all analyses we report both unadjusted associations and adjusted associations from

multiple variable regression models; Examining both unadjusted and adjusted associations

allowed us to see if the estimate of interest changes after controlling for other variables. For

adjusted models, we included hatch date and number of nestlings in a nest as potential

confounders. For all models, we z-score standardized numeric explanatory variables and

covariates to aid in model fit and comparison of effect sizes. All models included a random

intercept for nest ID to account for non-independence of measures taken at the same nest. We

originally included a random effect for site ID to account for non-independence of measures

taken at the same site, but this effect was removed because it explained zero variance in some

models. Removal of site ID from models in which it explained some variance did not

substantially affect the direction, magnitude, or significance of effects.

Model assumptions were checked using diagnostic plots. Specifically, we checked

whether each model met the assumptions of a Gaussian error distribution with homogenous

variance by examining plots of residuals versus fitted values, a histogram of the residuals, a

normal quantile-quantile plot, and Cook's distance and standardized residual influence plots,

using the package 'car', version 3.1.3 [49]. For some models, we detected influential outliers,

which we defined as having a Cook's distance greater than one or a standardized residual value

greater than three. When influential outliers were present, we conducted sensitivity analyses,

where we ran the model with and without the outlier(s) and compared the results.

For question one, we separately ran two sets of linear mixed models for each of our three

temperature variables (minimum, maximum, and variability) (S1 Fig): 1) In the first set of three

models, each temperature variable collected in early development (from hatch to the end of day

five) was included as the explanatory variable of interest and nestling mass on days 11-13 was

the outcome; 2) the second set of models were the same except that temperature variables were

quantified in late development (from day six to last nestling measures on days 11-13). We

conducted a sensitivity analysis to assess the robustness of our results to differences in the

selected cut-off for development of thermoregulatory independence. Specifically, we ran the

same set of models described above, except using cut-off ages of five days post-hatch and seven

days post-hatch, rather than six days post-hatch, for development of thermoregulatory

independence.

For questions two and three, we tested for effect modification of temperature effects on

nestling mass by relative nestling size and level of parental feeding, respectively. Effect

modification refers to a situation in which the effect of X on Y depends on a third variable, the

effect modifier [31]. In our analysis, relative nestling size (question two) and level of parental

feeding (question three) were treated as effect modifiers. To determine whether effect

modification was present, we ran analyses stratified by multiple levels of relative nestling size or

parental feeding. For question two, we ran models of the effect of each temperature variable on

mass stratified by nestling size, 'min' = the smallest nestling in the nest and 'other' = all other

nestlings. For question three, we modeled the effect of temperature on mass stratified by

parental feeding, which included three levels, 'low' = parental feeding below the ~33rd

percentile, 'med' = parental feeding between the ~33rd and ~67th percentile, and 'high' = parental

feeding above the ~67th percentile. S4 Fig displays the feeding rate BLUPs, grouped into three

strata. The results from these stratified models provide insight into the magnitude and

directionality of the effect of temperature on nestling mass for different relative nestling sizes or

levels of parental feeding.

All statistical analyses were carried out using R version 4.4.1 [50]. The 'tidyverse'

package, version 2.0.0 [51] was used for data cleaning and organization. The 'lme4' package,

version 1.1.35.5 [52] was used to fit models. Bootstrapped confidence intervals were obtained

for model coefficients using the package 'boot', version 1.3.30 [53,54]. P-values for two-tailed t-

tests for each coefficient were obtained using Satterthwaite degrees of freedom method, via the

package 'lmerTest' version 3.1.3 [55]. Goodness of fit was assessed using conditional and

marginal R-squared values, calculated using the package 'MuMIn', version 1.48.4 [56]. Results

were visualized using the package 'ggplot2' version 3.5.1 [57], 'ggeffects' version 1.7.2 [58],

and 'ggpubr' version 0.6.0 [59].

Results

Background characteristics

Of the 113 nestlings in this study, 108 survived until days eight to nine, and 106 survived

until days 11-13. Eggs hatched asynchronously over up to two days—the mean difference in age

between the first and last hatched nestling was 1.03 days (range = 0-2 days). For the near-nest

temperature measurements averaged across the nestling period, the mean ± SD was 13.10 ± 1.80

degrees C for minimum temperature, 36.90 ± 2.96 degrees C for maximum temperature, and

9.26 ± 2.43 degrees C for temperature variability. We categorized nestlings based on within nest

relative size on days eight to nine, measured in terms of right-wing length. At this time, the mean

± SD right wing length was 26.8 ± 5.55 mm for the smallest ('min') nestlings and 32.4 ± 4.26

mm for 'other' nestlings. The mean ± SD days eight to nine mass was 11.60 ± 2.61 grams for the

smallest ('min') nestlings and 13.60 ± 2.30 grams for 'other' nestlings. On days 11-13, the

timepoint at which the outcome was measured, the mean ± SD wing length was 49.20 ± 8.09 mm

for the smallest ('min') nestlings and 54.40 ± 4.76 mm for 'other' nestlings. The mean ± SD days

11-13 mass was 17.20 ± 2.75 grams for the smallest ('min') nestlings and 17.70 ± 1.91 grams for

'other' nestlings. For measures of parental care, the mean ± SD total feeding rate (counts/hour)

was 12.21 ± 6.92 on days three to four, 12.73 ± 8.08 on days eight to nine, and 21.62 ± 12.94 on

days 11-13. Feeding rate and proportion of time spent brooding during the observation were

negatively correlated (Spearman Rank Correlation: rho = -0.49, p < 0.001). Additional

background characteristics for the study population are provided in S2 Table. S5 Fig displays

daily minimum temperature, maximum temperature, and temperature variability across the

nestling period for each nest.

Question 1. Does the effect of temperature on nestling mass differ when the exposure is assessed during early versus late development?

Higher minimum temperatures in early development (β = 1.16 g per 1 SD ℃ [95% CI:

0.32, 2.00], P = 0.01) but not late development (β = 0.29 g per 1 SD ℃ [95% CI: -0.45, 1.06], P

= 0.46) were associated with higher nestling mass at last measure (S3 Table; Fig 3a). Higher

maximum temperatures in early (β = -0.66 g per 1 SD ℃ [95% CI: -1.32, -0.04], P = 0.05) and

late (β = -1.13 g per 1 SD ℃ [95% CI: -1.74, -0.52], 0.001) development were associated with

lower nestling mass (S3 Table; Fig 3b). Similarly, greater temperature variability in early (β = -

1.41 g per 1 SD ℃ [95% CI: -2.07, -0.77], P = 0.0003) and late (β = -1.33 g per 1 SD ℃ [95%

CI: -1.95, -0.70], P = 0.0003) development was associated with lower nestling mass (S3 Table;

Fig 3c).

Fig 3. Temperature effects in early and late development on the mass of 11–13-day old wild

barn swallow nestlings in Boulder County, CO. Predicted relationships of minimum

temperature (a), maximum temperature (b), and temperature variability (c) and nestling mass

from separate linear mixed models for each developmental stage—in early ('Early')

development, from hatch to end of day five days post-hatch, and late ('Late') development, from

six days post-hatch to final measures at 11-13 days post-hatch. Model predictions are displayed

as lines and raw data are displayed as points. Colors, line types, and shapes correspond to developmental stage.

The estimates of interest did not change substantively when influential outliers were excluded from analyses or when models were run using five days or seven days post-hatch, rather than six days post-hatch, as the cut-off for age for early versus late development.

Question 2. Does the effect of temperature on nestling mass depend on whether a nestling is the smallest in the brood?

In stratified analyses, higher minimum temperatures across the nestling period corresponded with higher nestling mass for both the smallest nestlings in the nest (β = 1.75 g per 1 SD ℃ [95% CI: 0.43, 3.14], P = 0.02; S4 Table; Fig 4a) and for other nestlings (β = 1.30 g per 1 SD ℃ [95% CI: 0.40, 2.21], P = 0.01; S4 Table; Fig 4a). Similarly, higher maximum temperatures and greater temperature variability across the nestling period were associated with lower nestling mass for the smallest nestlings in the nest (maximum: β = -1.42 g per 1 SD ℃ [95% CI: -2.48, -0.40], P = 0.01; variability: β = -1.95 g per 1 SD ℃ [95% CI: -2.98, -0.92], P = 0.001; S4 Table; Fig 4b-4c) and for other nestlings (maximum: β = -0.85 g per 1 SD ℃ [95% CI: -1.47, -0.27], P = 0.01; variability: β = - 1.50 g per 1 SD ℃ [95% CI: -2.15, -0.89], P = 0.0001; S4 Table; Fig 4b-4c).

Fig 4. Temperature effects on the mass of 11-13-day old wild barn swallow nestlings in Boulder County, CO that are the smallest in their brood versus other nestlings. Predicted

relationships of minimum temperature (a), maximum temperature (b), and temperature

variability (interquartile range) (c) across the nestling period and nestling mass from linear

mixed models, stratified by relative nestling size—the smallest nestling ('Smallest') and all other

nestlings ('Other')—at days eight to nine post-hatch. Model predictions are displayed as lines

and raw data are displayed as points. Colors, line types, and shapes correspond to relative

nestling size. Because relative nestling size was determined at days eight to nine and some

nestlings did not survive to final measures on days 11-13, for some nests, only 'Smallest' or

'Other' nestling points are present.

Question 3. Does the effect of temperature on nestling mass depend on the overall amount of care provided by parents?

In stratified analyses, higher minimum temperatures across the nestling period were

associated with higher nestling mass among nests that received low parental feeding, but the

confidence interval overlapped zero (β = 1.73 g per 1 SD ℃ [95% CI: -0.11, 3.53], P = 0.11; S5

Table; Fig 5a). Higher minimum temperatures were also associated with higher nestling mass

among nests that received medium parental feeding (β = 2.36 g per 1 SD ℃ [95% CI: 0.68, 3.92],

P = 0.03; S5 Table; Fig 5a). In contrast, we observed no clear association of minimum

temperatures with nestling mass among nests that received high parental feeding (β = -0.53 g per

1 SD ℃ [95% CI: -2.05, 0.94], P = 0.53; S5 Table; Fig 5a).

Fig 5. Temperature effects on nestling mass of 11–13-day old wild barn swallows in Boulder County, CO at nests receiving low, medium, or high parental feeding. Predicted relationships of minimum temperature (a), maximum temperature (b), and temperature variability (interquartile range) (c) across the nestling period and nestling mass from linear mixed models, stratified by level of parent feeding—low ('Low), medium ('Med'), and high ('High'). Model predictions are displayed as lines and raw data are displayed as points. Colors, line types, and shapes correspond to level of parent feeding.

Similarly, higher maximum temperatures and greater temperature variability across the nestling period were associated with lower nestling mass among nests that received low parental feeding, but confidence intervals included zero (maximum: β = -1.08 g per 1 SD ℃ [95% CI: -2.44, 0.34], P = 0.16; variability: β = -1.35 g per 1 SD ℃ [95% CI: -2.77, 0.14], P = 0.11; S5 Table; Fig 5b-c). Higher maximum temperatures and greater temperature variability were also associated with lower nestling mass among nests that received medium parental feeding (maximum: β = -1.22 g per 1 SD ℃ [95% CI: -2.34, -0.04], P = 0.07; variability: β = -1.80 g per 1 SD ℃ [95% CI: -2.78, -0.75], P = 0.01; S5 Table; Fig 5b-c). In contrast, we observed no clear association of maximum temperature or temperature variability with nestling mass among nests that received high parental feeding (maximum: β = 0.01 g per 1 SD ℃ [95% CI: -0.91, 0.92], P = 0.98; variability: β = -0.01 g per 1 SD ℃ [95% CI: -1.25, 1.28], P = 0.99; S5 Table; Fig 5b-c).

The magnitude, direction, and precision of the estimates of interest did not substantively change when influential outliers were excluded from analyses.

Discussion

In this study of 113 wild barn swallow nestlings in 31 nests, we investigated whether the

effects of minimum temperature, maximum temperature, and temperature variability on nestling

mass differ according to the timing of exposure across nestling ontogeny and two features of the

nestlings' social environment: the nestlings' size relative to brood mates and the level of parental

feeding the nest received. We found that exposure to lower minimum temperatures, higher

maximum temperatures, and greater temperature variability were associated with lower nestling

mass. Additionally, we found some evidence for heterogenous effects of temperature on

nestlings across developmental stages and social environments. When looking at the effects of

exposure to temperatures in early versus late development (before versus after putative

development of thermoregulatory independence), we found different relationships of minimum

temperature with nestling mass. Surprisingly, the association of temperature with nestling growth

did not vary with relative nestling size, suggesting that sibling size dynamics may not have a

large impact on nestling vulnerability to adverse temperatures. Finally, we found that the

relationship between temperature and nestling mass observed at medium levels of parental

feeding were absent from high parental feeding nests. Our work adds to relatively limited

knowledge of how temperature effects on nestlings vary with developmental constraints imposed

by timing of thermoregulatory development and aspects of early-life social environment, which

is critical to understanding factors that may ameliorate or exacerbate climate change impacts on

birds.

410   **Minimum temperature, maximum temperature, and temperature**

411   **variability have differential effects on nestling mass**

Our results align with literature reporting that exposure to cold, hot, and variable

temperatures may all lead to reduced nestling growth. We found that lower minimum

temperatures were generally associated with lower nestling mass, consistent with previous

studies finding that exposure to cold temperatures leads to reduced nestling growth [11,60,61].

On the other hand, higher maximum temperatures were generally associated with lower nestling

mass in our population as in other studies [62–64]. Greater temperature variability was generally

associated with lower nestling mass, consistent with evidence that nestling growth is reduced

when exposed to more variable than stable temperatures [65].

Notably, some studies report no effect of extreme or variable temperatures on nestling

growth [66–68]. Ecological and social factors, such as habitat, diet, and nest type, may shape

susceptibility to negative effects of temperature across species and populations [18,69]. For

example, insectivorous species may be more sensitive to extreme temperatures than granivorous

species because insect availability is more strongly influenced by weather conditions [70,71].

Additionally, open-cup nesting species may be more sensitive to extreme temperatures than

cavity-nesting species because open-cup nests may be less thermally buffered than cavity nests

[69]. Given that barn swallows are insectivorous and build open cup nests, it is perhaps

unsurprising that nestling mass is negatively affected by cold, hot, and variable temperatures in

this species.

430   Effects of temperature on nestling mass may differ across nestling

431   ontogeny

Consistent with our hypothesis, we found that lower minimum temperatures in early but

not late development--*before but not after* putative development of thermoregulatory

independence at six days post-hatch—were associated with lower nestling mass (Fig 3a).

Nestlings may be especially sensitive to cold temperatures before thermoregulatory

independence because colder ambient temperatures result in colder body temperatures at times

when the parents are away from the nest, and colder temperatures slow metabolic processes in

young nestlings [72]. In older nestlings, improved thermoregulation allows for a more constant

body temperature and metabolic rate increases when ambient temperatures are outside the

thermal neutral zone [6]. Additionally, due to their higher surface area to volume ratio and lack

of feathers, young nestlings may dissipate heat more quickly than older nestlings, which may be

a disadvantage under cold conditions [6,8,68].

Contrary to our hypothesis, higher maximum temperatures and greater temperature

variability in early and late development—*before and after* putative development of

thermoregulatory independence—were associated with lower nestling mass (Fig 3b-c). This

suggests that nestlings may be sensitive to heat and environmental instability regardless of

thermoregulatory status, though previous work indicates that the nature of thermal challenges

may vary across development. Early in development, nestlings may be susceptible to negative

impacts of extreme heat because they may gain heat more quickly than older nestlings due to

their high surface area to volume ratio and lack of feathers [6,8,68]. Late in development, after

development of thermoregulatory independence, nestlings may be sensitive to hot temperatures

because evaporative cooling is energetically costly and may quickly lead to dehydration [64,73].

In addition, nestlings may have limited ability to cool themselves [73], and this may be

exacerbated in older nestlings by reduced heat dissipation due to higher surface area to volume

ratio, feather development, and crowding in the nest (though as noted above, reduced heat

transfer may be an advantage when ambient temperatures exceed body temperature). Finally,

older nestlings have higher energetic requirements, which are likely near or in excess of the

limits of the food parents can provision [74,75], and they may be more strongly affected by

changes in food abundance or parental care behavior than younger nestlings [11].

460 Little evidence that effects of temperature on nestling mass differ by

461 relative size

Contrary to our hypothesis, we did not find evidence for effect modification by relative

nestling size on the relationship between temperature and nestling mass—lower minimum

temperatures, higher maximum temperatures, and greater temperature variability were associated

with lower nestling mass regardless of relative nestling size (Fig 4).

Our findings contrast with previous work suggesting that animals with a smaller body

size are more vulnerable to extreme temperatures because they are less able to regulate their

body temperature—they dissipate heat more quickly when ambient temperature is below body

temperature, and they absorb heat more quickly when ambient temperature exceeds body

temperature [6,23]. Consistent with this idea and in contrast to our findings, a study of tree

swallows found that the smallest nestling in a brood had a lower chance of surviving harsh cold

snaps than larger nestlings [26]. Similarly, in studies of black kites (*Milvus migrans*) and red-

winged blackbirds (*Agelaius phoeniceus*), higher maximum and mean temperatures were linked

to faster growth in the last hatched nestling, but not other nestlings [21,28].

Our findings are also surprising given that smaller nestlings may face a competitive

disadvantage compared to their brood mates. Larger nestlings often outcompete younger

nestlings for resources, which are crucial thermoregulation and other developmental processes

like growth [20,21]. For example, in a study of blue tits (*Cyanistes caeruleus*), higher levels of

479 rainfall had stronger positive impacts on the mass of early hatched nestlings than that of later

480 hatched nestlings, which the authors suggested may be due to the greater ability of early hatched

481 nestlings to secure food from parents than late hatched nestlings [27]. Additionally, competitive

482 dynamics may shift under adverse conditions—extreme temperatures may exacerbate the

483 competitive disadvantage of small nestlings [76]. Therefore, we might expect that larger

484 nestlings that more efficiently thermoregulate or that can acquire more resources are likely to be

485 less susceptible to extreme temperatures.

486  In contrast to this idea and findings previous studies, the lack of effect modification we

487 observed may reflect that the sibling size hierarchy does not strongly affect nestling vulnerability

488 to temperature in our study system. This may be because the differences in size among siblings

489 were relatively small—at the timepoint when relative size was determined in our study, the mean

490 $\pm$ SD right wing length was 26.8 ± 5.55 mm for the smallest ('min') nestlings and 32.4 ± 4.26

491 mm for 'other' nestlings.

Effects of temperature on nestling mass may differ by level of parental feeding

494  In support of our hypothesis, we found some evidence for effect modification by parental

495 feeding level for the association of temperature with nestling mass. Stratified models showed that

496 lower minimum temperatures, higher maximum temperatures, and greater temperature variability

497 were associated with lower nestling mass at nests receiving medium, but not high, levels of

498 parental feeding (Fig 5). There was also a pattern of cold, hot, and variable temperatures being

499 associated with lower nestling mass at nests receiving low levels of parental feeding, though the

500 confidence intervals for these associations overlapped zero.

These findings suggest that high levels of parental feeding may mitigate some of the

negative effects of adverse conditions on nestling development and are consistent with several

studies in non-avian species and cooperatively breeding birds. As previously mentioned,

maternal care in burying beetles reduced the adverse effects of cold temperatures on larvae.

Similarly, in sociable weavers (*Philetairus socius*), a cooperatively breeding bird species, having

additional helpers at a nest counteracted some negative effects of adverse weather conditions

(periods of low rainfall) on fledgling condition [18]. However, another study in the same species

reported that having larger groups feeding nestlings did not mitigate the negative effects of

adverse weather [29].

In addition to feeding, parental care may also involve other behaviors, like brooding, that

influence nestlings' responses to temperature exposure. For example, if parental feeding and

brooding positively covary, as some studies have reported [77,78], then the effect of parental

care on nestling mass maybe driven by active parental regulation of nest temperatures, in

addition to or instead of access to higher energetic resources via food provisioning. However, in

our study population (and other studies: [79,80]), feeding and brooding are negatively correlated,

suggesting that positive covariation among these behaviors is not driving our observed

differences in temperature effects across parental feeding levels.

Finally, it would be valuable for future work to tease apart the roles of average levels of

parental care across development, as we have done, in addition to considering changes in specific

bouts of parental care in response to temperature, in shaping effects of temperature on nestling

development. In other words, parental care may act as a modifier of temperature effects on

nestlings if these effects differ with average level of care, and as a mediator temperature effects

if extreme temperatures cause reduced care, leading to reduced nestling growth. An increasing

body of work documents the role of parental care as a mediator of temperature effects on

nestlings—parent provisioning rates may decrease in hot or cold temperatures due to reduced

availability of insect prey [11,81] and parents diverting more time and energy to

thermoregulation, instead of parental care [82,83]. These changes in parental provisioning

behavior during extreme weather conditions may then lead to reduced nestling survival and

growth [11,63]. However, studies considering parental care as a modifier of temperature effects

seem to be limited, and considering parental care as both an effect modifier and mediator

warrants further investigation.

Limitations and strengths

This study was not without limitations. First, our sample size may limit our ability to

detect weak associations and differences in magnitude among strata, which may have led to

some of our marginal and non-significant findings; further work with larger sample sizes is

needed to provide stronger quantitative evidence for these patterns. Second, we modeled three

temperature variables (minimum, maximum, and variability) separately due to limitations of our

statistical power; further work with a larger sample size is needed to determine the relative

strength of the effects of different temperature variables on nestlings or whether there are

cumulative effects of different types of adverse temperature exposures (see discussion in

[84,85]). Third, we examined differential effects of temperature across nestling ontogeny by

categorizing the nestling period into two time periods, before and after putative development of

thermoregulatory independence, based on previous studies of other species of swallows and

songbirds [14,44–46]. Further studies are needed to understand how vulnerability to adverse

temperatures changes during the gradual development of thermoregulatory independence

[46,72]. Fourth, because we were unable to track individual nestlings from hatch, further work is

needed to differentiate the effects of age and size in the relationship between relative nestling

size and vulnerability to extreme and variable temperatures, such as by manipulating hatch

asynchrony [86]. Additionally, degree of hatch asynchrony might be related to nest temperatures

during incubation, but we were unable to assess this because our loggers were not placed until

hatch. Ffith, in our sample, there was less temperature variability among nests with high feeding

levels than those with medium or low feeding levels. Finally, it is possible that parental care

behavior was affected by collection of morphological and physiological data before our

observations, though we attempted to minimize these effects by waiting until birds appeared to

resume normal behavior before beginning parental care data collection.

This study also has several strengths. In 113 wild nestling barn swallows, we investigated

fine-scale heterogeneity in temperature associations with nestling mass before fledging across

ontogeny and social environments, whereas most previous work has focused on characterizing

overall patterns in a population or on documenting broader-scale environmental dependence

(e.g., across climatic zones). We tracked temperature continuously at the nest-level, rather than at

site- or region-level, as many previous studies have done. The fine-scale resolution of our

temperature data enabled us to more precisely quantify variation in temperature associations with

mass among individuals within the same population, allowing us to investigate the role of

developmental timing and social environment in shaping temperature effects on nestlings.

Additionally, we characterized average nest feeding levels using detailed measures of parental

care behavior collected longitudinally across the nestling period, allowing us to thoroughly

characterize average nest-level differences in feeding and assess whether parental feeding can

buffer nestlings from the negative impacts of extreme and variable temperatures. In addition, by

investigating three temperature variables (minimum temperature, maximum temperature,

temperature variability) that capture different aspects of the thermal environment, our work

offers insight on the effects hot, cold, and variable temperatures on nestlings. We provide a more

complete picture of how different aspects of the thermal environment affect nestling growth,

while broadening our understanding of how developmental and social factors may exacerbate or

ameliorate the impacts of climate change on vulnerable individuals.

Summary and Conclusions

In a study of wild barn swallow nestlings, we found that the associations of temperature

with nestling mass differed among three temperature variables—lower minimum temperatures,

higher maximum temperatures, and greater temperature variability were associated with lower

nestling mass before fledging. Further, the associations of minimum temperature with nestling

mass differed before versus after putative development of thermoregulatory independence. The

associations of temperature with nestling mass did not vary with relative nestling size within a

brood. Finally, we found evidence that the associations of extreme and variable temperature with

nestling mass depended on level of parental feeding, with higher levels of parental feeding

ameliorating the negative effects of temperature. Although prior work has revealed broad-scale

environmental dependence of temperature effects on nestlings, our study unveils factors shaping

fine-scale heterogeneity in temperature effects on nestlings. We offer initial insight into how

early-life developmental constraints imposed by timing of thermoregulatory development,

competition with brood mates, and amount of care provided by parents shape the impacts of

extreme and variable temperatures on individual birds, providing the basis for future research to

tease apart the mechanisms through which climate change affects development.

Acknowledgements

We thank Aleea Pardue, Marina Ayala, Heather Kenny-Duddela, Avani Fachon, Grant Gonzalez, and Molly McDermott for assistance and/or advice regarding field work. We thank the private landowners who give us access to their property where barn swallows nest. Finally, we thank members of the Safran, Patricelli, and Karp labs for their constructive feedback on this project.

References

1. Intergovernmental Panel on Climate Change (IPCC), editor. Long-term Climate Change: Projections, Commitments and Irreversibility Pages 1029 to 1076. Climate Change 2013 – The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press; 2014. pp. 1029–1136. doi:10.1017/CBO9781107415324.024

2. Nord A, Giroud S. Lifelong Effects of Thermal Challenges During Development in Birds and Mammals. Front Physiol. 2020;11: 419. doi: 10.3389/fphys.2020.00419

3. Eyck HJF, Buchanan KL, Crino OL, Jessop TS. Effects of developmental stress on animal phenotype and performance: a quantitative review. Biol Rev Camb Philos Soc. 2019;94: 1143–1160. doi:10.1111/brv.12496

4. Lindström J. Early development and fitness in birds and mammals. Trends Ecol Evol. 1999;14: 343–348. doi:10.1016/s0169-5347(99)01639-0

5. Geiser F. Ontogeny and phylogeny of endothermy and torpor in mammals and birds. Comp Biochem Physiol A Mol Integr Physiol. 2008;150: 176–180. doi:10.1016/j.cbpa.2007.02.041

6.   Visser GH. Development of Temperature Regulation. Avian Growth and Development:

Evolution within the Altricial-Precocial Spectrum. Oxford: Oxford University Press; 1998.

pp. 117–156.

7.   Clutton-Brock TH. Parental Care in Birds and Mammals. The Evolution of Parental Care.

Princeton University Press; 1991. pp. 131–153.

8.   Starck JM, Ricklefs RE. Avian Growth and Development: Evolution Within the Altricial-

precocial Spectrum. Oxford University Press; 1998.

9.   Andreasson F, Nilsson J-Å, Nord A. Avian Reproduction in a Warming World. Front Ecol

Evol. 2020;8: 576331. doi: 10.3389/fevo.2020.576331

10. Sauve D, Friesen VL, Charmantier A. The Effects of Weather on Avian Growth and

Implications for Adaptation to Climate Change. Front Ecol Evol. 2021;9: 569741. doi:

10.3389/fevo.2021.569741

11. Mccarty JP, Winkler DW. Relative importance of environmental variables in determining

the growth off nestling Tree Swallows Tachycineta bicolor. Ibis. 1999;141: 286–296.

doi:10.1111/j.1474-919X.1999.tb07551.x

12. Cunningham SJ, Martin RO, Hojem CL, Hockey PAR. Temperatures in Excess of Critical

Thresholds Threaten Nestling Growth and Survival in A Rapidly-Warming Arid Savanna:

A Study of Common Fiscals. PLoS One. 2013;8: e74613.

doi:10.1371/journal.pone.0074613

13.   Marques-Santos F, Dingemanse NJ. Weather effects on nestling survival of great tits vary

according to the developmental stage. J Avian Biol. 2020;51: e02421.

doi:10.1111/jav.02421

14.   Dunn EH. The Timing of Endothermy in the Development of Altrical Birds. Ornithol Appl.

1975;77: 288–293. doi:10.2307/1366224

15.   Wingfield JC, Pérez JH, Krause JS, Word KR, González-Gómez PL, Lisovski S, et al. How

birds cope physiologically and behaviourally with extreme climatic events. Philos Trans R

Soc Lond B Biol Sci. 2017;372: 20160140. doi:10.1098/rstb.2016.0140

16.   Andreasson F, Nord A, Nilsson J-Å. Experimentally increased nest temperature affects

body temperature, growth and apparent survival in blue tit nestlings. J Avian Biol. 2018;49:

jav-01620. doi:10.1111/jav.01620

17.   Zaremba U, Golawski A, Kasprzykowski Z. Variation in growth patterns of Marsh Harrier

Circus aeruginosus nestlings: effects of hatching order, nestling subperiod, brood size and

weather conditions. J Ornithol. 2022;163: 457–467. doi:10.1007/s10336-022-01968-3

18.   Covas R, du Plessis MA, Doutrelant C. Helpers in colonial cooperatively breeding sociable

weavers Philetairus socius contribute to buffer the effects of adverse breeding conditions.

Behav Ecol Sociobiol. 2008;63: 103–112. doi:10.1007/s00265-008-0640-2

19.   Auer SK, Martin TE. Parental care mitigates carry-over effects of poor early conditions on

offspring growth. Behav Ecol. 2017;28: 1176–1182. doi:10.1093/beheco/arx082

20.   Clark AB, Wilson DS. Avian Breeding Adaptations: Hatching Asynchrony, Brood

Reduction, and Nest Failure. Q Rev Biol. 1981;56: 253–277.

21.   Forbes S, Glassey B, Thornton S, Earle L. The secondary adjustment of clutch size in red-

winged blackbirds (Agelaius phoeniceus). Behav Ecol Sociobiol. 2001;50: 37–44.

doi:10.1007/s002650100332

22.   Gibbons DW. Hatching Asynchrony Reduces Parental Investment in the Jackdaw. J Anim

Ecol. 1987;56: 403–414. doi:10.2307/5056

23.   Prinzinger R, Preßmar A, Schleucher E. Body temperature in birds. Comp Biochem Physiol

A Physiol. 1991;99: 499–506. doi:10.1016/0300-9629(91)90122-S

24.   Whitfield MC, Smit B, McKechnie AE, Wolf BO. Avian thermoregulation in the heat:

scaling of heat tolerance and evaporative cooling capacity in three southern African arid-

zone passerines. J Exp Biol. 2015;218: 1705–1714. doi:10.1242/jeb.121749

25.   McKechnie AE, Wolf BO. Climate change increases the likelihood of catastrophic avian

mortality events during extreme heat waves. Biol Lett. 2009;6: 253–256.

doi:10.1098/rsbl.2009.0702

26.   Griebel IA, Dawson RD. Predictors of nestling survival during harsh weather events in an

aerial insectivore, the Tree Swallow (Tachycineta bicolor). Can J Zool. 2019;97: 81–90.

doi:10.1139/cjz-2018-0070

27.   Mainwaring MC, Hartley IR. Local weather conditions have complex effects on the growth

of blue tit nestlings. J Therm Biol. 2016;60: 12–19. doi:10.1016/j.jtherbio.2016.05.005

28.   Hiraldo F, Veiga JP, Máñez M. Growth of nestling black kites Milvus migrans: effects of

hatching order, weather and season. J Zool. 1990;222: 197–214. doi:10.1111/j.1469-

7998.1990.tb05672.x

29.   D'Amelio PB, Ferreira AC, Fortuna R, Paquet M, Silva LR, Theron F, et al. Disentangling

climatic and nest predator impact on reproductive output reveals adverse high-temperature

effects regardless of helper number in an arid-region cooperative bird. Ecol Lett. 2022;25:

151–162. doi:10.1111/ele.13913

30.   Grew R, Ratz T, Richardson J, Smiseth PT. Parental care buffers against effects of ambient

temperature on offspring performance in an insect. Behav Ecol. 2019;30: 1443–1450.

doi:10.1093/beheco/arz100

31.   Laubach ZM, Murray EJ, Hoke KL, Safran RJ, Perng W. A biologist's guide to model

selection and causal inference. Proc R Soc Lond B Biol Sci. 2021;288: 20202815.

doi:10.1098/rspb.2020.2815

32.   Brown MB, Brown CR. Barn Swallow (Hirundo rustica), version 1.0. Birds of the World.

2020 [cited 8 Mar 2023]. doi:10.2173/bow.barswa.01

33.   Saino N, Incagli M, Martinelli R, Ambrosini R, Møller AP. Immunity, Growth and Begging

Behaviour of Nestling Barn Swallows Hirundo rustica in Relation to Hatching Order. J

Avian Biol. 2001;32: 263–270. doi: 10.1111/j.0908-8857.2001.320309.x

34.   Saino N, Ambrosini R, Martinelli R, Calza S, Møller AP, Pilastro A. Offspring sexual

dimorphism and sex-allocation in relation to parental age and paternal ornamentation in the

barn swallow. Mol Ecol. 2002;11: 1533–1544. doi:10.1046/j.1365-294x.2002.01542.x

35. Fernaz JM, Schifferli L, Grüebler MU. Ageing nestling Barn Swallows *Hirundo rustica* : an

illustrated guide and cautionary comments. Ringing & Migration. 2012;27: 65–75.

doi:10.1080/03078698.2012.747587

36. Laubach ZM, Madden SA, Pardue A, Safran RJ. Does Parental Care Modify the

Association of Early-Life Size and Growth with Physiology? Am Nat. 2025;205: 469–484.

doi:10.1086/734993

37. Negro JJ, Chastin A, Bird DM. Effects of Short-Term Food Deprivation on Growth of

Hand-Reared American Kestrels. Ornithol Appl. 1994;96: 749–760. doi:10.2307/1369478

38. Streby HM, Peterson SM, Lehman JA, Kramer GR, Vernasco BJ, Andersen DE. Do

digestive contents confound body mass as a measure of relative condition in nestling

songbirds? Wildl Soc Bull. 2014;38: 305–310. doi:10.1002/wsb.406

39. Masoero G, Dumas MN, Martin JGA, Bize P. Trait-specific sensitive developmental

windows: Wing growth best integrates weather conditions encountered throughout the

development of nestling Alpine swifts. Ecol Evol. 2024;14: e11491.

doi:10.1002/ece3.11491

40. Madden SA, McDermott MT, Safran RJ. Visitation rate, but not foraging range, responds to

brood size manipulation in an aerial insectivore. Behav Ecol Sociobiol. 2022;76: 139.

doi:10.1007/s00265-022-03244-z

41. Newton-Fisher NE. Animal Behavior Pro. Mobile app. 2023.

42.   Hartig F. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. R package version 0.4.6. 2022. Available: https://CRAN.R-

project.org/package=DHARMa

43.   Houslay TM, Wilson AJ. Avoiding the misuse of BLUP in behavioural ecology. Behav

Ecol. 2017;28: 948–952. doi:10.1093/beheco/arx023

44.   Stoner D. Temperature and Growth Studies on the Barn Swallow. Auk. 1935;52: 400–407.

doi:10.2307/4077512

45.   Dunn EH. Age of Effective Homeothermy in Nestling Tree Swallows According to Brood

Size. Wilson Bull. 1979;91: 455–457.

46.   Marsh RL. Development of Temperature Regulation in Nestling Tree Swallows. Ornithol

Appl. 1980;82: 461–463. doi:10.2307/1367576

47.   Halsey LG. The reign of the p-value is over: what alternative analyses could we employ to

fill the power vacuum? Biol Lett. 2019;15: 20190174. doi:10.1098/rsbl.2019.0174

48.   Muff S, Nilsen EB, O'Hara RB, Nater CR. Rewriting results sections in the language of

evidence. Trends Ecol Evol. 2022;37: 203–210. doi:10.1016/j.tree.2021.10.009

49.   Fox J, Weisberg S. An R Companion to Applied Regression. 2019. Available:

https://www.john-fox.ca/Companion/

50.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria:

R Foundation for Statistical Computing; 2024. Available: https://www.R-project.org/

51.   Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to

the Tidyverse. J Open Source Softw. 2019;4: 1686. doi:10.21105/joss.01686

52.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using

lme4. J Stat Softw. 2015;67: 1–48. doi:10.18637/jss.v067.i01

53.   Canty A, Ripley B. boot: Bootstrap R (S-Plus) Functions. R package version 1.3-30. 2024.

Available: https://cran.r-project.org/web/packages/boot/boot.pdf

54.   Davison AC, Hinkley DV. Bootstrap Methods and Their Applications. Cambridge:

Cambridge University Press; 1997.

55.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed

Effects Models. J Stat Softw. 2017;82: 1–26. doi:10.18637/jss.v082.i13

56.   Bartoń K. MuMIn: Multi-Model Inference. R package version 1.48.4. 2024. Available:

https://CRAN.R-project.org/package=MuMIn

57.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag;

2016.

58.   Lüdecke D. ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.

J Open Source Softw. 2018;3: 772. doi:10.21105/joss.00772

59.   Kassambara A. ggpubr: "ggplot2" Based Publication Ready. Available: https://CRAN.R-

project.org/package=ggpubr

60.   Dyrcz A. Factors Affecting the Growth Rate of Nestling Great Reed Warblers and Reed

Warblers at Milicz, Poland. Ibis. 1974;116: 330–339. doi:10.1111/j.1474-

919X.1974.tb00128.x

61.   Dawson RD, Lawrie CC, O'Brien EL. The importance of microclimate variation in

determining size, growth and survival of avian offspring: experimental evidence from a

cavity nesting passerine. Oecologia. 2005;144: 499–507. doi:10.1007/s00442-005-0075-7

62.   Murphy MT. Nestling Eastern Kingbird Growth: Effects of Initial Size and Ambient

Temperature. Ecology. 1985;66: 162–170. doi:10.2307/1941316

63.   Bourne AR, Ridley AR, Spottiswoode CN, Cunningham SJ. Direct and indirect effects of

high temperatures on fledging success in a cooperatively breeding bird. Behav Ecol. 2021

Nov 1;32: 1212-1213. doi: 10.1093/beheco/arab087

64.   Salaberria C, Celis P, López-Rull I, Gil D. Effects of temperature and nest heat exposure on

nestling growth, dehydration and survival in a Mediterranean hole-nesting passerine. Ibis.

2014;156: 265–275. doi:10.1111/ibi.12121

65.   Larson ER, Eastwood JR, Buchanan KL, Bennett ATD, Berg ML. How does nest box

temperature affect nestling growth rate and breeding success in a parrot? Emu. 2015;115:

247–255. doi:10.1071/MU14081

66.   Bradbury RB, Wilson JD, Moorcroft D, Morris AJ, Perkins AJ. Habitat and weather are

weak correlates of nestling condition and growth rates of four UK farmland passerines. Ibis.

2003;145: 295–306. doi:10.1046/j.1474-919X.2003.00142.x

67.   Gilroy JJ, Anderson GQA, Grice PV, Vickery JA, Watts PN, Sutherland WJ. Foraging

768   habitat selection, diet and nestling condition in Yellow Wagtails Motacilla flava breeding

769   on arable farmland. Bird Study. 2009;56: 221–232. doi:10.1080/00063650902792080

770   68.   Geiser S, Arlettaz R, Schaub M. Impact of weather variation on feeding behaviour, nestling

771   growth and brood survival in Wrynecks Jynx torquilla. J Ornithol. 2008;149: 597–606.

772   doi:10.1007/s10336-008-0305-y

773   69.   Lauck KS, Ke A, Olimpi EM, Paredes D, Hood K, Phillips T, et al. Agriculture and hot

774   temperatures interactively erode the nest success of habitat generalist birds across the

775   United States. Science. 2023;382: 290–294. doi:10.1126/science.add2915

776   70.   Wheelwright NT, Freeman-Gallant CR, Mauck RA. Nestling Savannah Sparrows and Tree

777   Swallows differ in their sensitivity to weather. Ornithology. 2022;139: ukac032.

778   doi:10.1093/ornithology/ukac032

779   71.   Colombo MA, Jauregui A, Segura LN. Weather influenced nestling growth of an

780   insectivorous but not a granivorous grassland passerine in Argentina. Avian Res. 2024;15:

781   100173. doi:10.1016/j.avrs.2024.100173

782   72.   Dawson WR, Evans FC. Relation of Growth and Development to Temperature Regulation

783   in Nestling Vesper Sparrows. Ornithol Appl. 1960;62: 329–340. doi:10.2307/1365163

784   73.   Diehl JN, Alton LA, White CR, Peters A. Thermoregulatory strategies of songbird nestlings

785   reveal limited capacity for cooling and high risk of dehydration. J Therm Biol. 2023;117:

786   103707. doi:10.1016/j.jtherbio.2023.103707

74.   Tinbergen JM, Verhulst S. A fixed energetic ceiling to parental effort in the great tit? J

Anim Ecol. 2000;69: 323–334. doi:10.1046/j.1365-2656.2000.00395.x

75.   Weathers WW. Scaling nestling energy requirements. Ibis. 1992;134: 142–153.

doi:10.1111/j.1474-919X.1992.tb08391.x

76.   Węgrzyn E, Węgrzyn W, Leniowski K. Hatching asynchrony as a parental reproductive

strategy in birds: a review of causes and consequences. J Ornithol. 2023;164: 477–497.

doi:10.1007/s10336-023-02066-8

77.   Wetzel DP, Westneat DF. Parental Care Syndromes in House Sparrows: Positive

Covariance Between Provisioning and Defense Linked to Parent Identity. Ethology.

2014;120: 249–257. doi:10.1111/eth.12198

78.   Teunissen N, Kingma SA, Peters A. Nest defence and offspring provisioning in a

cooperative bird: individual subordinates vary in total contribution, but no division of tasks

among breeders and subordinates. Behav Ecol Sociobiol. 2020;74: 94. doi:10.1007/s00265-

020-02877-2

79.   Johnson EJ, Best LB. Factors Affecting Feeding and Brooding of Gray Catbird Nestlings.

Auk. 1982;99: 148–156. doi:10.2307/4086031

80.   Rytkönen S, Koivula K, Orell M. Patterns of Per-Brood and Per-Offspring Provisioning

Efforts in the Willow Tit Parus montanus. J Avian Biol. 1996;27: 21–30.

doi:10.2307/3676957

81.   Riddell EA, Iknayan KJ, Wolf BO, Sinervo B, Beissinger SR. Cooling requirements fueled

807   the collapse of a desert bird community from climate change. Proc Natl Acad Sci U S A.

808   2019;116: 21609–21615. doi:10.1073/pnas.1908791116

809   82.   Cunningham SJ, Gardner JL, Martin RO. Opportunity costs and the response of birds and

810   mammals to climate warming. Front Ecol Environ. 2021;19: 300–307.

811   doi:10.1002/fee.2324

812   83.   du Plessis KL, Martin RO, Hockey PAR, Cunningham SJ, Ridley AR. The costs of keeping

813   cool in a warming world: implications of high temperatures for foraging, thermoregulation

814   and body condition of an arid-zone bird. Glob Chang Biol. 2012;18: 3063–3070.

815   doi:10.1111/j.1365-2486.2012.02778.x

816   84.   Bailey LD, van de Pol M. Tackling extremes: challenges for ecological and evolutionary

817   research on extreme climatic events. J Anim Ecol. 2016;85: 85–96. doi:10.1111/1365-

818   2656.12451

819   85.   Lawson CR, Vindenes Y, Bailey L, van de Pol M. Environmental variation and population

820   responses to global change. Ecol Lett. 2015;18: 724–736. doi:10.1111/ele.12437

821   86.   Kärkkäinen T, Teerikorpi P, Schuett W, Stier A, Laaksonen T. Interplays between pre- and

822   post-natal environments affect early-life mortality, body mass and telomere dynamics in the

823   wild. J Exp Biol. 2021;224: jeb231290. doi:10.1242/jeb.231290

Supporting information

S1 Fig. A directed acyclic graph displaying hypothesized causal relationships investigated

in question one: does the effect of temperature on nestling mass differ when the exposure is

assessed during early versus late development? The explanatory and response variables have

black boxes, while confounders, precision covariates, and random intercepts have gray boxes.

Below the DAG are the formulas for the linear mixed models we ran separately for each

temperature variable (minimum, maximum, and variability).

S2 Fig. A directed acyclic graph displaying hypothesized causal relationships investigated

in question two: does the effect of temperature on nestling mass depend on whether a

nestling is the smallest in the brood? The explanatory and response variables have black boxes,

while confounders, precision covariates, and random intercepts have gray boxes. Below the

DAG is the formula for the linear mixed models we ran separately for each temperature variable

(minimum, maximum, and variability).

S3 Fig. A directed acyclic graph displaying hypothesized causal relationships investigated

in question three: does the effect of temperature on nestling mass before fledging depend

on the average amount of feeding provided by parents? The explanatory and response

variables have black boxes, while confounders, precision covariates, and random intercepts have

gray boxes. Below the DAG is the formula for the linear mixed models we ran separately for

each temperature variable (minimum, maximum, and variability).

S4 Fig. Best unbiased linear predictions (BLUPs) for the feeding rate (visits/hour) at each

nest across development in wild barn swallows in Boulder County, CO. A boxplot of feeding

BLUPs, measured in visits per hour, is provided for each of the three levels of parental care

('low,' 'med,' 'high') used for stratified analyses in question three.

S5 Fig. Daily temperatures recorded at wild barn swallow nests in Boulder County, CO.

Daily minimum temperature (a), maximum temperature (b), and temperature variability

(interquartile range) (c) in degrees Celsius recorded by Govee thermometers near each barn

swallow nest during the nestling rearing period. Each color corresponds to one of seven breeding

sites. Lines connect daily measures for each individual nest.

S1 Table. Ethogram of barn swallow parental care behaviors. Each documented behavior has

a corresponding location, type (state versus event), description, and developmental context. This

ethogram was used for parental care observations.

S2 Table. Background characteristics of wild nestling barn swallows in Boulder County,

CO. Total number of feeding visits, nestling mass, and nestling wing length are separately

estimated for three time points (days three to four, days eight to nine, and days 11-13).

Temperature variability is defined as the interquartile range. The table provides the sample size

for each variable at the level at which it was collected (n), as well as the mean, standard

deviation (SD), minimum, and maximum.

S3 Table. Associations of three temperature variables in early and late development (before or after six days post-hatch) with nestling mass for wild barns swallows in Boulder County, CO. Temperature variability is defined as the interquartile range. For each temperature variable, results are provided for unadjusted and adjusted models. Sample size for each developmental stage is provided in the header (n). For each model, the table provides the effect size and 95% confidence interval for temperature effects (β (95% CI)), and the corresponding degrees of freedom, t-value, and p-value from a two-tailed t-test using a Satterthwaite degree of freedom estimation.

S4 Table. Associations of three temperature variables with nestling mass, assessed in separate models stratified by relative nestling size at days eight to nine measure (smallest vs. other), for wild barn swallows in Boulder County, CO. Temperature variability is defined as the interquartile range. For each temperature variable, results are provided for unadjusted and adjusted models. Sample size for each stratum is provided in the header (n). For each model, the table provides the effect size and 95% confidence interval for temperature effects (β (95% CI)), and the corresponding degrees of freedom, t-value, and p-value from a two-tailed t-test using a Satterthwaite degree of freedom estimation.

S5 Table. Associations of three temperature variables with nestling mass, assessed in separate models stratified by levels of parental feeding, for wild barn swallows in Boulder County, CO. Temperature variability is defined as the interquartile range. For each temperature

890   variable, results are provided for unadjusted and adjusted models. Sample size for each stratum is

891   provided in the header (n). For each model, the table provides the effect size and 95% confidence

892   interval for temperature effects (β (95% CI)), and the corresponding degrees of freedom, t-value,

893   and p-value from a two-tailed t-test using a Satterthwaite degree of freedom estimation.

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5