Trichoderma afroharzianum behaves differently with respect to the host transcriptome and microbial communities under varying iron availability in pea plants

Ahmad H. Kabir[1*], Asha Thapa[2], Shifat Ara Saiful[3], Shyamal K. Talukder[4,5]

[1]Department of Biology, Lamar University, Beaumont, TX 77705, USA
[2]Evolution, Behavior and Ecology Program, Department of Biology, Boston University, MA 02215, USA
[3]Biology Program, School of Sciences, University of Louisiana at Monroe, LA 71209, USA
[4]Texas A&M AgriLife Research Center, Beaumont, TX, 77713, USA
[5]Department of Soil and Crop Sciences, Texas A&M University, College Station, Texas, USA

[*]Correspondence: akabir2@lamar.edu

ABSTRACT

Trichoderma afroharzianum T22 is recognized for promoting plant growth and stress resilience. However, it remains unclear how its effects vary with Fe nutritional status in pea plants. In this study, we investigated the impact of T22 on pea grown under Fe-sufficient and Fe-deficient conditions by integrating physiological measurements with transcriptomic and microbiome analyses. Our results reveal that the benefits of T22 are highly context dependent, providing significant improvements in photosynthesis and Fe/N accumulation under Fe deficiency but minimal effects under sufficiency, consistent with conditional mutualism theory. RNA-seq analysis revealed that T22 inoculation induced 262 differentially expressed genes (DEGs) under Fe deficiency, whereas under Fe sufficiency, 555 DEGs were identified, largely linked to basal metabolic functions, suggesting colonization costs rather than adaptive responses. Particularly, T22 inoculation upregulated symbiosis-related genes (*Nodule-specific GRPs*, *Major facilitator, sugar transporter-like*), Fe transporters (*NRAMPs*, *HMAs*), and redox-associated genes (*Glutathione S-transferase*, *Glutathione peroxidase*) in the roots under Fe shortage, reflecting a coordinated response to enhance nutrient acquisition and stress tolerance. Microbiome profiling showed that under Fe deficiency, T22 reshaped the root-associated community by enriching *Pseudomonas*, *Variovorax*, *Mitsuaria*, and *Acidovorax*, along with increased fungal richness and diversity, whereas under Fe sufficiency, restructuring was minimal and largely limited to the enrichment of *Pararhizobium*. These enriched bacterial taxa may act as 'helpers' to T22 by providing complementary thereby amplifying its beneficial effects under Fe deficiency. These findings demonstrate that T22 acts in a context-dependent manner, exerting strong beneficial effects under Fe deficiency while having little influence when Fe is sufficient. These findings may establish T22 as a targeted bioinoculant for Fe-deficient soil and open avenues for designing microbial consortia to enhance legume resilience and sustainability in future agriculture.

Keywords: Helper microbes**,** Microbiome, Mutualism, Nodulation, Siderophores, Transcriptome.

1. Introduction

Iron (Fe) deficiency is a pervasive constraint in legume crops grown on calcareous and alkaline soils, where high pH and carbonate content reduce Fe solubility and drive leaf chlorosis and yield losses (Merry et al., 2022; Assefa et al., 2020). Fe is indispensable for photosynthesis, respiration, and redox homeostasis, and therefore for biomass production and crop quality (Briat 2015). In legumes, Fe nutrition carries additional significance because nodulation and symbiotic N☐ fixation require abundant Fe for nitrogenase, leghemoglobin, and other Fe cofactors (Liu et al., 2023; Brear et al., 2013). Consequently, Fe deficiency impairs nodule initiation and function and reduces whole-plant nitrogen status (Brear et al., 2013).

To acquire scarcely soluble Fe, dicots including legume deploy the Strategy I reduction mechanism acidifying the rhizosphere, reducing Fe(III) to Fe(II), and importing Fe(II), while grasses use Strategy II chelation with phytosiderophores; convergence and crosstalk between these strategies are increasingly recognized (Jeong and Guerinot 2009; Staiger, 2002). However, due to the lack of popular high-yield varieties with tolerance strategies, it remains a persistent issue for legume and other crops. Fe-deficiency-induced chlorosis management in crops typically relies on genotype selection and soil/seed treatments (Liberal et al., 2020). Yet, durable, biologically based solutions that enhance Fe bioavailability and maintain root–nodule performance remain a priority.

Harnessing the plant-associated microbiome could offer a promising alternative to conventional approaches (Compant et al., 2025). *Trichoderma* spp. are ubiquitous root-associated fungi widely used as biologicals that stimulate plant growth and resilience and changes to the rhizosphere microbiome (Thapa et al., 2025; Geng et al., 2025). Several *Trichoderma* strains produce and/or stimulate siderophores and organic acids that can increase Fe solubility for alleviating chlorosis induced by Fe deficiency (Thapa et al., 2025; Kabir and Bennetzen 2024; Tyśkiewicz et al., 2022). In general, plant–microbe mutualisms are context dependent; the magnitude and nature of the benefits vary with environmental conditions, host genotype, and microbial community composition (Thapa et al., 2025; Brown et al., 2020). However, these interactions are not static; under nutrient-rich conditions, the cost of maintaining the symbiont may outweigh its benefits, shifting the relationship away from mutualism (Kiers et al., 2003). Environmental stresses can further modulate mutualistic outcomes, often enhancing the reliance on microbial partners for nutrient acquisition and stress mitigation (Lamers et al., 2020; de Zelicourt et al., 2013). Understanding this context dependency is crucial for designing microbial inoculants that deliver consistent benefits under variable field conditions.

Despite the promise of *Trichoderma*, it remains unclear how Fe availability modulates the behavior of *Trichoderma afroharzianum* T22. Here we explicitly ask how T22 behaves differently in the presence versus absence of Fe deficiency at the root-soil interface. Using peas (*Pisum sativum*) as the host, we studied the differential effect of T22 on physiological responses paired with host transcriptomics and microbial community profiles in the rhizosphere of pea plants under Fe-sufficient and Fe-deficient conditions. This approach provides a systems view of how a well-characterized beneficial fungus recalibrates pea performance across Fe regimes.

2. Materials and methods

2.1. Plant cultivation and growth conditions

Garden pea (Sugar Snap variety) seeds were surface disinfected by soaking in a 1% sodium hypochlorite solution for 5 min, followed by thorough rinsing with sterile distilled water. Germination was performed in trays placed in an incubator at 25 °C for 2d Seedlings were transplanted into pots containing a 1:3 mixture of field soil and SunGro potting mix and grown in the growth chamber in two growth conditions (Fe-sufficient and Fe-deficient supplemented with 4.5 g $CaCO_3$ and 7.5 g $NaHCO_3$) with or without the inoculation of *Trichoderma afroharzianum* T22 (1×10^9 cfu/gram). The experiment was conducted following a randomized complete block design (RCBD) with three replicates per treatment for up to 5 weeks after transplantation. Plants were grown in a controlled growth chamber under a 10 h light/14 h dark photoperiod, with a light intensity of 250 µmol m^{-2} s^{-1} at ~25 °C.

2.2. Determination of morpho-physiological attributes

The plant height was measured from below ground root to the highest leaf using a measuring tape immediately before harvesting the plants. The aboveground portions of the plant were harvested, and fresh biomass weight was measured using a scale. Nodules were counted manually on the roots of pea plants after carefully washing off adhering soil under running water. SPAD score, an indicator of chlorophyll content in fully expanded leaves, was measured using a handheld SPAD meter (AMTAST, United States). Also, Fv/Fm (maximal photochemical efficiency of PSII) was measured on the uppermost fully expanded leaves after dark adaptation for 1 hour, using a FluorPen FP 110 (Photon Systems Instruments, Czech Republic).

2.3. Fe and N analysis of plant tissues

After harvest, the roots of pea plants were carefully excised and rinsed under running tap water to remove adhering soil. The roots were then immersed in 0.1 mM $CaSO_4$ for 10 min to displace surface ions, followed by a final rinse with deionized water. Roots and leaves were oven-dried at 75 °C for three days, and the dried samples were submitted to the Agricultural & Environmental Services Laboratories, University of Georgia, for iron quantification using inductively coupled plasma mass spectrometry (ICP-MS). Fe and N concentrations were determined by comparing ion signal intensities with those of certified calibration standards.
.

2.4. Molecular detection of T22 colonization in the roots

Colonization of T22 in pea root tips was assessed by quantifying the relative abundance of the T22 marker gene *TaAOX1* (L-amino acid oxidase) using DNA-based qPCR (Kabir and Bennetzen, 2024). Root tips (~0.2 g) were washed twice in sterile phosphate-buffered saline, vortexed briefly, rinsed twice with sterile water, and subjected to DNA extraction using the CTAB method (Clarke, 2009). DNA was quantified (NanoDrop ND-1000, Wilmington, USA) and normalized across samples. qPCR was performed on a CFX96 Touch Real-Time PCR System (Bio-Rad, USA) with *TaAOX1*-specific primers (forward: 5′-GTC GGT AGC TGA AAG GGG AT-3′; reverse: 5′-ATT AGG CCG GAA ACA CC-3′). All reactions were run with iTaq Universal SYBR Green Supermix (Bio-Rad, USA), and *PsGAPDH* was used as the internal control for relative quantification by the 2−ΔΔCT method (Livak and Schmittgen, 2001).

2.5. Determination of siderophore levels in the rhizosphere

Siderophore content in rhizosphere soil was quantified using the chrome azurol S (CAS) assay (Himpsl and Mobley, 2019). Briefly, soil samples were homogenized in 80% methanol and centrifuged at 10,000 rpm for 15 min. The supernatant (500 µL) was mixed with 500 µL CAS solution and incubated at room temperature for 5 min. Absorbance was recorded at 630 nm, with 1 mL CAS reagent serving as the reference. Siderophore units were calculated as %SU = [(Ar − As)/Ar] × 100, where Ar is the absorbance of the reference and As is the absorbance of the sample.

2.6. RNA-seq analysis in the roots

RNA-seq was performed in the root samples. Root samples were washed twice in sterile phosphate-buffered saline (PBS) by vortexing for 10 s, followed by two rinses with sterile water to remove surface contaminants. Samples were ground in liquid nitrogen with a pre-cooled mortar and pestle, and the resulting powder was used for RNA extraction with the SV Total RNA Isolation System (Promega, USA) according to the manufacturer's protocol. RNA integrity (RIN > 8) was confirmed, and 1 µg of total RNA per sample was used for library preparation with the KAPA HiFi HotStart Library Amplification Kit (Kapa Biosystems, USA). Libraries were sequenced on an Illumina NextSeq SE 100 bp platform, yielding >80% high-quality reads (Phred > 33 per sample).

Adaptor and low-quality sequences were trimmed using Trimmomatic v0.38 (Bolger et al., 2014). Filtered reads were aligned to the pea reference genome (Yang et al., 2022) with HISAT2 v2.0.5 (Kim et al., 2015). Gene-level read counts were obtained using HTSeq (Anders et al., 2014). Differentially expressed genes (DEGs) were identified with the Limma R package v3.19 (Law et al., 2016) following normalization with the FPKM method.

DEGs were defined as genes with log2FC ≥ 1 or ≤ −1 and false discovery rate (FDR) ≤ 0.05. Heatmaps were generated using the *pheatmap* R package. Pea gene ontology annotations were retrieved from Ensembl Plants and enrichment analysis was performed with ShinyGO v0.80.

2.7. Amplicon sequencing of 16S and ITS microbial communities

Microbial communities in pea roots were analyzed by amplicon sequencing of the 16S rRNA gene (bacteria) and ITS region (fungi). Roots were cleaned by vortexing in sterile phosphate-buffered saline (PBS) for 10 s, followed by two rinses with sterile water to remove surface contaminants. DNA was extracted from ~0.2 g of root tissue using the Wizard® Genomic DNA Purification Kit (Promega, USA), including RNase and Proteinase K treatments to remove RNA and protein impurities. The 16S rRNA gene library was prepared with primers 341F (CCT ACG GGNGGC WGC AG) and 805R (GAC TAC HVGGG TAT CTA ATC C), and sequencing was performed on an Illumina NovaSeq 6000 (PE250). Raw reads were quality trimmed with cutadapt (Martin, 2011) and processed using the DADA2 pipeline (Callahan et al., 2016). Sequences of mitochondrial and chloroplast origin were removed. Taxonomic classification of amplicon sequence variants (ASVs) was performed with the UNITE reference database (Nilsson et al., 2019) for ITS and Silva v138 (Quast et al., 2013) for 16S. Community structure was assessed by Principal Coordinate Analysis (PCoA) based on Bray–Curtis dissimilarity of Hellinger-transformed counts using the *phyloseq* and *vegan* R packages. Taxa abundance

differences were tested with the non-parametric Kruskal–Wallis test. Alpha diversity, beta diversity, relative abundance, and principal component analyses were conducted using *phyloseq* (McMurdie and Holmes, 2013), with significance determined at $p < 0.05$.

2.8. Statistical analysis

All measurements were conducted with a minimum of three independent biological replicates per treatment. Statistical significance ($p < 0.05$) was determined using Student's *t*-test in Microsoft Excel, and data visualization was performed with the ggplot2 package in R. Furthermore, a visual illustration was prepared using BioRender.

3. Results

3.1. Growth, photosynthesis, and nutrient status

Under Fe sufficiency, T22 had no significant effect on any measured trait (Fig. 1A–H, ns throughout). Plant height and biomass trended slightly upward with T22 (Fig. 1A–B, ns), and leaf SPAD and Fv/Fm values were comparable between −T22 and +T22 (Fig. 1C–D, ns). Root and leaf Fe concentrations were likewise unchanged (Fig. 1E–F, ns), and tissue N in roots and leaves remained statistically indistinguishable (Fig. 1G–H, ns). In contrast, under Fe deficiency, T22 conferred significant improvements across scales. T22-inoculated plants were taller and accumulated more biomass than uninoculated controls (Fig. 1A–B, significant). Photosynthetic performance recovered with T22: SPAD values increased and Fv/Fm rose relative to −T22 (Fig. 1C–D, significant). T22 also elevated Fe status, raising Fe concentrations in both roots and leaves (Fig. 1E–F, significant). Finally, T22 improved plant nitrogen status under Fe stress, increasing N% in roots and leaves compared with −T22 (Fig. 1G–H, significant). Together, these patterns show that T22's benefits are Fe-contingent—physiology and nutrient status are stabilized specifically under Fe limitation, while remaining non-significantly changed when Fe is sufficient.

3.2. T22 abundance, nodulation, and siderophore activity

Under Fe sufficiency, T22 inoculation significantly increased the relative abundance of T22 in the rhizosphere (Fig. 2A, *), while effects on nodules per plant and siderophore production were not significant (Fig. 2B–C, ns). In contrast, under Fe deficiency, all three responses improved with T22: T22 abundance increased significantly (Fig. 2A, **), nodulation rose significantly (Fig. 2B, **), and siderophore production was significantly higher than −T22 (Fig. 2C, *). These data indicate that T22 colonization and its Fe-scavenging output are Fe contingent—minimal under sufficiency but enhanced under Fe deficiency, where they are accompanied by increased nodulation.

3.3. Differential gene expression

Across conditions, T22 reshaped the pea root transcriptome in an Fe-dependent manner (Fig. 3A–C). Under Fe sufficiency, we detected 555 DEGs relative to −T22 (225 upregulated, 330 downregulated; FDR < 0.05). Under Fe deficiency, fewer genes responded (262 DEGs; 95 up, 167 down; FDR < 0.05). Although the DEG count was larger in Fe sufficiency, the direction and magnitude of key Fe- and stress-associated genes shifted most clearly under Fe deficiency: NRAMP transporters and HMA-domain metal-handling genes, a major facilitator/sugar-transporter–like gene, small auxin-up RNA (SAUR), phenylalanine ammonia-lyase (PAL), UDP-glucosyltransferase (UGT), and glutathione S-transferase (GST) all showed strong positive log FC with T22 specifically in Fe-deficient roots, whereas several of these were unchanged or

reduced (not significant) under Fe sufficiency. Glutathione peroxidase displayed a weaker/negative shift under Fe sufficiency. Together, these patterns indicate that T22 preferentially activates Fe transport, hormone signaling, phenylpropanoid metabolism, and detoxification modules under Fe limitation, with many of the same loci not significantly induced when Fe is adequate (Fig. 3C).

3.4. GO enrichment highlights distinct functional pathways

GO over-representation recapitulated these contrasts (Fig. 4A–B). In Fe-sufficient plants, T22-responsive genes were significantly enriched for molecular functions related to nucleotide/ATP/adenyl-ribonucleotide binding, copper/anion/ion binding, transferase activity, and membrane terms (FDR < 0.05), while structural and transport categories were not significant. Under Fe deficiency, the program shifted: the top signals were cell-wall–related processes—structural constituent of cell wall (largest fold enrichment, >100), plant-type cell wall organization/biogenesis, and cell wall organization/biogenesis—accompanied by inorganic-ion transmembrane transporter activity, oxidoreductase activity, and metal/cation binding (all significant, FDR < 0.05). GO terms outside these sets were not significantly enriched. Overall, T22 elicits a condition-specific transcriptional response, emphasizing cell-wall remodeling and ion transport under Fe stress, but nucleotide-binding and general metabolic functions when Fe is sufficient (Fig. 4).

3.5. Changes in bacterial community

Under Fe sufficiency, T22 did not alter alpha diversity: both Observed ASVs and Simpson index were non-significant (Fig. 5A; $P = 0.61$ and 0.30, respectively). Compositionally, class-level differences were significant for Actinobacteria and Alphaproteobacteria Fig. 5B), while the remaining classes were non-significant. At the genus level, *Variovorax* and *Pararhizobium* differed significantly between –T22 and +T22, whereas *Thermomonas*, *Shinella*, *Pseudomonas*, *Mycobacterium*, *Methylophilus*, *Massilia*, *Enterobacter*, and *Devosia* were non-significant (Fig. 5C). Under Fe deficiency, observed richness remained non-significant, but community evenness increased with T22: the Simpson index was significantly higher for +T22 (Fig. 5D; $P = 0.008$). At the class level, changes were modest and non-significant (Fig. 5E). In contrast, several genera showed significant shifts with T22, including *Variovorax*, *Shinella*, *Pseudomonas*, *Mitsuaria*, and *Acidovorax*. In addition, other genera (e.g., Pararhizobium, *Novosphingobium*, *Methylophilus*, *Flavobacterium*, *Comamonas*, *Cellvibrio*) were non-significant (Fig. 5F). Collectively, T22 exerted a stronger and more resolved impact on bacterial composition under Fe deficiency, accompanied by higher evenness (Fig. 5F).

3.6. Changes in the fungal community

T22 increased observed fungal richness under both Fe regimes (Fig. 6A-D; significant in Fe-sufficient, $P = 0.01$; significant in Fe-deficient, $P = 0.04$), while Simpson indices were non-significant (Fig. 6A-D; $P = 0.20$ and 0.40). Under Fe sufficiency, class-level composition showed a significant response of Sordariomycetes (Fig. 6B), with other classes non-significant. At the genus level, *Haptocillium* and *Fusarium* differed significantly between treatments, whereas *Pseudallescheria*, *Phialophora*, *Paecilomyces*, *Lasiobolidium*, *Halosphaeria*, *Cladosporium*, Chaetomium, and *Ascospheara* were non-significant (Fig. 6C). Under Fe deficiency, class-level shifts were non-significant (Fig. 6E), but several genera were significantly altered with T22 which includes *Pseudallescheria*, *Fusarium*, and *Ascospheara*, while others

(e.g., *Thermothielavioides*, *Paecilomyces*, *Lasiosphaeridium*, *Humicola*, *Haptocillium*, *Halosphaeria*, *Chaetomium*) were non-significant (Fig. 6F). Overall, T22 consistently elevated fungal richness and selectively restructured key genera, with clearer genus-level responses under Fe deficiency (Fig. 6F).

4. Discussion

4.1. Context-dependent benefits of Trichoderma afroharzianum T22

Our study demonstrates that the benefits of T22 are not universal, but highly dependent on Fe nutritional status. Under Fe sufficiency, T22 colonized pea roots but did not significantly affect plant health parameters or Fe and N concentrations. By contrast, under Fe deficiency, T22 showed efficiency in improving these physiological and nutrient traits, including recovery of photosynthesis and Fe/N accumulation. This confirms the hypothesis that *Trichoderma*-plant interactions are conditional mutualisms, where benefits are realized primarily under nutrient stress (Donoso et al., 2008; Kiers et al., 2003). Similar context-dependent outcomes have been observed in other legumes. For instance, soybean grown in calcareous soils showed greater reliance on microbial partners under Fe stress, but minimal response when Fe was sufficient (Roriz et al., 2023; Assefa et al., 2020). These findings underscore that the symbiotic relationship is maintained by a cost–benefit balance, where plants minimize investment in microbial interactions when internal Fe status is adequate.

Although the functional effects of T22 were dependent on Fe availability, our data indicate that T22 colonization itself was not constrained by Fe status. This suggests that colonization is a context-independent trait, reflecting the strong colonization capacity of *Trichoderma* to penetrate and persist in root tissues regardless of nutrient status (Compant et al., 2025; Tyśkiewicz et al., 2022). This separation between colonization capacity and functional outcomes emphasizes that while *Trichoderma* can establish robustly in diverse conditions, the ecological and physiological benefits to the host depend on stress context. Similar findings have been reported in sorghum and peas, where T22 colonized roots broadly but only conferred growth or nutrient benefits under stress conditions (Kabir & Bennetzen, 2024; Thapa et al., 2025).

4.2. Fe deficiency unlocks the functional potential of T22

While T22 colonization occurred regardless of Fe availability, its functional consequences were strongly amplified in Fe-limited plants. Specifically, T22 colonization, siderophore release, and nodulation all increased significantly under Fe deficiency but remained unchanged under sufficiency. This suggests that the root–soil environment under Fe limitation actively promotes microbial functions that mobilize Fe. *Trichoderma* is well documented to produce siderophores and organic acids, which solubilize otherwise insoluble Fe in alkaline soils (Kabir and Bennetzen, 2024; Sood et al., 2020). Kabir and Bennetzen (2024) demonstrated that T22 enhances siderophore activity in sorghum, directly increasing Fe uptake and conferring Fe-deficiency tolerance. Our pea data corroborate this mechanism, as T22-inoculated plants displayed elevated siderophore activity and restored Fe concentrations in roots and leaves. Fe is indispensable for rhizobial symbiosis, since both nitrogenase and leghemoglobin require high Fe inputs (Lindström and Mousav, 2020; Brear et al., 2013). By mobilizing Fe, T22 may indirectly sustain nodulation, which in turn improves N acquisition supported by higher N concentrations in roots and shoots. This interaction represents a tripartite synergy among *Trichoderma*,

Rhizobium, and the host, consistent with recent findings that T22 induces flavonoid-driven nodulation signals in pea under alkaline stress (Thapa et al., 2025). Moreover, studies suggest that plant growth and transcriptional responses to T22 are strongly influenced by substrate type and plant genotype (Thapa et al., 2025; Schmidt et al., 2020).

4.3. T22 Rewires root transcriptomes for Fe stress management

RNA-seq analysis revealed a distinct and Fe-dependent transcriptional reprogramming in pea roots colonized by *Trichoderma afroharzianum* T22. Under Fe deficiency, T22 strongly upregulated genes central to Fe uptake and transport, including *NRAMP* family members and heavy metal–associated (HMA) domain proteins. Similar transcriptional activation of NRAMPs has been reported in sorghum inoculated with T22, where enhanced Fe transport alleviated chlorosis symptoms (Kabir & Bennetzen, 2024). This activation aligns with the enhanced siderophore-mediated Fe mobilization observed in the rhizosphere, suggesting that T22 inoculation primes both fungal- and plant-mediated Fe acquisition strategies (Tyśkiewicz et al., 2022; Geng et al., 2025).

In addition, genes associated with phenylpropanoid metabolism (e.g., *phenylalanine ammonia-lyase*, PAL), hormone signaling (*small auxin-up RNA*, SAUR), and antioxidant defenses (*glutathione S-transferase*, GST; *glutathione peroxidase*, GPX) were activated. Phenylpropanoid and flavonoid pathways are widely implicated in signaling and microbial recruitment in legume roots (Thapa et al., 2025), while PAL-mediated phenylpropanoid flux has been shown to enhance both defense and symbiotic signaling under abiotic stress (Yadav et al., 2020). Similarly, GSTs and GPXs are well-established mediators of redox homeostasis in plants colonized by beneficial fungi, buffering oxidative stress generated during root colonization (Casatejada et al., 2023). The combined activation of these pathways suggests that T22 integrates nutrient uptake with redox regulation and stress signaling. Gene ontology enrichment further confirmed the preferential induction of pathways associated with cell-wall remodeling, oxidoreductase activity, and ion transport specifically under Fe limitation (Fig. 4B). These processes are consistent with reports that root cell-wall dynamics and oxidoreductase activity are central components of Strategy I Fe uptake responses in dicots (Hantzis et al., 2018; Jeong & Guerinot, 2009). Similarly, the induction of phenylpropanoid pathways and flavonoid metabolism has been widely linked to rhizosphere signaling and the recruitment of beneficial microbiota (Thapa et al., 2025). Upregulation of PAL and sugar transporter-like genes in our dataset suggests that T22 may promote enhanced root exudation, a mechanism that fosters microbial enrichment and nutrient exchange in the rhizosphere (Lei et al., 2025). Thus, T22 appears to modulate both direct nutrient transport and indirect exudate-mediated interactions to support Fe stress resilience.

In contrast, under Fe sufficiency, T22 inoculation triggered a greater number of DEGs (555 vs. 262 under deficiency), but these were largely associated with basal metabolic processes, such as nucleotide binding, protein turnover, and ion binding (Fig. 4A). These transcriptional changes may reflect the metabolic costs of colonization rather than an adaptive, stress-specific response. The lack of enrichment in Fe transport or redox-related pathways suggests that the host perceives T22 as an additional metabolic load rather than a symbiotic partner when Fe is abundant. This contrast highlights a hallmark of conditional mutualism: plants preferentially invest in transcriptional and metabolic shifts to support microbial partners only when external nutrient

availability is low (Blonde et al., 2025; Jones et al., 2019). Overall, these findings demonstrate that T22 functions as a molecular switch under Fe stress, activating targeted transcriptional programs that facilitate Fe uptake, redox homeostasis, and microbial recruitment.

4.4. Microbiome restructuring and selective enrichment of beneficial taxa

Microbiome profiling confirmed that *Trichoderma afroharzianum* T22 reshaped root-associated bacterial and fungal communities in a condition-specific manner. Under Fe deficiency, pea plants inoculated with T22 appear to actively recruit bacterial taxa with established roles in Fe mobilization, hormone signaling, and plant growth promotion. In particular, T22 selectively enriched *Pseudomonas, Mitsuaria, Variovorax,* and *Acidovorax*. These genera are well recognized for their roles in siderophore biosynthesis, indole-3-acetic acid (IAA) production, biofilm formation, and pathogen suppression (Compant et al., 2025). *Pseudomonas* species are widely reported to produce high-affinity siderophores such as pyoverdine, which mobilize Fe and improve Fe uptake under alkaline soils (Cornelis, 2010; Saha et al., 2016). *Variovorax* has been repeatedly identified as a keystone taxon in legume and cereal rhizospheres under nutrient stress, where it contributes to auxin metabolism, detoxification, and improved root architecture (Finkel et al., 2020; Sasse et al., 2018). *Mitsuaria* is increasingly associated with plant-beneficial consortia, particularly for phosphate solubilization and siderophore-mediated Fe acquisition (Zhang et al., 2019), while *Acidovorax* includes strains capable of producing siderophores and antifungal metabolites that enhance plant resilience (Mendes et al., 2011). The consistent enrichment of these taxa in Fe-starved, T22-inoculated plants strongly suggests that the fungus functions as a "microbial recruiter," promoting a rhizosphere microbiome specialized for Fe mobilization and nutrient cycling. On the fungal side, T22 consistently increased richness under both Fe conditions but induced stronger genus-level shifts under Fe deficiency, with enrichment of *Pseudallescheria, Fusarium,* and *Ascospheara* (Fig. 6F). While these genera may include opportunistic or weak pathogens, their enrichment under T22 inoculation likely reflects community restructuring that enhances microbial competition and antagonism, thereby limiting pathogen establishment (Abbas et al., 2022). Similar restructuring effects have been reported for *Trichoderma*-mediated rhizosphere modulation in cereals and legumes, where non-pathogenic fungal taxa occupy ecological niches to suppress disease outbreaks (López-Bucio et al., 2015; Harman et al., 2021).

In contrast, under Fe sufficiency, T22 exerted minimal influence on bacterial diversity, with only *Pararhizobium* showing significant enrichment. This limited restructuring corresponds with the weak physiological impact of T22 under Fe sufficiency. Although *Pararhizobium* may contribute to nodulation support, its enrichment in the absence of Fe stress likely reflects basal microbiome modulation without functional consequences for Fe acquisition. Taken together, these findings highlight that T22 acts as a microbial "hub" primarily under Fe deficiency, selectively enriching bacterial taxa that contribute to siderophore-mediated Fe cycling and nutrient mobilization (Fig. 7). Importantly, this restructuring is not a passive process but reflects a context-dependent reprogramming of the rhizosphere network, where T22 orchestrates microbial partnerships with complementary functions—including Fe scavenging (*Pseudomonas*), hormone modulation (*Variovorax*), and nutrient solubilization (*Mitsuaria, Acidovorax*) to sustain host nutrition and stress tolerance. These results align with the broader concept that plant-associated microbiomes are not static but are actively recruited by hosts and their microbial partners in response to nutrient limitations (Berendsen et al., 2012; Trivedi et al., 2020).

5. Conclusion

This study demonstrates that the benefits of T22 in pea are not universal, but strongly dependent on Fe nutritional status. While T22 colonization was observed under both Fe sufficiency and deficiency, its functional outcomes diverged sharply. Under Fe sufficiency, T22 had minimal effects on plant physiology, nutrient status, or microbial restructuring, indicating that colonization alone is not sufficient to drive host benefit. By contrast, under Fe deficiency, T22 significantly improved growth-related traits, photosynthesis, and Fe and N accumulation, underscoring the conditional nature of *Trichoderma*–plant mutualisms. T22 inoculation promoted siderophore release and sustained nodulation, supporting a tripartite synergy among *Trichoderma*, *Rhizobium*, and the host. Transcriptomic profiling revealed the activation of Fe uptake, phenylpropanoid metabolism, and antioxidant defenses processes. Simultaneously, microbiome profiling demonstrated that T22 restructured the rhizosphere community, selectively enriching *Pseudomonas, Variovorax, Mitsuaria,* and *Acidovorax* (Fig. 7). Together, our findings provide clear evidence that T22 functions as a molecular and microbial switch, conferring pronounced benefits only when plants face Fe starvation. Ultimately, these results establish a mechanistic foundation for deploying T22 in Fe-deficient or alkaline soils. More broadly, this work underscores the potential of Trichoderma-based inoculants as sustainable, microbiome-driven solutions to enhance crop resilience in nutrient-limited agroecosystems.

Author statement

We hereby certify that all authors have seen and approved the manuscript.

Declaration of Competing Interest

The authors have declared that no conflict of interest exists.

Data availability

Illumina sequencing data were submitted to NCBI under the following accession numbers: PRJNA1313627 (RNA-seq), PRJNA1314923 (16S) and PRJNA1314929 (ITS).

Acknowledgments

We express our gratitude to Louisiana Biomedical Research Network for the funding. This research was also supported by a startup grant from Lamar University.

Author contributions

AHK conceived the study, performed the molecular experiments, analyzed the data, performed bioinformatics analysis and drafted the manuscript. AT helped with RNA-seq data analysis and revised the manuscript. SAS assisted in amplicon data analysis and graphical representation. SKT interpreted the results and critically revised the manuscript.

REFERENCES

Abbas, A., Mubeen, M., Zheng, H., Sohail, M. A., Shakeel, Q., Solanki, M. K., Iftikhar, Y., Sharma, S., Kashyap, B.K., Hussain, S., Del Carmen Zuñiga Romano, M., Moya-Elizondo, E. A., Zhou, L., 2022. *Trichoderma* spp. Genes Involved in the Biocontrol Activity Against Rhizoctonia solani. Front. Microbiol. 13, 884469.

Anders, S., Pyl, P.T., Huber, W., 2015. HTSeq—a python framework to work with high throughput sequencing data. Bioinformatics 31(2), 166–169.

Assefa, T., Zhang, J., Chowda-Reddy, R.V., Moran Lauter, A.N., Singh, A., O'Rourke, J.A., Graham, M.A., Singh, A.K., 2020. Deconstructing the genetic architecture of iron deficiency chlorosis in soybean using genome-wide approaches. BMC Plant Biol. 20(1), 42.

Berendsen, R.L., Pieterse, C.M., Bakker, P.A., 2012. The rhizosphere microbiome and plant health. Trends Plant Sci. 17(8), 478–486.

Blonde, C., Caddeo, A., Nasser, W., Reverchon, S., Peyraud, R., Haichar, F.E.Z., 2025. New insights in metabolism modelling to decipher plant-microbe interactions. New Phytol. 246(4), 1485–1493.

Brear, E.M., Day, D.A., Smith, P.M., 2013. Iron: an essential micronutrient for the legume-rhizobium symbiosis. Front. Plant Sci. 4, 359.

Briat, J. F., Dubos, C., & Gaymard, F. (2015). Iron nutrition, biomass production, and plant product quality. Trends in plant science, 20(1), 33–40.

Brown, S.P., Grillo, M.A., Podowski, J.C., Heath, K.D., 2020. Soil origin and plant genotype structure distinct microbiome compartments in the model legume *Medicago truncatula*. Microbiome 8(1), 139.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nature Met. 13(7), 581–583.

Casatejada, A., Puerto-Galán, L., Pérez-Ruiz, J.M., Cejudo, F.J., 2023. The contribution of glutathione peroxidases to chloroplast redox homeostasis in Arabidopsis. Redox Biol. 63, 102731.

Compant, S., Cassan, F., Kostić, T., Johnson, L., Brader, G., Trognitz, F., Sessitsch, A., 2025. Harnessing the plant microbiome for sustainable crop production. Nat. Rev. Microbiol. 23(1), 9–23.

Cornelis, P., 2010. Iron uptake and metabolism in pseudomonads. Appl. Microbiol. Biotech. 86(6), 1637–1645.

de Zelicourt, A., Al-Yousif, M., Hirt, H., 2013. Rhizosphere microbes as essential partners for plant stress tolerance. Mol. Plant. 6(2), 242–245.

Donoso, E.P., Bustamante, R.O., Carú, M., Niemeyer, H.M., 2008. Water deficit as a driver of the mutualistic relationship between the fungus *Trichoderma harzianum* and two wheat genotypes. Appl. Environ. Microbiol. 74(5), 1412–1417.

Finkel, O.M., Salas-González, I., Castrillo, G., Spaepen, S., Law, T.F., Teixeira, P.J., Jones, C.D., Dangl, J.L., 2020. A single bacterial genus maintains root growth in a complex microbiome. Nat. 587(7832), 103–108.

Geng, Y., Chen, S., Lv, P., Li, Y., Li, J., Jiang, F., Wu, Z., Shen, Q., Zhou, R., 2025. Positive role of *Trichoderma harzianum* in increasing plant tolerance to abiotic stresses: a review. Antioxidants 14(7), 807.

Hantzis, L.J., Kroh, G.E., Jahn, C.E., Cantrell, M., Peers, G., Pilon, M., Ravet, K., 2018. A program for iron economy during deficiency targets specific Fe proteins. Plant Physiol. 176(1), 596–610.

Harman, G.E., Doni, F., Khadka, R.B., Uphoff, N., 2021. Endophytic strains of *Trichoderma* increase plants' photosynthetic capability. Int. J. Mol. Sci. 22(15), 8914.

Himpsl, S.D., Mobley, H.L.T., 2019. Siderophore detection using chrome azurol S and cross‐feeding assays. Met. Mol. Biol. 2021, 97–108.

Jeong, J., Guerinot, M.L., 2009. Homing in on iron homeostasis in plants. Trends Plant Sci. 14(5), 280–285.

Jones, P., Garcia, B.J., Furches, A., Tuskan, G.A., Jacobson, D., 2019. Plant host-associated mechanisms for microbial selection. Front. Plant Sci. 10, 862.

Kabir, A.H., Bennetzen, J.L., 2024. Molecular insights into the mutualism that induces iron deficiency tolerance in sorghum inoculated with *Trichoderma harzianum*. Microbiol. Res. 281, 127630.

Kiers, E.T., Rousseau, R.A., West, S.A., Denison, R.F., 2003. Host sanctions and the legume-rhizobium mutualism. Nat. 425(6953), 78–81.

Kim, D., Langmead, B., Salzberg, S.L., 2015. HISAT: a fast spliced aligner with low memory requirements. Nat. Methods 12(4), 357–360

Lamers, J., van der Meer, T., Testerink, C., 2020. How Plants Sense and Respond to Stressful Environments. Plant Physiol. 182(4), 1624–1635.

Law, C. W., Alhamdoosh, M., Su, S., Dong, X., Tian, L., Smyth, G. K., Ritchie, M.E., 2016. RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR. F1000Research, 5, ISCB Comm J-1408.

Lei, M., Wang, X., Chen, K., Wei, Q., Zhou, M., Chen, G., Su, S., Tai, Y., Zhuang, K., Li, D., Liu, M., Zhang, S., Wang, Y., 2025. Sugar transporters: mediators of carbon flow between plants and microbes. Front. Plant Sci. 16, 1536969.

Liberal, Â., Pinela, J., Vívar-Quintana, A.M., Ferreira, I.C.F.R., Barros, L., 2020.. Fighting iron-deficiency anemia: innovations in food fortificants and biofortification strategies. Foods, 9(12), 1871.

Lindström, K., Mousavi, S.A., 2020. Effectiveness of nitrogen fixation in rhizobia. Microbial. Biotech. 13(5), 1314–1335.

Liu, Y., Xiong, Z., Wu, W., Ling, H. Q., Kong, D., 2023. Iron in the Symbiosis of Plants and Microorganisms. Plants 12(10), 1958.

Livak, K.J., Schmittgen, T. D., 2001. CA: Analysis of relative gene expression data using real time quantitative PCR and the 2(Delta Delta C(T)) method. Methods 25, 402–408.

López-Bucio, J., Pelagio-Flores, R., Herrera-Estrella, A., 2015. *Trichoderma* as biostimulant: Exploiting the multi-level properties of a plant beneficial fungus. Sci. Hort. 196, 109–123.

McMurdie, P.J., Holmes, S., 2013. Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PloS one 8(4), e61217.

Mendes, R., Garbeva, P., Raaijmakers, J.M., 2011. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. FEMS Microbiol. Rev. 35(4), 635–663.

Merry, R., Espina, M.J., Lorenz, A.J., Stupar, R.M., 2022. Development of a controlled-environment assay to induce iron deficiency chlorosis in soybean by adjusting calcium carbonates, pH, and nodulation. Plant Met. 18(1), 36.

Nilsson, R.H., Larsson, K.H., Taylor, A.F.S., Bengtsson-Palme, J., Jeppesen, T.S., Schigel, D., Kennedy, P., Picard, K., Glöckner, F.O., Tedersoo, L., Saar, I., Kõljalg, U., Abarenkov, K., 2019. The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. Nucleic Acids Res. 47(D1), D259–D264.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O., 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res. 41, D590–D596.

Roriz, M., Pereira, S.I.A., Castro, P.M.L., Carvalho, S.M.P., Vasconcelos, M.W., 2023. Impact of soybean-associated plant growth-promoting bacteria on plant growth modulation under alkaline soil conditions. Heliyon 9(5), e14620.

Saha, M., Sarkar, S., Sarkar, B., Sharma, B. K., Bhattacharjee, S., Tribedi, P., 2016. Microbial siderophores and their potential applications: A review. Environ. Sci. Poll. Res. 23(5), 3984–3999.

Sasse, J., Martinoia, E., Northen, T., 2018. Feed your friends: Do plant exudates shape the root microbiome? Trends Plant Sci. 23(1), 25–41.

Schmidt, J., Dotson, B.R., Schmiderer, L., van Tour, A., Kumar, B., Marttila, S., Fredlund, K.M., Widell, S., Rasmusson, A.G., 2020. Substrate and plant genotype strongly influence the growth and gene expression response to Trichoderma afroharzianum T22 in sugar beet. Plants 9(8), 1005.

Shahid, I., Han, J., Hardie, D., Baig, D.N., Malik, K.A., Borchers, C.H., Mehnaz, S., 2021. Profiling of antimicrobial metabolites of plant growth promoting Pseudomonas spp. isolated from different plant hosts. 3 Biotech 11(2), 48.

Sood, M., Kapoor, D., Kumar, V., Sheteiwy, M. S., Ramakrishnan, M., Landi, M., Araniti, F., Sharma, A., 2020. *Trichoderma*: The "secrets" of a multitalented biocontrol agent. Plants 9(6), 762.

Staiger D., 2002. Chemical strategies for iron acquisition in plants. Angewandte Chemie (International ed. in English), 41(13), 2259–2264.

Thapa, A., Hasan, M.R., Kabir, A.H., 2025. *Trichoderma afroharzianum* T22 induces rhizobia and flavonoid-driven symbiosis to promote tolerance to alkaline stress in garden pea. Plant Cell Environ. https://doi.org/10.1111/pce.15581

Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T., Singh, B.K., 2020. Plant–microbiome interactions: From community assembly to plant health. Nat. Rev. Microbiol. 18(11), 607–621.

Tyśkiewicz, R., Nowak, A., Ozimek, E., Jaroszuk-Ściseł, J., 2022. *Trichoderma*: The current Status of its application in agriculture for the biocontrol of fungal phytopathogens and stimulation of plant growth. Int. J. Mol. Sci. 23(4), 2329.

Yadav, V., Wang, Z., Wei, C., Amo, A., Ahmed, B., Yang, X., Zhang, X., 2020. Phenylpropanoid pathway engineering: an emerging approach towards plant defense. Pathogens 9(4), 312.

Yang, T., Liu, R., Luo, Y., Hu, S., Wang, D., Wang, C., Pandey, M. K., Ge, S., Xu, Q., Li, N., Li, G., Huang, Y., Saxena, R. K., Ji, Y., Li, M., Yan, X., He, Y., Liu, Y., Wang, X., Xiang, C., … Zong, X., 2022. Improved pea reference genome and pan-genome highlight genomic features and evolutionary characteristics. Nat. Gen. 54(10), 1553–1563.

Zhang, J., Liu, Y. X., Zhang, N., Hu, B., Jin, T., Xu, H., Qin, Y., Yan, P., Zhang, X., Guo, X., Bai, Y., 2019. *NRT1.1B* is associated with root microbiota composition and nitrogen use in rice. Nat. Biotech. 37(6), 676–684.

Fig. 1. Effect of *Trichoderma afroharzianum* T22 under Fe-sufficient and Fe-deficient conditions on (A) Plant height, (B) plant biomass (fresh weight), (C) SPAD chlorophyll index, (D) maximum quantum efficiency of PSII (Fv/Fm), (E) Fe concentration in roots, (F) Fe concentration in leaves, (G) N concentration in roots, and (H) N concentration in leaves without (–T22) or with (+T22) inoculation. Boxplots show median (line), interquartile range (box), and range (whiskers); diamonds denote the mean. Pea plants were cultivated for 5 weeks; n = 3 independent biological replicates per treatment. Statistical differences between –T22 and +T22 within each Fe regime were assessed by two-tailed Student's *t*-test ($p < 0.05$ indicated by "*"; ns, not significant).

.

Fig. 2. Effect of *Trichoderma afroharzianum* T22 inoculation under Fe-sufficient and Fe-deficient conditions on (A) Relative abundance of T22, (B) number of nodules per plant, and (C) rhizosphere siderophore production (%) with (+T22) or without (–T22) inoculation in pea plants. Data represent mean ± SD from three independent biological replicates. Statistical differences between –T22 and +T22 within each genotype and Fe treatment were determined using a two-tailed Student's *t*-test ($p < 0.05$, *$p < 0.01$; ns, not significant).

Fig. 3. Transcriptomic changes induced by *Trichoderma afroharzianum* T22 inoculation under Fe-sufficient and Fe-deficient conditions. (A, B) Numbers of differentially expressed genes (DEGs) in roots of Fe-sufficient (A) and Fe-deficient (B) pea plants inoculated with T22 (+T22) compared to uninoculated controls (–T22). Genes with |log□ fold change| ≥ 1 and adjusted $p < 0.05$ were considered significantly upregulated (blue) or downregulated (orange). (C) Heatmap of selected DEGs associated with nutrient transport, stress response, and secondary metabolism in Fe-sufficient and Fe-deficient conditions. Color scale represents log□ fold change values. Data were obtained from three biological replicates per treatment.

Fig. 4. Gene Ontology (GO) enrichment analysis of differentially expressed genes in roots of plants inoculated with *Trichoderma afroharzianum* T22 (+T22) compared to uninoculated controls (–T22) under Fe-sufficient (A) and Fe-deficient (B) conditions in pea plants. The top enriched GO terms are shown for each condition, with fold enrichment on the x-axis. Dot size indicates the number of genes associated with each GO term, and dot color represents –log□□(FDR) values. GO terms are categorized into biological processes, molecular functions, and cellular components. Data were generated from three independent biological replicates per treatment.

Fig. 5. Impact of *Trichoderma afroharzianum* T22 inoculation on bacterial (16S) community diversity and composition in the roots of pea plants under Fe-sufficient and Fe-deficient conditions. (A, D) Alpha diversity measures (Observed ASVs and Simpson index) of root-associated bacterial communities under Fe-sufficient (A) and Fe-deficient (D) conditions for plants with (+T22) or without (–T22) inoculation. *P*-values were determined using Kruskal–Wallis tests (a non-parametric rank-based test). (B, E) Relative abundance of bacterial classes under Fe-sufficient (B) and Fe-deficient (E) conditions. (C, F) Relative abundance of bacterial genera under Fe-sufficient (C) and Fe-deficient (F) conditions. Taxa showing significant differences between +T22 and –T22 within each Fe condition are marked with an asterisk (*, *p* < 0.05). Data represents the mean of three independent biological replicates per treatment.

Fig. 6. Effect of *Trichoderma afroharzianum* T22 inoculation on fungal community diversity and composition in the roots of pea plants under Fe-sufficient and Fe-deficient conditions. (A, D) Alpha diversity measures (Observed ASVs and Simpson index) of root-associated fungal communities under Fe-sufficient (A) and Fe-deficient (D) conditions for plants with (+T22) or without (–T22) inoculation. *P*-values were determined using Kruskal–Wallis tests (a non-parametric rank-based test). (B, E) Relative abundance of fungal classes under Fe-sufficient (B) and Fe-deficient (E) conditions. (C, F) Relative abundance of fungal genera under Fe-sufficient (C) and Fe-deficient (F) conditions. Taxa showing significant differences between +T22 and –T22 within each Fe condition are marked with an asterisk (*, $p < 0.05$). Data represents the mean of three independent biological replicates per treatment.

Fig. 7. Model of *Trichoderma harzianum* T22–mediated effects on pea roots under Fe-deficient (Fe–) and Fe-sufficient (Fe+) conditions. (A) Under Fe deficiency, T22 inoculation leads to microbial enrichment of *Pseudomonas, Mitsuaria, Variovorax,* and *Acidovorax* along with *Rhizobium* induction. These changes promote siderophore release and nodulation, resulting in increased Fe and N availability. Transcriptomic reprogramming further supports these processes, including the induction of *NRAMP* and heavy metal–associated (HMA) transporters for Fe uptake, nodule-specific GRP-repeat genes for N metabolism, phenylalanine ammonia-lyase for defense and metabolic adjustments, and sugar transporter-like genes for nutrient allocation. Together, these microbial and molecular responses restore nutrient balance under Fe limitation. (B) Under Fe sufficiency, T22 inoculation results in minimal microbiome restructuring, characterized by enrichment of *Pararhizobium*. Siderophore activity is repressed, transcriptional changes are weak, and overall contributions to Fe and N balance are limited, leading to minimal physiological impact.