Maternal Influenza A Virus Infection Induces Antiviral and Immune Dysregulation in the Placenta and Fetus Without Vertical Transmission

Orlando Cervantes[1,2], Hazel Huang[2], Sidney Sun[2], Briana Del Rosario[2], Hong Zhao[2], Amanda Li[2,3], Andrew Vo[2,5], Gygeria Manuel[2], Roslyn Van Abel[2], Jeff Munson[4], John Cornelius[2], Raj P. Kapur[6,7], Miranda Li[2,5], Edmunda Li[2], Hanning Li[2], Alexandra Christodoulou[8], Melissa Berg[8], Britni Curtis[8], Elizabeth Miller[8], Shannon Ruff[8], Brenna Menz[8], Solomon Wangari[8], Audrey Germond[8], Audrey Baldessari[8], Chris English[8], W. McIntyre Durning[8], Thomas B. Lewis[9], Megan N. Fredericks[9], Mara Maughan[9], Austyn Orvis[10], Michelle Coleman[10], Lakshmi Rajagopal[1,10,11], Deborah H. Fuller[8,9], Kristina M. Adams Waldorf[1,2,8,*]

[1] Department of Global Health, University of Washington, Seattle, WA, 98195

[2] Department of Obstetrics and Gynecology, University of Washington, Seattle, WA, 98109

[3] Case Western Reserve School of Medicine, Cincinnati, OH

[4] Department of Psychiatry, University of Washington, Seattle, WA, 98195

[5] School of Medicine, University of Washington, Seattle, WA, 98195

[6] Department of Lab Medicine and Pathology, University of Washington, Seattle, WA, 98195

[7] Department of Pathology, Seattle Children's Hospital, Seattle, WA,

[8] Washington National Primate Research Center, Seattle, WA, 98195

[9] Department of Microbiology, University of Washington, Seattle, WA 98195

[10] Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle, WA

[11] Department of Pediatrics, University of Washington, Seattle, WA 98195

*Corresponding author: Dr. Kristina Adams Waldorf, 750 Republican St., Seattle, WA 98109; adamsk@uw.edu

Running Title: Placental-Fetal Immune Response to Influenza A Virus

Keywords: influenza, H1N1, placenta, interferon, fetus, pregnancy

Abstract Word Count (150 words): 148 words

Main Text Word Count (<4,000 words): 3,478

ABSTRACT

Influenza A virus (IAV) infection during pregnancy is associated with stillbirth and preterm birth, possibly by disrupting placental and fetal immunity. To investigate this, pregnant pigtail macaques were inoculated with IAV [A/California/07/2009 (H1N1)] and examined at necropsy 5 days post-infection (N=11) versus uninfected controls (N=16). Stillbirth occurred in 18% of infected pregnancies but not in controls. While vertical transmission was not observed, low levels of viral RNA were detected in two placentas. Maternal IAV infection was associated with increased placental IL-1β and IFN-β levels and an upregulated type I interferon and integrated stress transcriptional response. Fetuses exposed to IAV had greater frequencies of innate immune cells in lymph nodes and CD4+ T cells in lungs. These results suggest that placental and fetal immune environments undergo immune activation independent of the severity of maternal lung infection. Influenza vaccination during pregnancy may protect against potentially harmful effects on fetal development.

KEYWORDS: influenza A virus, H1N1, pregnancy, fetus, placenta, nonhuman primate, innate immunity

INTRODUCTION

Influenza A virus (IAV) infections remain a global public health threat, which is especially pronounced in vulnerable populations like pregnant women.[1,2] IAV is a negative-sense, single-stranded RNA virus composed of eight segments. Pandemics occur when reassortment of the viral segments occurs to produce new combinations of its surface glycoproteins, hemagglutinin (HA), and neuraminidase (NA). Epidemiologic evidence from several IAV pandemics (e.g., 1918, 1957, 1968, 2009) indicates that IAV infections in pregnancy impart a greater risk of maternal mortality and adverse pregnancy outcomes, such as stillbirth.[1-12] There may also be a long-term risk of neuropsychiatric disease in the children born to pregnant women infected with IAV or other infectious diseases, which has been reported in multiple populations.[13-20] Vaccination with the seasonal trivalent IAV vaccine has been associated with a reduced risk of stillbirth in some[21-25], but not all, studies.[9,26] Despite the public health recommendation for pregnant women to receive the seasonal IAV vaccine[27], only half of pregnant women in the U.S. typically receive the vaccination each year.[28]

Increasing vaccine skepticism since the COVID-19 pandemic requires a better public understanding of the importance of annual seasonal influenza vaccination to reduce the adverse impacts of IAV infections during pregnancy on placental and fetal health.[29,30] Reducing maternal mortality is an important benefit of vaccination, but in the absence of a pandemic, this rare outcome may not be perceived as a risk. A spectrum of adverse pregnancy and fetal outcomes has been reported in pregnant mice and rats[31-38], but these animal models are dissimilar to human pregnancy in many respects including a short gestation (3-5 weeks). Humans have a gestational period that is 8–10 times longer than these species, allowing for extended fetal brain development *in utero.* However, this long gestational period also increases the placenta and fetus' exposure to seasonal waves of IAV infection. Nonhuman primates (NHPs) share many similarities with human pregnancy, including a longer gestation, placental structure, and physiology of labor.[39,40] NHPs

are also frequently used in IAV disease modeling due to their similarities in human clinical and

pathological features.[41,42] A few studies using pregnant NHPs have modeled IAV infection but

were performed with the intent to understand complement activation and infant brain development

in African green monkeys and rhesus macaques, respectively.[43,44] The broader impact of IAV

infection during pregnancy on placental and fetal immunity and health was not characterized.

Our objective was to determine the impact of a maternal IAV infection with a 2009 pandemic

strain, on innate and adaptive immune responses in the placenta and fetus of the pigtail macaque

(*Macaca nemestrina*) model. There are comparatively few studies linking immune responses in

the placenta with those in the fetus.[32,36,45,46] A secondary objective was to account for how

differences in the severity of the maternal IAV infection might affect the placental immune

response, as part of the larger nexus of crosstalk between the mother, placenta, and fetus during

acute IAV infection.

RESULTS

Impact of Influenza A Viral Disease on Pregnancy Outcome and Lung and Placental

Histopathology

Eleven pregnant pigtail macaques were inoculated with an IAV H1N1 2009 pandemic strain

(A/CA/07/2009) in the third trimester (FLU cohort) using a combination of intranasal, intratracheal,

intraocular, and intraoral routes with 7.4×10^6 plaque-forming units (PFU). Results were compared

to 16 historical control uninfected pregnant pigtail macaques (CTRL cohort, Table S1). Five days

after inoculation, Cesarean section and necropsies of the dam and fetus were performed. Stillbirth

occurred in two of the eleven FLU pregnancies (18%, FLU7 and FLU9), but in no CTRL animals.

Clinical signs of maternal illness were documented using daily clinical scores and vital signs taken

at the time of sedations: before inoculation, 1- and 3-days post-inoculation (dpi), and on 5 dpi, the

experimental endpoint (Table 1). Coughing, wheezing, increased respiratory effort, nasal

discharge, and decreased appetite were common symptoms in 7 of 11 FLU animals (64%), while

the remaining four did not show clinical symptoms. Animals that exhibited the most severe

disease based on clinical scores were FLU8 and FLU9; of these animals, FLU9 experienced a

stillbirth. Clinical scores were highest between day 3 and day 5 post-inoculation. Uninfected CTRL

animals did not show signs of respiratory disease.

Histopathology of the maternal lungs and placentas was evaluated to determine if IAV inoculation

induced influenza-associated pathology in either organ (Fig. 1, Table S2). Lung histopathology

revealed pneumonia in all cases, including the two cases with stillbirth; one dam experiencing a

stillbirth had mild, multifocal pneumonia (FLU7), and the other had severe pneumonia with pleural

effusion and hilar lymphadenopathy (FLU9). Of the CTRL animals with histopathology data

available, none had pulmonary inflammation. Most placentas from FLU animals had evidence of

either placental infarction (coagulative villous necrosis), neutrophilic (necrotizing) villitis, or both

(8/11, 73%; Fig. 1B-C, Table S2). Infarction or active inflammation was observed less frequently

in the CTRL placentas (5/16, 31%). None of the placentas demonstrated active inflammation of

the chorioamniotic membranes. Placental histopathology in the two stillbirth cases (FLU7, FLU9)

overlapped with that observed in other IAV-infected animals, and some CTRL animals.

 Figure 1. Placental histopathology

Each pair of images shows a low-magnification image of the placental disc (fetal surface toward the top) on the left and a higher-magnification image of villous tissue on the right. (A, A') Normal placental anatomy from a control animal showing no inflammation or necrosis. (B, B') A focus of subacute infarction with coagulative necrosis and no active inflammation is shown from a representative animal in the FLU cohort. (C, C') A focus of necrotizing villitis (highlighted by arrows) with neutrophilic inflammation and villous necrosis is shown from a representative animal in the FLU cohort.

IAV H1N1 vRNA in Lung, Placenta, and Fetus

RT-qPCR was performed to measure IAV viral RNA (vRNA) in the maternal lungs, placenta, and

fetal organs (Table 1). At 5 dpi, all FLU animals had high levels of IAV vRNA in their lungs ranging

from 2,220 copies per milligram to more than 6 million copies per milligram. One exception was

FLU10, which had an almost undetectable lung viral load (3 copies/mg). IAV vRNA was detected

in the placental chorionic villous tissue of only two cases (2/11; 18%), both with a low copy number

in the placenta (FLU1 and FLU9; Table 1). Of these, FLU9 additionally experienced stillbirth. Fetal

tissue samples from the brain, brainstem, spinal cord, meninges, heart, lungs, liver, spleen, and

thymus were negative for IAV H1N1 vRNA across the entire FLU cohort (Table S3). Overall, the

maternal lung was characterized by a wide distribution of IAV viral load, infrequent detection of

IAV vRNA in placental tissues, and no evidence for congenital infection.

Cytokine Concentrations in the Placental Disc and Amniotic Fluid

Cytokine responses in the placental disc and amniotic fluid were evaluated to determine if there

was a correlation between maternal IAV infection status, viral load, or stillbirth. Chorionic villous

tissue from the placenta disc of FLU animals had significantly greater concentrations of some

inflammatory and antiviral cytokines than CTRL placentas, including IL-1β, IL-18, IFN-β, and GM-

CSF (all p≤0.03, Fig. 2). The highest concentrations of IFN-α, IFN-β, IL-6, TNF, and GM-CSF

within the FLU cohort were observed in both placentas with low IAV viral load, of which one was

a stillbirth case; however, the second stillbirth case had low cytokine concentrations. Several

cytokines with low expression (IL-2, IL-10, IL-12p70, IL-13, IL-17A, and IL-23) were significantly

higher in the CTRL versus FLU cohort (Fig. S1). However, in many cases, these cytokine values

fell below the working range of the Luminex assay.

Figure 2. Cytokine concentrations in the placental disc.

Each panel shows the concentration of a cytokine quantified by multiplex Luminex assay in chorionic villous tissue from the FLU (blue) and CTRL (black) cohorts. Values are either measured (dots) or estimated (x's) by extrapolation of the standard curve. The thick line indicates the median, and the bounding box indicates the 1st and 3rd quartiles. Red triangles marks samples from animals that experienced a stillbirth. A red circle marks samples from animals that had a low placental viral load. A red square marks samples from animals that experienced a stillbirth and had low placental viral load. The p-value for CTRL versus FLU cohorts is indicated at the top of each panel if less than 0.05.

In contrast to chorionic villous tissue, several cytokines in the amniotic fluid were significantly

higher in the CTRL versus FLU cohort (GM-CSF, IFN-β, IFN-γ, IL-1β, IL-4, IL-5, IL-13, IL-17A,

and IL-23; all $p<0.05$). In most cases, the concentration of the cytokines in both cohorts was very

low (Fig. S2). These results indicate that perturbations in cytokines associated with maternal IAV

infection were compartment-specific, and elevated cytokines in the chorionic villous tissue were

not transported into the amniotic fluid compartment.

Immune Cell Frequencies in the Placenta, Fetal Blood and Organs

To determine if a maternal IAV H1N1 infection was associated with changes in the innate and

adaptive immune cell composition of the placental disc, we performed an immunophenotyping

assay using flow cytometry. The frequency of classical monocytes among live leukocytes was

significantly decreased in the placental disc of the FLU versus CTRL cohort ($p=0.004$, Fig. 3).

Immune cell frequencies between the FLU and CTRL cohorts were variable in the chorionic villous

tissue, making it challenging to ascertain significant differences between the two groups with

modest group sizes. There was no relationship between stillbirth and the frequency of immune

cell populations in the placenta.

Figure 3. Immunophenotyping of the placental disc by flow cytometry

Each panel shows the frequency of the defined cell population as a percentage of live cells, in

chorionic villous tissue of the FLU (blue) and CTRL (black) cohorts. The thick line indicates the

median, and the bounding box indicates the 1st and 3rd quartiles. A red triangle marks samples

from animals that experienced stillbirth. A red circle marks samples from animals that had a low

placental viral load. A red square marks samples from animals that experienced a stillbirth and

had low placental viral load. The p-value for CTRL versus FLU cohorts is indicated at the top of

each panel if less than 0.05.

In fetal organs, there was a coordinated perturbation of innate immune cells and T cells in the

FLU versus CTRL cohorts. In the fetal lung, the frequencies of intermediate monocytes, non-

classical monocytes, plasmacytoid dendritic cells, and CD4+ T cells were significantly higher in

the FLU cohort compared to the CTRL cohort (p≤0.04, Fig. S3). A similar pattern was evident in

the fetal mesenteric lymph nodes, with a significantly higher frequency of intermediate monocytes

(p=0.02), and observably higher frequencies of NKT cells, non-classical monocytes, and

plasmacytoid dendritic cells in the FLU versus CTRL cohort (p=0.06, Fig. S4). In fetal whole blood,

there was a significantly higher frequency of NKT cells in the FLU cohort (p=0.03, Fig. S5). These

differences suggest that maternal IAV H1N1 infection perturbs populations of fetal innate and

adaptive immune cells in organs, lymphoid compartments, and blood even without evidence of

vertical transmission.

Placental Transcriptional Profile

To determine if a maternal IAV infection altered the transcriptional signature in the placental disc,

we performed total RNA-Seq on chorionic villous tissues and validated our results by direct gene

counting using a Nanostring nCounter assay. A multidimensional scaling plot of the normalized

RNA-Seq gene counts demonstrated a clear division between the two cohorts (Fig. 4A; CTRL,

blue dots) and FLU (red symbols). In the placental chorionic villous tissue, 185 genes were

upregulated, and 178 genes were downregulated relative to the gene expression of the CTRL

group (≥ or ≤ 1.5-fold, p<0.05; Fig. 4B). Inspection of the upregulated genes revealed a striking

upregulation of interferon (IFN)-stimulated genes (ISGs) including *IFIT3, MX1*, and *OAS2* in

placentas from FLU animals relative to CTRL placentas. A Gene Set Enrichment Analysis (GSEA)

revealed multiple significant, positively enriched pathways related to the type I IFN response and

the negative regulation of viral processes such as genome replication and viral entry in FLU

versus CTRL animals (Fig. 4C). Digital gene counting using a 770-plex Nanostring nCounter

panel supported these results with a similar striking upregulation of ISGs (Fig. S6), and strong

correlation with total RNA-Seq gene counts in both the FLU and CTRL groups (Fig. 4D; r=0.8,

p<2.2x10^{-16} for both).

Figure 4. Total RNA-Seq of the placental disc from FLU and CTRL animals.

(A) Multidimensional scaling (MDS) was performed on normalized log-transformed gene counts

using the Euclidean distance. Each point represents an individual sample, and the distance

between points reflects the relative similarity of their global expression profiles. Samples are

colored according to experimental group (FLU: red; CTRL: blue). Symbols denoted the following:

an open red circle for low placental IAV vRNA, an open red square for a stillbirth with undetectable

vRNA, and an open red triangle for a stillbirth with low vRNA. (B) A volcano scatter plot illustrates

the relationship between the log2 fold change of gene expression (x-axis) and statistical

significance [-log10 (p-value), y-axis] of differentially expressed genes between the FLU and

CTRL cohorts. Dot colors indicated significance and magnitude: red for genes with significant log2

fold change ≥1 or ≤–1; blue for genes with |log2 fold change| <1 but p<0.05; gold for non-

significant genes with |log2 fold change| ≥1; and black for non-significant genes with |log2 fold

change| <1. (C) A Gene Set Enrichment Analysis of significant Gene Ontology (GO) terms in FLU

versus CTRL placentas is shown (p<0.05). Colored dots indicate the Normalized Enrichment

Score (NES), which evaluates the degree to which a set of genes in a GO category is over-

represented in a ranked gene list. The FDR, or false discovery rate, is a statistical method used

to correct for multiple testing when analyzing many GO terms simultaneously. (D) This dot plot

illustrates the correlation between the placental log2-transformed gene counts per million

obtained from total RNA-Seq and the log10-transformed gene counts from the Nanostring

nCounter assay. Each dot represents a mean value of a gene count from either the FLU (red) or

CTRL (black) group, which has both nCounter and RNA-Seq data. A Pearson correlation test was

performed, and the r coefficient and p-value for each group are noted in the upper left-hand corner

of the plot. The line of best fit with standard error is shown.

Next, we evaluated whether the viral infection in the mother's lungs or cytokines produced in the

lung or placenta might be driving these findings. We performed unsupervised hierarchical

clustering of the placenta RNA-Seq data and initially ordered the heatmap according to the

maternal lung viral load or Type I IFN cytokine concentrations in the lung and placenta (Fig. 5A).

No clear pattern in the placental transcriptional response emerged regardless of how the heatmap

was ordered, whether it be by lung viral load, IFN-α (lung, placental disc) or IFN-β (placental disc)

concentrations. No correlation was observed between placental RNA-Seq gene counts and

placental viral load, which was undetectable in all but two cases.

Hierarchical clustering yielded two distinct clusters of differentially expressed genes: cluster 1

(downregulated genes, FLU vs. CTRL; Fig. 5B) and cluster 2 (upregulated genes, FLU vs. CTRL;

Fig. 5C). Overrepresentation Analysis (ORA) was performed to identify Gene Ontology (GO)

terms enriched in both clusters. No significant GO terms were identified in cluster 1. However,

inspection of genes in cluster 1 revealed downregulation in the FLU group of multiple genes

involved in ubiquitination, proteasome function, unfolded protein response, metabolism, and cell

cycle/DNA repair (Fig. 5B). To determine if this profile might reflect induction of a viral-associated

integrated stress response (ISR), we analyzed expression of two key genes in the pathway:

EIF2AK2 (ISR activator) and *ATF4* (ISR effector). Of these genes, *EIF2AK2* expression was

significantly upregulated in the placenta [log2 fold change=1.24, p<0.05; Fig. 5B]. In cluster 2,

significantly enriched GO terms in the FLU cohort were related to type I and II interferon signaling,

viral genome regulation, and antiviral defense processes that featured ISG genes (Fig. 5C; Fig.

S7). A notable outlier was FLU11, which had a transcriptional profile more similar to CTRL

animals. Placental transcriptomics of FLU3 and FLU4 were intermediate between the FLU and

CTRL cohorts. Collectively, transcriptional analysis of the placental chorionic villous tissue from

FLU animals revealed a striking upregulation of antiviral innate immune genes and

downregulation of critical cellular functions (cell cycle, metabolism, ubiquitination) consistent with

an antiviral and ISR response, regardless of the severity of disease in the maternal lung.

Figure 5. RNA-Seq analysis of placental chorionic villous tissue in FLU and CTRL groups

Heatmaps are shown of normalized counts from the placental total RNA-Seq data from the FLU

and CTRL cohorts. Each column represents an individual placental sample, and each row

represents an individual gene. The yellow-blue color intensity indicates the gene z-score which

was calculated based on the mean expression level of all animals for each gene. (A) An

unsupervised hierarchical clustering analysis was performed on the top 500 differentially

expressed genes in the placentas of the FLU cohort using the Euclidean distance metric and

Ward's method. The relative quantity of maternal lung viral load in the IAV-infected group is

indicated by the green color gradient. Lung IFN-α and IFN-β, and placenta IFN-β concentrations

are indicated by red, orange, and gray color gradients, respectively. (B) Cluster 1 genes from

panel A are shown, which have functions impacted by the viral-associated ISR, such as cell

cycle/DNA repair, metabolism, ubiquitination, unfolded protein response (UPR), and proteasome

activity. The viral-associated ISR activator (*EIF2AK2*) and mediator (*ATF4*) are shown at the top.

(C) An over-representation analysis (ORA) was performed on cluster 2 genes from panel A. Rows

are genes in the Molecular Signatures Database Gene Set: Hallmark Interferon Alpha Response.

DISCUSSION

Our results indicate that IAV infection in pregnant NHPs induces a placental stress response and alters the immune environment in both the placenta and fetus, even in the absence of detectable viral infection in these compartments. Interestingly, placental injury and abnormal gene expression was observed among all IAV-infected animals despite significant variability in viral load and type I interferon concentrations in the maternal lungs. To our knowledge, this is the first demonstration that maternal IAV lung infection, independent of disease severity, drives significant alterations in placental gene expression, emphasizing the placenta's function as an immunologically responsive organ to maternal systemic inflammation. This key finding suggests that pregnant women with even a mild course of IAV disease at the time of birth are as vulnerable to placental and fetal innate immune dysregulation as those with severe infection.

While direct placental infection and vertical transmission of IAV is rare, our data show that maternal infection can disrupt the fetal innate and adaptive immune environment. This cascading series of linked biological events, beginning in the mother's lungs and moving to the placenta, demonstrates how a pulmonary viral infection can influence fetal development and potentially prime the fetus for long-term health consequences. Dysregulation of the immune environment in the fetus may be subtle but could underlie the increased lifetime risk of neuropsychiatric conditions observed in children exposed to influenza and other infections *in utero*.[13,14,19,20,47,48] We recently reported granular cytoplasmic inclusions in astrocytes within the deep white matter of fetal pigtail macaques, a pathological finding that was more frequent in fetuses exposed to a maternal IAV infection.[49] The impact of perturbations in fetal immunity and subtle changes in the fetal brain associated with congenital infections are unknown, but it is a concerning potential risk factor for abnormal neurodevelopment and a spectrum of neuropsychiatric diseases.

A defining feature of our study was the identification of a robust type I interferon (IFN-β) response and inflammasome activation (IL-1β, IL-18) in the placenta. IAV is known to upregulate type I IFNs

in the lungs, but in the absence of placental infection, it has not been clear the degree to which

the virus in the lungs might induce an antiviral response in the placenta. Many of the upregulated

genes we observed are known to restrict IAV infection by sensing viral RNA (*DDX58*), restricting

viral entry (*MX1*) or replication (e.g., *IFIT2*, *EIF2AK2*), and destroying viral RNA (e.g., *OAS2,*

OAS3). Limited evidence suggests that IAV can replicate in the placenta and transfer to the fetus,

and documentation of this phenomenon or transplacental passage has been rare.[50,51] The

placenta's robust antiviral innate immune defense likely comes at a cost: type I IFNs and

inflammasome activation can lead to tissue injury, a known risk factor for stillbirth and preterm

birth.[52,53] The degree to which the placenta can balance the competing needs of viral clearance

without inciting tissue injury likely holds the key to preventing stillbirth.

NHPs in this study were not previously vaccinated against IAV and had no prior IAV infection. In

an IAV-naïve host, maternal CD8+ T cell activation is expected to begin within a few days, peak

in the draining lymph nodes between days 3-5, and become detectable by days 5-7, where they

aid in viral clearance.[54] Viral control in this early stage was likely exclusively a product of the

antiviral innate immune response in the maternal lung, which may have indirectly activated similar

responses in peripheral organs, including the placenta. In parallel, the viral-associated ISR may

equally impair critical cellular function in the placenta like DNA replication, enzymatic reactions,

and management of accumulating unfolded and misfolded proteins. An ISR is triggered by the

cell in response to a variety of stressors, including viral infections, and has significant interaction

with innate immune signaling pathways.[55,56] A critical requirement to activate the ISR, regardless

of the type of stressor, is the phosphorylation of eukaryotic translation initiation factor-2α

(EIF2α) by a variety of kinases, one of which is EIF2αK2.[57] Upregulation of *EIF2AK2*, and

downregulation of critical cell functions impacted by ISR (ubiquitination, metabolism, cell cycle,

DNA repair) provides strong evidence that the ISR pathway is induced in the placenta during a

maternal IAV infection. The virus exploits ISR to prevent the host cell from expending energy on

anything other than viral replication. If these cellular functions cannot recover, apoptosis can be

initiated by *ATF4* and may disrupt placental functioning; increased expression of ATF4 was seen

in some, but not all FLU cases.[56] The combination of a Type I IFN response and the viral-

associated ISR may underlie stillbirth risk in pregnant women infected with IAV.

The primary strength of our study lies in the translational nature of the NHP model, which closely

recapitulates human pregnancy, allowing one to model an IAV infection in the context of

pregnancy. NHPs are highly similar to humans in their genetics, anatomy, physiology, immune

responses, and pregnancies.[39] Macaques have a long gestational period and similar placental

architecture to humans. Another strength of the pregnant NHP model of IAV infection is the ability

to study antiviral immune responses at a defined time after infection onset, which is not possible

during human pregnancy. Further, this study was deliberately conducted in the third trimester of

pregnancy, a time when pregnant women experience the greatest maternal morbidity and

mortality.[58,59] These features of the pregnant NHP model of IAV infection demonstrate the

translational relevance of this model and the marked impact of our findings.

A spectrum of pathology was observed in the FLU placentas, which was also observed in some

normal human pregnancies[47,48] However, two stillbirths occurred in our study, which were

associated with placental infarctions in one case, and neutrophilic villitis, necrosis, and fibrin

deposition in the other. In human cases of stillbirth, it is routine to extensively sample the placenta

to survey histopathology across the entire disc. Greater sampling of the placenta in our NHP

cohorts for histopathological analysis would have helped render a clearer determination of the

underlying cause of the two stillbirths. Another important limitation was that we did not study

endpoints earlier or later than five days post-infection, preventing assessment of the trajectory of

the antiviral response in the maternal lungs and placenta.

Pandemic preparedness is critical to ensure that pregnant women have access to IAV vaccines,

and for preventing stillbirth and maternal mortality. Currently, there is concern that a highly

pathogenic IAV H5N1 strain may spill over into humans and become readily transmissible. IAV

H5N1 and other highly pathogenic strains have been associated with excess mortality and

stillbirth in human cohorts.[60,61] Our results suggest that a pandemic IAV strain infecting the

maternal lungs adversely affects placental and, ultimately, fetal health. This data provides new

evidence supporting IAV vaccination in pregnancy as an actionable step to prevent not only

maternal mortality, but also viral injury to the placenta, disordered immune responses in the fetus,

and stillbirth.

ONLINE METHODS

Research Animals and Ethics Statement

All animal experiments used in this study were approved by the University of Washington

Institutional Animal Care and Use Committee (protocol# 4165-03, last approved 07/22/2025), and

in compliance with the U.S. Department of Health and Human Services Guide for the Care and

Use of Laboratory Animals and Animal Welfare Act. All animal experiments were performed at

the Washington National Primate Research Center (WaNPRC).

Study Design

Two groups of female pigtail macaques *(Macaca nemestrina)* were inoculated using a

combination of intranasal, intratracheal, intraocular, and intraoral routes with $10^{7.4}$ PFU of IAV

[A/California/07/2009 (H1N1)], reflecting a dose previously used in a study of cynomolgus

macaques (*Macaca fascicularis*).[62] Experimental groups consisted of IAV H1N1-infected pregnant

(N=11) and uninfected pregnant controls (N=18; Table S1). IAV inoculation occurred in the early

third trimester to model the time in pregnancy when IAV H1N1 infections are thought to produce

the greatest maternal lung disease and pregnancy complications.[11,12] Uninfected controls in this

study consisted of pregnant adult pigtail macaques used in prior published studies.[63,64]

Macaques inoculated with IAV were observed daily for clinical symptoms of influenza disease,

which were scored on a 3-point scale to reflect responsiveness of the animal, nasal or ocular

discharge, cough, respiratory effort, food consumption, and fecal consistency; higher scores

reflected disease, with a 0 in each category indicating a normal assessment. We chose a

predefined study endpoint of 5 days post-inoculation (dpi) to capture immune responses near the

peak of disease.[35,36] A Cesarean section was performed at the endpoint, and placentas were

extracted to sample chorionic villous tissue. The dam was euthanized using a barbiturate

overdose while under anesthesia. The fetus was euthanized using a barbiturate overdose,

followed by exsanguination. These methods are approved by the 2007 American Veterinary

Medical Association Guidelines on Euthanasia. Necropsies were performed on both the dam and

the fetus to obtain samples for transcriptomics, immune assays, and histopathology. Random

sections of the placental membrane, chorionic disc, and umbilical cord were evaluated for

pathology.

H1N1 Virus Cultivation and Quantification

We used an IAV strain [A/California/07/2009 H1N1 influenza virus (CA09)] that was obtained from

the World Reference Center for Emerging Viruses and Arboviruses at the University of Texas

Medical Branch at Galveston. Before culturing the virus, Madin-Darby canine kidney (MDCK) cells

(ATCC, Manassas, Virginia, US) were cultured in Dulbecco's modified Eagle's medium (DMEM)

supplemented with 10% fetal bovine serum (FBS), 25 mM HEPES, 100 U/mL penicillin, and 100

μg/mL streptomycin. Cells were incubated at 37°C and 5% CO_2 and confirmed to be free of

mycoplasma with the MycoAlert™ mycoplasma detection kit (Lonza, LT07-318) before use for

virus cultivation. Sub-confluent monolayers of MDCK cells in T150 flasks were washed with PBS

and infected at a multiplicity of infection (MOI) of 0.5 (5 mL serum-free MEM) at 37°C and 5%

414 CO_2. After 1 hour, the media was removed, cells were washed with 1X PBS, and the media was

415 replaced. The supernatant was harvested after 36 hours and centrifuged at room temperature to

416 remove cellular debris (1800 rpm for 5 minutes, followed by 3000 rpm for 10 minutes). Virus stock

417 was frozen at −80°C. The viral titer of each stock was quantified using a TCID50 assay with

418 hemagglutination readout and converted to PFU/mL.[65] To transform TCID50/mL into PFU/ml,

419 TCID50/mL was multiplied by 0.7. Reagents were obtained from Thermo Fisher Scientific unless

420 otherwise stated (Waltham, MA).

421 *RNA Isolation and Reverse Transcription Quantitative Polymerase Chain Reaction (RT-qPCR)*

422 Tissue samples were kept in RNAlater™ Stabilization Solution (#15596018, Thermo Fisher

423 Scientific) prior to homogenization in Precellys homogenizing tubes (#1011152, Cayman

424 Chemical, Ann Arbor, MI, USA) with the Precellys Evolution machine (Bertin Technologies,

425 Montigny-le-Bretonneux, France). RNA was extracted from homogenates using the RNeasy Mini

426 Kit (Qiagen, #74106). RNA was quantified using the NanoDrop™ 2000 spectrophotometer

427 (Thermo Scientific, #ND-2000) and the Qubit 4 fluorometer (Invitrogen, #Q33238). RNA quality

428 was evaluated by measuring the RNA integrity number (RIN) with the 4200 Tapestation System

429 (Agilent, #G2991BA).

430 To quantify IAV vRNA levels from different sample origins, we used reverse transcription

431 quantitative polymerase chain reaction (RT-qPCR) and a WHO-approved primer/probe set.[66]

432 cDNA was generated from RNA isolates using an iScript™ Select cDNA synthesis kit (Bio-Rad,

433 #1708897) using H1N1 HA-specific forward primer following manufacturer's instructions. qPCR

434 was done with H1N1 HA-specific primers and probes (Forward: GGGTAGCCCCATTGCAT;

435 Reverse: AGAGTGATTCACACTCTGGATTTC; Probe: 5`-[FAM]

436 TGGGTAAATGTAACATTGCTGGCTGG [TAMRA]-3`) using the Taqman Fast Advanced Master

437 Mix (#444964, Thermo Fisher Scientific) and a QuantStudio 3 machine (#A28132, Thermo Fisher

Scientific) following manufacturer instructions. Samples were run in triplicate. Positive results

were defined as any sample with an average Ct value less than 38 cycles.

Cytokine Quantification

ProcartaPlex™ NHP multiplex kits were used to quantify cytokines from lysates or fluids (Thermo

Fisher Scientific). Tissue lysis and homogenization were performed using an Invitrogen™

ProcartaPlex™ Cell Lysis Buffer (#EPX-99999-000, Thermo Fisher Scientific) with the addition of

a protease inhibitor cocktail (Fisher Scientific, #PI78430) using the Precellys Evolution system,

as previously described. Type I Interferons [IFN; IFN-alpha (IFNα) and IFN-beta (IFNβ)] were

quantified using the Cynomolgus IFN-beta and Cynomolgus/Rhesus IFN-alpha ELISA kits

(#46415-1 and #46100-1, respectively; PBL Biosciences, Menlo Park, CA, USA). All protocols

followed the manufacturer's guidelines.

Immunophenotyping (IMP)

Immunophenotyping by flow cytometry was performed to determine the frequencies of immune

cells in the placenta and fetal lung, mesenteric fetal lymph node, and fetal blood. Tissues from

the placental disc were dissociated into single cell suspensions and assessed for viability using

Acridine Orange/Propidium Iodide (Nexcelom Bioscience LLC). Then, samples were stained with

a panel of antibodies in Brilliant Stain Buffer (BD Biosciences) for 20 minutes to identify immune

cells as previously described.[67] Antibodies used included BD Biosciences: CD3 (SP34-2)-BV650,

CD8 (RPA-T8)-BUV395, CD11b (ICRF44)-APC-Cy7, CD11c (SHCL-3)-PE, CD45 (D058-1283)-

PE-CF594, CD86 (IT2.2)-PE-Cy5, TCRγδ (B1)-FITC; BioLegend: CD4 (OKT4)-BV605, CD14

(M5E2)-BV785, CD16 (3G8)-Alexa Flour 700, CD20 (2H7)-BV570, HLA-DR (L243)-BV711;

Invitrogen: CD123 (6H6)-eFlour 450, LIVE/DEAD Fixable Aqua for 405nm excitation; Beckman

Coulter: CD159a (Z199)-PC7; and Miltenyi: CD66 (TET2)-APC. The staining protocol proceeded

in three steps: LIVE/DEAD Fixable Aqua, then all antibodies other than CD11c and CD123, and

finally CD11c and CD123. After staining, samples were fixed with 1% paraformaldehyde and

analyzed on a Symphony A3 flow cytometer (BD Biosciences) using FACS Diva 8 software (BD

Biosciences). Samples were analyzed using FlowJo 10.9.0 (FlowJo, LLC). All events were initially

gated on Forward Scatter (FSC) singlets, CD45+ leukocytes, live cells, and then on FSC-A and

Side Scatter (SSC)-A profiles. Populations were defined using canonical markers as described in

Table S4. Representative gating strategies are shown for granulocytes (Fig. S8), lymphoid cells

(Fig. S9), and monocytes (Fig. S10). Populations were excluded from analysis if they did not meet

the minimum threshold of >100 cells in their parent gate.

Total RNA-Seq Library Preparation and Analysis

Total RNA sequencing was performed on chorionic villous tissue from the placenta. We used

100ng of RNA with RIN>3 as input for RNA library prep using the KAPA RNA HyperPrep kit with

RiboErase HMR (#08098131702, Roche Diagnostics Corporation, Indianapolis, IN, USA). The kit

was designed for human, mouse, and rat samples, but it also effectively reduced ribosomal RNA

in our NHP RNA isolates. Libraries were prepared following the manufacturer's standard protocol.

Library quantity and quality were evaluated using Qubit Fluorometer (Thermo Fisher Scientific)

and TapeStation (Agilent Technologies, Santa Clara, CA, USA). Constructed libraries were

sequenced on a NovaSeq 6000 Sequencing System (#20012850, Illumina, San Diego, CA, USA),

which produced 2x100 nucleotide stranded paired-end reads.

Raw sequences were aligned to the *Macaca nemestrina* reference genome from Ensembl

(Mnem_1.0, INSDC Assembly GCA_000956065.1) utilizing STAR in Partek Flow, a bioinformatics

software suite (Partek, Chesterfield, MO, USA). Ensembl gene symbol conversions were

employed for consistency across all gene IDs. As the quality score was greater than 35 across all

samples, no bases were trimmed, and 100% of sequencing bases overlapped with read lengths

of the reference genome. The raw count matrix was downloaded from Partek. Subsequent

statistical processing and analysis of count data were performed within the R statistical computing

environment (R version 4.4.0). Gene counts were filtered by a row mean of 10 or greater, then

normalized using edgeR to implement TMM normalization. Counts were transformed into log2

counts using voom, a function in the limma package that modifies RNA-Seq data to create a

normalized count matrix.[39,40] Differential Expression (DE) analysis compared placental samples

from IAV-infected animals against controls from uninfected animals. Comparisons were based on

a linear model fit for each gene using limma. The criteria for all DE analyses included an absolute

fold change of 2 and a p-value of less than 0.05. Gene Set Enrichment Analysis was conducted

across the comparison group using fgsea with the Broad Institute Molecular Signatures

database.[41]

Hierarchical clustering analysis, an unsupervised machine learning approach, was performed to

group total RNA-Seq profiles based on their similarity in Euclidean distance using the hclust

function in R. Ward's linkage was used to minimize variance within each cluster. The optimal

number of clusters was determined using the elbow method. The resulting hierarchy was

visualized using a dendrogram. Finally, an overrepresentation analysis (ORA) was performed to

determine if predefined sets of genes in Gene Ontology (GO) pathways were enriched in the

cluster of differentially expressed genes (DEGs) obtained from the hierarchical clustering analysis

using the clusterProfiler package in R.

Nanostring nCounter®

To validate our results from bulk RNA sequencing, we chose a quantitative multiplex method of

direct digital counting of gene transcripts using the Nanostring nCounter® Analysis Systems

(Nanostring Technologies Inc., Seattle, WA, USA). RNA extracted from the chorionic villous

tissues of the placenta was analyzed using the Nanostring nCounter® NHP Immunology V2 gene

expression codeset (#11500276, Nanostring Technologies, Inc). RNA hybridization reactions

were performed according to the manufacturer's protocol. Probe-target RNA complexes from all

reactions were processed and immobilized on nCounter® cartridges using the nCounter® Prep

Station. Gene transcripts were quantified with the nCounter® Digital Analyzer (#MAN-C0035-07,

Nanostring Technologies, Inc).

Statistical Analyses

All statistical analyses were performed in RStudio (using R version 4.4.0) or nSolver (nCounter

data only), both of which use the R programming language. A p-value of less than 0.05 was

considered significant, and 0.5-0.1 was reported considering the modest sample size. Groups

were compared using non-parametric tests. Spearman's rank correlation coefficient was used to

calculate correlations between two variables. Data visualizations were done using ggplot2,

ggpmisc, and ComplexHeatmaps packages and their derivatives.[42,43]

DATA AVAILABILITY STATEMENT

The data are openly available on several platforms. The metadata, cytokine, RT-qPCR, and flow cytometry are available in the Dryad data repository at https://doi.org/10.5061/dryad.c866t1gjf. The placenta total RNA-Seq data is available through the NIH Gene Expression Omnibus (GEO) via the GSE289117 accession number. Finally, the code used for the computational analysis and to generate the plots is available in GitHub at https://github.com/kadamswaldorflab.

ACKNOWLEDGMENTS

This work was supported by funding from the National Institutes of Health grants R01AI164588 and R01AI176777 to K.A.W.; T32AI007509 to O.C. (PI: Lund); AOA Carolyn Kuckein Student Research Fellowship to M.L.; T32GM007266 to G.M.; and TL1TR002318 to G.M. This work was also supported by the P51OD010425, which is the core grant of the Washington National Primate Research Center. Additional support came from the U42OD011123, which funds the Washington National Primate Research Center breeding colony. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

541 **Table 1. IAV Viral Load in Lung, Placenta, and Clinical Findings**

ID	Clinical Symptoms	Maximum Clinical Score	Maternal Lung Viral Load (copies/mg)	Placenta Viral Load (copies/mg)	Stillbirth
FLU1	Diarrhea: D4, D5	3	2.43×10^6	48.3	No
FLU2	Mild Cough: D4 Dehydration: D5 Moderate Wheezing: D5	3	6.05×10^6	0	No
FLU3	None	0	8.58×10^5	0	No
FLU4	None	0	2.22×10^3	0	No
FLU5	Increased respiratory effort: D2 Mild cough: D3 Not eating: D3, D4	4	2.03×10^6	0	No
FLU6	None	0	5.69×10^5	0	No
FLU7	Decreased appetite: D2	1	2.01×10^4	0	Yes
FLU8	Sneezing: D2, D3 Nasal discharge: D2-D5 Increased respiratory effort: D2-D4 Decreased appetite: D4 Diarrhea: D3, D5	6	1.41×10^6	0	No
FLU9	Nasal discharge: D2, D4 Red nasal discharge: D3, D5 Increased respiratory effort: D2-D5 Decreased appetite: D4 Decreased responsiveness: D5	4	3.17×10^3	64.2	Yes
FLU10	None	0	3.49	0	No
FLU11	Increased respiratory effort: D2, D3 Nasal discharge: D3 Mild coughing: D3	3	8.94×10^5	0	No

542 Clinical symptoms were scored on a 3-point scale with higher scores indicating worse disease
543 and 0 indicating normal. The maximum clinical score was calculated by the sum of the highest
544 recorded scores for each of the 5 clinical areas. Abbreviations: D, day

REFERENCES

1. Chandra, S., Christensen, J., Mamelund, S.E. & Paneth, N. Short-Term Birth Sequelae of the 1918-1920 Influenza Pandemic in the United States: State-Level Analysis. *Am J Epidemiol* **187**, 2585-2595 (2018).
2. Nishiura, H. Excess risk of stillbirth during the 1918-1920 influenza pandemic in Japan. *Eur J Obstet Gynecol Reprod Biol* **147**, 115 (2009).
3. Hardy, J.M., Azarowicz, E.N., Mannini, A., Medearis, D.N., Jr. & Cooke, R.E. The effect of Asian influenza on the outcome of pregnancy, Baltimore, 1957-1958. *Am J Public Health Nations Health* **51**, 1182-1188 (1961).
4. Reid, A. The effects of the 1918-1919 influenza pandemic on infant and child health in Derbyshire. *Med Hist* **49**, 29-54 (2005).
5. Freeman, D.W. & Barno, A. Deaths from Asian influenza associated with pregnancy. *American journal of obstetrics and gynecology* **78**, 1172-1175 (1959).
6. Greenberg, M., Jacobziner, H., Pakter, J. & Weisl, B.A. Maternal mortality in the epidemic of Asian influenza, New York City, 1957. *Am J Obstet Gynecol* **76**, 897-902 (1958).
7. Pierce, M.*, et al.* Perinatal outcomes after maternal 2009/H1N1 infection: national cohort study. *BMJ* **342**, d3214 (2011).
8. Centers for Disease, C. & Prevention. Maternal and infant outcomes among severely ill pregnant and postpartum women with 2009 pandemic influenza A (H1N1)--United States, April 2009-August 2010. *MMWR Morb Mortal Wkly Rep* **60**, 1193-1196 (2011).
9. Haberg, S.E.*, et al.* Risk of fetal death after pandemic influenza virus infection or vaccination. *N Engl J Med* **368**, 333-340 (2013).
10. Rasmussen, S.A., Jamieson, D.J. & Uyeki, T.M. Effects of influenza on pregnant women and infants. *Am J Obstet Gynecol* **207**, S3-8 (2012).
11. Kosmak, G.W. The occurrence of epidemic influenza in pregnancy. *Am J Obstet Gynecol*, 239-251 (1919).
12. Harris, J.W. Influenza Occurring in Pregnant Women: A Statistical Study of Thirteen Hundred and Fifty Cases. *JAMA : the journal of the American Medical Association*, 978-980 (1919).
13. Al-Haddad, B.J.S.*, et al.* The fetal origins of mental illness. *American journal of obstetrics and gynecology* **221**, 549-562 (2019).
14. Barr, C.E., Mednick, S.A. & Munk-Jorgensen, P. Exposure to influenza epidemics during gestation and adult schizophrenia. A 40-year study. *Arch Gen Psychiatry* **47**, 869-874 (1990).
15. Canetta, S.E.*, et al.* Serological documentation of maternal influenza exposure and bipolar disorder in adult offspring. *Am J Psychiatry* **171**, 557-563 (2014).
16. Machon, R.A.*, et al.* Adult schizotypal personality characteristics and prenatal influenza in a Finnish birth cohort. *Schizophr Res* **54**, 7-16 (2002).
17. Machon, R.A., Mednick, S.A. & Huttunen, M.O. Adult major affective disorder after prenatal exposure to an influenza epidemic. *Arch Gen Psychiatry* **54**, 322-328 (1997).
18. Mednick, S.A., Huttunen, M.O. & Machon, R.A. Prenatal influenza infections and adult schizophrenia. *Schizophr Bull* **20**, 263-267 (1994).
19. Mednick, S.A., Machon, R.A., Huttunen, M.O. & Bonett, D. Adult schizophrenia following prenatal exposure to an influenza epidemic. *Arch Gen Psychiatry* **45**, 189-192 (1988).
20. Al-Haddad, B.J.S.*, et al.* Long-term Risk of Neuropsychiatric Disease After Exposure to Infection In Utero. *JAMA Psychiatry* (2019).
21. Regan, A.K.*, et al.* Seasonal Trivalent Influenza Vaccination During Pregnancy and the Incidence of Stillbirth: Population-Based Retrospective Cohort Study. *Clin Infect Dis* **62**, 1221-1227 (2016).

22. Bratton, K.N., Wardle, M.T., Orenstein, W.A. & Omer, S.B. Maternal influenza
immunization and birth outcomes of stillbirth and spontaneous abortion: a systematic
review and meta-analysis. *Clin Infect Dis* **60**, e11-19 (2015).
23. Jeong, S., Jang, E.J., Jo, J. & Jang, S. Effects of maternal influenza vaccination on
adverse birth outcomes: A systematic review and Bayesian meta-analysis. *PLoS One* **14**,
e0220910 (2019).
24. Getahun, D.*, et al.* Association between seasonal influenza vaccination with pre- and
postnatal outcomes. *Vaccine* **37**, 1785-1791 (2019).
25. Giles, M.L., Davey, M.A. & Wallace, E.M. Associations Between Maternal Immunisation
and Reduced Rates of Preterm Birth and Stillbirth: A Population Based Retrospective
Cohort Study. *Front Immunol* **12**, 704254 (2021).
26. Simoes, E.A.F.*, et al.* Trivalent influenza vaccination randomized control trial of pregnant
women and adverse fetal outcomes. *Vaccine* **37**, 5397-5403 (2019).
27. Influenza in Pregnancy: Prevention and Treatment: ACOG Committee Statement No. 7.
Obstet Gynecol **143**, e24-e30 (2024).
28. Razzaghi, H.*, et al.* Influenza, Tdap, and COVID-19 Vaccination Coverage and Hesitancy
Among Pregnant Women - United States, April 2023. *MMWR Morb Mortal Wkly Rep* **72**,
1065-1071 (2023).
29. Vaccine Confidence Falls as Belief in Health Misinformation Grows. (Annenberg Public
Policy Center, University of Pennsylvania, 2023).
30. Cunningham, M.C., Maxson A, Mowson R, Pagan D, Garofalini C, Hall K, Bobo M, McCall
T.C. 2023 Immunization Profile Study. (National Association of County & City Health
Officials, Washington, D.C., 2024).
31. Antonson, A.M.*, et al.* Moderately pathogenic maternal influenza A virus infection disrupts
placental integrity but spares the fetal brain. *Brain Behav Immun* **96**, 28-39 (2021).
32. Van Campen, H.*, et al.* Maternal Influenza A Virus Infection Restricts Fetal and Placental
Growth and Adversely Affects the Fetal Thymic Transcriptome. *Viruses* **12**(2020).
33. Jacobsen, H.*, et al.* Offspring born to influenza A virus infected pregnant mice have
increased susceptibility to viral and bacterial infections in early life. *Nat Commun* **12**, 4957
(2021).
34. Littauer, E.Q.*, et al.* H1N1 influenza virus infection results in adverse pregnancy outcomes
by disrupting tissue-specific hormonal regulation. *PLoS Pathog* **13**, e1006757 (2017).
35. Liong, S.*, et al.* Influenza A virus infection during pregnancy causes immunological
changes in gut-associated lymphoid tissues of offspring mice. *American journal of*
physiology. Gastrointestinal and liver physiology **325**, G230-G238 (2023).
36. Liong, S.*, et al.* Influenza A virus causes maternal and fetal pathology via innate and
adaptive vascular inflammation in mice. *Proc Natl Acad Sci U S A* **117**, 24964-24973
(2020).
37. Hall, O.J.*, et al.* Progesterone-Based Therapy Protects Against Influenza by Promoting
Lung Repair and Recovery in Females. *PLoS Pathog* **12**, e1005840 (2016).
38. Vermillion, M.S.*, et al.* Pregnancy preserves pulmonary function following influenza virus
infection in C57BL/6 mice. *Am J Physiol Lung Cell Mol Physiol* **315**, L517-l525 (2018).
39. Li, M.*, et al.* Non-human Primate Models to Investigate Mechanisms of Infection-
Associated Fetal and Pediatric Injury, Teratogenesis and Stillbirth. *Front Genet* **12**(2021).
40. Adams Waldorf, K.M., Rubens, C.E. & Gravett, M.G. Use of nonhuman primate models to
investigate mechanisms of infection-associated preterm birth. *BJOG* **118**, 136-144 (2011).
41. Bouvier, N.M. & Lowen, A.C. Animal Models for Influenza Virus Pathogenesis and
Transmission. *Viruses* **2**, 1530-1563 (2010).
42. Koday, M.T.*, et al.* Multigenic DNA vaccine induces protective cross-reactive T cell
responses against heterologous influenza virus in nonhuman primates. *PloS one* **12**,
e0189780 (2017).

43. Short, S.J.*, et al.* Maternal influenza infection during pregnancy impacts postnatal brain development in the rhesus monkey. *Biol Psychiatry* **67**, 965-973 (2010).

44. Mayer, A.E. & Parks, G.D. An AGM model for changes in complement during pregnancy: neutralization of influenza virus by serum is diminished in late third trimester. *PloS one* **9**, e112749 (2014).

45. Swieboda, D.*, et al.* Pregnancy Downregulates Plasmablast Metabolic Gene Expression Following Influenza Without Altering Long-Term Antibody Function. *Front Immunol* **11**, 1785 (2020).

46. Garcia-Flores, V.*, et al.* Maternal-Fetal Immune Responses in Pregnant Women Infected with SARS-CoV-2. *Res Sq* (2021).

47. Brown, A.S.*, et al.* Prenatal exposure to maternal infection and executive dysfunction in adult schizophrenia. *Am J Psychiatry* **166**, 683-690 (2009).

48. Khandaker, G.M., Zimbron, J., Lewis, G. & Jones, P.B. Prenatal maternal infection, neurodevelopment and adult schizophrenia: a systematic review of population-based studies. *Psychol Med* **43**, 239-257 (2013).

49. Kapur, R.P.*, et al.* Granular cytoplasmic inclusions in astrocytes and microglial activation in the fetal brain of pigtail macaques in response to maternal viral infection. *Acta Neuropathol Commun* **13**, 55 (2025).

50. Yawn, D.H., Pyeatte, J.C., Joseph, J.M., Eichler, S.L. & Garcia-Bunuel, R. Transplacental transfer of influenza virus. *JAMA : the journal of the American Medical Association* **216**, 1022-1023 (1971).

51. McGregor, J.A., Burns, J.C., Levin, M.J., Burlington, B. & Meiklejohn, G. Transplacental passage of influenza A/Bangkok (H3N2) mimicking amniotic fluid infection syndrome. *Am J Obstet Gynecol* **149**, 856-859 (1984).

52. Davidson, S., Maini, M.K. & Wack, A. Disease-promoting effects of type I interferons in viral, bacterial, and coinfections. *J Interferon Cytokine Res* **35**, 252-264 (2015).

53. Barrat, F.J. & Lu, T.T. Role of type I interferons and innate immunity in systemic sclerosis: unbalanced activities on distinct cell types? *Curr Opin Rheumatol* **31**, 569-575 (2019).

54. Lawrence, C.W., Ream, R.M. & Braciale, T.J. Frequency, Specificity, and Sites of Expansion of CD8+ T Cells during Primary Pulmonary Influenza Virus Infection1. *The Journal of Immunology* **174**, 5332-5340 (2005).

55. Kubelkova, K., Bostik, V., Joshi, L. & Macela, A. Innate Immune Recognition, Integrated Stress Response, Infection, and Tumorigenesis. *Biology (Basel)* **12**(2023).

56. Wu, Y., Zhang, Z., Li, Y. & Li, Y. The Regulation of Integrated Stress Response Signaling Pathway on Viral Infection and Viral Antagonism. *Front Microbiol* **12**, 814635 (2021).

57. Pakos-Zebrucka, K.*, et al.* The integrated stress response. *EMBO Rep* **17**, 1374-1395 (2016).

58. Siston, A.M.*, et al.* Pandemic 2009 influenza A(H1N1) virus illness among pregnant women in the United States. *JAMA : the journal of the American Medical Association* **303**, 1517-1525 (2010).

59. Neuzil, K.M., Reed, G.W., Mitchel, E.F., Simonsen, L. & Griffin, M.R. Impact of influenza on acute cardiopulmonary hospitalizations in pregnant women. *Am J Epidemiol* **148**, 1094-1102 (1998).

60. Liu, S.*, et al.* Avian influenza virus in pregnancy. *Rev Med Virol* **26**, 268-284 (2016).

61. Purcell, R., Giles, M.L., Crawford, N.W. & Buttery, J. Systematic Review of Avian Influenza Virus Infection and Outcomes during Pregnancy. *Emerg Infect Dis* **31**, 50-56 (2025).

62. Itoh, Y.*, et al.* In vitro and in vivo characterization of new swine-origin H1N1 influenza viruses. *Nature* **460**, 1021-1025 (2009).

63. Adams Waldorf, K.M.*, et al.* Congenital Zika virus infection as a silent pathology with loss of neurogenic output in the fetal brain. *Nat Med* **24**, 368-374 (2018).

696   64.   Boldenow, E., Gendrin, C., Ngo, L., Bierle, C., Vornhagen, J., Coleman, M., Merillat, S.,
697   Armistead, B., Whidbey, C., Alishetti, V., Santana-Ufret,V., Ogle, J., Gough, M.,
698   Srinouanprachanh, S., MacDonald, J., Bammler, T., Bansal, A., Liggitt, H., Rajagopal, L.,
699   Adams Waldorf, K. . Group B Streptococcus circumvents neutrophils and neutrophil
700   extracellular traps during amniotic cavity invasion and preterm labor. *Science Immunology*
701   **1**(2016).
702   65.   Reed, L.J. & Muench, H. A simple method of estimating fifty percent endpoints. *Am J Hyg*
703   **27**, 493-497 (1938).
704   66.   Yang, Y.*, et al.* Evaluation of Twelve Real-Time Reverse Transcriptase PCR Primer-Probe
705   Sets for Detection of Pandemic Influenza A/H1N1 2009 Virus. *Journal of Clinical*
706   *Microbiology* **49**, 1434-1440 (2011).
707   67.   O'Connor, M.A.*, et al.* Early cellular innate immune responses drive Zika viral persistence
708   and tissue tropism in pigtail macaques. *Nat Commun* **9**, 3371 (2018).

709