The ecological dynamics and consequences of phytoplasma in a spring ephemeral

Julia A. Boyle[1,2], Yanggaolan Yang[3], Amalia Caballero[4], Eleanor N. Hector[2,5], Alex Ellett,

Emily Glasgow[2], John R. Stinchcombe[2,6], Megan Bontrager[2]

1. Department of Ecology and Evolutionary Biology, University of Michigan, 1105 N University Ave, Ann Arbor, MI, 48109, USA

2. Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks St., Toronto, ON, M5S3B2, Canada

3. Department of Biology, Queen's University, 99 University Ave, Kingston, ON, K7L 3N6, Canada

4. Department of Molecular Genetics, University of Toronto, Medical Science Building, 1 King's College Cir Room 4386, Toronto, ON M5S 3K3, Canada

5. Department of Zoology, Biodiversity Research Centre, University of British Columbia, Vancouver, BC V6T 1Z4

6. Koffler Scientific Reserve, University of Toronto, Toronto, ON Canada, M5S3B2

Keywords: phytoplasma, Trillium grandiflorum, range, density dependent, pathogen, leafhoppers, vector transmission, frequency dependent, Empoasca, white trillium

Data accessibility: Data and code will be found on Dryad at https://doi.org/10.5061/dryad.fqz612k5d. All sequences are available on the NCBI Sequence Read Archive under BioProject PRJNA1321329.

Acknowledgements

We thank KSR Staff from 2018-2024 (Stephan Schneider, John Jensen, Kate Brown, Radana Molnarova, Alanah Joyce, and Julio de Almeida) at the Koffler Scientific Reserve for

their support of our field work, as well as Louisa Bartkovich for help in the field. We thank Chris

Carlson for thoughtful conversations about the manuscript. We also thank the lead course

instructors of EEB405: Temperate Field Ecology in 2018 and 2023, including Ben Gilbert, Art

Weis, and David Punzalan, who helped authors in the development and execution of their ideas.

We thank the Department of Ecology and Evolutionary Biology at the University of Toronto for

supporting the field course and providing staff time and equipment. We thank NSERC for

funding an Undergraduate Student Research Award to ENH, a Doctoral Postgraduate

Scholarship to JAB, and Discovery Grants to JRS and MB. Finally, we acknowledge the

contribution of iNaturalist observers, who are attributed in the supplemental material.

Abstract

Linking within-population dynamics to patterns of species occurrence and persistence is a central

goal of ecology. Theoretical models suggest pathogens have important implications for plant

populations and perhaps even limit plant ranges, though more empirical evidence is needed,

especially in wild plant populations. In the field, we characterized the infection dynamics of

phytoplasma, a vectored and sterilizing bacterial pathogen, in wild populations of white trillium

and identified the bacterial strain and local vectors. We then leveraged community science data

to reveal patterns and potential drivers of phytoplasma occurrence across the range of white

trillium. Within populations, we found that phytoplasma infection presence was more likely in

low host density plots, where insect vectors occupied a higher proportion of trillium, indicative

of an encounter-dilution effect. Leafhoppers in the genus *Empoasca* are the likely vectors of

Candidatus Phytoplasma pruni-related, group 16SrIII infections between trillium. Phytoplasma

was widely distributed across trillium's range, even at range edges, and probability of infection

was significantly reduced by hotter summer temperatures and increased by proximity to

cropland. Our results suggest vectored pathogens are sensitive to local host population densities

and able to persist across both the core and edge of host ranges.

49
50

Introduction

Sessile organisms face a unique challenge in response to transmissible disease, because they cannot disperse away from other infected individuals. As a consequence, for sessile organisms like plants, parasites can have an outsized impact on the persistence of populations and meta-populations through time and space (Gilbert 2002). Plant populations display density-dependent dynamics in interactions with competitors, pollinators, seed predators, herbivores, and others (Gilbert 2002, Gunton and Kunin 2009, Kim et al. 2013, Hegland 2014). The addition of phytopathogens may reinforce or disrupt a plant population's density-dependent processes (Parker and Gilbert 2018), which could lead to altered distribution patterns or demographic structure. Understanding how within-population dynamics scale up to affect species' metapopulation dynamics and geographic distributions has been a central goal of ecological research for decades (Harrison and Taylor 1997, Freckleton et al. 2005, Oldfather and Ackerly 2019), and understanding the impacts of pathogens on these large-scale processes remains challenging. Much of our understanding of plant pathogens comes from agricultural or managed systems, but natural populations may have fundamentally different infection dynamics due to environmental and demographic heterogeneity (Alexander 2010). Disease dynamics in natural populations are also important because wild plant species can act as reservoirs of pathogens (Webster et al. 2007, Petersen et al. 2019) and because disease outbreaks are a conservation concern, even for abundant plant species (Frankel 2008, Pautasso et al. 2013). Here, we characterise a pathogen in naturally-occurring plant populations and leverage community science data to evaluate whether population-level disease dynamics scale up across the range of the host species.

Pathogen success hinges on the properties of host populations, and host population

density often plays an important role at a local scale (Burdon and Chilvers 1982). Density-

dependent dynamics occur when transmission has a linear relationship with the absolute number

of infected individuals (Antonovics 2017); positive density-dependence has been observed in

many wild plant populations, often when pathogens are transmitted aerially or via direct contact

(Carlsson and Elmqvist 1992, Gilbert et al. 1994, Lively et al. 1995, Bell et al. 2006). Density-

dependent transmission may result in different proportions of individuals infected at different

host densities. In contrast, frequency-dependent dynamics occur when transmission increases

with the proportion of hosts that are infected, instead of correlating with host density

(Antonovics 2017, Antonovics et al. 2023). Over time, this may result in similar proportions of

individuals infected across populations of differing density. While vector-transmitted pathogens

are often considered frequency-dependent, they may possess qualities of both density-dependent

and frequency-dependent transmission because transmission can initially increase as host density

increases, but decrease at higher host density (Antonovics et al. 1995, 2017, 2023). Transmission

may decrease at high host density because vectors are limited by time, satiation, or abundance,

reducing the chance that a host interacts with a vector; this phenomenon is known as the

encounter-dilution effect (Mooring and Hart 1992, Krebs et al. 2014). Recently, Antonovics and

colleagues (2023) found that pollinators were more effective transmission vectors of anther-smut

fungus at lower host densities compared to higher host densities, providing evidence of the

encounter-dilution effect in plant-vector-pathogen interactions at relatively small spatial scales.

Scaling up phytopathogen dynamics from a local host population to explore patterns

across a whole host range is empirically challenging and as a consequence, rarely done (Bruns et

al. 2019a, Uricchio et al. 2023). Species ranges are often constrained by abiotic niche space (i.e.,

the climatic and resource parameters a species can tolerate) (Lee-Yaw et al. 2016), and it remains

an open question whether species interactions frequently expand or limit ranges and abiotic

niches (Afkhami et al. 2014, Harrison et al. 2018, Uricchio et al. 2023, Nathan et al. 2023).

Classic models with host-specific pathogens predict that regions of low host density, such as

some range edges, should also have low pathogen loads due to positive density-dependence,

forming a 'disease-free halo' (Antonovics 2009, Bruns et al. 2019a). However, vectored

pathogens conforming to patterns of frequency-dependent infection may also be found even in

areas of low host density, and may be a factor limiting a host's range. Bruns et al. (2019a) found

that a vectored disease was maintained at the range limits of host plant species and occurred at

higher prevalence at range edges compared to range centres. Further work indicates that this

vectored pathogen constrains the range of a host species, and may eventually lead to host range

contraction or extinction (Uricchio et al. 2023). The persistence of populations at range edges is

especially important for the future ability of a species to shift ranges and respond to climate

change (Rehm et al. 2015, Hargreaves and Eckert 2019), thus necessitating a deeper

understanding of how pathogens affect plant populations across host ranges.

Almost 150 years ago, Erwin Smith described a "monstrous form" of *Trillium*

grandiflorum (Smith 1879). Instead of the characteristic white blooms, Smith observed trillium

plants with petals marked with virescence (green pigmentation) and with their leaves more

numerous and held on longer petioles. While Smith suggested these phenotypically irregular

plants might be taxonomically distinct entities, we now know that these morphological variations

reflect symptoms of phytoplasma infection (Hooper et al. 1971). Phytoplasmas are an unusual

group of bacterial parasites that live obligately in the intracellular phloem tissue of plants and the

bodies of their insect vectors, which are often phloem-feeding insects in the order Hemiptera

(Sinha and Chiykowski 1980, Lee et al. 2000, Wang et al. 2024). Phytoplasmas have been found

in over 1000 plant species including those used as timber, ornamental plants, and vegetable, fruit,

and seed crops (Lee et al. 2000), yet their biology remains poorly understood (Liu et al. 2017).

Potential vectors are exposed to phytoplasmas when they consume the phloem of infected plants,

and vectors harbour transmissible levels of infection several weeks after exposure (Christensen et

al. 2005, Koinuma et al. 2020). In addition to inducing morphological changes, phytoplasmas

can sterilize their plant hosts and infection can be systemic and lifelong (Lee et al. 2000, Ustun et

al. 2023). The longevity of infected perennial plants is potentially problematic, as it means these

individuals may act as a source of infection for many years. *Trillium grandiflorum* is one such

long-lived species; individuals become reproductive only after 7-10 years, with a total lifespan of

over 30 years (Hanzawa and Kalisz 1993, Lamoureux and Larose 2002). Theoretical models

predict that sterilizing pathogens in small and long-lived host populations can drive host

populations to extinction more often than lethal pathogens (Anderson & May 1981; Boots &

Sasaki 2001; Gerber et al. 2005; Antonovics 2009). Thus, phytoplasma-induced sterility suggests

there could be grave consequences for trillium and other susceptible plants' population growth

and persistence over time. Smith's (1879) report of a trillium-dwelling phytoplasma was located

in Michigan, USA, suggesting this species interaction has long been established in northeastern

America. Although the first Canadian confirmation of phytoplasma infection in trillium occurred

in the last decade (Arocha-Rosete et al. 2016), observations of the symptoms of phytoplasma

infection in Ontario were documented long before (Gates 1917).

Here, we explore the links between phytoplasma infection and the performance, stage

structure, and density of wild populations of the spring ephemeral *Trillium grandiflorum*. We

further characterize the strain of phytoplasma and its insect vectors at our field site. We then

leverage community science data to examine how patterns in phytoplasma infection within plant

populations scale up across the range of *T. grandiflorum*. We hypothesized that phytoplasma

would reduce trillium fitness and reduce the relative abundance of earlier demographic stage

classes within populations. Given that phytoplasma is sterilizing and vector transmitted, we

hypothesized that its transmission would be primarily frequency-dependent, resulting in an equal

chance and rate of infection in populations regardless of host density. However, we had an

alternative hypothesis that the encounter-dilution effect could reduce transmission at high host

densities due to vector behaviour or limitation, resulting in a lower proportion of infected hosts,

and more consistent with negative density-dependence. Finally, we predicted that phytoplasma

infection would be prevalent even at range edges due to frequency dependent, vector-mediated

transmission. Our work characterizes infection dynamics of a fearsome and understudied plant

pathogen in a wild host population and across the host's range.

154

Figure 1. Diversity of phytoplasma infection symptoms in *Trillium grandiflorum* and two

identified vectors. Symptoms include phyllody and virescence of petals and larger sepals (A-E),

as well as other deformations including extra petals (C) and the absence of pollen (E).

Discolouration in an infected plant late in the flowering season (F). Using sticky traps, we

identified *Empoasca* spp. leafhoppers positive for phytoplasma (G, H). All photos by Julia

Boyle, images of *T. grandiflorum* were taken at the Koffler Scientific Reserve.

Methods

Study organisms and site

Trillium grandiflorum (white trillium), is a spring ephemeral native to the deciduous

woodlands of eastern North America. Its range spans from Minnesota to the Eastern coast, and

from Georgia to northern Ontario (Case 1993). *Trillium grandiflorum* reproduces by seed and

clonal reproduction by rhizomes is rare (Ohara 1989); seeds of this species are frequently

dispersed by ants over short distances (Kalisz et al. 1999). Trillium have four distinct

demographic stages: a seedling stage, a single leaf vegetative stage, a three leaf vegetative stage,

and a reproductive stage (Ohara 1989, Rooney and Gross 2003). Once a trillium has first

bloomed it often continues to bloom annually, but can also go dormant for one or more years

(persisting as a rhizome with no aboveground structures) or revert to a three-leaf non-

reproductive stage. In Ontario, white trillium begins flowering in mid-May and sets seed by July,

after which the aboveground parts of the plant die back. There are large populations of wild *T.*

grandiflorum at our study site (Figure S1), the Koffler Scientific Reserve (KSR,

www.ksr.utoronto.ca) in Ontario, Canada (44°01'48''N, 79°32'01''W). Symptoms of phytoplasma

infection have been observed in trillium populations at KSR since at least the early 2000s (A.

Weis, personal communication). Phytoplasma infection is most apparent when the white trillium

are flowering, as petals will range from having a streak of green to full virescence or phyllody

(Figure 1). The strain of phytoplasma previously identified in a different *T. grandiflorum*

population was *Candidatus* Phytoplasma pruni-related, group 16SrIII, otherwise known as

Western X-disease or peach X-disease (Arocha-Rosete et al. 2016).

Effects of phytoplasma infection on Trillium grandiflorum growth and reproduction

To investigate whether infection was associated with differences in plant size, we

quantified total plant height, leaf length, and sepal length in symptomatic (n = 32) and

asymptomatic (n = 30) trillium during peak flowering season (May 20-29, 2023) across 30

locations in an approximately 15 hectare section of mixed deciduous forest at KSR. We

determined infection status by visually inspecting petals for virescence; we suspect that some

infected individuals have no visible symptoms so this method likely underestimates the overall

infection rate of the population. To test for phenotypic differences in plant height, leaf length,

and sepal length associated with infection, we used linear mixed effects models with infection

status as a predictor and location as a random effect. We applied a natural log transformation on

responses as needed to improve normality. For statistical analysis, we used R v4.2.0 (R Core

Team 2022), with the *tidyverse* (Wickham et al. 2019), *lme4* (Bates et al. 2015), and *lmerTest*

(Kuznetsova et al. 2017) packages. All linear models were corrected using a type III ANOVA

with the *car* package (Fox and Weisberg Sanford 2019).

To quantify differences in seed production due to infection, we first characterized the

relationship between leaf length and seed number in asymptomatic trilliums. We haphazardly

sampled 100 individuals on 13 June 2024, aiming to maximize variation in plant size. On these

plants we measured leaf length and dissected fruits to count seed set. With these data we

estimated the relationship between leaf length and seed count using a generalized linear model

with a Poisson family. We also measured leaf length (May 20-28, 2024) and counted the number

of seeds (early to mid June 2024) produced by 81 haphazardly chosen symptomatic trillium. To

estimate potential reproductive loss due to infection in a given environment, we identified the

nearest asymptomatic trillium of similar size and estimated its predicted seed count from the

relationship between leaf length and seed number using the predict.glm function; logistical and

time constraints prevented us from also sampling seeds from these asymptomatic individuals and

determining this relationship directly. Then, we used a negative binomial generalized linear

mixed model with estimated seed count (rounded to the nearest whole number) as a response,

infection status as a fixed effect, and pair number as a random effect. To determine whether

symptomatic plants produced fewer seeds than expected based on their size, we predicted the

expected number of seeds symptomatic trillium would produce if they were asymptomatic using

the predict.glm function, then we used a negative binomial generalized linear mixed model with

rounded seed count as a response, whether the counts were modeled or counted as a fixed effect,

and plant ID as a random effect.

Within-population infection dynamics

We conducted surveys of trillium density and infection frequency in *T. grandiflorum*

populations of KSR in mid-late May 2018 and 2023 (populations indicated in Figure S2). In both

survey years, 1 m^2 plots were chosen haphazardly by throwing a flag into trillium populations

(Figure S1); all plots were >5 m apart and plot locations differed in the two sampling years. To

account for spatial proximity of plots, we categorized general sampling locations based on

geographic features (e.g. separate hill slopes). In 2018, we sampled 210 plots; we counted the

number of flowering *T. grandiflorum* in the plot and their infection status based on petal

virescence. In 2023, we sampled 64 additional plots with the aim of including more high density

populations and surveying potential vectors. We counted the number of non-flowering and

flowering *T. grandiflorum* individuals that had at least three leaves, noted whether phytophagous

Hemiptera nymphs were present on the plants, and noted the infection status of flowering

trillium. Across both surveys, we estimated the proportion of trillium infected by dividing the

number of symptomatic flowering plants by the total number of flowering trilliums (we used

only flowering individuals in the denominator because infection was determined by visual petal

symptoms). In addition, using data from our 2023 survey, we calculated the proportion of

trillium with phytophagous Hemipterans present by dividing the number of trillium with

Hemiptera by the number of trillium in the plot; we did this for total trilliums and also for only

flowering trillium.

To test whether the number of symptomatic trillium was dependent on the number of

flowering trillium in a patch and sampling year, we used zero-inflated poisson models with

(*GLMMadaptive*; Rizopoulos 2025) and without (*pscl*; Zeileis et al. 2008) a random effect of

categorical location. These zero-inflated models predict both the zeros (infection presence or

absence), and the counts (number of infected trillium) separately. We compared the log

likelihood of models with and without random effects and used the better fitting model. Then,

using data from 2023 only, we used the same models without the year predictor and added the

proportion (values between 0-1) of flowering trillium with Hemiptera as an additional predictor.

To further explore patterns of trillium density and potential vector occurrence in the 2023 data,

we used a binomial generalized linear model with the proportion of trillium with Hemiptera as

the response and the total number of trillium as the predictor. Additionally, we used a binomial

generalized linear model to test whether the proportion of symptomatic adults in a plot was

associated with the number of adult trillium with Hemiptera on them.

To determine how phytoplasma infection affected the age class distribution of KSR

trillium populations, we calculated plot level ratios of vegetative to reproductive trillium and

tested for correlations with infection frequency. We used ratios because they are agnostic to

trillium density. Using data from the 1 m^2 plots surveyed in 2023, we calculated the ratio of

vegetative to flowering three-leaved trillium and tested for a correlation with infection frequency

using cor.test (*stats*; R Core Team 2022). In 2024, we sampled 0.5 m^2 plots centered around the

asymptomatic and symptomatic individuals measured for seed production, and recorded the

number of one-leaved, three-leaved vegetative, and three-leaved flowering trillium. We

calculated the ratio of one-leaved to three-leaved flowering trillium, then tested for a correlation

between this ratio and the frequency of infection among three-leaved flowering trillium. Using

this 2024 data we also tested the correlation between the ratio of vegetative to flowering three-

leaved trillium and infection frequency.

Identification of vectors and phytoplasma strain

To identify the phytoplasma present in trillium, we sampled a leaf from a symptomatic

flowering plant and extracted DNA using QIAGEN's DNeasy® PowerSoil® Pro Kit. To assess

vector identity and abundance, we deployed yellow sticky traps (15.24 cm x 25.4 cm) attached to

bamboo poles just above trillium height at 0.5 m above ground (Figure S1). We placed 20 traps

across the sampled KSR trillium regions on June 6th 2023, collected them after 6 days, and then

stored traps at -20□. We counted and collected potential vectors from each trap, including

leafhopper, psyllid, aphid, and planthopper morphotypes. We chose 110 insects across potential

vector morphotypes and extracted their DNA using QIAGEN's DNeasy® Blood & Tissue Kit,

following the protocol for insect specimens. We detected and amplified the 16S rRNA region of

phytoplasma in insect and plant samples with a nested polymerase chain reaction (PCR) using

two universal phytoplasma primer pairs R16mF2/R1 and R16F2n/R2 (Gundersen & Lee 1996),

and visualised the presence of phytoplasma using gel electrophoresis. We amplified the

mitochondrial cytochrome oxidase subunit 1 region of insect samples positive for phytoplasma

using the lib220-1/LepR primers (Hajibabaei et al. 2006). Then, to identify the phytoplasma

present in plants and the insects that vector it, we performed Sanger sequencing on PCR products

(at the Center for Analysis of Genome Function and Evolution, University of Toronto) and

looked for similar sequences using the National Center for Biotechnology Information's

nucleotide basic local alignment search tool using the megablast algorithm (Altschul et al. 1990,

Chen et al. 2015).

Metapopulation infection dynamics

To characterize spatial patterns of phytoplasma infection of *T. grandiflorum* at a range-

wide scale, we collected data from user observations on iNaturalist. First, we annotated all

visually symptomatic *T. grandiflorum* observations with the observation field "Phytoplasma

Infection?=Yes". On October 7th, 2025, we downloaded all research-grade iNaturalist

observations of *T. grandiflorum* from the Global Biodiversity Information Facility (iNaturalist

contributors, iNaturalist 2025) and merged them with their associated metadata and observation

fields from iNaturalist, downloaded on the same day (data included in supplementary material).

In total, this dataset included 36,497 observations between 2012-2025 with 637 infected

individuals. We found that our results were qualitatively unchanged between when we used all

research-grade observations in iNaturalist and only those with only CC0, CC-BY, and CC-BY-

NC copyrights (Boyle, unpublished analyses). We removed observations with missing latitudes

or longitudes, observations with low or missing positional accuracy (< 10,000), and observations

likely to be erroneous with both a latitude above 20°N and a longitude below 65°W, resulting in

27,803 observations. To reduce user bias in the dataset, we limited the maximum number of

observations per user to 15, reducing the dataset to 24,103 observations. Then, we used

CoordinateCleaner (Zizka et al. 2019) to remove observations at country centroids or known

biological institutions, observations with equal latitudes and longitudes, observations with any

zeroes as coordinates, and observations with identical coordinates. After cleaning, the dataset

had 23,218 total observations and 384 infected observations. Users contributed a mean 1.89 total

observations and a median of 1 total observation. The infected observations were observed by

319 users. The vast majority (88.6%) of all observations in the dataset occurred after January 1st,

2020. We reprojected the coordinates of observations using ESRI:102003, an equal area

projection that is appropriate for *T. grandiflorum*'s range.

Next, we examined patterns of trillium observations across its range, and linked trillium

observation density, as well as other environmental and anthropogenic factors, to phytoplasma

infection using Bayesian spatial models. First, we assigned observations to hexagonal grid cells

drawn at a scale of 50 km between opposite sides to quantify trillium density across the range.

Then, we tested for significant spatial clustering of trillium observations and trillium infections

using Moran's I and the moran.mc function (*spdep*; Pebesma and Bivand 2023). We visualised

grids using the *sf* (Pebesma 2018), *mapview* (Appelhans et al. 2015), and *rnaturalearth*

(Massicotte and South 2017) packages. Then, for each observation, we used the *geodata* package

(Hijmans et al. 2025) to extract 30-seconds spatial resolution of worldclim bioclimatic variables,

cropland distribution (Zanaga et al. 2021), and human footprint circa 2009 (Venter et al. 2016).

Next, we used deterministic Bayesian Integrated Nested Laplace Approximation (INLA) models

to determine factors affecting trillium infection (Rue et al. 2009). INLA models can effectively

account for spatial autocorrelation of data and model continuous spatial processes like disease.

We used a model selection procedure to exclude predictors that increased model deviance

information criterion (DIC) by >2 (*ggregplot*; Albery 2025), but we did not exclude predictors

that we had an *a priori* reason to include, such as footprint and trillium observation density.

Then, we removed multicollinear variables using vifcor (*usdm*; Naimi et al. 2014), until all

predictors had a variance inflation factor below 5. We fit the base model using cropland,

footprint, total trillium number within the observation's 50 km grid cell, mean diurnal range,

temperature seasonality, mean temperature of wettest quarter, mean temperature of driest quarter,

mean temperature of warmest quarter, precipitation seasonality, and precipitation of wettest

quarter. Then, we fit a model incorporating spatial autocorrelation using a stochastic partial

differentiation equation (SPDE) random effect (Lindgren et al. 2011); we set priors following

Albery et al. (2020). When the 2.5% and 97.5% posterior estimates of predictors did not overlap

zero, we considered this strong evidence that the predictor affected infection. We used the

Efxplot and ggField functions (*ggregplot*; Albery 2025) to plot the INLA model fixed results and

the spatial field, which represents the marginal posterior mean effects of the spatial random

effects on infection.

Results

Trillium grandiflorum growth and reproduction is reduced by infection

In addition to altered petal colour and flower morphology (Figure 1), phytoplasma-

infected trillium also showed stunted growth and reproduction. Symptomatic trillium were on

average 15.4% shorter than asymptomatic trillium ($X^2 = 8.77$, $p = 0.003$; Figure S3, Table S1), a

difference of around 4.26 cm. While the length of leaves was not significantly different ($X^2 =$

0.491, $p = 0.484$; Table S1), sepals were on average 28% longer in symptomatic individuals (X^2

$= 25.6$, $p < 0.001$; Figure S3, Table S1). Phytoplasma infection significantly reduced trillium

seed set, with symptomatic individuals producing on average 68.4% fewer seeds than expected

for asymptomatic trillium ($X^2 = 34.7$, $p < 0.001$; Figure 2A, Table S2) and producing 64.6 %

fewer seeds than expected based on their size ($X^2 = 31.3$, $p < 0.001$; Figure 2A, Table S2). 63%

of symptomatic trillium produced no seeds.

Despite the effects of infection on seed production, the relative abundance of trillium

stage classes remained stable across patches varying in infection frequency. We found no

significant correlation between the infection frequency among flowering trillium in a plot and the

ratio of vegetative to flowering three-leaved trillium in 2023 ($r = 0.103$, $p = 0.412$) or 2024 ($r = -$

0.079, $p = 0.352$). Additionally, infection frequency did not significantly affect the ratio of one-

leaved trillium to flowering three-leaved trillium ($r = -0.082$, $p = 0.336$; Figure 2B).

Figure 2. Phytoplasma infection reduced trillium reproduction but the relative abundances of

stage classes remained intact. Symptomatic plants made fewer seeds than expected based on their

size and compared to nearby asymptomatic trillium (A). The boxplots show the median with the

lower and upper hinges corresponding to the 25th and 75th percentiles, and the upper and lower

whiskers represent the largest or smallest value, respectively, that is no further than 1.5 times the

inter-quartile range from the hinge. Individual data points represent either direct seed counts

("counted") or seed counts predicted based on an empirical relationship between leaf size and

seed set ("modeled"); these points are shown at lower opacity superimposed on the boxplots. The

ratios of one-leaved trillium to flowering three-leaved trillium were similar across plots of

varying infection rates in 2024, suggesting that local infection rate does not detectably alter

recruitment (B). The scatterplot shows data points and a linear model fit, statistical values are

from the correlation test.

Negative density-dependent infection within populations

Phytoplasma infection within populations was more likely at low densities of trillium,

indicating negative density-dependent dynamics (Figure 3A). Our survey plots included a wide

range of densities (2-74 reproductive trillium per $1m^2$) and infection prevalence of flowering

trillium ranged from 0% to 21% among plots. The zero-inflation component of the model

indicated that the presence of infection was significantly more likely when flowering trillium

were at low density (z-value = 2.56, p = 0.010; Table S3) and more likely in 2023 compared to

2018 (z-value = -2.07, p = 0.038; Table S3). The count component of the model found no

significant effect of year (z-value = 0.377, p = 0.706; Table S3) or flowering trillium density (z-

value = 1.81, p = 0.069; Table S3) on the number of infected trillium in a plot. Thus, both the

probability of infection presence and the proportion of infected individuals was higher at lower

densities. When we modelled data from only 2023 and included the prevalence of Hemiptera on

flowering trillium as a predictor in the model, we found that an increased prevalence of

Hemiptera was marginally associated with an increase in the number of symptomatic trillium (z-

value = 1.86, p = 0.063; Table S4). Consistent with our analysis of both years, we found that

symptomatic trillium were more likely to be present in plots with low densities of reproductive

stage trillium (z-value = 2.22, p = 0.026; Table S4). The zero-inflated model characterising

infection across both sampling years had a better fit with the inclusion of categorical location as

a random effect (log likelihood of -254.8 compared to -272.1), but for models using only 2023

data we excluded random effects of categorical location as this improved fit (log likelihood -82.6

compared to -83.3).

The density of trillium in patches influenced the prevalence of insect vectors on trillium

hosts. As total trillium density increased in a plot, the proportion of total trilliums with

Hemiptera nymphs decreased (z-value = -10.786, $p < 0.001$, Figure 3C, Table S5). As the

proportion of flowering trillium with Hemiptera increased in a plot, the proportion of

symptomatic trillium also increased (z-value = 4.65, $p < 0.001$, Figure 3D, Table S5). Hemiptera

were highly prevalent and occurred on 42% to 100% of flowering trillium (Figure 3D).

Figure 3. Density-dependent phytoplasma infection patterns within temperate forest populations

at Koffler Scientific Reserve (Ontario, Canada). Plot A uses a kernel density estimate to show

the density of sampled plots along different trillium densities, coloured by whether there were

one or more symptomatic trilliums in the plot. At lower trillium densities infection presence in a

plot is more likely, while at higher trillium densities, plots are less likely to contain infected

trillium (A); significance values are from the binomial component of a zero-inflated model.

When infection was present in a plot, the number of infected trillium was not significantly

predicted by trillium density or year (B); significance values are from the count component of a

zero-inflated model. In 2023, trillium in less dense patches were more likely to be found with

phytophagous Hemiptera nymphs (C), and rates of infection were higher in plots where more

plants had phytophagous Hemiptera (D). The scatterplots in C and D show data points and fits

from binomial generalised linear models.

Phytoplasma strain was vectored by multiple leafhopper morphotypes

The sequencing identified the strain of phytoplasma in *T. grandiflorum* plants at KSR as

Candidatus Phytoplasma pruni-related, group 16SrIII; the closest identity match was to the

phytoplasma previously identified in white trillium by Arocha-Rosete et al. (2016). In total, we

captured 906 insects in the Order Hemiptera from the sticky traps, including 14 morphotypes of

leafhoppers, 3 leafhopper nymph morphotypes, 2 psyllid morphotypes, 2 aphid morphotypes, 2

treehopper morphotypes, and whiteflies. Of the 110 insects tested for phytoplasma presence, we

found 4 individuals across 2 leafhopper morphotypes (Hemiptera: Cicadellidae) that carried

phytoplasma infection (Figure 1GH; Figure S4; Table S6). The sequences of the four

phytoplasma-positive insect specimens most closely matched leafhoppers in the *Empoasca*

genus. Further details are provided in the supplement (Figure S4; Table S6; Supplemental file)

and all sequences are available on the Sequence Read Archive (BioProject PRJNA1321329).

Phytoplasma is prevalent across Trillium grandiflorum range

415   Trillium observations and infection were spatially structured and clustered across the

landscape (Monte-Carlo simulation of Moran's I $p = 0.001$ for each, Figure 4AB). Infected

trillium observations were more clustered than trillium observations at large (Moran's I of 0.376

compared to 0.308), and infection was prevalent across Michigan in the United States of

America, and southern Ontario in Canada (Figure 4B). We found observations of phytoplasma

infection at most putative range edges of trillium, even when there were relatively few

observations in total (Figure 4B). The INLA model spatial field, representing the marginal

posterior mean effects of the spatial random effects on infection, indicated a high likelihood of

infection wherever trilliums were and across the range of *T. grandiflorum*, especially in the

central and western part of the range (Figure 4C; standard deviation shown in Figure S5). The

spatial model indicated that higher mean temperatures of the warmest quarter reduced infection

(mean effect -0.451, credible interval of -0.721 to -0.191) and cropland cover increased infection

(mean effect 1.49, credible interval of 0.833 to 2.14) (Figure 4D; Table S7). All other predictors

in the spatial model did not affect infection, including the number of trillium observations

nearby. All predictors had credible intervals non-overlapping with 0 in the base INLA model

without spatial structure, except observation density, precipitation seasonality, and mean

temperature of the wettest and driest quarters (Figure 4D; Table S8). INLA model fit was

substantially improved by incorporating spatial random effects (ΔDIC = 230).

Figure 4. Patterns and predictors of phytoplasma infection across the range of *Trillium*

grandiflorum. Observations span from 2012 to 2025. The dataset had 23,218 observations (A)

with 384 observations of phytoplasma-infected plants (B). The hexagonal grid in (A) and (B)

represents areas with at least one observation of *T. grandiflorum*; shading in (B) represents the

proportion of observations in that hexagon that show symptoms, and white hexagons had no

infected observations. Hexagons are drawn at a scale of 50 km between opposite sides. The

INLA spatial field (C) represents the marginal posterior mean effects of the spatial random

effects on infection (C), in other words, it shows the effect that spatial position has on infection

accounting for all other covariates in the model. Spatial effects on infection were generally

positive, particularly in the western part of the range (C). Cropland cover and mean temperature

of the warmest quarter significantly increased and decreased trillium infection respectively in the

model that accounted for spatial autocorrelation (SPDE) (D). Asterisks in (D) indicate that the

effect and its confidence intervals do not overlap zero in that model; the model is indicated by

asterisk colour, with light grey representing the base model and dark green representing the

spatial model. Abbreviations are as follows: min. = minimum, temp. = temperature, precip. =

precipitation, quart. = quarter.

Discussion

Ecologists have long sought to link within-population dynamics to patterns of species occurrence and persistence across the range. Theoretical models suggest both pathogens and their transmission mode have important implications for plant populations and perhaps even limit plant distributions, though more empirical evidence is needed. We studied the impacts of a sterilizing, vector-transmitted pathogen on white trillium populations and patterns of infection across trillium's range. Within populations, we found that phytoplasma infection presence was more likely in low host density plots, where insect vectors occupied a higher proportion of trillium. The absolute number of symptomatic trillium was similar across all plots that had infection present, such that low density plots had a greater proportion of infected individuals. Phytoplasma infection was widely distributed across trillium's range, even at range edges, and was influenced by temperature and proximity to cropland. Below, we discuss the ecological processes at work, and their potential consequences for white trillium and other plant populations.

Patterns and processes of phytoplasma within and across white trillium populations

One possible mechanism driving negative density-dependent patterns within populations is the encounter-dilution effect (Mooring and Hart 1992, Krebs et al. 2014), wherein vectors may satiate or saturate or are otherwise unable to occupy all hosts at high host densities (Gilbert 2002). Indeed, our results support vector limitation because Hemipteran nymphs occupied a lower proportion of trillium at higher host densities (Figure 3C), and nymph presence is likely a good direct and indirect proxy for vector visitation and contact. As suggested by theory and other recent empirical work (Antonovics et al. 1995, 2023, Bruns et al. 2019b), our results provide support for the idea that the encounter-dilution effect, a density-dependent process, affects the

outcome of vectored disease at ecologically realistic and unmanipulated plant densities. It is also

possible that infected trillium could experience higher mortality or regression to dormancy than

non-infected individuals (M. Bontrager, *personal observation*), leading to sparser patches over

time where transmission is high, and further compounding the encounter-dilution effect. Higher

mortality or regression did not lead to changes in stage class structure however, which we

discuss more below. If phytoplasmas are transmitted in a negative density-dependent manner,

they may be more difficult to eradicate from host populations, because efforts to reduce their

abundance by removing infected hosts can lead to the very conditions that promote the fastest

phytoplasma population growth, ultimately increasing their resilience (Churcher et al. 2006).

The community science data and the spatial model demonstrated that this vectored

pathogen is not constrained to specific regions of the host range, providing evidence that disease

can be maintained at both the core and edges (Bruns et al. 2019a) and potentially limit host

species ranges. There may be several contributing factors creating this pattern. First, if vectored

pathogens are more able to colonize small and low-density host populations compared to other

types of pathogens, that may lead to reservoirs of infection that bridge populations. Secondly,

while leafhoppers often disperse a few dozen meters a day (Zhou et al. 2003), they can

occasionally disperse long distances using wind-mediated migration (Santos et al. 2024),

meaning that this vectored pathogen may be less dispersal-limited compared to aerially or

contact transmitted pathogens. Finally, vectored pathogens also benefit from the intelligence of

their vector, as many insects are experts at seeking out their host plants in the environment

(Webster and Cardé 2017, Jones et al. 2022). At the landscape scale, vectors may find plant hosts

based on the frequency of the host relative to other plant species (Antonovics et al. 1995),

aligning with processes occurring in frequency-dependent transmission. We did not design our

study to explicitly test whether phytoplasma infection at the range scale correlates with trillium

density because our range-wide dataset consisted of observations—and observation density does

not necessarily correlate with trillium population size or density. Regardless, our work suggests

that phytoplasma transmission maintains infection even at range edges, and motivates more

experimental work to understand range-level drivers of vectored pathogens.

Several environmental factors predicted phytoplasma infection at the range-wide scale.

We found that higher mean temperatures of the warmest quarter correlated with reduced

infection, which was unexpected given the hypothesized relationship between heat stress and

plant infection susceptibility (Hector et al. 2022). High heat could correlate with some other

aspect of phytoplasma ecology, perhaps through vector activity levels or phenology, or cool

areas may coincide with historical regions of disease introduction. Higher temperatures

correlating with reduced infection may indicate that pathogen-host relationships will not respond

uniformly to warming. Additionally, we found that cropland cover increased infection in *T.*

grandiflorum. Phytoplasma is a major problem in agricultural crops (Munyaneza and Henne

2013, Rao et al. 2018), and our result suggests that infection may spread from farms and into

wild populations, and vice versa, and that the successful management of phytoplasma must take

into account both of these environments.

Caveats of spatial data

Combining community science data and hypothesis-driven field work is an especially

powerful way to understand disease dynamics and patterns in plants (Meentemeyer et al. 2015).

Using community science data allowed us to survey phytoplasma infection in trillium across an

otherwise intractable number of plants and locations. The observations showed spatial clustering

likely due to the real structure of trillium populations as well as biases like human population

density and location accessibility, however these biases did not manifest in the INLA model as

strong effects (e.g. through footprint or the number of nearby trillium observations). It is possible

that symptomatic trillium were more likely to be recorded as an iNaturalist observation, but this

works to our favour by increasing the likelihood of phytoplasma detection in low-density

populations. So, while there are limitations to what we can infer with our spatial data—for

example, infection frequency within populations across the range—the symptomatic phenotype

presence remains a strong indicator of phytoplasma presence across the majority of trillium

range.

Implications of phytoplasma on plant ecology and evolution

Given the high fitness cost accrued with phytoplasma infection and given that

leafhoppers transmit the parasite through a form of herbivory, we predict that plant traits

involved in herbivore attraction and defense are under selection (Mauricio and Rausher 1997,

Agrawal 2011). However, white trillium does not appear to have many physical defenses against

herbivory and its chemical defense remains unknown (Figure 1). Deer herbivory is known to

play a role in structuring trillium populations (Knight 2003, 2007), in contrast, insect herbivory

remains largely unstudied (but see Rathcke 1985). While it is clear phytoplasma affects fitness

components and thus has the potential to change the selective landscape of plants, more research

is needed to understand if there is any subsequent phytoplasma-mediated selection on plant traits

in trillium populations, and how this interacts with other important processes such as growth,

pollination, and seed dispersal.

Implications of phytoplasma on wild plant population persistence

Infection may have consequences for trillium population persistence. Because phytoplasma is sterilizing and prevalent across trillium's range, it may reduce white trillium population size in both core and edge populations (Antonovics 2009). Drift, which erodes genetic variation over time, is stronger in small populations and (Star and Spencer 2013), meaning that the adaptive capacity of populations may be negatively impacted by prevalent infection. If edge populations are smaller or more isolated than central populations, they may particularly struggle under further reduced population sizes, lower adaptability, and high risk of infection (Bahn et al. 2006). Edge populations can harbour genotypes adapted to a wider range of abiotic conditions (Rehm et al. 2015), and their preservation may be important for the ability of white trillium to expand its range and track shifting climates. Additional populations of conservation concern are the endangered species of trillium that overlap in range with *T. grandiflorum* (Meredith et al. 2022). We have reason to suspect many *Trillium* species are susceptible to phytoplasma because both *T. grandiflorum* and *T. erectum* were identified as hosts in the first disease report by Arocha-Rosete et al. (2016).

Despite theory predicting high rates of local extinction due to a sterilizing pathogen in a long-lived species (Anderson and May 1981, Boots et al. 2002, Gerber et al. 2005, Antonovics 2009), the trillium populations at KSR remain productive after hosting phytoplasma for more than 20 years, though the amount of change in spatial structure, absolute trillium abundance, and infection prevalence over that time remains unknown. Furthermore, at the patch level, the relative age class distribution of KSR trillium was not affected by infection as hypothesized. We may not see a shift in demographic structure because long-lived populations could experience high lag, or because the KSR populations remain large enough that the reduced seed input is not

yet materially affecting recruitment rates. Our plots were smaller than the seed dispersal kernel

of trillium (Kalisz et al. 1999, Vellend et al. 2003), so it's likely that plots with high infection

proportions receive seed subsidies from surrounding areas. It could also be possible that

reproductive infected trillium are dying or regressing to dormant stages at the same rate that

younger classes are not being recruited; Knight (2003) found that deer herbivory regressed

trillium to non-reproductive states, and a similar stress response may occur with phytoplasma

infection. Another alternative explanation is that the trillium continue to reproduce by rhizomes,

and that this masks the demographic consequences of plants producing fewer seeds. However,

more research should be done to ascertain whether infected trillium continue to produce

rhizomes and whether rhizomes perpetuate the systemic infection (Lee et al. 2000, Ustun et al.

2023). Multiple modes of transmission may compound the difficulties small populations may

face (Uricchio et al. 2023), and lead to an even higher likelihood of local extinction while

providing a veneer of stability. The full demographic consequences of phytoplasma infection

may have yet to materialize.

Literature cited

Afkhami, M. E., P. J. McIntyre, and S. Y. Strauss. 2014. Mutualist-mediated effects on species'

range limits across large geographic scales. Ecology Letters 17:1265–1273.

Agrawal, A. A. 2011. Current trends in the evolutionary ecology of plant defence. Functional

Ecology 25:420–432.

Albery, G. 2025. ggregplot: R package. https://github.com/gfalbery/ggregplot.

Albery, G. F., C. Newman, J. B. Ross, D. W. MacDonald, S. Bansal, and C. Buesching. 2020.

Negative density-dependent parasitism in a group-living carnivore. Proceedings of the Royal

Society B: Biological Sciences 287:20202655.

Alexander, H. M. 2010. Disease in Natural Plant Populations, Communities, and Ecosystems:

Insights into Ecological and Evolutionary Processes. Plant Disease 94:492–503.

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment

search tool. Journal of Molecular Biology 215:403–410.

Anderson, R. M., and R. M. May. 1981. The population dynamics of microparasites and their

invertebrate hosts. Philosophical Transactions of the Royal Society of London. B, Biological

Sciences 291:451–524.

Antonovics, J. 2009. The effect of sterilizing diseases on host abundance and distribution along

environmental gradients. Proceedings of the Royal Society B: Biological Sciences 276.

Antonovics, J. 2017. Transmission dynamics: critical questions and challenges. Philosophical

Transactions of the Royal Society B: Biological Sciences 372:20160087.

Antonovics, J., C. R. Amoroso, E. Bruns, and M. Hood. 2023. Host density shapes the relative

contribution of vector-based and aerial transmission of a pathogenic fungus. Ecology 104:e3970.

Antonovics, J., Y. Iwasa, and M. P. Hassell. 1995. A Generalized Model of Parasitoid, Venereal,

and Vector-Based Transmission Processes. The American Naturalist 145:661–675.

Antonovics, J., A. J. Wilson, M. R. Forbes, H. C. Hauffe, E. R. Kallio, H. C. Leggett, B.

Longdon, B. Okamura, S. M. Sait, and J. P. Webster. 2017. The evolution of transmission mode.

Philosophical Transactions of the Royal Society B: Biological Sciences 372:20160083.

Appelhans, T., F. Detsch, C. Reudenbach, and S. Woellauer. 2015, December 15. mapview:

Interactive Viewing of Spatial Data in R.

Arocha-Rosete, Y., N. P. Morales-Lizcano, A. Hasan, K. Yoshioka, W. Moeder, R. Michelutti,

E. Satta, A. Bertaccini, and J. Scott. 2016. First report of the identification of a 'Candidatus

Phytoplasma pruni'-related strain in *Trillium* species in Canada. New Disease Reports 34:19–19.

Bahn, V., R. J. O'Connor, and W. B. Krohn. 2006. Effect of dispersal at range edges on the

structure of species ranges. Oikos 115:89–96.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models

Using lme4. Journal of Statistical Software 67:1–48.

Bell, T., R. P. Freckleton, and O. T. Lewis. 2006. Plant pathogens drive density-dependent

seedling mortality in a tropical tree. Ecology Letters 9:569–574.

Boots, M., A. Sasaki, and E. J. Travis. 2002. Parasite☐Driven Extinction in Spatially Explicit

Host☐Parasite Systems. The American Naturalist 159:706–713.

Bruns, E. L., J. Antonovics, and M. Hood. 2019a. Is there a disease-free halo at species range

limits? The codistribution of anther-smut disease and its host species. Journal of Ecology 107:1–

11.

Bruns, E. L., I. Miller, M. E. Hood, V. Carasso, and J. Antonovics. 2019b. The role of infectious

disease in the evolution of females: evidence from anther-smut disease on a gynodioecious

alpine carnation. Evolution; international journal of organic evolution 73:497–510.

Burdon, J. J., and G. A. Chilvers. 1982. Host Density as a Factor in Plant Disease Ecology.

Annual Review of Phytopathology 20:143–166.

Carlsson, U., and T. Elmqvist. 1992. Epidemiology of anther-smut disease (Microbotryum

violaceum) and numeric regulation of populations ofSilene dioica. Oecologia 90:509–517.

Case, F. W. 1993. Trillium grandiflorum. Page Flora of North America North of Mexico

[Online]. New York and Oxford.

Chen, Y., W. Ye, Y. Zhang, and Y. Xu. 2015. High speed BLASTN: an accelerated

MegaBLAST search tool. Nucleic Acids Research 43:7762–7768.

Christensen, N. M., K. B. Axelsen, M. Nicolaisen, and A. Schulz. 2005. Phytoplasmas and their

interactions with hosts. Trends in Plant Science 10:526–535.

Churcher, T. S., J. A. N. Filipe, and M. G. Basáñez. 2006. Density dependence and the control of

helminth parasites. Journal of Animal Ecology 75.

Fox, J. and Weisberg Sanford. 2019. An R Companion to Applied Regression. Sage

Publications.

Frankel, S. J. 2008. Sudden oak death and Phytophthora ramorum in the USA: a management

challenge. Australasian Plant Pathology 37:19–25.

Freckleton, R. P., J. A. Gill, D. Noble, and A. R. Watkinson. 2005. Large□scale population

dynamics, abundance–occupancy relationships and the scaling from local to regional population

size. Journal of Animal Ecology 74:353–364.

Gates, R. R. 1917. A Systematic Study of the North American Genus Trillium, Its Variability,

and Its Relation to Paris and Medeola. Annals of the Missouri Botanical Garden 4:43–92.

Gerber, L. R., H. McCallum, K. D. Lafferty, J. L. Sabo, and A. Dobson. 2005. Exposing

extinction risk analysis to pathogens: Is disease just another form of density dependence?

Ecological Applications 15.

Gilbert, G. S. 2002. EVOLUTIONARY ECOLOGY OF PLANT DISEASES IN NATURAL

ECOSYSTEMS. Annual Review of Phytopathology 40:13–43.

Gilbert, G. S., R. B. Foster, and S. P. Hubbell. 1994. Density and distance-to-adult effects of a

canker disease of trees in a moist tropical forest. Oecologia 98:100–108.

Gunton, R. M., and W. E. Kunin. 2009. Density-dependence at multiple scales in experimental

and natural plant populations. Journal of Ecology 97:567–580.

Hajibabaei, M., M. A. Smith, D. H. Janzen, J. J. Rodriguez, J. B. Whitfield, and P. D. N. Hebert.

2006. A minimalist barcode can identify a specimen whose DNA is degraded. Molecular

Ecology Notes 6:959–964.

Hanzawa, F. M., and S. Kalisz. 1993. The Relationship between Age, Size, and Reproduction in

Trilliu grandiflorum (Liliaceae). American Journal of Botany 80:405–410.

Hargreaves, A. L., and C. G. Eckert. 2019. Local adaptation primes cold-edge populations for

range expansion but not warming-induced range shifts. Ecology Letters 22:78–88.

Harrison, S., and A. D. Taylor. 1997. 2 - Empirical Evidence for Metapopulation Dynamics.

Pages 27–42 *in* I. Hanski and M. E. Gilpin, editors. Metapopulation Biology. Academic Press,

San Diego.

Harrison, T. L., A. K. Simonsen, J. R. Stinchcombe, and M. E. Frederickson. 2018. More

partners, more ranges: generalist legumes spread more easily around the globe. Biology Letters

14:20180616.

Hector, T. E., K. L. Hoang, J. Li, and K. C. King. 2022. Symbiosis and host responses to heating.

Trends in Ecology & Evolution 37:611–624.

Hegland, S. J. 2014. Floral neighbourhood effects on pollination success in red clover are scale-

dependent. Functional Ecology 28:561–568.

Hijmans, R., M. Barbosa, A. Ghosh, and A. Mandel. 2025, May 31. geodata: Download

Geographic Data. https://github.com/rspatial/geodata.

Hooper, G. R., F. W. Case, and R. Myers. 1971. Mycoplasma-like bodies associated with a

flower greening disorder of a wild flower, Trillium grandiflorum. The Plant Diesase Reporter

55:1108–1110.

iNaturalist contributors, iNaturalist. 2025. iNaturalist Research-grade Observations.

iNaturalist.org. Occurrence dataset https://doi.org/10.15468/ab3s5x accessed via GBIF.org on

2025-10-07.

Jones, L. C., M. A. Rafter, and G. H. Walter. 2022. Host interaction mechanisms in herbivorous

insects – life cycles, host specialization and speciation. Biological Journal of the Linnean Society

137:1–14.

Kalisz, S., F. M. Hanzawa, S. J. Tonsor, D. A. Thiede, and S. Voigt. 1999. Ant-Mediated Seed

Dispersal Alters Pattern of Relatedness in a Population of Trillium Grandiflorum. Ecology

80:2620–2634.

Kim, T. N., N. Underwood, and B. D. Inouye. 2013. Insect herbivores change the outcome of

plant competition through both inter- and intraspecific processes. Ecology 94:1753–1763.

Knight, T. M. 2003. Effects of herbivory and its timing across populations of *Trillium*

grandiflorum (Liliaceae). American Journal of Botany 90:1207–1214.

Knight, T. M. 2007. Population-level consequences of herbivory timing in Trillium

grandiflorum. American Midland Naturalist 157.

Koinuma, H., K. Maejima, R. Tokuda, Y. Kitazawa, T. Nijo, W. Wei, K. Kumita, A. Miyazaki,

S. Namba, and Y. Yamaji. 2020. Spatiotemporal dynamics and quantitative analysis of

phytoplasmas in insect vectors. Scientific Reports 10:4291.

Krebs, B. L., T. K. Anderson, T. L. Goldberg, G. L. Hamer, U. D. Kitron, C. M. Newman, M. O.

Ruiz, E. D. Walker, and J. D. Brawn. 2014. Host group formation decreases exposure to vector-

borne disease: a field experiment in a 'hotspot' of West Nile virus transmission. Proceedings of

the Royal Society B: Biological Sciences 281:20141586.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests in

Linear Mixed Effects Models. Journal of Statistical Software 82:1–26.

Lamoureux, G., and R. Larose. 2002. Flore printanière: guide d'identification Fleurbec.

Fleurbec, Saint-Henri-de-Lévis [Québec].

Lee, I.-M., R. E. Davis, and D. E. Gundersen-Rindal. 2000. Phytoplasma: Phytopathogenic

Mollicutes1. Annual Review of Microbiology 54:221–255.

Lee-Yaw, J. A., H. M. Kharouba, M. Bontrager, C. Mahony, A. M. Csergő, A. M. E. Noreen, Q.

Li, R. Schuster, and A. L. Angert. 2016. A synthesis of transplant experiments and ecological

niche models suggests that range limits are often niche limits. Ecology Letters 19:710–722.

Lindgren, F., H. Rue, and J. Lindström. 2011. An Explicit Link between Gaussian Fields and

Gaussian Markov Random Fields: The Stochastic Partial Differential Equation Approach.

Journal of the Royal Statistical Society Series B: Statistical Methodology 73:423–498.

Liu, J., D. Gopurenko, M. J. Fletcher, A. C. Johnson, and G. M. Gurr. 2017. Phytoplasmas–The

"Crouching Tiger" Threat of Australian Plant Pathology. Frontiers in Plant Science 8.

Lively, C. M., S. G. Johnson, L. F. Delph, and K. Clay. 1995. Thinning Reduces the Effect of

Rust Infection on Jewelweed (Impatiens Capensis). Ecology 76:1859–1862.

Massicotte, P., and A. South. 2017, March 21. rnaturalearth: World Map Data from Natural

Earth.

Mauricio, R., and M. D. Rausher. 1997. EXPERIMENTAL MANIPULATION OF PUTATIVE

SELECTIVE AGENTS PROVIDES EVIDENCE FOR THE ROLE OF NATURAL ENEMIES

IN THE EVOLUTION OF PLANT DEFENSE. Evolution 51:1435–1444.

Meentemeyer, R. K., M. A. Dorning, J. B. Vogler, D. Schmidt, and M. Garbelotto. 2015. Citizen

science helps predict risk of emerging infectious disease. Frontiers in Ecology and the

Environment 13:189–194.

Meredith, C., A. Frances, A. Highland, L. Oliver, A. Floden, L. L. Gaddy, W. Knapp, D.

Leaman, S. Leopold, T. Littlefield, R. Raguso, E. Schilling, A. Schotz, A. Walker, and K.

Wayman. 2022. The Conservation Status of Trillium in North America. Mt. Cuba Center, and

New Mexico BioPark Society, Hockessin, DE and Albuquerque, NM.

Mooring, M. S., and B. L. Hart. 1992. Animal Grouping for Protection From Parasites: Selfish

Herd and Encounter-Dilution Effects. Behaviour 123:173–193.

Munyaneza, J. E., and D. C. Henne. 2013. Leafhopper and Psyllid Pests of Potato. Pages 65–102

Insect Pests of Potato. Elsevier.

Naimi, B., N. A. S. Hamm, T. A. Groen, A. K. Skidmore, and A. G. Toxopeus. 2014. Where is

positional uncertainty a problem for species distribution modelling? Ecography 37:191–203.

Nathan, P., E. P. Economo, B. Guénard, A. K. Simonsen, and M. E. Frederickson. 2023.

Generalized mutualisms promote range expansion in both plant and ant partners. Proceedings of

the Royal Society B: Biological Sciences 290:20231083.

Ohara, M. 1989. Life History Evolution in the Genus Trillium. Plant Species Biology 4:1–28.

Oldfather, M. F., and D. D. Ackerly. 2019. Microclimate and demography interact to shape

stable population dynamics across the range of an alpine plant. New Phytologist 222:193–205.

Parker, I. M., and G. S. Gilbert. 2018. Density-dependent disease, life-history trade-offs, and the

effect of leaf pathogens on a suite of co-occurring close relatives. Journal of Ecology 106:1829–

1838.

Pautasso, M., G. Aas, V. Queloz, and O. Holdenrieder. 2013. European ash (*Fraxinus excelsior*)

dieback – A conservation biology challenge. Biological Conservation 158:37–49.

Pebesma, E. 2018. Simple Features for R: Standardized Support for Spatial Vector Data. The R

Journal 10:439.

Pebesma, E., and R. Bivand. 2023. Spatial Data Science. Chapman and Hall/CRC, New York.

Petersen, S. M., C. Keith, K. Austin, S. Howard, L. Su, and W. Qiu. 2019. A Natural Reservoir

and Transmission Vector of Grapevine Vein Clearing Virus. Plant Disease 103:571–577.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for

Statistical Computing. Vienna, Austria.

Rao, G. P., E. Alvarez, and A. Yadav. 2018. Phytoplasma Diseases of Industrial Crops. Pages

91–121 Phytoplasmas: Plant Pathogenic Bacteria - I. Springer Singapore, Singapore.

Rathcke, B. 1985. Slugs as Generalist Herbivores: Tests of Three Hypotheses on Plant Choices.

Ecology 66:828–836.

Rehm, E. M., P. Olivas, J. Stroud, and K. J. Feeley. 2015. Losing your edge: climate change and

the conservation value of range-edge populations. Ecology and Evolution 5:4315–4326.

Rizopoulos, D. 2025. GLMMadaptive: Generalized Linear Mixed Models using Adaptive

Gaussian Quadrature. https://github.com/drizopoulos/GLMMadaptive.

Rooney, T. P., and K. Gross. 2003. A demographic study of deer browsing impacts on Trillium

grandiflorum. Plant Ecology 168:267–277.

Rue, H., S. Martino, and N. Chopin. 2009. Approximate Bayesian Inference for Latent Gaussian

models by using Integrated Nested Laplace Approximations. Journal of the Royal Statistical

Society Series B: Statistical Methodology 71:319–392.

Santos, A. A., J. Jacques, and E. Pérez-López. 2024. Potential impact of climate change on

Nearctic leafhopper distribution and richness in North America. npj Sustainable Agriculture

2:12.

Sinha, R. C., and L. N. Chiykowski. 1980. Transmission and morphological features of

mycoplasmalike bodies associated with peach X-disease. Canadian Journal of Plant Pathology

2:119–124.

Smith, E. T. 1879. A Michigan Trillium. Botanical Gazette 4:180–181.

Star, B., and H. G. Spencer. 2013. Effects of Genetic Drift and Gene Flow on the Selective

Maintenance of Genetic Variation. Genetics 194:235–244.

Uricchio, L. H., E. L. Bruns, M. E. Hood, M. Boots, and J. Antonovics. 2023. Multimodal

pathogen transmission as a limiting factor in host distribution. Ecology 104:e3956.

Ustun, N., M. G. Zamharir, and A. M. Al-Sadi. 2023. Chapter 7 - Updates on phytoplasma

diseases management. Pages 97–123 *in* A. K. Tiwari, K. Oshima, A. Yadav, S. A. Esmaeilzadeh-

Hosseini, Y. Hanboonsong, and S. Lakhanpaul, editors. Characterization, Epidemiology, and

Management. Academic Press.

Vellend, M., J. A. Myers, S. Gardescu, and P. L. Marks. 2003. Dispersal of Trillium Seeds by

Deer: Implications for Long-Distance Migration of Forest Herbs. Ecology 84:1067–1072.

Venter, O., E. W. Sanderson, A. Magrach, J. R. Allan, J. Beher, K. R. Jones, H. P. Possingham,

W. F. Laurance, P. Wood, B. M. Fekete, M. A. Levy, and J. E. M. Watson. 2016. Sixteen years

of change in the global terrestrial human footprint and implications for biodiversity conservation.

Nature Communications 7:12558.

Wang, R., B. Bai, D. Li, J. Wang, W. Huang, Y. Wu, and L. Zhao. 2024. Phytoplasma: A plant

pathogen that cannot be ignored in agricultural production—Research progress and outlook.

Molecular Plant Pathology 25:e13437.

Webster, B., and R. T. Cardé. 2017. Use of habitat odour by host-seeking insects. Biological

Reviews 92:1241–1249.

Webster, C. G., B. A. Coutts, R. a. C. Jones, M. G. K. Jones, and S. J. Wylie. 2007. Virus impact

at the interface of an ancient ecosystem and a recent agroecosystem: studies on three legume-

infecting potyviruses in the southwest Australian floristic region. Plant Pathology 56:729–742.

Wickham, H., M. Averick, J. Bryan, W. Chang, L. McGowan, R. François, G. Grolemund, A.

Hayes, L. Henry, J. Hester, M. Kuhn, T. Pedersen, E. Miller, S. Bache, K. Müller, J. Ooms, D.

Robinson, D. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K. Woo, and H. Yutani.

2019. Welcome to the Tidyverse. Journal of Open Source Software 4:1686.

Zanaga, D., R. Van De Kerchove, W. De Keersmaecker, N. Souverijns, C. Brockmann, R. Quast,

J. Wevers, A. Grosu, A. Paccini, S. Vergnaud, O. Cartus, M. Santoro, S. Fritz, I. Georgieva, M.

Lesiv, S. Carter, M. Herold, L. Linlin, N. Tsendbazar, F. Raimoino, and O. Arino. 2021. ESA

WorldCover 10 m 2020 v100. Meteosat Second Generation Evapotranspiration (MET).

Zeileis, A., C. Kleiber, and S. Jackman. 2008. Regression models for count data in R. Journal of

Statistical Software 27.

Zhou, L., C. W. Hoy, S. A. Miller, and L. R. Nault. 2003. Marking Methods and Field

Experiments to Estimate Aster Leafhopper (Macrosteles quadrilineaus) Dispersal Rates.

Environmental Entomology 32:1177–1186.

Zizka, A., D. Silvestro, T. Andermann, J. Azevedo, C. Duarte Ritter, D. Edler, H. Farooq, A.

Herdean, M. Ariza, R. Scharn, S. Svantesson, N. Wengström, V. Zizka, and A. Antonelli. 2019.

CoordinateCleaner: Standardized cleaning of occurrence records from biological collection

databases. Methods in Ecology and Evolution 10:744–751.