Oligo DNA-based quantum dot (QD) single-particle tracking for multicolor single-molecule imaging

Shigeo Sakuragi[1], Naoki Kato[1], Tomoya Uchida[1], Boxiao Zhao[1], Taro Katagiri[1], Miyu Enomoto[1], Rie Kato[1], Hideaki Yoshimura[2], Chisato Oyama[1], Iona Katayama[1], Ayano Chikuma[1], Yuji Teramura[3, 4], and Hiroko Bannai[1]

[1] *Department of Electrical Engineering and Biosciences, School of Advanced Science and Engineering, Waseda University, Shinjuku-Ku, Tokyo 162-0056, Japan*

[2] *Department of Chemistry, School of Science, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan*

[3] *Department of Bioengineering, Graduate School of Engineering, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-8656, Japan*

[4] ***Current address***: *Cellular and Molecular Biotechnology Research Institute (CMB), National Institute of Advanced Industrial Science and Technology (AIST), AIST Tsukuba Central 5, 1-1-1 Higashi, Tsukuba, Ibaraki 305-8565, Japan*

Quantum dot single-particle tracking (QD-SPT), a powerful tool for analyzing membrane domains that are critical to various cellular processes, is widely used in membrane molecular dynamics research. QDs, which possess both a broad excitation range and a narrow emission bandwidth, are inherently suited for multicolor imaging at various wavelengths. However, applying multicolor QD-SPT with multiple biomolecular targets has been challenging due to the limited methods for specifically conjugating QDs to biomolecules. Here, we propose a DNA hybridization-based QD labeling method that generates several specific interactions based on sequence. QD fused with 20-mer oligo DNA specifically labeled membrane lipid 1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine (DPPE) covalently bound to complementary oligo DNA, forming a stable label that is suitable for SPT. The combination of polyA–polyT sequence as the linker oligo caused more QDs to fuse to DPPE compared with a random sequence linker. Oligo DNA-based QD-SPT accurately reflected the diffusion dynamics of DPPE measured using the single-fluorescence tracking technique and was compatible with conventional QD-SPT utilizing secondary antibody Fab. Using different pairs of oligo DNA sequences, we successfully achieved multicolor QD-SPT that distinguishes the lateral diffusion of DPPE and a membrane protein, GABA$_A$ receptor (GABA$_A$R), within the same cell. The oligo DNA-based QD labeling method developed in this study is anticipated to substantially advance simultaneous multicolor QD-SPT of different living cell membrane molecules specific to DNA sequences.

Keywords: quantum dot, single-particle tracking, multicolor imaging, membrane molecule, diffusion

◀ *Significance* ▶

Quantum dot single-particle tracking (QD-SPT) is a powerful approach for investigating membrane dynamics; however, the limited conjugation methods available hinder multicolor labeling. This study introduces a DNA hybridization-based labeling strategy, where QDs conjugated with 20-mer oligos that specifically bind to complementary oligo-tagged lipids or antibodies. PolyA–polyT linkers improved labeling efficiency compared with random sequences. This method accurately captured 1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine (DPPE) diffusion, matched conventional QD-SPT, and enabled simultaneous multicolor tracking of DPPE and GABA$_A$ receptors in the same cell. This oligo-DNA-based approach enables multicolor QD-SPT of distinct membrane proteins.

Corresponding author: Hiroko Bannai, Waseda University, Shinjuku, Tokyo 162-0056, Japan. ORCID iD: https://orcid.org/0000-0002-0951-488X, e-mail: hiroko.bannai@waseda.jp

In the fluid-mosaic model proposed by Singer and Nicolson, biological membranes demonstrate fluid-like
properties, in which membrane proteins and lipids that are embedded in the lipid bilayer undergo lateral diffusion
across the plasma membrane [1]. However, high-resolution analyses at the sub-millisecond timescale have
revealed that even lipid molecules that diffuse freely are transiently confined within compartments of
approximately 13–100 nm in size, based on the actin cytoskeleton and transmembrane proteins anchored to it.
Lipid molecules display compartmentalized diffusion, characterized by restricted motion within individual
compartments and occasional hopping between adjacent compartments [2] [3]. Furthermore, specific membrane
proteins are localized in the membrane region on the nanometer to micrometer scale, generating functional
membrane domains [4]. These domains are developed through fine regulation and restriction of lateral diffusion
of membrane molecules and play crucial roles in a wide array of cellular processes, including signal transduction,
synaptic transmission, and immune responses. For example, neurotransmitter receptors accumulate at the synapse,
which serve as the fundamental units of inter-neuronal communication, and the number of receptors at the
synapse, ultimately, the efficacy of synaptic transmission, is regulated via lateral diffusion constraints caused by
interactions between receptors and scaffold proteins [5]. Further, transmembrane proteins and cytoskeletons
restrict the lateral diffusion of membrane lipids in the axon initial segment of neurons, underlying the physical
compartmentalization between neuronal input and output domains [6]. Observations of membrane molecule
dynamics are crucial to understanding such self-organizing principles of plasma membranes and revealing the
molecular mechanisms underlying their physiological functions.
Single-molecule imaging enables the observation of the dynamics of individual molecules, facilitating not only
the assessment of ensemble behavior but also detailed information about the behavior of molecules in confined
microdomains, stochastic movements, and heterogeneity within molecular populations. Single-particle tracking
(SPT) is one of the representative methods in single-molecule imaging, in which probes are specifically
conjugated to target molecules in live cells, causing the high-precision tracking of particle positions [7]. Among
the SPT methods that use diverse probes, quantum dot (QD)-SPT has become a widely used approach for
investigating the dynamics of membrane molecules in neurons. QDs are semiconductor nanocrystals that serve as
fluorescent probes, which overcome limitations associated with organic fluorophores caused by their superior
brightness and long fluorescence lifetimes [5]. Furthermore, their high brightness and significant Stokes shift
provide an excellent signal-to-noise ratio (SNR), enabling more straightforward and higher-resolution analysis of
fluorescence images compared with organic dyes [8]. However, QDs still have certain limitations, as the relatively
large size of QD bioconjugates hinders access to spatially restricted regions, such as synaptic clefts, due to
physical confinement [9] [10]. The blinking of QDs, i.e., intermittent fluorescence emission, hinders continuous
tracking and quantitative analysis [11]. However, the optical advantages of QDs, particularly their high SNR,
which helps improve spatial resolution, outweigh these drawbacks. Hence, QD-SPT has been extensively adopted
in neuroscience research to investigate membrane molecule dynamics at the single-molecule level.
The most distinctive feature of QDs is that their fluorescence emission wavelength is precisely identified based
on their particle size, leading to sharp and narrow emission spectra [12]. With their broad excitation range and
narrow emission bandwidth, QDs are inherently well-suited for multicolor imaging of various wavelengths.
Indeed, studies that used fixed tissues and cells reported multicolor imaging employing QDs [13] [14] [15].
Despite developing QD-SPT over 20 years ago, reports that utilize QDs for multicolor imaging, particularly those
involving labeling of multiple molecular species with QDs of distinct emission wavelengths, are limited. The lack
of robust methods for specifically labeling different target molecules in living cells with QDs with different
wavelengths of emission is one of the primary reasons. Currently, the commonly used QD labeling method
employs the specific interaction between the primary antibodies and secondary antibodies or biotin–avidin
interactions. However, these methods involve a small number of combinations of specific molecular interactions,
which limits the options for multicolor imaging that simultaneously targets multiple different molecules.
Therefore, novel QD-labeling strategies need to be developed to expand the applicability of multicolor QD-SPT in
the same cell.
In recent years, multicolor immunofluorescence techniques employing deoxyribonucleic acid (DNA) barcodes
have facilitated the visualization of multiple protein localizations within the same cell [16] [17]. In this study, we
apply the specific and stable hybridization between complementary DNA strands to achieve persistent conjugation
between membrane molecules and QDs. Unlike DNA-PAINT approaches that depend on rapid binding–unbinding
kinetics [18], the present strategy is designed to provide stable probe attachment over the time scale of single-
particle tracking. This enables sequence-specific targeting of distinct membrane molecules with QDs of defined
emission properties. Utilizing this approach, we demonstrate a practical method for multicolor QD-SPT in the

same cell, wherein multiple molecular species, the phospholipid and the neurotransmitter receptor, are labeled
with QDs through 20-mer DNA hybridization, each emitting at different wavelengths.
58
Materials and methods
Primary culture of rat hippocampal neurons
In this study, all experiments were conducted following the guidelines approved by the Animal Experimentation
Committee of Waseda University. Primary hippocampal neurons were prepared as previously described [19]. In
brief, hippocampi were dissected from embryonic day 20 (E20) Wistar/ST rats (Japan SLC) and treated with 0.125%
trypsin (Sigma-Aldrich) and 0.025 w/v% DNase (Sigma-Aldrich). The tissue was then dissociated in the plating
medium, composed of the minimum essential medium (MEM; Thermo Fisher Scientific) supplemented with 2%
MACS NeuroBrew-21 (Miltenyi Biotec), 2 mM L-glutamine (Nacalai Tesque), 1 mM sodium pyruvate (Nacalai
Tesque), and 5 units/mL of penicillin and 5 µg/mL of streptomycin (Nacalai Tesque). Dissociated cells were seeded
at a density of 1.4×10^5 cells per well on an 18-mm polyethyleneimine-coated glass coverslip and cultured in a CO_2
incubator (37°C, 5% CO_2). Four days after plating, the culture medium was replaced with the maintenance medium,
the Neurobasal-A (Thermo Fisher Scientific), supplemented with 2% MACS NeuroBrew-21, 2 mM L-glutamine,
and penicillin (5 units/mL)/streptomycin (5 µg/mL). Cultures were maintained at 37°C in a CO_2 incubator until the
experiments were performed at 14 days *in vitro* (DIV14) or later.
73
Membrane molecule labeling using ssDNA
We prepared 20-base-pair ssDNA probes comprising one pair
of complementary poly sequences (polyA and polyT) and two
pairs of complementary random sequences (random1S and
random1AS, random2S and random2AS). Supplementary Table
S1 shows the sequence. For the QD-SPT of phospholipids with
ssDNA-conjugated QDs, we prepared the 1,2-dipalmitoyl-sn-
glycero-3-phosphatidylethanolamine (DPPE) tagged with
ssDNA at the hydrophilic PEG end (ssDNA–DPPE) (Figure 1).
As previously described, ssDNA was attached to maleimide-
poly(ethylene glycol)-conjugated phospholipid DPPE through a
thiol–maleimide reaction (Supplementary Figure S1A) [20]. In
brief, DPPE and α-N-hydroxysuccinimidyl-ω-maleimidyl poly
(ethylene glycol) (NHS-PEG-Mal, Mw: 5000) were combined to
produce Mal-PEG-DPPE. Thiolated-labeled ssDNAs to the 5'
end were then reacted with Mal-PEG-DPPE to obtain the
ssDNA–DPPE and maintained in the phosphate-buffered saline
(PBS). Further, DPPE was conjugated with ssDNA via copper-
free click chemistry through dibenzocyclooctyl (DBCO)–azide
conjugation (ssDNA–DBCO–DPPE, Supplementary Figure
S1B) [21] [22]. DPPE–PEG (2000)–azide (10 µM; Avanti) was
reacted with 5' DBCO-C6-N$_2$-PolyA (20) (1 µM; Nihon Gene
Research Laboratories) in PBS for 60 min at room temperature.
Moreover, ssDNA and QD conjugates (ssDNA–QD) were
prepared by mixing Qdot 655 streptavidin conjugate (1 µM;
QD655; Thermo Fisher Scientific), ssDNA labeled with biotin at
the 5' end (10 µM), and Qdot Binding Buffer (QBB; 2.62 mM

Figure 1. Overview of the ssDNA labeling protocol used in this experiment
DPPE conjugated with a 20-mer ssDNA (ssDNA–DPPE), or an anti-GABA$_A$R antibody conjugated with DNA (ssDNA–Ab), and QDs modified with ssDNA via streptavidin–biotin interaction (ssDNA–QD) were prepared. Primary rat hippocampal neurons were incubated with ssDNA–DPPE or ssDNA–Ab for 5 min, followed by a 1-min incubation with ssDNA–QD, in a humid chamber maintained at 37°C.

sodium tetraborate decahydrate [Wako], 39.5 mM boric acid [Nacalai Tesque], 2 w/v% bovine serum albumin
[Sigma-Aldrich], 0.05 w/v% sodium azide) at a volume ratio of 1.0:0.5:8.5, followed by overnight incubation at
4°C. QD605 (Thermo Fisher Scientific) conjugated with ssDNA as described above was used for simultaneous
labeling of multiple membrane proteins in parallel with ssDNA–QD655. The imaging medium, MEM without
phenol red (Thermo Fisher Scientific), supplemented with 20 mM HEPES (Sigma Aldrich), 2 mM glutamine, 1
mM sodium pyruvate, and 33 mM glucose (Nacalai Tesque), was used for the dilution and the incubation of lipid
and antibody, cell washing, and live-cell imaging. QBB was supplemented with 215 mM of sucrose (QBB-suc) to
adjust the osmolarity when QDs were incubated with neurons.

To incorporate ssDNA–DPPE into the plasma membrane, ssDNA–DPPE (5 µg/mL) was incubated with primary
cultured rat hippocampal neurons for 5 min. After washing, the coverslips were incubated with 0.063-0.25 nM
ssDNA–QD in QBB-suc for 1 min, followed by washing and imaging in the imaging medium. Labeling processes,
which include lipid, antibody, and QD incubation, were performed in a humid chamber placed on a heating plate
maintained at 37°C [23].
For labeling $GABA_A$ receptors ($GABA_A$Rs) with ssDNA, we utilized a custom-made antibody that targets the
extracellular domain of the γ2 subunit of $GABA_A$R [24] conjugated with polyA (ssDNA–Ab; Figure 1). The
antibody was conjugated to ssDNA at a 1:5 molar ratio employing the Oligonucleotide Conjugation Kit (Abcam,
ab218260), which targets lysine residues via an amine-reactive ester. Endogenous $GABA_A$R in cultured rat
hippocampal neurons were labeled with ssDNA by incubating with 0.01 mg/mL of ssDNA–$GABA_A$R antibody for
5 min, followed by incubation with 1 nM ssDNA–QD in QBB-suc for 1 min in a humid chamber.
120
Conventional QD labeling using Fab fragment antibody
$GABA_A$R labeling using Fab fragment antibody as a linker for QD (conventional labeling) was performed
according to previous reports [25] [23]. Rat primary hippocampal neurons were incubated with a custom-made
anti-$GABA_A$R γ2 subunit antibody that targets the extracellular domain (0.01 mg/ml, 5 min) followed by
incubation with biotinylated Fab fragment antibody (10 µg/ml, 5 min; Biotin-Fab, Jackson ImmunoResearch).
After washing, $GABA_A$Rs were labeled via biotin–avidin binding by incubation with 1 nM QD655 in QBB-suc at
37°C for 1 min.
128
QD-SPT data acquisition
QD-SPT was conducted using an inverted microscope (IX73, Olympus) equipped with multi-wavelength LED
illumination (niji, Bluebox Optics), an oil-immersion objective lens (×60, NA 1.42, Olympus), and filter sets for
QD wavelength, i.e., an excitation filter (450/70 nm, e.g., Semrock, Rochester, NY; BrightLine® single-band band-
pass filter #FF01-450/70-25), dichroic beam splitters (DM) and emission filter (EM) based on each QD emission
wavelength (for Qdot655: 655/15 nm #FF01-655/15-25, DM #FF640-FDi01-25 × 36; for Qdot605: 605/15 nm
#FF01-605/15-25, DM #FF593-Di03), and the EM-CCD camera (ImagEM C9100-13, Hamamatsu Photonics).
MetaVue software (Molecular Devices) was used to control the excitation illumination and the EM-CCD camera.
A total of 100 consecutive frames were acquired within a 41.8 × 41.8 µm field of view at 13 Hz with a 75 ms
exposure time. During observation, the specimen and the microscope stage were maintained at 37°C using a heating
system (Tokai Hit). In the DNase experiment, 0.05 w/v% DNase (Sigma-Aldrich) in Hank's balanced salt solution
(Nacalai Tesque) was added to the observation chamber on the microscope stage and left for 3 min. Images were
then acquired in the same field of view after DNase administration.
142
Single-fluorescence tracking for DPPE
The ATTO647–DPPE (ATTO-TEC GmbH, AD647-151) was inserted in the cultured neurons by incubating the
coverslips with 0.1 µM ATTO647–DPPE diluted in the imaging medium for 10 min, within the humid chamber
placed in a CO_2 incubator. After washing three times, the coverslip was mounted in the observation chamber and
immersed in the imaging medium for observation.
Total internal reflection fluorescence microscopy (TIRFM) was used for single-molecule fluorescence imaging
of organic dye ATTO647 [26], based on an inverted microscope (IX81, Olympus) equipped with 640-nm diode
laser (CUBE, Coherent Inc.), 405/488/561/635-nm band excitation filter (Semrock Inc.), Dichroic mirror (FF662,
Semrock Inc.), 697/58-nm band-pass filter (Semrock Inc.), EM-CCD camera (ImagEM, Hamamatsu Photonics),
and an oil-immersion objective lens (×100, NA 1.49, Olympus). The imaging was performed in 256 × 256-pixel
subarray mode, recording 200 frames at 30 Hz (33 ms/frame), at the surface and the bottom focal planes in the
same field of view.
155
Tracking analysis of QD and organic dyes
A tracking analysis of QD-SPT was conducted using TI Workbench as previously described [23] [25]. The center
of each QD spot was defined based on cross-correlation of the fluorescent spots with a Gaussian model with a point
spread function. QD-trajectories are automatically assembled by linking the centers of fluorescent spots. Only QDs
with blinking were considered for the analysis to ensure that they are single particles, with the trajectories with >
30 points used for further analysis. The mean squared displacement (MSD) was calculated from the coordinates of

reconstructed trajectories. The MSD-Δt curve, which plots MSD versus time (Δt), was obtained using the following
equation (1) [26]:
164

165      $MSD(n\Delta t) = \frac{1}{N-n}\sum_{i=1}^{N-n}[(x((i+n)\Delta t) - x(i\Delta t))^2 + (y((i+n)\Delta t) - x(i\Delta t))^2]$ (1)
166

167   where Δt denotes the acquisition time (75 ms), N indicates the total number of frames, n represents the frame
168   number, and i denotes the positive integers. The diffusion coefficient (D) of QDs was obtained by fitting the first
169   1–4 points without origin of the MSD-$n\Delta t$ curves with the following equation (2):
170

171      $MSD(n\Delta t) = 4Dn\Delta t$ (2)
172

QDs with D of < 0.0004, equivalent to D of QDs adhered on the coverslips, were defined as "immobile" QDs
[23].
Single-fluorophore tracking (SFT) images were filtered using the dilate (circle, 4 pixels) function in MetaMorph
(Molecular Devices) to reduce spike-like noise, and the trajectories were composed employing TI Workbench
software. To ensure single-molecule fluorescence, the analysis included only bright spots that disappeared in a
single step during recording. The D values were calculated for trajectories tracked for more than 10 consecutive
frames.
180

Ca^{2+} imaging
Primary hippocampal neurons (DIV 14 or later), either with or without QD labeling for GABA$_A$R and DPPE,
were loaded with 0.5 μM Fluo-8 AM (AAT Bioquest) diluted with the imaging medium (MEM supplemented
with 20 mM HEPES [pH 7.4], 2 mM glutamine, and 1 mM sodium pyruvate) for 5 min at 37°C. After washing,
neurons were pre-incubated for 5 min in the imaging chamber with either muscimol (Sigma, 20 μM; GABA$_A$R
agonist) or imaging medium (mock control). The same conditions were maintained throughout the imaging
session. Time-lapse imaging was performed at 1 Hz for 180 s using the same imaging configuration as for QD-
SPT, with appropriate filter sets. Depolarization was induced by applying KCl (40 mM final concentration) 60 s
after the start of recording, and imaging continued for an additional 120 s.
MetaMorph software (Molecular Devices) or ImageJ (NIH) was used for imaging analysis. Regions of
interest (ROIs) were placed over 2–4 neuronal somata per experiment. Changes in intracellular Ca^{2+} levels were
expressed as $\Delta F/F_0$, calculated as $(F - F_0)/F_0$, where F_0 denotes the mean baseline fluorescence intensity before
KCl application. The maximum $\Delta F/F_0$ after KCl stimulation was calculated for each ROI and compared between
the muscimol-treated and mock-stimulated groups.
195

Statistics
Complementary binding by the four ssDNA–QDs was compared using a one-way analysis of variance with a
post hoc Tukey HSD test. The number of QDs before and after DNase application was compared using a Wilcoxon
signed-rank test. A Kolmogorov–Smirnov test was used to perform the cumulative distribution of the D. Differences
in D among three groups were analyzed using the Kruskal–Wallis test, followed by Dunn's post hoc test. A Mann–
Whitney U-test was used for other statistical analyses. The figure illustrates the P-values. Sample size (N) for each
experiment is described in the figure legend. Each experiment utilized at least three independent culture batches.
203

Result
QD Labeling of Membrane Molecules depending on annealing between complementary ssDNAs
First, we investigated whether the phospholipid DPPE tagged with ssDNA (ssDNA–DPPEs) was specifically
labeled with QDs conjugated with ssDNA (ssDNA–QDs) with a complementary sequence through the
hybridization. The 20-mer oligo DNA, composed of identical bases or random sequences, was developed as
ssDNA sequences (Supplementary Table S1) and added to QDs and target molecules. DPPEs conjugated with
polyA ssDNA (polyA–DPPE) via the thiol–maleimide reaction (Supplementary Figure S1A) were inserted into
the plasma membrane of primary rat hippocampal neurons and incubated with QDs conjugated with either polyA
or polyT sequences (Figure 2A). QDs with polyA (polyA–QDs), carrying the same sequence as the ssDNA on the
DPPE, did not remain on the surface of the neurons, whereas QDs with polyT (polyT–QDs), which carry the
complementary sequence, were observed on the neurons (Figure 2B). The median number of QDs bound to
neurons per image was 1 for polyA–QDs, whereas that for polyT–QDs was significantly higher at 241 per image

(Figure 2C, Supplementary Table S2). Comparable to the result of polyA–DPPEs, the DPPE tagged with polyA
by the copper-free click chemistry through the DBCO–azide reaction (polyA-DBCO-DPPE, Figure S1B) was
targeted by polyT–QDs, but not by polyA–QDs (Supplementary Figures S1C, S1D), indicating that the linker
between the DPPE and the ssDNA does not affect the DNA hybridization.
PolyA–QDs labeled the plasma membrane of neurons incubated with polyT–DPPE, whereas ssDNA–QD did
not bind to neurons without polyT–DPPE (Supplementary Figures S2A, S2B). To evaluate nonspecific binding,
neurons without polyT–DPPE were incubated with 0.063 nM polyA–QD or a 10-fold higher concentration (0.63
nM) (Supplementary Figure S2C). Fewer than one QD per field of view (41.8 × 41.8 μm) was detected at 0.063
nM, whereas the 10-fold higher concentration yielded 2–7 QDs per cell (4.4 ± 0.3, mean ± SEM) (Supplementary
Figure S2D). Considering that more than 20 QDs (median) per field of view were typically detected under
specific binding conditions (Supplementary Table S2), the number of non-specifically bound QDs remained
substantially lower than the specific signal even at higher ssDNA–QD concentration. Furthermore, polyA–DPPE
was rarely targeted with ssDNA–QD that carries two random sequences, random 1AS and random 2AS
(Supplementary Figures S2E, S2F), indicating the high dependence of QD labeling via ssDNA on the DNA
sequence complementarity.
231

Figure 2. QD labeling of membrane lipids via complementary DNA sequences
(A) Schematic illustration of QD labeling of DPPE via complementary DNA hybridization. PolyA–DPPE was incorporated into the neuronal plasma membrane and incubated with either polyA–QDs or polyT–QDs. (B) Representative fluorescence images after labeling with polyA–QDs (left) or polyT–QDs (right). Scale bar indicates 10 μm. (C) Number of bound QDs per image. Solid lines indicate medians (N = 10 images per group). (D) Specificity test using random ssDNA sequences. Random1S–DPPE was probed with polyA–QD, polyT–QD, random1AS–QD (complementary), or random2AS–QD. (E) Representative images of the four ssDNA conditions. Scale bar indicates 10 μm. (F) Number of QDs per field of view. Solid lines indicate medians (N = 10 images per group). (G) DNase treatment time course. Images were acquired before and 3 min after adding 0.05% DNase or HBSS (mock). (H) Representative images of the same field before and after treatment. Scale bar indicates 10 μm. (I) Number of QDs per field of view (N = 11 images per group). Statistics: (C) Mann–Whitney U test; (F) one-way ANOVA with Tukey's HSD; (I) Wilcoxon signed-rank test.

To further confirm the binding specificity of QD labeling with ssDNA, we inserted a random ssDNA
sequence-conjugated DPPE (random1S–DPPE) into the plasma membrane of cultured neurons and evaluated
whether QDs with four different ssDNA sequences (polyA, polyT, random1AS, and random2AS) could bind to
random1S–DPPE (Figure 2D). QDs with polyA, polyT, or random2AS, all of which are not complementary to
random1S, did not remain on the cell surface. In contrast, a substantial amount of QDs with the complementary
sequence random1AS bound to random1s–DPPE (Figure 2E, Supplementary Table S2). Statistical analysis
revealed that random1AS–QDs were significantly more observed in the plasma membrane (Figure 2F,
Supplementary Table S2). These results indicate that hybridization between 20-base ssDNAs with complementary
sequences exhibits sufficient binding specificity for QD labeling, even when using random sequences. Further, the
binding between ssDNA–DPPEs and ssDNA–QDs was sensitive to DNase treatment (Figure 2G). Specifically,
0.05% DNase treatment of neurons labeled with random1S–DPPE and random1AS–QDs reduced the number of

QDs in the same field, whereas the mock group remained unchanged before and after the addition of DNase
(Figures 2H, 2I, Supplementary Table S2). This further emphasizes that the QD labeling was not nonspecific but
instead oligo DNA-dependent. Collectively, the hybridization of ssDNAs with complementary sequences applies
to the specific labeling of membrane molecules with QDs.
248
Measurement of the diffusion coefficient (D) of DPPE by oligo DNA-dependent QD-SPT
The suitability of DPPE labeled with QDs was investigated through DNA hybridization for SPT. The diffusion
of membrane molecules at the lower (coverslip-facing) membrane in some cell types is reduced compared with
the upper (nonadherent) membrane [27] [28]. TIRFM was utilized to measure the diffusion coefficient (D) of the
DPPE fused with the organic dye ATTO647 (ATTO647–DPPE) in the primary neurons at the upper and lower
membranes of the same cell [26] (Figure 3A). ATTO647–DPPE trajectories at the upper and lower membranes
demonstrated different diffusion properties (Figure 3B). The D of ATTO647–DPPE at the lower membrane
(median: 1.15×10^{-1} µm²/s) and upper membrane (median: 2.21×10^{-1} µm²/s) were significantly larger than that
of glass-attached ATTO647–DPPE (median: 2.40×10^{-2} µm²/s, Figure 3C and Supplementary Table S2).
Furthermore, the D at the lower membrane was significantly lower than that at the upper membrane (Figures 3C
and 3D, Supplementary Table S2), indicating the reduced lateral mobility of DPPE at the lower membrane of the
primary cultured neuron.
261

262
263

Figure 3. Diffusion dynamics of the DPPE measured using single-fluorophore imaging and oligo DNA-based QD-SPT
(A) ATTO647–DPPE labeling scheme and representative upper-membrane image. Scale bar, 5 µm. (B) Representative trajectories of ATTO647–DPPE at the upper and lower membranes. Scale bar, 1 µm. (C) D of ATTO647–DPPE at the glass-attached (gray), lower (blue), and upper (red) membranes. (D) Cumulative distributions of D at the upper and lower membranes. (E) OligoDNA-based QD labeling scheme and representative upper-membrane image. Scal bar indicates 5 µm. (F) Representative QD trajectories at the upper and lower membranes. Scale bars indicate 1 µm (main) and 0.2 µm (inset). (G) Fraction of immobile QDs per field of view. (H, I) Dot plot (H) and cumulative distribution (I) of D of mobile QDs. Statistical analysis: (C) Kruskal–Wallis test with Dunn's post hoc test; (G, H) Mann–Whitney U test (Solid lines indicate medians); (D, I) Kolmogorov–Smirnov test (*Dks,* KS distance). N is reported in parentheses as spots (C, D), images (G), and QDs (H, I) (see Supplementary Table S2).

264
265　The diffusion of polyT–DPPE labeled with polyA–QDs was then measured at the upper and lower membranes
266 of cultured neurons (Figure 3E). QD–polyA/polyT–DPPE complex trajectories exhibited lateral diffusion on the

plasma membrane (Figure 3F), indicating the sufficient stability of QD labeling via oligo DNA for tracking membrane diffusion dynamics. The percentage of immobile QDs, QDs with D of less than 4.00×10^{-4} µm²/s [23], was significantly larger at the lower membrane (median: 17.3%) than at the upper membrane (median: 2.16%) (Figure 3G, Supplementary Table S2). Furthermore, the D of mobile QD–ssDNA–DPPE remained smaller at the lower membrane (median: 0.82×10^{-1} µm²/s) than at the upper membrane (median: 1.86×10^{-1} µm²/s), with a greater proportion of low D at the lower membrane (Figures 3H and 3I, Supplementary Table S2). These results indicate a suppressed diffusion of the QD–oligo DNA–DPPE complex at the lower membrane of cultured neurons, in coordination with the finding of ATTO647-conjugated DPPE whose molecular size is smaller than that of QDs (Figures 3C and 3D). However, a more immobile QD fraction and a smaller D at the lower membrane indicate the existence of steric hindrance due to the QD's relatively large size compared with organic dyes. These results indicate that molecules in the upper membrane layer, where the effect of physical diffusion suppression is minimal, should be considered in subsequent experiments measuring D.

Impacts of ssDNA Sequence on QD labeling efficacy and diffusion coefficient (D)

We examined whether ssDNA sequence affects the QD binding affinity and D with the ssDNA sequence pair of polyA and polyT (melting temperature [Tm] 52.17°C) and random1S/1AS (Tm: 64.47°C) (Figure 4A). The same ssDNA–DPPE (5 µg/mL) and ssDNA–QD (0.25 nM) concentration resulted in a higher number of QDs per image in the polyA/polyT ssDNA pair group than the random1S/1AS pair (Figure 4B, Supplementary Table S2). Despite having the lower Tm, polyA/polyT combination detected ~10 fold more QDs (median:193 QDs/image), compared with random1S/1AS (median: 20 QDs/image) (Figure 4C, Supplementary Table S2). This indicates that the poly ssDNA sequence has higher labeling efficacy compared with the random sequence under current experimental conditions. QD-SPT should be performed at a sufficiently low density so that each QD can be identified as a single particle optically without overlapping the trajectory [9]. Therefore, ssDNA–DPPE and ssDNA–QD concentrations were adjusted based on the oligo DNA pair sequence in experiments conducted thereafter; thus, fewer than ~50 QDs are found in one field of view (polyA/polyT pair: 1 µg/mL ssDNA–DPPE and 0.063 nM ssDNA–QD; random sequence pair: 20 µg/mL ssDNA–DPPE and 0.25 nM ssDNA–QD).

Figure 4. Impact of oligo DNA sequence on the diffusion dynamics of QD–oligo DNA–DPPE
(A) QD labeling via complementary sequences of polyA/polyT or random ssDNA. (B) Fluorescence images of neuron surface labeled with 0.25 nM polyT–QD and 5 µg/mL of polyA–DPPE (top) or 0.25 nM random1AS–QD and 5 µg/mL of random1S–DPPE (bottom). Scale bar indicates 10 µm. (C) Number of QDs per field. The line indicates the median. N = 5 images for both groups. (D) Example of trajectories of QD labeled via polyA/polyT (top) or random ssDNA (bottom). Scale bar indicates 1 µm. (E, F) The dot plot (E) and cumulative probability distribution (F) of D for QDs labeled with polyA/polyT or random sequence oligo DNA. Solid lines on the plot indicate medians; N = 220 (poly) and 257 (random) QDs. Mann–Whitney U-test for (C and E) and Kolmogorov–Smirnov test for (F).

Under our experimental conditions, each QD is expected to carry more than one ssDNA. To investigate whether such structural multivalency might promote clustering of ssDNA-DPPE on the plasma membrane, we conducted a quantitative binding analysis for the random1S/1AS pair and an independent random sequence (random2S/2AS) (Supplementary Figure S3). In this assay, the QD–ssDNA concentration was held constant at 0.25 nM, whereas the amount of ssDNA-DPPE applied to cells varied over a wide range (0–100 µg/mL). Binding increased with ligand concentration and reached a clear plateau (Bmax = 208.6 for random1S/1AS; 224.8 for random2S/2AS, Supplementary Figures S3B and S3D), consistent with a finite number of accessible membrane

binding sites. Under the experimental conditions utilized throughout this study, the observed number of bound
QDs per cell (~50) was well below this saturation level (<25% of Bmax). Further, the working concentration of
ssDNA-DPPE (20 μg/mL) remained below the half-maximal binding concentration ($C_{1/2}$ = 24.73 μg/mL for
random1S/1AS; 27.98 μg/mL for random2S/2AS, Supplementary Figures S3B and S3D), indicating that
membrane occupancy was maintained in a sub-saturating regime. Under these sparse labeling conditions, the
likelihood that a single QD bridges multiple spatially separated ssDNA-DPPE molecules is expected to be low.
We cannot formally exclude rare multivalent interactions at the single-particle level; however, the quantitative
binding behavior argues against substantial QD-induced aggregation under our imaging conditions.
To investigate the association of ssDNA sequences with the diffusion dynamics of target molecules, we
compared the diffusion dynamics of DPPE labeled with QDs conjugated to either polyA/polyT pair or
random1S/1AS pair. The trajectories of QD–oligo DNA–DPPE complex of both polyA/polyT pair and
random1S–AS pair exhibited similar diffusion behavior on the plasma membrane (Figure 4D). No significant
difference in the median and distribution of D was found between the polyA/polyT pair and the random1S/1AS
pair (Figures 4E, 4F, Supplementary Table S2). These results indicate that the sequence of linker ssDNA affects
labeling efficacy but has little effect on the D of oligo DNA-based QD-SPT.
317

DNA-based QD-SPT for membrane proteins comparable to conventional QD-SPT

We further applied oligoDNA-dependent QD labeling to the diffusion analysis of a membrane protein. Here,
for the proof of concept, we selected an inhibitory neurotransmitter receptor GABA$_A$R, whose diffusion behavior
on the neuron has been analyzed using conventional QD-SPT [25] [29] [30] [31]. PolyA ssDNA was conjugated
to the anti-GABA$_A$R antibody (polyA-Ab) and targeted to the endogenous GABA$_A$Rs in the neuronal plasma
membrane, followed by the incubation with polyA–QD or polyT–QD (Figure 5A). Only polyT–QD, which are
complementary to the polyA sequence, bound to polyA–GABA$_A$R (Figure 5B). The median number of QDs per
image was 2 for polyA–QD and 28 for polyT–QD (Figure 5C, Supplementary Table S2). These results indicate
that QDs can be specifically targeted to the protein of interest via oligo DNA hybridization.
To investigate whether QD labeling affects GABA$_A$R function at the cellular level, we conducted Ca^{2+}
imaging experiments under muscimol stimulation. The application of the GABA$_A$R agonist muscimol in
unlabeled neurons significantly suppressed the increase in intracellular Ca^{2+} induced by 40 mM KCl application
that induced the activation of voltage-dependent Ca^{2+} channels, consistent with GABA$_A$R-mediated
hyperpolarization (Supplementary Figure S4A). Neurons labeled with QD-ssDNA-GABA$_A$Rs (Supplementary
Figure S4B), as well as cells with QD-ssDNA-DPPE (Supplementary Figure S4C), demonstrated a similar
suppression of KCl-induced Ca^{2+} elevation in the presence of muscimol treatment (Supplementary Figure S4D).
These results indicate that GABA$_A$R-mediated inhibitory signaling is preserved at the cellular level after QD
labeling.
We then compared oligo DNA-based QD-SPT with conventional QD-SPT, which targets QDs via secondary
antibody Fab. PolyA-Abs on the GABA$_A$R were labeled with polyT–QDs (oligo DNA) or with a conventional
method using biotinylated Fab antibodies and streptavidin–QDs (Fab) (Figure 5D). In both methods, QD-labeled
GABA$_A$Rs demonstrated diffusive behavior on the neuronal membrane (Figure 5E), with no statistically
significant difference in the median D (Figure 5F, oligo DNA: 6.49 × 10^{-2} μm²/s; Fab: 5.19 × 10^{-2} μm²/s).
However, analysis of the distribution of D revealed a tendency for D-values to be high in the slow fraction of
QDs–oligo DNA–GABA$_A$R (Figure 5G; Kolmogorov–Smirnov test, p = 0.012, D_{KS} = 0.080). Collectively, oligo
DNA-based labeling functions as an effective alternative method to secondary antibody Fab labeling in QD-SPT
of membrane proteins. However, differences in the linkers between QD and protein may affect the distribution of
D.
346

Figure 5. Analysis of membrane proteins with oligo DNA-based QD-SPT and conventional QD-SPT

(A) Labeling of QD with antibodies against $GABA_AR$ γ2 subunit conjugated with polyA ssDNA (polyA–Ab). PolyT–QD and polyA–QD were targeted to the polyA–Ab. (B) Representative QD images on neurons with polyA–Ab, after incubating with polyA–QD (top) or polyT–QD (bottom). Scale bar indicates 10 μm. (C) Number of QDs per field. Solid lines indicate the median. n = 30 images for both groups. (D) Schematic diagram of $GABA_AR$ QD labeling with oligo DNA-based labeling (oligo DNA) and conventional labeling with secondary antibody Fab fragments (Fab). (E) QD trajectories for oligo DNA-based labeling (left) and the conventional labeling (right). Scale bar indicates 1 μm. (F, G) Dot plot with the median solid lines (E) and cumulative probability distribution (F) of D for oligo-DNA-based or conventional DNA labeling. N = 969 (oligo DNA) and 686 (Fab). Mann–Whitney U-test for (C and F) and Kolmogorov–Smirnov test (G).

Multicolor imaging of the dynamics of lipids and proteins with oligo DNA-based QD-SPT

348
The results obtained so far demonstrate that oligo DNA-based QD-SPT can be applied to both membrane
lipids and membrane proteins, yielding results equivalent to those obtained using conventional labeling methods.
Finally, we investigated whether oligonucleotide-based QD-SPT enables simultaneous measurement of the
diffusion of multiple membrane molecules within the same neuron using spectrally distinct QDs.
Before labeling cells, we verified that QDs of different wavelengths could be reliably distinguished under
our imaging conditions. The QD605 and QD655 probes were mixed, immobilized on glass, and imaged within the
same field of view using the same acquisition settings applied to both probes. Under these conditions, QD605-
labeled particles were not detectably observed in the QD655 detection channel. Conversely, QD655 signals were
not detected under QD605 imaging conditions (Figure 6A). This indicates negligible channel cross-talk in our
dual-color imaging setup (Figure 6A).
$GABA_AR$ were then targeted with polyA–Ab and polyT–QD605 ($GABA_AR$–QD605), and random1S–DPPE
were labeled with random1AS–QD655 (DPPE–QD655) (Figure 6B). To assess potential cross-effects between
QD labeling of different molecular species, we compared single- and double-label conditions using neurons from
the same batch (Supplementary Figure S5A and S5B). The trajectories and D distributions for DPPE under
double-label conditions (DPPE + $GABA_AR$) were indistinguishable from those in single-label experiments
(Supplementary Figure S5C–S5E). Similarly, $GABA_AR$ trajectories and D distributions were unchanged under
double-label conditions (Supplementary Figure S5F–S5H). No statistically significant differences were observed
between single- and double-label conditions for either molecule (Mann–Whitney U test and Kolmogorov–
Smirnov test; n ≥ 298 trajectories per condition; $p > 0.05$). These results indicate that labeling one molecular
species does not measurably change the diffusion dynamics of the other within the same cellular context.
Finally, trajectories of $GABA_AR$–QD605 and DPPE–QD655 were analyzed. Representative 100-frame
trajectories (~8 s) of $GABA_AR$–QD605 appeared shorter than those of DPPE–QD655 (Figure 6C), suggesting
different diffusion behaviors. DPPE–QD655 (median: 1.15×10^{-1} μm²/s) exhibited a significantly higher D
than $GABA_AR$–QD605 (median: 2.32×10^{-2} μm²/s) (Figure 6D and Supplementary Table S2), and their D
distributions were clearly distinct (Figure 6E; Kolmogorov–Smirnov test, $p < 0.001$, $Dks = 0.300$). The median
D and interquartile ranges obtained using single-color QD-SPT were comparable to those obtained with
multicolor QD-SPT (Supplementary Table S2). These results demonstrate that two-color QD-SPT using ssDNA
enables simultaneous targeting and discrimination of two distinct membrane molecules based on QD emission
wavelength.
378

Discussion

Figure 6. DNA-mediated two-color QD-SPT of and DPPE.
(A) Images of QD605 and QD655 attached to glass, detected using wavelength-specific filters. (B) Strategy for multicolor QD labeling of GABA$_A$R with QD605 and DPPE with QD655 using oligoDNAs. (C) Representative trajectories of QD605–GABA$_A$R (left) and QD655–DPPE (right). Bars, 1 μm. (D, E) Distribution of D for QD605–GABA$_A$R and QD655–DPPE shown as a dot plot (solid lines, medians) (D) and cumulative distribution (E). n = 419 (QD605–GABA$_A$R) and 412 (QD655–DPPE). Statistical analysis: (D) Mann–Whitney U test; (E) Kolmogorov–Smirnov test.

In the present study, we demonstrated a method for labeling membrane molecules with QDs using the hybridization between ssDNAs with complementary sequences. This oligo DNA-based QD labeling was stable enough for tracking the dynamics of membrane molecules and compatible with conventional QD-SPT using a secondary antibody Fab. Furthermore, using different ssDNA sequences to label multiple types of membrane proteins with QDs of various wavelengths, we established multicolor QD-SPT, which enables observation of the dynamics of multiple membrane molecules, i.e., DPPE and GABA$_A$Rs, within the same cell.

QD labeling of membrane molecules via oligo DNA and dependence on the sequence

QDs exhibit a broad absorption spectrum and a narrow fluorescence spectrum, enabling excitation with a single light source and separation of multiple fluorescence signals. These QD characteristics are desirable properties for multicolor imaging, which enables the observation of different targets at various wavelengths in the same cell. Studies have labeled multiple types of molecules with QDs of different wavelengths in fixed cells [13] [14] [15]; however, most targets of QD-SPT in live cells were a single type of molecule. One of the reasons for this limitation was the combination of linkers connecting the membrane molecules and QD, such as the animal species of antibodies, and the number of protein-ligand interactions. In this study, we aimed to overcome this limitation of the number of combinations by applying hybridization between ssDNA, which generates numerous specific interactions based on the sequence. All 20 base oligo DNAs designed in this study, polyA/polyT pair, random1S/1AS pair, and random2S/2AS pairs, provided sufficient binding affinity and specificity, enabling hybridization-dependent labeling with QDs to DPPE molecules inserted in the plasma membrane. However, the number of QDs bound differed according to the sequence of linker oligo DNA. More QD bindings were observed when poly ssDNA was used as a linker, compared with random sequence. Under our experimental condition at 37°C, over 10-fold more QDs were found with polyA/polyT pair without guanine–cytosine pair than with random1S/1AS pair which has a higher Tm, although lower DNA duplex stability was expected for polyA/polyT combination (Tm: 52.17°C) compared with random1S/1AS pair (Tm: 64.47°C) [32]. One explanation for the higher binding of QDs by polyA/polyT pair is the high degree of freedom in sequence overlap, enabling stable binding even if only part of the 20-mer hybridizes. Further, ssDNAs with a random sequence can form secondary structures through base pairing. Indeed, secondary structure prediction of random ssDNA sequences using mFold [33] indicated the formation of secondary structures, which may inhibit the access to the complementary strand (Supplementary Figure S6), resulting in fewer observed QD counts. The minimum free energy change (ΔG) for these secondary structure formations is −1.74 kcal/mol or higher. Conversely, the free energy change during hybridization of complementary strands of random sequences calculated using DINAmelt [34] was −20 kcal/mol or higher, indicating that hybridization between complementary strands is more stable compared with secondary structure within the ssDNA. The DNA hybridization time was set to 1 min in the present study; however, extending the incubation time to reach equilibrium could potentially yield different results for the observed QD counts. However, the diffusion parameters, such as D, were not affected by the ssDNA sequence under conditions

where the number of QD bindings per field of view was adjusted to be suitable for SPT. These results indicate that
both poly ssDNA and random ssDNA are available for oligo DNA-based QD-SPT linkers.
Under the experimental conditions described here, the most crucial parameter for achieving an optimal
QD labeling density was the surface density of ssDNA moieties that are directly associated with the plasma
membrane (e.g., ssDNA-DPPE or ssDNA-conjugated anti-GABA$_A$R antibody). When the ssDNA-QD
concentration is kept constant, and the number of detected particles approaches the binding plateau, reducing the
membrane-associated ssDNA density is an effective first step to prevent over-labeling and minimize potential
multivalent crosslinking. Further, appropriate sequence design can tune hybridization efficiency, as random
sequences typically demonstrate lower binding efficiency than polyA–polyT pairs. Thus, selecting the ssDNA
sequence provides an additional means to modulate labeling density. Further reduction of ssDNA-QD
concentration can help fine-tune particle density for single-particle tracking if membrane ssDNA density remains
high after these adjustments.
427
Pros and cons of oligo DNA-based QD-SPT against SFT
As reported in other cultured cells [27] [28], the diffusion dynamics of phospholipids vary between the
surface and bottom layers of primary cultured rat hippocampal neurons. The D of DPPE labeled with ATTO647
was lower in the bottom layer than in the surface layer. Further, the D at the cell bottom obtained by oligo DNA-
based QD-SPT was significantly lower than that in the surface layer, with an increased proportion of immobile
QDs. Although limited to mobile QD fractions, the D of DPPE measured in the bottom layer was significantly
lower than that in the surface layer. These results indicate that oligo DNA-based QD-SPT can reliably detect
differences in diffusion dynamics at the subcellular level. However, only QD–oligo DNA–DPPE exhibited lower
D ($< 10^{-3}$ μm²/s) and immobile fractions (D < 0.0004 μm²/s) in the bottom layer, which were not observed in the
ATTO647–DPPE group. The size of organic dyes is approximately 1–2 nm, whereas QDs with functionalization
layers are approximately 15–30 nm [35]. Therefore, interactions between the QD as a probe and glycans or the
extracellular matrix may affect the lateral diffusion of cell membrane molecules. The diffusion of QD-conjugated
membrane proteins, calcium-activated potassium channels and B cell receptors, is significantly reduced at the
bottom layer of the cell compared with proteins labeled with organic dyes [36] [37]. The difference in diffusion
motion observed in this study between ATTO647-labeled DPPE and QD-labeled DPPE at the bottom layer of the
cell may indicate the existence of steric hindrance that affects phospholipid diffusion. However, notably, the
localization accuracy for single molecules decreases in single-fluorescence tracking due to the low photon
count[38]. Indeed, the apparent D of ATTO647-DPPE immobilized on glass was 2.4×10^{-2} μm²/s. These
molecules are physically immobilized; thus, this nonzero value reflects the localization precision and analytical
limit of the single-fluorophore tracking method. These results indicate that D at or below this value cannot be
reliably distinguished from measurement noise in the SFT approach. The main advantages of QDs are their bright
fluorescence, which provides a high SNR, and their long fluorescence lifetime, which enables stable tracking.
QD-SPT is a practical choice in environments where small spatial obstacles are expected, and when high
localization accuracy and long trajectories are required.
452
Effect of linkers on the protein diffusion measured by QD-SPT
We used the GABA$_A$R for the example of the membrane proteins to compare oligo DNA-based QD-SPT
with the widely used conventional QD-SPT using Fab fragments. The QD labeling achieved through ssDNA
hybridization also maintained sufficient specificity and stability for SPT, such as labeling using the interaction
between primary and secondary antibodies. The D of GABA$_A$R showed no significant difference in their median
values between the oligo DNA-based QD-SPT and the conventional QD-SPT. However, in the oligo DNA-QD-
SPT method, the proportion of QDs showed significantly higher D in the fraction with low D values (K–S test, p
$= 0.012$) compared with QD conjugated through Fab. The impact on the overall distribution of D was small (K–S
test D_{KS} value $= 0.080$); however, the potential influence of the linker between QDs and target molecules should
be fully considered when combining the DNA method and the antibody approach for membrane proteins.
The labeling using annealing between complementary ssDNA is an effective method for increasing the
number of specific intermolecular bonds; however, the process of fusing oligoDNAs to primary antibodies has
been a major challenge. Advancements in copper-free click chemistry, which was utilized for the DPPE label in
this study, have profoundly transformed the landscape of the field of bioconjugation. The increased availability of
modified oligoDNAs and reagents for click chemistry enabled the fusion of oligonucleotides to antibodies and

biomolecules in a few steps at room temperature via chemical reactions in biological laboratories [21]. Therefore,
protein labeling using oligonucleotides as linkers is expected to be an option in the future.
The hydrodynamic radius of the Fab antibody is approximately 2.5 nm [39], and the hydrodynamic radius
of the 20-base double-strand DNA is estimated to be 4–6 nm [40], indicating the comparability of the overall size
of the Fab-QD and DNA-QD probes. The root mean square deviation calculated by molecular dynamics
simulations is of the same order (1–5 Å), [41] [42] [43] [44], suggesting that differences in the size or rigidity of
the linker molecules cannot explain the difference in diffusion dynamics. The mechanism by which differences in
the linker result in variations in the D remains to be clarified in future studies.
476

Prospects for Multicolor QD-SPT

QDs with sharp fluorescent spectra are ideal probes for multicolor imaging. Labeling the same membrane
molecules with QDs of eight wavelengths helps avoid trajectory overlap in the same field of view, enabling
tracking of the dynamics of more QDs, thereby significantly increasing the throughput of QD-SPT [45]. In the
present study, we aimed to develop a new labeling method that utilizes the sequence specificity of ssDNA
hybridization to target QDs of different wavelengths to specific targets. We successfully fused QD655 and QD605
to different targets, namely lipids (DPPE) and membrane proteins (GABA$_A$R), respectively. The biotin–
streptavidin interaction used for linking QD and oligoDNA forms a firm non-covalent bond (dissociation constant
Kd $=10^{-15}$ M) [46], and the possibility of dissociation is expected to be extremely low. The biotin–streptavidin
binding was sufficiently stable not to allow cross-talk between QDs, considering the finding that DPPE and
GABA$_A$R diffusion, both of which have significantly different D values, could be distinguished using QD655 and
QD605.
Here, we performed a two-color QD-SPT as a proof of concept for the availability of oligo DNA for live-
cell multicolor imaging. Previous studies using fixed cells or fixed tissues have reported the application of QDs to
multicolor detection methods. Flow cytometry was performed using simultaneous detection of 17 fluorescent
probes, utilizing seven QDs and ten organic dyes [47]. Further, in single-cell imaging, a study revealed the
staining of human prostate cancer cells using five types of biomolecules with five different QDs (QD525, QD565,
QD605, QD655, and QD705) [48] [49]. This method requires dividing the labeling process into multiple steps and
repeating the blocking of the antigenic sites of the primary antibody with a secondary antibody at saturation levels
as many times as required. This approach is not ideal for SPT of live cells, which involves labeling to be
performed quickly and at low density. In contrast, oligo DNA-based QD-SPT uses oligo DNA as linkers, which
generates several combinations due to their sequence diversity, thereby enabling simultaneous labeling of multiple
target molecules with different-colored QDs in a short time (several minutes). In principle, multicolor QD-SPT
with three or more colors in the same cell is considered feasible; however, experimental verification remains a
future challenge.
To perform multicolor QD-SPT, the optical system needs to be optimized using the sharp spectrum of
QDs. In this study, the same field of view was recorded separately at each wavelength. Still, an optical system that
simultaneously records multiple colors of QDs is desirable to be developed to minimize observation time. An
image-splitting optical system can be employed to split the emission into two wavelengths and focus them onto a
single camera, thereby enabling simultaneous imaging of two colors [50]. Furthermore, using two cameras and an
image-splitting optical system, four colors of QDs within the same field is possible to be simultaneously imaged.
508

Conclusion

510   In this study, we revealed that a linker using hybridization between complementary ssDNA can be
511 utilized for the analysis of membrane molecular dynamics with QD-SPT. The DNA-based QD labeling method
512 developed in this study is expected to help target multiple types of molecules with different wavelength QDs in an
513 oligo DNA sequence-specific manner. It is considered to significantly enable simultaneous multicolor QD-SPT of
514 living cell membrane molecules.
515

Conflict of interest

517
518   The authors declare no conflict of interest related to this publication.
519

Author contributions

522 SS, TK, NK, BZ, TU, ME, RK, HY, HB conducted the experiment. SS, TK, NK, TU, BZ, ME, RK, CO, IK, and AC
523 performed the analysis. YT synthesized the ssDNA–DPPE. HB designed the study and supervision of this work. All
524 authors discussed the results and commented on the manuscript.
525
526 **Data availability**
527
528 The evidence data generated and/or analyzed during the current study are available from the corresponding author upon
529 reasonable request.
530
531 **Acknowledgments**
532

The authors thank Dr. Hiromu Kashida (Nagoya University) for designing the random oligo DNA sequences, Prof.
Akihiro Kusumi (OIST) for advice on single-fluorophore imaging of DPPE, Prof. Hiroaki Tateno for advice on
nucleotide conjugation to antibodies, and Dr. Atushi Okazawa (Waseda University) for the support of figure
preparation. This research was supported by grants from JST PRESTO JPMJPR15F8, JSPS Grants-in-Aid for Scientific
Research (23K08410 to SS; 23K18116 and 23K21299 to HB), and grants from Naito Foundation and Takeda Science
Foundation to HB. DeepL, ChatGPT 4o, Grammarly, and Enago, the editing brand of Crimson Interactive Pvt. Ltd
supported the English editing of this article.

540
541 **References**
542

[1] Singer, S.J., Nicolson, G.L. The fluid mosaic model of the structure of cell membranes. Science 175,
720-731 (1972). https://doi.org/10.1126/science.175.4023.720
[2] Fujiwara, T., Ritchie, K., Murakoshi, H., Jacobson, K., Kusumi, A. Phospholipids undergo hop
diffusion in compartmentalized cell membrane. J Cell Biol 157, 1071-1081 (2002).
https://doi.org/10.1083/jcb.200202050
[3] Fujiwara, T.K., Tsunoyama, T.A., Takeuchi, S., Kalay, Z., Nagai, Y., Kalkbrenner, T., et al. Ultrafast
single-molecule imaging reveals focal adhesion nano-architecture and molecular dynamics. J Cell Biol
222 (2023). https://doi.org/10.1083/jcb.202110162
[4] Poveda, J.A., Fernandez, A.M., Encinar, J.A., Gonzalez-Ros, J.M. Protein-promoted membrane
domains.   Biochim   Biophys   Acta   1778,   1583-1590   (2008).
https://doi.org/10.1016/j.bbamem.2008.01.021
[5] Choquet,   D.,   Triller,   A.   The   dynamic   synapse.   Neuron   80,   691-703   (2013).
https://doi.org/10.1016/j.neuron.2013.10.013
[6] Nakada, C., Ritchie, K., Oba, Y., Nakamura, M., Hotta, Y., Iino, R., et al. Accumulation of anchored
proteins forms membrane diffusion barriers during neuronal polarization. Nat Cell Biol 5, 626-632
(2003). https://doi.org/10.1038/ncb1009
[7] Alcor, D., Gouzer, G., Triller, A. Single-particle tracking methods for the study of membrane receptors
dynamics. Eur J Neurosci 30, 987-997 (2009). https://doi.org/10.1111/j.1460-9568.2009.06927.x
[8] Resch-Genger, U., Grabolle, M., Cavaliere-Jaricot, S., Nitschke, R., Nann, T. Quantum dots versus
organic   dyes   as   fluorescent   labels.   Nat   Methods   5,   763-775   (2008).
https://doi.org/10.1038/nmeth.1248
[9] Groc, L., Lafourcade, M., Heine, M., Renner, M., Racine, V., Sibarita, J.B., et al. Surface trafficking
of neurotransmitter receptor: Comparison between single-molecule/quantum dot strategies. J Neurosci
27, 12433-12437 (2007). https://doi.org/10.1523/jneurosci.3349-07.2007
[10] Cai, E., Ge, P., Lee, S.H., Jeyifous, O., Wang, Y., Liu, Y., et al. Stable small quantum dots for synaptic
receptor tracking on live neurons. Angew Chem Int Ed Engl 53, 12484-12488 (2014).
https://doi.org/10.1002/anie.201405735
[11] Ha, T., Tinnefeld, P. Photophysics of fluorescent probes for single-molecule biophysics and super-
resolution imaging. Annu Rev Phys Chem 63, 595-617 (2012). https://doi.org/10.1146/annurev-
physchem-032210-103340
[12] Medintz, I.L., Uyeda, H.T., Goldman, E.R., Mattoussi, H. Quantum dot bioconjugates for imaging,
labelling and sensing. Nat Mater 4, 435-446 (2005). https://doi.org/10.1038/nmat1390

[13] Arnspang, E.C., Brewer, J.R., Lagerholm, B.C. Multi-color single particle tracking with quantum dots.
PLoS One 7, e48521 (2012). https://doi.org/10.1371/journal.pone.0048521
[14] Zrazhevskiy, P., Gao, X. Quantum dot imaging platform for single-cell molecular profiling. Nature
Communications 4, 1619 (2013). https://doi.org/10.1038/ncomms2635
[15] Wang, W., Liu, Z., Lan, X. Quantum dot-based simultaneous multicolor imaging. Mol Imaging Biol
22, 820-831 (2020). https://doi.org/10.1007/s11307-019-01432-4
[16] Goltsev, Y., Samusik, N., Kennedy-Darling, J., Bhate, S., Hale, M., Vazquez, G., et al. Deep profiling
of mouse splenic architecture with codex multiplexed imaging. Cell 174, 968-981 e915 (2018).
https://doi.org/10.1016/j.cell.2018.07.010
[17] Zhong, S., Wang, R., Gao, X., Guo, Q., Lin, R., Luo, M. Modular DNA barcoding of nanobodies
enables multiplexed in situ protein imaging and high-throughput biomolecule detection. Elife 14
(2025). https://doi.org/10.7554/eLife.105225
[18] Jungmann, R., Avendano, M.S., Woehrstein, J.B., Dai, M., Shih, W.M., Yin, P. Multiplexed 3d
cellular super-resolution imaging with DNA-paint and exchange-paint. Nat Methods 11, 313-318
(2014). https://doi.org/10.1038/nmeth.2835
[19] Bannai, H., Hirose, M., Niwa, F., Mikoshiba, K. Dissection of local ca2+ signals in cultured cells by
membrane-targeted ca2+ indicators. J Vis Exp (2019). https://doi.org/10.3791/59246
[20] Teramura, Y. Cell surface modification with ssdna-peg-lipid for analysing intercellular interactions
between   different   cells.   Biomaterials   48,   119-128   (2015).
https://doi.org/10.1016/j.biomaterials.2015.01.032
[21] Maerle, A.V., Simonova, M.A., Pivovarov, V.D., Voronina, D.V., Drobyazina, P.E., Trofimov, D.Y.,
et al. Development of the covalent antibody-DNA conjugates technology for detection of ige and igm
antibodies   by   immuno-pcr.   PLoS   One   14,   e0209860   (2019).
https://doi.org/10.1371/journal.pone.0209860
[22] Sun, N., Tran, B.V., Peng, Z., Wang, J., Zhang, C., Yang, P., et al. Coupling lipid labeling and click
chemistry enables isolation of extracellular vesicles for noninvasive detection of oncogenic gene
alterations. Adv Sci (Weinh) 9, e2105853 (2022). https://doi.org/10.1002/advs.202105853
[23] Bannai, H., Inoue, T., Hirose, M., Niwa, F., Mikoshiba, K. Synaptic function and neuropathological
disease revealed by quantum dot-single-particle tracking. in Single molecule microscopy in
neurobiology. (Yamamoto, N., Okada, Y. eds.), pp. 131-155 Springer US, New York, NY (2020).
[24] Niwa, F., Bannai, H., Arizono, M., Fukatsu, K., Triller, A., Mikoshiba, K. Gephyrin-independent
gaba(a)r mobility and clustering during plasticity. PLoS One 7, e36148 (2012).
https://doi.org/10.1371/journal.pone.0036148
[25] Bannai, H., Niwa, F., Sherwood, M.W., Shrivastava, A.N., Arizono, M., Miyamoto, A., et al.
Bidirectional control of synaptic gabaar clustering by glutamate and calcium. Cell Rep 13, 2768-2780
(2015). https://doi.org/10.1016/j.celrep.2015.12.002
[26] Nishiguchi, T., Yoshimura, H., Kasai, R.S., Fujiwara, T.K., Ozawa, T. Synergetic roles of formyl
peptide receptor 1 oligomerization in ligand-induced signal transduction. ACS Chem Biol 15, 2577-
2587 (2020). https://doi.org/10.1021/acschembio.0c00631
[27] Tournier, J.F., Lopez, A., Tocanne, J.F. Effect of cell substratum on lateral mobility of lipids in the
plasma membrane of vascular endothelial cells. Exp Cell Res 181, 105-115 (1989).
https://doi.org/10.1016/0014-4827(89)90186-9
[28] Makaremi, S., Rose, M., Ranjit, S., Digman, M.A., Bowdish, D.M.E., Moran-Mirabal, J.M. Lateral
diffusion of cd14 and tlr2 in macrophage plasma membrane assessed by raster image correlation
spectroscopy   and   single   particle   tracking.   Scientific   Reports   10,   19375   (2020).
https://doi.org/10.1038/s41598-020-76272-2
[29] Muir, J., Arancibia-Carcamo, I.L., MacAskill, A.F., Smith, K.R., Griffin, L.D., Kittler, J.T. Nmda
receptors regulate gabaa receptor lateral mobility and clustering at inhibitory synapses through serine
327 on the γ2 subunit. Proc Natl Acad Sci U S A 107, 16679-16684 (2010).
https://doi.org/10.1073/pnas.1000589107
[30] Merlaud, Z., Marques, X., Russeau, M., Saade, U., Tostain, M., Moutkine, I., et al. Conformational
state-dependent regulation of gaba(a) receptor diffusion and subsynaptic domains. iScience 25,

105467 (2022). https://doi.org/10.1016/j.isci.2022.105467

[31] Battaglia, S., Renner, M., Russeau, M., Côme, E., Tyagarajan, S.K., Lévi, S. Activity-dependent
inhibitory synapse scaling is determined by gephyrin phosphorylation and subsequent regulation of
gaba_a receptor diffusion. eneuro 5, ENEURO.0203-0217.2017 (2018).
https://doi.org/10.1523/eneuro.0203-17.2017

[32] Breslauer, K.J., Frank, R., Blocker, H., Marky, L.A. Predicting DNA duplex stability from the base
sequence. Proc Natl Acad Sci U S A 83, 3746-3750 (1986). https://doi.org/10.1073/pnas.83.11.3746

[33] Zuker, M. Mfold web server for nucleic acid folding and hybridization prediction. Nucleic Acids Res
31, 3406-3415 (2003). https://doi.org/10.1093/nar/gkg595

[34] Markham, N.R., Zuker, M. Dinamelt web server for nucleic acid melting prediction. Nucleic Acids
Res 33, W577-581 (2005). https://doi.org/10.1093/nar/gki591

[35] Manzo, C., Garcia-Parajo, M.F. A review of progress in single particle tracking: From methods to
biophysical insights. Rep Prog Phys 78, 124601 (2015). https://doi.org/10.1088/0034-
4885/78/12/124601

[36] Nechyporuk-Zloy, V., Dieterich, P., Oberleithner, H., Stock, C., Schwab, A. Dynamics of single
potassium channel proteins in the plasma membrane of migrating cells. Am J Physiol Cell Physiol 294,
C1096-1102 (2008). https://doi.org/10.1152/ajpcell.00252.2007

[37] Abraham, L., Lu, H.Y., Falcão, R.C., Scurll, J., Jou, T., Irwin, B., et al. Limitations of qdot labelling
compared to directly-conjugated probes for single particle tracking of b cell receptor mobility.
Scientific Reports 7, 11379 (2017). https://doi.org/10.1038/s41598-017-11563-9

[38] Ober, R.J., Ram, S., Ward, E.S. Localization accuracy in single-molecule microscopy. Biophys J 86,
1185-1200 (2004). https://doi.org/10.1016/s0006-3495(04)74193-4

[39] Shatz, W., Hass, P.E., Mathieu, M., Kim, H.S., Leach, K., Zhou, M., et al. Contribution of antibody
hydrodynamic size to vitreal clearance revealed through rabbit studies using a species-matched fab.
Mol Pharm 13, 2996-3003 (2016). https://doi.org/10.1021/acs.molpharmaceut.6b00345

[40] Fernandes, M.X., Ortega, A., López Martínez, M.C., García de la Torre, J. Calculation of
hydrodynamic properties of small nucleic acids from their atomic structure. Nucleic Acids Res 30,
1782-1788 (2002). https://doi.org/10.1093/nar/30.8.1782

[41] Young, M.A., Ravishanker, G., Beveridge, D.L. A 5-nanosecond molecular dynamics trajectory for
b-DNA: Analysis of structure, motions, and solvation. Biophys J 73, 2313-2336 (1997).
https://doi.org/10.1016/s0006-3495(97)78263-8

[42] Dixit, S.B., Ponomarev, S.Y., Beveridge, D.L. Root mean square deviation probability analysis of
molecular dynamics trajectories on DNA. J Chem Inf Model 46, 1084-1093 (2006).
https://doi.org/10.1021/ci0504925

[43] Li, T., Tracka, M.B., Uddin, S., Casas-Finet, J., Jacobs, D.J., Livesay, D.R. Rigidity emerges during
antibody evolution in three distinct antibody systems: Evidence from qsfr analysis of fab fragments.
PLoS Comput Biol 11, e1004327 (2015). https://doi.org/10.1371/journal.pcbi.1004327

[44] Tang, J., Zhang, C., Castillo, N.C., Lalaurie, C.J., Gao, X., Dalby, P.A., et al. Crystal structures and
molecular dynamics simulations of a humanised antibody fragment at acidic to basic ph. Scientific
Reports 13, 16281 (2023). https://doi.org/10.1038/s41598-023-42698-7

[45] Cutler, P.J., Malik, M.D., Liu, S., Byars, J.M., Lidke, D.S., Lidke, K.A. Multi-color quantum dot
tracking using a high-speed hyperspectral line-scanning microscope. PLoS One 8, e64320 (2013).
https://doi.org/10.1371/journal.pone.0064320

[46] Green, N.M. Avidin. in Advances in protein chemistry. (Anfinsen, C.B., Edsall, J.T., Richards, F.M.
eds.) vol. 29, pp. 85-133 Academic Press, (1975).

[47] Chattopadhyay, P.K., Perfetto, S.P., Yu, J., Roederer, M. The use of quantum dot nanocrystals in
multicolor flow cytometry. Wiley Interdisc Rev Nanomed Nanobiotechnol 2, 334-348 (2010).
https://doi.org/10.1002/wnan.75

[48] Xing, Y., Smith, A.M., Agrawal, A., Ruan, G., Nie, S. Molecular profiling of single cancer cells and
clinical tissue specimens with semiconductor quantum dots. Int J Nanomedicine 1, 473-481 (2006).
https://doi.org/10.2147/nano.2006.1.4.473

[49] Xing, Y., Chaudry, Q., Shen, C., Kong, K.Y., Zhau, H.E., Chung, L.W., et al. Bioconjugated quantum

dots for multiplexed and quantitative immunohistochemistry. Nat Protoc 2, 1152-1165 (2007).
https://doi.org/10.1038/nprot.2007.107
[50]   Juette, M.F., Terry, D.S., Wasserman, M.R., Zhou, Z., Altman, R.B., Zheng, Q., et al. The bright future
of single-molecule fluorescence imaging. Curr Opin Chem Biol 20, 103-111 (2014).
https://doi.org/10.1016/j.cbpa.2014.05.010
684

Table S1 ssDNA nucleotide sequences.

List of ssDNA sequences used and melting temperatures (Tm) estimated by nearest neighbor base pairing (Breslauer et al., 1986). PolyT and PolyA, random1S and 1AS, random2S and 2AS, are complementary pairs.

ssDNA name	Sequence	Tm (°C)
polyA	5'-AAAAAAAAAAAAAAAAAAAA-3'	52.17
polyT	5'-TTTTTTTTTTTTTTTTTTTT-3'	52.17
Random1S	5'-ATGAGGTCGTAAGCGTTCCA-3'	64.47
Random1AS	5'-TGGAACGCTTACGACCTCAT-3'	64.47
Random2S	5'-GCTACCTATGCCGTCAATGG-3'	64.21
Random2AS	5'-CCATTGACGGCATAGGTAGC-3'	64.21

Table S2: Summary of the median and Interquartile Range (IQR) for the data
The median and IQR of each data sets are shown here.

Figure	group	Median (spots/image)	IQR (25% – 75%) (spots/image)	N(images)
Figure 2C	polyA-QD	1	0 – 2	10
	polyT-QD	241	96 – 400	10
Figure 2F	polyA	3	2 – 4	10
	polyT	3	2 – 4	10
	random1AS	40	26 – 49	10
	random2AS	1	0 – 2	10
Figure 2I	mock (0min)	25	17 – 29	11
	mock (3min)	27	20 – 30	
	DNase (0min)	31	24 – 33	11
	DNase (3min)	10	4 – 11	
Figure 4C	polyA-polyT	193	169 – 206	5
	random1S-1AS	20	17 – 25	5
Figure 5C	polyA-QD	2	1 – 3	30
	polyT-QD	28	22 – 44	30
Figure S1D	polyA-QD	1	0 – 1	29
	polyT-QD	22	15 – 26	30
Figure S2B	polyT-DPPE + polyA-QD	23	20 – 24	6
	polyA-QD	0	0 – 0	6
Figure S2D	0.063 nM	0	0 – 0	19
	0.63 nM	4	3 – 5	21
Figure S2F	random1AS	1	1 – 3	10
	random2AS	1	0 – 3	10

Figure	group	Median ($\mu m^2/s$)	IQR (25% – 75%) ($\mu m^2/s$)	N
Figure 3C **3D**	glass attached	2.40×10^{-2}	$1.66 - 3.03 \times 10^{-2}$	152 spots
	lower	1.15×10^{-1}	$5.70 - 26.6 \times 10^{-2}$	368 spots
	upper	2.21×10^{-1}	$1.26 - 3.43 \times 10^{-1}$	584 spots
Figure 3H **3I**	lower	8.23×10^{-2}	$0.387 - 20.6 \times 10^{-2}$	879 QDs
	upper	1.86×10^{-1}	$9.13 - 31.5 \times 10^{-2}$	815 QDs
Figure 4E **4F**	polyA-polyT	1.38×10^{-1}	$7.74 - 22.6 \times 10^{-2}$	219 QDs
	random1S-1AS	1.47×10^{-1}	$7.86 - 23.0 \times 10^{-2}$	253 QDs
Figure 5F **5G**	ssDNA	6.49×10^{-2}	$1.41 - 12.3 \times 10^{-2}$	969 QDs
	Fab	5.19×10^{-2}	$0.841 - 12.8 \times 10^{-2}$	686 QDs
Figure 6D **6E**	GABA$_A$R	2.32×10^{-2}	$0.300 - 9.56 \times 10^{-2}$	419 QDs
	DPPE	1.15×10^{-1}	$0.826 - 24.5 \times 10^{-2}$	412 QDs
Figure S6D **S6E**	DPPE (alone)	9.74×10^{-2}	$1.14 - 21.6 \times 10^{-2}$	451 QDs
	DPPE (+GABA$_A$R)	9.18×10^{-2}	$1.78 - 22.9 \times 10^{-2}$	611 QDs
Figure S6G **S6H**	GABA$_A$R (alone)	2.38×10^{-2}	$0.19 - 11.1 \times 10^{-2}$	298 QDs
	GABA$_A$R (+DPPE)	2.72×10^{-2}	$0.29 - 10.5 \times 10^{-2}$	300 QDs

Figure	group	Median (%)	IQR (25% – 75%) (%)	N (images)
Figure 3G	lower	17.3	4.53 – 25.1	28
	upper	2.16	0 – 7.80	28

Figure	group	Median (max $\Delta F/F_0$)	IQR (25% – 75%) (max $\Delta F/F_0$)	N (images)
Figure S4D	Unlabeled (–muscimol)	2.58	2.34 – 2.86	9
	Unlabeled (+muscimol)	0.48	0.24 – 0.67	14
	$GABA_AR$ (–muscimol)	2.03	1.30 – 2.34	14
	$GABA_AR$ (+muscimol)	0.18	0.14 – 0.35	12
	DPPE (–muscimol)	2.84	1.30 – 3.45	9
	DPPE (+muscimol)	0.16	0.13 – 0.18	11

Supplementary Figure S1

Effect of the linker between DPPE and ssDNA on QD-labeling specificity
(A) PolyA–DPPE generated via thiol–maleimide coupling. (B) PolyA–DPPE–DBCO
generated via DBCO–azide click chemistry. (C) Representative images of polyA–DPPE–
DBCO labeled with polyA–QD (left) or polyT–QD (right). Bar, 10 μm. (D) Number of QDs
per field (41.8 × 41.8 μm; solid lines, medians; n = 29 images for polyA–QD and n = 30
images for polyT–QD). Mann–Whitney U test.

Supplementary Figure S2

Specificity of ssDNA–QD binding to the DPPE

A) Representative images of neurons incubated with polyA–QD in the presence (left) or absence
(right) of polyT–DPPE. (B) Number of QDs per field (41.8 × 41.8 µm; solid lines, medians).
(C) Representative images of neurons incubated with 0.063 or 0.63 nM polyA–QD. (D) Number of
QDs per field (solid lines, medians). (E) Representative images of neurons expressing polyT–DPPE
labeled with QDs carrying random sequences (random1AS or random2AS). (F) Number of QDs per
field (solid lines, medians).
Bars in (A), (C), and (E), 10 µm. Statistical analysis: (B, D) Mann–Whitney U test. N is shown in
parentheses (see Table S2).

Supplementary Figure S3

Number of QDs at different concentrations of random 1S-DPPE and random 2S-DPPE.
(A, C) Representative QD images for random 1S-DPPE + random 1AS-QD (A) and random 2S-DPPE + random 2AS-QD (C) at the indicated DPPE concentrations (0, 2, 5, 20, 50, 100 µg/mL). Bar, 10 µm.
(B, D) Number of QDs per field of view (41.8 × 41.8 µm) for random 1S-DPPE (B) and random 2S-DPPE (D). Each point represents one field. n = 5 images (0 µg/mL) and 7 images (others) for random 1S-DPPE; n = 5 images (0 µg/mL), 7 images (2–50 µg/mL), and 9 images (100 µg/mL) for random 2S-DPPE.

Comparison of KCl-induced intracellular Ca²⁺ responses in the presence or absence of muscimol under QD-labeled and unlabeled conditions.

(A–C) QD images, Fluo-8 images, pseudocolor $\Delta F/F_0$ images before and after 40 mM KCl application, and representative $\Delta F/F_0$ traces of neurons with or without preincubation with 20 μM muscimol. Bar, 20 μm. (A) Unlabeled cells; (B) GABA$_A$R-labeled cells; (C) DPPE-labeled cells.

(D) Maximum $\Delta F/F_0$ with (+) or without (–) muscimol under unlabeled, GABA$_A$R-labeled, and DPPE-labeled conditions (solid lines, medians). n = 9 (unlabeled), 14 (unlabeled + muscimol), 14 (GABA$_A$R-labeled), 12 (GABA$_A$R-labeled + muscimol), 9 (DPPE-labeled), and 11 (DPPE-labeled + muscimol). Statistical analysis: Mann–Whitney U test.

Supplementary Figure S5

Comparison of diffusion coefficients under single-color and multicolor QD-labeling conditions.
(A) Strategy for single-color and multicolor QD labeling of GABA$_A$R (QD605) and DPPE (QD655) using oligoDNAs. (B) Representative QD-labeling images. Bars, 10 μm. (C) Representative trajectories of DPPE under single-color and multicolor labeling. Bar, 1 μm. (D, E) Distribution of D for DPPE shown as a dot plot (D) and cumulative distribution (E) (solid lines, medians). n = 451 (DPPE alone) and 611 (DPPE + GABA$_A$R). (F) Representative trajectories of QD-GABA$_A$R under single-color and multicolor labeling. Bar, 1 μm. (G, H) Distribution of D for GABA$_A$R shown as a dot plot (G) and cumulative distribution (H) (solid lines, medians). n = 298 (GABA$_A$R alone) and 300 (GABA$_A$R + DPPE). Statistical analysis: (D, G) Mann–Whitney U test; (E, H) Kolmogorov–Smirnov test.

Supplementary Figure S6

Secondary structures predicted by mfold [33] for the random ssDNAs used in this experiment. Red lines indicate canonical base pairs of Watson–Crick type, which contribute significantly to the thermodynamic stability of the structure. Green lines indicate non-canonical or weak base pairing; ΔG denotes free energy change [kcal/mol].