Matrix Matters: Context-Driven Metabolic Shifts in *Bacillus cereus* and *Bacillus subtilis*

Emilie Viktoria Skriver[1,a], Tessa Suzanne Canoy[1,a], Yuandong Sha[1], Morten Arendt Rasmussen[1,2], Bekzod Khakimov[1], Susanne Knøchel[1], Henriette Lyng Røder*[1]

1: Department of Food Science, University of Copenhagen, Copenhagen, Denmark

2: Copenhagen Studies on Asthma in Childhood, Herlev-Gentofte Hospital, University of Copenhagen, Copenhagen, Denmark

[a] Emilie Viktoria Skriver and Tessa Suzanne Canoy contributed equally to this work and share first authorship.

*Corresponding author: h.lyng.roeder@food.ku.dk
            Section for Microbiology and Fermentation
            Rolighedsvej 26
            DK-1958 Frederiksberg

16   Abstract

Spore-forming *Bacillus* species, including pathogenic *Bacillus cereus* and spoilage-associated *Bacillus*

subtilis, are major contributors to foodborne illness and product degradation. Understanding their

metabolic behaviour in diverse food matrices is essential for improving risk assessment, spoilage

prediction, and fermentation control. This study integrates isothermal microcalorimetry and targeted

metabolomics to characterize the metabolic activity of *B. cereus* and *B. subtilis* in five nutrient sources:

Brain Heart Infusion (BHI) medium, oat drink, milk, pea hydrolysate, and a combined oat-pea matrix.

Metabolic heat production was monitored for 24 hours at 30°C. In BHI, *B. cereus* exhibited a shorter

lag phase (mean ± sd: 4.3 hours ± 0.8) than *B. subtilis* (7.9 hours ± 1.0) but produced less total heat.

Across all food matrices, *B. subtilis* consistently generated more heat. The oat-pea matrix supported the

highest growth rates, surpassing oat or pea alone, and showed sugar depletion and accumulation of

organic acids, indicating enhanced carbohydrate metabolism. Free amino acid release was matrix- and

species-specific: *B. subtilis* had increased levels in oat, while *B. cereus* did so in pea. While *B. cereus*

was metabolically active in all matrices, cereulide levels were matrix-dependent: 47.3 ± 1.7 ng/mL in

oat, 3.0 ± 0.1 ng/mL in oat–pea, and undetectable in pea. These findings reveal clade-specific and

matrix-driven metabolic strategies. This is the first study to combine calorimetry and metabolomics to

evaluate *Bacillus* activity in plant-based and dairy matrices. This approach enhances our understanding

of microbial physiology in complex food systems and provides a foundation for developing targeted

strategies to improve food safety, stability, and product design.

Keywords: *Bacillus cereus*, *Bacillus subtilis*, isothermal calorimetry, microbial metabolism, cereulide,

food matrix, plant-based food, metabolomics

1. Introduction

Microbial contamination is a key concern in the food industry (Snyder et al., 2024). In 2023, the United Nations estimated that 19% of all food available at the consumer level was wasted, which has a significant impact on the economy, environment, and public health (UNEP, 2024). The Gram-positive, spore-forming, facultative aerobe *Bacillus* spp. are ubiquitous in nature and include species such as *B. cereus* and *B. subtilis*. *Bacillus* species are usually introduced into the food production system through raw ingredients (Ehling-Schulz et al., 2015). *B. cereus* and *B. subtilis* both produce highly resistant spores, complicating their control with conventional techniques like heat, acid, and cleaning practices (Earl et al., 2008; Rahnama et al., 2023).

Bacillus cereus is an important foodborne pathogen, capable of causing diarrhoea due to enterotoxins or inducing vomiting due to the pre-formed heat- and acid-stable emetic toxin, cereulide (Ehling-Schulz et al., 2004). The *B. cereus* group, also known as *B. cereus sensu lato* (hereafter referred to as *B. cereus*), contains many genetically closely related species, including *B. cereus, B. mycoides,* and *B. thuringiensis*. *B. cereus* taxonomy has undergone many changes over the years, especially after the introduction of genome sequencing, and numerous nomenclatural frameworks have been used to describe *B. cereus* taxonomy (Carroll et al., 2022).

Foodborne disease caused by *B. cereus* is believed to be grossly underreported. Nevertheless, in 2022, *B. cereus* toxins (both emetic and enterotoxins) caused 306 registered foodborne outbreaks, 3192 cases of illness, 66 hospitalizations, and 2 deaths in the EU and UK alone (EFSA, 2023). *B. cereus* is typically isolated from plant materials (Messelhäusser et al., 2014), and given the increasing development of plant-based ingredients for food production (Kyrylenko et al., 2023). There is a strong need to monitor the prevalence and toxicity of *B. cereus* in plant matrices.

B. subtilis is a key driver of many traditional fermentations based on legumes, seeds, and tubers (Li et al., 2023). A high occurrence of *B. cereus* has been observed in *B. subtilis*-fermented food products, and its presence poses a significant concern for food safety (Owusu-Kwarteng et al., 2022). *B. subtilis* is also associated as a spoilage agent in foods, such as slime formation in rice products and ropiness in bread (Dong et al., 2023; Valerio et al., 2008). Consequently, understanding the metabolic characteristics, growth dynamics, and responses of *B. cereus* and *B. subtilis* is essential for assessing food safety and quality risks, particularly for plant-based food products.

An attractive tool for fast detection of microbial activity is isothermal microcalorimetry (IMC). IMC is a non-invasive method based on real-time, accurate measurements of heat flow generated by the metabolic activity of biological systems, especially in opaque or complex matrices where optical methods are limited. IMC equipment has become less costly and has demonstrated improved sensitivity

and accuracy (Braissant et al., 2010). IMC has recently been successfully applied to improve bacterial

detection time in a clinical context (Cichos et al., 2023) and to investigate the metabolic activity of

bacteria living in biofilms (Lichtenberg et al., 2022). Despite its potential relevance in non-invasive risk

assessment of food, IMC has not yet been explored as a tool for monitoring the growth of pathogenic

and non-pathogenic *Bacillus spp*.

While IMC provides valuable insights into the overall metabolic activity of microorganisms, it offers

limited specificity regarding the biochemical processes underlying bacterial metabolism. Monitoring

changes in key metabolites such as sugars, organic acids, and amino acids enables a deeper

understanding of microbial nutrient assimilation while also revealing biochemical changes in the food

matrices that are relevant to safety and quality. Sugars serve as primary energy sources for microbial

growth, while organic acids are key intermediates in carbon metabolism (Schilling et al., 2007). Free

amino acids are an important nitrogen source and can be released either through proteolytic activity or

produced through de novo synthesis (Sonenshein, 2007). Furthermore, quantifying the heat-stable

emetic toxin cereulide is essential for evaluating the safety risks associated with food products

contaminated by *B. cereus*.

To advance understanding of *Bacillus* physiology in food systems, this study integrates isothermal

microcalorimetry (IMC) with targeted metabolomic profiling. We investigate the metabolic heat

production of *B. cereus* and *B. subtilis* in a rich laboratory medium and in dairy- and plant-based

matrices, while quantifying key metabolites—sugars, organic acids, free amino acids, and cereulide—

at selected time points. This combined approach offers a more nuanced view of microbial activity and

its implications for food safety and quality.

2. Materials and methods

2.1 Bacterial strains

Sixteen strains from the *B. cereus* group and eight *B. subtilis* strains were selected based on diversity in

toxicity, genetic characteristics, and origin (Table 1). Five of the included *B. subtilis* strains were

previously isolated from traditional fermentations in Ghana and Burkina Faso (Kpikpi et al., 2014)

(Parkouda et al., 2010; Wiedenbein et al., 2023). Two of the *B. cereus* strains were isolated from African

fermented condiments (Thorsen et al., 2010). Strains were maintained in a -80°C glycerol stock and

streaked onto Brain Heart Infusion (BHI) agar plates before being cultured in BHI broth.

TABLE 1: Characteristics of the *Bacillus* strains used in this study, based on sequencing and ANIm data for the *Bacillus cereus* group obtained by Skriver et

al. (2025) and for the *Bacillus subtilis* strains obtained by Wiedenbein et al. (2023). Asterisk (*) denotes values below species threshold (<96% ANIm). The

letter T refers to the type strain.

Strain	Toxin production	Origin isolation	Previously assigned taxonomy (ANIm in %)	Taxonomy reference	Best taxonomic match (ANIm in %)	Assembly size (basepairs)	GC content
F4433/73	Enterotoxin	Meatloaf, USA	*Bacillus cereus* (91.68*)	(Midura et al., 1970)	*Bacillus tropicus* (96.26)	5,433,211	35.5
O-216	Unknown	Milk, Denmark	*Bacillus cereus* (91.89*)	(Larsen & Jørgensen, 1999)	*Bacillus pacificus* (99.94)	5,655,403	35.4
LWL3	Enterotoxin	Foodborne outbreak	*Bacillus cereus* (91.68*)	(Dufrenne et al., 1995)	*Bacillus wiedmannii* (96.43), *Bacillus pretiosus* (97.1%)	5,686,534	35.3
59	Unknown	Cream	*Bacillus cereus* (91.72*)	(Stenfors & Granum, 2001)	*Bacillus wiedmannii* (96.56), *Bacillus pretiosus* (96.7%)	5,516,290	35.5
MC67	Emetic	Sandy soil, Denmark	*Bacillus weihenstephanensis[a]* (95*)	(Hendriksen et al., 2006)	*Bacillus mycoides* (95*)	5,802,073	35.5
A536	Enterotoxin	Milk, silotank	*Bacillus cereus* (89.96*)	(Svensson et al., 2004)	*Bacillus mycoides* (97.23)	6,122,672	35.3
NVH 0075-95	Enterotoxin	Foodborne outbreak in Norway	*Bacillus cereus* (92.08*)	(T. Lund & Granum, 1996)	*Bacillus paranthracis* (97.61)	6,044,751	35.3
F4810/72	Emetic	Foodborne outbreak, UK	*Bacillus cereus* (92.09*)	(Turnbull et al., 1979)	*Bacillus paranthracis* (97.62)	5,584,855	35.5
NS117	Emetic	Spruce tree, Norway	*Bacillus cereus* (92.04*)	(Hallaksella et al., 1991)	*Bacillus paranthracis* (97.61)	5,832,990	35.3
LKT1/1	Emetic	Filler material of a building with moisture damage, Finland	*Bacillus cereus* (92.06*)	(Apetroaie et al., 2005)	*Bacillus paranthracis* (97.62)	5,690,232	35.5
Y48H10	Emetic	Iru, Africa	*Bacillus cereus* (92.09*)	(Thorsen et al., 2010)	*Bacillus paranthracis* (97.62)	5,549,004	35.5

Ba18H2	Emetic	Sonru, Africa	*Bacillus cereus* (92.09*)	(Thorsen et al., 2010)	*Bacillus paranthracis* (97.62)	5,598,436	35.5
SDA GR177	Emetic + Enterotoxin	Milk	*Bacillus cereus* (92.09*)	Svensk Mjolk, Swedish Dairy Association	*Bacillus paranthracis* (97.62)	5,634,738	35.5
SDA MA60	Enterotoxin	Milk	*Bacillus cereus* (92.06*)	Svensk Mjolk, Swedish Dairy Association	*Bacillus paranthracis* (97.62)	5,644,507	35.5
F4430/73	Emetic	Foodborne outbreak, UK	*Bacillus cereus* (92.06*)	(Spira & Goepfert, 1975)	*Bacillus paranthracis* (97.62)	5,661,860	35.5
ISI2489	Emetic	Unknown	*Bacillus cereus* (92.08*)	Arla culture collection	*Bacillus paranthracis* (97.64)	5,541,441	35.5
DSM13019		Boiled hay, Denmark	*Bacillus subtilis* (98.53)	(Wiedenbein et al., 2023)	*Bacillus subtilis* (98.53)	4,044,377	43.6
DSM10 T		Unknown ("Marburg")	*Bacillus subtilis* (99.99)	(Lilge et al., 2021; Wiedenbein et al., 2023)	*Bacillus subtilis* (99.99)	4,299,854	43.4
DSM21393		potato tubers, Tatarstan, Russian Federation	*Bacillus subtilis* (98.8)	(Wiedenbein et al., 2023)	*Bacillus subtilis* (98.8)	4,100,908	43.9
PRO35		Kantong, Nyankpala, Ghana	*Bacillus subtilis* (98.4)	(Kpikpi et al., 2014; Wiedenbein et al., 2023)	*Bacillus subtilis* (98.4)	4,032,381	43.4
PRO31		Kantong, Nyankpala, Ghana	*Bacillus subtilis* (98.35)	(Kpikpi et al., 2014; Wiedenbein et al., 2023)	*Bacillus subtilis* (98.35)	4,272,877	43.2
PRO64		Kantong, Nyankpala, Ghana	*Bacillus subtilis* (97.9)	(Wiedenbein et al., 2023)	*Bacillus subtilis* (97.9)	4,113,761	43.6
PRO99		Kantong, Navrongo, Ghana	*Bacillus subtilis* (98.2)	(Wiedenbein et al., 2023)	*Bacillus subtilis* (98.2)	3,963,991	43.4
PRO115		Maari, Mansila, Burkina Faso	*Bacillus subtilis* (100)	(Wiedenbein et al., 2023)	*Bacillus subtilis* (100)	4,140,503	43.8

103   Footnote *(a): B. mycoides* has been used for ANIm comparison, as *B. mycoides* is a later synonym of *B. weihenstephanensis* (Liu et al., 2018).

2.2 Whole genome sequencing

Bacillus subtilis whole genomes included in this work were previously characterized in Wiedenbein et al. (2023), while the *Bacillus cereus* whole genomes were obtained from Skriver et al. (2025).

2.3 Phylogenetic tree construction

Phylogenetic trees were constructed with the up-to-date bacterial core gene (UBCG) method based on the concatenation of 92 core genes found in all taxonomic ranks of Bacteria (Na et al., 2018). FastTree was selected to generate the maximum-likelihood phylogenetic trees (Price et al., 2010). The phylogenetic trees were visualized with Molecular Evolutionary Genetics Analysis version 11 software (MEGA 11) (Tamura et al., 2021). *Halobacillus halophilus* DSM 2266 (type strain) (GenBank: GCA_000284515.1) was selected as the outgroup to root the trees.

2.4 Isothermal calorimetric assessment of *B. cereus* and *B. subtilis* in BHI medium

All sixteen *B. cereus* and eight *B. subtilis* strains (Table 1) were cultivated in BHI broth (Merck). To measure the heat flow of the *Bacillus* strains in BHI medium, the protocol of Beilharz et al. (2023) was followed with a few modifications. Strains were cultured in BHI broth overnight (30°C, 200 rpm) and then diluted with BHI broth to an optical density at 600 nm (OD600) of 0.2. The inoculum was mixed with fresh BHI medium, with an inoculation rate of 1% and a total volume of 200 µL. The inoculated mixtures were transferred to plastic inserts (calWells™, Symcel AB, Sweden) in the titanium vials of a 48-well calPlate™ (Symcel AB, Sweden). The vials were pre-incubated for 60 min to reach temperature equilibrium. The plates were then incubated in the Calscreener (Symcel AB, Sweden) at 30°C for 20h.

2.5 Isothermal calorimetric assessment of *B. cereus* and *B. subtilis* in food matrices

Oat drinks (Naturli, Denmark) and UHT whole milk (Geia Food A/S, Czech Republic) were purchased commercially in Danish supermarkets. For the mixed oat-pea matrix, 5% w/w pea protein hydrolysate (NUTRALYS® PEA F853M-EXP, Roquette, France) was transferred to store-bought oat drink and autoclaved (121°C, 20 min). For the pea matrix, 5% w/w was added to MilliQ water and autoclaved (121°C, 20 min). A selection of three *B. cereus* strains (Y48H10, SDA GR177, MC67) and three *B. subtilis* strains (PRO99, DSM10, DSM21393) were cultivated in the food matrices.

Cells were cultured in BHI broth overnight (30°C, 200 rpm), centrifuged at 4500 *g* for 12 minutes, and washed twice in 0.9% NaCl solution. Washed cells were adjusted to OD600 of 0.2 before being transferred with an inoculation rate of 1% and a total volume of 200 µL into the different food matrices. Similarly to section 2.5, the inoculated mixtures were transferred to plastic inserts in the titanium vials

136 of the 48-well calPlate and pre-incubated for 60 min to reach temperature equilibrium. The calPlate was
137 then incubated in the Calscreener at 30°C for 24 hours.

2.6 Cultivation in oat, pea, and oat-pea matrices for metabolic profiling

One strain from each species (*B. cereus* Y48H10 and *B. subtilis* DSM21393) was selected for in-depth
metabolic profiling. Washed cells were prepared as in section 2.5 and then inoculated in oat, pea, and
oat-pea matrices. For this experiment, two biological replicates, each consisting of four technical
replicates, were used. Biological replicates were defined as individual colonies streaked from the same
BHI plate, while technical replicates originated from the same biological replicate after two rounds of
pre-cultivation on BHI broth. Samples were taken at maximum calorimetric growth rate, approximately
8 hours for *B. cereus* and 11 hours for *B. subtilis*, and at the end-metabolic point, 24 hours for both *B.*
cereus and *B. subtilis*. Directly after sampling, technical replicates were pooled to obtain a final volume
of approximately 800 μL. This pooled sample was analysed for sugars and organic acids (sections 2.7),
free amino acids (section 2.8), and the *B. cereus* samples also for cereulide (section 2.9).

2.7 Quantification of simple sugars and organic acids produced by HPLC-RID

Pooled technical replicates of food samples were centrifuged at 12000 *g* for 15 minutes, and the
supernatant was separated and filtered through PTFE membranes with a pore size of 0.45 μm directly
into the HPLC vials and stored at -21°C until analysis.

An Agilent 1260 Infinity II LC system with refractive index detection (Agilent Technologies, Santa
Clara, CA, U.S.A.) was used to quantify simple sugars and organic acids at 35°C. The mobile phase
was water, with a flow rate of 0.5 mL/min. A Cation H Bio-Rad Micro-Guard column (304.6 mm)
followed by an Aminex HPX-87H carbohydrate analysis column (300 × 7.8 mm) (Bio-Rad
Laboratories, Inc.; Hercules, CA, U.S.A.) was used to separate organic acids and simple sugars
isocratically at 35°C. The injection volume of the sample was 10 μL and the run time was 15 minutes.

The chromatographic data were acquired, monitored, and processed using OpenLAB CDS LC
Chemstation M8307AA. Quantification and assignment of peaks were based on the peak areas and
retention times of known standard curves for purified external calibration curves prepared using seven
standard solutions for sugars (glucose, fructose, arabinose, galactose, and sucrose) and organic acids
(lactic-, acetic-, and citric acid) in concentrations ranging from 0.05 – 10 g/L.

2.8 Free Amino Acid analysis

Samples were freeze-dried (Lyovapor L-200, Buchi) for two days. To each freeze-dried sample, 1 mL
of 0.1 M HCl was added, and the mixture was ultrasonicated (Branson 2210, Emerson) for five minutes.
To the 200 μL sonicated sample, 200 μL of an ice-cold solution of 12.5% trichloroacetic acid was added.

After incubation for 2 hours at 4°C, the sample was centrifuged at 10,000 g for 20 minutes. The
supernatant was neutralized with 45 µL of freshly prepared 1 M NaOH and then mixed with 195 µL of
50 µM aminocaproic acid (internal standard). Free amino acids were quantified using HPLC, with
mobile phase compositions and HPLC settings as in Rehlund et al. (2025). All standard free amino acids
were quantified, except cysteine and proline. HPLC data were pre-processed using Chromeleon 7.2.7
(Thermo Scientific, MA, USA).

2.9 Quantification of cereulide produced by strain Y48H10 using HPLC-MS/MS

Quantification of cereulide from strain Y48H10 was performed in two biological duplicates. Four
technical replicates were pooled together for each sampling time point. Cereulide analysis was
performed using ISO 18465 standard (Anonymous, 2017) method with few modifications. Briefly,
samples were diluted in 1:10 ratio with HPLC-grade acetonitrile (VWR Chemicals, Poland). Labelled
cereulide (^{13}C6-cereulide) was added as an internal standard in each sample at the concentration of 10
ng/ml. The cereulide was extracted by a rotating shaker (300 rpm) for 1 hour followed by resting for 30
minutes. The solution was decanted twice (first 4500 rcf for 15 minutes and second 12,000 rcf for 15
minutes). A 200 µL of the supernatant was transferred into HPLC vial for LC-MS/MS analysis.

Samples were randomized and analysed on an Elute UPLC system coupled to an Impact II QTOF-MS
instrument (Bruker Daltonik, Germany). Chromatographic separation was performed on a reverse-
phase Supelco Discovery C18 column (10 cm × 2.1 mm, 5 µm particle size) maintained at 36 °C. The
mobile phase consisted of an isocratic elution with 5% phase A (10 mM ammonium formate in water
containing 0.1% formic acid) and 95% phase B (acetonitrile with 0.1% formic acid). The flow rate was
set to 0.4 mL/min with an injection volume of 5 µL. Mass spectra were acquired in the range of m/z
1000–1300 at 12 Hz using electrospray ionization (ESI) in positive-ion mode. Prior to analysis, the
instrument was externally calibrated with 10 mM sodium formate in 50% isopropanol and tuned using
the ESI-L Low Concentration Tuning Mix (Agilent Technologies, CA, USA). ESI parameters were as
follows: end plate offset, –500 V; capillary voltage, 4500 V; nebulizer pressure, 2.2 bar; dry gas flow,
10 L/min; and drying temperature, 220 °C. MS settings included funnel 1 and 2 RF, 200 Vpp; hexapole,
50 Vpp; ion energy, 5 eV; collision energy, 5 eV; and pre-pulse storage, 5 µs. Peak areas of natural
cereulide and isotope-labelled cereulide were extracted using ion traces corresponding to their precursor
ions at m/z 1170–1171 and 1176–1177, respectively.

2.10 Data analysis

Isothermal microcalorimetric data were analysed from 1 to 24 hours after inoculation, as the first hour
was needed for thermal equilibration of the instrument. Baseline correction was applied using the heat
flow data collected before plate insertion. For 2 out of 96 experiments, an alternative baseline heat flow
value was selected to achieve a better fit after plate insertion at 50 minutes. Data from one replicate

experiment corresponding to strain *B. subtilis* PRO31 was excluded, due to negative heat flow values
at the beginning of the experiment, which was attributed by the instrument manufacturer to moisture
evaporation from the vial.

The heat curves (heat accumulated as a function of time) were fitted with the Gompertz model to
estimate microbial growth characteristics (Zwietering et al., 1990), based on two assumptions (1) that
cells remain in the medium, and (2) that heat flow data present themselves with one central peak or a
closely similar shape (e.g. peak with a shoulder) (Braissant et al., 2010). Data analysis of the isothermal
calorimetry was conducted using RStudio (R Core Team, 2023). Curve fitting and parameter estimation
were performed on each experiment using the grofit R package, based on Gompertz modelling, to
estimate the maximum heat production (mJ), the length of the calorimetric lag time (h), and the
maximum calorimetric growth rate (mW · h^{-1}) (Kahm et al., 2010). The lag time was corrected for the
one-hour pre-incubation period. Statistical tests were performed using the Gompertz parameters related
to species, with a significance level of 0.05. Group comparisons between *B. cereus* and *B. subtilis* and
between the interspecies clades in BHI medium were performed using unpaired T-tests, and
phylogenetic comparisons between *B. cereus* clades were performed using Kruskal-Wallis with Dunn's
post-hoc test. Comparison between heat curves obtained in BHI media and food matrices of selected *B.*
cereus (Y48H10, SDA GR177, MC67) and *B. subtilis* strains (PRO99, DSM10, DSM21393) were
conducted using paired t-test with Bonferroni correction.

A two-way analysis of variance (ANOVA) was performed to examine the effects of *Bacillus* clade (*B.*
cereus and *B. subtilis*), food (milk, oat, pea, and oat-pea), and their interaction on the square-root
transformed Gompertz parameters. A separate two-way ANOVA was performed to test the influence
of all matrices (foods and BHI medium), *Bacillus* clade, and their interaction on square-root transformed
Gompertz parameters. All ANOVA models were fitted using the lmerTest package, which extends lme4
by providing p-values for mixed effects. Post-hoc analyses were conducted using estimated marginal
means (EMM) via the emmeans package. For the metabolomics data, a three-way ANOVA was
performed to evaluate the influence of *Bacillus* clade, metabolic time-point (peak vs. end), and food
matrix, as well as their interactions, on the content of individual sugars, individual organic acids, and
total free amino acids. Biological replicates were included as a random effect using lme4, and
significance testing was carried out with lmerTest. Post-hoc analyses were conducted using EMM.
Additionally, a two-way ANOVA was conducted to investigate the effects of fermentation (non-
fermented vs. fermented) and matrix on total sugars. Tukey post-hoc comparisons were performed
between non-fermented and fermented samples within each matrix (oat, pea, and oat–pea). Non-
detected compounds were assigned a value equal to half the lowest detected concentration to enable
statistical comparisons. Principal component analysis (PCA) was applied to investigate potential

interactions between the metabolites of *Bacillus* spp. in food matrices using RStudio (R Core Team,
2023), and a biplot was created using the ggplot2 package.

The quantification of cereulide produced by *B. cereus* strain Y48H10 in food matrices was based on
mass spectrometric peak area ratios derived from baseline-corrected chromatograms, using a signal-to-
noise ratio (SNR) threshold of 3. The limit of detection (LOD) and limit of quantification (LOQ) were
determined from the standard deviation of the noise in 0.1 ng/mL samples, which represented the lowest
concentration at which cereulide was not detectable. Inspection and processing of obtained cereulide
data were performed using MATLAB 2023b.

3. Results

3.1 Taxonomy and phylogeny

A maximum-likelihood phylogenetic tree was constructed using the UBCG pipeline (Na et al., 2018)
from the whole-genome sequences of the *B. cereus* and *B. subtilis* strains. Species names were drawn
from the best taxonomic (ANIm) matches rather than the previously recognized taxonomy of the strains
(Table 1). For strains BC59 and LWL3, previously classified as *B.* cereus, ANIm values exceeded the
species threshold (>96%) for both *Bacillus wiedmannii* and *Bacillus pretiosus*. However, we adopted
B. wiedmannii as the taxonomic designation, in line with (Mozhaitseva et al., 2025), who reported
genomic similarity between these species and grouped them within a shared cluster. Moreover, *B.*
pretiosus remains an effectively published but not validly recognized name under current bacterial
nomenclature rules (Oren et al., 2023; Robas Mora et al., 2023), further supporting the use of *B.*
wiedmannii. Five different clades were identified (Figure 1), differentiating the *B. cereus* group strains
into (Clade A) *B. paranthracis, B. pacificus,* and *B. tropicus*, (Clade B) *B. wiedmannii/B. pretiosus*, and
(Clade C) *B. mycoides*, and distinguishing the *B. subtilis* group into (Clade D) PRO31, PRO35, PRO64,
PRO99, and (Clade E) DSM13019, DSM21393, PRO115, and DSM10. In short, we observed a diverse
range of *B. cereus* (encompassing five different genomic subspecies) and *B. subtilis* isolates (two
distinct clades) to facilitate comparisons between phylogeny and microcalorimetric analysis of
metabolic activity.

Figure 1: Rooted maximum likelihood phylogenetic tree of Bacillus cereus *group strains (Clades A, B,*

and C) and Bacillus subtilis strains (Clades D and E) using the up-to-date bacterial core gene (UBCG)

pipeline, drawn by MEGA11. The names of the species were drawn from the best taxonomic (ANIm)

matches, as displayed in Table 1. Halobacillus halophilus *DSM2266 was used to root the tree.*

Percentage bootstrap values are indicated at branching points, and the scale bar represents 0.01

substitutions per site.

3.2 Metabolic activity of *Bacillus* cultivated in BHI and food matrices

To investigate the metabolic activity of *Bacillus* strains under controlled conditions, we monitored the

metabolic heat production profiles of 16 *B. cereus* strains and 8 *B. subtilis* strains. Cultivation was first

carried out in BHI medium at 30°C for 20 hours, until the metabolic activity reached a steady-state

plateau (Figure 2B). During 20 hours of cultivation, *B. cereus* and *B. subtilis* exhibited a typical

274 sigmoidal pattern in the heat production curves (Figure 2A). The heat flow profiles of both *B. subtilis*
275 and *B. cereus* exhibit one main peak followed by a second shoulder peak (Figure 2B).

2A
Accumulated heat over time

2B
Heat flow over time

Group — *cereus* — *subtilis*

Figure 2: Metabolic heat characteristics of 16 B. cereus *(red) and 8* B. subtilis *(blue) strains cultivated*
in BHI medium, based on four biological replicates divided between two different experiments (2A)
Mean values (curve) and standard deviation (error bars) of accumulated heat production over time
(2B) Mean values (curve) and standard deviation (error bars) of heat flow over time of 16 B. cereus
(red) and 8 B. subtilis *(blue) strains.*

To gain a better understanding of the metabolic heat profiles, fitted parameters from the Gompertz
model, lag time (h), the maximum calorimetric growth rate (mJ*h^{-1}), and total heat (mJ), were plotted
for each *Bacillus* group (*cereus* vs. *subtilis*) (Figure 3A) and the phylogenetic clades within *B. cereus*
and *B. subtilis* groups (A, B vs. C; D vs. E) (Figure 3B). Between the *B. subtilis* and *B. cereus* groups,
significant differences were observed across all parameters. *B. subtilis* released a significantly higher

amount of total heat (2732 mJ ± 462) than *B. cereus* (1941 mJ ± 566) ($p < 10^{-10}$). *B. subtilis* displayed

a higher maximum metabolic activity (378 h^{-1} ± 140) than *B. cereus* (240 h^{-1} ± 106) ($p < 10^{-5}$). Finally,

a longer lag time was observed for *B. subtilis* (7.9 hours ± 1.0) than for *B. cereus* (4.3 hours ± 0.8) ($p <$

10^{-16}).

In addition, differences in metabolic activity were observed within the *B. subtilis* and *B. cereus* groups.

Between the *B. cereus* clades, *B. mycoides* strains from clade C produced a significantly lower amount

of total heat (971 mJ ± 292) than *B. paranthracis*, *B. pacificus,* and *B. tropicus* strains from clade A

(2119 mJ ± 464, $p < 10^{-5}$) and *B. wiedmannii* strains from clade B (1839mJ ± 212, p = 0.01). In addition,

the maximum calorimetric growth rate differed significantly between Clade A and B ($p < 10^{-5}$) and

between Clade A and C (p <10-5). Between the *B. subtilis* clades, clade D strains exhibited a longer lag

time (8.6 hours ± 0.8) than Clade E (7.2 hours ± 0.7) ($p < 10^{-5}$).

*Figure 3: Differences in heat flow characteristics, including calorimetric lag time (h), maximum calorimetric growth rate (mJ*h⁻¹), and total heat (mJ). Calorimetric lag time was corrected for 1 h pre-incubation period. (3A): Differences in heat flow characteristics between the subtilis and cereus groups. (3B): Differences in heat flow characteristics between the phylogenetic clades as defined in Figure 1, between Clade A (*B. paranthracis, B. pacificus *and* B. tropicus*), B (*B. wiedmannii*), C (*B. mycoides*), D (*B. subtilis PRO31, PRO35, PRO64, PRO99) and E (*B. subtilis DSM13019, DSM21393, PRO115 and DSM10)*

In addition to BHI, four strains of *B. cereus* (Y48H10, SDA GR177, MC67) and three *B. subtilis*
(PRO99, DSM10, DSM21393) were cultured in food matrices (milk, oat, pea, and oat-pea) at 30°C.
Their metabolic heat activity was monitored for 24 hour periods. The pea and oat-pea matrices were
autoclaved prior to inoculation. In contrast, the oat was not autoclaved based on preliminary findings:
B. cereus did not metabolize autoclaved oat, but autoclaving was necessary to solubilize the pea
hydrolysate. Gompertz parameters were subsequently derived from the calorimetric data (Table 2).

Table 2: Gompertz calorimetric growth parameters (mean (± sd)) of *B. cereus* Y48H10 and *B. subtilis*
DSM21393 cultivation in milk, oat, pea, and oat-pea matrices.

Species	Food	Calorimetric lag time [hours]	Max calorimetric growth rate [mJ/h]	Max accumulated heat [mJ]
B. cereus	Milk	7.4 (± 0.5)	150 (± 8)	2040 (± 120)
	Oat	14.5 (±12.8)	184 (± 97)	1696 (± 865)
	Pea	7.7 (± 0.3)	216 (± 65)	1663 (± 714)
	Oat-pea	9.8 (±1.7)	407 (± 103)	1880 (± 665)
B. subtilis	Milk	12.4 (± 2.7)	109 (± 30)	2057 (± 334)
	Oat	11.6 (± 3.1)	211 (± 43)	2452 (± 212)
	Pea	10.4 (±1.3)	275 (± 40)	2065 (± 301)
	Oat-pea	10.9 (± 0.4)	419 (± 28)	2403 (± 232)

B. subtilis exhibited a more consistent calorimetric lag time across food matrices, with a longer lag time
in milk (12.4 hours ± 2.7) compared to *B. cereus* (7.35 hours ± 0.5). Maximum calorimetric growth
rates varied significantly by substrate and were higher in the combined oat-pea matrix compared to oat
($p = 0.002$, EMM), pea ($p = 0.01$, EMM), and milk ($p < 10^{-4}$, EMM). Across all food matrices, *B. subtilis*
produced more metabolic heat than *B. cereus* ($p = 0.02$, EMM), consistent with observations on BHI
medium. When comparing *Bacillus* metabolism between BHI and the food matrices, the calorimetric
lag time was significantly lower in BHI than in oat ($p = 0.02$, EMM), while no significant differences
were observed between the other matrices.

3.3 Metabolic profiling of *Bacillus* growth in oat, pea, and oat-pea matrices

Targeted metabolomic profiling was performed to assess how matrix composition influences *Bacillus*
metabolic activity, with particular focus on the enhanced growth observed in the combined oat-pea
matrix. One *B. subtilis* (DSM21393) and one *B. cereus* strain (Y48H10) were selected for cultivation
on the food matrices and sampled at two time-points: at peak metabolic activity (8 hours for *B. cereus,*
11 hours for *B. subtilis*) and at minimum metabolic activity (24 hours for both).

Sugars, organic acids, and free amino acids were quantified in all samples. To interpret the multifactorial data generated from targeted metabolomic profiling (simple sugars, organic acids, and free amino acids), principal component analysis (PCA) was employed to visualize trends in metabolite synthesis and consumption by *B. subtilis* and *B. cereus* cultured in plant-based matrices over 24 hours. PCA revealed that matrix composition was the main driver of sample separation (Figure 4). PC1 (41% variance) was associated with free amino acids, galactose, and fructose. In comparison, PC2 (25% variance) separated the pea matrix from the oat and oat-pea, reflecting lower sugar levels (sucrose, glucose) and higher citrate content.

Figure 4: PCA biplot of Bacillus cereus *Y48H10 (red) and* Bacillus subtilis *DSM21393 (blue) metabolizing free amino acids, simple sugars, and organic acids in oat (circles), pea (squares), and oat-pea (triangles). Ellipses represent 95% of the population for each food matrix.*

To gain insight into the factors driving PCA separation, we analysed temporal changes in sugars, organic acids, and free amino acids. Matrix composition strongly influenced sugar and organic acid profiles. Total sugar content, as quantified by HPLC-RID, was significantly higher in the uninoculated oat (49.4 ± 1.8 g/L) and oat-pea (52.6 ± 3.6 g/L, p = 0.001, Tukey, Figure 5A) matrices compared to pea

(0.74 ± 0.00 g/L, p = 0.001, Tukey, Figure 5). *Bacillus* fermentation significantly reduced the total

amount of sugars in the oat-pea matrix (p = 0.01, two-way ANOVA, EMM). Citrate was uniquely

present in the pea matrix and absent from both oat and oat-pea (Figure 5B), contributing to its distinct

separation in PCA (Figure 4). Cultivation of *B. subtilis* and *B. cereus* led to an increase in total organic

acids from peak to end metabolic phase (p = 10^{-5}, three-way ANOVA). The oat-pea matrix showed the

most pronounced increase in organic acids, with both species sharply increasing lactic acid levels (p =

10^{-4}, EMM, Figure 5B). In contrast, the changes in organic acid levels in oat and pea were more modest

(p > 0.05, EMM). During *B. subtilis* growth on pea, citrate was consumed, and acetic acid was produced.

B. cereus produced more acetic acid than *B. subtilis* after 24 hours (p = 10^{-4}, EMM).

(A) Sugars

Sugar ■ Galactose ■ Arabinose ■ Fructose ■ Glucose ■ Sucrose

(B) Organic acids

Organic acid ☐ Citric acid ■ Acetic acid ■ Lactic acid

Figure 5: Metabolic profiling of Bacillus cereus *Y48H10 and* Bacillus subtilis *DSM21393 cultivated in oat, pea, and oat-pea matrices. Error bars are based on two biological replicates, each of which consists of four pooled technical replicates (A) sugar composition, (B) organic acid composition.*

Free amino acids were observed in all uninoculated matrices, with the highest levels detected in the

(non-autoclaved) oat drink (Figure 6). *B. cereus* and *B. subtilis* had distinct effects on the free amino

acid profiles depending on the matrix. The most pronounced effects were observed during *B. subtilis*

growth on oat and *B. cereus* growth on pea, both increasing free amino acid levels, as also reflected in

the PCA (Figure 4). These effects differed in the types of free amino acids released and the timing of

their accumulation. For *B. cereus* in pea, total free amino acids increased from 59 µmol/g ± 1 (8 hours)

to 219 µmol/g ± 54 (24 hours) (p = 0.003, EMM), primarily due to accumulation of Gly, Phe, Ala, Ser,

Leu and Tyr. In contrast, *B. subtilis* displayed higher total free amino acid levels in oat than in pea (p =

10^{-4}, EMM), peaking at 227 µmol/g ± 38 at maximum metabolic activity after 11 hours, with elevated

levels of Asp, Gly and Ala, and a reduction in Asn (Figure 4). In the oat-pea matrix, *Bacillus* did not

significantly alter total free amino acid levels over time (p = 1, EMM).

Figure 6: Free amino acid profile of Bacillus cereus *Y48H10 and* Bacillus subtilis *DSM21393*
cultivated in oat, pea, and oat-pea matrices. Error bars are based on two biological replicates, each
consisting of four pooled technical replicates.

Cereulide was exclusively quantified in the uninoculated matrices and in the *B. cereus* samples, since it was assumed that *B. subtilis* is incapable of producing cereulide. Cereulide was below the detection limit (0.1 ng/mL) in the pea matrix. In contrast, concentrations of 3.0 and 2.9 ng/mL were detected in the oat-pea matrix, while the oat drink exhibited substantially higher concentrations of 46 and 48 ng/mL in the two biological replicates.

4. Discussion

To examine the metabolic heat production of *B. cereus* and *B. subtilis*, a total of 24 *Bacillus* strains were selected to represent diversity in origin, pathogenic potential, and genetic background. The pathogenic potential - reflected in the ability to produce enterotoxins and/or emetic toxins (Table 1) - was partially aligned with phylogenetic clustering, suggesting a link between genetic lineage and toxin production capacity. All emetic strains showed the closest ANIm resemblance to *B. paranthracis*, except for the emetic psychrotolerant strain, *B. mycoides* M67. Enterotoxin-producing strains were identified in all three clades of the *B. cereus* group and could not be distinguished by phylogeny or metabolic profiles. The classification of the strains within the *B. cereus* group (Table 1) reflects some of the recent changes in the nomenclature of *Bacillus cereus* taxonomy. *B. wiedmannii* was proposed as a new species in 2016 (Miller et al., 2016). *B. paranthracis, B. pacificus, and B. tropicus* were proposed in 2017 (Liu et al., 2017), and *B. weihenstephanensis* merged with *B. mycoides* in 2018 (Liu et al., 2018).

Clade-specific growth dynamics were observed in all Gompertz parameters during cultivation in BHI at 30°C. *B. subtilis* exhibited a higher model-estimated total heat output (2745 mJ) than the observed value (2473 mJ), indicating sustained metabolic activity beyond the 20-hour endpoint. In contrast, *B. cereus* reached a metabolic plateau earlier, with modelled (1941 mJ) and measured (2000 mJ) values closely aligned. Both species displayed a shoulder peak following the main heat flow peak (Figure 2B), likely reflecting a shift in substrate utilization. Notably, *B. cereus* strains had a significantly shorter lag phase (4.3 h ± 0.8) compared to *B. subtilis* (7.9 h ± 1.0), with no overlap observed between clades.

The observed differences in metabolic growth profiles may reflect the distinct ecological strategies of each species in a highly nutrient-rich context, such as BHI medium. It is plausible that *B. subtilis* exhibited more sustained metabolic activity and higher heat output in BHI due to a broader capacity for peptide and carbohydrate catabolism, potentially supported by a diverse enzymatic repertoire. In contrast, the rapid onset and early saturation of metabolic activity observed in *B. cereus* may reflect a faster environmental adaptation coupled with a less energetically intensive growth strategy in nutrient-

rich conditions. Phylogenetic variation within each species further influenced metabolic heat production

in the BHI medium. *B. mycoides* (clade B) strains, for instance, released significantly less heat than

other *B. cereus* clades. This behaviour is consistent with the known psychrotolerant phenotype of one

of its members (MC67) and reduced growth at mesophilic temperatures (Guinebretière et al., 2008).

The influence of substrate composition on *Bacillus* growth dynamics became more apparent when

strains were cultivated in food matrices. To evaluate whether clade-specific metabolic traits persisted

across different food matrices, three strains of each of *Bacillus cereus* and *Bacillus subtilis* were

cultivated in milk, oat, pea, and a combined oat-pea matrix. As observed in BHI medium, *B. subtilis*

generally exhibited longer lag times than *B. cereus*, most notably in milk (12.4 hours vs. 7.4 hours).

However, species-level distinctions were less pronounced in the food matrices compared to BHI,

suggesting that substrate composition exerts a stronger influence on physiological responses than

phylogenetic background. The oat-pea matrix stood out as particularly supportive of *Bacillus* activity,

with the highest maximum calorimetric growth rate observed among all tested substrates. This suggests

that the combination of oat and pea provides a more favourable nutrient environment than either

substrate alone.

To elucidate the biochemical mechanisms underlying matrix-dependent growth dynamics, we analysed

sugar, organic acid, and free amino acid profiles during cultivation of one representative strain of each

B. cereus and *B. subtilis* in oat, pea, and oat-pea matrices. Despite the controlled experimental setup,

direct comparison of metabolic profiles across matrices was complicated by differences in thermal

treatment. The oat drink was not autoclaved, as preliminary experiments showed that heat treatment

inhibited *B. cereus* growth, likely due to degradation of heat-sensitive nutrients such as free amino

acids. In contrast, autoclaving was necessary to solubilize the pea hydrolysate in the pea and oat-pea

matrices. This discrepancy introduces a confounding variable, as thermal processing can trigger

Maillard reactions that reduce the availability of amino acids and simple sugars and alter the redox

balance (Lund & Ray, 2017). These effects may explain the elevated free amino acid levels observed

in the non-autoclaved oat matrix compared to the autoclaved pea-based matrices.

Despite these limitations, clear trends emerged in how matrix composition shaped metabolic activity.

The pea matrix, characterised by its low sugar content and the unique presence of citrate, showed clear

separation from oat and oat-pea in PCA. During cultivation on pea, both *B. cereus* and *B. subtilis*

produced and consumed organic acids, but their metabolic patterns diverged. *B. cereus* produced more

acetic acid than *B. subtilis* at the end of metabolism (24 hours). While citrate levels did not significantly

change during *B. cereus* metabolism in pea. *B. subtilis* reduced citrate levels, consistent with adaptive

citrate fermentation under nutrient stress or acidic conditions (Schilling et al., 2007). These species-

specific responses suggest that sugar limitation in pea may drive differential organic acid turnover. The

oat-pea matrix, which combines carbohydrate-rich and peptide-rich components, supported the most

pronounced accumulation of organic acids, including lactic acid and acetic acid, across both *Bacillus*

species. This suggests that the combined matrix could promote more efficient carbohydrate metabolism,

likely due to synergistic interactions between sugars and peptides, and may account for the higher

calorimetric growth observed in oat-pea.

Distinct species-specific strategies also emerged in the dynamics of free amino acids, revealing how *B.*

cereus and *B. subtilis* differentially interact with their nutrient environments. *B. subtilis* showed

transient free amino acid release in oat, consistent with its capacity to metabolize complex

carbohydrates and synthesize amino acids under nutrient-rich conditions. In contrast, *B. cereus*

accumulated free amino acids in the pea matrix, likely reflecting proteolytic activity and amino acid

catabolism under nutrient or oxygen stress. The limited changes of free amino acid levels in the oat-pea

matrix, despite high organic acid production, may indicate a metabolic shift away from amino acid

turnover toward carbohydrate fermentation. These observations align with previous findings that *B.*

cereus preferentially utilizes proteinaceous substrates under limited oxygen conditions (Chang et al.,

2021).

Cereulide, the heat- and acid-stable emetic toxin produced by *B. cereus*, was quantified across the food

matrices. The toxin was detected in both oat and oat-pea matrices, but not in pea, suggesting that matrix

composition influences toxin production—likely through differences in nutrient availability.

Interestingly, although *B. cereus* Y48H10 exhibited the highest heat production in the oat-pea matrix,

cereulide levels were lower than in oat alone. Ellouze et al. (2021), using *B. cereus* F4810/72, similarly

observed faster cereulide formation in a cereal-based matrix (rice) compared to a vegetable matrix

(vegetables purée) despite slower bacterial growth in the cereal-based matrix, supporting the notion that

cereulide production is matrix-dependent. However, in contrast to their findings on the vegetable

matrix, our study detected no quantifiable cereulide in pea at 30°C within 24 hours. High sugar and acid

levels in oats may enhance ATP availability, thereby favouring cereulide production (Marxen et al.,

2015). Previous studies have shown that cereulide expression is highly sensitive to environmental

factors, including pH, nutrient profile, and redox state (Dommel et al., 2010; Ehling-Schulz et al., 2015).

The absence of cereulide in pea and lower cereulide levels in oat-pea, despite microbial activity, may

reflect an indirect inhibitory effect of matrix composition on regulatory networks involved in cereulide

biosynthesis.

Whether components in pea actively suppress cereulide biosynthesis or limit precursor availability

remains unresolved and warrants further investigation. While cereulide is typically associated with

starchy foods (Messelhäusser et al., 2014), outbreaks have been reported in legume-based products such

as fermented soybeans (Inatsu et al., 2020) and soaked navy beans (Nicholls et al., 2016), indicating

that low-carbohydrate environments can support toxin synthesis under permissive conditions. Thorsen

et al. (2011) found no emetic strains in a survey of African bread snacks made from legumes, whereas

Kyrylenko et al. (2023) reported that 9% of *B. cereus* isolates from plant-based dairy alternative
ingredients carried cereulide-producing genes. These findings highlight the variability in risk across
substrates and reinforce the importance of matrix-specific evaluations when assessing cereulide-
associated hazards.

Metabolic heat profiling provided complementary insights into *Bacillus* physiology, distinguishing
species- and clade-level behaviour under controlled conditions. Our findings reveal that *B. subtilis*
consistently produces more metabolic heat than *B. cereus*. Differences in lag time and growth rate were
matrix-dependent, highlighting the limitations of extrapolating from defined media, such as BHI.
Although calorimetry alone cannot resolve underlying metabolic pathways, its integration with
metabolomic profiling offers a robust framework for assessing microbial activity, adaptation, and
potential safety risks in complex food systems. While the metabolic capacities of *B. subtilis* and *B.*
cereus are well-characterized in defined media, their proteomic responses in food matrices remain
underexplored (Wicke et al., 2023). Future studies integrating calorimetry, metabolomics, and
proteomics could help uncover matrix-specific regulatory mechanisms that govern spoilage activity and
beneficial fermentation traits in *B. subtilis*, as well as toxin production and contamination risks
associated with *B. cereus*. Such approaches are essential for understanding *Bacillus* behaviour in diverse
food environments.

5. Conclusion

This study demonstrates that *Bacillus cereus* and *Bacillus subtilis* exhibit distinct metabolic behaviour
across food-relevant matrices. In BHI at 30°C, *B. subtilis* showed longer lag times and higher total heat
production than *B. cereus*. However, in food matrices, substrate composition had a more pronounced
effect on metabolic activity than species identity, although *B. subtilis* consistently generated more heat
overall.

The combined oat-pea matrix supported the highest calorimetric growth rates, likely driven by enhanced
carbohydrate metabolism, as evidenced by reduced sugar levels and increased organic acids. In contrast,
the pea matrix- characterised by low sugar and the unique presence of citrate, triggered post-metabolic
consumption of citrate by *B. subtilis*. Free amino acid profiles reflected matrix-specific strategies, *B.*
subtilis released free amino acids in oat, while *B. cereus* predominantly did so in pea. Interestingly,
despite high calorimetric activity, the oat-pea matrix showed limited change in free amino acid levels,
suggesting a shift toward carbohydrate-driven metabolism.

Cereulide production by *B. cereus* was strongly matrix-dependent, with high levels detected in oat,
moderate levels in oat-pea, and none in pea, highlighting the role of substrate composition in toxin
synthesis. Together, these findings underscore the need for matrix-specific microbial risk assessment

tools. They also demonstrate that integration of isothermal microcalorimetry with metabolomics
provides a more nuanced understanding of microbial behaviour in complex food environments. Similar
heat profiles can mask divergent metabolic strategies; thus, this combined approach offers a valuable
tool for evaluating spoilage and contamination risks, particularly in the development of safer plant-
based food products.

Acknowledgment

514 The authors wish to acknowledge the Villum Foundation (HLR:34434) and the joint alliance grant
515 between the Danish Dairy Research Foundation, Arla Foods amba, Novonesis, and the University of
516 Copenhagen (SK: SaFeMix) for financial support.

Conflict of interest

518 The authors declare that they have no known competing financial interests or other conflicts of
519 interest.

Author contribution

Emilie V. Skriver: Conceptualization, Methodology, Formal analysis, Investigation, Visualization,
Writing – original draft, Writing – review and editing; Tessa S. Canoy: Conceptualization,
Methodology, Formal analysis, Investigation, Visualization, Writing – original draft, Writing – review
and editing; Yuandong Sha: Formal analysis, Investigation, Visualization; Morten A. Rasmussen:
Software, Formal analysis; Bekzod Khakimov: Methodology, Formal analysis, Supervision; Susanne
Knøchel: Writing – review and editing, Resources, Supervision, Project administration, Funding
acquisition; Henriette L. Røder; Conceptualization, Methodology, Resources, Writing – review and
editing, Supervision, Project administration, Funding acquisition.

References

Anonymous. (2017). ISO18465: *Microbiology of the food chain-Quantitative determination of emetic*
toxin (cereulide) using LC-MS/MS COPYRIGHT PROTECTED DOCUMENT. www.iso.org
Beilharz, K., Kragh, K. N., Fritz, B., Kirkegaard, J. B., Tolker-Nielsen, T., Bjarnsholt, T., &
Lichtenberg, M. (2023). Protocol to assess metabolic activity of Pseudomonas aeruginosa by
measuring heat flow using isothermal calorimetry. *STAR Protocols*, *4*(2).
https://doi.org/10.1016/j.xpro.2023.102269
Braissant, O., Wirz, D., Göpfert, B., & Daniels, A. U. (2010). Use of isothermal microcalorimetry to
monitor microbial activities [Article]. *FEMS Microbiology Letters*, *303*(1), 1–8.
https://doi.org/10.1111/j.1574-6968.2009.01819.x
Carroll, L. M., Cheng, R. A., Wiedmann, M., & Kovac, J. (2022). Keeping up with the Bacillus cereus
group: taxonomy through the genomics era and beyond. In *Critical Reviews in Food Science and*

542 *Nutrition* (Vol. 62, Issue 28, pp. 7677–7702). Taylor and Francis Ltd.
543 https://doi.org/10.1080/10408398.2021.1916735

544 Chang, J. D., Vaughan, E. E., Liu, C. G., Jelinski, J. W., Terwilliger, A. L., & Maresso, A. W. (2021).
545 Metabolic profiling reveals nutrient preferences during carbon utilization in Bacillus species
546 [Article]. *Scientific Reports*, *11*(1), 23917–23917. https://doi.org/10.1038/s41598-021-03420-7

547 Cichos, K. H., Ruark, R. J., & Ghanem, E. S. (2023). Isothermal microcalorimetry improves accuracy
548 and time to bacterial detection of periprosthetic joint infection after total joint arthroplasty
549 [Article]. *Journal of Clinical Microbiology*, *61*(12), e0089323–e0089323.
550 https://doi.org/10.1128/jcm.00893-23

551 Dommel, M. K., Frenzel, E., Strasser, B., Bloechinger, C., Scherer, S., & Ehling-Schulz, M. (2010).
552 Identification of the Main Promoter Directing Cereulide Biosynthesis in Emetic Bacillus cereus
553 and Its Application for Real-Time Monitoring of ces Gene Expression in Foods [Article].
554 *Applied and Environmental Microbiology*, *76*(4), 1232–1240.
555 https://doi.org/10.1128/AEM.02317-09

556 Dong, H., Zhu, H., He, H., & Yi, C. (2023). Identification and evaluation of spoilage potential of four
557 Bacillus strains isolated from slimy rice noodles [Article]. *Food Microbiology*, *110*, 104160.
558 https://doi.org/10.1016/j.fm.2022.104160

559 Earl, A. M., Losick, R., & Kolter, R. (2008). Ecology and genomics of Bacillus subtilis [Article].
560 *Trends in Microbiology (Regular Ed.)*, *16*(6), 269–275.
561 https://doi.org/10.1016/j.tim.2008.03.004

562 EFSA. (2022). The European Union One Health 2021 Zoonoses Report. *EFSA Journal*, *20*(12).
563 https://doi.org/10.2903/j.efsa.2022.7666

564 Ehling-Schulz, M., Frenzel, E., & Gohar, M. (2015). Food-bacteria interplay: Pathometabolism of
565 emetic Bacillus cereus. *Frontiers in Microbiology*, *6*(JUN).
566 https://doi.org/10.3389/fmicb.2015.00704

567 Ehling-Schulz, M., Fricker, M., & Scherer, S. (2004). Bacillus cereus, the causative agent of an
568 emetic type of food-borne illness [Article]. *Molecular Nutrition & Food Research*, *48*(7), 479–
569 487. https://doi.org/10.1002/mnfr.200400055

570 Ellouze, M., Buss Da Silva, N., Rouzeau-Szynalski, K., Coisne, L., Cantergiani, F., & Baranyi, J.
571 (2021). Modeling Bacillus cereus Growth and Cereulide Formation in Cereal-, Dairy-, Meat-,
572 Vegetable-Based Food and Culture Medium. *Frontiers in Microbiology*, *12*.
573 https://doi.org/10.3389/fmicb.2021.639546

574 Guinebretière, M. H., Thompson, F. L., Sorokin, A., Normand, P., Dawyndt, P., Ehling-Schulz, M.,
575 Svensson, B., Sanchis, V., Nguyen-The, C., Heyndrickx, M., & De Vos, P. (2008). Ecological
576 diversification in the Bacillus cereus Group. *Environmental Microbiology*, *10*(4), 851–865.
577 https://doi.org/10.1111/j.1462-2920.2007.01495.x

578 Inatsu, Y., Chotiko, A., & Nanchaipattana, C. (2020). Contaminated Bacillus cereus in Lao and Thai
579 fermented soybean "Tua Nao." *Japan Agricultural Research Quarterly*, *54*(1), 47–51.
580 https://doi.org/10.6090/jarq.54.47

581 Kahm, M., Hasenbrink, G., Lichtenberg-Frate, H., Ludwig, J., & Kschischo, M. (2010). grofit: Fitting
582 Biological Growth Curves with R [Article]. *Journal of Statistical Software*, *33*(7), 1–21.
583 https://doi.org/10.18637/jss.v033.i07

584 Kpikpi, E. N., Thorsen, L., Glover, R., Dzogbefia, V. P., & Jespersen, L. (2014). Identification of
585 Bacillus species occurring in Kantong, an acid fermented seed condiment produced in Ghana
586 [Article]. *International Journal of Food Microbiology*, *180*, 1–6.
587 https://doi.org/10.1016/j.ijfoodmicro.2014.03.028

588 Kyrylenko, A., Eijlander, R. T., Alliney, G., de Bos, E. L. van, & Wells-Bennik, M. H. J. (2023).
589 Levels and types of microbial contaminants in different plant-based ingredients used in dairy

alternatives. *International Journal of Food Microbiology*, *407*.
https://doi.org/10.1016/j.ijfoodmicro.2023.110392

Li, Z., Zheng, M., Zheng, J., & Gänzle, M. G. (2023). Bacillus species in food fermentations: an
underappreciated group of organisms for safe use in food fermentations [Article]. *Current*
Opinion in Food Science, *50*, 101007. https://doi.org/10.1016/j.cofs.2023.101007

Lichtenberg, M., Kvich, L., Larsen, S. L. B., Jakobsen, T. H., & Bjarnsholt, T. (2022). Inoculum
Concentration Influences Pseudomonas aeruginosa Phenotype and Biofilm Architecture
[Article]. *Microbiology Spectrum*, *10*(6). https://doi.org/10.1128/spectrum.03131-22

Liu, Y., Du, J., Lai, Q., Zeng, R., Ye, D., Xu, J., & Shao, Z. (2017). Proposal of nine novel species of
the Bacillus cereus group [Article]. *International Journal of Systematic and Evolutionary*
Microbiology, *67*(8), 2499–2508. https://doi.org/10.1099/ijsem.0.001821

Liu, Y., Lai, Q., & Shao, Z. (2018). Genome analysis-based reclassification of Bacillus
weihenstephanensis as a later heterotypic synonym of Bacillus mycoides [Article]. *International*
Journal of Systematic and Evolutionary Microbiology, *68*(1), 106–112.
https://doi.org/10.1099/ijsem.0.002466

Lund, M. N., & Ray, C. A. (2017). Control of Maillard Reactions in Foods: Strategies and Chemical
Mechanisms [Article]. *Journal of Agricultural and Food Chemistry*, *65*(23), 4537–4552.
https://doi.org/10.1021/acs.jafc.7b00882

Marxen, S., Stark, T. D., Frenzel, E., Rütschle, A., Lücking, G., Pürstinger, G., Pohl, E. E., Scherer,
S., Ehling-Schulz, M., & Hofmann, T. (2015). Chemodiversity of cereulide, the emetic toxin of
Bacillus cereus. *Analytical and Bioanalytical Chemistry*, *407*(9), 2439–2453.
https://doi.org/10.1007/s00216-015-8511-y

Messelhäusser, U., Frenzel, E., Blöchinger, C., Zucker, R., Kämpf, P., & Ehling-Schulz, M. (2014).
Emetic Bacillus cereus are more volatile than thought: Recent foodborne outbreaks and
prevalence studies in bavaria (2007-2013). *BioMed Research International*, *2014*.
https://doi.org/10.1155/2014/465603

Miller, R. A., Beno, S. M., Kent, D. J., Carroll, L. M., Martin, N. H., Boor, K. J., & Kovac, J. (2016).
Bacillus wiedmannii sp nov., a psychrotolerant and cytotoxic Bacillus cereus group species
isolated from dairy foods and dairy environments [Article]. *International Journal of Systematic*
and Evolutionary Microbiology, *66*(11), 4744–4753. https://doi.org/10.1099/ijsem.0.001421

Mozhaitseva, K., Pairaud, S., Firmesse, O., & Bonis, M. (2025). Population structure of Bacillus
cereus sensu lato associated with foodborne outbreaks in France between 2004 and 2023
[Document]. *BioRxiv*. https://doi.org/10.1101/2025.03.24.644875

Na, S.-I., Kim, Y. O., Yoon, S.-H., Ha, S., Baek, I., & Chun, J. (2018). UBCG: Up-to-date bacterial
core gene set and pipeline for phylogenomic tree reconstruction [Article]. *The Journal of*
Microbiology, *56*(4), 280–285. https://doi.org/10.1007/s12275-018-8014-6

Nicholls, M., Purcell, B., Willis, C., Amar, C. F. L., Kanagarajah, S., Chamberlain, D., Woolridge, D.,
Morgan, J., McLauchlin, J., Grant, K. A., Harvey-Vince, L., Padfield, M., Mearkle, R., & Chow,
J. Y. (2016). Investigation of an outbreak of vomiting in nurseries in South East England, May
2012 [Article]. *Epidemiology and Infection*, *144*(3), 582–590.
https://doi.org/10.1017/S0950268815001491

Oren, A., Arahal, D. R., Göker, M., Moore, E. R. B., Rossello-Mora, R., & Sutcliffe, I. C. (2023).
International Code of Nomenclature of Prokaryotes. Prokaryotic Code (2022 Revision) [Article].
International Journal of Systematic and Evolutionary Microbiology, *73*(5a).
https://doi.org/10.1099/ijsem.0.005585

Owusu-Kwarteng, J., Parkouda, C., Adewumi, G. A., Ouoba, L. I. I., & Jespersen, L. (2022).
Technologically relevant Bacillus species and microbial safety of West African traditional

alkaline fermented seed condiments [Article]. *Critical Reviews in Food Science and Nutrition*,
62(4), 871–888. https://doi.org/10.1080/10408398.2020.1830026

Parkouda, C., Thorsen, L., Compaoré, C. S., Nielsen, D. S., Tano-Debrah, K., Jensen, J. S., Diawara,
B., & Jakobsen, M. (2010). Microorganisms associated with Maari, a Baobab seed fermented
product [Article]. *International Journal of Food Microbiology*, *142*(3), 292–301.
https://doi.org/10.1016/j.ijfoodmicro.2010.07.004

Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2-Approximately Maximum-Likelihood
Trees for Large Alignments [Article]. *PloS One*, *5*(3), e9490–e9490.
https://doi.org/10.1371/journal.pone.0009490

Rahnama, H., Azari, R., Yousefi, M. H., Berizi, E., Mazloomi, S. M., Hosseinzadeh, S., Derakhshan,
Z., Ferrante, M., & Conti, G. O. (2023). A systematic review and meta-analysis of the
prevalence of Bacillus cereus in foods [Article]. *Food Control*, *143*, 109250.
https://doi.org/10.1016/j.foodcont.2022.109250

Rehlund, A. S. G., Canoy, T. S., Nielsen, D. S., Lund, M. N., & Poojary, M. M. (2025). Protein
hydrolysis for amino acid analysis revisited. *Food Chemistry*, 146224.

Robas Mora, M., Fernández Pastrana, V. M., Oliva, L. L. G., Lobo, A. P., & Jiménez Gómez, P. A.
(2023). Plant growth promotion of the forage plant Lupinus albus Var. Orden Dorado using
Pseudomonas agronomica sp. nov. and Bacillus pretiosus sp. nov. added over a valorized
agricultural biowaste [Article]. *Frontiers in Microbiology*, *13*, 1046201.
https://doi.org/10.3389/fmicb.2022.1046201

Schilling, O., Frick, O., Herzberg, C., Ehrenreich, A., Heinzle, E., Wittmann, C., & Stülke, J. (2007).
Transcriptional and Metabolic Responses of Bacillus subtilis to the Availability of Organic
Acids: Transcription Regulation Is Important but Not Sufficient To Account for Metabolic
Adaptation [Article]. *Applied and Environmental Microbiology*, *73*(2), 499–507.
https://doi.org/10.1128/AEM.02084-06

Skriver, E. V., Canoy, T. S., Reilly, E. T., Gkogka, E., Lucena, D., Knøchel, S., & Lyng Røder, H.
(2025). Whole genome sequences of 20 Bacillus cereus strains isolated from various
environments. *Submitted to Publication*.

Snyder, A. B., Martin, N., & Wiedmann, M. (2024). Microbial food spoilage: impact, causative agents
and control strategies [Article]. *Nature Reviews. Microbiology*, *22*(9), 528–542.
https://doi.org/10.1038/s41579-024-01037-x

Sonenshein, A. L. (2007). Control of key metabolic intersections in Bacillus subtilis [Article]. *Nature*
Reviews. Microbiology, *5*(12), 917–927. https://doi.org/10.1038/nrmicro1772

Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: Molecular Evolutionary Genetics Analysis
Version 11 [Article]. *Molecular Biology and Evolution*, *38*(7), 3022–3027.
https://doi.org/10.1093/molbev/msab120

Thorsen, L., Abdelgadir, W. S., Rønsbo, M. H., Abban, S., Hamad, S. H., Nielsen, D. S., & Jakobsen,
M. (2011). Identification and safety evaluation of Bacillus species occurring in high numbers
during spontaneous fermentations to produce Gergoush, a traditional Sudanese bread snack
[Article]. *International Journal of Food Microbiology*, *146*(3), 244–252.
https://doi.org/10.1016/j.ijfoodmicro.2011.02.028

Thorsen, L., Azokpota, P., Hansen, B. M., Hounhouigan, D. J., & Jakobsen, M. (2010). Identification,
genetic diversity and cereulide producing ability of Bacillus cereus group strains isolated from
Beninese traditional fermented food condiments [Article]. *International Journal of Food*
Microbiology, *142*(1), 247–250. https://doi.org/10.1016/j.ijfoodmicro.2010.06.004

UNEP. (2024). *Food Waste Index Report 2024. Think Eat Save: Tracking Progress to Halve Global*
Food Waste. https://wedocs.unep.org/20.500.11822/45230.

684 Valerio, F., De Bellis, P., Lonigro, S. L., Visconti, A., & Lavermicocca, P. (2008). Use of
685   Lactobacillus plantarum fermentation products in bread-making to prevent Bacillus subtilis ropy
686   spoilage [Article]. *International Journal of Food Microbiology*, *122*(3), 328–332.
687   https://doi.org/10.1016/j.ijfoodmicro.2008.01.005
688 Wicke, D., Meißner, J., Warneke, R., Elfmann, C., & Stülke, J. (2023). Understudied proteins and
689   understudied functions in the model bacterium Bacillus subtilis—A major challenge in current
690   research [Article]. *Molecular Microbiology*, *120*(1), 8–19. https://doi.org/10.1111/mmi.15053
691 Wiedenbein, E. S., Canoy, T. S., Hui, Y., Parkouda, C., Compaoré, C. S., Ametefe, E., Jakobsen, M.,
692   Jespersen, L., & Nielsen, D. S. (2023). Complete and Draft Genome Sequences of 52 Bacillus
693   and Priestia Strains Isolated from West African Fermentations and 26 Reference Strains from a
694   Public Culture Collection . *Microbiology Resource Announcements*, *12*(7).
695   https://doi.org/10.1128/mra.00394-23
696 Zwietering, M. H., Jongenburger, I., Rombouts, F. M., & Vantriet, K. (1990). Modeling of the
697   Bacterial Growth Curve [Article]. *Applied and Environmental Microbiology*, *56*(6), 1875–1881.
698   https://doi.org/10.1128/aem.56.6.1875-1881.1990
699

Bacillus paranthracis 7595

Bacillus paranthracis F481072

Bacillus paranthracis GR117

Bacillus paranthracis NS117

100 — Bacillus paranthracis ISI2489

100 Bacillus paranthracis F443073

Bacillus paranthracis LKT11

Bacillus paranthracis PA21

100 Bacillus paranthracis PA24

100 Bacillus paranthracis MA60

100 Bacillus pacificus O-216

Bacillus tropicus F443373

Clade A

100 — Bacillus wiedmannii BC59

100 100 — Bacillus wiedmannii LWL3

Clade B

Bacillus mycoides A536

100 — Bacillus mycoides MC67

Clade C

Bacillus subtilis PRO64

100 Bacillus subtilis PRO31

98 Bacillus subtilis PRO35

100 Bacillus subtilis PRO89

Clade D

Bacillus subtilis DSM13019

65 Bacillus subtilis DSM21393

100 Bacillus subtilis PRO115

100 Bacillus subtilis DSM10

Clade E

Halobacillus halophilus DSM2266

(A) Sugars

Sugar ■ Galactose ■ Arabinose ■ Fructose ■ Glucose ■ Sucrose

(B) Organic acids

Organic acid □ Citric acid ▨ Acetic acid ■ Lactic acid

2A
Accumulated heat over time

2B
Heat flow over time

Group — *cereus* — *subtilis*

A

Comparison of Gompertz heat parameters between *Bacillus cereus* and *Bacillus subtilis*

Combined Metabolite Analysis