Mechanistic insight into the oligomerisation of *Arabidopsis* CRY1 and its inhibition by BIC1

Alicia Just[1], Nils Niemann[2], Petra Gnau[3], Dennis Kock[2], Thomas Heimerl[4], Lars-Oliver Essen[3], Alfred Batschauer[2] & Nina Morgner[1*]

[1]Institute of Physical and Theoretical Chemistry, Johann Wolfgang Goethe University, Frankfurt/Main, Germany

[2]Department of Biology, Philipps University Marburg, Marburg, Germany

[3]Department of Chemistry, Philipps University Marburg, Marburg, Germany

[4]SYNMIKRO Research Center and Department of Chemistry, Philipps University Marburg, Marburg, Germany

*Corresponding author: Nina Morgner, e-mail: morgner@chemie.uni-frankfurt.de

Abstract

Cryptochromes (CRY) convert blue light signals into biological responses, however, the molecular processes underlying their activation are not fully understood. In this study, we uncover the molecular mechanism underlying the blue-light activation of *Arabidopsis* CRY1 using time-resolved native mass spectrometry combined with kinetic modelling – an approach that allows us to monitor light-driven complex formation with temporal and molecular resolution. We show that CRY1 activation follows a defined, reversible assembly pathway in which monomers rapidly form dimers that then assemble into tetramers. A quantitative two-step model captures the dynamic interplay between light-driven assembly and thermal disassembly. Strikingly, ATP accelerates tetramer formation and stabilises oligomers by tuning the underlying photochemistry of the FAD chromophore. In contrast, the Blue-light Inhibitor of Cryptochromes 1 (BIC1) acts as a potent antagonist. We found that BIC1 binds to CRY1 even in the dark, with a significant increase in binding strength under blue-light conditions. BIC1 not only blocks CRY1 oligomerisation, but also actively dismantles pre-assembled tetramers. This disassembly process is light-independent and occurs regardless of CRY1's redox state. Together, these findings reveal a finely balanced regulatory system in which ATP and BIC1 act as opposing regulators to control CRY1 activation. This work provides a kinetic and mechanistic framework for reversible cryptochrome signalling and highlights how blue light responses can be precisely modulated at the molecular level.

Introduction

Plants rely on light not only as an energy source but also as a critical environmental signal that guides nearly every aspect of their growth and development[1–3]. To perceive and translate light information across the spectrum, plants have evolved a sophisticated network of photoreceptors, including phytochromes, phototropins, the UV-B photoreceptor UVR8, and cryptochromes (CRYs)[1,4–6]. Among these, CRYs are evolutionarily conserved flavoproteins that absorb blue and ultraviolet-A light via a flavin adenine dinucleotide (FAD) chromophore, which is bound non-covalently within the photolyase homology region (PHR) domain[7–10]. The PHR domain is complemented by the cryptochrome C-terminal extension (CCE) that is much less

conserved and varies in length and sequence among cryptochromes[11]. In *Arabidopsis*
thaliana, two major cryptochromes, CRY1 and CRY2, operate in partially overlapping
but distinct physiological contexts[12]. CRY1 predominantly controls the early seedling
stage, where it suppresses hypocotyl elongation, whereas CRY2 regulates flowering
time by acting as a photoperiod sensor[12–16]. Activation of plant CRYs is initiated by
light-driven photoreduction of FAD[17,18], which induces conformational changes that
promote oligomerisation[19–22] and binding to downstream signalling partners[23].
Although CRY1 and CRY2 are structurally related[24], their functional specialization
suggests that they may differ in their activation dynamics, signalling interactions, and
mechanisms of regulation. Structural studies have demonstrated that both CRY1 and
CRY2 undergo blue light–induced oligomerisation[13,15–17], and in the case of CRY2,
tetrameric assemblies have been linked to signalling activity[25]. CRY1 likewise forms
light-induced oligomers, as revealed by structural analyses[26], yet the dynamics and
regulation of this process are poorly understood. While oligomerisation is mediated by
the PHR domain, the CCEs are engaged in liquid–liquid phase separation (LLPS)[11].
To distinguish specific oligomerisation from the formation of dynamic size-variable
condensates, *in vitro* studies of the isolated PHR domain are essential to uncover the
intrinsic mechanism of cryptochrome oligomerisation without the interfering influence
of LLPS. In particular, the molecular mechanism underlying CRY1 oligomerisation as
well as the influence of cellular cofactors or inhibitory proteins on this process, remains
poorly understood.

A major class of negative regulators of CRYs are the Blue-light Inhibitors of
Cryptochromes (BICs)[27]. BICs interact with CRY2 in its PHR domain and efficiently
block tetramerization, thereby suppressing signalling[21,24]. CRY1 has also been
identified as a target of BIC1[21,25–27], but the mechanistic details of this inhibition are
not resolved. Previous studies demonstrated that BIC1 may associate with CRY1 in
both dark and light conditions[28], raising the question whether BIC1 primarily prevents
oligomer formation or whether it can actively dismantle pre-assembled CRY1
complexes.

Another layer of potential regulation arises from cellular metabolites[29–31]. ATP is known
to bind the PHR domain of plant cryptochromes[17,18] and a variety of studies reported,
that ATP accelerates FAD photoreduction and stabilizes reduced states[30,32–34].

However, whether ATP directly influences CRY1 oligomerisation dynamics has
remained unclear.

Building on our previous work, where we demonstrated time-resolved investigation of
light-induced conformational changes in the monomeric animal-like cryptochrome
from *Chlamydomonas reinhardtii* (*Cra*CRY)[35], we now take one step further and
explore the oligomerisation process that follows photoactivation in CRY1 from
Arabidopsis thaliana. We combined different native mass spectrometry techniques,
absorption spectroscopy and negative-stain transmission electron microscopy, to
unravel the light-induced assembly and regulation of *Arabidopsis* CRY1 PHR domain
in vitro. We show that CRY1 oligomerisation proceeds via a sequential
monomer → dimer → tetramer pathway that can be quantitatively described by a
reversible two-step kinetic model. ATP accelerates this process, boosting the
oligomerisation rates and stabilizing the oligomeric states. Furthermore, we
demonstrate that BIC1 binds to CRY1 both in darkness and under blue light
illumination, with significantly different affinities. Importantly, BIC1 not only prevents
tetramer assembly but also disassembles preformed CRY1 oligomers, a property that
extends to the constitutively tetrameric CRY1^{L407F} mutant[36,37]. Finally, we show that
BIC1 inhibition does not interfere with photoreduction of the FAD chromophore,
indicating that BIC1 regulates CRY1 through direct competition with CRY1-CRY1
contacts rather than through modulation of chromophore photochemistry.

Together, our findings establish a quantitative kinetic framework for CRY1
oligomerisation and reveal dual regulatory inputs: metabolic tuning by ATP and active
disassembly by BIC1. These results uncover fundamental principles of CRY1
regulation that distinguish it from CRY2 and highlight new mechanisms by which plant
cells integrate metabolic state and the effect of inhibitory proteins to fine-tune
cryptochrome signalling.

Results

Oligomerisation mechanism of *Arabidopsis* CRY1. To gain mechanistic insight into
Arabidopsis CRY1 oligomerisation, we investigated the oligomeric states of the
recombinant CRY1 PHR domain (CRY1-PHR), thereafter named CRY1, *in vitro* by

nano electrospray ionization mass spectrometry (nESI-MS). Consistent with previous

reports[26] CRY1 exists predominantly as a monomer in the dark (m/z = 3500-4800,

Fig. 1a) with only a small fraction present as dimers (m/z = 5000-5700). Exposure to

blue light promotes its oligomerisation into a tetrameric form (m/z = 6000-8500)

(Fig. 1b).

To elucidate the underlying mechanism of this light-induced oligomerisation, we next

examined whether tetramer assembly occurs exclusively via dimer intermediates or

involves trimeric species as transitional state. To address this, we performed time-

resolved mass spectrometry at room temperature, allowing us to detect and quantify

transient oligomeric species under blue light conditions. The time-resolved data

revealed a well-defined kinetic progression of CRY1 oligomerisation (Fig. 1e-g). In the

initial dark-adapted state, the cryptochrome was predominantly monomeric

(Fig. 1a, e). Upon exposure to blue light, we observed a rapid decrease in monomer

abundance (Fig. 1e), accompanied by the appearance of dimers (Fig. 1f). Tetramer

formation was observed after a short delay, consistent with a stepwise assembly

pathway (Fig. 1g). Dimer levels peaked after approximately 3 seconds (Fig. 1f), while

tetramer accumulation progressed more slowly and reached a stationary plateau at

120 seconds (Fig. 1g). Notably, the monomeric species did not fully disappear

(Fig. 1b, e), suggesting that oligomerisation is opposed by a concurrent thermal

dissociation process. This observation shows that even under sustained light

activation, reversible oligomer assembly and disassembly are in kinetic competition.

To quantitatively describe the oligomerisation mechanism, we propose a reversible

kinetic model comprising two steps as shown in Fig. 1l. This kinetic model assumes

sequential assembly, from monomer via dimer to tetramer (M→D→T), with no direct

conversion from monomer to tetramer and no trimeric species. Fitting our data with

this model (Fig. 1e-g, Extended Data Fig. 2a) yielded association and dissociation rate

constants listed in Fig. 1k. The model accurately described the experimental time

courses of all three oligomeric states at room temperature. The relatively low value of

$k_{M \to D}$ = (3.7 ± 0.3) × 10^{-3} $\mu M^{-1} s^{-1}$ indicates slow dimerization, while the larger

association rate $k_{D \to T}$ = (75 ± 4) × 10^{-3} $\mu M^{-1} s^{-1}$ reflects a more efficient tetramerization

step once dimers are present. The dissociation rates further reveal distinct stabilities:

the dimer dissociates rapidly with $k_{D \to M}$ = (560 ± 70) × 10^{-3} s^{-1}, whereas the tetramer

is more stable ($k_{T \to D}$ = (6.3 ± 0.6) × 10^{-3} s^{-1}). Together these rate constants collectively

explain the observed accumulation of tetramers and the persistence of monomers in
a dynamic light-induced stationary state. These results demonstrate that the
experimentally observed oligomerisation dynamics of CRY1 can be effectively
described by a two-step reversible reaction model and provide quantitative insight into
the kinetics of light-induced assembly.

ATP boosts oligomer formation. Given previous reports that ATP influences the
FAD photoreduction kinetics of cryptochromes[30,32–34], we next investigated whether
ATP also affects the light-induced oligomerisation of CRY1. Under identical conditions
in presence and absence of ATP, we observed a pronounced ATP-dependent
enhancement of tetramer formation (Fig. 1c, d). In the presence of 100 µM ATP, CRY1
assembled into tetramers more efficiently and reached markedly higher tetramer levels
upon blue light exposure (Fig. 1d, Extended Data Fig. 2b). Moreover, the rate of
tetramer oligomerisation almost quadrupled, while the subsequent thermal
dissociation in darkness was slowed approximately twofold for the tetramer and even
more than tenfold for the dimer (Fig. 1h-k). To further assess the concentration
dependence of this effect, we also tested a concentration of 200 µM ATP. Under these
conditions, we detected a substantial increase in the dimeric CRY1 population already
in the dark-adapted state compared to samples without ATP or in presence of 100 µM
ATP (Extended Data Fig. 1a-c). This observation suggests that higher ATP
concentrations promote or stabilize dimer formation even in the absence of blue light.
These findings show that ATP enhances and stabilizes dimer formation, which we
observed to be the rate limiting step for tetramer formation under blue light conditions,
revealing a broader modulatory role of ATP in tuning cryptochrome oligomerisation
dynamics. Absorption spectra also show the boosting effect of ATP on the kinetics of
FAD photoreduction, as has already been shown in previous studies[30,32–34]. In our
measurements the addition of ATP accelerated the photoreduction of FAD_{ox} to $FADH°$
by approximately 12%, while slowing down the thermal decay of $FADH°$ back to FAD_{ox}
by nearly 60% (Extended Data Fig. 1d-h).

BIC1 binds CRY1 in both darkness and under blue light. While BIC1 is known to
bind CRY1, it remains unresolved whether this interaction blocks CRY1

oligomerisation or dismantles existing assemblies. We therefore directly investigated

the mechanism of CRY1 inhibition by BIC1 using Laser-Induced Liquid Bead Ion

Desorption mass spectrometry (LILBID-MS)[38]. In line with the nESI-MS data, LILBID-

MS analysis further confirmed that CRY1 exists predominantly as a monomer in the

dark (~ 59 kDa), with only a minor dimeric population (~ 120 kDa) detectable (Fig. 2a).

Blue-light illumination shifted the oligomeric distribution rapidly toward tetrameric

assemblies (~ 240 kDa), underscoring light-dependent higher-order organization of

CRY1 (Fig. 2b). A small trimeric CRY1 population was also detected, arising from

partial dissociation of the tetramer induced by the energy input during LILBID-MS

analysis (Extended Data Fig. 3a). Upon addition of BIC1 (~ 20 kDa) in both the dark

and under blue light conditions, we observed a discrete charge distribution

corresponding to the mass of CRY1-BIC1 heterodimer (~ 79 kDa), indicating direct

binding between the two proteins (Fig. 2c, d). While CRY1 exhibited reduced binding

to BIC1 in the dark (Fig. 2c), we observed a significant increase in CRY1-BIC1

heterodimer formation under blue light conditions (Fig. 2d). Notably, in light-exposed

samples, the CRY1 tetramer signal was absent, suggesting that BIC1 binding prevents

oligomer formation. To quantify the interaction strength of CRY1-BIC1 heterodimer in

the dark and under blue light conditions, we employed quantitative LILBID-MS

(qLILBID-MS), a label-free technique capable of measuring dissociation constants

under native-like conditions[39,40]. In the dark, BIC1 bound CRY1 with moderate affinity

($K_D \approx 1.2\ \mu M$) (Fig. 2g), whereas under blue light, the interaction became more stable

($K_D \approx 0.13\ \mu M$) (Fig. 2f). This ten-fold increase in affinity suggests that light-induced

conformational changes in CRY1 enhance its recognition by BIC1. This demonstrates

that BIC1 binds CRY1 in both its inactive and active state but with a strong preference

for the light-activated form, consistent with a mechanism in which BIC1 selectively

inhibits photoactivated CRY1.

BIC1 reverses CRY1 oligomerisation. Given that CRY1 tetramerizes upon blue light

exposure (Fig. 2b), and that BIC1 binds with high affinity under blue light (Fig. 2d), the

question arises whether BIC1 merely blocks tetramer formation or can even actively

dismantle pre-existing CRY1 oligomers as it was shown for CRY2 and BICs[21,24].

Therefore, we added BIC1 to pre-illuminated, tetrameric CRY1 under constant

illumination with blue light. The mass spectra indicate that BIC1 binds sequentially to

individual CRY1 protomers within the tetramer (~ 260-320 kDa) (Fig. 2e), displacing

them from the tetrameric complex and a formation of a dominant BIC1–CRY1

heterodimer (~ 79 kDa). These data suggest that BIC1 not only prevents tetramer

formation but also actively dismantles existing CRY1 tetramers by occupying all

subunits.

Negative-stain transmission electron microscopy (TEM) corroborates the mass

spectrometry data. In blue light-treated samples, CRY1 formed doughnut-like ring

structures that were absent in dark-adapted preparations (Fig. 2h, i), consistent with

the formation of tetrameric assemblies. The addition of BIC1 markedly reduced the

abundance of these oligomeric particles. Even immediately after 30 minutes of

illumination, CRY1 samples incubated with BIC1 displayed few ring-like structures

(Fig. 2k). Moreover, following transfer to darkness, CRY1 oligomers disassembled

significantly faster in the presence of BIC1, with near-complete monomerization

observed at the earliest time point (Fig. 2j).

BIC1 disassembles CRY1 tetramers independently of light activation. To

examine whether BIC1-mediated inhibition of CRY1 tetramers requires light activation,

we analyzed the CRY1[L407F] mutant, which has been observed to be hyperactive and

results in a constitutive photomorphogenic phenotype[36,37]. Analytical size exclusion

chromatography (SEC) revealed, that CRY1[L407F] shows already a tetrameric retention

time in darkness compared to the wild-type CRY1 (Fig. 3d). LILBID-MS confirmed the

presence of tetramers for CRY1[L407F] in darkness (Fig. 3a) and absorption

spectroscopy has shown that the leucine to phenylalanine replacement at position 407

does not impair photoreduction of the chromophore (Fig. 3h).

Upon addition of BIC1, the CRY1[L407F] tetramers were efficiently disassembled into

BIC1–CRY1[L407F] heterodimers (Fig. 3b, Extended Data Fig. 3b). In contrast to the wild-

type CRY1 (Fig. 2c, d), exposure to blue light did not further enhance the formation of

the CRY1[L407F]-BIC1 heterodimer (Fig. 3c) or have any other significant effect on the

observed complexes. qLILBID measurements revealed that BIC1 binds CRY1[L407F]

independently of illumination with high affinity (K_D = 0.10 ± 0.03 µM), comparable to

the affinity of BIC1 for light-activated wild-type CRY1 (K_D = 0.13 ± 0.09 µM). This

result demonstrates that the L407F mutation induces a conformational change in

CRY1 that mimics its light-activated state with respect to BIC1 recognition.

Collectively, these findings suggest that BIC1 disrupts CRY1 oligomerisation by
binding individual CRY1 protomers with higher affinity than the protomer-protomer
interactions, effectively outcompeting CRY1-CRY1 contacts regardless of the
photoreduction state of the chromophore.

BIC1 does not inhibit photoreduction of the CRY1 FAD cofactor. The remaining
open question is whether BIC1 suppresses CRY1 oligomerisation by interfering with
its photoactivation, as previously demonstrated for CRY2 inhibition by BIC2[24]. To
address this, we examined the photoreduction state of the FAD chromophore in the
presence and absence of BIC1. Absorption spectra of purified CRY1 recorded before
and after blue light exposure showed the expected spectral changes associated with
FAD photoreduction (Fig. 3f, Extended Data Fig. 1d, e). In the dark-adapted state,
CRY1 displayed absorption maxima characteristic of oxidized flavin (FAD_{ox}) at 450 nm
and a shoulder near 480 nm. Upon blue light illumination, these peaks decreased in
intensity, accompanied by the appearance of a broad band between 550-600 nm,
consistent with the formation of the neutral semiquinone radical ($FADH°$)[17,18,41]. These
spectral changes are characteristic for the light-driven redox transition of the CRY1
chromophore. Identical spectral features were observed when CRY1 or CRY1[L407F]
were incubated with BIC1 under the same conditions (Fig. 3g, i, Extended Data
Fig. 4a-c), indicating that BIC1 does not impair the light-induced redox transition of the
FAD_{ox} to FADH°. This suggests that BIC1 does not block the initial photoactivation
step of CRY1. Instead, it acts downstream of photoreduction, targeting the light-
induced oligomeric states of CRY1.

Discussion

Light perception in plants is a dynamic process influenced by multiple molecular
factors. Our kinetic analysis of CRY1 reveals that oligomerisation proceeds through
discrete assembly steps and is modulated by both metabolic signals and inhibitory
interactions. These findings suggest that CRY1 activity can be finely adjusted in
response to changing conditions, enabling a more nuanced regulation of light
signalling.

Our time-resolved native mass spectrometry data establish a clear kinetic pathway for

CRY1 oligomerisation (Fig. 4). The sequential monomer-to-dimer-to-tetramer

assembly we observed is consistent with a two-step reversible kinetic model (Fig. 1l),

identifying dimer formation as the rate-limiting step in CRY1 activation. In comparison,

FAD photoreduction occurs within microseconds[42], much faster than oligomerisation,

indicating that the light-induced activation of the flavin cofactor does not contribute any

significant kinetic delay. The relative stabilities of the oligomers differ substantially,

with dimers being highly transient while tetramers display significantly increased

stability. These light dependent equilibria, suggests that CRY1 signalling is a highly

dynamic process rather than a simple on-off switch. To directly compare our *in vitro*

oligomerisation kinetics with *in vivo* observations made by Liu *et al*. (2020)[22], we fitted

the tetramer formation data using a single-exponential model and determined a half-

life of approximately $t_{1/2}$ = 18 s (Extended Data Fig. 2c). *In vivo*, the corresponding

half-life at a fluence rate of 30 µmol $m^{-2} s^{-1}$ is about five times longer, reflecting

additional cellular constraints such as regulatory interactions (e. g. BIC1 inhibition),

metabolites, different reducing environment, temperature and, of course, different

protein concentrations. Despite these differences, the timescales remain within the

same order of magnitude, underscoring the physiological relevance of the

oligomerisation process captured by time-resolved mass spectrometry. Importantly,

our *in vitro* approach allows disentangling the influence of individual factors that cannot

easily be resolved in the cellular context, thereby complementing *in vivo*

measurements. Beyond this biological relevance, such a detailed kinetic analysis also

opens opportunities for engineering CRY1-based optogenetic tools with tuneable

temporal precision, complementing existing CRY2-based optogenetic systems[43–46].

In addition to intrinsic kinetic insights, our findings highlight how cellular metabolites

can modulate this process. We show that ATP dramatically enhances CRY1

oligomerisation upon illumination (Fig. 4). Increasing ATP concentrations significantly

enhance the formation of CRY1 dimer (Extended Data Fig. 1a-c) which we showed to

be the rate limiting step in oligomer assembly. Then tetramer formation proceeds

rapidly, resulting in more stable higher-order oligomers.

This is particularly intriguing given that ATP is not only a fundamental energy currency

but has also been reported to bind the PHR domain of plant cryptochromes[21,28] and

affect FAD photoreduction kinetics[30,32–34]. Our data support the notion that ATP acts

as a cofactor, stabilizing the oligomeric state and promoting the conformational
changes required for stable oligomer formation. This mechanism suggests a novel link
between the cellular energy status and CRY1 activity, allowing a plant to modulate its
light response in accordance with its metabolic state. For instance, it is conceivable
that under conditions of high energy availability, CRY1 signalling may be sensitized to
support energy-intensive processes such as development and growth.

Our data further elucidate the inhibitory role of BIC1, revealing an elegant mechanism
of regulation (Fig. 4). We demonstrate that BIC1 binds to CRY1 in both dark and light
conditions (Fig. 2c, d), but with an enhanced affinity for the light-activated form
(Fig. 2f, g). This light-dependent increase in binding affinity indicates that light-induced
conformational changes in CRY1 significantly promote its interaction with BIC1. This
light-dependent increase in affinity is a crucial finding, as it provides a mechanism for
the selective inhibition of active CRY1. Moreover, we provide clear evidence that BIC1
is an active disassembler of CRY1 oligomers, rather than just a passive competitor
(Fig. 2e). Given that the constitutively active CRY1^{L407F} mutant, which mimics the
tetrameric light-activated state, shows a high affinity for BIC1 (Fig. 3e), strongly
suggests that BIC1 recognizes a specific conformational state of CRY1, independent
of the FAD redox state. Our result that BIC1 does not interfere with FAD
photoreduction (Fig. 3g, i) differs from recent reports on the CRY2-BIC2 interaction[24],
suggesting that the inhibitory mechanisms are distinct for CRY1 and CRY2. This
supports the idea that while cryptochromes share a conserved photochemical
activation step, their downstream regulation can vary, allowing for functional
specificity.

In summary, our work establishes a quantitative kinetic model for CRY1
oligomerisation. By showing that ATP acts as an oligomerisation accelerator and BIC1
as an active disassembler, we provide a more detailed and nuanced understanding of
how CRY1 activity is regulated beyond simple light perception. These insights not only
shed light on the functional specialization of CRY1 but also highlight a broader
principle of plant signalling, where a photoreceptor's output is shaped by a complex
interplay between light, metabolites, and dedicated regulatory proteins.

Methods

Cloning of constructs. The expression vectors encoding the PHR domain (residues 1–509) of *Arabidopsis thaliana* CRY1 and the CRY1[L407F] mutant, both cloned into the pACYCDuet-1 vector have been described previously[26]. To prevent disulfide bond-mediated dimerization observed in the *Arabidopsis thaliana* wild-type BIC1 protein (Extended Data Fig. 5a, b), a synthetic gene encoding a BIC1 variant carrying three cysteine-to-alanine substitutions (C99A, C105A, C136A) was generated. This triple mutant, named BIC1 throughout the text, was synthesized and cloned into the pET28a(+) vector (*BioCat GmbH*). Recombinant proteins carried an N-terminal His$_6$-tag.

Overproduction and purification of recombinant proteins. For overexpression of CRY1 and the CRY1[L407F] mutant as well as BIC1[3CA], competent *E. coli* BL21 Star (DE3) cells (*Thermo Fisher*) were used. For CRY1 and CRY1[L407F] overproduction, 20 mL overnight cultures were inoculated into 2 L LB medium supplemented with 34 µg/mL chloramphenicol in 5 L baffled flasks. Cultures were incubated at 37 °C with shaking (110 rpm) until an $OD_{600\,nm}$ of ~1.0 was reached. Cells were placed on ice for 30 min, and expression was induced with 0.25 mM IPTG. All further steps were carried out under red light. Cultures were incubated for 3 days at 16 °C. Cells were harvested by centrifugation (5,330g, 15 min, 4 °C) and resuspended in lysis buffer (50 mM MOPS, pH 7.4, 300 mM NaCl, 10% glycerol, 1% Triton X-100). Lysozyme (0.2 mg/mL), PMSF (0.4 mM) and DNase I were added, and the suspensions were incubated for 10 min at room temperature before disruption using a French press. Lysates were clarified by centrifugation (39,000g, 30 min) and filtered. The supernatant was applied onto a 1 mL Protino Ni-NTA column (*Macherey-Nagel*), followed by sequential washing steps with lysis buffer, buffer containing 3% Triton X-100, and lysis buffer without TX-100 until a stable UV baseline was reached. Non-specifically bound proteins were removed with 10% elution buffer (50 mM MOPS, pH 7.4, 300 mM NaCl, 10% glycerol, 250 mM imidazole). Recombinant CRYs were eluted with 100% elution buffer. CRY1 concentrations were kept below 1 mg/mL to prevent oligomerisation. Final purification was performed by size-exclusion chromatography (Superdex 200 pg 16/60, *Cytiva*).

BIC1 purification followed a similar protocol. Expression was induced with 0.5 mM
IPTG, and cultures were incubated for 18 h at 25 °C. Cells were harvested, washed
with PBS, and resuspended in lysis buffer (50 mM HEPES, pH 7.0, 500 mM NaCl, 10%
glycerol). Cell disruption was performed as described above, using a protease inhibitor
mix (*Carl Roth, Cat. no. 3760.1*) in place of PMSF. The soluble fraction was loaded
onto a Ni-NTA column and washed with protein buffer (50 mM HEPES, pH 7.0,
300 mM NaCl, 10% glycerol), followed by a wash containing 3% Triton X-100 and
protein buffer. Additional washes were performed with 10% and 20% elution buffer
(50 mM HEPES, pH 7.0, 300 mM NaCl, 250 mM imidazole, 10% glycerol). BIC1 was
eluted with 100% elution buffer, concentrated, and further purified via size-exclusion
chromatography (Superdex 75 pg 16/60, *Cytiva*).

Size exclusion chromatography (SEC). For analysis of blue light-dependent
oligomerisation, samples containing 30 µM CRY1 and 10 mM ß-mercaptoethanol in
100 µl volumes were separated on a Superdex 200 10/300 GL column (*GE
Healthcare*) equilibrated at 10 °C in buffer D containing 20 mM ß-mercaptoethanol.
CRY1 was pre-illuminated with blue light (465 ± 5 nm, 115 µmol m^{-2} s^{-1}) for 20 min.
Three blue light LEDs were used to illuminate the column resin. Dark controls were
kept in red safe light. Calibration of the column was done using gel-filtration standards
(*Sigma Aldrich*).

Photoreduction and dark reversion of CRY1. Absorbance changes of bound flavin
were measured by UV-Vis spectroscopy (*Specord 600, Analytik Jena*). Before
photoreduction, the CRY1 solution was centrifuged (10 min, 16873 g, 4 °C) to pellet
and remove aggregated protein. A protein solution of 10 µM CRY1 was prepared, and
dithiothreitol (DTT) added to a final concentration of 10 mM. For photoreduction,
samples were irradiated with a 455 nm LED (M455L2, *Thorlabs*) at 100 µmol m^{-2} s^{-1}.
At defined time points, the absorption between 200 nm – 900 nm was recorded. The
dark conversion was monitored with the same photoreduced sample by leaving it in
the dark until full recovery and absorbance recording. During all steps, cuvettes were
chilled at 20 °C.

Nano Electrospray ionization mass spectrometry. nESI-MS was carried out using
a Synapt G2-S instrument (*Waters Corporation, Wilmslow, UK*) equipped with a high-
mass quadrupole modification. Nano electrospray emitters are produced in-house by

pulling borosilicate glass capillaries with a Flaming/Brown micropipette puller (P-1000,

Sutter Instrument Co.) and subsequently coated with a gold layer via sputtering. CRY1

(15 µM) was analyzed at room temperature in positive ion mode under a capillary

voltage of 1.1 kV. The ion source was operated with a cone voltage of 150 V, an offset

of 100 V, and maintained at 20 °C. The scan time was set to 1 scan per second.

Calibration is performed using a standard cesium iodide (CsI) solution. Sample

illumination was carried out by irradiation of the nano electrospray emitters with a

455 nm LED (M455L4, *Thorlabs*) at 30-50 µmol $m^{-2}\,s^{-1}$ (Extended Data Fig. 6a), using

an illumination setup similar to that described by Camacho *et al*. (2019).[47] Acquisition

and data processing are achieved using MassLynx™ V4.1 SCN901. Prior to

measurements, CRY1 was buffer-exchanged in 200 mM ammonium acetate, 10 mM

DTT and pH 7.4, using Zeba Micro Spinbuffer exchangers (7 kDa MWCO, *Thermo*

Fisher Scientific).

nESI-MS based rate constant determination. The measurements of the rate

constants were performed following the same procedure as for the nESI-MS

measurements except for the scan time, which was set to 10 scans per second. The

intensity (counts) of each species (monomer, dimer, tetramer) during the

measurement period was estimated by generating an extracted ion chromatogram

using MassLynx™ V4.1 SCN901. Initial concentrations of each species (monomer,

dimer, tetramer) were determined from the measured signal intensities, assuming a

total sample concentration of 15 µM relative to the molecular mass of the monomer.

Kinetic data were analyzed by using DynaFit[48] software and a reversible two-step

kinetic model (Fig. 1l) based on the differential equations 1-3.

$$\frac{d[M]}{dt} = -2k_{M\to D}[M][M] + 2k_{D\to M}[D] \tag{1}$$

$$\frac{d[D]}{dt} = +k_{M\to D}[M][M] - k_{D\to M}[D] - 2k_{D\to T}[D][D] + 2k_{T\to D}[T] \tag{2}$$

$$\frac{d[T]}{dt} = +k_{D\to T}[D][D] - k_{T\to D}[T] \tag{3}$$

LILBID-MS. Prior to analysis by LILBID-MS[38], CRY1 and BIC1[3CA] were buffer-

exchanged as described for nESI-MS. All measurements were performed at 20 °C,

using 4 µL of buffer-exchanged and degassed sample for each analysis. Microdroplets

with a diameter of 50 µm are generated using a piezo-driven droplet generator (MD-

K-130, *Microdrop Technologies GmbH,* Norderstedt, Germany) operating at 10 Hz.

The droplets are transferred into a vacuum of about 10^{-5} mbar and irradiated with a

pulsed infrared laser (*Innolas Spitlight 400, Continuum PowerLite 8000*) tuned to the

vibrational absorption maximum of water (2.8 µm). Absorption of laser energy (10-

23 mJ per pulse) by water molecules causes an explosive expansion of the droplets,

resulting in the release of analyte ions into the gas phase. The ions are accelerated

using a pulsed electric field and analysed with a home-built time-of-flight mass

spectrometer in negative ion mode. Each mass spectrum represents an average

signal from 1,000 individual microdroplets. Mass spectra are calibrated against a

10 µM aqueous bovine serum albumin standard and subsequently processed by

smoothing and background subtraction. The sample was exposed to blue light

(455 nm LED, M455L4, *Thorlabs*) at 100 µmol $m^{-2}\,s^{-1}$ inside the droplet generator via

a coupled optical fibre that was installed on the generator (Extended Data Fig. 6b).

For qLILBID-MS[39,40] experiments the laser energy transferred into the droplets is

correlated to the degree of complex dissociation. Therefore, the explosion of the

irradiated droplets is visualized by illuminating the explosion with a green laser (Minilite

I, *Continuum*, San Jose, USA) and the resulting images are captured using a high-

speed camera (DFK 23UP031, *Imaging Source*, Bremen, Germany). Each individual

droplet yields both a mass spectrum and a corresponding image of the explosion. The

explosion width for each droplet is determined by analysing the recorded images with

OpenCV, using a custom written Python script. The resulting dissociation plots allow

K_D determination.

TEM negative staining. Carbon coated copper grids (400 mesh) were hydrophilized

by glow discharging (*PELCO easiGlow*, Ted Pella, USA). 5 µl of a protein suspension

with a concentration of 30 µg/ml was applied onto the hydrophilized grids and stained

with 2% uranyl acetate after a short washing step with filtered H_2O_{bidest}. If applicable,

samples were prepared under blue (465 ± 5 nm, 115 µmol $m^{-2}s^{-1}$) or safe red light,

respectively. Samples were analysed with a JEOL JEM-2100 transmission electron

microscope using an acceleration voltage of 120 kV. For image acquisition a F214

FastScan CCD camera (*TVIPS, Gauting*) was used.

Data Availability

All data generated in this study are provided as Source Data with this paper. Gene accessions codes are At3G52740 (*BIC1*) and At4G08920 (*CRY1*).

References

1. Franklin, K. A., Larner, V. S. & Whitelam, G. C. The signal transducing photoreceptors of plants. *Int. J. Dev. Biol.* **49**, 653–664 (2005).

2. Halliday, K. J., Martínez-García, J. F. & Josse, E.-M. Integration of Light and Auxin Signaling. *Cold Spring Harb. Perspect. Biol.* **1**, a001586 (2009).

3. Heijde, M. & Ulm, R. UV-B photoreceptor-mediated signalling in plants. *Trends Plant Sci.* **17**, 230–237 (2012).

4. Fankhauser, C. The Phytochromes, a Family of Red/Far-red Absorbing Photoreceptors *. *J. Biol. Chem.* **276**, 11453–11456 (2001).

5. Christie, J. M. & Briggs, W. R. Blue Light Sensing in Higher Plants *. *J. Biol. Chem.* **276**, 11457–11460 (2001).

6. Yadav, A. *et al.* Light signaling and UV-B-mediated plant growth regulation. *J. Integr. Plant Biol.* **62**, 1270–1292 (2020).

7. Lin, C. *et al.* Association of Flavin Adenine Dinucleotide with the *Arabidopsis* Blue Light Receptor CRY1. *Science* **269**, 968–970 (1995).

8. Malhotra, K., Kim, S.-T., Batschauer, A., Dawut, L. & Sancar, A. Putative Blue-Light Photoreceptors from Arabidopsis thaliana and Sinapis alba with a High Degree of Sequence Homology to DNA Photolyase Contain the Two Photolyase Cofactors but Lack DNA Repair Activity. *Biochemistry* **34**, 6892–6899 (1995).

9. Brautigam, C. A. *et al.* Structure of the photolyase-like domain of cryptochrome 1 from Arabidopsis thaliana. *Proc. Natl. Acad. Sci.* **101**, 12142–12147 (2004).

10.   Hanić, M. *et al.* Dimerization of European Robin Cryptochrome 4a. *J. Phys.*

Chem. B **127**, 6251–6264 (2023).

11.   Wang, X. & Lin, C. The two action mechanisms of plant cryptochromes.

Trends Plant Sci. **30**, 775–791 (2025).

12.   Yang, Z. *et al.* Cryptochromes Orchestrate Transcription Regulation of

Diverse Blue Light Responses in Plants. *Photochem. Photobiol.* **93**, 112–127

(2017).

13.   Ahmad, M. & Cashmore, A. R. HY4 gene of A. thaliana encodes a protein with

characteristics of a blue-light photoreceptor. *Nature* **366**, 162–166 (1993).

14.   Guo, H., Yang, H., Mockler, T. C. & Lin, C. Regulation of Flowering Time by

Arabidopsis Photoreceptors. *Science* **279**, 1360–1363 (1998).

15.   Wang, J., Du, X., Pan, W., Wang, X. & Wu, W. Photoactivation of the

cryptochrome/photolyase superfamily. *J. Photochem. Photobiol. C Photochem.*

Rev. **22**, 84–102 (2015).

16.   Wang, Q. & Lin, C. Mechanisms of Cryptochrome-Mediated Photoresponses

in Plants. *Annu. Rev. Plant Biol.* **71**, 103–129 (2020).

17.   Bouly, J.-P. *et al.* Cryptochrome Blue Light Photoreceptors Are Activated

through Interconversion of Flavin Redox States*. *J. Biol. Chem.* **282**, 9383–9391

(2007).

18.   Banerjee, R. *et al.* The Signaling State of Arabidopsis Cryptochrome 2

Contains Flavin Semiquinone *. *J. Biol. Chem.* **282**, 14916–14922 (2007).

19.   Sang, Y. *et al.* N-Terminal Domain–Mediated Homodimerization Is Required

for Photoreceptor Activity of Arabidopsis CRYPTOCHROME 1. *Plant Cell* **17**,

1569–1584 (2005).

20.   Rosenfeldt, G., Viana, R. M., Mootz, H. D., von Arnim, A. G. & Batschauer, A.

Chemically Induced and Light-Independent Cryptochrome Photoreceptor

Activation. *Mol. Plant* **1**, 4–14 (2008).

21.   Wang, Q. *et al.* Photoactivation and inactivation of Arabidopsis cryptochrome

2. *Science* **354**, 343–347 (2016).

22.   Liu, Q. *et al.* Photooligomerization Determines Photosensitivity and

Photoreactivity of Plant Cryptochromes. *Mol. Plant* **13**, 398–413 (2020).

23.   Ponnu, J. & Hoecker, U. Signaling Mechanisms by Arabidopsis

Cryptochromes. *Front. Plant Sci.* **13**, 844714 (2022).

24.   Ma, L. *et al.* Structural insights into BIC-mediated inactivation of Arabidopsis

cryptochrome 2. *Nat. Struct. Mol. Biol.* **27**, 472–479 (2020).

25.   Palayam, M. *et al.* Structural insights into photoactivation of plant

Cryptochrome-2. *Commun. Biol.* **4**, 28 (2021).

26.   Shao, K. *et al.* The oligomeric structures of plant cryptochromes. *Nat. Struct.*

Mol. Biol. **27**, 480–488 (2020).

27.   Wang, X. *et al.* A CRY–BIC negative-feedback circuitry regulating blue light

sensitivity of Arabidopsis. *Plant J. Cell Mol. Biol.* **92**, 426–436 (2017).

28.   Miao, L. *et al.* Arabidopsis cryptochrome 1 undergoes COP1 and LRBs-

dependent degradation in response to high blue light. *New Phytol.* **234**, 1347–

1362 (2022).

29.   El-Esawi, M. *et al.* Cellular metabolites modulate in vivo signaling of

Arabidopsis cryptochrome-1. *Plant Signal. Behav.* **10**, e1063758 (2015).

30.   Engelhard, C. *et al.* Cellular metabolites enhance the light sensitivity of

Arabidopsis cryptochrome through alternate electron transfer pathways. *Plant Cell*

26, 4519–4531 (2014).

31.   Zhao, X. *et al.* A Study of Gibberellin Homeostasis and Cryptochrome-

Mediated Blue Light Inhibition of Hypocotyl Elongation. *Plant Physiol.* **145**, 106–

118 (2007).

32.   Immeln, D., Schlesinger, R., Heberle, J. & Kottke, T. Blue Light Induces

Radical Formation and Autophosphorylation in the Light-sensitive Domain of

Chlamydomonas Cryptochrome*. *J. Biol. Chem.* **282**, 21720–21728 (2007).

33.   Müller, P. *et al.* ATP Binding Turns Plant Cryptochrome Into an Efficient

Natural Photoswitch. *Sci. Rep.* **4**, 5175 (2014).

34.   Eckel, M., Steinchen, W. & Batschauer, A. ATP boosts lit state formation and

activity of Arabidopsis cryptochrome 2. *Plant J.* **96**, 389–403 (2018).

35.   Zangl, R. *et al.* Time-Resolved Ion Mobility Mass Spectrometry to Solve

Conformational Changes in a Cryptochrome. *J. Am. Chem. Soc.* **146**, 14468–

14478 (2024).

36.   Exner, V. *et al.* A Gain-of-Function Mutation of Arabidopsis

CRYPTOCHROME1 Promotes Flowering. *Plant Physiol.* **154**, 1633–1645 (2010).

37.   Orth, C., Niemann, N., Hennig, L., Essen, L.-O. & Batschauer, A. Hyperactivity

of the Arabidopsis cryptochrome (cry1) L407F mutant is caused by a structural

alteration close to the cry1 ATP-binding site. *J. Biol. Chem.* **292**, 12906–12920

(2017).

38.   Peetz, O. *et al.* LILBID and nESI: Different Native Mass Spectrometry

Techniques as Tools in Structural Biology. *J. Am. Soc. Mass Spectrom.* **30**, 181–

191 (2019).

39.   Schulte, J., Tants, J.-N., von Ehr, J., Schlundt, A. & Morgner, N.

Determination of dissociation constants via quantitative mass spectrometry. *Front.*

Anal. Sci. **3**, (2023).

40.   Young, P., Hense, G., Immer, C., Wöhnert, J. & Morgner, N. LILBID laser

dissociation curves: a mass spectrometry-based method for the quantitative

assessment of dsDNA binding affinities. *Sci. Rep.* **10**, 20398 (2020).

41.   Liu, H., Zhong, D. & Lin, C. Searching for a photocycle of the cryptochrome

photoreceptors. *Curr. Opin. Plant Biol.* **13**, 578–586 (2010).

42.   Langenbacher, T., Immeln, D., Dick, B. & Kottke, T. Microsecond Light-

Induced Proton Transfer to Flavin in the Blue Light Sensor Plant Cryptochrome. *J.*

Am. Chem. Soc. **131**, 14274–14280 (2009).

43.   Taslimi, A. *et al.* An optimized optogenetic clustering tool for probing protein

interaction and function. *Nat. Commun.* **5**, 4925 (2014).

44.   Che, D. L., Duan, L., Zhang, K. & Cui, B. The Dual Characteristics of Light-

Induced Cryptochrome 2, Homo-oligomerization and Heterodimerization, for

Optogenetic Manipulation in Mammalian Cells. *ACS Synth. Biol.* **4**, 1124–1135

(2015).

45.   Park, H. *et al.* Optogenetic protein clustering through fluorescent protein

tagging and extension of CRY2. *Nat. Commun.* **8**, 30 (2017).

46.   Hallett, R. A., Zimmerman, S. P., Yumerefendi, H., Bear, J. E. & Kuhlman, B.

Correlating in Vitro and in Vivo Activities of Light-Inducible Dimers: A Cellular

Optogenetics Guide. *ACS Synth. Biol.* **5**, 53–64 (2016).

47.   Camacho, I. S. *et al.* Native mass spectrometry reveals the conformational

diversity of the UVR8 photoreceptor. *Proc. Natl. Acad. Sci.* **116**, 1116–1125

(2019).

48.   Kuzmic, P. Program DYNAFIT for the analysis of enzyme kinetic data:

application to HIV proteinase. *Anal. Biochem.* **237**, 260–273 (1996).

Acknowledgements

We thank Konstantinos Stamatakis for his support and fruitful discussions during the kinetic modelling process, and Josef Wachtveitl for providing access to the absorption spectrometer. We also acknowledge Jeanette Schermuly for technical assistance and Stephan Kiontke for his contributions to the recombinant expression of BIC1 and for helpful discussions. Funding was provided by the German Research Foundation (DFG): A.J. was supported through the Collaborative Research Center CRC 1507 "Membrane-Associated Protein Assemblies, Machineries and Supercomplexes", N.M. through DFG Grant 557111829, L.-O.E. through DFG grant ES152/18 and A.B. through DFG grant BA985/15-1.

Author contributions

A.J., N.M., L.-O.E., A.B. designed and coordinated the experiments. P.G., A.J., N.N. and D.K. expressed and purified the proteins. A.J. performed mass spectrometry, UV-Vis spectroscopy and kinetic modelling. N.N. and T.H. acquired and interpreted the TEM data. N.N. acquired and interpreted the SEC data. A.J. wrote the original draft. A.J., N.M., A.B., P.G., L.-O.E. edited the manuscript.

Competing interests

The authors declare no competing interests.

Figures and Legends

Fig. 1: Oligomerisation kinetics of CRY1 under blue light and the influence of ATP. nESI-MS spectrum of CRY1 in the dark (grey line) (**a, c**) and under illumination with blue light (blue line) (**b, d**) in absence (**a, b**) and presence of 100 µM ATP (**c, d**). Exposure to blue light leads to oligomerisation of CRY1, and the addition of ATP promotes this process. Time-resolved mass spectrometry measurements of the CRY1 monomer (**e**), dimer (**f**) and tetramer (**g**) in absence and presence of ATP (**h-j**) and the fit (pink line) obtained from the two-step reversible

Fig. 2: BIC1 inhibits CRY1 oligomerisation in both dark and blue light conditions with higher affinity of BIC1 to CRY1 under blue light conditions. LILBID-MS spectrum of CRY1 shows a monomer (pink dot) (**a**) in darkness, which forms tetramers upon illumination with blue light (**b**). CRY1 trimers are caused by laser dissociation. **c,** Incubation of CRY1 monomer and BIC1 (purple dot) in an equimolar ratio in darkness results in a mixture of CRY1-BIC1 heterodimer and CRY1 monomer. **d,** Upon blue light exposure, the equilibrium shifts towards a higher proportion of the CRY1-BIC1 heterodimer relative to the CRY1 monomer (highlighted in grey). **e,** Under blue light conditions, addition of BIC1 to the CRY1 tetramer results in binding to each protomer, thereby dissociating the tetramer into CRY1-BIC1 heterodimers. **f,** qLILBID-MS measurements reveal that the CRY1-BIC1 heterodimer exhibits a ten-fold higher affinity under blue light compared to its dark-state counterpart (**g**). TEM images of CRY1 particles visualised by negative staining of probes kept in darkness (**h**), or treated with blue light (450 nm, 20 min, 115 µmol m^{-2} s^{-1}) (**i**) showing light-induced formation of doughnut-like ring structures of CRY1. Scale bars 10 nm. **j,** Kinetics of CRY1 monomerization (n = 3 ± SD) after transfer of samples from blue light to darkness in absence and presence of BIC1 or *E. coli* GroEL as negative control. **k,** Effect of BIC1 on kinetics of

 decrease in number of CRY1 particles showing monomerization of CRY1 (n = 3 ± SD). Blue light (465 nm, fluence

 rate 115 µmol m^{-2} s^{-1}) treatment was for 20 min.

Fig. 3: BIC1 disassembles constitutively active CRY1 tetramers and does not inhibit photoreduction of the FAD cofactor. a, LILBID-MS spectrum of the constitutively active CRY1^{L407F} mutant shows a tetramer in darkness. CRY1 trimers are caused by laser dissociation. **b**, Addition of BIC1 (purple dot) to CRY1^{L407F} in an equimolar ratio in darkness results in binding to each protomer, thereby dissociating the CRY1^{L407F} tetramer into CRY1^{L407F}-BIC1 heterodimer and monomeric CRY1^{L407F}. **c**, Exposure to blue light does not shift the equilibrium toward the CRY1^{L407F}-BIC1 heterodimer, as has been shown for wild-type CRY1. **d**, SEC elution profiles of wild-type CRY1 kept in darkness or irradiated with blue light, and of hyperactive CRY1^{L407F} mutant kept in darkness. **e**, qLILBID-MS measurement of the CRY1^{L407F}-BIC1 heterodimer under dark conditions revealed that BIC1 binds CRY1^{L407F} with high affinity. **f**, Absorption spectra of wild-type CRY1 before (grey line) and after (blue line) blue light exposure indicate FAD photoreduction, with a decrease in FAD$_{ox}$ peak at 450 nm, and the emergence of a broad band at 550-600 nm, characteristic for the FADH° semiquinone. **g**, Addition of BIC1 to wild-type CRY1 does not inhibit the photoreduction of CRY's FAD chromophore. **h**, Absorption spectra of the CRY1^{L407F} mutant indicates that chromophore photoreduction remains unaffected by the mutation. **i**, Photoreduction of the FAD chromophore in the CRY1^{L407F} mutant remains unaffected by BIC1 addition, as observed for wild-type CRY1.

Fig.4: Schematic model of CRY1 oligomerisation dynamics, ATP modulation, and BIC1-mediated inhibition.
Upon blue light exposure, CRY1 transitions from a monomeric to a tetrameric state via a short-lived dimeric
intermediate. This oligomerisation follows a sequential, reversible two-step mechanism. ATP enhances the
association rates for dimer and tetramer formation, while slowing down the dissociation rates of these oligomers.
BIC1 binds CRY1 in both dark and light conditions, with significantly higher affinity for the light-activated
conformation. This interaction prevents oligomer formation and actively disassembles pre-formed tetramers by
binding to individual CRY1 protomers even under continuous blue-light illumination.