Mechanistic insight into the oligomerization of *Arabidopsis* CRY1 and its inhibition by BIC1

Alicia Just[1], Nils Niemann[2], Petra Gnau[3], Dennis Kock[2], Thomas Heimerl[4], Lars-Oliver Essen[3], Alfred Batschauer[2] & Nina Morgner[1*]

[1]Institute of Physical and Theoretical Chemistry, Johann Wolfgang Goethe University, Frankfurt/Main, Germany

[2]Department of Biology, Philipps University Marburg, Marburg, Germany

[3]Department of Chemistry, Philipps University Marburg, Marburg, Germany

[4]SYNMIKRO Research Center and Department of Chemistry, Philipps University Marburg, Marburg, Germany

*Corresponding author: Nina Morgner, e-mail: morgner@chemie.uni-frankfurt.de

Abstract

Cryptochromes (CRY) convert blue light signals into biological responses, however, the molecular processes underlying their activation are not fully understood. In this study, we elucidate the mechanism by which *Arabidopsis* CRY1 forms signalling-active tetramers in response to blue light using time-resolved native mass spectrometry combined with kinetic modelling. We found that oligomerization proceeds via a defined, reversible pathway in which monomers rapidly form dimers that then assemble into tetramers. A quantitative two-step model captures the dynamic interplay between light-driven assembly and thermal disassembly. Strikingly, ATP amplifies this response, accelerating tetramer formation, stabilising oligomers and tuning the underlying photochemistry of the FAD chromophore. In contrast, the Blue-light Inhibitor of Cryptochromes 1 (BIC1) binds to CRY1 with nanomolar affinity under blue light conditions, acting as a potent antagonist by not only blocking oligomerization, but also actively dismantling pre-assembled tetramers. This process is light-independent and occurs regardless of CRY1's redox state. Together, these findings reveal a finely balanced regulatory system in which ATP and BIC1 act as opposing regulators to control CRY1 activation. This work provides a kinetic and mechanistic framework for reversible cryptochrome signalling and highlights how blue light responses can be precisely modulated at the molecular level.

Introduction

Plants rely on light not only as an energy source but also as a critical environmental signal that guides nearly every aspect of their growth and development[1,2]. To perceive and translate light information across the spectrum, plants have evolved a sophisticated network of photoreceptors, including phytochromes, phototropins, the UV-B photoreceptor UVR8, and cryptochromes (CRYs)[1,3–5]. Among these, CRYs are evolutionarily conserved flavoproteins that absorb blue and ultraviolet-A light via a flavin adenine dinucleotide (FAD) chromophore, which is bound non-covalently within the photolyase homology region (PHR) domain[6–8]. The PHR domain is complemented by the cryptochrome C-terminal extension (CCE) that is much less conserved and varies in length and sequence among cryptochromes[9]. In *Arabidopsis thaliana*, two major cryptochromes, CRY1 and CRY2, operate in partially overlapping but distinct

physiological contexts[10]. CRY1 predominantly controls the early seedling stage, where it suppresses hypocotyl elongation, whereas CRY2 regulates flowering time by acting as a photoperiod sensor[10–14]. Activation of plant CRYs is initiated by light-driven photoreduction of FAD[15,16], which induces conformational changes that promote oligomerization[17–20] and binding to downstream signalling partners[21]. Although CRY1 and CRY2 are structurally related[22], their functional specialization suggests that they may differ in their activation dynamics, signalling interactions, and mechanisms of regulation. Structural studies have demonstrated that both CRY1 and CRY2 undergo blue light–induced oligomerization[11,13–15], and in the case of CRY2, tetrameric assemblies have been linked to signalling activity[23]. CRY1 likewise forms light-induced oligomers, as revealed by structural analyses[24], yet the dynamics and regulation of this process are poorly understood. While oligomerization is mediated by the PHR domain, the CCEs are engaged in liquid–liquid phase separation (LLPS)[9]. To distinguish specific oligomerisation from the formation of dynamic size-variable condensates, *in vitro* studies of the isolated PHR domain are essential to uncover the intrinsic mechanism of cryptochrome oligomerization without the interfering influence of LLPS. In particular, the molecular mechanism underlying CRY1 oligomerization as well as the influence of cellular cofactors or inhibitory proteins on this process, remains poorly understood.

A major class of negative regulators of CRYs are the Blue-light Inhibitors of Cryptochromes (BICs)[25]. BICs interact with CRY2 in its PHR domain and efficiently block tetramerization, thereby suppressing signalling[19,22]. CRY1 has also been identified as a target of BIC1[19,23–25], but the mechanistic details of this inhibition are not resolved. Previous studies demonstrated that BIC1 may associate with CRY1 in both dark and light conditions[26], raising the question whether BIC1 primarily prevents oligomer formation or whether it can actively dismantle pre-assembled CRY1 complexes.

Another layer of potential regulation arises from cellular metabolites[27–29]. ATP is known to bind the PHR domain of plant cryptochromes[17,18] and a variety of studies reported, that ATP accelerates FAD photoreduction and stabilizes reduced states[28,30–32]. However, whether ATP directly influences CRY1 oligomerization dynamics has remained unclear.

Here, we combined different native mass spectrometry techniques, absorption
spectroscopy and negative-stain transmission electron microscopy, to unravel the
light-induced assembly and regulation of *Arabidopsis* CRY1 PHR domain *in vitro*. We
show that CRY1 oligomerization proceeds via a sequential
monomer → dimer → tetramer pathway that can be quantitatively described by a
reversible two-step kinetic model. ATP accelerates this process, boosting the
oligomerization rates and stabilizing the oligomeric states. Furthermore, we
demonstrate that BIC1 binds to CRY1 both in darkness and under blue light
illumination, with significantly different affinities. Importantly, BIC1 not only prevents
tetramer assembly but also disassembles preformed CRY1 oligomers, a property that
extends to the constitutively tetrameric CRY1^{L407F} mutant[33,34]. Finally, we show that
BIC1 inhibition does not interfere with photoreduction of the FAD chromophore,
indicating that BIC1 regulates CRY1 through direct competition with CRY1-CRY1
contacts rather than through modulation of chromophore photochemistry.

Together, our findings establish a quantitative kinetic framework for CRY1
oligomerization and reveal dual regulatory inputs: metabolic tuning by ATP and active
disassembly by BIC1. These results uncover fundamental principles of CRY1
regulation that distinguish it from CRY2 and highlight new mechanisms by which plant
cells integrate metabolic state and the effect of inhibitory proteins to fine-tune
cryptochrome signalling.

Results

Oligomerization mechanism of *Arabidopsis* CRY1. To gain mechanistic insight into
Arabidopsis CRY1 oligomerization, we investigated the oligomeric states of the
recombinant CRY1 PHR domain (CRY1-PHR), thereafter named CRY1, *in vitro* by
nano electrospray ionization mass spectrometry (nESI-MS). Consistent with previous
reports[24] CRY1 exists predominantly as a monomer in the dark (m/z = 3500-4800,
Fig. 1a) with only a small fraction present as dimers (m/z = 5000-5700). Exposure to
blue light promotes its oligomerization into a tetrameric form (m/z = 6000-8500)
(Fig. 1b).

To elucidate the underlying mechanism of this light-induced oligomerization, we next
examined whether tetramer assembly occurs exclusively via dimer intermediates or
involves trimeric species as transitional state. To address this, we performed time-
resolved mass spectrometry at room temperature, allowing us to detect and quantify
transient oligomeric species under blue light conditions. The time-resolved data
revealed a well-defined kinetic progression of CRY1 oligomerization (Fig. 1e-g). In the
initial dark-adapted state, the cryptochrome was predominantly monomeric
(Fig. 1a, e). Upon exposure to blue light, we observed a rapid decrease in monomer
abundance (Fig. 1e), accompanied by the appearance of dimers (Fig. 1f). Tetramer
formation was observed after a short delay, consistent with a stepwise assembly
pathway (Fig. 1g). Dimer levels peaked after approximately 3 seconds (Fig. 1f), while
tetramer accumulation progressed more slowly and reached a stationary plateau at
120 seconds (Fig. 1g). Notably, the monomeric species did not fully disappear
(Fig. 1b, e), suggesting that oligomerization is opposed by a concurrent thermal
dissociation process. This observation shows that even under sustained light
activation, reversible oligomer assembly and disassembly are in kinetic competition.

To quantitatively describe the oligomerization mechanism, we propose a reversible
kinetic model comprising two steps as shown in Fig. 1l. This kinetic model assumes
sequential assembly, from monomer via dimer to tetramer (M→D→T), with no direct
conversion from monomer to tetramer and no trimeric species. Fitting our data with
this model (Fig. 1e-g, Extended Data Fig. 2a) yielded association and dissociation rate
constants listed in Fig. 1k. The model accurately described the experimental time
courses of all three oligomeric states at room temperature. The relatively low value of
$k_{M \to D} = (3.7 \pm 0.3) \times 10^{-3} \, \mu M^{-1} s^{-1}$ indicates slow dimerization, while the larger
association rate $k_{D \to T} = (75 \pm 4) \times 10^{-3} \, \mu M^{-1} s^{-1}$ reflects a more efficient tetramerization
step once dimers are present. The dissociation rates further reveal distinct stabilities:
the dimer dissociates rapidly with $k_{D \to M} = (560 \pm 70) \times 10^{-3} \, s^{-1}$, whereas the tetramer
is more stable ($k_{T \to D} = (6.3 \pm 0.6) \times 10^{-3} \, s^{-1}$). Together these rate constants collectively
explain the observed accumulation of tetramers and the persistence of monomers in
a dynamic light-induced stationary state. These results demonstrate that the
experimentally observed oligomerization dynamics of CRY1 can be effectively
described by a two-step reversible reaction model and provide quantitative insight into
the kinetics of light-induced assembly.

ATP boosts oligomer formation. Given previous reports that ATP influences the FAD photoreduction kinetics of cryptochromes[28,30–32], we next investigated whether ATP also affects the light-induced oligomerization of CRY1. Under identical conditions in presence and absence of ATP, we observed a pronounced ATP-dependent enhancement of tetramer formation (Fig. 1c, d). In the presence of 100 μM ATP, CRY1 assembled into tetramers more efficiently and reached markedly higher tetramer levels upon blue light exposure (Fig. 1d, Extended Data Fig. 2b). Moreover, the rate of tetramer oligomerization almost quadrupled, while the subsequent thermal dissociation in darkness was slowed approximately twofold for the tetramer and even more than tenfold for the dimer (Fig. 1h-k). To further assess the concentration dependence of this effect, we also tested a concentration of 200 μM ATP. Under these conditions, we detected a substantial increase in the dimeric CRY1 population already in the dark-adapted state compared to samples without ATP or in presence of 100 μM ATP (Extended Data Fig. 1a-c). This observation suggests that higher ATP concentrations promote or stabilize dimer formation even in the absence of blue light. These findings show that ATP enhances and stabilizes dimer formation, which we observed to be the rate limiting step for tetramer formation under blue light conditions, revealing a broader modulatory role of ATP in tuning cryptochrome oligomerization dynamics. Absorption spectra also show the boosting effect of ATP on the kinetics of FAD photoreduction, as has already been shown in previous studies[28,30–32]. In our measurements the addition of ATP accelerated the photoreduction of FAD_{ox} to FADH° by approximately 12%, while slowing down the thermal decay of FADH° back to FAD_{ox} by nearly 60% (Extended Data Fig. 1d-h).

BIC1 binds CRY1 in both darkness and under blue light. While BIC1 is known to bind CRY1, it remains unresolved whether this interaction blocks CRY1 oligomerization or dismantles existing assemblies. We therefore directly investigated the mechanism of CRY1 inhibition by BIC1 using Laser-Induced Liquid Bead Ion Desorption mass spectrometry (LILBID-MS)[35]. In line with the nESI-MS data, LILBID-MS analysis further confirmed that CRY1 exists predominantly as a monomer in the dark (~ 59 kDa), with only a minor dimeric population (~ 120 kDa) detectable (Fig. 2a).

Blue-light illumination shifted the oligomeric distribution rapidly toward tetrameric
assemblies (~ 240 kDa), underscoring light-dependent higher-order organization of
CRY1 (Fig. 2b). A small trimeric CRY1 population was also detected, arising from
partial dissociation of the tetramer induced by the energy input during LILBID-MS
analysis (Extended Data Fig. 3a). Upon addition of BIC1 (~ 20 kDa) in both the dark
and under blue light conditions, we observed a discrete charge distribution
corresponding to the mass of CRY1-BIC1 heterodimer (~ 79 kDa), indicating direct
binding between the two proteins (Fig. 2c, d). While CRY1 exhibited reduced binding
to BIC1 in the dark (Fig. 2c), we observed a significant increase in CRY1-BIC1
heterodimer formation under blue light conditions (Fig. 2d). Notably, in light-exposed
samples, the CRY1 tetramer signal was absent, suggesting that BIC1 binding prevents
oligomer formation. To quantify the interaction strength of CRY1-BIC1 heterodimer in
the dark and under blue light conditions, we employed quantitative LILBID-MS
(qLILBID-MS), a label-free technique capable of measuring dissociation constants
under native-like conditions[36,37]. In the dark, BIC1 bound CRY1 with moderate affinity
($K_D \approx 1.2$ µM) (Fig. 2g), whereas under blue light, the interaction became more stable
($K_D \approx 0.13$ µM) (Fig. 2f). This ten-fold increase in affinity suggests that light-induced
conformational changes in CRY1 enhance its recognition by BIC1. This demonstrates
that BIC1 binds CRY1 in both its inactive and active state but with a strong preference
for the light-activated form, consistent with a mechanism in which BIC1 selectively
inhibits photoactivated CRY1.

BIC1 reverses CRY1 oligomerization. Given that CRY1 tetramerizes upon blue light
exposure (Fig. 2b), and that BIC1 binds with high affinity under blue light (Fig. 2d), the
question arises whether BIC1 merely blocks tetramer formation or can even actively
dismantle pre-existing CRY1 oligomers as it was shown for CRY2 and BICs[19,22].
Therefore, we added BIC1 to pre-illuminated, tetrameric CRY1 under constant
illumination with blue light. The mass spectra indicate that BIC1 binds sequentially to
individual CRY1 protomers within the tetramer (~ 260-320 kDa) (Fig. 2e), displacing
them from the tetrameric complex and a formation of a dominant BIC1–CRY1
heterodimer (~ 79 kDa). These data suggest that BIC1 not only prevents tetramer
formation but also actively dismantles existing CRY1 tetramers by occupying all
subunits.

Negative-stain transmission electron microscopy (TEM) corroborates the mass spectrometry data. In blue light-treated samples, CRY1 formed doughnut-like ring structures that were absent in dark-adapted preparations (Fig. 2h, i), consistent with the formation of tetrameric assemblies. The addition of BIC1 markedly reduced the abundance of these oligomeric particles. Even immediately after 30 minutes of illumination, CRY1 samples incubated with BIC1 displayed few ring-like structures (Fig. 2k). Moreover, following transfer to darkness, CRY1 oligomers disassembled significantly faster in the presence of BIC1, with near-complete monomerization observed at the earliest time point (Fig. 2j).

BIC1 disassembles CRY1 tetramers independently of light activation. To examine whether BIC1-mediated inhibition of CRY1 tetramers requires light activation, we analyzed the CRY1^{L407F} mutant, which has been observed to be hyperactive and results in a constitutive photomorphogenic phenotype[33,34]. Analytical size exclusion chromatography (SEC) revealed, that CRY1^{L407F} shows already a tetrameric retention time in darkness compared to the wild-type CRY1 (Fig. 3d). LILBID-MS confirmed the presence of tetramers for CRY1^{L407F} in darkness (Fig. 3a) and absorption spectroscopy has shown that the leucine to phenylalanine replacement at position 407 does not impair photoreduction of the chromophore (Fig. 3h).

Upon addition of BIC1, the CRY1^{L407F} tetramers were efficiently disassembled into BIC1–CRY1^{L407F} heterodimers (Fig. 3b, Extended Data Fig. 3b). In contrast to the wild-type CRY1 (Fig. 2c, d), exposure to blue light did not further enhance the formation of the CRY1^{L407F}-BIC1 heterodimer (Fig. 3c) or have any other significant effect on the observed complexes. qLILBID measurements revealed that BIC1 binds CRY1^{L407F} independently of illumination with high affinity ($K_D = 0.10 \pm 0.03$ µM), comparable to the affinity of BIC1 for light-activated wild-type CRY1 ($K_D = 0.13 \pm 0.09$ µM). This result demonstrates that the L407F mutation induces a conformational change in CRY1 that mimics its light-activated state with respect to BIC1 recognition. Collectively, these findings suggest that BIC1 disrupts CRY1 oligomerization by binding individual CRY1 protomers with higher affinity than the protomer-protomer interactions, effectively outcompeting CRY1-CRY1 contacts regardless of the photoreduction state of the chromophore.

BIC1 does not inhibit photoreduction of the CRY1 FAD cofactor. The remaining open question is whether BIC1 suppresses CRY1 oligomerization by interfering with its photoactivation, as previously demonstrated for CRY2 inhibition by BIC2[22]. To address this, we examined the photoreduction state of the FAD chromophore in the presence and absence of BIC1. Absorption spectra of purified CRY1 recorded before and after blue light exposure showed the expected spectral changes associated with FAD photoreduction (Fig. 3f, Extended Data Fig. 1d, e). In the dark-adapted state, CRY1 displayed absorption maxima characteristic of oxidized flavin (FAD_{ox}) at 450 nm and a shoulder near 480 nm. Upon blue light illumination, these peaks decreased in intensity, accompanied by the appearance of a broad band between 550-600 nm, consistent with the formation of the neutral semiquinone radical ($FADH°$)[15,16,38]. These spectral changes are characteristic for the light-driven redox transition of the CRY1 chromophore. Identical spectral features were observed when CRY1 or $CRY1^{L407F}$ were incubated with BIC1 under the same conditions (Fig. 3g, i, Extended Data Fig. 4a-c), indicating that BIC1 does not impair the light-induced redox transition of the FAD_{ox} to $FADH°$. This suggests that BIC1 does not block the initial photoactivation step of CRY1. Instead, it acts downstream of photoreduction, targeting the light-induced oligomeric states of CRY1.

Discussion

Light perception in plants is a dynamic process influenced by multiple molecular factors. Our kinetic analysis of CRY1 reveals that oligomerization proceeds through discrete assembly steps and is modulated by both metabolic signals and inhibitory interactions. These findings suggest that CRY1 activity can be finely adjusted in response to changing conditions, enabling a more nuanced regulation of light signalling.

Our time-resolved native mass spectrometry data establish a clear kinetic pathway for CRY1 oligomerization (Fig. 4). The sequential monomer-to-dimer-to-tetramer assembly we observed is consistent with a two-step reversible kinetic model (Fig. 1l), identifying dimer formation as the rate-limiting step in CRY1 activation. In comparison, FAD photoreduction occurs within microseconds[39], much faster than oligomerization, indicating that the light-induced activation of the flavin cofactor does not contribute any

281　significant kinetic delay. The relative stabilities of the oligomers differ substantially, with dimers being highly transient while tetramers display significantly increased stability. These light dependent equilibria, suggests that CRY1 signalling is a highly dynamic process rather than a simple on-off switch. To directly compare our *in vitro* oligomerization kinetics with *in vivo* observations made by Liu *et al.* (2020)[20], we fitted the tetramer formation data using a single-exponential model and determined a half-life of approximately $t_{1/2} = 18$ s (Extended Data Fig. 2c). *In vivo*, the corresponding half-life at a fluence rate of 30 µmol m^{-2} s^{-1} is about five times longer, reflecting additional cellular constraints such as regulatory interactions (e. g. BIC1 inhibition), metabolites, different reducing environment, temperature and, of course, different protein concentrations. Despite these differences, the timescales remain within the same order of magnitude, underscoring the physiological relevance of the oligomerization process captured by time-resolved mass spectrometry. Importantly, our *in vitro* approach allows disentangling the influence of individual factors that cannot easily be resolved in the cellular context, thereby complementing *in vivo* measurements. Beyond this biological relevance, such a detailed kinetic analysis also opens opportunities for engineering CRY1-based optogenetic tools with tuneable temporal precision, complementing existing CRY2-based optogenetic systems[40–43].

299　In addition to intrinsic kinetic insights, our findings highlight how cellular metabolites can modulate this process. We show that ATP dramatically enhances CRY1 oligomerization upon illumination (Fig. 4). Increasing ATP concentrations significantly enhance the formation of CRY1 dimer (Extended Data Fig. 1a-c) which we showed to be the rate limiting step in oligomer assembly. Then tetramer formation proceeds rapidly, resulting in more stable higher-order oligomers.

305　This is particularly intriguing given that ATP is not only a fundamental energy currency but has also been reported to bind the PHR domain of plant cryptochromes[19,26] and affect FAD photoreduction kinetics[28,30–32]. Our data support the notion that ATP acts as a cofactor, stabilizing the oligomeric state and promoting the conformational changes required for stable oligomer formation. This mechanism suggests a novel link between the cellular energy status and CRY1 activity, allowing a plant to modulate its light response in accordance with its metabolic state. For instance, it is conceivable that under conditions of high energy availability, CRY1 signalling may be sensitized to support energy-intensive processes such as development and growth.

Our data further elucidate the inhibitory role of BIC1, revealing an elegant mechanism
of regulation (Fig. 4). We demonstrate that BIC1 binds to CRY1 in both dark and light
conditions (Fig. 2c, d), but with an enhanced affinity for the light-activated form
(Fig. 2f, g). This light-dependent increase in binding affinity indicates that light-induced
conformational changes in CRY1 significantly promote its interaction with BIC1. This
light-dependent increase in affinity is a crucial finding, as it provides a mechanism for
the selective inhibition of active CRY1. Moreover, we provide clear evidence that BIC1
is an active disassembler of CRY1 oligomers, rather than just a passive competitor
(Fig. 2e). Given that the constitutively active CRY1^{L407F} mutant, which mimics the
tetrameric light-activated state, shows a high affinity for BIC1 (Fig. 3e), strongly
suggests that BIC1 recognizes a specific conformational state of CRY1, independent
of the FAD redox state. Our result that BIC1 does not interfere with FAD
photoreduction (Fig. 3g, i) differs from recent reports on the CRY2-BIC2 interaction[22],
suggesting that the inhibitory mechanisms are distinct for CRY1 and CRY2. This
supports the idea that while cryptochromes share a conserved photochemical
activation step, their downstream regulation can vary, allowing for functional
specificity.

In summary, our work establishes a quantitative kinetic model for CRY1
oligomerization. By showing that ATP acts as an oligomerization accelerator and BIC1
as an active disassembler, we provide a more detailed and nuanced understanding of
how CRY1 activity is regulated beyond simple light perception. These insights not only
shed light on the functional specialization of CRY1 but also highlight a broader
principle of plant signalling, where a photoreceptor's output is shaped by a complex
interplay between light, metabolites, and dedicated regulatory proteins.

Methods

Cloning of constructs. The expression vectors encoding the PHR domain (residues
1–509) of *Arabidopsis thaliana* CRY1 and the CRY1^{L407F} mutant, both cloned into the
pACYCDuet-1 vector have been described previously[26]. To prevent disulfide bond-
mediated dimerization observed in the *Arabidopsis thaliana* wild-type BIC1 protein
(Extended Data Fig. 5a, b), a synthetic gene encoding a BIC1 variant carrying three
cysteine-to-alanine substitutions (C99A, C105A, C136A) was generated. This triple

mutant, named BIC1 throughout the text, was synthesized and cloned into the
pET28a(+) vector (*BioCat GmbH*). Recombinant proteins carried an N-terminal His_6-
tag.

Overproduction and purification of recombinant proteins. For overexpression of
CRY1 and the $CRY1^{L407F}$ mutant as well as $BIC1^{3CA}$, competent *E. coli* BL21 Star
(DE3) cells (*Thermo Fisher*) were used. For CRY1 and $CRY1^{L407F}$ overproduction,
20 mL overnight cultures were inoculated into 2 L LB medium supplemented with
34 µg/mL chloramphenicol in 5 L baffled flasks. Cultures were incubated at 37 °C with
shaking (110 rpm) until an $OD_{600 nm}$ of ~1.0 was reached. Cells were placed on ice for
30 min, and expression was induced with 0.25 mM IPTG. All further steps were carried
out under red light. Cultures were incubated for 3 days at 16 °C. Cells were harvested
by centrifugation (5,330g, 15 min, 4 °C) and resuspended in lysis buffer (50 mM
MOPS, pH 7.4, 300 mM NaCl, 10% glycerol, 1% Triton X-100). Lysozyme (0.2 mg/mL),
PMSF (0.4 mM) and DNase I were added, and the suspensions were incubated for
10 min at room temperature before disruption using a French press. Lysates were
clarified by centrifugation (39,000g, 30 min) and filtered. The supernatant was applied
onto a 1 mL Protino Ni-NTA column (*Macherey-Nagel*), followed by sequential
washing steps with lysis buffer, buffer containing 3% Triton X-100, and lysis buffer
without TX-100 until a stable UV baseline was reached. Non-specifically bound
proteins were removed with 10% elution buffer (50 mM MOPS, pH 7.4, 300 mM NaCl,
10% glycerol, 250 mM imidazole). Recombinant CRYs were eluted with 100% elution
buffer. CRY1 concentrations were kept below 1 mg/mL to prevent oligomerization.
Final purification was performed by size-exclusion chromatography (Superdex 200 pg
16/60, *Cytiva*).

BIC1 purification followed a similar protocol. Expression was induced with 0.5 mM
IPTG, and cultures were incubated for 18 h at 25 °C. Cells were harvested, washed
with PBS, and resuspended in lysis buffer (50 mM HEPES, pH 7.0, 500 mM NaCl, 10%
glycerol). Cell disruption was performed as described above, using a protease inhibitor
mix (*Carl Roth, Cat. no. 3760.1*) in place of PMSF. The soluble fraction was loaded
onto a Ni-NTA column and washed with protein buffer (50 mM HEPES, pH 7.0,
300 mM NaCl, 10% glycerol), followed by a wash containing 3% Triton X-100 and
protein buffer. Additional washes were performed with 10% and 20% elution buffer
(50 mM HEPES, pH 7.0, 300 mM NaCl, 250 mM imidazole, 10% glycerol). BIC1 was

eluted with 100% elution buffer, concentrated, and further purified via size-exclusion

chromatography (Superdex 75 pg 16/60, *Cytiva*).

Size exclusion chromatography (SEC). For analysis of blue light-dependent

oligomerization, samples containing 30 µM CRY1 and 10 mM ß-mercaptoethanol in

100 µl volumes were separated on a Superdex 200 10/300 GL column (*GE*

Healthcare) equilibrated at 10 °C in buffer D containing 20 mM ß-mercaptoethanol.

CRY1 was pre-illuminated with blue light (465 ± 5 nm, 115 µmol m^{-2} s^{-1}) for 20 min.

Three blue light LEDs were used to illuminate the column resin. Dark controls were

kept in red safe light. Calibration of the column was done using gel-filtration standards

(*Sigma Aldrich*).

Photoreduction and dark reversion of CRY1. Absorbance changes of bound flavin

were measured by UV-Vis spectroscopy (*Specord 600, Analytik Jena*). Before

photoreduction, the CRY1 solution was centrifuged (10 min, 16873 g, 4 °C) to pellet

and remove aggregated protein. A protein solution of 10 µM CRY1 was prepared, and

dithiothreitol (DTT) added to a final concentration of 10 mM. For photoreduction,

samples were irradiated with a 455 nm LED (M455L2, *Thorlabs*) at 100 µmol m^{-2} s^{-1}.

At defined time points, the absorption between 200 nm – 900 nm was recorded. The

dark conversion was monitored with the same photoreduced sample by leaving it in

the dark until full recovery and absorbance recording. During all steps, cuvettes were

chilled at 20 °C.

Nano Electrospray ionization mass spectrometry (nESI-MS). nESI-MS was

carried out using a Synapt G2-S instrument (*Waters Corporation, Wilmslow, UK*)

equipped with a high-mass quadrupole modification. Nano electrospray emitters are

produced in-house by pulling borosilicate glass capillaries with a Flaming/Brown

micropipette puller (P-1000, *Sutter Instrument Co.*) and subsequently coated with a

gold layer via sputtering. CRY1 (15 µM) was analyzed at room temperature in positive

ion mode under a capillary voltage of 1.1 kV. The ion source was operated with a cone

voltage of 150 V, an offset of 100 V, and maintained at 20 °C. The scan time was set

to 1 scan per second. Calibration is performed using a standard cesium iodide (CsI)

solution. Sample illumination was carried out by irradiation of the nano electrospray

emitters with a 455 nm LED (M455L4, *Thorlabs*) at 30-50 µmol m^{-2} s^{-1} (Extended Data

Fig. 6a). Acquisition and data processing are achieved using MassLynx™ V4.1

SCN901. Prior to measurements, CRY1 was buffer-exchanged in 200 mM ammonium
acetate, 10 mM DTT and pH 7.4, using Zeba Micro Spinbuffer exchangers (7 kDa
MWCO, *Thermo Fisher Scientific*).

nESI-MS based rate constant determination. The measurements of the rate
constants were performed following the same procedure as for the nESI-MS
measurements except for the scan time, which was set to 10 scans per second. The
intensity (counts) of each species (monomer, dimer, tetramer) during the
measurement period was estimated by generating an extracted ion chromatogram
using MassLynx™ V4.1 SCN901. Initial concentrations of each species (monomer,
dimer, tetramer) were determined from the measured signal intensities, assuming a
total sample concentration of 15 µM relative to the molecular mass of the monomer.
Kinetic data were analyzed by using DynaFit[44] software and a reversible two-step
kinetic model (Fig. 1l) based on the differential equations 1-3.

$$\frac{d[M]}{dt} = -2k_{M \to D}[M][M] + 2k_{D \to M}[D] \tag{1}$$

$$\frac{d[D]}{dt} = +k_{M \to D}[M][M] - k_{D \to M}[D] - 2k_{D \to T}[D][D] + 2k_{T \to D}[T] \tag{2}$$

$$\frac{d[T]}{dt} = +k_{D \to T}[D][D] - k_{T \to D}[T] \tag{3}$$

LILBID-MS. Prior to analysis by LILBID-MS[35], CRY1 and BIC1[3CA] were buffer-
exchanged as described for nESI-MS. All measurements were performed at 20 °C,
using 4 µL of buffer-exchanged and degassed sample for each analysis. Microdroplets
with a diameter of 50 µm are generated using a piezo-driven droplet generator (MD-
K-130, *Microdrop Technologies GmbH,* Norderstedt, Germany) operating at 10 Hz.
The droplets are transferred into a vacuum of about 10^{-5} mbar and irradiated with a
pulsed infrared laser (*Innolas Spitlight 400, Continuum PowerLite 8000*) tuned to the
vibrational absorption maximum of water (2.8 µm). Absorption of laser energy (10-
23 mJ per pulse) by water molecules causes an explosive expansion of the droplets,
resulting in the release of analyte ions into the gas phase. The ions are accelerated
using a pulsed electric field and analysed with a home-built time-of-flight mass
spectrometer in negative ion mode. Each mass spectrum represents an average
signal from 1,000 individual microdroplets. Mass spectra are calibrated against a

10 µM aqueous bovine serum albumin standard and subsequently processed by
smoothing and background subtraction. The sample was exposed to blue light
(455 nm LED, M455L4, *Thorlabs*) at 100 µmol m^{-2} s^{-1} inside the droplet generator via
a coupled optical fibre that was installed on the generator (Extended Data Fig. 6b).

For qLILBID-MS[36,37] experiments the laser energy transferred into the droplets is
correlated to the degree of complex dissociation. Therefore, the explosion of the
irradiated droplets is visualized by illuminating the explosion with a green laser (Minilite
I, *Continuum*, San Jose, USA) and the resulting images are captured using a high-
speed camera (DFK 23UP031, *Imaging Source*, Bremen, Germany). Each individual
droplet yields both a mass spectrum and a corresponding image of the explosion. The
explosion width for each droplet is determined by analysing the recorded images with
OpenCV, using a custom written Python script. The resulting dissociation plots allow
K_D determination.

TEM negative staining. Carbon coated copper grids (400 mesh) were hydrophilized
by glow discharging (*PELCO easiGlow*, Ted Pella, USA). 5 µl of a protein suspension
with a concentration of 30 µg/ml was applied onto the hydrophilized grids and stained
with 2% uranyl acetate after a short washing step with filtered H_2O_{bidest}. If applicable,
samples were prepared under blue (465 ± 5 nm, 115 µmol m^{-2}s^{-1}) or safe red light,
respectively. Samples were analysed with a JEOL JEM-2100 transmission electron
microscope using an acceleration voltage of 120 kV. For image acquisition a F214
FastScan CCD camera (*TVIPS, Gauting*) was used.

Data Availability

461   All data generated in this study are provided as Source Data with this paper. Gene
462   accessions codes are At3G52740 (*BIC1*) and At4G08920 (*CRY1*).

References

1. Franklin, K. A., Larner, V. S. & Whitelam, G. C. The signal transducing

photoreceptors of plants. *Int. J. Dev. Biol.* 49, 653–664 (2005).

2. Halliday, K. J., Martínez-García, J. F. & Josse, E.-M. Integration of Light and Auxin

Signaling. *Cold Spring Harb. Perspect. Biol.* 1, a001586 (2009).

3. Fankhauser, C. The Phytochromes, a Family of Red/Far-red Absorbing

Photoreceptors *. *J. Biol. Chem.* 276, 11453–11456 (2001).

4. Christie, J. M. & Briggs, W. R. Blue Light Sensing in Higher Plants *. *J. Biol.*

Chem. 276, 11457–11460 (2001).

5. Yadav, A. *et al.* Light signaling and UV-B-mediated plant growth regulation. *J.*

Integr. Plant Biol. 62, 1270–1292 (2020).

6. Lin, C. *et al.* Association of Flavin Adenine Dinucleotide with the *Arabidopsis* Blue

Light Receptor CRY1. *Science* 269, 968–970 (1995).

7. Malhotra, K., Kim, S.-T., Batschauer, A., Dawut, L. & Sancar, A. Putative Blue-

Light Photoreceptors from Arabidopsis thaliana and Sinapis alba with a High

Degree of Sequence Homology to DNA Photolyase Contain the Two Photolyase

Cofactors but Lack DNA Repair Activity. *Biochemistry* 34, 6892–6899 (1995).

8. Brautigam, C. A. *et al.* Structure of the photolyase-like domain of cryptochrome 1

from Arabidopsis thaliana. *Proc. Natl. Acad. Sci.* 101, 12142–12147 (2004).

9. Wang, X. & Lin, C. The two action mechanisms of plant cryptochromes. *Trends*

Plant Sci. 30, 775–791 (2025).

10.   Yang, Z. *et al.* Cryptochromes Orchestrate Transcription Regulation of

Diverse Blue Light Responses in Plants. *Photochem. Photobiol.* 93, 112–127

(2017).

11.   Ahmad, M. & Cashmore, A. R. HY4 gene of A. thaliana encodes a protein with

characteristics of a blue-light photoreceptor. *Nature* 366, 162–166 (1993).

12.   Guo, H., Yang, H., Mockler, T. C. & Lin, C. Regulation of Flowering Time by

Arabidopsis Photoreceptors. *Science* 279, 1360–1363 (1998).

13.   Wang, J., Du, X., Pan, W., Wang, X. & Wu, W. Photoactivation of the

cryptochrome/photolyase superfamily. *J. Photochem. Photobiol. C Photochem.*

Rev. 22, 84–102 (2015).

14.   Wang, Q. & Lin, C. Mechanisms of Cryptochrome-Mediated Photoresponses

in Plants. *Annu. Rev. Plant Biol.* 71, 103–129 (2020).

15.   Bouly, J.-P. *et al.* Cryptochrome Blue Light Photoreceptors Are Activated

through Interconversion of Flavin Redox States*. *J. Biol. Chem.* 282, 9383–9391

(2007).

16.   Banerjee, R. *et al.* The Signaling State of Arabidopsis Cryptochrome 2

Contains Flavin Semiquinone *. *J. Biol. Chem.* 282, 14916–14922 (2007).

17.   Sang, Y. *et al.* N-Terminal Domain–Mediated Homodimerization Is Required

for Photoreceptor Activity of Arabidopsis CRYPTOCHROME 1. *Plant Cell* 17,

1569–1584 (2005).

18.   Rosenfeldt, G., Viana, R. M., Mootz, H. D., von Arnim, A. G. & Batschauer, A.

Chemically Induced and Light-Independent Cryptochrome Photoreceptor

Activation. *Mol. Plant* 1, 4–14 (2008).

19.   Wang, Q. *et al.* Photoactivation and inactivation of Arabidopsis cryptochrome

2. *Science* 354, 343–347 (2016).

20.   Liu, Q. *et al.* Photooligomerization Determines Photosensitivity and

Photoreactivity of Plant Cryptochromes. *Mol. Plant* 13, 398–413 (2020).

21.   Ponnu, J. & Hoecker, U. Signaling Mechanisms by Arabidopsis

Cryptochromes. *Front. Plant Sci.* 13, 844714 (2022).

22.   Ma, L. *et al.* Structural insights into BIC-mediated inactivation of Arabidopsis

cryptochrome 2. *Nat. Struct. Mol. Biol.* 27, 472–479 (2020).

516 23. Palayam, M. *et al.* Structural insights into photoactivation of plant

517   Cryptochrome-2. *Commun. Biol.* 4, 28 (2021).

518 24. Shao, K. *et al.* The oligomeric structures of plant cryptochromes. *Nat. Struct.*

519   *Mol. Biol.* 27, 480–488 (2020).

520 25. Wang, X. *et al.* A CRY–BIC negative-feedback circuitry regulating blue light

521   sensitivity of Arabidopsis. *Plant J. Cell Mol. Biol.* 92, 426–436 (2017).

522 26. Miao, L. *et al.* Arabidopsis cryptochrome 1 undergoes COP1 and LRBs-

523   dependent degradation in response to high blue light. *New Phytol.* 234, 1347–

524   1362 (2022).

525 27. El-Esawi, M. *et al.* Cellular metabolites modulate in vivo signaling of

526   Arabidopsis cryptochrome-1. *Plant Signal. Behav.* 10, e1063758 (2015).

527 28. Engelhard, C. *et al.* Cellular metabolites enhance the light sensitivity of

528   Arabidopsis cryptochrome through alternate electron transfer pathways. *Plant Cell*

529   26, 4519–4531 (2014).

530 29. Zhao, X. *et al.* A Study of Gibberellin Homeostasis and Cryptochrome-

531   Mediated Blue Light Inhibition of Hypocotyl Elongation. *Plant Physiol.* 145, 106–

532   118 (2007).

533 30. Immeln, D., Schlesinger, R., Heberle, J. & Kottke, T. Blue Light Induces

534   Radical Formation and Autophosphorylation in the Light-sensitive Domain of

535   *Chlamydomonas* Cryptochrome*. *J. Biol. Chem.* 282, 21720–21728 (2007).

536 31. Müller, P. *et al.* ATP Binding Turns Plant Cryptochrome Into an Efficient

537   Natural Photoswitch. *Sci. Rep.* 4, 5175 (2014).

538 32. Eckel, M., Steinchen, W. & Batschauer, A. ATP boosts lit state formation and

539   activity of Arabidopsis cryptochrome 2. *Plant J.* 96, 389–403 (2018).

33.   Exner, V. *et al.* A Gain-of-Function Mutation of Arabidopsis

CRYPTOCHROME1 Promotes Flowering. *Plant Physiol.* 154, 1633–1645 (2010).

34.   Orth, C., Niemann, N., Hennig, L., Essen, L.-O. & Batschauer, A. Hyperactivity

of the Arabidopsis cryptochrome (cry1) L407F mutant is caused by a structural

alteration close to the cry1 ATP-binding site. *J. Biol. Chem.* 292, 12906–12920

(2017).

35.   Peetz, O. *et al.* LILBID and nESI: Different Native Mass Spectrometry

Techniques as Tools in Structural Biology. *J. Am. Soc. Mass Spectrom.* 30, 181–

191 (2019).

36.   Schulte, J., Tants, J.-N., von Ehr, J., Schlundt, A. & Morgner, N.

Determination of dissociation constants via quantitative mass spectrometry. *Front.*

Anal. Sci. 3, (2023).

37.   Young, P., Hense, G., Immer, C., Wöhnert, J. & Morgner, N. LILBID laser

dissociation curves: a mass spectrometry-based method for the quantitative

assessment of dsDNA binding affinities. *Sci. Rep.* 10, 20398 (2020).

38.   Liu, H., Zhong, D. & Lin, C. Searching for a photocycle of the cryptochrome

photoreceptors. *Curr. Opin. Plant Biol.* 13, 578–586 (2010).

39.   Langenbacher, T., Immeln, D., Dick, B. & Kottke, T. Microsecond Light-

Induced Proton Transfer to Flavin in the Blue Light Sensor Plant Cryptochrome. *J.*

Am. Chem. Soc. 131, 14274–14280 (2009).

40.   Taslimi, A. *et al.* An optimized optogenetic clustering tool for probing protein

interaction and function. *Nat. Commun.* 5, 4925 (2014).

41.   Che, D. L., Duan, L., Zhang, K. & Cui, B. The Dual Characteristics of Light-

Induced Cryptochrome 2, Homo-oligomerization and Heterodimerization, for

Optogenetic Manipulation in Mammalian Cells. *ACS Synth. Biol.* 4, 1124–1135

(2015).

42.   Park, H. *et al.* Optogenetic protein clustering through fluorescent protein

tagging and extension of CRY2. *Nat. Commun.* 8, 30 (2017).

43.   Hallett, R. A., Zimmerman, S. P., Yumerefendi, H., Bear, J. E. & Kuhlman, B.

Correlating in Vitro and in Vivo Activities of Light-Inducible Dimers: A Cellular

Optogenetics Guide. *ACS Synth. Biol.* 5, 53–64 (2016).

44.   Kuzmic, P. Program DYNAFIT for the analysis of enzyme kinetic data:

application to HIV proteinase. *Anal. Biochem.* 237, 260–273 (1996).

Acknowledgements

We thank Konstantinos Stamatakis for his support and fruitful discussions during the kinetic modelling process, and Josef Wachtveitl for providing access to the absorption spectrometer. We also acknowledge Jeanette Schermuly for technical assistance and Stephan Kiontke for his contributions to the recombinant expression of BIC1 and for helpful discussions. Funding was provided by the German Research Foundation (DFG): A.J. was supported through the Collaborative Research Center CRC 1507 "Membrane-Associated Protein Assemblies, Machineries and Supercomplexes", N.M. through DFG Grant 557111829, L.-O.E. through DFG grant ES152/18 and A.B. through DFG grant BA985/15-1.

Author contributions

A.J., N.M., L.-O.E., A.B. designed and coordinated the experiments. P.G., A.J., N.N. and D.K. expressed and purified the proteins. A.J. performed mass spectrometry, UV-Vis spectroscopy and kinetic modelling. N.N. and T.H. acquired and interpreted the TEM data. N.N. acquired and interpreted the SEC data. A.J. wrote the original draft. A.J., N.M., A.B., P.G., L.-O.E. edited the manuscript.

Competing interests

The authors declare no competing interests.

Figures and Legends

Fig. 1: Oligomerization kinetics of CRY1 under blue light and the influence of ATP. nESI-MS spectrum of
CRY1 in the dark (grey line) (**a, c**) and under illumination with blue light (blue line) (**b, d**) in absence (**a, b**) and
presence of 100 µM ATP (**c, d**). Exposure to blue light leads to oligomerization of CRY1, and the addition of ATP
promotes this process. Time-resolved mass spectrometry measurements of the CRY1 monomer (**e**), dimer (**f**) and
tetramer (**g**) in absence and presence of ATP (**h-j**) and the fit (pink line) obtained from the two-step reversible

Fig. 2: BIC1 inhibits CRY1 oligomerization in both dark and blue light conditions with higher affinity of BIC1 to CRY1 under blue light conditions. LILBID-MS spectrum of CRY1 shows a monomer (pink dot) (**a**) in darkness, which forms tetramers upon illumination with blue light (**b**). CRY1 trimers are caused by laser dissociation. **c,** Incubation of CRY1 monomer and BIC1 (purple dot) in an equimolar ratio in darkness results in a mixture of CRY1-BIC1 heterodimer and CRY1 monomer. **d,** Upon blue light exposure, the equilibrium shifts towards a higher proportion of the CRY1-BIC1 heterodimer relative to the CRY1 monomer (highlighted in grey). **e,** Under blue light conditions, addition of BIC1 to the CRY1 tetramer results in binding to each protomer, thereby dissociating the tetramer into CRY1-BIC1 heterodimers. **f,** qLILBID-MS measurements reveal that the CRY1-BIC1 heterodimer exhibits a ten-fold higher affinity under blue light compared to its dark-state counterpart (**g**). TEM images of CRY1 particles visualised by negative staining of probes kept in darkness (**h**), or treated with blue light (450 nm, 20 min, 115 µmol m^{-2} s^{-1}) (**i**) showing light-induced formation of doughnut-like ring structures of CRY1. Scale bars 10 nm. **j,** Kinetics of CRY1 monomerization (n = 3 ± SD) after transfer of samples from blue light to darkness in absence and presence of BIC1 or *E. coli* GroEL as negative control. **k,** Effect of BIC1 on kinetics of decrease in number of

CRY1 particles showing monomerization of CRY1 (n = 3 ± SD). Blue light (465 nm, fluence rate 115 µmol m^{-2} s^{-1})
treatment was for 20 min.

Fig. 3: BIC1 disassembles constitutively active CRY1 tetramers and does not inhibit photoreduction of the
FAD cofactor. a, LILBID-MS spectrum of the constitutively active CRY1^{L407F} mutant shows a tetramer in darkness.
CRY1 trimers are caused by laser dissociation. **b**, Addition of BIC1 (purple dot) to CRY1^{L407F} in an equimolar ratio
in darkness results in binding to each protomer, thereby dissociating the CRY1^{L407F} tetramer into CRY1^{L407F}-BIC1
heterodimer and monomeric CRY1^{L407F}. **c**, Exposure to blue light does not shift the equilibrium toward the
CRY1^{L407F}-BIC1 heterodimer, as has been shown for wild-type CRY1. **d**, SEC elution profiles of wild-type CRY1
kept in darkness or irradiated with blue light, and of hyperactive CRY1^{L407F} mutant kept in darkness. **e**, qLILBID-
MS measurement of the CRY1^{L407F}-BIC1 heterodimer under dark conditions revealed that BIC1 binds CRY1^{L407F}
with high affinity. **f**, Absorption spectra of wild-type CRY1 before (grey line) and after (blue line) blue light exposure
indicate FAD photoreduction, with a decrease in FAD$_{ox}$ peak at 450 nm, and the emergence of a broad band at
550-600 nm, characteristic for the FADH° semiquinone. **g**, Addition of BIC1 to wild-type CRY1 does not inhibit the
photoreduction of CRY's FAD chromophore. **h**, Absorption spectra of the CRY1^{L407F} mutant indicates that
chromophore photoreduction remains unaffected by the mutation. **i**, Photoreduction of the FAD chromophore in
the CRY1^{L407F} mutant remains unaffected by BIC1 addition, as observed for wild-type CRY1.

Fig.4: Schematic model of CRY1 oligomerization dynamics, ATP modulation, and BIC1-mediated inhibition. Upon blue light exposure, CRY1 transitions from a monomeric to a tetrameric state via a short-lived dimeric intermediate. This oligomerization follows a sequential, reversible two-step mechanism. ATP enhances the association rates for dimer and tetramer formation, while slowing down the dissociation rates of these oligomers. BIC1 binds CRY1 in both dark and light conditions, with significantly higher affinity for the light-activated conformation. This interaction prevents oligomer formation and actively disassembles pre-formed tetramers by binding to individual CRY1 protomers even under continuous blue-light illumination.