PHYTOCHROME INTERACTING FACTOR 7 moderates the activity and the cytosolic-

nuclear shuttling of PHYTOCHROME A

Mengke Zhou[1*], Andrés Romanowski[2], Karen J. Halliday[1*]

1. Institute for Molecular Plant Sciences, School of Biological Sciences, University of

Edinburgh, Edinburgh EH9 3BF, United Kingdom.

2. Timing of Environmental Signaling lab, Molecular Biology Group, Plant Sciences,

Wageningen University & Research, Wageningen, The Netherlands.

ORCID IDs: 0000-0003-1043-2766 (MK. Z); 0000-0003-0737-2408 (A.R.); 0000-0003-0467-

104X (K.J.H.).

*Corresponding authors:

Mengke Zhou and Karen J. Halliday

Institute for Molecular Plant Sciences, School of Biological Sciences, University of Edinburgh,

Edinburgh EH9 3BF, United Kingdom

+44-0131-651-9083

mengke.zhou@ed.ac.uk

karen.halliday@ed.ac.uk

ABSTRACT

In nature, plants sense and navigate complex vegetative environments using the phytochrome (phy) photoreceptor-signaling system. The canopy adaptation strategy, elicited by phyA, and the shade avoidance syndrome, regulated by phyB, allow plants to adapt to diverse forms of vegetation shade. The phyA High Irradiance Response (HIR) mode, involving photocycle-coupled nuclear shuttling, is essential for detecting persistent far-red rich conditions typical of canopy or occlusive shade. Our research identifies PHYTOCHROME INTERACTION FACTOR 7 (PIF7) as a critical regulator of phyA function. We demonstrate PIF7ox can effectively block phyA action. In persistent shade, PIF7 and phyA both accumulate and interact within nuclear bodies, with binding facilitated through the prion-like domain of PIF7 and the C-terminal PAS-PAS domain of phyA. PIF7 also interacts with FHY1 and FHL chaperones, interfering with phyA cytosolic-nuclear shuttling. This strategic molecular targeting by PIF7 provides an effective strategy for modulating the phyA HIR in response to shade signals.

INTRODUCTION

Light sensing is fundamental for plant adaptation and survival in vegetative shade. This is particularly important during seedling establishment, a stage when plants are most vulnerable. Plants have evolved distinct developmental strategies to cope with different types of shade that are prevalent in nature. The light receptor phytochrome B (phyB), is able to detect FR-rich light from neighboring plants and elicit the Shade Avoidance Syndrome (SAS), which induces elongation of seedling hypocotyls, repositions leaves to improve light capture, and triggers early flowering[1,2]. Under persistent canopy shade, plants adopt alternative strategies, such as the phyA-mediated Canopy Adaptation Strategy (CAS), which suppresses hypocotyl elongation and conserves resources, optimizing growth when shade is unyielding[3]. As vegetative habitats can change considerably through a season, the ability to adjust photoreceptor activity offers adaptive flexibility, ensuring successful seedling establishment and progression through the life cycle.

In Arabidopsis, phytochromes A-E, form a family of photoreceptors that principally absorb light in the red (R) and far-red (FR) regions of the spectrum[4]. Absorption of R photoconverts phytochrome into its active Pfr form, while FR photoconverts Pfr to the inactive Pr conformer.

When activated phytochromes move from the cytosol to the nucleus where they aggregate in

sub-nuclear photobodies to perform a range of functions[5,6]. While all phytochromes possess

similar photosensory properties, they fall into two classes based on their distinct modes of

action. Phytochromes B-E, which operate in the low fluence response (LFR) mode, are

activated by R and deactivated by FR light. Consequently, in FR-rich vegetation shade

environments, these phytochromes are less active[4,7]. A key distinction of phyA is that it is

activated by FR-rich light through its high irradiance response (HIR) mode of action[7]. This

property arises partly from its photobiology and partly from the shuttling mechanism that

concentrates phyA in the nucleus[8]. PhyA lacks a nuclear localization signal and primarily relies

on the chaperone proteins FHY1/FHL to facilitate its transport into the nucleus. These proteins

specifically recognize phyA-Pfr and transport it to the nucleus, where it is converted into phyA-

Pr, allowing FHY1/FHL to dissociate and be recycled to the cytoplasm for further shuttling[8,9,10].

Nuclear located phyA regulates a number of processes, including expression of target genes,

through association with transcription factors such as PHYTOCHROME INTERACTING

FACTORS (PIFS) and ELONGATED HYPOCOTYL 5 (HY5)[11]. The central role of

FHY1/FHL shuttling in the phyA HIR, identifies this mechanism as a potential target for

modifying phyA action and the adaptive response[8]. Indeed, HY5 has been reported to be

involved in negative-feedback regulation of phyA signaling through repressing FHY1/FHL

expression[11]. Here, HY5 directly interacts with the DNA-binding regions of FAR-RED

ELONGATED HYPOCOTYLS 3 (FHY3) and FAR-RED IMPAIRED RESPONSE 1 (FAR1)

to prevent their transcriptional activation of FHY1/FHL.

PIF transcription factors are well-studied negative regulators of phytochrome signaling. They

function by directly binding to G/PBE-boxes in the promoters of their target genes[12]. In

Arabidopsis, there are eight PIFs (PIF1, PIF2/PIL1-PIF8), each with an active phytochrome B

(APB) domain that facilitates their interaction with phyB. Only PIF1 and PIF3 contain an active

phytochrome A (APA) domain, which directly binds the C-terminus of phyA, comprising two

Period-Arnt-SIM (PAS) and a histidine kinase-related (HKRD) domains[13]. In FR light, phyA

directly interacts with PIF1 and PIF3, triggering their proteolytic destruction and/or

sequestration from target promoters[13,14]. Although PIF4, PIF5, PIF8 and PIF7 lack an APA

domain, they have been shown to control aspects of phyA signaling[15,16,17]. Interestingly, levels

of PIF8 protein are boosted by phyA. In darkness, CONSTITUTIVE

PHOTOMORPHOGENESIS 1 (COP1) promotes PIF8 protein degradation through the 26S

proteasome, thereby reducing its abundance. Conversely, under FR light, phyA inhibition of

COP1 stabilizes the PIF8 protein, forming a negative feedback loop that limits phyA action[15].

Notably, PIF7, a close homologue of PIF8, is a potent regulator of SAS. In unshaded conditions,

PIF7 interacts with active phyB (Pfr) through its APB domain, and both proteins localize to

sub-nuclear photobodies. This interaction prevents PIF7 from binding to the promoters of target

genes, effectively inhibiting PIF7-regulated gene expression[18,19]. Under shade conditions, the

deactivation of phyB releases PIF7 from this repression, leading to the activation of shade-

responsive genes and triggering the physiological SAS response[20,21]. Furthermore, exposure to

shade triggers the dephosphorylation and nuclear accumulation of PIF7, processes that enhance

the activation of SAS[20]. Given its strong promotory role in SAS, it is notable that PIF7 has

recently been demonstrated to increase *PHYA* transcript abundance, an action that could

potentially limit SAS. This PIF7 action can be inhibited by *PHYA UTR ANTISENSE RNA*

(PUAR), which binds to PIF7 and prevents its interaction with *PHYA's* 5' UTR, thereby reducing

PHYA transcript levels[17]. Collectively, this research underscores the importance of PIF7 as a

key agent in executing and moderating the molecular response to shade.

Here, we show that PIF7 modulates phyA signaling under vegetative shade by controlling phyA

subcellular localization and activity. PIF7 overexpression (PIF7ox) abolishes phyA-mediated

suppression of seedling hypocotyl elongation under persistent shade. Under these conditions,

PIF7 and phyA co-localize in nuclear photobodies and interact directly via the PIF7 prion-like

domain and the phyA C-terminus. Additionally, our data shows that PIF7 governs phyA

subcellular localization by modulating FHY1/FHL-dependent nuclear import. These insights

clarify how PIF7 effectively restricts both the nuclear transport and signaling activities of phyA,

illuminating the intricate molecular mechanisms at play.

RESULTS

PIF7ox blocks phyA action

PIF7, which functions downstream of phyB, also shares significant sequence similarity with PIF8, a regulator of phyA responses. This similarity led us to investigate the potential genetic interaction between PIF7 and phyA. To achieve this, we measured the hypocotyl length of seedlings grown under two conditions: 8h white light (WL): 16h dark (D), and 8h low R:FR ratio (0.2) light (WLFR). Both photoperiods started one day after germination to strongly elicit a phyA response (Fig. 1a). As expected, wild type (WT, Col-0) seedlings were slightly shorter in WLFR compared to WL, while *phyA-211* had longer hypocotyls in WLFR due to reduced control of shade-avoidance elongation (Fig. 1b-c). In contrast, the phyAox line displayed a constitutively short hypocotyl and retained WLFR-responsivity. In WL, the *pif7-1, pif7-2* loss-of-function alleles were marginally shorter than the WT but retained a response to WLFR that was absent in the *pif7-1*; *phyA-211* double mutant. In keeping with PIF7's function in promoting the phyB-mediated shade response, under WLFR, *pif7-1* diminishes the impact of *phyA-211* on hypocotyl length. Conversely, hypocotyls of *p35S::PIF7-FLASH* (PIF7ox) seedlings were elongated in WL and exhibited further elongation under WLFR. As hypocotyl lengths were similar in PIF7ox and PIF7ox;*phyA-211,* this suggests PIF7ox facilitates increased shade-avoidance elongation while simultaneously suppressing phyA action (Fig. 1c)[18]. This observation infers that PIF7 may participate in phyB to phyA cross-regulation.

Given the capacity of PIF7ox to inhibit phyA activity, we sought to investigate the possibility of functional redundancy with other PIFs. As anticipated, PIF8ox also lacked WLFR-mediated growth inhibition, but PIF3ox, PIF4ox and PIF5ox all retained this response (Fig. S1a-b)[15]. Our data therefore illustrate that even though PIF3, PIF4 and PIF5 have been implicated in *PHYA* regulation and signaling, their functions diverge from PIF7 and PIF8[16,22,23]. We found the hypocotyl lengths of *pif8-1* and *pif8-1;phyA-211* were qualitatively similar to *pif7-1, pif7-2*, and *pif7-1;phyA-211*, respectively (Fig. S1c). However, comparison of the *PIF7ox;phyA-211* and *PIF8ox;phyA-211* lines reveals distinct differences in PIF7 and PIF8. While PIF7ox is epistatic to *phyA-211*, this genetic relationship shifts for PIF8, particularly under WLFR, where we show *phyA-211* is epistatic to PIF8ox. This finding aligns well with evidence showing PIF8 functions

downstream of phyA[15].

These results show PIF7 has a distinct role in antagonizing the suppression of hypocotyl growth

by phyA, a phenomenon particularly evident in the PIF7ox lines, which effectively block this

response.

PIF7 moderates phyA protein abundance

PIF7 is known to promote shade induction of *PHYA* expression by binding to its 5' UTR[17].

Concurring with this study, we found that when sampled one hour before dusk (ZT7) in an

8L:16D photoperiod, WLFR enhanced *PHYA* gene expression in the WT, whereas this response

was diminished in the *pif7-1* mutant. Likewise, PIF7ox exhibited elevated *PHYA* levels in

control conditions, which were further enhanced in WLFR (Fig. 1d). Immunoblot analysis

using an antibody against native phyA, reported phyA levels to be consistently low in WL,

across all genotypes. We observed that WLFR led to an increase in phyA protein levels in WT,

pif7-1 and PIF7ox, which qualitatively corresponded with the patterns seen in *PHYA* mRNA

data (Fig. 1d-f). Further, we found that PIF7ox had a negligible effect on the *pPHYA::PHYA-*

nLUC (phyA-nLUC) in WL, which is known to gradually deplete under these conditions. In-

line previous studies, we observed higher levels of daytime phyA-nLUC in WLFR, which was

notably elevated by PIF7ox (Fig. 1g)[3]. Collectively, these findings demonstrate that under

WLFR conditions, PIF7 regulates *PHYA* expression and the abundance of phyA protein.

Interestingly, despite PIF7ox accumulating significantly higher levels of phyA under WLFR,

PIF7ox seedlings fail to elicit a phyA response. This indicates that the regulatory effect of PIF7

on phyA activity does not stem from changes in phyA abundance.

To examine the *in vivo* dynamics of PIF7 under our experimental conditions, we generated

pPIF7::PIF7-nLUC (PIF7-nLUC) lines. We discovered that PIF7-nLUC exhibits rhythmic

patterns, with higher levels during the day compared to the night (Fig. S2a). WLFR boosted

daytime levels of PIF7-nLUC levels, with the *phyA-211* mutation elevating PIF7-nLUC toward

the end of the period, especially in WLFR (Fig. S2b). Therefore, our data indicated that similar

to phyA, WLFR increases the abundance of PIF7, and that phyA plays a minor role in reducing

PIF7 production toward the end of the photoperiod.

PIF7 can interact with phyA in nuclear photobodies

To establish whether PIF7 and phyA interact *in planta* we first used *p35S::PIF7-sfGFP* (PIF7-

GFP) and *p35S::PHYA-RFP* (phyA-RFP) transiently expressed in *N. benthamiana* leaves to

establish whether PIF7 are found in the same cellular location. We found that following WLFR

treatment PIF7-GFP and phyA-RFP co-localize in nuclei and in sub-nuclear foci, or

photobodies (Fig. 2a). Quantitative fluorescence measurements along a nuclear transect,

intersecting two photobodies, clearly show this co-localization (Fig. 2a). Next, we created split

bimolecular fluorescence complementation (BiFC) constructs *p35S::PIF7-nYFP* (PIF7-nYFP)

and *p35S::PHYA-cYFP* (phyA-cYFP*)*, to test whether PIF7 and phyA physically interact *in vivo*.

In WLFR, BiFC assays yielded strong nuclear and sub-nuclear foci signals, but not in control

lines expressing *p35S::PIF7-nYFP* and *p35S::EV-cYFP*. The data suggest PIF7 and phyA are

indeed able to interact in nuclei and within photobodies under WLFR (Fig. 2b).

To gain further evidence for this interaction we also generated split nanoLUC (nLUC) luciferase

reporter constructs, *p35S::PIF7-lgbitnLUC* (PIF7-lgbit) and *p35S::PHYA-smbitnLUC* (phyA-

smbit), with *p35S::PHYB-smbitnLUC* (phyB-smbit) and *p35S::PHYA-nLUC* (phyA-nLUC) as

controls (Fig. 2c). As expected, we found a strong signal in the PIF7-lgbit+phyB-smbit in WL

which is diminished in WLFR, which is consistent with the known light conditional interaction

between PIF7 and phyB[18]. Again, as expected we observed a strong signal from phyA-nLUC

in WLFR, which is known to increase phyA abundance, while negative controls (phyA-smbit+

phyB-smbit) had low signal in both conditions. In WL conditions, the signal from PIF7-lgbit +

phyA-smbit was low, similar to that of the negative control, whereas in WLFR conditions, the

signal was strong, indicating an interaction. Further, using an antibody to native phyA we

demonstrated co-immunoprecipitation of phyA with PIF7ox, in WLFR but not WL conditions

(Fig. 2d). Thus, our data provides evidence that PIF7 and phyA directly interact within

photobodies after exposure to WLFR, which promotes their respective nuclear

accumulation[20,24].

187   **Prion-like Domain of PIF7 interacts with phyA PAS-PAS Domain**

188   Earlier work demonstrated that phyA binds directly with PIF1 and PIF3. The phyA C-terminal

189   output module (OPM) is known to be important for this interaction, as is the active phyA-

190   binding domains (APA) of PIF1 and PIF3[13,22]. To identify PHYA regions that required for the

191   molecular interaction we utilized *p35S::PIF7-nYFP* (PIF7-nYFP) and *p35S::PHYA-cYFP*

192   (phyA-cYFP) split YFP assay, creating truncated PHYA constructs as follows: C-terminal

193   (PHYA-C), comprising HINGE (H), PAS-PAS DOMAIN (PAS), and HISTIDINE KINASE

194   RELATED DOMAIN (HKRD) regions [*H+PAS+HKRD-cYFP*], PHYA-C1 [*PAS+HKRD-*

195   *cYFP*], PHYA-P [*PAS-cYFP*], and PHYA-H [*HKRD-cYFP*] (Fig. 3a). In WLFR we obtained

196   nuclear fluorescence signals for PIF7-nYFP and the full-length phyA-cYFP control, as well as

197   the truncated PHYA-C, PHYA-C1, and PHYA-P constructs. No signal was observed for PHYA-

198   H and the negative EV control. The strong signal detected for PHYA-P, which includes PAS-

199   PAS domain motifs, highlights the significance of this molecular region in facilitating the

200   interaction with PIF7. To identify the region of PIF7 that mediated the interaction with phyA,

201   we generated PIF7-N which contains the ACTIVE PHYTOCHROME BINDING domain

202   [*PIF7-N-nYFP*], PIF7-B, containing the BASIC HELIX-LOOP-HELIX motif [*PIF7-B-nYFP*],

203   and PIF7-C which contains a Prion-like domain (PrLD) [*PIF7-C-nYFP*] which has a polyQ-

204   rich sequence is located at PrLD center (Fig. 3b, Fig. S6). Interestingly we found that only the

205   PIF7 full length control and PIF7-C gave nuclear fluorescence in WLFR. Thus, our data point

206   to the prion-like domain of PIF7 as being important for its interaction with phyA. Since PIF7

207   lacks an APA domain, it seems that prion region is able to fulfill this role.

208   **PIF7 regulates the subcellular localization of phyA**

209   Our data show PIF7ox elevates phyA protein levels, yet PIF7ox lines are unable to mount a

210   phyA response in WLFR conditions. Given that WLFR stimulates the phyA HIR and its nuclear

211   accumulation, we sought to analyze the subcellular localization more closely. As anticipated,

212   confocal microscopy revealed strong cytosolic and weaker nuclear signals for phyA-GFP

213   (*pPHYA::PHYA-sGFP*), and for GFP-PIF7 (*p35S::GFP-PIF7*) when in WL (Fig. S3). Exposure

214   to WLFR notably enhanced nuclear relative to cytosolic signals, confirming the nuclear

accumulation of both PIF7 and phyA under our shade conditions (Fig. S3a)[19,20,25]. We then

aimed to establish if PIF7 affected the subcellular localization of phyA by introgressing phyA-

GFP into *pif7-1* and PIF7ox backgrounds. In WL, we found phyA-GFP to be predominantly in

the cytosol in *Col-0*, *pif7-1* and PIF7ox lines (Fig. 4a). Under WLFR, we recorded a higher

proportion of nuclear phyA-GFP in WT, with an even greater increase in the *pif7-1* mutant,

whereas significantly lower levels were detected in PIF7ox (Fig. 4b-c). Thus, PIF7 appears to

influence the cytoplasmic to nuclear partitioning of phyA-GFP. To verify our results, we

performed immunoblot assays that differentiated between nuclear and cytosolic fractions. This

method yielded qualitatively similar results showing increased nuclear phyA in *pif7-1* and

predominately cytosolic PHYA in PIF7ox (Fig. 4d-e). Thus, our findings indicate that PIF7

modifies subcellular phyA localization. The notably reduced nuclear presence of phyA in

PIF7ox could partially account for the disrupted phyA signaling observed in PIF7ox.

PIF7ox blocks phyA regulation of target genes

To further interrogate the mechanism through which PIF7 inhibits phyA action we initially

tested whether PIF7 alters the expression of genes (*CHS, BIN2, FHY3, FHY1*, *FHL* and *IAA29)*

known to be directly regulated by phyA (Fig. 5a)[26,27]. As determined by qPCR all the genes,

except *IAA29*, were upregulated in WT in WLFR conditions, otherwise, the expression patterns

fell into two categories. In WLFR conditions, *CHS, BIN2,* and *FHY3* exhibited decreased

expression in both *phyA-211* and PIF7ox, while their expression was increased in *pif7-1*. This

gene set is upregulated by phyA in WLFR, and their expression is repressed by PIF7. On the

other hand, *FHY1, FHYL* and *IAA29* form a group that are upregulated in *phyA-211* and PIF7ox,

and for *FHY1* and *FHYL*, expression is lowered in *pif7-1*. These genes are downregulated by

phyA in WLFR, and their expression is promoted by PIF7. The common factor for all these

genes is that phyA and PIF7 exert opposing control of their regulation, particularly in WLFR.

We then sought to establish if the antagonistic action of PIF7 was mediated through direct

repression of phyA at target gene promoters. Here we performed chromatin

immunoprecipitation (ChIP) using a native phyA antibody, and qPCR using to detect phyA

enrichment at *CHS, FHY1*, and *FHL* promoters (Fig. 5b). For all three genes we found WLFR

stimulated phyA enrichment at promoter regions sequences containing G-box motifs (two in

CHS), but not in G-box-free regions (Control, coding region). This enrichment was eliminated

in the presence of PIF7ox, suggesting the interaction between PIF7 and phyA is able to block

phyA regulation of target genes.

PIF7 interacts with FHY1 and FHL

In WLFR phyA cytosolic to nuclear shuttling is controlled by FHY1 and FHL chaperones[9]. The

central importance of this mechanism for phyA function is evident in the *fhy1-3;fhl-1* double

mutant hypocotyl response which is phenotypically similar to *phyA-211* (Fig. S4). An intriguing

finding of our study is that PIF7 appears to control the subcellular localization of phyA. We

therefore wanted to establish if this arose through a process involving PIF7 interaction with

FHY1 and FHL. Using the *N. benthamiana* transient expression system, we recorded nuclear

interactions between the BiFC constructs *p35S::FHY1-cYFP*, *p35S::FHL-cYFP*, and

p35S::PIF7-nYFP under WLFR conditions (Fig. 6a). We also observed PIF7 interactions with

FHY1 and FHL, but not the negative control using a split nLUC assay (Fig. 6b). It is worth

noting that in this assay the interaction signal strength between the shuttle proteins and PIF7

was lower than that between PIF7 and phyA (Fg. 6b). However, we were able to verify these

interactions through co-immunoprecipitation using *p35S::FHY1/FHL-nLUC* (FHY1/FHL-

nLUC) and *p35S::PIF7-sfGFP* (PIF7-GFP) (Fig. 6c). As our data shows that PIF7 can interact

with phyA, FHY1 and FHL, we postulated that these proteins may form a complex under

inductive WLFR conditions. Supporting this hypothesis, we found that when FHY1-YFP or

FHL-YFP are co-expressed with PIF7-GFP and phyA-RFP in *N. benthamiana* leaves, these

constructs localize to the same nuclear regions under WLFR (Fig. 6d-e).

We also observed increased levels of *ARABIDOPSIS SUMO PROTEASE 1 (ASP1)*, a crucial

factor in SUMOylation and the maintenance of FHY1 levels under FR light conditions[28], in

PIF7ox compared to WT in WL. In WT, *ASP1* levels are increased by WLFR, a response that

is further amplified by *pif7-1* and suppressed by PIF7ox (Fig. 6f). Thus, PIF7 regulation of

ASP1 may play a role in its control of the shuttling mechanism.

To determine if PIF7 primarily acts by sequestering phyA or whether the PIF7 interactions with

FHY1/FHL are also important, we employed *pPHYA::PHYA-sGFP-NLS* (phyA-NLS), where

phyA is constitutively nuclear (Fig. S7)[29]. In line with earlier findings, under our experimental

conditions, phyA-NLS restores the *phyA-201* elongated hypocotyl phenotype to WT length (Fig.

6g). We hypothesized that if PIF7 regulation of phyA was solely due to its sequestration of

phyA, then PIF7ox should diminish the function of phyA-NLS through this mechanism[29].

However, our data show phyA-NLS exhibits strong epistasis over PIF7ox, for hypocotyl length,

and the expression of *CHS* and *IAA29* under WLFR (Fig. 6g, S5). Consequently, it seems that

the mechanism of phyA sequestration alone likely cannot explain the potent effects of PIF7ox

on phyA action. Previous research demonstrated that phyA-NLS completely restores phyA

control of hypocotyl growth in *fhy1-3;fhl-1 and phyA-211;fhy1-3;fhl-1*, indicating that phyA-

NLS can bypass the requirement for nuclear shuttling[30]. Consequently, our results imply that

the interaction between PIF7 and FHY1/FHL may be required to suppress phyA action. Under

WLFR, the transport of phyA into the nucleus is highly dynamic. Therefore, PIF7 may interact

with phyA-FHY1/FHL complexes soon after their nuclear entry and before the complexes

disassemble. This scenario is consistent with our co-localization data (Fig. 6d)[9]. Collectively,

our data indicates that PIF7 forms inactive complexes with phyA and FHY1/FHL, which

simultaneously impedes phyA shuttling and signaling.

DISCUSSION

The operating properties of phyA are distinct from other phytochromes. Under shaded

conditions, phyA requires a cytosolic to nuclear shuttling mechanism to operate effectively, a

process facilitated by the chaperone proteins FHY1 and FHL. In this study, we have shown

PIF7 is a potent regulator of this process, limiting the availability of phyA for signaling. The

impact on phyA is most extreme in the PIF7ox line, where WLFR-induced phyA action is

blocked. Our research illustrates that PIF7 moderates phyA intracellular movement and

signaling through molecular sequestration.

Studies have shown that both PIF7 and phyA accumulate in the nucleus under FR-rich shade

conditions[31,18,29]. Our findings further reveal that PIF7-GFP and phyA-RFP co-localise to

specific sub-nuclear foci, where they physically interact as evidenced by BiFC (Fig. 2a-b).

Additional confirmation of this interaction was provided by split nLUC assays employing PIF7-

lgbit and phyA-smbit, along with co-immunoprecipitation of PIF7ox and phyA.

While all PIFs have an active phytochrome B (APB) domain that enables binding to phyB, only

PIF1 and PIF3 have active phytochrome A (APA) motifs, through which they interact with the

phyA C-terminus[15,32]. This binding is known to facilitate the proteolysis of PIF1 and PIF3, and

to inhibit the activity of PIF3 by isolating it from target promoters[15,32,33]. Our research has

demonstrated that although PIF7 lacks an APA domain, it can still directly interact with phyA.

This binding is facilitated through the PAS domain of phyA and the C-terminal region of PIF7,

which contains a prion-like domain. PIF7 moderates the activity of phyA by inhibiting its ability

to bind to target promoters (Fig. 5b). For the genes *CHS*, *FHY3*, and *BIN2*, which are activated

by phyA in WLFR, and for *FHY1*, *FHL*, *IAA29*, which are repressed by phyA, PIF7 counteracts

the effects of phyA.

PIF7 is particularly effective in FR-rich shade, which triggers its dephosphorylation, increases

its abundance, and enhances its accumulation in nuclear photobodies (Fig. S3b)[20]. Shade also

deactivates phyB, which typically limits the DNA-binding capacity of PIF7[19]. Since continuous

shade also stimulates phyA activity, it creates conducive conditions for the interaction between

PIF7 and phyA.

We have demonstrated that PIF7 and phyA interact within sub-nuclear structures known as

photobodies[5]. These photobodies are essentially molecular condensates that form through

liquid-liquid phase separation (LLPS), accompanied by a nucleation step. Nucleation aids

sequestration, particularly at low concentrations, and targets the formation of photobodies to

specific sub-nuclear sites[6,34]. Across organisms, the spatial organization of biomolecules into

distinct cellular locations is essential for enabling specialized functions[35].

PIF7 is unique amongst PIF proteins as it contains a prion-like domain with a central polyQ

stretch of 17 consecutive glutamine residues. PIF8 is the closest in sequence similarity to PIF7,

and while does possess a short polyQ sequence in its C-terminus, this domain exhibits low

prion-like predictability (Fig. S6). Prion-like domains (PrLD) typically comprise low-

complexity or intrinsically disordered regions that facilitate protein condensation and LLPS.

Indeed, PIF7 is able to form condensates with LLPS characteristics *in vitro*, which is proposed

to facilitate the formation of PIF7-phyB photobodies[19]. Substituting polyQ with polyA

diminishes the capacity of PIF7 to form photobodies, highlighting the importance of polyQ in

this process. Interestingly, phyB has also been shown to possess LLPS characteristics, for its

C-terminus, and disordered N-terminal extension have both been shown to phase-separate in

mammalian cells[36]. We have shown that in WLFR, PIF7 interacts with the phyA, through its

PrLD, which results in the formation of PIF7-phyA-containing photobodies. Within these

biomolecular condensates, phyA-mediated signaling and control of hypocotyl growth are

suppressed. In the PIF7ox line, this inhibition results in excessive elongation of the hypocotyl

under WLFR, since PIF7 action is no longer constrained by phyA.

The movement of phyA to the nucleus in WLFR is strongly contingent on the chaperone

proteins FHY1 and FHL7. Through the use of BiFc, split nanoLUC and co-

immunoprecipitation assays, we demonstrated that PIF7 also interacts with FHY1 and FHL

under WLFR. Fluorescent tagged versions of PIF7, phyA and FHY1/FHL illustrated that these

proteins co-localize within nuclei, suggesting the formation of PIF7-phyA-FHY1/FHL

complexes. Nuclear shuttling of phyA is a highly dynamic process that operates continuously

under WLFR conditions[9,37]. Earlier work indicates that dissociation of the phyA-FHY1/FHL

complex is required to both recycle the shuttle proteins and to facilitate phyA signaling[31,37]. We

therefore propose that PIF7 may serve to trap phyA and FHY1/FHL in inactive complexes,

which simultaneously perturbs the nuclear transport of phyA and disconnects it from

downstream signaling. Consistent with this interpretation, we find *pif7-1* and PIF7ox have

higher proportions of nuclear, and cytosolic phyA, respectively (Fig. 4b-e). Further, our genetic

data shows that phyA-NLS, which is constitutively nuclear, and bypasses the FHY1/FHL

shuttling mechanism, can restore phyA function to PIF7ox in WLFR, in terms of hypocotyl

phenotype and *CHS* and *IAA29* gene expression (Fig. 6g, Fig. S5). We also discovered that PIF7

reduces the expression of *ARABIDOPSIS SUMO PROTEASE 1 (ASP1)*, which aids the

deSUMOylation of FHY1 under FR light, thereby preventing its degradation. *ASP1* suppression

may, therefore, represent an alternative mechanism through which PIF7 modulates phyA

shuttling. Previously, our research demonstrated that under simulated shade, PIF7 interacts with and suppresses the transcriptional activity of ANGUSTIFOLIA 3 (AN3), a promoter of leaf cell proliferation[38]. In this case, PIF7 operates by evicting and substituting for AN3 at target promoters. Thus, in shade conditions, PIF7 assembles different nuclear complexes which execute distinct functions.

In summary, our study outlines the central role PIF7 plays in regulating phyA activity. Its direct binding to phyA, and the FHY1/FHL chaperones, likely represents a rapid, energy-efficient, and highly effective way to stem phyA action. The compartmentalization of phyA and FHY1/FHL into PIF7 photobodies constitutes a dynamic regulatory mechanism that responds to changes in environmental light quality. This subcellular partitioning enables PIF7 to drive hypocotyl elongation while simultaneously reducing the suppressive influence of phyA. Since the modulation of PIF7 by phyB is well-established, our findings position PIF7 at the molecular intersection of phyB and phyA, underscoring its pivotal role in adjusting phyA activity to promote SAS elongation under vegetative shade[19].

METHODS

Plant material and growth conditions

The wild-type of *Arabidopsis thaliana* seedlings are Columbia-0 (Col-0). The *pif7-1*, *pif7-2*[39], *pif8-1*, *pif8-2* mutants, and the construction of *pif8-1;phyA-211*, PIF8ox;*phyA-211* and PIF8ox[15], PIF7ox-FLASH[40], *p35S::GFP-PIF7*[20], *p35S::PIF3-LUC*[41], *p35S::PIF5-HA*, *p35S::PIF4-LUC*[16], *pPHYA::PHYA-sGFP, pPHYA::PHYA-sGFP-NLS*[25], and *pPHYA::PHYA-nanoLUC-3×FLAG-10×His* (phyA-nLUC)[3] have been described previously. The *phyA-211* mutant was obtained from NASC collection. *pPIF7::PIF7-nanoLUC-3×FLAG-10×His* (PIF7-nLUC) (Basta) were transformed into the *pif7-1* background. The *pif7-1*;phyA-GFP-NLS, and PIF7ox;phyA-GFP-NLS, *pif7-1;phyA-211* and PIF7ox;*phyA-211* lines were generated by genetic crossing. Likewise, the *pif7-1;*phyA-nLUC, PIF7ox;phyA-nLUC, PIF7-nLUC; *phyA-211* lines were made through genetic crossing. All the cross lines were verified using PCR or sequencing.

For the hypocotyl growth experiments seeds were first sterilized and then re-suspended in 0.1%
(w/v) agar and stored in darkness at 4 °C for 3 d to stratify. Following this, seeds were
individually sown onto plates containing half-strength Murashige and Skoog basal salts mixture
(Duchefa Biochemie, Haarlem, Netherlands) dissolved in 1% (w/v) agar (1/2 MS media; pH
5.8). The plates were then transferred to Percival I30BLL growth chambers at 22 °C (Plant
Climatics, Wertingen, Germany). To promote germination seeds were exposed to a 4 h period
of white light (WL) (85 ± 5 µmol m^{-2} s^{-1}), and then darkness for 20 h, before transferring to
experimental photoperiods. For all experiments, seedlings were grown under short day (8L:16D)
conditions in either WL or WL with supplementary FR (WLFR) to provide a R:FR ratio of 0.20,
PAR=85 ± 5 µmol m^{-2} s^{-1}.

RNA extraction and qPCR

For RNA extraction seedlings (n = 36 ± 5 seedling numbers), were harvested at Zeitgeber time
7 (ZT), one hour before dusk. Total RNA was extracted using a phenol-based method[38], 300 µL
RNA extraction buffer (PH=8.0) was added to each sample, followed by 300 µL 1:1 acidic
phenol: chloroform. The supernatant was separated from the pellet by centrifuging for 15
minutes at 13,000 rpm under 4 °C. Up to 300 µL of the supernatant was transferred to collection
tubes with 240 µL isopropanol and 30 µL 3 M sodium acetate. Samples were precipitated at -
80 °C for 15 minutes, then centrifuged for 30 minutes at 13,000 rpm. Pellets were washed with
300 µL of 70% ethanol and then suspended in nuclease-free water and stored at -80 °C .
Complementary DNA (cDNA) synthesis was performed using the qScript cDNA SuperMix
(Quanta Biosciences) as described by the manufacturer. A Lightcycler 480 system (Roche) was
employed for qPCR, with 10 µL reactions that incorporated the SYBR Green Master Mix
(Roche). The qPCR primers are shown in **Supplementary Table 2**. Results were analyzed
using Light Cycler 480 (Roche) followed by the Δ ΔCt method.

Hypocotyl measurements and statistical analysis

Images of hypocotyls were taken after 6 days of growth. Hypocotyl length was analyzed using
ImageJ (https://imagej.nih.gov/ij/). Data were analyzed by Excel and GraphPad Prism 10. Data
are presented as mean values ± s.e.m. Two-way ANOVA (α=0.05) was conducted to assess
statistical significance across light conditions for each genotype; asterisks denote significance
relative to control treatments: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), P < 0.0001
(****). Representative data from three independent replicates are shown.

Construct assembly

The full-length CDS of *PHYA* and nanoLUC tag (*nanoLUC-3×FLAG-10×His*; NL3F10H) have been described previously[3]. The CDS of *PIF7* were synthesized by the GeneArt Gene Synthesis service (Invitrogen, MA, USA). The split nanoLUC tags were made by splitting nanoLUC to lgbit (158 aa) and smbit (11 aa). Mobius Assembly[42] was used to insert the constructs into a mUAV plasmid (L0, CamR) with *AraI* restriction site and plasmid assembly was performed and subsequently verified by the Edinburgh Genome Foundry. The full-length CDS of *FHY1* and *FHL* parts were amplified with Q5 High-Fidelity DNA polymerase (New England Biolabs, MA, USA) from the Col-0 genome by gene-specific primers. The *PHYA-C*, *PHYA-C1*, *PHYA-P*, *PHYA-H*, *PIF7-N*, *PIF7-B* and *PIF7-C* part cloned on the CDS region of *PHYA* and *PIF7* (L0 parts). These parts were assembled into a mUAV plasmid and sequenced by Source Bioscience (Cambridge, UK). All L0 parts will assemble in pMAP B L1 (KamR) vector with *BsaI* restriction site. All the constructs for the tobacco (*Nicotiana benthamiana*) assay used a CaMV35S promoter, which is synthesized by the GeneArt Gene Synthesis service (Invitrogen, MA, USA).

To construct *pEXP601-pPIF7::PIF7-nanoLUC-3×FLAG-10×His*, a 2740 bp genomic fragment was generated, encompassing the PIF7 promoter region (1015 bp), 5'UTR (161 bp), and 1564 bp ORF, excluding the stop codon of exon 6 of the AT5G61270.1 gene model. This fragment was PCR-amplified using specific primers with Phusion High-Fidelity DNA Polymerase (NEB, M0530S). The resulting amplicon was subsequently recombined into pDONR221 using BP II clonase (Invitrogen, 12535029), and its sequence verified by Sanger sequencing (Genepool, Edinburgh Genomics, Edinburgh, UK). The genomic regions were then recombined into *pGWB601:nanoLUC-3×FLAG-10×Hiss* using LR Clonase (Invitrogen)[43]. The final construct was transformed into Agrobacterium AGL1. Primers used for construct assembly are listed in **Supplementary Table 1.**

The Tobacco Transient Expression Assay

For this assay, constructs were transferred into the Agrobacterium AGL1 strain and cultured at 28 ºC for two days, before infiltration into 4-week-old *Nicotiana benthamiana* leaves with P19. The transient expression solution contained 10 mM MES buffer (pH 5.6), 10 mM $MgCl_2$ and 200 mM aceto-syringone. Following infiltration plants were grown for 3 days in short day (8L:16D, at 22 ºC) conditions in either WL or WLFR to provide a R:FR ratio of 0.20. Samples were analyzed at ZT7 on day 3.

Confocal Imaging, Colocalization, and BiFC

Images of GFP, YFP and RFP fluorescence were obtained with a Leica confocal micro-scope

(Leica SP8) using 488 nm and 552 nm lasers. For BiFC and co-localization assay all images

were taken at ZT7 on day 3 after WL/WLFR treatments. The co-localization signals were

analyzed by LAS X Office 1.4.7.2982 and plotted by GraphPad Prism 10. *pPHYA::PHYA-*

sGFP(NLS), *pif7-1*;phyA-GFP, PIF7ox;phyA-GFP and *p35S::GFP-PIF7* transgenic lines

images were taken at ZT7 within 30 minutes after 6 days of WL/WLFR treatments. The Nuclear

and cytosolic signals were quantified from cells in the mid-hypocotyl region within a 20 × 1.36

objective lens. Nucleus and cytoplasmic signals were quantified using Image J, after eliminating

background interference. Nuclear/cytoplasmic ratios were calculated and plotted in GraphPad

Prism 10[26]. Imaging experiments were comprised of three biological replicates, with 6 seedlings

per replicate.

Immunoblot Analysis and Co-immunoprecipitation Assays

Each sample comprised 0.5-1 g of Arabidopsis seedling tissue, harvested on day 6 at ZT7.

Samples were added to lysis buffer (125 mM Tris-HCl (pH 7.5); 0.25mM EDTA (pH 7.5), 150

mM NaCl, 5% glycerol, 0.1% Triton X-100, 0.5% NP-40), which contained 1× Protease

inhibitor (Sigma-Aldrich, P9599) and 1× HALT-Phosphatase inhibitor (Thermo Scientific).

Samples were centrifuged at 12,000 rpm for 10 mins at 4 ºC and then transferred to lysate mix

with 1× SDS loading buffer (containing 0.1 M DTT) at a 1:1 volume ratio. Samples were

incubated at 80 ºC for 10 mins, followed by a centrifugation step, before western blotting.

For Co-immunoprecipitation, after lysis an aliquot of the total protein was saved as an input

sample. Then, other total protein extracts were incubated with Protein A/G beads (MCE

Magnetic Agarose) for 1 h at 4 °C to remove adventitious binding proteins. Then anti-

FLAG/His beads (Anti-FLAG M2 agarose gel Sigma-Aldrich, Cat.No. A2220; Anti-His

Magnetic Beads, Cat. 209, MCE) were added into treated protein extracts and kept overnight

at 4 ºC. The antibody/protein complex was washed with wash buffer (125 mM Tris-HCl [pH

7.5]; 0.25mM EDTA [pH 7.5], 150 mM NaCl, 5% glycerol, 0.1 % Triton X-100) 5 times. Then,

50 µL of 1× SDS loading buffer were added to each sample and incubated at 80 ºC for 8 mins.

To detect native phyA protein, an anti-Phytochrome A antibody (PHYTOAB, PHY1907) was

used at 1:1,000. To detect the PIF7ox-FLASH and *p35S::FHY1/FHL-nanoLUC × 3FLAG×*

10His an anti-FLAG M2 antibody (Sigma-Aldrich, F1804) was used at 1:2,000. To detect the

p35S::PIF7-sfGFP an anti-GFP anti-body (Roche, ab290) was used at 1:3,000. Anti-RPN6

antibody (Agrisera, AS121858) was used at 1:3,000 as internal control. Secondary antibodies

(1:3,000, anti-mouse IgG (cell signaling, 7076s); 1:4,000 Goat anti-rabbit-IgG (Abcam,

AB6013) were used for the ECL Prime Western Blotting System. The signals were detected by

Azure 300 Imager with ECL prime western blotting detection reagents (Cytiva, RPN2232).

NanoLUC and split nanoLUC assays

NanoLUC bioluminescence assays were conducted as previously described[3]. Single Arabidopsis seedlings were grown in individual wells, within a 96-well plate, each containing 200 µl 1/2 MS media. Plates were sealed with cling film and placed under 8L:16D (WL) for 6 days. Before imaging, 50 µl of 1:50 water dilution of Nano-Glo buffer (Promega, Nano-Glo®Luciferase Assay) was added to the seedlings in each well. Plates were sealed and moved to experimental light conditions. Bioluminescence was recorded every hour (for 1.5 s) for several days using the TriStar2 LB942 multimode microplate reader (Berthold Technologies, Germany). In **Fig. 1g**, the data were analyzed by averaging individual seedling measurements and then calculating the fold change relative to the ZT0 data for each biological replicate. Subsequently, the results were plotted using GraphPad Prism 10. A total of 9 biological replicates were conducted, with each replicate comprising at least 6 seedlings. The data was analyzed by BioDare2 software[44], applying amp&baseline detrending and normalization steps, before plotting in GraphPad Prism 10 (**Fig. S2**)

The split nanoLUC assay utilized the *N. benthamiana* transient expression system described above. Following WL/WLFR treatments for 3 days, 1cm x 1cm leaf samples (*n* =12) were transferred to 96 wells plates containing 100 µl diluted (1:50) Nano-Glo for imaging with a plate reader. The data analyzed by Excel and GraphPad Prism 10. Experiments repeated at least 3 times.

Chromatin Immunoprecipitation (ChIP)-qPCR

We employed ChIP-qPCR methods described previously[45]. For each sample 0.5-1 g of tissue was extracted from 6 day -old seedlings at ZT7. Cross-linking was induced with 1% formaldehyde under vacuum infiltration for 15 mins (up to -70 kPa), then 2.5 ml of 2 M glycine was added to quench crosslinking. DNA was fragmented into an average size of ~150 bp in Diagenode BioRuptor Plus, (30 cycles of high power 30s ON / 30s OFF). Anti- PHYA antibody (PHYTOAB, PHY1907,1:300) incubated with protein A/G beads (MCE Magnetic Agarose) in 4 ºC overnight. And the following day, samples were incubated with protein A/G bead/PHYA antibody complex for 6 hours at 4 ºC. DNA was extracted using phenol/chloroform (1:1, vol/vol). Protein enrichment at DNA regions was quantified using qPCR with the primers detailed in **Supplementary Table 2.** Enrichment was determined by %Input method (%Input=2 ^ [CtInput-CtChIP] × IDF × 100%).

Nuclear-Cytoplasmic Fractionation

Nuclear fractionation was performed essentially as previously described with the following

modifications[46]. Briefly, 0.5-1 g of Arabidopsis seedlings ground into fine powder in liquid nitrogen and then homogenized in double the volume of pre-cooled (4 °C) lysis buffer (20 mM Tris-HCl, pH 7.4, 25% Glycerol, 20 mM KCl, 2 mM EDTA, 2.5 mM $MgCl_2$, 250 mM Sucrose and 5 mM DTT, add 1× protease inhibitor cocktail (Roche) and 1× HALT-Phosphatase inhibitor added in fresh). The sample was filtered twice through two layers of Mira cloth (Merck Millipore). The flow-through was spun 10 min at 2000 g, 4 °C. Supernatant, consisting of the cytoplasmic fraction, was centrifuged 20 min at 13,000 g, 4 °C and collected. The pellet, containing the nuclear fraction, was washed six times with 2.5 mL of nuclear resuspension buffer NRBT1 (20 mM Tris-HCl, pH 7.4, 25% Glycerol, 2.5 mM $MgCl_2$, 0.2% Triton X-100) and centrifuged 3 min at 1,500 g, 4 °C. The pellet was resuspended in 300 µL of pre-cooled NRBT2 (250 mM Sucrose, 10 mM Tris-HCl, pH 7.5, 10 mM $MgCl_2$, 1% Triton X-100, 0.035% β-mercaptoethanol, and 1× protease inhibitor cocktail added in fresh), and carefully overlaid on top of 300 µL NRBT3(1.7 M Sucrose, 10 mM Tris-HCl, pH 7.5, 2 mM $MgCl_2$, 0.15% Triton X-100, 0.035% β-mercaptoethanol, and 1× protease inhibitor cocktail added in fresh). These were centrifuged at 16,000 g for 55 min at 4 °C, and the final nuclear pellet was resuspended in 2× SDS loading buffer. As quality controls for the fractionation, RbCL (Agrisera, AS03037, 1:3,000) protein was used as the cytoplasmic control, and histone H3 (Abcam, ab18521, 1:3,000) was used as the nuclear control in immunoblot assays.

ACKNOWLEDGEMENTS

This research was supported by University of Edinburgh Darwin Trust Scholarship awarded to MK. Z and Leverhulme Trust grant RPG-2024-082 awarded to K.J.H. We thank Dr. Rist Van de Weyer who generated PIF7ox;*phyA-211* and first observed the PIF7ox hypocotyl phenotype in WLFR. We thank Prof. Lin Li, Prof. Dr. Andreas Hiltbrunner and Dr. Gabriela Toledo-Ortiz who kindly shared *p35S::GFP-PIF7*, *p35S::PHYA-RFP* and *pPHYA::PHYA-sGFP*(NLS) seeds and thank Prof. Giltsu Choi for sharing *pif8-1;phyA-211*, *PIF8ox;phyA-211* and PIF8ox lines.

AUTHOR CONTRIBUTIONS

MK.Z and K.J.H. planned and designed the research. MK.Z conducted the experiments, which were analyzed by K.J.H. and MK.Z. A.R. generated the *pPIF7::PIF7-nanoLUC* line. The manuscript was prepared by MK.Z and K.J.H, subsequent editing performed by K.J.H and A.R.

REFERENCES

1   Liu, Y., Jafari, F. & Wang, H. Y. Integration of light and hormone signaling pathways in the regulation of plant shade avoidance syndrome. *Abiotech* **2**, 131-145 (2021). https://doi.org/10.1007/s42994-021-00038-1.

2   Franklin, K. A. Shade avoidance. *New Phytol* **179**, 930-944 (2008).
https://doi.org:10.1111/j.1469-8137.2008.02507.x.

3   Butlin, P., Valdivia-Cabrera, M., Grima, R. & Halliday, K. J. Arabidopsis thaliana
Phytochrome A Sensory Properties in Canopy Shade. *bioRxiv* (2025).
https://doi.org:10.1101/2025.05.14.654025.

4   Mathews, S. & Sharrock, R. A. Phytochrome gene diversity. *Plant Cell Environ* **20**, 666-
671 (1997). https://doi.org:10.1046/j.1365-3040.1997.d01-117.x.

5   Cheng, M. C., Kathare, P. K., Paik, I. & Huq, E. Phytochrome Signaling Networks. *Annu*
Rev Plant Biol **72**, 217-244 (2021). https://doi.org:10.1146/annurev-arplant-080620-
024221.

6   Du, J., Kim, K. & Chen, M. Distinguishing individual photobodies using Oligopaints
reveals thermo-sensitive and -insensitive phytochrome B condensation at distinct
subnuclear locations. *Nature Communications* **15** (2024). https://doi.org:10.1038/s41467-
024-47789-1.

7   Casal, J. J., Luccioni, L. G., Oliverio, K. A. & Boccalandro, H. E. Light, phytochrome
signalling and photomorphogenesis in. *Photoch Photobio Sci* **2**, 625-636
(2003).https://doi.org:10.1039/b300094j.

8   Rausenberger, J. *et al.* Photoconversion and Nuclear Trafficking Cycles Determine
Phytochrome A's Response Profile to Far-Red Light. *Cell* **146**, 813-825 (2011).
https://doi.org:10.1016/j.cell.2011.07.023.

9   Hiltbrunner, A. *et al.* Nuclear accumulation of the phytochrome A photoreceptor requires
FHY1. *Curr Biol* **15**, 2125-2130 (2005). https://doi.org:10.1016/j.cub.2005.10.042.

10   Genoud, T. *et al.* FHY1 Mediates Nuclear Import of the Light-Activated Phytochrome A
Photoreceptor. *Plos Genet* **4** (2008). https://doi.org:10.1371/journal.pgen.1000143.

11   Li, J. G. *et al.* Transcription Factor ELONGATED HYPOCOTYL5 Plays a Role in the
Feedback Regulation of Phytochrome A Signaling. *Plant Cell* **22**, 3634-3649 (2010).
https://doi.org:10.1105/tpc.110.075788.

12   Pham, V. N., Kathare, P. K. & Huq, E. Phytochromes and Phytochrome Interacting Factors.
Plant Physiol **176**, 1025-1038 (2018). https://doi.org:10.1104/pp.17.01384.

13   Jeong, J., Lee, Y. & Choi, G. Both phytochrome A and phyB interact with
PHYTOCHROME-INTERACTING FACTORs through an evolutionary conserved
phy(OPM)-APA interaction. *Nat Commun* **16**, 3946 (2025).
https://doi.org:10.1038/s41467-025-59327-8.

14   Al-Sady, B., Ni, W., Kircher, S., Schafer, E. & Quail, P. H. Photoactivated phytochrome
induces rapid PIF3 phosphorylation prior to proteasome-mediated degradation. *Mol Cell*
23, 439-446 (2006). https://doi.org:10.1016/j.molcel.2006.06.011.

15   Oh, J., Park, E., Song, K., Bae, G. & Choi, G. PHYTOCHROME INTERACTING
FACTOR8 Inhibits Phytochrome A-Mediated Far-Red Light Responses in Arabidopsis.
Plant Cell **32**, 186-205 (2020). https://doi.org:10.1105/tpc.19.00515.

16   Seaton, D. D. *et al.* Dawn and photoperiod sensing by phytochrome A. *P Natl Acad Sci*
USA **115**, 10523-10528 (2018). https://doi.org:10.1073/pnas.1803398115.

17   Zhu, T., Yang, C., Xie, Y., Huang, S. & Li, L. Shade-induced lncRNA PUAR promotes
shade response by repressing PHYA expression. *EMBO Rep* **24**, e56105 (2023).
https://doi.org:10.15252/embr.202256105.

18 Leivar, P. *et al.* The Arabidopsis phytochrome-interacting factor PIF7, together with PIF3
and PIF4, regulates responses to prolonged red light by modulating phyB levels. *Plant*
Cell **20**, 337-352 (2008). https://doi.org:10.1105/tpc.107.052142.

19 Xie, Y. *et al.* Phytochrome B inhibits the activity of phytochrome-interacting factor 7
involving   phase   separation.   *Cell   Rep*   **42**   (2023).
https://doi.org:10.1016/j.celrep.2023.113562.

20 Huang, X. *et al.* Shade-induced nuclear localization of PIF7 is regulated by
phosphorylation   and   14-3-3   proteins   in   Arabidopsis.   *Elife*   **7**   (2018).
https://doi.org:10.7554/eLife.31636.

21 Park, E. *et al.* Phytochrome B inhibits binding of phytochrome-interacting factors to their
target promoters. *Plant Journal* **72**, 537-546 (2012). https://doi.org:10.1111/j.1365-
313X.2012.05114.x.

22 Ni, M., Tepperman, J. M. & Quail, P. H. PIF3, a phytochrome-interacting factor necessary
for normal photoinduced signal transduction, is a novel basic helix-loop-helix protein. Cell
95, 657-667 (1998). https://doi.org:10.1016/S0092-8674(00)81636-0.

23 Lorrain, S., Trevisan, M., Pradervand, S. & Fankhauser, C. Phytochrome interacting
factors 4 and 5 redundantly limit seedling de-etiolation in continuous far-red light. *Plant*
Journal **60**, 449-461 (2009). https://doi.org:10.1111/j.1365-313X.2009.03971.x.

24 Kevei, E., Schafer, E. & Nagy, F. Light-regulated nucleo-cytoplasmic partitioning of
phytochromes. *J Exp Bot* **58**, 3113-3124 (2007). https://doi.org:10.1093/jxb/erm145.

25 Toledo-Ortiz, G. *et al.* Subcellular sites of the signal transduction and degradation of
phytochrome   A.   *Plant   Cell   Physiol*   **51**,   1648-1660   (2010).
https://doi.org:10.1093/pcp/pcq121.

26 Chen, F. *et al.* Phosphorylation of FAR-RED ELONGATED HYPOCOTYL1 is a key
mechanism defining signaling dynamics of phytochrome A under red and far-red light in
Arabidopsis. *Plant Cell* **24**, 1907-1920 (2012). https://doi.org:10.1105/tpc.112.097733.

27 Wang, H. & Deng, X. W. Arabidopsis FHY3 defines a key phytochrome A signaling
component directly interacting with its homologous partner FAR1. *EMBO J* **21**, 1339-
1349 (2002). https://doi.org:10.1093/emboj/21.6.1339.

28 Qu, G. P. *et al.* Reversible SUMOylation of FHY1 Regulates Phytochrome A Signaling in
Arabidopsis. *Mol Plant* **13**, 879-893 (2020). https://doi.org:10.1016/j.molp.2020.04.002.

29 Toledo-Ortiz, G. *et al.* Subcellular sites of the signal transduction and degradation of
phytochrome   A.   *Plant   Cell   Physiol*   **51**,   1648-1660   (2010).
https://doi.org:10.1093/pcp/pcq121.

30 Menon, C., Klose, C. & Hiltbrunner, A. Arabidopsis FHY1 and FHY1-LIKE Are Not
Required for Phytochrome A Signal Transduction in the Nucleus. *Plant Commun* **1**,
100007 (2020). https://doi.org:10.1016/j.xplc.2019.100007.

31 Rausenberger, J. *et al.* Photoconversion and Nuclear Trafficking Cycles Determine
Phytochrome A's Response Profile to Far-Red Light. *Cell* **146**, 813-825 (2011).
https://doi.org:10.1016/j.cell.2011.07.023.

32 Al-Sady, B., Ni, W., Kircher, S., Schafer, E. & Quail, P. H. Photoactivated phytochrome
induces rapid PIF3 phosphorylation prior to proteasome-mediated degradation. *Mol Cell*
23, 439-446 (2006). https://doi.org:10.1016/j.molcel.2006.06.011.

33 Shen, H. *et al.* Light-induced phosphorylation and degradation of the negative regulator

PHYTOCHROME-INTERACTING FACTOR1 from depend upon its direct physical
interactions with photoactivated phytochromes. *Plant Cell* **20**, 1586-1602 (2008).
https://doi.org:10.1105/tpc.108.060020.

34   Kim, C. *et al.* Phytochrome B photobodies are comprised of phytochrome B and its
primary and secondary interacting proteins. *Nature Communications* **14** (2023).
https://doi.org:10.1038/s41467-023-37421-z.

35   Zhang, X. Y. *et al.* Liquid-liquid phase separation in diseases. *Medcomm* **5** (2024).
https://doi.org:e64010.1002/mco2.640.

36   Chen, D. *et al.* Integration of light and temperature sensing by liquid-liquid phase
separation of phytochrome B. *Mol Cell* **82**, 3015-+ (2022).
https://doi.org:10.1016/j.molcel.2022.05.026.

37   Genoud, T. et al. FHY1 Mediates Nuclear Import of the Light-Activated Phytochrome A
Photoreceptor. Plos Genet 4 (2008).
https://doi.org:100014310.1371/journal.pgen.1000143.

38   Hussain, E., Romanowski, A. & Halliday, K. J. PIF7 controls leaf cell proliferation through
an AN3 substitution repression mechanism. *P Natl Acad Sci USA* **119** (2022).
https://doi.org:10.1073/pnas.2115682119.

39   Leivar, P. *et al.* Multiple Phytochrome-Interacting bHLH Transcription Factors Repress
Premature Seedling Photomorphogenesis in Darkness. *Curr Biol* **18**, 1815-1823 (2008).
https://doi.org:10.1016/j.cub.2008.10.058.

40   Li, L. *et al.* Linking photoreceptor excitation to changes in plant architecture. *Genes Dev*
26, 785-790 (2012). https://doi.org:10.1101/gad.187849.112.

41   Viczián, A. *et al.* Functional characterization of phytochrome interacting factor 3 for the
circadian clockwork. *Plant and Cell Physiology* **46**, 1591-1602 (2005).
https://doi.org:10.1093/pcp/pci175.

42   Andreou, A. I. & Nakayama, N. Mobius Assembly: A versatile Golden-Gate framework
towards universal DNA assembly. Plos One 13 (2018). https://doi.org:
10.1371/journal.pone.0189892.

43   Urquiza-García, U. & Millar, A. J. Expanding the bioluminescent reporter toolkit for plant
science with NanoLUC. Plant Methods 15 (2019). https://doi.org:10.1186/s13007-019-
0454-4.

44   Zielinski, T., Moore, A. M., Troup, E., Halliday, K. J. & Millar, A. J. Strengths and
Limitations of Period Estimation Methods for Circadian Data. *Plos One* **9** (2014).
https://doi.org:10.1371/journal.pone.0096462

45   Wang, Z. S. *et al.* Proteasome-associated ubiquitin ligase relays target plant hormone-
specific transcriptional activators. *Sci Adv* **8** (2022).
https://doi.org:10.1126/sciadv.abn4466.

46   Zhu, D. M. *et al.* Biochemical Characterization of Complexes Containing
CONSTITUTIVELY PHOTOMORPHOGENIC1 and SUPPRESSOR OF PHYA Proteins
in Light Control of Plant Development. *Plant Cell* **20**, 2307-2323 (2008).
https://doi.org:10.1105/tpc.107.056580.

47   Rosler, J., Klein, I. & Zeidler, M. double mutant reveals a distinct cytoplasmic action of
phytochrome A. *P Natl Acad Sci USA* **104**, 10737-10742 (2007).
https://doi.org:10.1073/pnas.0703855104.

Fig. 1 PIF7ox influences phyA Function in WLFR

a, Schematic of experimental growth conditions. Seeds were sown on 1/2 MS plates and cultivated for one day under white light, followed by 4 hours (h) of light and 20 h of dark. Seedlings were subsequently grown for 6 days in white light (WL) photoperiods with or without supplementary far-red (WLFR), lowering the R:FR to 0.20 (red arrow). For all experiments, light regimes were 8 h light at PAR = 85 ± 5 μmol m^{-2} s^{-1}, 16 h dark, at 22°C.

b, Representative images of *Arabidopsis thaliana* WT (*Col-0*), *pif7-1*, *pif7-2*, and PIF7ox seedlings grown in WL or WLFR for 6 days. Scale bar = 1 cm.

c, PIF7ox blocks phyA action in WLFR. Hypocotyl length (mm) of WT (*Col-0*), *pif7-1*, *pif7-2*, PIF7ox, *phyA-211*, *pif7-1;phyA-211*, *PIF7ox;phyA-211* and phyAox under WL and WLFR. Representative images from three independent replicates are shown.

d, In WLFR, PIF7 enhances *PHYA* expression at Zeitgeber time 7 (ZT). Real-time qPCR was used to quantify *PHYA* mRNA levels, with *PP2A* as internal control. Samples were collected at ZT7 (1 hour before dusk) from 6d-old seedlings grown in WL or WLFR.

e,f, In WLFR, PIF7 increases PHYA protein accumulation. PHYA protein levels were determined by immunoblotting. Anti-PHYA was used to detect native phyA, while anti-RPN6 served as an internal control. The bar chart (**f**) displays the relative levels of phyA, normalised to RPN6, based on data from four biological replicates collected at ZT7.

g, PIF7ox boosts PHYA protein levels in diel conditions. The phyA-nLUC line was generated by expressing (p*PHYA::PHYA-nLUC*) in *phyA-211*, while the PIF7ox;phyA-nLUC was generated by introgressing p*PHYA::PHYA-nLUC* into the PIF7ox line. Bioluminescence signals were recorded from day 6.

For **c,d,f,g**, data are presented as mean values ± s.e.m., *n*=25 (seedling numbers) in **c**; with *n*=3,4,9 (biological repeats) in **d,f,g**, respectively. For **c,d,f**, a Two-way ANOVA was conducted to assess statistical significance across light conditions for each genotype (α=0.05); asterisks denote significance relative to WL treatments: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), $P < 0.0001$ (****).

Fig. 2 PIF7 and phyA interact in sub-nuclear photobodies under WLFR.

a, *p35S::PIF7-sfGFP* and *p35S::PHYA-RFP* co-localise in nuclei and photobodies in WLFR. Co-localisation is evident from the chart (right) which traces the GFP and RFP channel signals along the white bar shown in merged nucleus image (left).

b, Bimolecular Fluorescence Complementation (BiFC) assay confirms PIF7 and phyA interact in nuclear photobodies. BiFC was detected between *p35S::PIF7-nYFP* and *p35S::PHYA*-cYFP (PIF7-nYFP + phyA-cYFP) in WLFR, but not with the *p35S::EV-cYFP* (EV-cYFP, empty vector) control.

c, Split NanoLUC (nLUC) assay demonstrates PIF7 and phyA physically interact. *p35S::PIF7-lgbitnLUC* (PIF7-lgbit), *p35S::PHYA-smbitnLUC* (phyA-smbit), *p35S::PHY*B-smbitnLUC (phyB-smbit) and *p35S::PHY*A-nLUC (phyA-nLUC) reporter constructs were expressed in *Nicotiana benthamiana*. A WLFR-dependent interaction was demonstrated for PIF7-lgbit+phyA-smbit. PhyA-nLUC was used as light conditional controls, and phyB-smbit+phyA-smbit as the negative control.

d, Co-immunoprecipitation assay (Co-IP) shows that phyA interacts with PIF7 in WLFR *in vivo*. Anti-His beads were used to pull down PIF7 from PIF7ox (*p35S::PIF7-FLASH*; 9×Myc-6×His-3×FLAG) or the WT(*Col-0*) negative control. Anti-PHYA and anti-FLAG used for detecting phyA and PIF7 respectively, in IP/Input. Samples were collected at ZT7 from 6-d old seedlings exposed to WL or WLFR.

For **a,b**, 488nm and 552nm laser detected the GFP/YFP and RFP signals. Scale Bar=15μm. For **c**, data are presented as mean values ± s.e.m., *n*=12 (leaf numbers). A Two-way ANOVA test was used to obtain statistical significance (α=0.05); asterisks denote statistical significance comparing negative to positive test or tested interaction treatments: 0.1234 (ns), 0.0332(*), 0.0021 (**), 0.0002(***), P<0.0001 (****). Representative images from three independent replicates are shown in all figs.

Fig. 3 The PIF7-phyA Interaction Depends on the PIF7 Prion like domain

a-b, BIFC assays using truncated constructs identify the PHYA PAS domain and the PIF7 prion-like domain, as important for the phyA-PIF7 interaction. The constructs image depicted the complete PHYA/PIF7 protein structures. Full length and truncated contructs were fused to C-terminal nYFP (PIF7) or cYFP (PHYA) BIFC constructs. PIF7-nYFP+phyA-cYFP was used as positive control and *p35S::EV-cYFP/p35S::EV-nYFP* (EV-cYFP/EV-nYFP) as used as negative controls. 488nm lasers were used to detect the YFP signals. Scale Bar=15μm. Representative images from three independent replicates are shown.

Fig. 4 PIF7 Influences the Sub-cellular Location of phyA

a-c, In WLFR, PIF7ox promotes accumulation of phyA-GFP in the cytosol. Lines of *pif7-1*;phyA-GFP and PIF7ox;phyA-GFP were generated by introgressing *pPHYA::PHYA-sGFP* into *pif7-1* and PIF7ox. Close-up images under WLFR are shown in **b**, white arrows and N label indicate nuclear, while red arrows and C label indicate cytosolic signals. In **c**, nuclear/cytosolic (N/C) ratios (relative to WT (*Col-0*), WL control) were calculated following quantification using ImageJ.

d-e, PIF7ox enhances PHYA protein accumulation in the cytosol under WLFR conditions. Immunoblot assays were conducted using an anti-PHYA antibody, using anti-RbCL as the cytosolic control, and anti-Histone 3 (H3) as the nuclear control. In **e**, cytoplasmic and nuclear phyA was normalised to RbCL and H3, respectively. The N/C ratio for each genotype is shown relative to WT (*Col-0*), WL control. Six-day-old seedlings were sampled as ZT7.

For **a-b** a wavelength of 488nm was used to detect GFP signals. Scale bar=25μm. Representative images from three independent replicates are shown. For **c,e**, data are presented as mean values ± s.e.m., *n*=6 (seedling numbers) in **c**; *n*=4 (biological repeats) in **e**. A two-way ANOVA test assessed statistical significance between genotypes under WL/WLFR condition (α=0.05); asterisks denote significance compared to the phyA-GFP (phyA N/C) values for WL/WLFR condition(**c**) or phyA protein (phyA N/C) values in WT with WL/WLFR condition(**e**): 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), $P < 0.0001$ (****).

Fig. 5. PIF7 Inhibits phyA-Mediated Regulation of Target Genes

a, qPCR analysis of *CHS, FHY1, FHL, FHY3, BIN2,* and *IAA29* expression in WT (*Col-0*), *pif7-1*, PIF7ox and *phyA-211*. Transcript levels were normalised to the *PP2A* internal control.

b, PIF7ox reduces phyA enrichment at *CHS, FHY1,* and *FHL* promoters under WLFR. Chromatin immunoprecipitation (ChIP)-qPCR was conducted with 6-day-old WT (*Col-0*) or PIF7ox seedlings sampled at ZT7. Native phyA was immobilised on protein A/G beads using an anti-phyA antibody, and promoter enrichment quantified by qPCR at Gbox promoter regions and coding regions, which served as negative controls. Data shown is normalised to input samples.

For **a-b**, data are presented as mean values ± s.e.m., *n*=3 (biological repeats). A two-way ANOVA test was used to assess statistical significance between the WL/WLFR for each genotype (**a**) or different genotypes under the same conditions (**b**) (α=0.05); asterisks indicate statistical significance compared to control: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), $P < 0.0001$ (****).

Fig. 6. PIF7 interacts with FHY1/FHL in WLFR

a, The BiFC assay demonstrates PIF7 interaction with FHY1 and FHL. Interaction fluorescence was detected when *p35S::FHY1-cYFP* (FHY1-cYFP) or *p35S::FHL-cYFP* (FHL-cYFP) were with expressed *p35S::PIF7-nYFP* (PIF7-nYFP) in *N. benthamiana* under WLFR.

b, Interactions between PIF7 and FHY1/FHL are detected using a split NanoLUC assay. Strong fluorescence signals were detected between *p35S::FHY1-smbitnLUC* (FHY1-smbit), *p35S::FHL-lgbitnLUC* (FHL-lgbit) and *p35S::PIF7-lgbit(smbit)nLUC* (PIF7-lgbit/smbit) in the transient *N. benthamiana* expression system. The positive controls were phyA-nLUC and PIF7-lgbit+phyA-smbit, while PIF7-smbit+phy-Asmbit was used as a negative control.

c, An immunoprecipitation assay demonstrates PIF7 interacts with FHY1/FHL *in vivo*. *p35S::FHY1/FHL-nanoLUC × 3FLAG × 10His* and *p35S::PIF7-sfGFP* constructs were transiently expressed in *N. benthamiana* leaves. Anti FLAG beads were used to pull down FHY1/FHL and its target proteins by Co-IP. Anti GFP antibodies revealed GFP-tagged PIF7 co-IP with FHY1/FHL in WLFR.

d-e, Fluorescence assays illustrate that PIF7, phyA and FHY1/FHL co-localise in nuclei and photobodies. *p35S::PIF7-sfGFP*, *p35S::FHY1/FHL-EYFP*, and *p35S::PHYA-RFP* constructs were transiently expressed in *N. benthamiana* leaves. The merged image (**d**) signifies co-localisation, and **e**, displays GFP, YFP and RFP channel signals relative to white bar shown in the merged nuclear image.

f, PIF7 reduces *ASP1* mRNA levels under WLFR. qPCR analysis of *ASP1* expression in WT, *pif7-1*, PIF7ox, *PP2A* was used as internal control.

g, phyA-NLS is epistatic to PIF7ox. Hypocotyl length (mm) is shown for WT (Col-0), *phyA-211, phyAox;phyA-211,* WT (Ler), *phyA-201, phyA-NLS;phyA-201*, *PIF7ox;phyA-NLS;phyA-201.* Double and triple mutant lines were generated through genetic.

h, Schematic depicting the cellular model for PIF7 control of phyA action in canopy shade. In WLFR PIF7 accumulates in the nucleus, where it interacts with phyA and FHL/FHY1 in photobodies. This sequestration not only blocks phyA from regulating downstream genes but also inhibits its transport into the nucleus. Consequently, PIF7 is able to amplify SAS-mediated hypocotyl elongation by blocking the antagonistic action of phyA. The image created in BioRender (https://BioRender.com).

For **a,d**, 488nm and 552nm lasers were used for detecting GFP/YFP and RFP signals. Scale bar=15μm. For **b,f,g**, data are presented as mean values ± s.e.m., *n*=12 (leaf numbers) in **b**; *n*=3 (biological repeats) in **f**; *n*=24 (seedling numbers) in **g**. A two-way ANOVA test was conducted to determine statistical significance (α=0.05), asterisks denote significance when comparing negative to positive controls and tested interactions (**b**) or compared to WL treatments in each genotype (**f-g**): 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), $P < 0.0001$ (****).