PHYTOCHROME INTERACTING FACTOR 7 moderates the activity and the cytosolic-

nuclear shuttling of PHYTOCHROME A

Mengke Zhou[1*], Andrés Romanowski[2], Karen J. Halliday[1*]

1. Institute for Molecular Plant Sciences, School of Biological Sciences, University of

Edinburgh, Edinburgh EH9 3BF, United Kingdom.

2. Timing of Environmental Signalling lab, Molecular Biology Group, Plant Sciences,

Wageningen University & Research, Wageningen, The Netherlands.

ORCID IDs: 0000-0003-1043-2766 (MK. Z); 0000-0003-0737-2408(A.R.); 0000-0003-0467-

104X (K.J.H.).

*Corresponding authors:

Mengke Zhou and Karen J. Halliday

Institute for Molecular Plant Sciences, School of Biological Sciences, University of Edinburgh,

Edinburgh EH9 3BF, United Kingdom

+44-0131-651-9083

mengke.zhou@ed.ac.uk

karen.halliday@ed.ac.uk

ABSTRACT

In nature, plants sense and navigate complex vegetative environments using the phytochrome (phy) photoreceptor-signalling system. The canopy adaptation strategy, elicited by phyA, and the shade avoidance syndrome, regulated by phyB, allow plants to adapt to diverse forms of vegetation shade. The phyA High Irradiance Response (HIR) mode, involving photocycle-coupled nuclear shuttling, is essential for detecting persistent far-red rich conditions typical of canopy or occlusive shade. Our research identifies PHYTOCHROME INTERACTION FACTOR 7 (PIF7) as a critical regulator of phyA function. We demonstrate PIF7ox can effectively block phyA action. In persistent shade, PIF7 and phyA both accumulate and interact within nuclear bodies, with binding facilitated through the prion-like domain of PIF7 and the C-terminal PAS-PAS domain of phyA. PIF7 also interacts with FHY1 and FHL chaperones, interfering with phyA cytosolic-nuclear shuttling. This strategic molecular targeting by PIF7 provides an effective strategy for modulating the phyA HIR in response to shade signals.

INTRODUCTION

Light sensing is fundamental for plant adaptation and survival in vegetative shade. This is particularly important during seedling establishment, a stage when plants are most vulnerable. Plants have evolved distinct developmental strategies to cope with different types of shade that are prevalent in nature. The light receptor phytochrome B (phyB), is able to detect FR-rich light from neighbouring plants and elicit the Shade Avoidance Syndrome (SAS), which induces elongation of seedling hypocotyls, repositions leaves to improve light capture, and triggers early flowering [1,2]. Under persistent canopy shade, plants adopt alternative strategies, such as the phyA-mediated Canopy Adaptation Strategy (CAS), which suppresses hypocotyl elongation and conserves resources, optimising growth when shade is unyielding[3]. As vegetative habitats can change considerably through a season, the ability to adjust photoreceptor activity offers adaptive flexibility, ensuring successful seedling establishment and progression through the life cycle.

In Arabidopsis, phytochromes A-E, form a family of photoreceptors that principally absorb light in the red (R) and far-red (FR) regions of the spectrum[4]. Absorption of R, photoconverts phytochrome into its active Pfr form, while FR photoconverts Pfr to the inactive Pr conformer.

When activated phytochromes move from the cytosol to the nucleus where they aggregate in sub-nuclear photobodies to perform a range of functions[5,6]. While all phytochromes possess similar photosensory properties, they fall into two classes based on their distinct modes of action. Phytochromes B-E, which operate in the low fluence response (LFR) mode, are activated by R and deactivated by FR light. Consequently, in FR-rich vegetation shade environments, these phytochromes are less active[4,7]. A key distinction of phyA is that it is activated by FR-rich light through its high irradiance response (HIR) mode of action[7]. This property arises partly from its photobiology and partly from the shuttling mechanism that concentrates phyA in the nucleus[8]. PhyA lacks a nuclear localisation signal and primarily relies on the chaperone proteins FHY1/FHL to facilitate its transport into the nucleus. These proteins specifically recognise phyA-Pfr and transport it to the nucleus, where it is converted into phyA-Pr, allowing FHY1/FHL to dissociate and be recycled to the cytoplasm for further shuttling[8,9,10]. Nuclear located phyA regulates a number of processes, including expression of target genes, through association with transcription factors such as PHYTOCHROME INTERACTING FACTORS (PIFS) and ELONGATED HYPOCOTYL 5 (HY5)[11]. The central role of FHY1/FHL shuttling in the phyA HIR, identifies this mechanism as a potential target for modifying phyA action and the adaptive response[8]. Indeed, HY5 has been reported to be involved in negative-feedback regulation of phyA signalling through repressing FHY1/FHL expression[11]. Here HY5 directly interacts with the DNA-binding regions of FAR-RED ELONGATED HYPOCOTYLS 3 (FHY3) and FAR-RED IMPAIRED RESPONSE 1 (FAR1) to prevent their transcriptional activation of FHY1/FHL.

PIF transcription factors are well-studied negative regulators of phytochrome signalling that. They function by directly binding to G/PBE-boxes in the promoters of their target genes[12]. In Arabidopsis, there are eight PIFs (PIF1, PIF2/PIL1-PIF8), each with an active phytochrome B (APB) domain that facilitates their interaction with phyB. Only PIF1 and PIF3 contain an active phytochrome A (APA) domain, which directly binds the C-terminus of phyA, comprising two Period-Arnt-SIM (PAS) and a histidine kinase-related (HKRD) domains[13]. In FR light phyA directly interacts with PIF1 and PIF3 triggering their proteolytic destruction and/or sequestration from target promoters[13,14]. Although PIF4, PIF5, PIF8 and PIF7 lack an APA

domain, they have been shown to controls aspects of phyA signalling[15,16,17]. Interestingly, levels

of PIF8 protein are boosted by phyA. In darkness COP1 (CONSTITUTIVE

PHOTOMORPHOGENESIS 1) promotes PIF8 protein degradation through the 26S

proteasome, thereby reducing its abundance. Conversely, under FR light, phyA inhibition of

COP1 stabilises the PIF8 protein, forming a negative feedback loop that limits phyA action[15].

Notably, PIF7, a close homologue of PIF8, is a potent regulator of SAS. In unshaded conditions,

PIF7 interacts with active phyB (Pfr) through its APB domain, and both proteins localise to

sub-nuclear photobodies. This interaction prevents PIF7 from binding to the promoters of target

genes, effectively inhibiting PIF7-regulated gene expression[18,19]. Under shade conditions, the

deactivation of phyB releases PIF7 from this repression, leading to the activation of shade-

responsive genes and triggering the physiological SAS response[20,21]. Furthermore, exposure to

shade triggers the dephosphorylation and nuclear accumulation of PIF7, processes that enhance

the activation of SAS[20]. Given its strong promotory role in SAS, it is notable that PIF7 has

recently been demonstrated to increase *PHYA* transcript abundance, an action that could

potentially limit SAS. This PIF7 action can be inhibited by *PHYA UTR ANTISENSE RNA*

(PUAR), which binds to PIF7 and prevents its interaction with the 5' UTR, thereby reducing

PHYA transcript levels[17]. Collectively, this research underscores the importance of PIF7 as a

key agent in executing and moderating the molecular response to shade.

Our study substantially deepens our understanding of PIF7's role in modulating phyA signalling.

We have demonstrated that overexpression of PIF7 (PIF7ox) can completely negate the

suppressive effects of phyA on seedling hypocotyl elongation in persistent shade. Under these

conditions, PIF7 and phyA co-localize in nuclear photobodies, where they directly interact. This

interaction is facilitated by the prion-like domain of PIF7 and the C-terminus of phyA.

Additionally, our results reveal that PIF7 crucially governs the subcellular localisation of phyA,

a process mediated by the chaperone proteins FHY1 and FHL. These findings elucidate the

molecular mechanism whereby PIF7 concurrently limits phyA nuclear transport and signalling.

RESULTS

PIF7ox blocks phyA action

PIF7, which functions downstream of phyB, also shares significant sequence similarity with PIF8, a regulator of phyA responses. This similarity led us to investigate the potential genetic interaction between PIF7 and phyA. To do this we assessed hypocotyl length in seedlings grown in 8h white light (WL): 16h dark (D) or in 8h low R:FR ratio (0.2) light (WLFR) photoperiods, which commenced one day after germination to elicit a strong phyA response (Fig. 1a). As expected, wild type (WT, *Col-0*) seedlings were slightly shorter in WLFR compared to WL, while *phyA-211* had longer hypocotyls in WLFR due to reduced control of shade-avoidance elongation (Fig. 1b-c). In contrast, the phyAox line displayed a constitutively short hypocotyl and retained WLFR-responsivity. In WL the *pif7-1, pif7-2* loss-of-function alleles were marginally shorter than the WT but retained a response to WLFR that was absent in the *pif7-1; phyA-211* double mutant. In keeping with PIF7's function in promoting the phyB-mediated shade response, under WLFR, *pif7-1* diminishes the impact of *phyA-211* on hypocotyl length. Conversely, hypocotyls of 35S:PIF7-FLASH (PIF7ox) seedlings were elongated in WL and exhibited further elongation under WLFR. As hypocotyl lengths were similar in PIF7ox and PIF7ox;*phyA-211,* this suggests PIF7ox facilitates increased shade-avoidance elongation while simultaneously suppressing phyA action (Fig. 1c)[18]. This observation infers PIF7 may participate in phyB to phyA cross-regulation.

Given the capacity of PIF7ox to inhibit phyA activity, we sought to investigate the possibility of functional redundancy with other PIFs. As anticipated, PIF8ox also lacked WLFR-mediated growth inhibition, but PIF3ox, PIF4ox and PIF5ox all retained this response (Fig. S1a-b)[15]. Our data therefore illustrate that even though PIF3, PIF4 and PIF5 have been implicated in *PHYA* regulation and signalling, their functions diverge from PIF7 and PIF8[16,22,23]. We found the hypocotyl lengths of *pif8-1* and *pif8-1;phyA-211* were qualitatively similar to *pif7-1, pif7-2*, and *pif7-1;phyA-211*, respectively (Fig. S1c). However, comparison of the *PIF7ox;phyA-211* and *PIF8ox;phyA-211* lines reveal distinct differences in PIF7 and PIF8. While PIF7ox is epistatic to *phyA-211*, this genetic relationship shifts for PIF8, particularly under WLFR, where we show *phyA-211* is epistatic to PIF8ox. This finding aligns well with evidence showing PIF8 functions

downstream of phyA[15].

In summary, our findings show PIF7 has a distinct role in antagonising the suppression of

hypocotyl growth by phyA, a phenomenon particularly evident in the PIF7ox lines, which

effectively block this response.

PIF7 moderates phyA protein abundance

PIF7 is known to promote shade induction of *PHYA* expression by binding to 5' UTR[17].

Concurring with this study, we found that when sampled one hour before dusk (ZT7) in an

8L:16D photoperiod, WLFR enhanced *PHYA* expression in the WT, whereas this response was

diminished in the *pif7-1* mutant. Likewise, PIF7ox exhibited elevated *PHYA* levels in control

conditions which were further enhanced in WLFR (Fig. 1d). Immunoblot analysis using an

antibody against native phyA, reported phyA levels to be consistently low in WL, across all

genotypes. We observed that WLFR led to an increase in phyA protein levels in WT, *pif7-1* and

PIF7ox, which qualitatively corresponded with the patterns seen in *PHYA* mRNA data (Fig. 1e-

f). Further, we found that PIF7ox had a negligible effect on the phyApr:phyA-nLUC (phyA-

nLUC) in WL, which is known to gradually deplete under these conditions. In-line previous

studies, we observed higher levels of daytime phyA-nLUC in WLFR, which was notably

elevated by PIF7ox (Fig. 1g)[3]. Collectively, these findings demonstrate that under WLFR

conditions, PIF7 regulates *PHYA* expression and the abundance of phyA protein. Interestingly,

despite PIF7ox accumulating significantly higher levels of phyA under WLFR, PIF7ox

seedlings fail to elicit a phyA response. This indicates that the regulatory effect of PIF7 on phyA

activity does not stem from changes in phyA abundance.

To examine the *in vivo* dynamics of PIF7 under our experimental conditions we generated

PIF7pr:PIF7-nLUC (PIF7-nLUC) lines. We discovered that PIF7-nLUC exhibits rhythmic

patterns, with higher levels during the day compared to the night (Fig. S2 a). WLFR boosted

daytime levels of PIF7-nLUC levels, with the *phyA-211* mutation elevating PIF7-nLUC toward

the end of the period, especially in WLFR (Fig. S2 b). Therefore, our data indicate that similar

to phyA, WLFR increases the abundance of PIF7, and that phyA plays a minor role in reducing

PIF7 production toward the end of the photoperiod.

PIF7 can interact with phyA in nuclear photobodies

To establish whether PIF7 and phyA interact *in planta* we first used 35S:PIF7-sfGFP (PIF7-

GFP) and 35S:phyA-RFP (phyA-RFP) transiently expressed in *N. bethamina* leaves to establish

whether PIF7 are found in the same cellular location. We found that following WLFR treatment

PIF7-GFP and phyA-RFP co-localise in nuclei and in sub-nuclear foci, or photobodies (Fig. 2a).

Quantitative fluorescence measurements along a nuclear transect, intersecting two photobodies,

clearly shows this co-localisation (Fig. 2a). Next, we created split bimolecular fluorescence

complementation (BiFC) constructs 35S:PIF7-nYFP and 35S:phyA-cYFP, to test whether PIF7

and phyA physically interact *in vivo*. In WLFR, BiFC assays yielded strong nuclear, and sub-

nuclear foci signals, but not in control lines expressing 35S:PIF7-nYFP and 35S:EV-cYFP. The

data suggest PIF7 and phyA are indeed able to interact in nuclei and within photobodies under

WLFR (Fig. 2b).

To gain further evidence for this interaction we also generated split nanoLUC (nLUC) luciferase

reporter constructs, 35S:PIF7-lgbitnLUC (PIF7-lgbit) and 35S:phyA-smbitnLUC (phyA-

smbit), with 35S:phyB-smbitnLUC (phyB-smbit) and 35S:phyA-nLUC (phyA-nLUC) as

controls (Fig. 2c). As expected, we found a strong signal in the PIF7-lgbit+phyB-smbit in WL

which is diminished in WLFR, which is consistent with the known light conditional interaction

between PIF7 and phyB[18]. Again, as expected we observed a strong signal from phyA-nLUC

in WLFR, which is known to increase phyA abundance, while negative controls (phyA-smbit+

phyB-smbit) had low signal in both conditions. In WL conditions, the signal from PIF7-lgbit +

phyA-smbit was low, similar to that of the negative control, whereas in WLFR conditions, the

signal was strong, indicating an interaction. Further, using an antibody to native phyA we

demonstrated co-immunoprecipitation of phyA with PIF7ox, in WLFR but not WL conditions

(Fig. 2d). Thus, our data provides evidence that PIF7 and phyA directly interact in photobodies

after exposure to WLFR which promotes their respective nuclear accumulation[20,24].

Prion-like Domain of PIF7 interacts with phyA PAS-PAS Domain

Earlier work demonstrated that phyA binds directly with PIF1 and PIF3. The phyA C-terminal

output module (OPM) is known to be important for this interaction, as is the active phyA-

binding domains (APA) of PIF1 and PIF3[13,22]. To identify PHYA regions that required for the

molecular interaction we utilised PIF7-nYFP and phyA-cYFP split YFP assay, creating

truncated PHYA constructs as follows: C-terminal (phyA-C), comprising HINGE (H), PAS-

PAS DOMAIN (PAS), and HISTIDINE KINASE RELATED DOMAIN (HKRD) regions

[H+PAS+HKRD-cYFP], phyA-C1 [PAS+HKRD-cYFP], phyA-P [PAS-cYFP], and phyA-H

[HKRD-cYFP] (Fig. 3a). In WLFR we obtained nuclear fluorescence signals for PIF7-nYFP

and the full-length phyA-cYFP control, as well as the truncated phyA-C, phyA-C1, and phyA-

P constructs. No signal was observed for phyA-H and the negative EV control. The strong signal

detected for phyA-P, which includes PAS-PAS domain motifs, highlights the significance of

this molecular region in facilitating the interaction with PIF7. To identify the region of PIF7

that mediated the interaction with phyA, we generated PIF7-N which contains the ACTIVE

PHYTOCHROME BINDING domain [PIF7-N-nYFP], PIF7-B, containing the BASIC

HELIX-LOOP-HELIX motif [PIF7-N-nYFP], and PIF7-C which contains a Prion-like domain

(PrLD) [PIF7-C-nYFP] which has a polyQ-rich sequence is located at PrLD center (Fig. 3b,

Fig. S6). Interestingly we found that only the PIF7 full length control and PIF7-C gave nuclear

fluorescence in WLFR. Thus, our data point to the prion-like domain of PIF7 as being important

for its interaction with phyA. Since PIF7 lacks an APA domain, it seems that prion region is

able to fulfill this role.

PIF7 regulates the subcellular localisation of phyA

Our data show PIF7ox elevates phyA protein levels, yet PIF7ox lines are unable to mount a

phyA response in WLFR conditions. As WLFR drives the phyA HIR and its nuclear

accumulation, we sought to analyse cellular location more closely. As anticipated, confocal

microscopy revealed strong cytosolic and weaker nuclear signals for phyA-GFP

(phyApr:phyA-sGFP), and for GFP-PIF7 (35S:GFP-PIF7) when in WL (Fig. S3). Exposure to

WLFR notably enhanced nuclear relative to cytosolic signals, confirming the nuclear

accumulation of both PIF7 and phyA under our shade conditions (Fig. S3a)[19,20,25]. We then

aimed to establish if PIF7 affected the location of phyA by introgressing phyA-GFP into *pif7-*
1 and PIF7ox backgrounds. In WL we found phyA-GFP to be predominantly in the cytosol in
Col-0, *pif7-1* and PIF7ox lines (Fig. 4a). Under WLFR, we recorded a higher proportion of
nuclear phyA-GFP in WT, especially in *pif7-1*, while significantly lower levels were observed
in PIF7ox (Fig. 4b-c). Thus, PIF7 appears to influence the cytoplasmic to nuclear partitioning
of phyA-GFP. To verify our results, we performed immunoblot assays that differentiated
between nuclear and cytosolic fractions. This method yielded qualitatively similar results
showing increased nuclear phyA in *pif7-1* and predominately cytosolic PHYA in PIF7ox (Fig.
4d-e). Thus, our findings indicate that PIF7 modifies subcellular phyA localization. The notably
reduced nuclear presence of phyA in PIF7ox could partially account for the disrupted phyA
signaling observed in PIF7ox.

PIF7ox blocks phyA regulation of target genes

To further interrogate the mechanism through which PIF7 inhibits phyA action we initially
tested whether PIF7 alters the expression of genes (*CHS, BIN2, FHY3, FHY1, FHL* and *IAA29)*
known to be directly regulated by phyA (Fig. 5a)[26,27]. As determined by qPCR all the genes,
except *IAA29*, were upregulated in WT in WLFR conditions, otherwise, the expression patterns
fell into two categories. In WLFR conditions, *CHS, BIN2,* and *FHY3* exhibited decreased
expression in both *phyA-211* and PIF7ox, while their expression was increased in *pif7-1*. This
gene set is upregulated by phyA in WLFR, and their expression is repressed by PIF7. On the
other hand, *FHY1, FHYL* and *IAA29* form a group that are upregulated in *phyA-211* and PIF7ox,
and for *FHY1* and *FHYL*, expression is lowered in *pif7-1*. These genes are downregulated by
phyA in WLFR, and their expression is promoted by PIF7. The common factor for all these
genes is that phyA and PIF7 exert opposing control of their regulation, particularly in WLFR.

We then sought to establish if the antagonistic action of PIF7 was mediated through direct
repression of phyA at target gene promoters. Here we performed chromatin
immunoprecipitation (ChIP) using a native phyA antibody, and qPCR using to detect phyA
enrichment at *CHS, FHY1*, and *FHL* promoters (Fig. 5b). For all three genes we found WLFR
stimulated phyA enrichment at promoter regions sequences containing G-box motifs (two in

CHS), but not in G-box-free regions (Control, coding region). This enrichment was eliminated

in the presence of PIF7ox, suggesting the interaction between PIF7 and phyA is able to block

phyA regulation of target genes.

PIF7 interacts with FHY1 and FHL

In WLFR phyA cytosolic to nuclear shuttling is controlled by FHY1 and FHL chaperones[9]. The

central importance of this mechanism for phyA function is evident in the *fhy1-3;fhl-1* double

mutant hypocotyl response which is phenotypically similar to *phyA-211* (Fig. S4). An intriguing

finding of our study is that PIF7 appears to control the subcellular location of phyA. We

therefore wanted to establish if this arose through a process involving PIF7 interaction with

FHY1 and FHL. Using the *N. benthamiana* transient expression system, we recorded nuclear

interactions between the BiFC constructs 35S:FHY1-cYFP, 35S:FHL-cYFP, and 35S:PIF7-

nYFP under WLFR conditions (Fig. 6a). We also observed PIF7 interactions with FHY1 and

FHL, but not the negative control using a split nLUC assay (Fig. 6b). It is worth noting that in

this assay the interaction signal strength between the shuttle proteins and PIF7 was lower than

that for PIF7 and phyA (Fg. 6b). However, we were able to verify these interactions through

co-immunoprecipitation using 35S:FHY1/FHL-nLUC and 35S:PIF7-sfGFP (Fig 6c). As our

data shows PIF7 is able to interact with phyA, FHY1 and FHL, we postulated that these proteins

may form a complex under inductive WLFR conditions. Supporting this hypothesis, we found

that when FHY1-YFP or FHL-YFP are co-expressed with PIF7-sfGFP and phyA-RFP in *N.*

benthamiana leaves, these constructs localise to the same nuclear regions under WLFR (Fig.

6d-e).

Another factor worth considering is *ARABIDOPSIS SUMO PROTEASE 1 (ASP1)*, which plays

an important role in SUMOylation and maintaining FHY1 levels under FR light conditions[28].

We observed increased levels of *ASP1* expression in PIF7ox compared to WT in WL. We found

ASP1 levels are enhanced by WLFR, a response that is further amplified by *pif7-1* and

suppressed by PIF7ox (Fig. 6f). Thus, PIF7 regulation of *ASP1* may contribute to its regulation

of the shuttling mechanism.

To determine if PIF7 primarily acts by sequestering phyA or whether the PIF7 interactions with

FHY1/FHL are also important, we employed phyApr:phyA-sGFP-NLS (phyA-NLS), where

phyA is constitutively nuclear (Fig S7)[29]. In line with earlier findings, under our experimental

conditions, phyA-NLS restores the *phyA-201* elongated hypocotyl phenotype to WT length (Fig.

6g). We hypothesised that if PIF7 regulation of phyA was solely due to its sequestration of

phyA, then PIF7ox should diminish the function of phyA-NLS through this mechanism[29].

However, our data show phyA-NLS exhibits strong epistasis over PIF7ox, for hypocotyl length,

and the expression of *CHS* and *IAA29* under WLFR (Fig. 6g, S5). Consequently, it seems that

the mechanism of phyA sequestration alone likely cannot explain the potent effects of PIF7ox

on phyA action. Previous research demonstrated that phyA-NLS completely restores phyA

control of hypocotyl growth in *fhy1-3;fhl-1 and phyA-211;fhy1-3;fhl-1*, indicating that phyA-

NLS can bypass the requirement for nuclear shuttling[30]. Consequently, our results imply that

the interaction between PIF7 and FHY1/FHL may be required to suppress phyA action. Under

WLFR, the transport of phyA into the nucleus is highly dynamic. Therefore, PIF7 may interact

with phyA-FHY1/FHL complexes soon after their nuclear entry and before the complexes

disassemble. This scenario is consistent with our co-localisation data (Fig.6d)[9]. Collectively,

our data indicates that PIF7 forms inactive complexes with phyA and FHY1/FHL, which

simultaneously impedes phyA shuttling and signalling.

DISCUSSION

The operating properties of phyA are distinct from other phytochromes. Under shaded

conditions, phyA requires a cytosolic to nuclear shuttling mechanism to operate effectively, a

process facilitated by the chaperone proteins FHY1 and FHL. In this study, we have shown

PIF7 is a potent regulator of this process, limiting the availability of phyA for signalling. The

impact on phyA is most extreme in the PIF7ox line, where WLFR-induced phyA action is

blocked. Our research illustrates that PIF7 moderates phyA intracellular movement and

signalling through molecular sequestration.

Studies have shown that both PIF7 and phyA accumulate in the nucleus under FR-rich shade

conditions[31,18,29]. Our findings further reveal that PIF7-GFP and phyA-RFP co-localise to

specific sub-nuclear foci, where they physically interact as evidenced by BiFC (Fig.2a-b).

Additional confirmation of this interaction was provided by split nLUC assays employing PIF7-

lgbit and phyAsmbit, along with co-immunoprecipitation of PIF7ox and phyA.

While all PIFs have an active phytochrome B (APB) domain that enables binding to phyB, only

PIF1 and PIF3 have active phytochrome A (APA) motifs, through which they interact with the

phyA C-terminus[15,32]. This binding is known to facilitate the proteolysis of PIF1 and PIF3, and

to inhibit the activity of PIF3 by isolating it from target promoters[15,32,33]. Our research has

demonstrated that although PIF7 lacks an APA domain, it can still directly interact with phyA.

This binding is facilitated through the PAS domain of phyA and the C-terminal region of PIF7,

which contains a prion-like domain. PIF7 moderates the activity of phyA by inhibiting its ability

to bind to target promoters (Fig. 5b). For the genes *CHS*, *FHY3*, and *BIN2*, which are activated

by phyA in WLFR, and for *FHY1*, *FHL*, *IAA29*, which are repressed by phyA, PIF7 counteracts

the effects of phyA.

PIF7 is particularly effective in FR-rich shade, which triggers its dephosphorylation, increases

its abundance, and enhances its accumulation in nuclear photobodies (Fig. S3b)[20]. Shade also

deactivates phyB, which typically limits the DNA-binding capacity of PIF7[19]. Since continuous

shade also stimulates phyA activity, it creates conducive conditions for the interaction between

PIF7 and phyA.

We have demonstrated that PIF7 and phyA interact within sub-nuclear structures known as

photobodies[5]. These photobodies are essentially molecular condensates that form through

liquid-liquid phase separation (LLPS), accompanied by a nucleation step. Nucleation aids

sequestration, particularly at low concentrations, and targets the formation of photobodies to

specific sub-nuclear sites[6,34]. Across organisms, the spatial organisation of biomolecules into

distinct cellular locations is essential for enabling specialised functions[35].

PIF7 is unique amongst PIF proteins as it contains a prion-like domain with a central polyQ

stretch of 17 consecutive glutamine residues. PIF8 is the closest in sequence similarity to PIF7,

and while does possess a short polyQ sequence in its C-terminus, this domain exhibits low

prion-like predictability (Fig.S6). Prion-like domains (PrLD) typically comprise low-

complexity or intrinsically disordered regions that facilitate protein condensation and LLPS.

Indeed, PIF7 is able to form condensates with LLPS characteristics *in vitro*, which is proposed

to facilitate the formation of PIF7-phyB photobodies[19]. Substituting polyQ with polyA

diminishes the capacity of PIF7 to form photobodies, highlighting the importance of polyQ in

this process. Interestingly, phyB has also been shown to possess LLPS characteristics, for its

C-terminus, and disordered N-terminal extension have both been shown to phase-separate in

mammalian cells[36]. We have shown that in WLFR, PIF7 interacts with the phyA, through its

PrLD, which results in the formation of PIF7-phyA-containing photobodies. Within these

biomolecular condensates, phyA-mediated signalling and control of hypocotyl growth are

suppressed. In the PIF7ox line, this inhibition results in excessive elongation of the hypocotyl

under WLFR, since PIF7 action is no longer constrained by phyA.

The movement of phyA to the nucleus in WLFR is strongly contingent on the chaperone

proteins FHY1 and FHL7. Through the use of BiFc, split nanoLUC and co-

immunoprecipitation assays, we demonstrated that PIF7 also interacts with FHY1 and FHL

under WLFR. Fluorescent tagged versions of PIF7, phyA and FHY1/FHL illustrated that these

proteins co-localise within nuclei, suggesting the formation of PIF7-phyA-FHY1/FHL

complexes. Nuclear shuttling of phyA is a highly dynamic process that operates continuously

under WLFR conditions[9,37]. Earlier work indicates that dissociation of the phyA-FHY1/FHL

complex is required to both recycle the shuttle proteins and to facilitate phyA signalling[31,37].

We therefore propose that PIF7 may serve to trap phyA and FHY1/FHL in inactive complexes,

which simultaneously perturbs the nuclear transport of phyA and disconnects it from

downstream signalling. Consistent with this interpretation, we find *pif7-1* and PIF7ox have

higher proportions of nuclear, and cytosolic phyA, respectively (Fig. 4b-e). Further, our genetic

data shows that phyA-NLS, which is constitutively nuclear, and bypasses the FHY1/FHL

shuttling mechanism, can restore phyA function to PIF7ox in WLFR, in terms of hypocotyl

phenotype and *CHS* and *IAA29* gene expression (Fig. 6g, Fig. S5). We also discovered that PIF7

reduces the expression of *ARABIDOPSIS SUMO PROTEASE 1 (ASP1)*, which aids the

deSUMOylation of FHY1 under FR light, thereby preventing its degradation. *ASP1* suppression

may, therefore, represent an alternative mechanism through which PIF7 modulates phyA

shuttling. Previously, our research demonstrated that under simulated shade, PIF7 interacts with and suppresses the transcriptional activity of ANGUSTIFOLIA 3 (AN3), a promoter of leaf cell proliferation[38]. In this case, PIF7 operates by evicting and substituting for AN3 at target promoters. Thus, in shade conditions, PIF7 assembles different nuclear complexes which execute distinct functions.

In summary, our study outlines the central role PIF7 plays in regulating phyA activity. Its direct binding to phyA, and the FHY1/FHL chaperones, likely represents a rapid, energy-efficient, and highly effective way to stem phyA action. The compartmentalisation of phyA and FHY1/FHL into PIF7 photobodies constitutes a dynamic regulatory mechanism that responds to changes in environmental light quality. This subcellular partitioning enables PIF7 to drive hypocotyl elongation while simultaneously reducing the suppressive influence of phyA. Since the modulation of PIF7 by phyB is well-established, our findings position PIF7 at the molecular intersection of phyB and phyA, underscoring its pivotal role in adjusting phyA activity to promote SAS elongation under vegetative shade[19].

METHODS

Plant material and growth conditions

The wild-type of *Arabidopsis thaliana* seedlings are Columbia-0 (*Col-0*). The *pif7-1*, *pif7-2*[39], *pif8-1*, *pif8-2* mutants, and the construction of *pif8-1;phyA-211*, *PIF8ox;phyA-211* and PIF8OX[15], PIF7OX-FLASH[40], 35S:GFP-PIF7[20], 35S:PIF3-LUC[41], 35S:PIF5-HA, 35S:PIF4-LUC[16], phyApr:phyA-sGFP, phyApr:phyA-sGFP-NLS[25], and phyApr:phyA-nanoLUCx3FLAGx10His (phyA-nLUC)[3] have been described previously. The *phyA-211* mutant was obtained from NASC collection. PIF7pr:PIF7-nanoLUCx3FLAGx10His (PIF7-nLUC) (Basta) were transformed into the *pif7-1* background. The *pif7-1*;phyA-sGFP-NLS, and PIF7ox;phyA-sGFP-NLS, *pif7-1;phyA-211* and PIF7ox;*phyA-211* lines were generated by genetic crossing. Likewise, the *pif7-1;*phyA-nLUC, PIF7ox;phyA-nLUC, PIF7pr:PIF7-nLUC; *phyA-211* lines were made through genetic crossing. All the cross lines were verified using PCR or sequencing.

For the hypocotyl growth experiments seeds were first sterilized and then re-suspended in 0.1% (w/v) agar and stored in darkness at 4°C for 3 d to stratify. Following this, seeds were individually sown onto plates containing half-strength Murashige and Skoog basal salts mixture (Duchefa Biochemie, Haarlem, Netherlands) dissolved in 1% (w/v) agar (1/2 MS media; pH 5.8). The plates were then transferred to Percival I30BLL growth chambers at 22 °C (Plant Climatics, Wertingen, Germany). To promote germination seeds were exposed to a 4 h period of white light (WL) (85 ± 5 μmol m^{-2} s^{-1}), and then darkness for 20 h, before transferring to experimental photoperiods. For all experiments seedlings were grown under short day (8L:16D) conditions in either WL or WL with supplementary FR (WLFR) to provide a R:FR ratio of 0.20.

RNA extraction and q-PCR

For RNA extraction seedlings (n = 36 ± 5 seedling numbers), were harvested at Zeitgeber time 7 (ZT) which is one hour before dark. Total RNA was extracted using a phenol-based method[38], 300 μL RNA extraction buffer (PH=8.0) was added to each sample, followed by 300 μL 1:1 acidic phenol: chloroform. The supernatant was separated from the pellet by centrifuging for 15 minutes at 13,000 rpm under 4°C. Up to 300 μL of the supernatant was transferred to collection tubes with 240 μL isopropanol and 30 μL 3 M sodium acetate. Samples were precipitated at -80°C for 15 minutes, then centrifuged for 30 minutes at 13,000 rpm. Pellets were washed with 300 μL of 70% ethanol and then suspended in nuclease free water and stored at -80°C. Complementary DNA (cDNA) synthesis was performed using the qScript cDNA SuperMix (Quanta Biosciences) as described by the manufacturer. A Lightcycler 480 system (Roche) was employed for real time-qPCR, with 10 μL reactions that incorporated the SYBR Green Master Mix (Roche). The q-PCR primers are shown in **Supplementary table 2**. Results were analysed using Light Cycler 480 (Roche).

Hypocotyl measurements and statistical analysis

Images of hypocotyls were taken after a period of 6 days growth, and hypocotyl length was analysed using ImageJ (https://imagej.nih.gov/ij/). Data analysed by Excel and GraphPad Prism 10. Data are presented as mean values ± s.e.m. Two-way ANOVA (α=0.05) was conducted to assess statistical significance across light conditions for each genotype; asterisks denote significance relative to control treatments: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), P < 0.0001 (****). Representative data from three independent replicates are shown.

Construct assembly

The full-length CDS of *phyA* and nanoLUC tag (NLUCx3FLAGx10His; NL3F10H) have been described previously[3]. The CDS of *PIF7* were synthesised by the GeneArt Gene Synthesis

service (Invitrogen, MA, USA). The split nanoLUC tags were made by splitting nanoLUC to lgbit (158 aa) and smbit (11 aa). Mobius Assembly[42] was used to insert the constructs into a mUAV plasmid (L0, CamR) with *AraI* restriction site and plasmid assembly was performed and subsequently verified by the Edinburgh Genome Foundry. The full-length CDS of *FHY1* and *FHL* parts were amplified with Q5 High-Fidelity DNA polymerase (New England Biolabs, MA, USA) from the *Col-0* genome by gene-specific primers. The phyA-C, phyA-C1, phyA-P, phyA-H, PIF7-N, PIF7-B and PIF7-C part cloned on the CDS region of *phyA* and *PIF7* (L0 parts). These parts were assembled into a mUAV plasmid and sequenced by Source Bioscience (Cambridge, UK). All L0 parts will assemble in pMAP B L1 (KamR) vector with *BsaI* restriction site. All the constructs for the tobacco (*Nicotiana benthamiana*) assay used a CaMV35S promoter, which is synthesized by the GeneArt Gene Synthesis service (Invitrogen, MA, USA).

To construct pEXP601- PIF7pr:PIF7-nanoLUCx3FLAGx10His, a 2740 bp genomic fragment was generated, encompassing the PIF7 promoter region (1015 bp), 5'UTR (161 bp), and 1564 bp ORF, excluding the stop codon of exon 6 of the AT5G61270.1. This fragment was PCR-amplified using specific primers with Phusion High-Fidelity DNA Polymerase (NEB, M0530S). The resulting amplicon was subsequently recombined into pDONR221 using BP II clonase (Invitrogen, 12535029), and its sequence verified by Sanger sequencing (Genepool, Edinburgh Genomics, Edinburgh, UK). The genomic regions were then recombined into pGWB601: NanoLUC-3FLAG-10His sing LR Clonase (Invitrogen)[43]. The final construct was transformed into Agrobacterium AGL1. Primer used for construct assembly are listed in **Supplementary table 1.**

The Tobacco Transient Expression Assay

The CaMV35S constructs were transferred into the Agrobacterium AGL1 strain and cultured at 28℃ for two days, before infiltration into 4-week-old *Nicotiana benthamiana* leaves. Agrobacterium expression construsts was infiltrated into *Nicotiana benthamiana* leaves with P19. The solution used for transient expression experiments contained 10 mM MES buffer (pH 5.6), 10 mM $MgCl_2$ and 200 mM aceto-syringone. Plants were grown under short day (8L:16D, at 22℃) conditions in either WL or WL with supplementary FR (WLFR) to provide a R:FR ratio of 0.20. Tobacco will used for next step after 3 days treatments.

Confocal Imaging, Colocalization, and BiFC

The fluorescence images of GFP, YFP and RPP expression were obtained with a Leica confocal micro-scope (Leica SP8) at 488 and 552 nm laser. For BiFC and co-localization assay all images

were taken at ZT7 on day 3 after WL/WLFR treatments. The co-localization signals were analysed by LAS X Office 1.4.7.2982 and plotted by GraphPad Prism 10. phyApr:phyA-sGFP(NLS), *pif7-1*;phyA-GFP, PIF7ox;phyA-GFP and 35S:GFP-PIF7 transgenic lines images were taken at ZT7 within 30 minutes after 6 days of WL/WLFR treatments. The Nuclear and cytosolic signals were quantified from cells in the mid-hypocotyl region within a 20 x 1.36 objective lens. Nucleus and cytoplasmic signals were quantified using Image J and plotted in GraphPad Prism 10. Imaging experiment comprised three biological replicates, with 6 seedlings in each replicate.

Immunoblot Analysis and Co-immunoprecipitation Assays

Each sample comprised 0.5-1 g of Arabidopsis seedling tissue, harvested on day 6 at ZT7. Samples were added to lysis buffer (125 mM Tris-HCl (pH 7.5); 0.25mM EDTA (pH 7.5), 150 mM NaCl, 5% glycerol, 0.1% Triton X-100, 0.5% NP-40), which contained 1X Protease inhibitor (Sigma-Aldrich, P9599) and 1x HALT-Phosphatase inhibitor (Thermo Scientific). Samples were centrifuged at 12,000 rpm for 10 mins at 4ºC and then transferred to lysate mix with 1x SDS loading buffer (containing 0.1 M DTT) at a 1:1 volume ratio. Samples were incubated at 80ºC for 10 mins, followed by a centrifugation step, prior to western blotting.

For Co-immunoprecipitation, after lysis an aliquot of the total protein was saved as an input sample. Then other total protein extracts were incubated with Protein A/G beads (MCE Magnetic Agarose) for 1 h at 4 C to remove adventitious binding proteins. Then anti-FLAG/His beads (Anti-FLAG M2 agarose gel Sigma-Aldrich, Cat.No. A2220; Anti-His Magnetic Beads, Cat. 209, MCE) added into treated protein extracts over night at 4ºC. The antibody/protein complex was washed with wash buffer (125 mM Tris-HCl [pH 7.5]; 0.25mM EDTA [pH 7.5], 150 mM NaCl, 5% glycerol, 0.1 % Triton X-100) 5 times. Then added 50 μL of 1x SDS loading buffer to each sample and incubate at 80ºC for 8 mins.

To detect phyA protein, an anti-Phytochrome A antibody (PHYTOAB, PHY1907) was used at 1:1,000. To detect the PIF7OX-FLASH and 35S:FHY1/FHL-nanoLUC x3FLAGx10His an anti-FLAG M2 antibody (Sigma-Aldrich, F1804) was used at 1:2,000. To detect the 35S:PIF7-sfGFP an anti-GFP anti-body (Roche, ab290) was used at 1:3,000. Anti-RPN6 antibody (Agrisera, AS121858) was used at 1:3,000 as internal control. Secondary antibodies (1:3,000, anti-mouse IgG (cell signalling, 7076s); 1:4,000 Goat anti-rabbit-IgG (Abcam, AB6013) were used for the ECL Prime Western Blotting System. The signals were detected by Azure 300 Imager with ECL prime western blotting detection reagents (Cytiva, RPN2232).

NanoLUC and split nanoLUC assays

NanoLUC bioluminescence assays were conducted as previously described[3]. Add in each well

of. Single Arabidopsis seedlings were grown in individual wells, in a 96 well plate, each

containing 200 µl 1/2 MS media. Plates were sealed with cling film and placed under 8L:16D

(WL) for 6 days. Before imaging 50 µl diluted (1:50) of Nano-Glo (Promega, Nano-

Glo®Luciferase Assay) was added to the seedlings in each well. Plates were sealed and moved

to experimental light conditions. Bioluminescence was recorded every hour (for 1.5 s) for

several days using the TriStar2 LB942 multimode microplate reader (Berthold Technologies,

Germany). The data was analysed and plotted using BioDare2 software[44] and Excel. Typically,

9 biological repeats were conducted with at least 6 seedlings per replicate in **Fig.1g**.

The split nanoLUC assay is like BIFC assay method. Through Agrobacterial infiltrated into

Nicotiana benthamiana leaves. After 3 days WL/WLFR treatments, the 1cm x 1cm size leaves

will be used in 96 wells plates (*n* =12 leaf numbers) for collecting signals. Next add 100 µl

diluted (1:50) Nano-Glo on the top of the leaves to each well. Then the signals will collect by

plate reader. The data analysed by Excel and GraphPad Prism 10. The experiments repeated at

least 3 times and had similar results.

Chromatin Immunoprecipitation (ChIP)-qPCR

We employed ChIP-qPCR methods described by previews work, as follows[45]. For each sample

0.5-1g of tissue was extracted from 6 day -old seedlings at ZT7. Cross-linking was induced

with 1% formaldehyde under vacuum infiltration for 15 mins (up to -70 kPa), then 2.5 ml of 2

M glycine was added to quench crosslinking. DNA was fragmented into an average size of

~150 bp in Diagenode BioRuptor Plus, (30 cycles of high power 30s ON / 30s OFF). Anti-

phyA antibody (PHYTOAB, PHY1907,1:300) incubated with protein A/G beads (MCE

Magnetic Agarose) in 4℃ overnight. And the following day samples were incubated with

protein A/G bead/phyA antiboday complex for 6 hours at 4℃. DNA was extracted using

phenol/chloroform (1:1, vol/vol). Protein enrichment at DNA regions was quantified using RT-

qPCR using primers detailed in **Supplementary table 2.** Enrichment was determined

by %Input method (%Input=2 ^ [CtInput-CtChIP] × IDF × 100%).

Nuclear-Cytoplasmic Fractionation

Nuclear fractionation was performed essentially as previously described with the following

modifications[46]. Briefly, 0.5-1 g of Arabidopsis seedlings ground into fine powder in liquid

nitrogen and then homogenised in double the volume of pre-cooled (4℃) lysis buffer (20 mM

Tris-HCl, pH 7.4, 25% Glycerol, 20 mM KCl, 2 mM EDTA, 2.5 mM MgCl2, 250 mM Sucrose

and 5 mM DTT, add 1×protease inhibitor cocktail (Roche) and 1 x HALT-Phosphatase inhibitor

added in fresh). The sample was filtered twice through two layers of Mira cloth (Merck

Millipore). The flow-through was spun 10 min at 2000g, 4°C. Supernatant, consisting of the cytoplasmic fraction, was centrifuged 20 min at 13,000g, 4°C and collected. The pellet, containing the nuclear fraction, was washed six times with 2.5 mL of nuclear resuspension buffer NRBT1 (20 mM Tris-HCl, pH 7.4, 25% Glycerol, 2.5 mM $MgCl_2$, 0.2% Triton X-100) and centrifuged 3 min at 1,500g, 4°C. The pellet was resuspended in 300 µL of pre-cooled NRBT2 (250 mM Sucrose, 10 mM Tris-HCl, pH 7.5, 10 mM MgCl2, 1% Triton X-100, 0.035% β-mercaptoethanol ,and 1× protease inhibitor cocktail added in fresh), and carefully overlaid on top of 300 µL NRBT3(1.7 M Sucrose, 10 mM Tris-HCl, pH 7.5, 2 mM MgCl2, 0.15% Triton X-100, 0.035% β-mercaptoethanol, and 1× protease inhibitor cocktail added in fresh). These were centrifuged at 16,000g for 55 min at 4°C, and the final nuclear pellet was resuspended in 2× SDS loading buffer. As quality controls for the fractionation, RbCL (Agrisera, AS03037, 1:3,000) protein was used as the cytoplasmic control, and histone H3 (Abcam, ab18521, 1:3,000)) was used as the nuclear control in immunoblot assays.

ACKNOWLEDGEMENTS

This research was supported by University of Edinburgh Darwin Trust Scholarship awarded to MK. Z and Leverhulme Trust grant RPG-2024-082 awarded to K.J.H. We thank Dr. Rist Van de Weyer who generated PIF7ox;*phyA-211* and first observed the PIF7ox hypocotyl phenotype in WLFR. We thank Prof. Lin Li, Prof. Dr. Andreas Hiltbrunner and Dr. Gabriela Toledo-Ortiz who kindly shared 35S:GFP-PIF7, 35S:phyA-RFP and phyApr:phyA-sGFP(NLS) seeds and thank Prof. Giltsu Choi for sharing *pif8-1;phyA-211*, *PIF8ox;phyA-211* and PIF8OX lines.

AUTHOR CONTRIBUTIONS

MK. Z and K.J.H. planned and designed the research. MK. Z performed and analysed all experiments with K.J.H. A.R. generated the PIF7pr:PIF7-nanoLUC line. MK. Z and K.J.H. both contributed to manuscript preparation.

REFERENCE

1 Liu, Y., Jafari, F. & Wang, H. Y. Integration of light and hormone signaling pathways in the regulation of plant shade avoidance syndrome. *Abiotech* **2**, 131-145 (2021). https://doi.org/10.1007/s42994-021-00038-1.

2 Franklin, K. A. Shade avoidance. *New Phytol* **179**, 930-944 (2008). https://doi.org/10.1111/j.1469-8137.2008.02507.x.

3 Butlin, P., Valdivia-Cabrera, M., Grima, R. & Halliday, K. J. Arabidopsis thaliana Phytochrome A Sensory Properties in Canopy Shade. *bioRxiv* (2025). https://doi.org/10.1101/2025.05.14.654025.

4 Mathews, S. & Sharrock, R. A. Phytochrome gene diversity. *Plant Cell Environ* **20**, 666-

671 (1997). https://doi.org/10.1046/j.1365-3040.1997.d01-117.x.

5   Cheng, M. C., Kathare, P. K., Paik, I. & Huq, E. Phytochrome Signaling Networks. *Annu*
Rev Plant Biol **72**, 217-244 (2021). https://doi.org/10.1146/annurev-arplant-080620-
024221.

6   Du, J., Kim, K. & Chen, M. Distinguishing individual photobodies using Oligopaints
reveals thermo-sensitive and -insensitive phytochrome B condensation at distinct
subnuclear locations. *Nature Communications* **15** (2024). https://doi.org/10.1038/s41467-
024-47789-1.

7   Casal, J. J., Luccioni, L. G., Oliverio, K. A. & Boccalandro, H. E. Light, phytochrome
signalling and photomorphogenesis in. *Photoch Photobio Sci* **2**, 625-636
(2003).https://doi.org/10.1039/b300094j.

8   Rausenberger, J. *et al.* Photoconversion and Nuclear Trafficking Cycles Determine
Phytochrome A's Response Profile to Far-Red Light. *Cell* **146**, 813-825 (2011).
https://doi.org/10.1016/j.cell.2011.07.023.

9   Hiltbrunner, A. *et al.* Nuclear accumulation of the phytochrome A photoreceptor requires
FHY1. *Curr Biol* **15**, 2125-2130 (2005). https://doi.org/10.1016/j.cub.2005.10.042.

10   Genoud, T. *et al.* FHY1 Mediates Nuclear Import of the Light-Activated Phytochrome A
Photoreceptor. *Plos Genet* **4** (2008). https://doi.org/10.1371/journal.pgen.1000143.

11   Li, J. G. *et al.* Transcription Factor ELONGATED HYPOCOTYL5 Plays a Role in the
Feedback Regulation of Phytochrome A Signaling. *Plant Cell* **22**, 3634-3649 (2010).
https://doi.org/10.1105/tpc.110.075788.

12   Pham, V. N., Kathare, P. K. & Huq, E. Phytochromes and Phytochrome Interacting Factors.
Plant Physiol **176**, 1025-1038 (2018). https://doi.org/10.1104/pp.17.01384.

13   Jeong, J., Lee, Y. & Choi, G. Both phytochrome A and phyB interact with
PHYTOCHROME-INTERACTING FACTORs through an evolutionary conserved
phy(OPM)-APA interaction. *Nat Commun* **16**, 3946 (2025).
https://doi.org/10.1038/s41467-025-59327-8.

14   Al-Sady, B., Ni, W., Kircher, S., Schafer, E. & Quail, P. H. Photoactivated phytochrome
induces rapid PIF3 phosphorylation prior to proteasome-mediated degradation. *Mol Cell*
23, 439-446 (2006). https://doi.org/10.1016/j.molcel.2006.06.011.

15   Oh, J., Park, E., Song, K., Bae, G. & Choi, G. PHYTOCHROME INTERACTING
FACTOR8 Inhibits Phytochrome A-Mediated Far-Red Light Responses in Arabidopsis.
Plant Cell **32**, 186-205 (2020). https://doi.org/10.1105/tpc.19.00515.

16   Seaton, D. D. *et al.* Dawn and photoperiod sensing by phytochrome A. *P Natl Acad Sci*
USA **115**, 10523-10528 (2018). https://doi.org/10.1073/pnas.1803398115.

17   Zhu, T., Yang, C., Xie, Y., Huang, S. & Li, L. Shade-induced lncRNA PUAR promotes
shade response by repressing PHYA expression. *EMBO Rep* **24**, e56105 (2023).
https://doi.org/10.15252/embr.202256105.

18   Leivar, P. *et al.* The Arabidopsis phytochrome-interacting factor PIF7, together with PIF3
and PIF4, regulates responses to prolonged red light by modulating phyB levels. *Plant*
Cell **20**, 337-352 (2008). https://doi.org/10.1105/tpc.107.052142.

19   Xie, Y. *et al.* Phytochrome B inhibits the activity of phytochrome-interacting factor 7
involving phase separation. *Cell Rep* **42** (2023).
https://doi.org/10.1016/j.celrep.2023.113562.

20 Huang, X. *et al.* Shade-induced nuclear localization of PIF7 is regulated by
phosphorylation and 14-3-3 proteins in Arabidopsis. *Elife* **7** (2018).
https://doi.org:10.7554/eLife.31636.
21 Park, E. *et al.* Phytochrome B inhibits binding of phytochrome-interacting factors to their
target promoters. *Plant Journal* **72**, 537-546 (2012). https://doi.org:10.1111/j.1365-
313X.2012.05114.x.
22 Ni, M., Tepperman, J. M. & Quail, P. H. PIF3, a phytochrome-interacting factor necessary
for normal photoinduced signal transduction, is a novel basic helix-loop-helix protein. Cell
95, 657-667 (1998). https://doi.org:10.1016/S0092-8674(00)81636-0.
23 Lorrain, S., Trevisan, M., Pradervand, S. & Fankhauser, C. Phytochrome interacting
factors 4 and 5 redundantly limit seedling de-etiolation in continuous far-red light. *Plant
Journal* **60**, 449-461 (2009). https://doi.org:10.1111/j.1365-313X.2009.03971.x.
24 Kevei, E., Schafer, E. & Nagy, F. Light-regulated nucleo-cytoplasmic partitioning of
phytochromes. *J Exp Bot* **58**, 3113-3124 (2007). https://doi.org:10.1093/jxb/erm145.
25 Toledo-Ortiz, G. *et al.* Subcellular sites of the signal transduction and degradation of
phytochrome A. *Plant Cell Physiol* **51**, 1648-1660 (2010).
https://doi.org:10.1093/pcp/pcq121.
26 Chen, F. *et al.* Phosphorylation of FAR-RED ELONGATED HYPOCOTYL1 is a key
mechanism defining signaling dynamics of phytochrome A under red and far-red light in
Arabidopsis. *Plant Cell* **24**, 1907-1920 (2012). https://doi.org:10.1105/tpc.112.097733.
27 Wang, H. & Deng, X. W. Arabidopsis FHY3 defines a key phytochrome A signaling
component directly interacting with its homologous partner FAR1. *EMBO J* **21**, 1339-
1349 (2002). https://doi.org:10.1093/emboj/21.6.1339.
28 Qu, G. P. *et al.* Reversible SUMOylation of FHY1 Regulates Phytochrome A Signaling in
Arabidopsis. *Mol Plant* **13**, 879-893 (2020). https://doi.org:10.1016/j.molp.2020.04.002.
29 Toledo-Ortiz, G. *et al.* Subcellular sites of the signal transduction and degradation of
phytochrome A. *Plant Cell Physiol* **51**, 1648-1660 (2010).
https://doi.org:10.1093/pcp/pcq121.
30 Menon, C., Klose, C. & Hiltbrunner, A. Arabidopsis FHY1 and FHY1-LIKE Are Not
Required for Phytochrome A Signal Transduction in the Nucleus. *Plant Commun* **1**,
100007 (2020). https://doi.org:10.1016/j.xplc.2019.100007.
31 Rausenberger, J. *et al.* Photoconversion and Nuclear Trafficking Cycles Determine
Phytochrome A's Response Profile to Far-Red Light. *Cell* **146**, 813-825 (2011).
https://doi.org:10.1016/j.cell.2011.07.023.
32 Al-Sady, B., Ni, W., Kircher, S., Schafer, E. & Quail, P. H. Photoactivated phytochrome
induces rapid PIF3 phosphorylation prior to proteasome-mediated degradation. *Mol Cell*
23, 439-446 (2006). https://doi.org:10.1016/j.molcel.2006.06.011.
33 Shen, H. *et al.* Light-induced phosphorylation and degradation of the negative regulator
PHYTOCHROME-INTERACTING FACTOR1 from depend upon its direct physical
interactions with photoactivated phytochromes. *Plant Cell* **20**, 1586-1602 (2008).
https://doi.org:10.1105/tpc.108.060020.
34 Kim, C. *et al.* Phytochrome B photobodies are comprised of phytochrome B and its
primary and secondary interacting proteins. *Nature Communications* **14** (2023).
https://doi.org:10.1038/s41467-023-37421-z.

35   Zhang, X. Y. *et al.* Liquid-liquid phase separation in diseases. *Medcomm* **5** (2024).
https://doi.org:e64010.1002/mco2.640.
36   Chen, D. *et al.* Integration of light and temperature sensing by liquid-liquid phase
separation of phytochrome B. *Mol Cell* **82**, 3015-+ (2022).
https://doi.org:10.1016/j.molcel.2022.05.026.
37   Genoud, T. et al. FHY1 Mediates Nuclear Import of the Light-Activated Phytochrome A
Photoreceptor. Plos Genet 4 (2008).
https://doi.org:100014310.1371/journal.pgen.1000143.
38   Hussain, E., Romanowski, A. & Halliday, K. J. PIF7 controls leaf cell proliferation through
an AN3 substitution repression mechanism. *P Natl Acad Sci USA* **119** (2022).
https://doi.org:10.1073/pnas.2115682119.
39   Leivar, P. *et al.* Multiple Phytochrome-Interacting bHLH Transcription Factors Repress
Premature Seedling Photomorphogenesis in Darkness. *Curr Biol* **18**, 1815-1823 (2008).
https://doi.org:10.1016/j.cub.2008.10.058.
40   Li, L. *et al.* Linking photoreceptor excitation to changes in plant architecture. *Genes Dev*
26, 785-790 (2012). https://doi.org:10.1101/gad.187849.112.
41   Viczián, A. *et al.* Functional characterization of phytochrome interacting factor 3 for the
circadian clockwork. *Plant and Cell Physiology* **46**, 1591-1602 (2005).
https://doi.org:10.1093/pcp/pci175.
42   Andreou, A. I. & Nakayama, N. Mobius Assembly: A versatile Golden-Gate framework
towards universal DNA assembly. Plos One 13 (2018). https://doi.org:
10.1371/journal.pone.0189892.
43   Urquiza-García, U. & Millar, A. J. Expanding the bioluminescent reporter toolkit for plant
science with NanoLUC. Plant Methods 15 (2019). https://doi.org:10.1186/s13007-019-
0454-4.
44   Zielinski, T., Moore, A. M., Troup, E., Halliday, K. J. & Millar, A. J. Strengths and
Limitations of Period Estimation Methods for Circadian Data. *Plos One* **9** (2014).
https://doi.org:10.1371/journal.pone.0096462
45   Wang, Z. S. *et al.* Proteasome-associated ubiquitin ligase relays target plant hormone-
specific transcriptional activators. *Sci Adv* **8** (2022).
https://doi.org:10.1126/sciadv.abn4466.
46   Zhu, D. M. *et al.* Biochemical Characterization of Complexes Containing
CONSTITUTIVELY PHOTOMORPHOGENIC1 and SUPPRESSOR OF PHYA Proteins
in Light Control of Plant Development. *Plant Cell* **20**, 2307-2323 (2008).
https://doi.org:10.1105/tpc.107.056580.
47   Rosler, J., Klein, I. & Zeidler, M. double mutant reveals a distinct cytoplasmic action of
phytochrome A. *P Natl Acad Sci USA* **104**, 10737-10742 (2007).
https://doi.org:10.1073/pnas.0703855104.

Fig. 1 PIF7ox influences phyA Function in WLFR

a, Schematic of experimental growth conditions. Seeds were sown on 1/2 MS plates and cultivated for one day under white light, followed by 4 hours (h) of light and 20 h of dark. Seedlings were subsequently grown for 6 days in white light (WL) photoperiods with or without supplementary far-red (WLFR), lowering the R:FR to 0.20 (red arrow). For all experiments, light regimes were 8 h light at PAR = 85 ± 5 μmol m^{-2} s^{-1}, 16 h dark, at 22°C.

b, Representative images of *Arabidopsis thaliana* WT (*Col-0*), *pif7-1*, *pif7-2*, and PIF7ox seedlings grown in WL or WLFR for 6 days. Scale bar = 1 cm.

c, PIF7ox blocks phyA action in WLFR. Hypocotyl length (mm) of WT (*Col-0*), *pif7-1*, *pif7-2*, PIF7ox, *phyA-211, pif7-1;phyA-211, PIF7ox;phyA-211* and phyAox under WL and WLFR. Representative images from three independent replicates are shown.

d, In WLFR, PIF7 enhances *PHYA* expression at Zeitgeber time 7 (ZT). Real-time qPCR was used to quantify *PHYA* mRNA levels, with *PP2A* as internal control. Samples were collected at ZT7 (1 hour before dusk) from 6d-old seedlings grown in WL or WLFR.

e,f, In WLFR, PIF7 increases phyA protein accumulation. PhyA protein levels were determined by immunoblotting. Anti-phyA was used to detect native phyA, while anti-RPN6 served as an internal control. The bar chart (**f**) displays the relative levels of phyA, normalised to RPN6, based on data from four biological replicates collected at ZT7.

g, PIF7ox boosts phyA protein levels in diel conditions. The phyA-nLUC line was generated by expressing (*PHYApr:phyA-nLUC*) in *phyA-211*, while the PIF7ox;phyA-nLUC was generated by introgressing *PHYApr:phyA-nLUC* into the PIF7ox line. Bioluminescence signals were recorded from day 6.

For **c,d,f,g**, data are presented as mean values ± s.e.m., *n*=25 (seedling numbers) in **c**; with *n*=3,4,9 (biological repeats) in **d,f,g**, respectively. For **c,d,f**, a Two-way ANOVA was conducted to assess statistical significance across light conditions for each genotype (α=0.05); asterisks denote significance relative to WL treatments: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), *P* < 0.0001 (****).

Fig. 2 PIF7 and phyA interact in sub-nuclear photobodies under WLFR.

a, 35S:PIF7-sfGFP and 35S:phyA-RFP co-localise in nuclei and photobodies in WLFR. Co-localisation is evident from the chart (right) which traces the GFP and RFP channel signals along the white bar shown in merged nucleus image (left).

b, Split NanoLUC (nLUC) assay demonstrates PIF7 and phyA physically interact. 35S:PIF7-lgbitnLUC (PIF7-lgbit), 35S:phyA-smbitnLUC (phyA-smbit), 35S:phyB-smbitnLUC (phyB-smbit) and 35S:phyA-nLUC (phyA-nLUC) reporter constructs were expressed in *Nicotiana benthamiana*. A WLFR-dependent interaction was demonstrated for PIF7-lgbit+phyA-smbit. PhyA-nLUC was used as light conditional controls, and phyB-smbit+phyA-smbit as the negative control.

c, Bimolecular Fluorescence Complementation (BiFC) assay confirms PIF7 and phyA interact in nuclear photobodies. BiFC was detected between 35S:PIF7-nYFP and 35S:phyA-cYFP (PIF7-nYFP + phyA-cYFP) in WLFR, but not with the 35S:EV-cYFP (EV-cYFP) control.

d, Co-immunoprecipitation assay (Co-IP) shows that phyA interacts with PIF7 in WLFR *in vivo*. Anti-His beads were used to pull down PIF7 from PIF7ox (35S:*PIF7-FLASH*; 9xMyc-6xHis-3xFlag) or the WT(*Col-0*) negative control. Anti-phyA and anti-FLAG used for detecting phyA and PIF7 respectively, in IP/Input. Samples were collected at ZT7 from 6-d old seedlings exposed to WL or WLFR.

For **a**,**c**, 488nm and 552nm laser detected the GFP/YFP and RFP signals. Scale Bar=15μm. For **b**, data are presented as mean values ± s.e.m., *n*=12 (leaf numbers). A Two-way ANOVA test was used to obtain statistical significance (α=0.05); asterisks denote statistical significance comparing negative to positive test or tested interaction treatments: 0.1234 (ns), 0.0332(*), 0.0021 (**), 0.0002(***), P<0.0001 (****). Representative images from three independent replicates are shown in all figs.

Fig. 3 The PIF7-phyA Interaction Depends on the PIF7 Prion like domain

a-b, BIFC assays using truncated constructs identify the phyA PAS domain and the PIF7 prion-like domain, as important for the phyA-PIF7 interaction. The constructs image depicted the complete phyA/PIF7 protein structures. Full length and truncated contructs were fused to C-terminal nYFP (PIF7) or cYFP (phyA) BIFC constructs. PIF7-nYFP+phyA-cYFP was used as positive control and 35S:EV-cYFP/35S:EV-nYFP (EV-cYFP/EV-nYFP) as used as negative controls. 488nm lasers were used to detect the YFP signals. Scale Bar=15μm. Representative images from three independent replicates are shown.

Fig. 4 PIF7 Influences the Sub-cellular Location of phyA

a-c, In WLFR, PIF7ox promotes accumulation of phyA-GFP in the cytosol. Lines of *pif7-1*;phyA-GFP and PIF7ox;phyA-GFP were generated by introgressing *PHYApr*:phyA-sGFP into *pif7-1* and PIF7ox. Close-up images under WLFR are shown in **b**, white arrows and N label indicate nuclear, while red arrows and C label indicate cytosolic signals. In **c**, nuclear/cytosolic (N/C) ratios (relative to WT (*Col-0*), WL control) were calculated following quantification using ImageJ.

d-e, PIF7ox enhances phyA protein accumulation in the cytosol under WLFR conditions. Immunoblot assays were conducted using an anti-phyA antibody, using anti-RbCL as the cytosolic control, and anti-Histone 3 (H3) as the nuclear control. In **e**, cytoplasmic and nuclear phyA was normalised to RbCL and H3, respectively. The N/C ratio for each genotype is shown relative to WT (*Col-0*), WL control. Six-day-old seedlings were sampled as ZT7.

For **a-b** a wavelength of 488nm was used to detect GFP signals. Scale bar=25μm. Representative images from three independent replicates are shown. For **c,e**, data are presented as mean values ± s.e.m., *n*=6 (seedling numbers) in **c**; *n*=4 (biological repeats) in **e**. A two-way ANOVA test assessed statistical significance between genotypes under WL/WLFR condition (α=0.05); asterisks denote significance compared to the phyA-GFP (phyA N/C) values for WL/WLFR condition(**c**) or phyA protein (phyA N/C) values in WT with WL/WLFR condition(**e**): 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), $P < 0.0001$ (****).

Fig. 5. PIF7 Inhibits phyA-Mediated Regulation of Target Genes

a, qPCR analysis of *CHS, FHY1, FHL, FHY3, BIN2,* and *IAA29* expression in WT (*Col-0*), *pif7-1*, PIF7ox and *phyA-211*. Transcript levels were normalised to the *PP2A* internal control.

b, PIF7ox reduces phyA enrichment at *CHS, FHY1,* and *FHL* promoters under WLFR. Chromatin immunoprecipitation (ChIP)-qPCR was conducted with 6-day-old WT (*Col-0*) or PIF7ox seedlings sampled at ZT7. Native phyA was immobilised on protein A/G beads using an anti-phyA antibody, and promoter enrichment quantified by qPCR at Gbox promoter regions and coding regions, which served as negative controls. Data shown is normalised to input samples.

For **a-b**, data are presented as mean values ± s.e.m., *n*=3 (biological repeats). A two-way ANOVA test was used to assess statistical significance between the WL/WLFR for each genotype (**a**) or different genotypes under the same conditions (**b**) (α=0.05); asterisks indicate statistical significance compared to control: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), $P < 0.0001$ (****).

Fig. 6. PIF7 interacts with FHY1/FHL in WLFR

a, The BiFC assay demonstrates PIF7 interaction with FHY1 and FHL. Interaction fluorescence was detected when 35S:FHY1-cYFP (FHY1-cYFP) or 35S:FHL-cYFP (FHL-cYFP) were with expressed 35S:PIF7-nYFP (PIF7-nYFP) in *N. benthamiana* under WLFR.

b, Interactions between PIF7 and FHY1/FHL are detected using a split NanoLUC assay. Strong fluorescence signals were detected between 35S:FHY1-smbitnLUC (FHY1-smbit), 35S:FHL-lgbitnLUC (FHL-lgbit) and 35S:PIF7-lgbit(smbit)nLUC (PIF7-lgbit/smbit) in the transient *N. benthamiana* expression system. The positive controls were phyA-nLUC and PIF7-lgbit+phyA-smbit, while PIF7-smbit+phy-Asmbit was used as a negative control.

c, An immunoprecipitation assay demonstrates PIF7 interacts with FHY1/FHL *in vivo*. 35S:FHY1/FHL-nLUC:3xFLAG-10xHis and 35S:PIF7-sfGFP constructs were transiently expressed in *N. benthamiana* leaves. Anti FLAG beads were used to pull down FHY1/FHL and its target proteins by Co-IP. Anti GFP antibodies revealed GFP-tagged PIF7 co-IP with FHY1/FHL in WLFR.

d-e, Fluorescence assays illustrate that PIF7, phyA and FHY1/FHL co-localise in nuclei and photobodies. 35S driven PIF7-sfGFP, FHY1/FHL-EYFP, and phyA-RFP constructs were transiently expressed in *N. benthamiana* leaves. The merged image (**d**) signifies co-localisation, and **e**, displays GFP, YFP and RFP channel signals relative to white bar shown in the merged nuclear image.

f, PIF7 reduces *ASP1* mRNA levels under WLFR. qPCR analysis of *ASP1* expression in WT, *pif7-1*, PIF7ox, *PP2A* was used as internal control.

g, phyA-NLS is epistatic to PIF7ox. Hypocotyl length (mm) is shown for WT (Col-0), *phyA-211, phyAox;phyA-211,* WT (Ler), *phyA-201, phyA-NLS;phyA-201, PIF7ox;phyA-NLS;phyA-201.* Double and triple mutant lines were generated through genetic.

h, Schematic depicting the cellular model for PIF7 control of phyA action in canopy shade. In WLFR PIF7 accumulates in the nucleus, where it interacts with phyA and FHL/FHY1 in photobodies. This sequestration not only blocks phyA from regulating downstream genes but also inhibits its transport into the nucleus. Consequently, PIF7 is able to amplify SAS-mediated hypocotyl elongation by blocking the antagonistic action of phyA. The image created in BioRender (https://BioRender.com).

For **a,d,** 488nm and 552nm lasers were used for detecting GFP/YFP and RFP signals. Scale bar=15μm. For **b,f,g**, data are presented as mean values ± s.e.m., *n*=12 (leaf numbers) in **b**; *n*=3 (biological repeats) in **f**; *n*=24 (seedling numbers) in **g**. A two-way ANOVA test was conducted to determine statistical significance (α=0.05), asterisks denote significance when comparing negative to positive controls and tested interactions (**b**) or compared to WL treatments in each genotype (**f-g**): 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), *P* < 0.0001 (****).