Opposing CTCF and GATA4 activities set the pace of chromatin topology remodeling during cardiomyogenesis

Silvia Becca[1], Sara Bianchi[1], Elisa M. Hahn[1#], Kirsten E. Snijders[1], Lukasz Truszkowski[1],
Anna Krepelova[2], Francesco Neri[2], Davide Cacchiarelli[3], Sasha Mendjan[4], Elisa Balmas[1],
and Alessandro Bertero[1,5*]

[1]Molecular Biotechnology Center "Guido Tarone", Department of Molecular Biotechnology and Health Sciences, University of Torino, Italy
[2]Molecular Biotechnology Center "Guido Tarone", Department of Life Sciences and Systems Biology, University of Torino, Italy
[3]Telethon Institute of Genetics and Medicine, Pozzuoli, Italy
[4]Institute of Molecular Biotechnology of the Austrian Academy of Sciences (IMBA), Vienna BioCenter, Austria
[5]Lead contact
[#]Current affiliation: European Molecular Biology Laboratory, EMBL Barcelona, Spain
[*]Correspondence: alessandro.bertero@unito.it

SUMMARY

Reorganization of the three-dimensional chromatin structure is a critical feature of human embryonic development. Yet, the mechanisms regulating integrative remodelling of local structures (e.g., loops) and global architecture (e.g., A/B compartmentalization) remain unclear. Here, we investigate this aspect in the context of cardiomyogenesis, characterized by pronounced B-to-A remodelling of several cardiac-specific genes such as *TTN*. We focus on the roles of the pioneer transcription factor GATA4 and the architectural protein CTCF. Using an inducible knockdown system during human induced pluripotent stem cell differentiation, we show that GATA4 is essential for timely topological activation of key cardiac genes, while partial depletion of CTCF, anticipating physiological downregulation during development, enhances this process. Deletion of a single CTCF binding site on *TTN* leads to modest gene decompaction and transcriptional activation. Bulk and single-cell RNA sequencing of chamber-specific cardiac organoids reveals that loss of GATA4 delays differentiation and sustains proliferation of early cardiomyocytes, whereas premature CTCF depletion accelerates yet confounds cardiomyocyte maturation. These findings suggest that CTCF and GATA4 have antagonistic roles on chromatin dynamics during cardiomyogenesis, forming a rheostat that maintains accurate developmental tempo. Disruption of this mechanism may contribute to congenital heart defects caused by mutations in these factors.

INTRODUCTION

Chromatin architecture dynamically shapes transcriptional programs during development and in human disease[1], yet the underlying mechanisms remain incompletely understood and are an active area of investigation[2]. During differentiation, pluripotent stem cells (PSCs) undergo extensive chromatin reorganization and transcriptional remodeling to establish lineage-specific gene expression programs. This involves coordinated changes in chromatin accessibility[3], histone modifications[4], and three-dimensional (3D) genome organization, a higher-order layer of epigenetic regulation that has recently emerged as a key orchestrator of developmental gene expression[5]. This multi-layered regulation spans from short- and long-range chromatin loops, to topologically associating domains (TADs) hundreds of kilobases long, and up to the segregation of megabases-sized chromosomal regions into so-called A (active) and B (inactive) compartments[6]. Together, these structures mediate inter- and intra-chromosomal interactions that shape chromosome territorialization and radial nuclear positioning.

Cardiac development provides a striking example of developmental 3D chromatin remodeling, as changes in loops, TADs, A/B compartments, and inter-chromosomal networks facilitate the temporal and spatial control of cardiac gene expression (reviewed in[7]). Disruption of these processes has been linked to genetic dilated cardiomyopathy (DCM) caused by lamin A/C mutations[8,9], and mutations in chromatin regulators are strongly enriched in *de novo* congenital heart defects (CHD)[10], underscoring their pathogenic relevance[11].

Unbiased genome-wide analyses have revealed a remarkable pattern in which large cardiac genes transition from the lamina-associated B compartment to the A compartment within the nuclear interior, where they engage in higher-order interactions and enhanced transcription[12–14]. These topological changes establish stable gene regulatory networks essential for cardiac lineage commitment. However, the molecular mechanisms driving chromatin reorganization in early cardiomyogenesis remain poorly understood, and it is unclear how their dysregulation disrupts cardiac development and function, ultimately leading to CHD and DCM.

Pioneer factors such as GATA4 can access and open closed chromatin regions, enabling recruitment of transcriptional machinery to activate lineage-specific programs[15]. GATA4 is particularly crucial in cardiac development, where it regulates genes controlling mesoderm specification, first and second heart field formation, and cardiomyocyte maturation[16,17]. Mutations in GATA4 have been identified in Mendelian forms of isolated CHD, underscoring its central role in early cardiac morphogenesis[18,19].

Architectural proteins such as CTCF (CCCTC-binding factor), a ubiquitously expressed insulator, play a key role in defining TADs and mediating enhancer-promoter looping[20]. By partnering with cohesin, CTCF stabilizes chromatin loops that shape gene regulation[21]. Traditionally viewed as a boundary factor preventing aberrant interactions[22], CTCF is now recognized as a dynamic regulator of developmental transitions and lineage specification[23,24]. *De novo* CTCF mutations cause a syndromic developmental disorder that includes CHD, implicating its dysfunction in cardiac malformations[25,26].

The mechanism of action of GATA4 and CTCF and their combined effects on chromatin reorganization in cardiomyogenesis remain largely unexplored. We addressed this question using cardiomyocytes derived from human PSCs and PSC-derived cardiac organoids[27], which recapitulate chamber-specific identities. Our study reveals opposing roles for GATA4 and CTCF, respectively promoting and restraining early cardiac differentiation.

RESULTS

The *TTN* locus models B-to-A compartment switching during cardiomyogenesis

We and others previously reported that *TTN*, encoding the large sarcomere protein titin, transitions from the B to A compartment during PSC differentiation into cardiomyocytes (CMs), accompanied by a marked repositioning from the nuclear periphery toward the interior[12,13]. This was evident from shallow-sequenced DNAse Hi-C data from both female RUES2 human embryonic stem cell-derived CMs (hESC-CMs) and male WTC11 human induced pluripotent stem cell-derived CMs (hiPSC-CMs) differentiated for 2 weeks using a biphasic WNT modulation protocol, though with different kinetics (Figure S1A). We observed the same phenomenon by re-analysing deeply sequenced *in situ* Hi-C during a CM differentiation time course of female H9 hESCs up to day 80 (Figure 1A). In all datasets, the compartment switch coincided with progressive reduction of promoter-gene body interactions, indicative of gene unpacking. Thus, *TTN* offers a model to study topological rearrangement that is reproducible across different types of hPSCs, cell lines, and laboratories.

Figure 1. Reciprocal dynamics of the *TTN* locus regulators GATA4 and CTCF

(A) Integrative view of local and global chromatin topology remodelling at *TTN* during H9 hESC-CM differentiation, as assessed by Hi-C (data plotted from quantile normalized average of n = 2 differentiations)[28], ATAC-seq (data plotted from average of n = 2 differentiations)[13], and CTCF[29] and GATA4 ChIP-seq[17,30]. Virtual 4C extracted from Hi-C data showing *TTN* DNA interaction with its promoter proximal region. D: day; PS: primitive streak; MES: mesoderm; CP: cardiac progenitors; CM: cardiomyocyte; eCM: early CM; vCM: ventricular CM.

(B) Expression of *GATA4* and *CTCF* from single-cell RNA-seq of human early embryo development[31–33] (normalized read counts). Cells were developmentally ranked based on pseudotime analysis (Figure S1B). n = 88 pre-implantation embryos, n = 1 gastrulating embryo, and n = 18 fetal heart samples

(C) Bulk RNA-seq time course of H9 hESC-CM differentiation[34]. Average z-scores of n = 3 – 4 differentiations.

(D) Representative western blot during WTC11 hiPSC-CM differentiation. CTCF expression was normalized to vinculin and to the day 0 sample across n = 3 differentiations; p-value by one-sample t test against μ = 1. POU5F1 (OCT4) and ITGB1BP2 (Melusin) are included as established hiPSC and CM markers, respectively[35].

TTN experienced clear changes in chromatin accessibility during CM differentiation: ATAC-seq identified two CM-specific peaks that correlated with ChIP-seq signal for the cardiac pioneer TF GATA4 at the two promoters, one driving full length titin and the other the shorter cronos isoform. In contrast, six regions became less accessible in CMs, correlating with ChIP-seq signal for the ubiquitous architectural protein CTCF (Figure 1A). Notably, four out of five putative CTCF binding sites within the *TTN* gene body displayed a convergent orientation with a promoter-proximal CTCF peak. According to the chromatin loop extrusion model, such an arrangement is predicted to favor CTCF-mediated cohesin arrest, stabilizing promoter-gene body interactions that may interfere with transcription initiation and/or elongation[21,36].

Intrigued by these findings, we integrated publicly available single cell RNA sequencing (scRNA-seq) datasets of early human embryo development to determine the expression patterns of *GATA4* and *CTCF* during cardiomyogenesis (Figure 1B, Figure S1B). In line with mouse development[37], *GATA4* began to be expressed at the nascent mesoderm stage (Figure 1B); this corresponded to ~5-6 days of directed differentiation towards left ventricle (LV) in monolayer cultures of all three aforementioned hPSC lines (Figure 1C, Figure S1C), and to ~3.5-4.5 days of WTC11 hiPSC specification into self-organizing LV-, right ventricle-, and atrial-like three-dimensional cardiac organoids (cardioids[27,38]; Figure S1D). Unexpectedly, the increase in *GATA4* expression was mirrored by a decrease in *CTCF* expression, which was approximately halved after mesoderm specification in all models examined (Figures 1B–1C, Figures S1C–S1D). Accordingly, we observed significantly lower CTCF protein levels in WTC11 hiPSC-CMs, the system we selected for further mechanistic studies (Figure 1D).

The decrease in CTCF protein levels may explain its selective detachment from *TTN* during cardiomyogenesis. Supporting this, the core CTCF binding motifs at *TTN* had lower predicted scores than a constitutively-bound positive control, with the weakest motifs corresponding to the most differentially-bound sites at the 5' and 3' end of the gene (Figure S1E). In addition, DNA methylation at the 5' site was low and remained unchanged throughout differentiation, arguing against methylation as a major factor limiting CTCF binding (Figure S1F). Thus, opposite changes in GATA4 and CTCF expression represent a conserved feature of both *in vivo* and *in vitro* cardiac development and correlate with dynamic occupancy at *TTN*.

GATA4 and CTCF oppositely regulate cardiac B-to-A compartment switching

The correlation between changes in DNA binding and the expression dynamics of GATA4 and CTCF prompted us to test their functional roles in genome topology during cardiomyogenesis. To this end, we generated polyclonal titin-mEGFP reporter hiPSC lines carrying tetracycline (TET)-inducible short hairpin RNAs (shRNAs) for conditional knockdown (KD) of either GATA4 or CTCF, or a scrambled (SCR) shRNA control[39]. We validated at least one shRNA per gene that achieved >70% KD in hiPSC-derived cardiac progenitors (CPs; Figure S2A). Strong depletion of GATA4 throughout hiPSC-CM differentiation almost completely blocked cardiac induction, as assessed by loss of cardiac troponin T (cTnT; Figure S2C). We therefore selected a less effective shRNA to model a more moderate impairment of GATA4, which may better reflect CHD patients carrying heterozygous mutations (Figures 2A–2B). This resulted in a ~50% decrease in cTnT expression and a ~8-fold reduction in titin-mEGFP levels (Figure 2C). In contrast, CTCF KD during differentiation significantly increased titin-mEGFP expression by more than twofold, while cTnT levels remained comparable to controls (Figures 2A–2C).

Figure 2. Inducible knockdown of GATA4 or CTCF during hiPSC-CM differentiation

(A) RT-qPCR validation of gene knockdown in day 25 hiPSC-CMs. N = 3 differentiations. p-values by two-way ANOVA with Holm–Sidak corrected pairwise comparisons. CTR: no tetracycline control; TET: tetracycline.

(B) Western blot validation of gene knockdown at protein level. VNC: vinculin loading control.

(C) Representative flow cytometry quantification of cardiac differentiation efficiency, and quantifications from n =3 differentiations. Statistical analyses as in panel A.

(D) ImmunoFISH to quantify the distance of the *TTN* locus from the nuclear lamina (LMNB1). Violin plots report nuclear-normalized distances from the indicated number of cells. p-values one-way ANOVA with with Sidak corrected pairwise comparisons. On the right panel representative images for GATA4 CTR and TET.

(E) RT-qPCR analyses for a subset of cardiac B to A genes. Average fold-change TET/CTR from n = 3 differentiations.

(F) Gene set enrichment analyses (GSEA) of an extended cardiac B to A gene list in bulk RNA-seq from n = 2 differentiations. p-values by Adaptive Monte-Carlo Permutation test.

To connect these expression changes with nuclear topology, we examined the proximity of the *TTN* locus to the nuclear lamina. Following GATA4 KD, *TTN* was more closely associated with lamin B1, consistent with impaired B-to-A compartment transition (Figure 2D). By contrast, CTCF KD did not produce a detectable change in *TTN* positioning (Figure 2D, Figure S2D). Notably, nuclear repositioning of *TTN* was less pronounced in WTC11 hiPSC-CMs than in RUES2 hESC-CMs[13], consistent with a slower B-to-A transition kinetics for this line (Figure S1A).

To assess whether GATA4 and CTCF have broader roles at other loci undergoing B-to-A transition during cardiomyogenesis, we performed reverse transcription quantitative PCR (RT-qPCR) for 13 such genes we had previously identified during RUES2 hESC-CM differentiation[13]. Most of these genes were downregulated after GATA4 KD, whereas several were upregulated after CTCF KD (Figure 2E). We next broadened the analysis with bulk RNA sequencing (RNA-seq), which was consistent among biological replicates (Figures S2E–S2F). We examined a larger set of B-to-A genes derived from Hi-C developmental kinetics in H9 hESC-CMs (Figure S2G). Gene set enrichment analyses (GSEA) revealed significant downregulation of B-to-A genes following GATA4 KD and upregulation after CTCF KD (Figure 2F). Consistently, expression of genes in the KEGG cardiac muscle contraction pathway was reduced after GATA4 KD but increased after CTCF KD (Figure S2H). Altogether, these results suggest that GATA4 and CTCF function as positive and negative regulators of B-to-A compartment switching, likely acting through global and local chromatin remodeling, respectively.

CTCF promotes repressive intergenic looping at *TTN*

GATA4 is an established master cardiac TF[17]. By contrast, CTCF has not previously been implicated as a negative modulator of cardiomyogenesis. We therefore focused our attention on this factor, aiming to gain mechanistic insights. Specifically, we perturbed CTCF binding *via* deletion of its cognate DNA sites and examined the effects on chromatin architecture and gene expression (Figure 3A). We first selected the CTCF binding site (CBS) at the gene 5' end (CBS1) because it is convergent with four other sites within the gene body. As discussed above, the loop extrusion model predicts that CBS1 may stabilize repressive CTCF-mediated loops between the promoter-proximal region and the gene body. We also selected the last site at the gene 3' end (CBS6), considering that it could contribute to a topologically associating domaing (TAD) boundary isolating *TTN* from the downstream A compartment in hPSCs.

We targeted the core CTCF binding motifs by CRISPR/Cas9-mediated homology-directed repair (HDR) using single-stranded DNA (ssDNA) donor oligos, combined with a co-targeting with selection strategy to increase positive clone recovery[40] (Figure 3A). We isolated two edited lines: one with homozygous disruption of CBS1 and compound heterozygous editing at CBS6 (Δ2CBS), and another with homozygous editing only at CBS1 (Δ1CBS; Figures S3A–S3B). In the Δ2CBS line, CBS6 carried the intended edit on one allele and a 2-bp deletion on the other (Figure S3C). CTCF ChIP-qPCR in hiPSCs confirmed complete loss of CTCF binding at the targeted sites, while occupancy at other CBSs remained unaffected (Figure 3B).

We performed circular chromosome conformation capture sequencing (4C-seq) using the promoter proximal region of *TTN* as the viewpoint. The interaction profiles across the *TTN* gene body were very similar for Δ2CBS and Δ1CBS hiPSCs, and each differed clearly from wild-type (WT) controls (Figure 3C). To place these changes in developmental context, we compared our 4C data with virtual 4C generated from published Hi-C of RUES2 hESCs across

Figure 3. Precise disruption of CTCF binding sites at *TTN*

(A) Schematic of the CRISPR/Cas9 strategy to edit selected CTCF binding sites (CBS) on the *TTN* locus in hiPSCs. The core CTCF binding motif was replaced with an unrelated EcoRI sequence.

(B) CTCF ChIP-qPCR in hiPSCs validating genome editing. Enrichment of all *TTN* CBSs was normalized to four negative control regions and compared to a positive control site outside *TTN*. Δ1CBS: homozygous deletion of CBS1; Δ2CBS: homozygous deletion of CBS1 plus compound heterozygous disruption of CBS6; WT: wild-type. n = 3 cultures, p-values by two-way ANOVA with Holm–Sidak corrected multiple comparisons; only significant comparisons are annotated.

(C) 4C-seq in hiPSCs using the promoter-proximal region of *TTN* as viewpoint. Data are also shown as the differences between pairs of conditions. Right inset: zoom-in of the CBS4-5 region. Mean of n = 3 cultures.

(D) Differential 4C-seq AUC between pairs of conditions for the indicated genomic intervals. p-values by one-sample t-test against μ = 0.

(E) RT-qPCR of *TTN* mRNA expression (left) and flow cytometry quantification of titin-positive cells and titin median fluorescence intensity (MFI; middle and right) in day 13 hiPSC-CMs. n = 6 differentiations, p-values by two-way RM ANOVA with Holm–Sidak corrected multiple comparisons.

CM differentiation. Interaction patterns in WT hiPSCs correlated most closely with the pluripotent stage, as expected, whereas Δ2CBS and Δ1CBS clones aligned more closely with cardiac progenitors (Figure S3D). Closer inspection of specific regions revealed two opposing effects in both Δ2CBS and Δ1CBS: promoter interactions increased up to CBS2, but decreased beyond CBS4, particularly between CBS4 and CBS5 (Figures 3C–3D, Figures S3E–S3F). These observations suggest that CBS1 deletion is sufficient to mimic some developmental chromatin topology changes by disrupting long-range intergenic looping between the *TTN* promoter and gene body, favoring shorter-range interactions typical of a less constrained chromatin polymer.

We next asked whether these architectural changes influenced *TTN* expression. RT-qPCR revealed a significant increase in *TTN* mRNA by 50-75% in both Δ2CBS and Δ1CBS hiPSC-CMs, despite comparable efficiency of CM specification (Figure 3E). Titin protein levels, however, remained unchanged, likely due to compensatory mechanisms buffering the modest gene expression increase (Figure 3E). The absence of further changes in Δ2CBS compared to Δ1CBS is consistent with the 4C data and indicates that CBS6 is redundant in this context.

Overall, deletion of a single CTCF binding motif produced only moderate effects on *TTN* chromatin organization and transcription, consistent with the redundancy of CTCF sites at large loci[41,42]. Nevertheless, these findings support a model in which CTCF-mediated intergenic looping at *TTN* contributes to transcriptional restraint, providing a mechanistic link to the broader derepression of B-to-A genes observed upon partial depletion of CTCF.

GATA4 and CTCF exert antagonistic control over second heart field development

To explore the physiological relevance of GATA4- and CTCF-dependent chromatin topology, we turned to a more complex three dimensional (3D) model: self-assembling, multi-lineage cardioids[27,38]. Using scRNA-seq, we validated protocols to generate WTC11 hiPSC-derived 3D cardioids representing the three primary heart chambers: the first heart field (FHF)-derived left ventricle (LV), and the second heart field (SHF)-derived right ventricle (RV) and atria (A; (Figure S4A). Each cardioid type contained the major cardiac cell populations, including cardiomyocytes (*TNNT2*+ and *TTN*+), fibroblasts (*COL3A1*+), and endothelial cells (*ERG*+), while maintaining chamber-specific identities (e.g., *IRX4*+ LV CMs, *IRX1*+ RV CMs, and *N2RF2*+ A CMs; Figures S4A–S4B). In addition, we identified proliferating cells (*MKI67*+), mesodermal-stage cells (*MESP1*+), and small fractions of undifferentiated (*POU5F1*+) and epithelial-like (*KRT19*+) cells (Figure S4A).

We investigated the effects of GATA4 and CTCF silencing in this model by differentiating inducible KD hiPSCs into FHF- and SHF-derived cardioids for 7.5 days. RT-qPCR indicated that KD did not alter chamber identity (Figure S4C). GATA4 KD cardioids showed upregulation of both FHF and SHF cardiac progenitor markers (*HAND1*, *HAND2*, *NKX2-5*), together with increased expression of the fibroblast marker *COL1A1* (Figure S4C). In contrast, CTCF KD caused downregulation of *COL1A1*(Figure S4C). Consistent with findings in monolayer hiPSC-CMs, *TTN* expression was decreased by GATA4 KD in all cardioid types and increased by CTCF KD in RV and A cardioids, though intriguingly not in LV ones (Figure 4A). The fraction of titin-positive cells largely mirrored the mRNA changes, though differences were more modest (Figure 4B), possibly because titin expression was more accelerated and homogeneous in cardioids compared with monolayer hiPSC-CMs (Figure 2C).

Figure 4. Inducible knockdown during FHF and SHF cardioid differentiation
Legend on the next page

Because the effects of GATA4 and CTCF silencing were more evident in SHF derivatives,
we performed bulk RNA-seq on day 7.5 RV and A cardioids in biological duplicate, which
showed good reproducibility and clustering by KD type (Figure S4D). Expression analysis of
B-to-A genes revealed a general downregulation after GATA4 KD and an upregulation after
CTCF KD in RV organoids, while effects in atrial organoids were modest (Figure 4C). We next
compared global transcriptional changes between conditions using rank-rank hypergeometric
overlap (RRHO) analysis[43], which demonstrated a strong anti-correlation between GATA4 and
CTCF KD profiles, particularly in RV organoids (Figure 4D). Finally, GSEA of heart process-
related genes showed significant downregulation after GATA4 KD in both RV and A cardioids,
and conversely, a significant upregulation after CTCF KD in RV ones (Figure 4E). Together,
these observations indicate that GATA4 and CTCF exert opposing control over B-to-A gene
expression and, more broadly, cardiomyocyte development of the SHF, particularly the RV.

GATA4 and CTCF regulate the timing and quality of first heart field differentiation

Because CTCF silencing produced limited changes in LV cardioids when assessed in bulk, we
investigated this chamber type with higher resolution using scRNA-seq. To capture potential
early differences, we analyzed LV cardioids at both day 4.5 and day 7.5 of differentiation, each
in biological duplicate. We reasoned that LV differentiation may progress so rapidly that early
transcriptional changes could be compensated by later stages. This may explain some of the
differences between LV *versus* and SHF-derived cardioids, which in our protocol enter the final
stage of differentiation one day later, to mimic the slower timing of SHF development[27,38].

Day 4.5 LV cardioids from inducible KD lines maintained in control conditions were highly
similar to each other and to TET-treated SCR controls, as expected (Figure S5A). These car-
dioids were composed primarily of CMs (~50%) and their progenitors (~10%), fibroblasts (~25%),
and a smaller fraction of earlier developmental populations (Figures 5A–5B and Figure S5B).
Biological replicates also showed strong reproducibility (Figure S5A). GATA4 KD resulted in
an expansion of undifferentiated and mesodermal cells accompanied by a reduction in CMs
(Figure 5B). In contrast, CTCF did not alter the overall CM fraction. Notably, both perturbations
markedly altered CM identity, generating clusters that were virtually unique to GATA4 or CTCF
KD. Pseudotime analysis ranked these CM populations as the least mature (GATA4 KD) and
most mature (CTCF KD), respectively, with the trajectory correlating with increasing expres-
sion of B-to-A genes such as *TTN* and *RYR2* (Figure 5C). Milo-based differential abundance
testing[44] confirmed that the shifts in cell populations after GATA4 and CTCF KD were robust
across replicates and at the neighborhood level (Figure 5D).

Figure 5. Single-cell RNA-seq of differentiating inducible knockdown LV cardioids
Legend on the next page

Figure 5. Single-cell RNA-seq of differentiating inducible knockdown LV cardioids (continued)

(A) UMAP representation of scRNA-seq data from two differentiations of day 4.5 FHF-derived LV cardioids with inducible knockown.

(B) Cell type composition, shown as percentage of cells per cluster. C: control; T: tetracycline.

(C) Gene expression in cardiac progenitors and CM subtypes at day 4.5, ranked by pseudotime.

(D) Differential cluster abundance analysis at day 4.5 using graph neighborhoods (KNN). Significantly different clusters (*) are identified by an absolute median logFC > 0.05, a fraction of significant cells > 0.75, and a minimal spatial FDR < 0.05.

(E) UMAP representation of scRNA-seq data from two differentiations of day 7.5 FHF-derived LV cardioids with inducible knockown.

(F) Cell type composition, shown as percentage of cells per cluster.

(G) Gene expression in cardiac progenitors and CM subtypes at day 7.5, ranked by pseudotime.

(H) Differential cluster abundance analysis at day 7.5 using graph neighborhoods (KNN). Significant changes defined as described in panel D..

By day 7.5, control LV cardioids again showed strong similarity (Figure S5A). Their overall
cellular composition was comparable to day 4.5, though CMs appeared more mature, as ex-
pected (Figures S5C–S5D). At this stage, GATA4 KD produced a distinct cluster of immature
CMs with reduced B-to-A gene expression (Figures 5E–5G). In contrast, CTCF KD no longer
yielded a distinct CM cluster but instead showed a modestly lower CM fraction, consistent with
flow cytometry results (Figure 4B). This reduction was accompanied by the expansion of a
poorly defined population co-expressing fibroblast and cardiac markers (Figure 5H and Fig-
ure S5C). Together, these findings indicate that while CTCF functions as a general brake on
CM differentiation, it is nonetheless required for proper LV chamber development.

GATA4 promotes cardiomyocyte cell cycle exit

To assess the functional consequences of GATA4 and CTCF silencing on cardioid morphogene-
sis, we quantified LV cardioid size from day 2.5 to 7.5 of differentiation. GATA4 KD cardioids be-
came significantly larger, whereas CTCF KD had no marked effect (Figure 6A and Figure S6A).
Because scRNA-seq had indicated an accumulation of immature cardiomyocytes in GATA4 KD
cardioids, we hypothesized that this enlargement might reflect increased proliferation. To test
this, we simultaneously measured DNA synthesis and DNA content in day 4.5 LV cardioids, an
assay commonly used to assess cell cycle distribution. GATA4 KD resulted in a significantly
lower proportion of titin$^+$ CMs in G1 and a higher fraction in S and G2 phases (Figures 6B–6C
and Figure S6B). Non-CMs also appeared slightly more proliferative (Figure S6C). Histological
staining of MKI67, a marker of DNA synthesis, supported these findings (Figure S6D).
Because maturing CMs can undergo endoreplication leading to polyploidy, DNA synthesis
and G2-like DNA content alone cannot conclusively demonstrate proliferation. We therefore
assessed phospho-histone H3, a more specific marker of late G2/M, which was also increased
in GATA4 KD cardioids (Figure 6D). To obtain a direct measure of cytokinesis, we labeled car-
dioids at day 3.5 with a cell tracer and monitored its dilution through day 6.5. This confirmed an
increased proliferation rate, particularly in titin$^+$ CMs (Figure 6E and Figures S6E–S6F). Finally,
bioinformatic inference of cell cycle phase from scRNA-seq data further supported increased
proliferation of GATA4 KD CMs (Figure 6F). Altogether, these data show that GATA4 promotes
cardiomyocyte cell cycle exit, whereas CTCF silencing, despite promoting cardiac maturation,
has little effect on proliferation, underscoring their distinct roles in LV development.

Figure 6. Proliferation dynamics in inducible knockdown LV cardioids

(A) Quantification of LV cardioid size progression. n = 3 differentiation, 8 cardioids each. p-values by two-way ANOVA with Holm-Sidak corrected pairwise multiple comparisons.

(B) Representative flow cytometry plots of cell cycle distribution in titin[+] CMs at day 4.5, based on 5-ethynyl-2′-deoxyuridine (EdU) incorporation and DNA content.

(C) Quantification of the experiment shown in panel B. n = 3 differentiations, p-values by two-way RM ANOVA with Holm-Sidak corrected pairwise multiple comparisons.

(D) Immunofluorescence of day 4.5 LV cardioids stained for α-actinin 2 (ACTC2), nuclei (DAPI) and phospho-histone H3. Scale bar: 100 μm.

(E) Proliferation assay with CellTrace dye. Cardioids were labelled at day 3.5 and analyzed by flow cytometry at day 6.5. Data show fold dilution of CellTrace signal in titin[+] CMs, normalized to baseline (day 3.5). n = 3 differentiations, p-value by one-sample t-test against μ = 1.

(F) Bioinformatic inference of G2/M phase from scRNA-seq data of day 4.5 cardioids. p-value by Wilcoxon test.

DISCUSSION

It is well established that genome architecture is highly dynamic during embryonic development. Here, we leveraged monolayer cardiomyocyte differentiation and chamber-specific multilineage 3D organoids to shed light on the mechanisms driving lineage-specific gene activation during a clinically relevant developmental trajectory. Indeed, CHD is the most common birth defect, affecting ~1% of all live births and ~0.4% of adults[24]. Our working model is that GATA4 plays a pivotal role in the early chromatin remodeling required to activate cardiomyogenesis, while CTCF puts a brake on progression towards cardiomyocyte maturation. We propose that these concerted activities ensure a sufficient pool of early cardiomyocytes before cell cycle exit and maturation, and that disruption of either checkpoint compromises chamber formation (Figure 7).

Figure 7. Working model of GATA4 and CTCF function in cardiomyocyte differentiation
During cardiomyocyte differentiation, GATA4 acts as a pioneer transcription factor that drives decompaction and transcriptional activation of cardiac genes undergoing B-to-A switching, exemplified by *TTN*. Reduced GATA4 expression leaves cardiac progenitors in an immature, proliferative state, marked by low cardiac gene expression and persistent association of these loci with the nuclear lamina. Conversely, CTCF, which is highly expressed in pluripotent cells, restricts cardiac gene accessibility and prevents premature activation. As differentiation progresses, declining CTCF levels allow timely chromatin remodeling. Premature CTCF depletion facilitates cardiac maturation but disrupts proper chamber morphogenesis.

In line with a previous hypothesis on *ENPP3* regulation in HEK239T cells[45], our data indicate that CTCF contributes to the compact organization of *TTN* in the undifferentiated state through repressive loops formed at convergent binding sites in its 5' region. These loops restrict *TTN* accessibility when not required and may represent a general mechanism for other B-to-A cardiac genes regulated by CTCF. Notably, CTCF appears to act as a local chromatin remodeller without affecting *TTN* radial positioning, consistent with previous reports[46]. During differentiation, CTCF levels decrease to about half of those in hPSCs, which may be sufficient to release weaker binding sites at *TTN*. Other "epigenetic brakes" have also been described in differentiating cortical neurons[47], suggesting a broader role for such mechanisms in development.

CTCF detachment coincides with activation of the pioneer factor GATA4, rendering *TTN* accessible to transcription at the appropriate time. We previously observed that GATA4 binding is enriched at newly accessible chromatin regions in both cardiac progenitors and cardiomyocytes, whereas NKX2-5 and TBX5, key cardiac TFs which lack pioneer activity, are enriched only at cardiomyocyte-specific accessible sites[13]. In TBX5 haploinsufficient cardiomyocytes, reduced TBX5 dosage increased CTCF binding at cardiac genes[14]. Together, these findings support a model in which GATA4-triggered chromatin remodeling enables recruitment of other cardiac transcription factors, stabilizing a CTCF-free chromatin environment productive for transcription.

We confirmed the importance of GATA4 in driving cardiac specification and maturation[48,49]. In particular, GATA4 binding to cardiac genes is fundamental for their B-to-A compartment transition, which is coupled with increased expression key for cardiomyocyte differentiation and maturation. Consistent with rat models that implicated GATA4 as a repressor of cardiac fibroblast genes[50], GATA4 knockdown in our system led to fibroblast accumulation. In contrast to mouse studies[51], we observed an increase in immature cardiomyocytes in a proliferative state in left ventricle organoids, suggesting a human-specific mechanism.

Much less is known about CTCF in heart development. Conditional CTCF knockout in cardiac progenitors causes severe cardiac malformation and embryonic death[52]. Interestingly, these studies reported premature arrest of cardiac development with concomitant promotion of cardiomyocyte maturation, consistent with our left ventricle single-cell data showing upregulation of electrophysiologically relevant genes such as *RYR2*. A recent mouse study found that a homozygous CTCF point mutation (R567W) severely impairs cardiac development, reducing ventricular cavity size and downregulating genes involved in sarcomere assembly and calcium handling[53]. This mutation affects the zinc finger 11, required for upstream motif recognition, and disrupts CTCF binding. In contrast, our inducible knockdown approach mimics and anticipates the physiological decrease of CTCF expression observed in the human heart and reproduced across hESC and hiPSC models. We propose that this partial and timed CTCF reduction recapitulates developmental regulation without broadly disrupting TADs and chromatin loops. Thus, whereas strong CTCF disruption is pathological, partial downregulation may represent a physiological mechanism during human cardiomyogenesis.

GATA4 and CTCF are mutated in patients with isolated CHD or in syndromes that include CHD[18,25,26,54]. The mechanism we propose may therefore provide new insight into CHD pathogenesis, though patient-specific hiPSCs will be needed to test this directly. More broadly, our work illustrates how differentiating hPSCs into human-relevant organoids that recapitulate distinct heart chambers can be leveraged to dissect CHD pathophysiology and clarify the causal links between nuclear architecture dysregulation and human disease.

Data, Code, and Material Availability

Raw NGS data for bulk and scRNA-seq have been deposited in ArrayExpress under accession numbers E-MTAB-15684, E-MTAB-15693, E-MTAB-15710, and E-MTAB-15711, while processed data is available on Zenodo (https://doi.org/10.5281/zenodo.17256128). 4C-seq data have been deposited in ENA under accession number PRJEB97970. All custom code is available at GitHub (https://github.com/alessandro-bertero/Becca_et_al_2025). Computational analyses were conducted within containerized environments to ensure reproducibility: Hi-C (hedgelab/hic_pipeline:image12), bulk RNA-seq (hedgelab/bulk_image:image3), single-cell RNA-seq (hedgelab/cellranger8:latest for preprocessing; hedgelabrstudio_image:iPS2seq for integration and downstream analysis), and 4C-seq (hedgelab4c:image1). All other data, code, and material used for the study is detailed in the Key Resources Table.

Acknowledgements

We thank Giulia Savorè, Federica Sozza, Sveva Bottini, Matteo Sorge, Maddalena Arigoni, Alessia Labate, Paola Circosta, and Luca Ponzone for experimental assistance; Manuel Rosa Garrido for advice on CTCF ChIP; Estela Mancheno Juncosa for advice on cardioids; Antonio Grimaldi for MGI sequencing; and the Vienna BioCenter Core Facility for Illumina sequencing. We are also grateful to the UniTo facility staff, including Francesca Anselmi (NGS), Federica Antico (histology), Laura Conti (FACS), and Marta Gai (microscopy). This work was supported by the FEBS Excellence Award 2022 (A.B.), the Additional Ventures Single Ventricle Research Fund 2022 (A.B. and S.M.), and the Giovanni Armenise-Harvard Foundation Career Development Award 2021 (A.B.).

Author Contributions

Conceptualization: S.Be., A.B.; Methodology: S.Be., E.M.H.; Investigation: S.Be., E.M.H., K.E.S., L.T., A.K., E.B., A.B.; Formal analysis: S.Bi.; Data curation: E.B.; Software: S.Bi., E.B.; Visualization: S.Be., S.Bi., A.B.; Resources: F.N., D.C., S.M., A.B.; Supervision: S.M., E.B., A.B.; Project administration: A.B.; Funding acquisition: A.B.; Writing – original draft: S.Be., A.B.; Writing – review & editing: all authors.

Competing Interests

The authors declare no competing interests.

METHODS

Stem cell models

hiPSC culture

hiPSCs from an apparently healthy male donor (WTC11)[55] and WTC11 titin-mEGFP reporter hiPSCs[56] were cultured on Geltrex or Matrigel in Essential 8 or mTeSR Plus. Colonies were passaged as small clumps using Versene (0.5 mM EDTA in PBS) and replated in media containing 2 µM thiazovivin for 24 h. Genomic stability was verified by microarray-based karyotyping.

hiPSC-CM differentiation

Monolayer cardiac differentiation was performed according to a biphasic WNT signaling protocol[13]. On day –2, cells ($1.5–3 \times 10^5$ per well in 12-well plates) were seeded with 2 µM thiazovivin. On day –1, they were primed with 1 µM CHIR99021 in pluripotency medium. On day 0, differentiation was induced in RBA medium [RPMI 1640 supplemented with 0.5 mg mL^{-1} BSA and 213 µg mL^{-1} L-ascorbic acid 2-phosphate] with 5 µM CHIR99021. On day 2, 2 µM WNT-C59 was added to direct cardiac mesoderm formation. On day 4, the medium was replaced with fresh RBA, and from day 6 cells were maintained in RPMI + B27 every other day.

Cardiac organoids

Cardiac organoids (cardioids) were generated following the protocol of Schmidt *et al.*[27] with minor modifications. On day –1, hiPSCs (5×10^4 per well) were seeded in 24-well plates with 5 µM Y-27632. After 24 h, the medium was replaced with mesoderm (Mes) formulation. After 36 h, cells were detached with TrypLE Express, resuspended in cardiac mesoderm (CardMes) medium containing 5 µM Y-27632, and seeded into ultra-low-attachment 96-well plates (1.5×10^4 cells per well), then centrifuged at 140 g for 4 min to promote aggregation. The following days (from day 2.5 onward), spheroids were fed with fresh CardMes medium without Y-27632 until day 5.5, after which they were fed with Cardio medium. All media were prepared in chemically defined medium (CDM; 50% IMDM and 50% Ham's F12 supplemented with 0.4% BSA, 1% lipids, 0.004% monothioglycerol, and 15 µg mL^{-1} transferrin). Growth factors and small molecules were adjusted according to chamber identity. For FHF–LV, Mes contained 6 ng mL^{-1} FGF2, 5 µM LY294002, 5 ng mL^{-1} Activin A, 8 ng mL^{-1} BMP4, and 5 µM CHIR99021; CardMes contained 8 ng mL^{-1} BMP4, 1.6 ng mL^{-1} FGF2, 10 µg mL^{-1} insulin, 2 µM WNT-C59, and 50 nM retinoic acid; Cardio contained 8 ng mL^{-1} BMP4, 1.6 ng mL^{-1} FGF2, and 10 µg mL^{-1} insulin. For SHF-RV and SHF-A, Mes contained 6 ng mL^{-1} FGF2, 5 µM LY294002, 50 ng mL^{-1} Activin A, 10 ng mL^{-1} BMP4, and 6 µM CHIR99021. During CardMes (days 1.5–4.5), both RV and A were first cultured (days 1.5–2.5) with 1 µg mL^{-1} insulin (Day 1.5 only) and 10 µM SB431542; 2 µM WNT-C59 was added for RV and 5 µM XAV-939 plus 500 nM retinoic acid for A. From days 3.5–4.5, both received 10 ng mL^{-1} BMP4, 1.6 ng mL^{-1} FGF2, 10 µg mL^{-1} insulin, and 500 nM retinoic acid, maintaining 2 µM WNT-C59 for RV and 5 µM XAV-939 for A. The final Cardio (from day 5.5) contained 10 ng mL^{-1} BMP4, 1.6 ng mL^{-1} FGF2, and 10 µg mL^{-1} insulin.

Genome editing

Inducible knock-down

hiPSCs were engineered with tetracycline-inducible shRNAs targeted to the *AAVS1* genomic safe harbour using an established approach[57]. Validated shRNAs from the TRC library (GATA4: TRCN0000020424; CTCF: TRCN000021849) and a scrambled control (SCR: SHC016) were cloned into pAAV-Puro_siKD by ligating annealed oligonucleotides with BglII and SalI over-hangs (Table S1). A 5' terminal G was added to ensure efficient transcription from the H1 promoter. Cells (3×10^5 per well) were seeded in 6-well plates and transfected with Gene-Juice and 2 µg total plasmid DNA [pAAV-Puro_siKD, pZFN-AAVS1_ELD, pZFN-AAVS1_KKR; equimass ratio] in Opti-MEM. Medium was refreshed after 24 h, and genome-edited cells were selected with 1 µg mL^{-1} puromycin from days 3–10, with 10 µM Y-27632 added on days 3–5. To avoid clonal bias and given the >90% on-target efficiency of this method[39], all surviving hiPSCs were pooled for experiments. Knock-down was induced by addition of 1 µg mL^{-1} tetracycline, starting 3 days prior to differentiation, with daily medium changes.

CTCF binding site (CBS) editing

To identify candidate CBS within the *TTN* locus, ChIP-seq data for hESCs and hESC-derived CMs were downloaded from the ENCODE Data Portal. BigWig data were visualized on the GRCh38 genome using IGV to define six CTCF binding regions. Core motifs within each region were predicted using CTCFBSDB2.0 and confirmed with FIMO (MEME Suite). A CTCF bind-ing site score was computed with Bioconductor to estimate the degree of consensus sequence match, which served as a proxy for binding strength. To delete CBS1 and CBS6, WTC11 hiP-SCs were genome edited using single guide RNA (sgRNA)-directed Cas9 cleavage and HDR with ssODN templates. sgRNAs were designed using GuideScan2 and IDT's CRISPR-Cas9 design checker, selecting guides with no off-targets in coding regions expressed in hiPSCs or hiPSC-CMs (CBS1: 5'-CACCGTATTCCCTCTTTGACCACTA -3'; CBS2: 5'-CACCGTATTCCC-TCTTTGACCACTA-3'). sgRNAs with BbsI overhangs were cloned as annealed double stranded oligos into pX330-U6-Chimeric_BB-CBh-hSpCas9 following standard protocols based on T4 PNK and T4 DNA ligase. The ssODN for CBS6 was synthesized on the strand opposite to the sgRNA, with 50 bp homology arms, while the ssODN for CBS1 was on the same strand, with 80 bp and 30 bp arm. Both replaced the core CTCF motif with an EcoRI site to facilitate genotyping[36]. hiPSCs were detached with Accutase and 2×10^6 cells were electroporated with 2 µg pX330-gRNA plasmids (1 µg per site), 100 pmol ssODN per target, 2 µg AAVS1 SA-2A-puro-pA donor, and 2 µg each of pZFN-AAVS1_KKR and pZFN-AAVS1_ELD using the P3 Primary Cell 4D-Nucleofector Kit (CA-137 program). Cells were recovered in E8 + CEPT (50 nM chroman 1, 5 µM emricasan, 1:1000 polyamine supplement, 0.7 µM trans-ISRIB) overnight. Puromycin selection was applied to enrich HDR-competent cells[40], followed by clonal isolation by single-cell sorting into 96-well plates. Genomic DNA was extracted, and targeted loci were amplified by PCR, digested with EcoRI to detect edited alleles, and visualized by agarose gel electrophoresis. Positive clones were validated by PCR to exclude random plasmid integra-tion (Table S2), followed by Sanger sequencing of the target regions. Heterozygous edits were resolved by TOPO-TA subcloning prior to sequencing. Selected clones were karyotyped by low-resolution microarray to confirm genomic stability.

Functional assays

Flow cytometry

hiPSC-CMs were dissociated for FACS by incubating them in 0.25% trypsin for 10–15 min. The reaction was stopped with RPMI-B27 supplemented with 20% FBS, and cells were pelleted at 190 g for 5 min. After a PBS wash, cells were resuspended in PBS containing Fixable Viability Dye (eFluor 780 or 450) and incubated in the dark for 10 min. Following two washes in FACS buffer (PBS + 5% FBS), live cells were either directly analyzed for titin-mEGFP reporter activity or fixed for antibody staining. For antibody staining, cells were washed in FACS buffer twice and permeabilized during centrifugation (250 g, 5 min) in permeabilization buffer (FACS buffer + 0.75% saponin), then incubated for 50 min with either mouse anti-cardiac Troponin T (Alexa Fluor 647, 1:100) or rabbit anti-titin (targeting the Z-disc Z1Z2 region, 1:500). Titin staining was followed by a 1 h incubation with Alexa Fluor 488 goat anti-rabbit secondary antibody (5 µg mL^{-1}). After washing, cells were resuspended in FACS buffer and analyzed on a FACSVerse cell analyzer, acquiring $\sim$30,000 events per condition. Events were gated with FlowJo v10.10.0 to exclude debris, doublets, and dead cells. For cardioid dissociation, up to 8 organoids per condition were pooled, washed twice with 0.5 mM EDTA, and incubated in 70 µL of 0.5% trypsin in EDTA at 37°C for 10–20 min, mixing every 5 min. The reaction was stopped with CDM + 10% FBS, and cells were pelleted at 190 g for 5 min. Live titin-mEGFP recordings were performed as for hiPSC-CMs, except that FACS buffer consisted of PBS + 1% BSA.

Morphometry

Cardioid images were acquired daily from day 2.5 to day 7.5 of differentiation using the Incucyte SX5 Live-Cell Analysis System (Sartorius) with the spheroid analysis module and a 4$\times$ objective. Size measurements were obtained by applying filters for cardioid identification, including a minimum object area of $3 \times 10^4 \mu m^2$ and a maximum eccentricity of 0.9. To exclude debris, quantification was restricted to the largest object per field.

Cell cycle assay

DNA synthesis was assessed using the Click-iT^{TM} EdU Alexa Fluor 647 Flow Cytometry Assay. On day 4.5, cardioids (8 per condition) were incubated with 10 µM EdU in culture medium for 2 h at 37°C. After incubation, cardioids were dissociated as described above and stained for dead cells with Fixable Viability Dye (eFluor 780). Cells were then fixed with 50 µL Click-iT$^{®}$ fixative (Component D) for 15 min at room temperature, washed with 1% BSA in PBS, and resuspended in Click-iT$^{®}$ saponin-based permeabilization and wash reagent. Following 15 min permeabilization, the Click reaction cocktail was added and incubated for 30 min at room temperature in the dark. Cells were washed again with saponin-based buffer and subsequently stained with anti-titin primary and secondary antibodies (as described for flow cytometry) in the same buffer. Finally, DNA content was measured by adding FxCycle Violet Stain (1:1,000 in Click-iT$^{®}$ saponin-based buffer) and incubating for 30 min at room temperature in the dark. Samples were analyzed on a FACSVerse cell analyzer.

Histology

Organoids were collected using wide-bore tips, fixed in 4% PFA at room temperature for 30 min with gentle shaking, and washed twice with PBS. Samples were incubated overnight at 4°C in 30% sucrose/PBS, embedded in O.C.T., and cryosectioned at 12 µm using a Leica CM1950 cryostat. Sections were stored at –20°C and thawed before staining. For staining, sections were washed in PBS for 15 min at room temperature, then permeabilized in PBST-D1 (4% goat serum, 1% Triton-X in PBS). Primary antibodies were diluted in PBST-D1 and applied within a hydrophobic barrier, incubating in the dark for 3 h at room temperature. The following primary antibodies were used: mouse anti-Ki67 (1:200), rabbit anti-α-actinin 2 (1:200), and mouse anti-pHH3 (1:50). After incubation, slides were washed in PBST2 (PBS + 0.1% Tween-20) for 15–30 min. Secondary antibodies (goat anti-rabbit Alexa Fluor 488, 1:500; goat anti-mouse Alexa Fluor 633, 1:500; goat anti-mouse Alexa Fluor 568, 1:500) were diluted in PBST2 and applied for 1 h at room temperature in the dark within a humidified chamber. After washing in PBST2, nuclei were counterstained with Hoechst 33342 (1:2000 in PBS, 5 min), then washed again. Slides were air-dried and mounted with ProLong Glass Antifade Mountant. Confocal imaging was performed using a Leica SP8 microscope with a 63X objective and the following laser settings: excitation/emission 488/520 nm (Alexa Fluor 488), 631/650 nm (Alexa Fluor 633), and 579/603 nm (Alexa Fluor 568).

Proliferation assay

Cell division was assessed using the CellTrace^{TM} Far Red Cell Proliferation Kit for flow cytometry. On day 3.5, 16 cardioids per condition were incubated with 2 µM CellTrace dye for 1 h at 37°C in PBS containing Ca^{2+}/Mg^{2+}. After incubation, cardioids were washed twice with basal medium and returned to culture in complete medium. Half of the labeled cardioids were immediately dissociated and stained for live/dead discrimination (as described for flow cytometry), and basal CellTrace fluorescence was acquired on a FACSVerse analyzer. The remaining cardioids were maintained in culture under standard conditions until day 6.5, then dissociated, stained for live/dead cells, and analyzed by flow cytometry. Cell division was quantified by calculating the median fluorescence intensity (MFI) of titin$^+$ cells at day 6.5, normalized to the MFI at day 3.5. The resulting ratio for tetracycline-treated samples was then normalized to the matched untreated controls.

Gene expression assays

Reverse transcription quantitative PCR (RT-qPCR)

Total RNA was extracted using the Quick-RNA MiniPrep Kit according to the manufacturer's instructions. cDNA synthesis was performed with the High-Capacity cDNA Reverse Transcription Kit. qPCR reactions were prepared with 10 ng cDNA, 250 nM primer mix, and either PowerUp SYBR Green 2X Master Mix or Luna® Universal qPCR Master Mix. All reactions were run in technical duplicates on a QuantStudio 6 Flex Real-Time PCR System. Gene expression was normalized to the geometric mean of *RPLP0* and *PBGD* or to *HPRT1*, as indicated. Primer sequences are listed in Table S3.

Bulk RNA sequencing (RNA-seq)

RNA-seq for monolayer hiPSC-CMs was performed on biological duplicates using the Zymo-Seq RiboFree Total RNA Library Prep Kit with 650 ng input RNA per sample. Ribodepletion was carried out for 1 h, and libraries were amplified with 11 PCR cycles. Library quality was assessed by Qubit DNA HS and acrylamide gel electrophoresis, and sequencing was performed on an Illumina NextSeq 550 (75 bp paired-end). RNA-seq for cardiac organoids was performed on biological duplicates using the Illumina TruSeq Stranded mRNA Library Prep with 400 ng RNA input, 11 PCR cycles, and library QC by Agilent Tapestation. Sequencing was performed on an MGI T7 platform (100 bp paired-end). After demultiplexing with bcl2fastq, reads were trimmed with Trim Galore (v0.6.10) and aligned with STAR v2.7.11b using the ENSEMBL GRCh38 (release-112) genome and annotation. Gene counts were imported into R and processed with DESeq2 and the RNAseqQC package for quality control, filtering (protein-coding genes with ≥ 3 counts in at least two samples), and variance stabilization. Batch effects were corrected with the `removeBatchEffect` function (limma)[58]. QC included analysis of library complexity, gene detection, replicate correlation, and PCA. For expression normalization, raw counts were converted to TPM using ADImpute with gene lengths from `gtftools`. TPM values were batch-corrected and used for WGCNA network analysis[59], with GO enrichment of modules performed via `goana` and `topGO` (limma)[58]. Differential expression was conducted on raw filtered counts normalized with the `voom` function (limma), followed by contrasts testing, moderated statistics (`eBayes`), and reporting with `topTreat`. Genes were ranked by $\log_2$ fold change and analyzed with `gseGO` (ClusterProfiler)[60] for biological process enrichment. For cardiac organoids, trimmed and aligned reads were analyzed as above. Heatmaps of B-to-A genes were generated from the average $\log_2$ fold change TPM values after batch correction, using the `pheatmap` package. Differential expression comparisons were performed separately for atrial and right ventricle genotypes. Enrichment of cardiac processes was visualized with `gseaplot` (enrichplot). Rank-rank hypergeometric overlap (RRHO) analysis was performed with RedRibbon, comparing $\log_2$ fold changes of CTCF (TET vs. no TET) and GATA4 (TET vs. no TET) in atrial and ventricular samples. RRHO maps were visualized with the `ggRedRibbon` function.

Single-cell RNA-seq (scRNA-seq)

FHF-LV cardioids at days 4.5 and 7.5 were analyzed in biological duplicates by microfluidics-based scRNA-seq (10X Genomics). Sixteen cardioids per condition were pooled, dissociated with 0.5% trypsin/0.5 mM EDTA, quenched with RPMI-B27 + 20% FBS, centrifuged (200 g, 5 min, 4°C), and resuspended in PBS/1% BSA. Cells were counted and labeled with barcoded lipid-conjugated cell multiplexing oligos (CMOs; 0.5 million cells/condition, 100 µL, 5 min, RT), washed, stained with Fixable Viability Dye, and live-sorted on a Sony SH800S (100 µm chip, targeted mode, single-cell 3-drop). Controls (2.5×10^5 cells each) and tetracycline-treated samples (7.5×10^5 cells) were pooled, and 5.0×10^5 total cells were loaded per channel on the Chromium Next GEM Chip G. Libraries were prepared with the Chromium Next GEM Single Cell 3' Reagent Kit v3.1 and sequenced on an Illumina NovaSeq 6000 (S2 100 cycles, 18+90) to a depth of $\sim$4 billion reads ($\sim$18,000 singlets/condition, $\sim$50,000 reads/GEX, $\sim$7,500 reads/CMO per cell). A similar workflow was applied to pooled FHF-LV, SHF-LV, and SHF-A cardioids (3.0×10^5 cells total), sequenced on an Illumina NextSeq 1000 (P2 100 cycles). Raw

fastq files were processed with Cell Ranger v8.0.1 (multi, aggr without normalization) against GRCh38. Count matrices were imported with `Read10X` and analyzed with Seurat. Filtering criteria included ≥200 expressed genes per cell, genes expressed in ≥3 cells, mitochondrial content ≤15%, and removal of outliers (lowest 5% / highest 1% by UMI or gene counts). Data were normalized with `NormalizeData` (log, scale factor 10,000), scaled with `ScaleData`, and scored for cell cycle phases using `CellCycleScoring`. Filtered matrices were imported into Monocle3 for trajectory analysis. Dimensionality reduction was performed with PCA and UMAP; clustering with `cluster_cells` (partition q-value 0.05); and pseudotime inference with `learn_graph` and `order_cells`, rooting at the branch enriched for pluripotency markers. Cluster stability was assessed by resampling (10 iterations, 1,000 cells removed each). Marker genes were identified with `top_markers`, and GO enrichment was performed with `goana` and `topGO`. Differential expression was tested at the cluster level (`fit_models`, Poisson; GSEA with `gseGO`) and along pseudotime (`graph_test`, q < 0.05). Cardiomyocyte clusters were further sub-analyzed by pairwise comparisons (`fit_models`, negative binomial), followed by enrichment with `gseGO` and `gseKEGG`. For condition-specific comparisons, untreated controls (all genotypes pooled) were compared to each tetracycline-treated genotype using Fisher's exact test with multiple-testing correction (`p.adjust`). Finally, the differential abundance of cells per cluster was assessed using MiloR. kNN graphs were built with `buildGraph` (k = 30, 30 dimensions), and overlapping neighborhoods were generated with `makeNhoods`. Neighborhood distances were computed with `calcNhoodDistance` (d = 30), and differential abundance was tested with `testNhoods`, using shRNA-matched CTR as the reference condition. Results were summarized at the cluster level by computing the median logFC, the fraction of significant neighborhoods, direction consistency, and the minimum spatialFDR. Clusters were considered significant when minSpatialFDR ≤ 0.05 with median logFC > 0.5 or < –0.5 and frac_sig > 0.75. Cluster ID transferring was performed by importing GSE263193 scRNAseq counts into Seurat, filtering for genes expressed in both datasets, performing $\log_2$ normalization and data processing with the default Seurat pipeline. Transferring anchors were identified with the `FindTransferAnchors` function and transferred with the `TransferData` function.

Chromatin and nuclear architecture assays

Chromatin immunoprecipitation (ChIP)-qPCR

Chromatin immunoprecipitation (ChIP) for CTCF was performed with minor modifications of a published protocol[61]. Cells were crosslinked in 1% formaldehyde for 15 min at room temperature with gentle agitation, and the reaction was quenched with 125 mM glycine for 10 min. All subsequent steps were performed on ice unless otherwise indicated. After washing with cold PBS, cells were scraped, pelleted (250 g, 5 min, 4°C), and lysed in cell lysis buffer (10 mM Tris-HCl pH 8.0, 10 mM NaCl, 0.2% NP-40) supplemented with protease inhibitors. Nuclei were isolated by centrifugation (600 g, 5 min), then lysed in nuclear lysis buffer (50 mM Tris-HCl pH 8.0, 10 mM EDTA, 1% SDS). Chromatin was diluted in buffer (20 mM Tris-HCl pH 8.0, 2 mM EDTA, 150 mM NaCl, 0.01% SDS, 1% Triton X-100) and sheared with a Bioruptor Pico (7 cycles of 30 s ON/30 s OFF). The lysate was clarified (16,000 g, 10 min, 4°C), and an aliquot was saved as input. The remaining solution was incubated overnight at 4°C with a mix of 2 µL rabbit anti-CTCF and 8 µL mouse anti-CTCF antibodies, or with matched rabbit IgG and

mouse IgG1κ isotype controls. Immunocomplexes were captured for 1 h at room temperature with Protein A/G magnetic beads, which had been pre-washed in wash buffer. Beads were sequentially washed twice with buffer I (20 mM Tris-HCl pH 8.0, 2 mM EDTA, 50 mM NaCl, 0.1% SDS, 1% Triton X-100), once with buffer II (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 0.25 M LiCl, 1% NP-40, 1% sodium deoxycholate), and twice with TE (10 mM Tris-HCl pH 8.0, 1 mM EDTA). DNA was eluted twice in 100 mM sodium bicarbonate, 1% SDS, for 15 min each at room temperature. Eluates and input were treated in parallel with 1 µg RNase A and 300 mM NaCl, then incubated 5 h at 65°C to degrade RNA and reverse crosslinks. Samples were then treated overnight with 60 µg Proteinase K at 45°C and purified with the ChIP DNA Clean & Concentrator kit. qPCR was performed on target regions using 4 µL purified DNA, 5 µL 2X Luna Universal qPCR Master Mix, and 2.5 µM primer mix (Table S4). Ct values were normalized to inputs and then to the average Ct of four negative control regions lacking CTCF binding and compared to two constitutively-bound positive controls on the same chromosome (hg38 chr2:176140583-176140677 and chr2:176140649-176140792).

Circularized Chromosome Conformation Capture with sequencing (4C-seq)

4C was performed following an established protocol[62] with minor modifications. hiPSCs were detached, pelleted (100 g, 5 min), and resuspended at 2×10^6 cells/mL in isolation buffer (10% FBS in PBS). Fixation was carried out by adding an equal volume of 4% formaldehyde in isolation buffer and incubating for 10 min at room temperature with shaking, followed by quenching with 0.13 M glycine for 5 min. Cells were pelleted (500 g, 5 min, 4°C), washed in PBS, and lysed in ice-cold lysis buffer (50 mM Tris-HCl pH 8.0, 0.5% NP-40, 1% Triton X-100, 150 mM NaCl, 5 mM EDTA) supplemented with protease inhibitors. Nuclei were pelleted (500 g, 5 min, 4°C) and resuspended in 1.2× RE1 buffer with 0.3% SDS, incubated for 1 h at 37°C with shaking, then treated with 2.5% Triton X-100 for 1 h under the same conditions. Digestion was performed with 100 U DpnII for 3 h at 37°C, followed by an additional 100 U overnight. Digestion efficiency was verified on agarose gel. After heat inactivation, samples were ligated overnight at 16°C in 2 mL reactions containing 50 U T4 ligase. Crosslinks were reversed with 75 µg/mL proteinase K, and DNA was purified by isopropanol precipitation and ethanol washes. DNA was eluted in nuclease-free water and subjected to a second digestion with Csp6I (overnight at 37°C, 20 min at 65°C, then 4°C), followed by a second ligation (25 µg DNA template in 5 mL reaction, 50 U ligase, overnight at 16°C). DNA was purified again and quantified. PCR was performed in two steps: (1) first-round amplification with LongAmp Taq polymerase (16 cycles, annealing 50°C) using primers with partial Illumina overhangs (reading primer: 5'-TACACGACGCTCTTCCGATCTNNNNNAACCAGCTTAAATTGATC-3'; non-reading primer: 5'-ACTGGAGTTCAGACGTGTGCTCTTCCGATCTTTCAGAACAGAATAGTCATG-3'); (2) second-round amplification with NEBNext High-Fidelity PCR Mix and Illumina dual indexes. PCR products were size-selected with SPRI beads (0.8× ratio). The viewpoint was defined at *chr2:178,805,901–178,806,592*, and 800 ng DNA was used for amplification. Libraries were sequenced on an Illumina NextSeq 1000 (100 cycles kit, 58 cycles read 1). Bioinformatic analysis followed the 4C-seq pipeline[62] within a Docker container (hedgelab/4c:image1). Primer sequences, restriction enzymes (DpnII: GATC; Csp6I: GTAC), and genome reference (hg38) were specified in the viewpoint configuration file. Reads were demultiplexed by primer, trimmed at the 5'-end (DpnII site), mapped to hg38 with bowtie2, and filtered for overlaps with

the in silico digested genome. After read counting and smoothing, wig files were converted to bed format (wig2bed, bedops) and imported into R. Batch correction was performed with `removeBatchEffect` (limma), normalizing z-scores across replicates. Data were restricted to bins overlapping *TTN*, excluding the viewpoint. Area-under-curve (AUC) values between and under CTCF ChIP peaks (ENCFF332TNJ) were computed with `AUC` (DescTools). Difference curves were generated by reciprocal subtraction between conditions with the pyGenomeTracks module[63], and the difference in each region was tested with a one sample t-test (mu = 0). Comparison with virtual 4C tracks (RUES2 hESCs, hESC-MES, hESC-CPs, and hESC-CMs Hi-C data) was performered by binning the 4C tracks at 40kb, filtering for bins on chromosome2, rescaling each track on the library size, $\log_2$ normalizing and fitting on a 20-degree polynomial curve with the `lm` function. For each pair of conditions it was evaluated the Pearson correlation of the polynomial coefficients of the derivatives with the `rcorr` function.

Immunofluorescence paired to DNA fluorescence *in situ* hybridization (Immuno-FISH)

ImmunoFISH following a published protocol[13] with minor modifications. The *TTN* locus-spanning CH17-275G10 BAC (BACPAC Resources Center) was purified with the Qiagen Plasmid Midi Kit, with modified centrifugation steps to preserve large BAC DNA. The product was validated by PCR using two primer pairs amplifying the expected region (5'-AGTAAGGCGGGTGGCTTTAT-3' – 5'-CCACATCACAACCGTGAAAG-3', T_a 56°C; 5'-AGGCT-GGCTGAAAAATAGGTC-3' – 5'-TAATACAACGCATTGGCCCTC-3', T_a 57°C). Probes were labeled by nick translation with Alexa Fluor 488-dUTP (4 h at 15°C, inactivation at 70°C, 10 min), purified, and precipitated with salmon sperm DNA, human Cot-1 DNA, sodium acetate, and ethanol overnight at –20°C. The pellet was resuspended in hybridization buffer (50% formamide, 2$\times$ SSC, 10% dextran sulfate). hiPSC-derived cardiomyocytes (day 24) underwent a 45 min heat shock at 42°C in RPMI + HEPES, were returned to 37°C overnight, and replated (40,000 cells/slide) on geltrex-coated HistoBond slides. After 4 h, cells were fixed in 4% PFA (10 min), quenched with 0.1 M Tris-HCl pH 7.4 (10 min), and cryoprotected in 20% glycerol (20 min) followed by 50% glycerol overnight at –20°C. Slides were permeabilized and blocked in 0.1% Triton X-100 + 1% goat serum in PBS (1 h), incubated overnight at 4°C with rabbit anti-Lamin B1 (1:350) and mouse anti-α-actinin (1:100), and washed. Secondary antibodies (goat anti-rabbit Alexa Fluor 405, 1:500; goat anti-mouse Alexa Fluor 633, 1:500) were applied for 1 h at room temperature. Post-fixation was performed with 3% PFA in PBS (10 min). Slides were treated with 0.1 M HCl + 0.7% Triton X-100 (10 min, 4°C), washed in SSC, and DNA was denatured in 50% formamide/2$\times$ SSC (30 min, 80°C). Denatured probes (10 µL/slide) were applied under coverslips sealed with rubber cement, and hybridized overnight at 42°C. Slides were washed in 50% formamide/2$\times$ SSC (3$\times$, 42°C), then in 2$\times$ SSC at 42°C and room temperature. Slides were mounted in ProLong Diamond antifade, dried 2 h, sealed, and imaged within a few days. Images were acquired on a Zeiss Elyra 7 microscope (63$\times$, NA 1.4, HiLo mode). Z-stacks encompassed the Lamin B1 signal. At least 50 cardiomyocytes per condition were imaged. Analysis was performed in Imaris v10.2: nuclei were segmented with a surface filter (0.5 µm), Lamin B1 and FISH background removed with size filters (9 µm and 1 µm, respectively), and only α-actinin$^+$ cells retained. Distances between FISH spot centers and nuclear boundaries were computed with the distance transformation function, normalized to the average cube root of the nuclear volume, and batch processed with a custom Python script. Non-diploid nuclei

695 (≠2 spots) were excluded. The number of cardiomyocytes analyzed per condition is reported
696 in the figures.

Bisulfite pyrosequencing

Bisulfite conversion of genomic DNA was performed using the Zymo Methylation-Lightning Kit
following the manufacturer's instructions. Bisulfite-treated DNA was amplified with the Pyro-
Mark PCR Kit, and PCR products were sequenced on a PyroMark Q48 Autoprep using Pyro-
Mark Q48 Advanced CpG Reagents. DNA methylation levels were quantified with the PyroMark
Q48 Autoprep Software. Primers were designed with the PyroMark Assay Design Software
2.0. The forward primer was 5'-GAATTTGGTTTTTTATTATTGGTAAGATTG-3', the reverse
primer (biotinylated) was 5'-CTCCAAATAAATTACTCCATAAATCATCTA-3', and the sequenc-
ing primer was 5'-GTTTTTTATTATTGGTAAGATTGTA-3'.

Secondary data integration and analysis

Hi-C data analysis

DNase Hi-C datasets for RUES2 hESCs, hESC-MES, hESC-CP and hESC-CMs were down-
loaded from the 4DN Data Portal. Biological replicates were kept separate, while technical
replicates were concatenated using the `cat` function. Reads were aligned with HiC-Pro v3.1.0[64]
to the hg38 genome, indexed with `bowtie2-build` (UCSC FASTA and chromosome sizes files,
excluding chrX, chrY, and chrM). Default parameters were used, except `GENOME_FRAGMENT` and
`LIGATION_SITE`, which were left empty as required for DNase Hi-C. Differential compartment
analysis was performed with the dcHiC pipeline[65] using ICE-normalized matrices binned at 150
kb. PCA was first computed per chromosome (`pcatype-cis`), the optimal component was se-
lected (`pcatype-select`), and differential analysis was conducted (`pcatype-analyze`) to gener-
ate IGV tracks (`pcatype-viz`). Raw HiC-Pro matrices at 40 kb resolution were converted to `h5`
format with HiCExplorer v3.4.1[66] (`hicConvertFormat.py`). Biological replicates were summed
(`hicSumMatrices.py`), filtered and ICE-normalized (`hicCorrectMatrix.py`), then converted
to `mcool` format and imported into R with `HiCExperiment`. Virtual 4C was performed with
`virtual4C` (HiContacts package)[67] using the *TTN* promoter (hg38 chr2:178,806,793–178,808,363)
as viewpoint. BedGraph outputs were visualized in IGV and final tracks plotted with pyGenome-
Tracks. For H9 hESC-CM differentiation time-course Hi-C, raw `hic` matrices were downloaded
from GEO and converted to HiC-Pro–like sparse matrices with `preprocess.py` (dcHiC envi-
ronment). Chromosomes X, Y, and M were excluded, and resolution was set to 50 and 100
kb. ICE normalization was performed with HiC-Pro v3.1.0 (`step = ice_norm`, default settings).
Differential compartments were identified with dcHiC using 100 kb bins. Virtual 4C was per-
formed at 50 kb resolution as above, with the viewpoint lifted from hg19 to hg38 (UCSC LiftOver;
hg19 coordinates: chr2:179,671,520–179,673,090). For WTC11 hESC-CM, normalized `mcool`
matrices were downloaded from the 4DN DataPortal[14] and converted to HiC-Pro–like sparse
matrices with `preprocess.py` at 50kb and 100kb resolutions. Differential compartment analysis
(100kb) and virtual4C analysis (50kb) were performed as described above.

ChIP-seq data analysis

Publicly available GATA4 ChIP-seq datasets in hESCs and CMs(aligned to hg38) were pro-
cessed as previously described[13]. For CTCF, hESC and CM datasets were downloaded from
ENCODE in `bed` (pseudo-replicated IDR-thresholded peaks) and `BigWig` (fold-change over con-
trol) formats, using the ENCODE4 v1.5.1 GRCh38 annotation. Each `BigWig` file was visualized
together with its corresponding `bed` file using pyGenomeTracks, following manual customiza-
tion of tracks generated with the `make_tracks_file` function. The bin size was set to 10 bp in
the region of interest to enable high-resolution visualization.

ATAC-seq data analysis

Publicly available ATAC-seq datasets were downloaded from GEO in `BigWig` format. For each
condition, the signal was averaged across the two biological replicates and visualized with
pyGenomeTracks. Binning was set to 10 bp within the region of interest.

Single-cell data integration

Single-cell transcriptomic datasets were obtained from human preimplantation embryos (day 3–
7, ArrayExpress), human gastrula (day 16, ArrayExpress), and human fetal hearts (week 5–25,
GEO). Raw counts were downloaded and only undifferentiated and mesendoderm-committed
cells (for early datasets) were retained. A unified count matrix was generated using the intersec-
tion of gene names across datasets. Cells with fewer than 200 expressed genes were excluded,
and the filtered matrix was imported into a Monocle3 `cell_dataset` object. Data were nor-
malized with `preprocess_cds` (PCA, log normalization, 50 dimensions), then batch-corrected
using `align_cds`. Dimensionality reduction was performed with UMAP (`reduce_dimension`,
`umap.min_dist = 1`), followed by clustering (`cluster_cells`), trajectory inference (`learn_graph`),
and pseudotime ordering (`order_cells`), rooting at the branch containing day 3 preimplantation
cells. Because trajectories showed reduced linearity, data were further subdivided by cham-
ber identity (RA, RV, LA, LV) based on the fetal heart dataset classification. Each subset was
reprocessed through the same Monocle3 pipeline. UMAPs and pseudotime trajectories were vi-
sualized with `plot_cells` and `plot_genes_in_pseudotime` to examine chamber-specific gene
expression dynamics.

Cardiac B-to-A gene selection

Cardiac B-to-A genes were identified from H9 cell Hi-C compartmentalization data, which pro-
vided the deepest sequencing coverage and the widest time-course (day 0 to day 80 of car-
diomyocyte differentiation). Bins located in the B compartment at day 0 and in the A com-
partment at day 80 were selected. Each bin was plotted based on the absolute difference
in principal component (PC) value between the two timepoints and the $\log_{10}$ of the adjusted
p-value from dcHiC analysis (as described above). Significantly switching bins were filtered
using a hyperbolic cutoff (y = 2x/(x − 2)), corresponding to an adjusted p-value $\leq$ 0.01 and a
PC difference $\geq$ 2. B-to-A genes were then defined as those with transcription start sites (TSS)
located within these bins. To assess expression dynamics, bulk RNA-seq data associated with
the H9 Hi-C dataset were analyzed. RPKM-normalized counts were downloaded from GEO

and converted to TPM values. TPM counts were batch-corrected across replicates using the `removeBatchEffect` function (limma). Genes were classified as early, mid, or late expressed based on the timepoint of peak z-score (averaged across replicates): day 0–2 (early), day 5–7 (mid), or day 15–80 (late). Gene set enrichment analysis (GSEA) was performed with `fgsea`[68], using the three temporal gene groups as pathways and genes ranked by t-statistics from differential expression analysis (tetracycline vs. isogenic control). Enrichment curves were plotted with `plotEnrichment` for late-expressed genes in each comparison.

Statistics

Statistical analyses were performed using GraphPad Prism v10. The number of biological replicates, statistical test applied, and exact p-values are reported in the figure legends. Unless otherwise specified, data are presented as mean $\pm$ SEM.

Key resources table

Reagent or Resource	Source	Identifier
Antibodies		
Mouse Anti-Cardiac Troponin T, Alexa Fluor 647 (clone 13-11)	BD Pharmingen	#565744
Rabbit anti-Titin (Z-disc, Z1Z2)	Myomedix	#TTN-1
Goat anti-Rabbit IgG, Alexa Fluor 488	Thermo Fisher Scientific	#A11008
Mouse anti-Ki67	BD Biosciences	#550609
Rabbit anti-α-actinin 2	Invitrogen	#701914
Mouse anti-pHH3	Santa Cruz Biotechnology	#sc-374669
Goat anti-Mouse IgG, Alexa Fluor 633	Thermo Fisher Scientific	#A21052
Goat anti-Mouse IgG, Alexa Fluor 568	Thermo Fisher Scientific	#A11004
Rabbit anti-Lamin B1	Invitrogen	#PA5-19468
Mouse anti-α-actinin	Abcam	Abcam #ab9465
Goat anti-Rabbit IgG, Alexa Fluor 405	Thermo Fisher Scientific	#A31556
Rabbit anti-CTCF	Active Motif	#V13-61932
Mouse anti-CTCF	BD Biosciences	#612148
Rabbit IgG Isotype Control	BD Biosciences	#550875
Mouse IgG1κ Isotype Control	BD Biosciences	#554121
Baterial and virus strains		
BAC clone CH17-275G10 (*TTN* probe)	BACPAC Resources Center	#CH17-275G10
Biological samples		
Human iPSC line WTC11 (male donor)	J. David Gladstone Institutes; Bruce Conklin	RRID: CVCL_Y803
WTC11 titin-mEGFP reporter line	Allen Institute for Cell Science; distributed via Coriell	RRID: CVCL_UD16
Chemicals, peptides, and recombinant proteins		
Geltrex, ESC-qualified	Gibco	#A1413302
Matrigel, hESC-qualified	Corning	#354277
Essential 8 Medium	Gibco	#A1517001
mTeSR Plus Medium	STEMCELL Technologies	#100-0276
Versene (0.5 mM EDTA in PBS)	Calbiochem	#4055100
Thiazovivin	Cayman Chemical	#004CA14245
CHIR99021	Cayman Chemical	#004CA13122
RPMI 1640	Gibco	#11875093
BSA	Sigma-Aldrich	#A8412-100ML
L-ascorbic acid 2-phosphate	Fujifilm	#321-44823
WNT-C59	Cayman Chemical	#16644
B-27 Supplement	Thermo Fisher Scientific	#17504044
IMDM	Gibco	#21980065
Ham's F-12 with GlutaMAX	Gibco	#31765068
Concentrated Lipids	Gibco	#11905031
Monothioglycerol	Sigma-Aldrich	#M6145-100ML
Optiferrin (human transferrin)	Invitria	#777TRF029-1G
Y-27632	Cayman Chemical	#004CA10005583-10
FGF2-G3	Qkine	#Qk035
LY294002	Selleckchem	#1105
Activin A	Qkine	#Qk001
BMP4	Qkine	#Qk038
Insulin	Thermo Fisher Scientific	#A11382II
All-trans Retinoic Acid	Sigma-Aldrich	#R2625
SB431542	Selleckchem	#S1067
XAV-939	Selleckchem	#S1180
Trypsin 0.25%	Euroclone	#ECB3051D
Fetal Bovine Serum (FBS)	Life Technologies	#10270106
Saponin	Sigma-Aldrich	#8047-15-2
Hoechst 33342	Thermo Fisher Scientific	#H1399
ProLong Glass Antifade Mountant	Invitrogen	#P36982
Fixable Viability Dye eFluor 780	Invitrogen	#65-0865-14
Fixable Viability Dye eFluor 450	Invitrogen	#65-0863-14
ProLong Diamond/Glass Mountants	Invitrogen	#P36982
Formamide	Fisher Scientific	#10050110
Cot-1 DNA (Human)	Thermo Fisher Scientific	#15279011

Reagent or Resource	Source	Identifier
Salmon Sperm DNA	Thermo Fisher Scientific	#AM9680
Alexa Fluor 488-5-dUTP	Thermo Fisher Scientific	#C11397
DpnII restriction enzyme	NEB	#R0543S
Csp6I restriction enzyme	Thermo Fisher Scientific	#ER0211
T4 DNA Ligase	Thermo Fisher Scientific	#EL0014
RNase A	Thermo Fisher Scientific	#R1253
Proteinase K	Thermo Fisher Scientific	#4333793
Protease inhibitor cocktail	Roche	#11697498001
Critical commercial assays		
TOPO™ TA Cloning™ Kit for Sequencing	Thermo Fisher Scientific	#450030
Click-iT EdU Alexa Fluor 647 Flow Cytometry Assay Kit	Invitrogen	#C10424
CellTrace Far Red Cell Proliferation Kit	Invitrogen	#C34564
Quick-RNA MiniPrep Kit	Zymo Research	#ZYR1055
High-Capacity cDNA Reverse Transcription Kit	Applied Biosystems	#4368813
PowerUp SYBR Green Master Mix (2X)	Applied Biosystems	#A25743
Luna Universal qPCR Master Mix	New England Biolabs (NEB)	#BM3003E
Pierce Protein A/G Magnetic Beads	Thermo Scientific	#88802
PyroMark PCR Kit	Qiagen	#978801
PyroMark Q48 Advanced CpG Reagents	Qiagen	#974022
Methylation-Lightning Kit	Zymo Research	#D5030
Zymo-Seq RiboFree Total RNA Library Prep Kit	Zymo Research	#R3000S
Zymo-See Index Primer Set	Zymo Research	#D3007
Illumina TruSeq Stranded mRNA Library Prep	Illumina	#20020594
IDT for Illumina – TruSeq RNA UD Indexes	Illumina	#20020591
Illumina Stranded mRNA Prep, Ligation	Illumina	#20040534
NEBNext High-Fidelity 2X PCR Mix	New England Biolabs (NEB)	#BM0541
LongAmp Taq DNA Polymerase	New England Biolabs (NEB)	#M0323
Illumina Dual Index Kit TT, Set A	Illumina	#PN-1000215
SPRIselect Beads	Beckman Coulter	#B23318
Nick Translation Kit	Roche	#10976776001
ChIP DNA Clean & Concentrator Kit	Zymo Research	#D5201
Deposited data		
RNA-seq data - 2D KD hiPSC-CMs	This paper; ArrayExpress	E-MTAB-15684
RNA-seq data - SHF KD cardioids	This paper; ArrayExpress	E-MTAB-15710
sc-RNA-seq data - SHF and FHF cardioids	This paper; ArrayExpress	E-MTAB-15693
sc-RNA-seq data - FHF KD cardioids	This paper; ArrayExpress	E-MTAB-15711
4C-seq data	This paper; ENA	PRJEB97970
Processed NGS data	This paper; Zenodo	https://doi.org/10.5281/-zenodo.17256128
Hi-C - RUES2 and WTC11 hPSC-CM time course	GEO	GSE106687
Hi-C -H9 hESC-CM time course	GEO	GSE116862
GATA4 ChIP-seq - hESC	GEO	GSE61475
GATA4 ChIP-seq - hiPSC-CM	GEO	GSE85628
CTCF ChIP-seq - hESCs	ENCODE	ENCFF764RHO; ENCFF332TNJ
CTCF ChIP-seq - hESCs-CMs	ENCODE	ENCFF777TNC; ENCFF466BIT
ATAC-seq - RUES2 hESC-CM time course	GEO	GSE106689
RNA-seq - RUES2 hPSC-CM time course	GEO	GSE106688
RNA-seq - H9 hESC-CMS time course	GEO	GSE203375
scRNA-seq - human heart	GEO	GSE106118
scRNA-seq - human preimpl. embryo	ArrayExpress	E-MTAB-3929
scRNA-seq - Human gastrula	ArrayExpress	E-MTAB-9388

Reagent or Resource	Source	Identifier
Oligonucleotides		
Primers and shRNA sequences	This paper	Tables S1–S4
Recombinant DNA		
pAAV-Puro_siKD	Addgene	#86695
pZFN-AAVS1_ELD	Addgene	#159297
pZFN-AAVS1_KKR	Addgene	#159298
pX330-U6-Chimeric_BB-CBh-hSpCas9	Addgene	#42230
AAVS1 SA-2A-puro-pA donor	Addgene	#22075
Software and algorithms		
Code repository	This paper; GitHub	https://github.com/alessandro-bertero/Becca_et_al_2025
Single-cell analysis environment	Docker Hub (hedgelab)	Docker image: hedge-lab/rstudio_image:iPS2seq
Hi-C analysis environment	Docker Hub (hedgelab)	Docker image: hedge-lab/hic_pipeline:image12
Bulk RNA-seq analysis environment	Docker Hub (hedgelab)	Docker image: hedge-lab/bulk_image:image3
Cell Ranger environment	Docker Hub (hedgelab)	Docker image: hedge-lab/cellranger8:latest
4C-seq analysis environment	Docker Hub (hedgelab)	Docker image: hedge-lab/4c:image1
IGV (Integrative Genomics Viewer)	Broad Institute	https://software.broadinstitute.org/software/igv/
FIMO (MEME Suite)	MEME Suite	http://meme-suite.org/
CTCFBSDB 2.0	University of Massachusetts	http://insulatordb.uthsc.edu/
FlowJo v10.10.0	BD Biosciences	https://www.flowjo.com/
Imaris v10.2	Oxford Instruments	https://imaris.oxinst.com/
HiC-Pro v3.1.0	Institut Curie	https://github.com/nservant/HiC-Pro
dcHiC	GitHub	https://github.com/ay-lab/dcHiC
HiCExplorer v3.4.1	HiCExplorer	https://hicexplorer.readthedocs.io/
HiContacts / HiCExperiment (R packages)	Bioconductor	https://bioconductor.org/
pyGenomeTracks	GitHub	https://github.com/deeptools/pyGenomeTracks
UCSC LiftOver	UCSC Genome Browser	https://genome.ucsc.edu/
bowtie2	Johns Hopkins University	http://bowtie-bio.sourceforge.net/bowtie2/
STAR v2.7.11b	GitHub	https://github.com/alexdobin/STAR
trim-galore v0.6.10	Babraham Institute	https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/
bcl2fastq	Illumina	https://support.illumina.com/
cellranger v8.0.1	10x Genomics	https://www.10xgenomics.com/
Seurat	Satija Lab	https://satijalab.org/seurat/
monocle3	Trapnell Lab	https://cole-trapnell-lab.github.io/monocle3/
limma	Bioconductor	https://bioconductor.org/packages/limma/
clusterProfiler	Bioconductor	https://bioconductor.org/packages/clusterProfiler/
edgeR	Bioconductor	https://bioconductor.org/packages/edgeR/
miloR	Bioconductor	https://bioconductor.org/packages/release/bioc/html/miloR.html
RNAseqQC	GitHub	https://github.com/getzlab/rnaseq_qc

Reagent or Resource	Source	Identifier
Software and algorithms		
ADImpute	Bioconductor	https://bioconductor.org/-packages/ADImpute/
gtftools	GitHub	https://github.com/zyndagj/-gtftools
WGCNA	UCLA / Horvath Lab	https://horvath.genetics.-ucla.edu/html/Coexpress-ionNetwork/Rpackages/WGCNA/
fgsea	Bioconductor	https://bioconductor.org/-packages/fgsea/
RRHO2	GitHub	https://github.com/Caleb-Huo/RRHO2
FlowJo v10.10	BD	Commercial software
GraphPad Prism v10	GraphPad	Commercial software
Imaris v10.2	Oxford Instruments	Commercial software
PyroMark Q48 Autoprep Software	QIAGEN	Commercial software
PyroMark Assay Design Software 2.0	QIAGEN	Commercial software
Other		
FACSVerse Flow Cytometer	BD Biosciences	Instrument
Sony SH800S Cell Sorter	Sony Biotechnology	Instrument
Leica SP8 Confocal Microscope (63x)	Leica Microsystems	Instrument
Zeiss Elyra 7 Microscope	Zeiss	Instrument
Incucyte SX5 Live-Cell Analysis System	Sartorius	Instrument
Leica CM1950 cryostat	Leica Microsystems	Instrument
Bioruptor Pico sonication device	Diagenode	Instrument
PyroMark Q48 Autoprep	Qiagen	Instrument
QuantStudio 6 Flex Real-Time PCR System	Applied Biosystems	Instrument
Chromium X Controller + Next GEM Chip G	10x Genomics	Instrument
Illumina NovaSeq 6000	Illumina	Instrument
Illumina NextSeq 1000/550	Illumina	Instrument
MGI T7 sequencer	MGI Tech	Instrument
Molecular Karyotyping	Life & Brain GmBH	Service
Sanger sequencing service	Eurofins Genomics	Service

References

1. Anania, C. and Lupiáñez, D. G. (2020). Order and Disorder: Abnormal 3D Chromatin Organization in Human Disease. Briefings in Functional Genomics *19*, 128–138. https://doi.org/10.1093/bfgp/elz028.

2. Dekker, J., Alber, F., Aufmkolk, S., Beliveau, B. J., Bruneau, B. G., Belmont, A. S., Bintu, L., Boettiger, A., Calandrelli, R., Disteche, C. M., et al. (2023). Spatial and Temporal Organization of the Genome: Current State and Future Aims of the 4D Nucleome Project. Molecular Cell *83*, 2624–2640. https://doi.org/10.1016/j.molcel.2023.06.018.

3. Meshorer, E., Yellajoshula, D., George, E., Scambler, P. J., Brown, D. T., and Misteli, T. (2006). Hyperdynamic Plasticity of Chromatin Proteins in Pluripotent Embryonic Stem Cells. Developmental cell *10*, 105–16. https://doi.org/10.1016/j.devcel.2005.10.017.

4. Paige, S. L., Thomas, S., Stoick-Cooper, C. L., Wang, H., Maves, L., Sandstrom, R., Pabon, L., Reinecke, H., Pratt, G., Keller, G., et al. (2012). A Temporal Chromatin Signature in Human Embryonic Stem Cells Identifies Regulators of Cardiac Development. Cell *151*, 221–232. https://doi.org/10.1016/j.cell.2012.08.027.

5. Dixon, J. R., Jung, I., Selvaraj, S., Shen, Y., Antosiewicz-Bourget, J. E., Lee, A. Y., Ye, Z., Kim, A., Rajagopal, N., Xie, W., et al. (2015). Chromatin Architecture Reorganization during Stem Cell Differentiation. Nature *518*, 331–336. https://doi.org/10.1038/nature14222.

6. Lieberman-Aiden, E., van Berkum, N. L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B. R., Sabo, P. J., Dorschner, M. O., et al. (2009). Comprehensive Mapping of Long Range Interactions Reveals Folding Principles of the Human Genome. Science (New York, N.Y.) *326*, 289–293. https://doi.org/10.1126/science.1181369.

7. Bertero, A. and Rosa-Garrido, M. (2021). Three-Dimensional Chromatin Organization in Cardiac Development and Disease. Journal of Molecular and Cellular Cardiology *151*, 89–105. https://doi.org/10.1016/j.yjmcc.2020.11.008.

8. Shah, P. P., Lv, W., Rhoades, J. H., Poleshko, A., Abbey, D., Caporizzo, M. A., Linares-Saldana, R., Heffler, J. G., Sayed, N., Thomas, D., et al. (2021). Pathogenic LMNA Variants Disrupt Cardiac Lamina-Chromatin Interactions and de-Repress Alternative Fate Genes. Cell Stem Cell *28*, 938–954.e9. https://doi.org/10.1016/j.stem.2020.12.016.

9. Bertero, A., Fields, P. A., Smith, A. S., Leonard, A., Beussman, K., Sniadecki, N. J., Kim, D.-H., Tse, H.-F., Pabon, L., Shendure, J., et al. (2019a). Chromatin Compartment Dynamics in a Haploinsufficient Model of Cardiac Laminopathy. Journal of Cell Biology *218*, 2919–2944. https://doi.org/10.1083/jcb.201902117.

10. Zaidi, S., Choi, M., Wakimoto, H., Ma, L., Jiang, J., Overton, J. D., Romano-Adesman, A., Bjornson, R. D., Breitbart, R. E., Brown, K. K., et al. (2013). De Novo Mutations in Histone-Modifying Genes in Congenital Heart Disease. Nature *498*, 220–223. https://doi.org/10.1038/nature12141.

11. Salvarani, N., Crasto, S., Miragoli, M., Bertero, A., Paulis, M., Kunderfranco, P., Serio, S., Forni, A., Lucarelli, C., Dal Ferro, M., et al. (2019). The K219T-Lamin Mutation Induces

Conduction Defects through Epigenetic Inhibition of SCN5A in Human Cardiac Laminopathy. Nature Communications *10*, 2267. https://doi.org/10.1038/s41467-019-09929-w.

12. Poleshko, A., Shah, P. P., Gupta, M., Babu, A., Morley, M. P., Manderfield, L. J., Ifkovits, J. L., Calderon, D., Aghajanian, H., Sierra-Pagán, J. E., et al. (2017). Genome-Nuclear Lamina Interactions Regulate Cardiac Stem Cell Lineage Restriction. Cell *171*, 573–587.e14. https://doi.org/10.1016/j.cell.2017.09.018.

13. Bertero, A., Fields, P. A., Ramani, V., Bonora, G., Yardimci, G. G., Reinecke, H., Pabon, L., Noble, W. S., Shendure, J., and Murry, C. E. (2019b). Dynamics of Genome Reorganization during Human Cardiogenesis Reveal an RBM20-dependent Splicing Factory. Nature Communications *10*, 1538. https://doi.org/10.1038/s41467-019-09483-5.

14. Grant, Z. L., Kuang, S., Zhang, S., Horrillo, A. J., Rao, K. S., Kameswaran, V., Joubran, C., Lau, P. K., Dong, K., Yang, B., et al. (2025). *Dose-Dependent Sensitivity of Human 3D Chromatin to a Heart Disease-Linked Transcription Factor*. https://doi.org/10.1101/2025.01.09.632202.

15. Iwafuchi-Doi, M. and Zaret, K. S. (2014). Pioneer Transcription Factors in Cell Reprogramming. Genes & Development *28*, 2679–2692. https://doi.org/10.1101/gad.253443.114.

16. Molkentin, J. D., Lin, Q., Duncan, S. A., and Olson, E. N. (1997). Requirement of the Transcription Factor GATA4 for Heart Tube Formation and Ventral Morphogenesis. Genes & Development *11*, 1061–1072. https://doi.org/10.1101/gad.11.8.1061.

17. Ang, Y. S., Rivas, R. N., Ribeiro, A. J., Srivas, R., Rivera, J., Stone, N. R., Pratt, K., Mohamed, T. M., Fu, J. D., Spencer, C. I., et al. (2016). Disease Model of GATA4 Mutation Reveals Transcription Factor Cooperativity in Human Cardiogenesis. Cell *167*, 1734–1749.e22. https://doi.org/10.1016/j.cell.2016.11.033.

18. Garg, V., Kathiriya, I. S., Barnes, R., Schluterman, M. K., King, I. N., Butler, C. A., Rothrock, C. R., Eapen, R. S., Hirayama-Yamada, K., Joo, K., et al. (2003). GATA4 Mutations Cause Human Congenital Heart Defects and Reveal an Interaction with TBX5. Nature *424*, 443–447. https://doi.org/10.1038/nature01827.

19. Rajagopal, S. K., Ma, Q., Obler, D., Shen, J., Manichaikul, A., Tomita-Mitchell, A., Boardman, K., Briggs, C., Garg, V., Srivastava, D., et al. (2007). Spectrum of Heart Disease Associated with Murine and Human GATA4 Mutation. Journal of Molecular and Cellular Cardiology *43*, 677–685. https://doi.org/10.1016/j.yjmcc.2007.06.004.

20. Ong, C.-T. and Corces, V. G. (2014). CTCF: An Architectural Protein Bridging Genome Topology and Function. Nature Reviews Genetics *15*, 234–246. https://doi.org/10.1038/nrg3663.

21. Li, Y., Haarhuis, J. H. I., Sedeño Cacciatore, Á., Oldenkamp, R., van Ruiten, M. S., Willems, L., Teunissen, H., Muir, K. W., de Wit, E., Rowland, B. D., et al. (2020). The Structural Basis for Cohesin–CTCF-anchored Loops. Nature *578*, 472–476. https://doi.org/10.1038/s41586-019-1910-z.

22. Dixon, J. R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J. S., and Ren, B. (2012). Topological Domains in Mammalian Genomes Identified by Analysis of Chromatin Interactions. Nature *485*, 376–380. https://doi.org/10.1038/nature11082.

23. Narendra, V., Bulajić, M., Dekker, J., Mazzoni, E. O., and Reinberg, D. (2016). CTCF-mediated Topological Boundaries during Development Foster Appropriate Gene Regulation. Genes & Development *30*, 2657–2662. https://doi.org/10.1101/gad.288324.116.

24. Liu, Y., Wan, X., Li, H., Chen, Y., Hu, X., Chen, H., Zhu, D., Li, C., and Zhang, Y. (2023). CTCF Coordinates Cell Fate Specification via Orchestrating Regulatory Hubs with Pioneer Transcription Factors. Cell Reports *42*, 113259. https://doi.org/10.1016/j.celrep.2023.113259.

25. Bastaki, F., Nair, P., Mohamed, M., Malik, E. M., Helmi, M., Al-Ali, M. T., and Hamzeh, A. R. (2017). Identification of a Novel CTCF Mutation Responsible for Syndromic Intellectual Disability - a Case Report. BMC medical genetics *18*, 68. https://doi.org/10.1186/s12881-017-0429-0.

26. Gregor, A., Oti, M., Kouwenhoven, E. N., Hoyer, J., Sticht, H., Ekici, A. B., Kjaergaard, S., Rauch, A., Stunnenberg, H. G., Uebe, S., et al. (2013). De Novo Mutations in the Genome Organizer CTCF Cause Intellectual Disability. American Journal of Human Genetics *93*, 124–131. https://doi.org/10.1016/j.ajhg.2013.05.007.

27. Schmidt, C., Deyett, A., Ilmer, T., Haendeler, S., Torres Caballero, A., Novatchkova, M., Netzer, M. A., Ceci Ginistrelli, L., Mancheno Juncosa, E., Bhattacharya, T., et al. (2023). Multi-Chamber Cardioids Unravel Human Heart Development and Cardiac Defects. Cell *186*, 5587–5605.e27. https://doi.org/10.1016/j.cell.2023.10.030.

28. Zhang, Y., Li, T., Preissl, S., Amaral, M. L., Grinstein, J. D., Farah, E. N., Destici, E., Qiu, Y., Hu, R., Lee, A. Y., et al. (2019). Transcriptionally Active HERV-H Retrotransposons Demarcate Topologically Associating Domains in Human Pluripotent Stem Cells. Nature Genetics *51*, 1380–1388. https://doi.org/10.1038/s41588-019-0479-7.

29. ENCODE Project Consortium (2012). An Integrated Encyclopedia of DNA Elements in the Human Genome. Nature *489*, 57–74. https://doi.org/10.1038/nature11247.

30. Tsankov, A. M., Gu, H., Akopian, V., Ziller, M. J., Donaghey, J., Amit, I., Gnirke, A., and Meissner, A. (2015). Transcription Factor Binding Dynamics during Human ES Cell Differentiation. Nature *518*, 344–349. https://doi.org/10.1038/nature14233.

31. Tyser, R. C. V., Mahammadov, E., Nakanoh, S., Vallier, L., Scialdone, A., and Srinivas, S. (2021). Single-Cell Transcriptomic Characterization of a Gastrulating Human Embryo. Nature *600*, 285–289. https://doi.org/10.1038/s41586-021-04158-y.

32. Petropoulos, S., Edsgärd, D., Reinius, B., Deng, Q., Panula, S. P., Codeluppi, S., Plaza Reyes, A., Linnarsson, S., Sandberg, R., and Lanner, F. (2016). Single-Cell RNA-Seq Reveals Lineage and X Chromosome Dynamics in Human Preimplantation Embryos. Cell *165*, 1012–1026. https://doi.org/10.1016/j.cell.2016.03.023.

33. Cui, Y., Zheng, Y., Liu, X., Yan, L., Fan, X., Yong, J., Hu, Y., Dong, J., Li, Q., Wu, X., et al. (2019). Single-Cell Transcriptome Analysis Maps the Developmental Track of the Human Heart. Cell Reports *26*, 1934–1950.e5. https://doi.org/10.1016/j.celrep.2019.01.079.

34. Dark, N., Cosson, M.-V., Tsansizi, L. I., Owen, T. J., Ferraro, E., Francis, A. J., Tsai, S., Bouissou, C., Weston, A., Collinson, L., et al. (2023). Generation of Left Ventricle-like Cardiomyocytes with Improved Structural, Functional, and Metabolic Maturity from Human Pluripotent Stem Cells. Cell Reports Methods *3*, 100456. https://doi.org/10.1016/j.crmeth.2023.100456.

35. Brancaccio, M., Guazzone, S., Menini, N., Sibona, E., Hirsch, E., De Andrea, M., Rocchi, M., Altruda, F., Tarone, G., and Silengo, L. (1999). Melusin Is a New Muscle-Specific Interactor for Beta(1) Integrin Cytoplasmic Domain. The Journal of Biological Chemistry *274*, 29282–29288. https://doi.org/10.1074/jbc.274.41.29282.

36. de Wit, E., Vos, E. S. M., Holwerda, S. J. B., Valdes-Quezada, C., Verstegen, M. J. A. M., Teunissen, H., Splinter, E., Wijchers, P. J., Krijger, P. H. L., and de Laat, W. (2015). CTCF Binding Polarity Determines Chromatin Looping. Molecular cell *60*, 676–84. https://doi.org/10.1016/j.molcel.2015.09.023.

37. Rojas, A., De Val, S., Heidt, A. B., Xu, S.-M., Bristow, J., and Black, B. L. (2005). Gata4 Expression in Lateral Mesoderm Is Downstream of BMP4 and Is Activated Directly by Forkhead and GATA Transcription Factors through a Distal Enhancer Element. Development *132*, 3405–3417. https://doi.org/10.1242/dev.01913.

38. Hofbauer, P., Jahnel, S. M., Papai, N., Giesshammer, M., Deyett, A., Schmidt, C., Penc, M., Tavernini, K., Grdseloff, N., Meledeth, C., et al. (2021). Cardioids Reveal Self-Organizing Principles of Human Cardiogenesis. Cell *184*, 3299–3317.e22. https://doi.org/10.1016/j.cell.2021.04.034.

39. Bertero, A., Pawlowski, M., Ortmann, D., Snijders, K., Yiangou, L., Cardoso de Brito, M., Brown, S., Bernard, W. G., Cooper, J. D., Giacomelli, E., et al. (2016). Optimized Inducible shRNA and CRISPR/Cas9 Platforms for in Vitro Studies of Human Development Using hPSCs. Development (Cambridge, England) *143*, 4405–4418. https://doi.org/10.1242/dev.138081.

40. Mitzelfelt, K. A., McDermott-Roe, C., Grzybowski, M. N., Marquez, M., Kuo, C.-T., Riedel, M., Lai, S., Choi, M. J., Kolander, K. D., Helbling, D., et al. (2017). Efficient Precision Genome Editing in iPSCs via Genetic Co-targeting with Selection. Stem Cell Reports *8*, 491–499. https://doi.org/10.1016/j.stemcr.2017.01.021.

41. Barutcu, A. R., Maass, P. G., Lewandowski, J. P., Weiner, C. L., and Rinn, J. L. (2018). A TAD Boundary Is Preserved upon Deletion of the CTCF-rich Firre Locus. Nature Communications *9*, 1444. https://doi.org/10.1038/s41467-018-03614-0.

42. Kang, J., Kim, Y. W., Park, S., Kang, Y., and Kim, A. (2021). Multiple CTCF Sites Cooperate with Each Other to Maintain a TAD for Enhancer–Promoter Interaction in the β-Globin Locus. The FASEB Journal *35*, e21768. https://doi.org/10.1096/fj.202100105RR.

43. Plaisier, S. B., Taschereau, R., Wong, J. A., and Graeber, T. G. (2010). Rank–Rank Hypergeometric Overlap: Identification of Statistically Significant Overlap between Gene-

Expression Signatures. Nucleic Acids Research *38*, e169. https://doi.org/10.1093/nar/gkq636.

44. Dann, E., Henderson, N. C., Teichmann, S. A., Morgan, M. D., and Marioni, J. C. (2022). Differential Abundance Testing on Single-Cell Data Using k-Nearest Neighbor Graphs. Nature Biotechnology *40*, 245–253. https://doi.org/10.1038/s41587-021-01033-z.

45. Zuin, J., Dixon, J. R., van der Reijden, M. I. J. A., Ye, Z., Kolovos, P., Brouwer, R. W. W., van de Corput, M. P. C., van de Werken, H. J. G., Knoch, T. A., van IJcken, W. F. J., et al. (2014). Cohesin and CTCF Differentially Affect Chromatin Architecture and Gene Expression in Human Cells. Proceedings of the National Academy of Sciences *111*, 996–1001. https://doi.org/10.1073/pnas.1317788111.

46. Hyle, J., Zhang, Y., Wright, S., Xu, B., Shao, Y., Easton, J., Tian, L., Feng, R., Xu, P., and Li, C. (2019). Acute Depletion of CTCF Directly Affects MYC Regulation through Loss of Enhancer–Promoter Looping. Nucleic Acids Research *47*, 6699–6713. https://doi.org/10.1093/nar/gkz462.

47. Ciceri, G., Baggiolini, A., Cho, H. S., Kshirsagar, M., Benito-Kwiecinski, S., Walsh, R. M., Aromolaran, K. A., Gonzalez-Hernandez, A. J., Munguba, H., Koo, S. Y., et al. (2024). An Epigenetic Barrier Sets the Timing of Human Neuronal Maturation. Nature *626*, 881–890. https://doi.org/10.1038/s41586-023-06984-8.

48. Sepulveda, J. L., Belaguli, N., Nigam, V., Chen, C.-Y., Nemer, M., and Schwartz, R. J. (1998). GATA-4 and Nkx-2.5 Coactivate Nkx-2 DNA Binding Targets: Role for Regulating Early Cardiac Gene Expression. Molecular and Cellular Biology *18*, 3405–3415.

49. Sepulveda, J. L., Vlahopoulos, S., Iyer, D., Belaguli, N., and Schwartz, R. J. (2002). Combinatorial Expression of GATA4, Nkx2-5, and Serum Response Factor Directs Early Cardiac Gene Activity*. Journal of Biological Chemistry *277*, 25775–25782. https://doi.org/10.1074/jbc.M203122200.

50. Gao, H., Pathan, S., Dixon, B. R. E. A., Pugazenthi, A., Mathison, M., Mohamed, T. M. A., Rosengart, T. K., and Yang, J. (2024). Sall4 and Gata4 Induce Cardiac Fibroblast Transition towards a Partially Multipotent State with Cardiogenic Potential. Scientific Reports *14*, 24182. https://doi.org/10.1038/s41598-024-73975-8.

51. Misra, C., Sachan, N., McNally, C. R., Koenig, S. N., Nichols, H. A., Guggilam, A., Lucchesi, P. A., Pu, W. T., Srivastava, D., and Garg, V. (2012). Congenital Heart Disease-Causing Gata4 Mutation Displays Functional Deficits in Vivo. PLoS Genetics *8*, e1002690. https://doi.org/10.1371/journal.pgen.1002690.

52. Gomez-Velazquez, M., Badia-Careaga, C., Lechuga-Vieco, A. V., Nieto-Arellano, R., Tena, J. J., Rollan, I., Alvarez, A., Torroja, C., Caceres, E. F., Roy, A. R., et al. (2017). CTCF Counter-Regulates Cardiomyocyte Development and Maturation Programs in the Embryonic Heart. PLoS genetics *13*, e1006985. https://doi.org/10.1371/journal.pgen.1006985.

53. Ren, H., Zhong, H., Zhang, J., Lu, Y., Hu, G., Duan, W., Ma, N., and Yao, H. (2024). CTCF Point Mutation at R567 Disrupts Mouse Heart Development via 3D Genome Rearrangement and Transcription Dysregulation. Cell Proliferation *58*, e13783. https://doi.org/10.1111/cpr.13783.

54. Pehlivan, T., Pober, B. R., Brueckner, M., Garrett, S., Slaugh, R., Van Rheeden, R., Wilson, D. B., Watson, M. S., and Hing, A. V. (1999). GATA4 Haploinsufficiency in Patients with Interstitial Deletion of Chromosome Region 8p23.1 and Congenital Heart Disease. American Journal of Medical Genetics *83*, 201–206.

55. Kreitzer, F. R., Salomonis, N., Sheehan, A., Huang, M., Park, J. S., Spindler, M. J., Lizarraga, P., Weiss, W. A., So, P.-L., and Conklin, B. R. (2013). A Robust Method to Derive Functional Neural Crest Cells from Human Pluripotent Stem Cells. American Journal of Stem Cells *2*, 119–131.

56. Roberts, B., Hendershott, M. C., Arakaki, J., Gerbin, K. A., Malik, H., Nelson, A., Gehring, J., Hookway, C., Ludmann, S. A., Yang, R., et al. (2019). Fluorescent Gene Tagging of Transcriptionally Silent Genes in hiPSCs. Stem Cell Reports *12*, 1145–1158. https://doi.org/10.1016/j.stemcr.2019.03.001.

57. Bertero, A., Yiangou, L., Brown, S., Ortmann, D., Pawlowski, M., and Vallier, L. (2018). Conditional Manipulation of Gene Function in Human Cells with Optimized Inducible shRNA. Current Protocols in Stem Cell Biology *44*. https://doi.org/10.1002/cpsc.45.

58. Law, C. W., Alhamdoosh, M., Su, S., Dong, X., Tian, L., Smyth, G. K., and Ritchie, M. E. (2016). RNA-seq Analysis Is Easy as 1-2-3 with Limma, Glimma and edgeR. F1000Research *5*, ISCB Comm J–1408. https://doi.org/10.12688/f1000research.9005.3.

59. Langfelder, P. and Horvath, S. (2008). WGCNA: An R Package for Weighted Correlation Network Analysis. BMC Bioinformatics *9*, 559. https://doi.org/10.1186/1471-2105-9-559.

60. Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: An R Package for Comparing Biological Themes Among Gene Clusters. OMICS : a Journal of Integrative Biology *16*, 284–287. https://doi.org/10.1089/omi.2011.0118.

61. Bertero, A., Madrigal, P., Galli, A., Hubner, N. C., Moreno, I., Burks, D., Brown, S., Pedersen, R. A., Gaffney, D., Mendjan, S., et al. (2015). Activin/Nodal Signaling and NANOG Orchestrate Human Embryonic Stem Cell Fate Decisions by Controlling the H3K4me3 Chromatin Mark. Genes & Development *29*, 702–717. https://doi.org/10.1101/gad.255984.114.

62. Krijger, P. H., Geeven, G., Bianchi, V., Hilvering, C. R., and de Laat, W. (2019). 4C-seq from Start to End: A Detailed Protocol for Sample Preparation and Data Analysis. Methods, epub July 26. https://doi.org/10.1016/j.ymeth.2019.07.014.

63. Lopez-Delisle, L., Rabbani, L., Wolff, J., Bhardwaj, V., Backofen, R., Grüning, B., Ramírez, F., and Manke, T. (2021). pyGenomeTracks: Reproducible Plots for Multivariate Genomic Datasets. Bioinformatics (Oxford, England) *37*, 422–423. https://doi.org/10.1093/bioinformatics/btaa692.

64. Servant, N., Varoquaux, N., Lajoie, B. R., Viara, E., Chen, C.-J., Vert, J.-P., Heard, E., Dekker, J., and Barillot, E. (2015). HiC-Pro: An Optimized and Flexible Pipeline for Hi-C Data Processing. Genome Biology *16*, 259. https://doi.org/10.1186/s13059-015-0831-x.

65. Chakraborty, A., Wang, J. G., and Ay, F. (2022). dcHiC Detects Differential Compartments
across Multiple Hi-C Datasets. Nature Communications *13*, 6827. https://doi.org/10.1038/
s41467-022-34626-6.

66. Wolff, J., Rabbani, L., Gilsbach, R., Richard, G., Manke, T., Backofen, R., and Grüning,
B. A. (2020). Galaxy HiCExplorer 3: A Web Server for Reproducible Hi-C, Capture Hi-
C and Single-Cell Hi-C Data Analysis, Quality Control and Visualization. Nucleic Acids
Research *48*, W177–W184. https://doi.org/10.1093/nar/gkaa220.

67. Serizay, J., Matthey-Doret, C., Bignaud, A., Baudry, L., and Koszul, R. (2024). Orches-
trating Chromosome Conformation Capture Analysis with Bioconductor. Nature Commu-
nications *15*, 1072. https://doi.org/10.1038/s41467-024-44761-x.

68. Wiebe, D. S., Omelyanchuk, N. A., Mukhin, A. M., Grosse, I., Lashin, S. A., Zemlyanskaya,
E. V., and Mironova, V. V. (2020). Fold-Change-Specific Enrichment Analysis (FSEA):
Quantification of Transcriptional Response Magnitude for Functional Gene Groups. Genes
11, 434. https://doi.org/10.3390/genes11040434.