A regulatory loop regulating signaling between LRRK2 and PP2A in cellular models of Parkinson's disease

Panagiotis S. Athanasopoulos[1], Anna Memou[2], Franz Y. Ho[1], Ahmed Soliman[1,3], Henderikus Pots[1], Vassiliki Papadopoulou[2], Felix von Zweydorf[4,5], Saiganesh Sriraman[1], Alexander Marc Thouin[1], Laurine Vandewynckel[6], William Sibran[6], Marie-Christine Chartier-Harlin[6], R. Jeremy Nichols[7], Elisa Greggio[8,9], Jean-Marc Taymans[6], Christian Johannes Gloeckner[4,5], Hardy J. Rideout[2*], and Arjan Kortholt[1,10*]

[1]Department of Cell Biochemistry, University of Groningen, Nijenborgh 7, 9747 AG, Groningen, The Netherlands

[2]Center for Clinical, Experimental Surgery, and Translational Research; Biomedical Research Foundation of the Academy of Athens; Athens, Greece

[3]Division of Engineering in Medicine, Department of Medicine, Brigham Women's Hospital, Harvard Medical School, Boston, MA 02115, USA

[4]German Center for Neurodegenerative Diseases, 72076 Tübingen, Germany

[5]Institute for Ophthalmic Research, Center for Ophthalmology, University of Tübingen, Tübingen, Germany

[6]Univ. Lille, Inserm, CHU Lille, UMR-S 1172 - LilNCog - Lille Neuroscience & Cognition, F-59000 Lille, France.

[7] Department of Structural Biology, St Jud Children's Research Hospital, Memphis TN, USA

[8]University of Padova, Department of Biology, Padova, Italy

[9]Centro Studi per la Neurodegenerazione (CESNE), University of Padova, Italy

[10]YETEM-Innovative Technologies Application and Research Centre Suleyman Demirel University West Campus, Isparta, Turkey

*Corresponding authors:

Arjan Kortholt, Nijenborgh 7, 9747 AG, Groningen, The Netherlands, a.kortholt@rug.nl

Hardy J. Rideout, BRFAA, Soranou Efessiou 4, 115 27, Athens, Greece, hrideout@bioacademy.gr

Key words: LRRK2, PP2A, Parkinson's disease, phosphatase, kinase

Abstract

Mutations in Leucine-rich repeat kinase 2 (LRRK2) are the most frequent cause of late-onset familial and idiopathic Parkinson's disease (PD), known to date. Importantly, recent data from post-mortem tissue as well as biomarker studies suggest that independent of mutations, increased kinase activity of LRRK2 plays an essential role in idiopathic PD pathogenesis. Despite extensive research on LRRK2, its activation mechanism(s) and how the various mutations result in increased kinase activity and neuronal death is still not completely understood. Accumulating evidence points to LRRK2 phospho-regulation, both auto-phosphorylation and phosphorylation by other kinases, as one potential molecular trigger of its activation. LRRK2 activation and localization is regulated by phosphatases such as Protein phosphatase 1 (PP1) and Protein phosphatase 2A (PP2A), however the exact mechanism of this phospho-regulation is not known. Our data reveal that PP2A dephosphorylates sites within the RocCOR-GTPase domain of LRRK2 and as a result de-stabilizes LRRK2 dimers, with consequent reduction of its kinase activity. Strikingly, our data further highlight that LRRK2 in turn phosphorylates the catalytic subunit of the PP2A holoenzyme PPP2CA at its critical residue T304, crucial for both holoenzyme formation and catalytic activity. Importantly, expression of WT-PPP2CA protects from LRRK2-G2019S induced neuronal cell death, while PPP2CA-T304 mutants fail to do so, suggesting that impaired PP2A holoenzyme formation might be detrimental for LRRK2-PD.

Running title: A LRRK2 and PP2A feedback loop

Significance statement

Increased kinase activity of LRRK2 is a hallmark of Parkinson's disease. Accumulating evidence points to LRRK2 phospho-regulation as one potential molecular trigger of its activation. However, the exact regulation of phosphorylation was not well understood. Our data reveal a feedback loop between LRRK2 and PP2A (a phosphatase) activity, which explains the neuroprotective properties of PP2A's enzymatic activity in LRRK2-induced parkinsonism. Interestingly, PP2A activity is downregulated in alpha-synuclein PD models, having the same net-effect on PP2A activity as with hyper-active mutant LRRK2, and the methylation status of the C-terminal of PPP2CA plays an essential role in this regulation and progression of the disease. Understanding the regulation of PP2A, by post-translational modifications such as phosphorylation and methylation, can thus give important insight into the onset and progression of PD in general.

Introduction

PD is the second most common neurodegenerative disease after Alzheimer's disease, affecting 1–2% of the Western world's population older than 60 years (1). From a pathological point of view, it is characterized by the progressive death of dopaminergic neurons in the ventral midbrain associated with the formation of fibrillar aggregates that are α-synuclein (αSyn) enriched in most forms of PD. Mutations in leucine-rich-repeat kinase 2 (LRRK2) have been identified to be the main cause of late-onset hereditary Parkinson's disease (PD) (2). From a clinical perspective, it is largely indistinguishable from the more common idiopathic PD; however, there are some neuropathological differences. LRRK2 is a large multidomain protein consisting of two enzymatic domains, namely a RocCOR GTPase domain, belonging to the Roco protein family of G proteins, which is followed by a Serine/Threonine kinase domain (3–7). Several pathogenic point mutations located in the GTPase domain (such as N1437H, R1441C/G/H/S) and in the kinase domain (G2019S, I2020T), as well as the COR domain (Y1699C), all lead to increased kinase activity and subsequently play an important role in PD causality (8–10). Moreover, LRRK2 kinase activity is not only increased in LRRK2 familial PD, but also in idiopathic patients in the absence of a mutation (11) suggesting that LRRK2 (dys)function plays a broader role in PD. Therefore, the regulation of LRRK2 kinase activity is considered a key therapeutic target in familial and sporadic PD.

Despite a vast amount of research and the identification of several LRRK2 substrates and interaction partners, the exact cellular function of LRRK2, the underlying mechanism that results in increased LRRK2 activity and how this subsequently results in the progression of the disease is still not completely understood (12). Importantly, Rab proteins have been identified as both upstream regulators and as key physiological substrates of LRRK2 (13–15). Accumulating evidence suggests that this LRRK2/Rab pathway functions at the interface of vesicular trafficking and autophagy. LRRK2 is able to phosphorylate a subset of Rab GTPases within their Switch-II motif thereby controlling interaction with effectors and cellular trafficking. The PPM1H phosphatase is able to dephosphorylate this motif in the substrate Rabs (16, 17) thereby counteracting LRRK2 signaling.

Phospho-regulation plays an important role in the LRRK2 activation cycle (18–21). The N-terminal LRRK2 phosphorylation cluster (S910, S935, S955, S973) regulates binding to 14-3-3, which is important for LRRK2 localization, activity and neurotoxicity (21–28). Furthermore, phosphomutant studies at the heterologous phosphosites of LRRK2 (around S910 and S935) showed that phosphodead LRRK2 displays hyperactivatable kinase activity while phosphomimick LRRK2 shows reduced kinase activity markers, pointing to a role in LRRK2 heterophosphosites in regulating LRRK2 kinase activation (21), however so far, the role of LRRK2 autophosphorylation sites in LRRK2 kinase activation and toxicity is unknown. Phosphorylation of LRRK2 is regulated by both upstream kinases and its own kinase domain. PKA has

been reported to be an upstream regulator of S935 phosphorylation. (24, 29, 30). Furthermore, the PD R1441C/G/H mutation also impairs PKA-induced S1444 LRRK2 phosphorylation, and disrupts its interaction with 14-3-3 (31). Dephosphorylation of S910/S935/S955 and S973 is regulated by Protein Phosphatase 1 (PP1) and thereby stimulates the dissociation of 14-3-3 proteins from LRRK2 (25). In addition, several auto-phosphorylation sites within the Roc domain have an impact on GTPase and kinase activity (7, 31–37). We have recently revealed an intramolecular feedback regulation of the LRRK2 Roc G domain by a LRRK2 kinase dependent mechanism (36, 37). Biochemical analysis showed that auto-phosphorylation of the Roc domain reduced the GTPase activity, most likely by regulating the monomer-dimer equilibrium of LRRK2 (36, 37). Furthermore, the phosphorylation status of the T1503 residue has been shown to be essential for the kinase activity as well as for the GTP binding properties of LRRK2 (35)while S1444 phosphorylation has been shown to influence the kinase activity of LRRK2 (31) .

We have previously reported subunits of the PP1 and PP2A classes of phosphatases as well as the PAK6 kinase as regulators of LRRK2 dephosphorylation (25, 38–42). However, so far, no phosphatases have been identified that regulate the phosphorylation status of the Roc domain. We have previously identified PP2A as an interacting partner of the LRRK2 RocCOR-tandem domain (43). Furthermore, activation and overexpression of PP2A results in neuroprotection in SH-SY5Y cells expressing LRRK2-R1441C, primary cortical neurons expressing LRRK2-G2019S, and in a LRRK2-*Drosophila* model (43, 44). In addition, we have shown that PAK6-mediated phosphorylation of the PP2A subunit PPP2R2C regulates recruitment of PPP2R2C to the LRRK2 complex and PPP2R2C subcellular localization (45). Taken together, this shows that PP2A plays an important role in regulating LRRK2 activation and functioning.

In this study, we demonstrate that PP2A dephosphorylates LRRK2 at Threonine 1503 (T1503) of the Roc domain and thereby regulates LRRK2 dimerization and kinase activity. Surprisingly, our data reveal that LRRK2 can in turn also phosphorylate Threonine 304 (T304) of the catalytic subunit of PP2A (PPP2CA). PPP2CA-T304 phosphorylation results in impaired PP2A holoenzyme assembly and reduced enzymatic phosphatase activity. Together, our data reveal a feedback loop between LRRK2 autophosphorylation and PP2A activity, which might explain the neuroprotective properties of PP2A's enzymatic activity in LRRK2-induced parkinsonism.

Results

PP2A is able to reduce LRRK2 dimerization, as well as its kinase activity.

To explain the previously observed neuroprotective properties of PP2A's enzymatic activity in LRRK2-PD neuronal cell lines (43, 44), it is crucial to characterize how PP2A impacts LRRK2 kinase function. Since the neurotoxicity of PD-mutant LRRK2 is tightly associated with its dimerization dependent kinase activity (46–52), we tested whether dimerization and kinase activity are altered by PPP2CA expression by using a method to label and purify dimeric LRRK2 from cells (51). Over-expression of PPP2CA in HEK293T cells together with the proximity biotinylation LRRK2 construct pairs reduced LRRK2 dimerization (Fig. 1A). Moreover, the *in vitro* kinase activity of the isolated dimers (phosphorylation of NICtide) was also reduced in cells co-expressing WT-PPP2CA (Fig. 1B). In a cellular setting, the over-expression of WT-PPP2CA reduced the auto-phosphorylation of LRRK2 at S1292 induced by the G2019S mutation or exposure to H_2O_2 (Fig. 1C); as well as phosphorylation of the LRRK2 substrate Rab10 (T73). This reduction in LRRK2 cellular activity by PPP2CA was evident both in total cellular extracts as well as in isolated dimers of LRRK2. These data suggest that PP2A expression can regulate the dimerization, and thus kinase activity of LRRK2.

PP2A is able to dephosphorylate LRRK2 *in vitro* and in A549 cells

To further understand the mechanism by which PP2A regulates LRRK2 dimerization and kinase activity we next tried to identify the PP2A target sites. Previously, Liu et al., developed an assay in which they phosphorylated the human LRRK2 Roc domain using the kinase domain of the *Dictyostelium* LRRK2 orthologue Roco4 (53). Since we have shown that the COR domain plays an important role in LRRK2 dimerization (52, 54) we used a similar approach, but with purified recombinant purified LRRK2 RocCOR instead of a Roc-only protein (55). The recombinant human RocCOR fragment of LRRK2 was incubated with recombinant purified Roco4 kinase in reaction buffer containing ATP. Consistent with Liu et al., we see that incubation of LRRK2 RocCOR with the Roco4 kinase domain results in phosphorylation of the RocCOR domain. Besides multiple Roco4 autophosphorylation sites, confirming that active kinase was used, we were able to detect significant phosphorylation at T1503 of the RocCOR fragment by a DIA-based MS approach (Supplementary Fig. 1).

Since PP2A is interacting with LRRK2 via the Roc domain as a docking site (43), and T1503 residue has been shown to be important for the kinase and GTP binding properties of LRRK2, we tested with a commercial phospho-T1503 antibody, whether PP2A has an impact on LRRK2 phosphorylation status at this residue. Addition of the purified PPP2CA catalytic domain, results in significantly reduced phosphorylation of the RocCOR fragment of LRRK2 at T1503 (Fig. 2A + B). A549 lung carcinoma cell

line have high expression of LRRK2 and are therefore commonly used to measure the activation of LRRK2 in cells (56–58). We used chloroquine-induced lysosomal stress to increase LRRK2 activity (59). Additional treatment of A549 cells with Okadaic acid (OA), an inhibitor more selective for PP2A, resulted in the upregulation of the pT1503 signal from endogenously immunoprecipitated LRRK2, compared to the pT1503 signal of DMSO-treated (control) A549 cells (Fig. 2C + D). We next investigated whether the phospho-dead mutant version T1503A, also has an effect on LRRK2 activity and dimerization. Indeed, homo-dimers of T1503A-LRRK2 showed decreased dimerization and kinase activity, compared to wild-type LRRK2 (Fig. 3A + B); whereas hetero-dimers of WT/T1503A fail to de-stabilize dimers (Fig. 3C), yet have slightly reduced kinase activity (Fig. 3D). Together our *in vitro* and cellular data show that PP2A can dephosphorylate LRRK2 at T1503 and that this is important for the regulation of LRRK2 dimerization and kinase activity.

LRRK2 phosphorylates PPP2CA at T304 and regulates PP2A holoenzyme formation and its catalytic activity.

Surprisingly, using phospho-mass spectrometry we observed that LRRK2 can also phosphorylate PPP2CA at its critical T304 residue (phospho-Mass Spectrometry, Supplementary Fig 2). The T304 residue in the PPP2CA sequence is followed by a proline residue. Previous work showed that LRRK2 does have the potential to phosphorylate ThrPro sites (60). Unfortunately, there is no commercial pT304-PPP2CA antibody available. Therefore, we used a pThrPro antibody to detect and confirm LRRK2-mediated phosphorylation of PPP2CA-T304. We co-expressed LRRK2-G2019S with either N-terminally twintrep-FLAG (NSF) tagged WT-PPP2CA or the phosphodead T304A-PPP2CA mutant and measured the phosphorylation status after streptavidin pull down (Fig. 4A). We observed very clear phosphorylation of PPP2CA, while the T304A mutant showed a strongly reduced signal, confirming that we can use the pThrPro antibody to measure pT304. Next, purified human full length LRRK2 and PPP2CA were tested in an *in vitro* kinase assay (Fig. 4B). In the presence of ATP, LRRK2 is phosphorylating its auto-phosphorylation site S1292 and PPP2CA at T304, while in the presence of MLi-2, a potent and selective LRRK2 inhibitor, no phosphorylation was observed. To test LRRK2-mediated phosphorylation of PPP2CA under endogenous conditions, we immunoprecipitated the catalytic PPP2CA and PPP2CB subunits using a PP2Ac specific antibody. The pThrPro signal of the immunoprecipitated catalytic PP2A subunits was significantly reduced in the KO-LRRK2 RAW 264.7 cells, compared to WT RAW 264.7 (Fig. 4C +D). Consistently, the PP2A catalytic subunits are phosphorylated in A549 cells, however when the LRRK2 kinase activity was inhibited with MLi-2, the pT304 signal was diminished (Supplementary Fig. 3). Together this thus shows that PPP2CA is a LRRK2 substrate.

Previous data have shown that the C-terminal region of PPP2CA is essential for binding to its regulatory B subunit. C-terminally deleted PPP2CA mutants failed to interact with different regulatory B subunits ((61, 62)Supplementary Fig. 4), while the phosphomimicking mutant T304D-PPP2CA, loses its interaction with PPP2R2B (B55β1) and PPP2R2A (B55α) subunits (62). Consistently, our pull-down assays in HEK293 cell lysates showed that the interaction of PPP2R2B with T304D-PPP2CA is reduced compared to WT-PPP2CA or T304A-PPP2CA (Fig. 5A). Previous studies (43, 44) have shown that PP2A is neuroprotective against mutant LRRK2-induced cell death; and that this is regulated via dephosphorylation of S6K and requires binding of PPP2R5C (B56γ3) PP2A to the holoenzyme (63). Therefore, we next determined the interaction between PPP2CA phospho-mutants and PPP2R5C. Pull down experiments in HEK293 cells show that both T304A-PPP2CA and T304D-PPP2CA have strongly reduced interaction with PPP2R5C, compared to WT-PPP2CA (Fig. 5B + C). Therefore, we conclude that the de-regulation of T304 phosphorylation status of PPP2CA regulates holoenzyme formation. Next, we investigated whether T304A-PPP2CA or T304D-PPP2CA mutants exhibit alterations in their enzymatic phosphatase activity. By overexpressing and purifying WT/T304A/T304D-PPP2CA proteins from HEK293 cells and subsequently performing an *in vitro* phosphatase assay using a phosphopeptide and malachite green, we found that T304A-, as well as T304D-PPP2CA, exhibit lower phosphatase activity towards a phospho-Threonine peptide (K-R-pT-I-R-R), compared to the wild-type PPP2CA (Fig. 5D). Together, these data show that LRRK2 phosphorylates PPP2CA at position T304 and that this site is important for the regulation of holoenzyme complex formation and its catalytic activity.

Previous studies have shown that either overexpression of PPP2CA or addition of sodium selenate, as a PP2A activator, was able to reverse G2019S-LRRK2-induced neurotoxicity in primary cortical neurons (43). Similar data were shown in *Drosophila*, where overexpression of different single PP2A subunits were able to block G2019S-LRRK2-induced cell death in dopaminergic neurons (44). Consistently, our current data show that co-expression of WT-PPP2CA and G2019S-LRRK2 in primary mouse embryonic cortical neurons significantly decreases the neurotoxicity of G2019S-LRRK2. However, when T304A-PPP2CA is expressed, it fails to protect from G2019S-LRRK2-induced neurotoxicity (Fig. 6), suggesting that phosphorylation of PPP2CA at this residue, and thus its phosphatase activity, is critical for its neuroprotective effects in this primary neuronal model.

Discussion

The importance of phosphatases in mutant LRRK2-induced parkinsonism has already been established by numerous studies. PPP1CA regulates the important N-terminal phosphorylation cluster, including S910 and S935. Interestingly, while PPP2CA alone does not have a significant direct effect on regulating S910/S935-LRRK2 phosphorylation (25), a phosphatome wide RNAi study revealed that PPP2CA associated to its regulatory subunit PPP2R2A/B was a potent regulator of those sites (42). Related to this, LRRK2 autophosphorylation is reported to be sensitive to okadaic acid, suggesting that PP2A is also involved in regulating LRRK2 autophosphorylation sites (41). In addition, PPM1H has been shown not to dephosphorylate LRRK2 directly, but to counteract LRRK2 function, by dephosphorylating LRRK2's downstream targets, namely the small Rab GTPases (16). Although we previously have shown that PP2A can bind to the ROC domain and that PP2A has neuroprotective effects in mutant LRRK2-induced cell death in SH-SY5Y neuroblastoma cells (43), and primary neurons (the present study), the molecular mechanism by which PP2A regulates LRRK2 signaling remained unclear.

In this study, we revealed a novel regulatory loop between LRRK2 and PP2A. Using *in vitro* and cellular models, we could identify the T1503 residue as one of the phospho-sites in the RocCOR domain of LRRK2, which can be dephosphorylated by PP2A. Both the LRRK2 T1503A mutant and overexpression of the catalytic subunit of the PP2A holoenzyme, PPP2CA, resulted in decreased LRRK2 dimerization and kinase activity. This suggests that the specific autophosphorylation-sites in LRRK2 modified by PP2A are critical in maintaining dimer stability, and by extension – kinase activity. Surprisingly, our data additionally revealed that LRRK2 could also phosphorylate PP2A in its catalytic subunit at the T304 residue. Impaired T304 phosphorylation results in impaired holoenzyme formation and reduced phosphatase activity. Previously it has been shown that de-regulation of T304 phosphorylation alters methylation of the C-terminus and induces the dissociation of PPP2CA from the B regulatory subunits, resulting in changes in the subcellular localization of the PP2A holoenzyme and its substrate repertoire (61). This bi-directional regulation between LRRK2 and PP2A can regulate subcellular localization of active LRRK2 and PP2A, thus regulating localized signaling within the cell. Importantly, expression of WT-PPP2CA protects from G2019S-LRRK2 induced neuronal cell death, while PPP2CA phospho-mutants at the LRRK2 regulated site T304 fail to do so. Interestingly, PP2A activity is downregulated in αSyn PD models, having the same net-effect on PP2a activity as with hyper-active mutant LRRK2 (64, 65). Furthermore, in postmortem brain samples (substantia nigra-SN and frontal cortex (FC) from PD or dementia with Lewy body (DLB) patients, PP2A methylation in PD and DLB samples was greatly decreased, while Leucine carboxyl methyltransferase (LCMT-1), PP2A's methylating enzyme, was significantly reduced in the SN of PD patients and protein phosphatase methylesterase (PME-1), a demethylating enzyme of PP2A, was elevated

in the SN and FC of PD and DLB patients (64). In addition, impaired activation of the PP2A phosphatase has been linked with autosomal recessive early-onset parkinsonism with intellectual disability (66).

Together, this shows that the regulation of the C-terminal segment of PPP2CA is crucial for the progression of PD in general. In addition to acting on LRRK2, PP2A has been shown to dephosphorylate αSyn at S129, which is believed to play a role in the pathogenesis of αSyn PD (67, 68). Moreover, it has been shown that PP2A can also dephosphorylate Tau *in vitro* (69–72). Tau plays a crucial role in AD pathology and in addition has been identified as a potential PD factor in Genome wide associate studies (73). Interestingly, it has been shown that PP2A activity is downregulated in AD brains (74, 75). Understanding the regulation of PP2A, by post-translational modifications such as phosphorylation and methylation, can thus give important insight into the onset and progression of PD in general.

In summary, our data show that PP2A can dephosphorylate LRRK2, leading to de-stabilization of LRRK2 dimers and a reduction in its kinase activity. Conversely, LRRK2 can also phosphorylate the PP2A catalytic subunit. This latter event results in changes in PP2A holoenzyme formation leading to the reduction of PP2A dephosphorylation activity. Together, this suggests LRRK2 and PP2A regulate the activity of one another in a bi-directional manner. Our data suggest that this cycle involves PP2A dephosphorylating of LRRK2 at autophosphorylation sites and inhibiting LRRK2 activity (via de-stabilization of dimers), or LRRK2- phosphorylating and blocking PP2A, allowing mutant forms of LRRK2 to trigger a cell death pathway. Importantly, our study also suggests that boosting PP2A activity by stimulating complex formation might be beneficial for mutant LRRK2-mediated PD and PD in general.

Materials and Methods

Cell Culture, pharmacological treatment, transfection and cell lysis

HEK293 cells were cultured in DMEM high glucose (#11960-044, Gibco), supplemented with 10% fetal bovine serum (FBS) (A31604-01, Gibco) and penicillin/streptomycin/glutamine (#10378-016, Gibco). HEK293 cells were transfected with jetPEI (#101-10, Westburg) according to the manufacturer's protocol, or by incubating with DNA: CaPO4 precipitates. Protein expression was validated 48hrs after transfection with fluorescence microscopy and/or western blotting. Murine RAW 264.7 cells, parental (#sc-6003, ATCC) or Knock-out (KO) (# sc-6004, ATCC), were cultured in DMEM high glucose (#20-2002, ATCC), supplemented with 10% fetal bovine serum (FBS) (#20-2020, ATCC) and penicillin/streptomycin (#15070-063, Gibco). A549 cells were obtained from Prof. Amalia Dolga, (University of Groningen, the Netherlands) and were cultured in RPMI-160 (#12-702F, Lonza), supplemented with 10% FBS (#A31604-01, Gibco), and penicillin/streptomycin (#15070-063, Gibco). As a general lysis buffer protocol, cells were washed three times with wash buffer (50 mM Tris-HCl 7.5, 150 mM NaCl, 1mM EDTA, 2.5 mM EGTA). Then they were lysed (wash buffer supplemented with 0.1% NP-40, protease inhibitors (p2714-1btl, Sigma) (76), and phosphatase inhibitors (1mM Na3VO4, 1mM NaF, 5 mM β-glycerol phosphate).

In order to increase LRRK2 kinase activity and subsequently its autophosphorylation, 50 μM of chloroquine (#C6628, Sigma Aldrich) was used for treatment of A549 cells overnight (77). For inhibition of the endogenous PP2A holoenzyme, 100 nM of okadaic acid (#O7885, Sigma Aldrich) was used according to the Lobbestael et al. 2013 study, (25) while DMSO was used as a vehicle control, 60 minutes before the cells got harvested.

Primary Antibodies

We used the following antibodies in this study: anti-PP2Ac (detects PPP2CA and PPP2CB proteins) (#2038, Cell Signaling), anti-pThreonine-Proline (#9391, Cell signaling), anti-pS1292-LRRK2 (#203181, Abcam), anti-GST (#27457701, GE Healthcare), anti-His (#ab1187, Abcam), anti-LRRK2 (clone c41-2, Abcam), anti-Strep (#34850, Qiagen), anti-myc (#9E10:sc-40, Santa Cruz Biotechnology), and anti-pT1503 (#154423, Abcam).

Purification of dimeric LRRK2

To purify LRRK2 dimers to be used in our *in vitro* assessments of kinase function, we relied on the proximity biotinylation technique previously described (51). Briefly, two cDNAs were created encoding LRRK2 fusions with biotin ligase (BirA; N-term, Flag-tagged) and an acceptor peptide (AP, N-term; c-Myc tagged), and over-expressed in HEK293T cells grown in biotin-depleted medium (OptiMEM+2%

FBS). After 48hr following transfection, the cells were extensively washed in PBS, given a brief biotin pulse (50 μM, 5 min, 37°C), followed by another 3X washes in PBS, centrifuged and the pellet snap frozen in a dry-ice/MeOH bath. Following lysis, extracts were diluted in TBST/BSA (10 mM Tris HCl, pH 7.6; 100 mM NaCl; 0.1% Triton X-100; 1% BSA) and 2.5 μg of protein loaded in parallel ELISA plates, coated with streptavidin (SA; to capture biotinylated LRRK2 dimers) and anti-LRRK2 (to quantify LRRK2 over-expression). To detect and quantify dimeric LRRK2, SA-coated plates were incubated with HRP-conjugated anti-Flag antibodies (1hr, room temperature). Since the biotin tag is only present on AP-LRRK2 fusions, and the Flag epitope tag is located on the BirA-LRRK2 fusion, by using HRP-Flag as our detector reagent, we are specifically labeling dimeric LRRK2 present in the ELISA plates. On parallel anti-LRRK2 coated plates (clone c41-2), total over-expressed LRRK2 is quantified using HRP-LRRK2 antibodies (clone N241), and used to normalize the relative amounts of dimeric LRRK2. We assessed the following LRRK2 dimers: WT/WT and G2019S/G2019S homo-dimers; as well as LRRK2 with the T1503 site mutated to Ala, in hetero- and homo-dimer conformations.

We co-expressed PPP2CA (WT, T304A, or T304D) with WT or mutant G2019S-LRRK2 dimer construct pairs. Here, the ratio of expression of LRRK2 to PPP2CA was maintained at 3:1. Dimeric LRRK2 was determined as above with the following modification: LRRK2 dimers were estimated in SA-coated ELISA plates using HRP-conjugated anti-LRRK2 (clone N241; NeuroMab). We measured *in vitro* kinase activity of captured dimers using NICtide as a peptide substrate as described previously (51). To assess phosphorylation at Ser935 in dimeric LRRK2, we used anti-pS935 LRRK2 conjugated to HRP (Expedion/Abcam) as our detector reagent, as recently described (78).

Preparation of primary mouse neuronal cultures, and assessment of neuronal death.

Embryonic day 16 (E16) pregnant C57BL mice were used in this study, with primary cortical neurons prepared as described (79). Briefly, under aseptic conditions, cortices were removed and cut into small pieces before enzymatic digestion (trypsin 0.05%) and mechanical dissociation. Cells were centrifuged and counted and plated at a density of 150,000/cm^2 in BrainPhys neuronal culture medium (StemCell Technologies) supplemented with SM1 Neuronal Supplement (StemCell Technologies), L-glutamine (0.5 mM) and penicillin/streptavidin. After 3-4 DIV, neurons were transfected using Lipofectamine 2000 (ThermoScientific) as per the manufacturer's instructions. Neurons were transfected with WT or mutant (G2019S) human LRRK2 (in pcms-EGFP vectors), with EGFP co-expressed under a separate promoter. Co-transfections with WT, T304A, or T304D-PPP2CA, or control plasmid, were performed at a ratio of 3:1 with excess LRRK2 plasmid. After three days following transfection, the coverslips were washed in PBS and fixed in 3.7% paraformaldehyde for 20 min at 4°C. The neurons were processed for immunofluorescence labeling with the following antibodies: GFP (chicken; Abcam), Flag (M2 mouse;

Sigma-Aldrich), active caspase-3 (rabbit; R&D Systems), and DAPI nuclear stain. Mounted coverslips were imaged on a Leica TSP5 multi-photon confocal microscope, and the Z-stacks processed in ImageJ, and Adobe Photochop. For quantification of apoptotic neuronal profiles, we used the approach described by Antoniou and colleagues (79).

Immunoprecipitation of endogenous PPP2CA/ LRRK2 and streptavidin pull down

To purify the endogenous PP2A catalytic subunits from RAW264.7 cells, the anti PP2Ac antibody (#2038, Cell signalling), detecting both PPP2CA and PPP2CB, p, together with Protein A-conjugated magnetic beads (#161-4013, Biorad). For the immunoprecipitation of endogenous LRRK2 from A549 cells, the Abcam LRRK2 antibody (#c41-2, Abcam) was used with Protein A-conjugated magnetic beads. In both cases, the following lysis/wash buffer was used: 10 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5 % NP-40. To purify overexpressed NSF-PPP2CA, WT and T304A, as well as NSF-PPP2R2B and NSF-PPP2R5C, the cells were lysed with 50 mM Tris-HCl 7.5, 150 mM NaCl, 1mM EDTA, 2.5 mM EGTA, 1% NP-40, protease inhibitors (p2714, Sigma) and phosphatase inhibitors (1mM Na3VO4, 1mM NaF, 5 mM beta glycerol phosphate) and incubated with streptavidin beads (#2-1206-025, IBA Lifesciences) for 4 hrs. Finally, they were washed 3 times with lysis buffer.

Phosphatase activity assay

Phosphatase activity of purified NSF-PPP2CA (WT and T304A) was measured using the Ser/Thr Phosphatase assay kit (#17-127, Merck; Fig. 5D). Over-expressed PPP2CA proteins were purified on streptavidin beads as described above, and the activity determined with proteins bound to the beads. The protocol for the phosphatase assay and representation of the data has been previously described (80).

Purification of proteins, *in vitro* assays and western blot assay

The lysis buffer used for LRRK2 purification from HEK293 cells consisted of 30 mM Tris pH 8.0, 150 mM NaCl, 5 mM MgCl2, 3 μM DTT, 5 % glycerol, 0.1 mM nucleotide, 0.5% NP-40), while the wash buffer is similar to the lysis buffer (excluding NP40). Finally, the streptavidin beads were used (#2-1206-025, IBA Lifesciences), while the 10x elution buffer (#2-1000-025, IBA Lifesciences) was diluted in wash buffer. The RocCOR domain of human LRRK2 fused with His and MBP tags at the N-terminus was expressed in *E. coli* BL21 (DE3). Cells were lysed in buffer consisted of 50 mM HEPES (pH 8), 150 mM NaCl, 10 mM MgCl2, 10% Glycerol, 0.5 mM GDP, 5 mM β-mercaptoethanol and complete EDTA free protease inhibitor cocktail by sonication, and then purified using MBP-Trap affinity chromatography column (GE Healthcare). The protein was eluted in the presence of 10mM maltose. For the *in vitro* kinase-phosphatase assay, the reaction buffer consisted of 25 mM Tris pH 7.5, 30 mM MgCl2, 150 mM NaCl, and

2mM DTT. 18 µg *Dictyostelium discoideum* 18 µg GST-Roco4-kinase (purified as previously described in (81)) 4 µg of His-MBP-RocCOR and 50 µM ATP were incubated at 25 degrees for 30 minutes. In order to block the Roco4 kinase activity and subsequently the phosphorylation of the RocCOR fragment, 10 mM of H1152 (#2414, Tocris Bioscience) was added. Finally, 1 µg of commercially available GST-PP2Ac (#128557, Abcam) was added to the solution for 105 minutes. The buffer of the *in vitro* kinase assay (Fig. 4B) consisted of 25 mM Tris pH 7.5, 15 mM MgCl2, 150 mM NaCl, 2 mM DTT. Recombinant purified full-length wild-type LRRK2 (3 µg) (82), 0.5 µg commercially available GST-PPP2CA (#128557, Abcam), and 50 µM ATP were incubated at 25°C for 30 minutes. In one of the samples, 1µM MLi-2 (#5756, Tocris) was included as a positive control for LRRK2 kinase inhibition.

Mass spectrometry

Phosphopeptide enrichment followed by LC-MSMS has been performed as previously described (83). Briefly, after *in vitro* phosphorylation, protein samples were precipitated with chloroform methanol and re-dissolved in 100 mM ammonium bicarbonate. The proteins were subsequently alkylated by DTT/ Idoacetaminde treatment and submitted to tryptic proteolysis over-night. The phosphopeptides have been enriched by TiO2 prior to their analysis on an Orbitrap Fusion mass spectrometer (Thermo-Fisher Scientific) coupled to a nanoflow HPLC system (Ultimate 3000 RSLC; Thermo-Fisher Scientific) which was equipped with a µPac C18 reverse phase column (Thermo-Fisher Scientific). Full scans were acquired at 120K resolution (mass range: m/z 400-1000). MS2 Spectra were acquired via DIA in the Orbitrap (15K resolution), using window-wise CID fragmentation in 31 m/z windows of 20 Da width (1Da overlap) and 2h gradients. For spectral library creation, LRRK2-G2019S was pre-phosphorylated *in vitro* and processed as described above and analyzed via DDA to generate a spectral library. Instead, Roco4 MS2 spectra were directly extracted from the DIA files by DIAUmpire (2.0) (84). For this purpose, MS2 spectra were searched in Mascot (v.2.5.1)(84) using carbamidomethyl cysteine as fixed modification and oxidizes methionine and phospho-STY as variable modifications. The search was performed against the human subset of the SwissProt database (release 2016_08) supplemented by the Roco4 sequence (total: 20178 entries). The peptide tolerance was set to 10 ppm, the MSMS tolerance was set to 20 mmu. Library creation from Mascot .dat files and final data analysis was performed in Skyline (v21.2.0.568) (85).

Data availability

Source data will be made available via Zenodo.org

Author contributions

P.S.A performed most of the biochemical and cellular experiments with the help of S.S. and A.M.T.; A.M. H.J.R. and V.P. performed the dimerization assays, kinase assays and primary neuronal cell death assays; A.S. and H.P. assisted P.S.A. with cloning of different constructs; F.Y.H. assisted with the purification of proteins; F.v.Z. and C.J.G. performed the Mass spectrometry experiments/analysis; L.V., W.S., M.C.C.H, R.J.N., E.G. and J.M.T. interpreted experiments; P.S.A., A.K. and H.J.R. conceived the study; A.K. and H.J.R. supervised the experiments; P.S.A., A.K., and H.J.R. wrote the manuscript with the help and input of all other authors.

Acknowledgments

We gratefully acknowledge Prof. Chi-Wu Chiang (National Cheng Kung University, Taiwan) for providing us with the PPP2CA, PPP2R5C and PPP2R2B sequences. We thank the staff of the Core Facility for Medical Proteomics at the University of Tubingen for technical assistance.

Funding

We have received funding from the Michael J. Fox Foundation for Parkinson's Research (project 6709.3. to A.K, JMT, MCCH, R.JN, EG).

Competing interest statement

The authors declare no conflict of interest

References

1. A. J. Lees, J. Hardy, T. Revesz, Parkinson's disease. *Lancet* **373**, 2055–66 (2009).
2. T. Gasser, J. Hardy, Y. Mizuno, Milestones in PD genetics. *Movement Disorders* **26**, 1042–1048 (2011).
3. M. R. Cookson, O. Bandmann, Parkinson's disease: insights from pathways. *Hum Mol Genet* **19**, R21-7 (2010).
4. B. K. Gilsbach, A. Kortholt, Structural biology of the LRRK2 GTPase and kinase domains: implications for regulation. *Front Mol Neurosci* **7** (2014).
5. A. Biosa, *et al.*, GTPase activity regulates kinase activity and cellular phenotypes of Parkinson's disease-associated LRRK2. *Hum Mol Genet* **22**, 1140–1156 (2013).
6. C. J. Gloeckner, *et al.*, The Parkinson disease causing LRRK2 mutation I2020T is associated with increased kinase activity. *Hum Mol Genet* **15**, 223–232 (2006).
7. A. B. West, *et al.*, Parkinson's disease-associated mutations in leucine-rich repeat kinase 2 augment kinase activity. *Proceedings of the National Academy of Sciences* **102**, 16842–16847 (2005).
8. A. Zimprich, *et al.*, Mutations in LRRK2 Cause Autosomal-Dominant Parkinsonism with Pleomorphic Pathology. *Neuron* **44**, 601–607 (2004).
9. C. Paisán-Ruíz, *et al.*, Cloning of the Gene Containing Mutations that Cause PARK8-Linked Parkinson's Disease. *Neuron* **44**, 595–600 (2004).

10. D. R. Alessi, E. Sammler, LRRK2 kinase in Parkinson's disease. *Science (1979)* **360**, 36–37 (2018).

11. R. Di Maio, *et al.*, LRRK2 activation in idiopathic Parkinson's disease. *Sci Transl Med* **10** (2018).

12. J.-M. Taymans, *et al.*, Perspective on the current state of the LRRK2 field. *NPJ Parkinsons Dis* **9**, 104 (2023).

13. D. A. Roosen, M. R. Cookson, LRRK2 at the interface of autophagosomes, endosomes and lysosomes. *Mol Neurodegener* **11**, 73 (2016).

14. M. Steger, *et al.*, Phosphoproteomics reveals that Parkinson's disease kinase LRRK2 regulates a subset of Rab GTPases. *Elife* **5**, 1–28 (2016).

15. T. Komori, T. Kuwahara, An Update on the Interplay between LRRK2, Rab GTPases and Parkinson's Disease. *Biomolecules* **13**, 1645 (2023).

16. K. Berndsen, *et al.*, PPM1H phosphatase counteracts LRRK2 signaling by selectively dephosphorylating Rab proteins. *Elife* **8**, 1–37 (2019).

17. A. Adhikari, A. Tripathi, C. Y. Chiang, P. Sherpa, S. R. Pfeffer, Allosteric regulation of the Golgi-localized PPM1H phosphatase by Rab GTPases modulates LRRK2 substrate dephosphorylation in Parkinson's disease. *Journal of Biological Chemistry* 110679 (2025). https://doi.org/10.1016/j.jbc.2025.110679.

18. N. Dzamko, *et al.*, LRRK2 levels and phosphorylation in Parkinson's disease brain and cases with restricted Lewy bodies. *Movement Disorders* **32**, 423–432 (2017).

19. S. Wang, *et al.*, Elevated LRRK2 autophosphorylation in brain-derived and peripheral exosomes in LRRK2 mutation carriers. *Acta Neuropathol Commun* **5**, 86 (2017).

20. P. S. Athanasopoulos, R. Heumann, A. Kortholt, The role of (auto)-phosphorylation in the complex activation mechanism of LRRK2. *Biol Chem* **399**, 643–647 (2018).

21. A. Marchand, *et al.*, A Phosphosite Mutant Approach on LRRK2 Links Phosphorylation and Dephosphorylation to Protective and Deleterious Markers, Respectively. *Cells* **11**, 1018 (2022).

22. N. Dzamko, *et al.*, Inhibition of LRRK2 kinase activity leads to dephosphorylation of Ser(910)/Ser(935), disruption of 14-3-3 binding and altered cytoplasmic localization. *Biochem J* **430**, 405–13 (2010).

23. R. J. Nichols, *et al.*, 14-3-3 binding to LRRK2 is disrupted by multiple Parkinson's disease-associated mutations and regulates cytoplasmic localization. *Biochem J* **430**, 393–404 (2010).

24. E. A. Doggett, J. Zhao, C. N. Mork, D. Hu, R. J. Nichols, Phosphorylation of LRRK2 serines 955 and 973 is disrupted by Parkinson's disease mutations and LRRK2 pharmacological inhibition. *J Neurochem* **120**, 37–45 (2012).

25. E. Lobbestael, *et al.*, Identification of protein phosphatase 1 as a regulator of the LRRK2 phosphorylation cycle. *Biochemical Journal* **456**, 119–128 (2013).

26. M. Blanca Ramírez, *et al.*, GTP binding regulates cellular localization of Parkinson's disease-associated LRRK2. *Hum Mol Genet* **26**, 2747–2767 (2017).

27. N. J. Lavalley, S. R. Slone, H. Ding, A. B. West, T. A. Yacoubian, 14-3-3 Proteins regulate mutant LRRK2 kinase activity and neurite shortening. *Hum Mol Genet* **25**, 109–122 (2016).

28. E. Giusto, T. A. Yacoubian, E. Greggio, L. Civiero, Pathways to Parkinson's disease: a spotlight on 14-3-3 proteins. *NPJ Parkinsons Dis* **7**, 85 (2021).

29. A. B. West, *et al.*, Parkinson's disease-associated mutations in LRRK2 link enhanced GTP-binding and kinase activities to neuronal toxicity. *Hum Mol Genet* **16**, 223–232 (2007).

30. E. Greggio, L. Bubacco, I. Russo, Cross-talk between LRRK2 and PKA: implication for Parkinson's disease? *Biochem Soc Trans* **45**, 261–267 (2017).

31. K. Muda, *et al.*, Parkinson-related LRRK2 mutation R1441C/G/H impairs PKA phosphorylation of LRRK2 and disrupts its interaction with 14-3-3. *Proceedings of the National Academy of Sciences* **111** (2014).

32. J.-M. Taymans, *et al.*, LRRK2 Kinase Activity Is Dependent on LRRK2 GTP Binding Capacity but Independent of LRRK2 GTP Binding. *PLoS One* **6**, e23207 (2011).

33. E. Greggio, *et al.*, Kinase activity is required for the toxic effects of mutant LRRK2/dardarin. *Neurobiol Dis* **23**, 329–341 (2006).

34. V. S. Anand, *et al.*, Investigation of leucine-rich repeat kinase 2. *FEBS J* **276**, 466–478 (2009).

35. P. J. Webber, *et al.*, Autophosphorylation in the Leucine-Rich Repeat Kinase 2 (LRRK2) GTPase Domain Modifies Kinase and GTP-Binding Activities. *J Mol Biol* **412**, 94–110 (2011).

36. B. K. Gilsbach, *et al.*, Intramolecular feedback regulation of the LRRK2 Roc G domain by a LRRK2 kinase-dependent mechanism. *Elife* **12** (2024).

37. G. Guaitoli, *et al.*, Biophysical analysis reveals autophosphorylation as an important negative regulator of LRRK2 dimerization. [Preprint] (2023). Available at: http://biorxiv.org/lookup/doi/10.1101/2023.08.11.549911.

38. J.-M. Taymans, "Regulation of LRRK2 by Phosphatases" in *Advances in Neurobiology*, (2017), pp. 145–160.

39. L. Civiero, *et al.*, Leucine-rich repeat kinase 2 interacts with p21-activated kinase 6 to control neurite complexity in mammalian brain. *J Neurochem* **135**, 1242–1256 (2015).

40. S. Cogo, *et al.*, The Roc domain of LRRK2 as a hub for protein-protein interactions: a focus on PAK6 and its impact on RAB phosphorylation. *Brain Res* **1778**, 147781 (2022).

41. A. Reynolds, E. A. Doggett, S. M. Riddle, C. S. Lebakken, R. J. Nichols, LRRK2 kinase activity and biology are not uniformly predicted by its autophosphorylation and cellular phosphorylation site status. *Front Mol Neurosci* **7** (2014).

42. M. Drouyer, *et al.*, Protein phosphatase 2A holoenzymes regulate leucine-rich repeat kinase 2 phosphorylation and accumulation. *Neurobiol Dis* **157**, 105426 (2021).

43. P. S. Athanasopoulos, *et al.*, Identification of protein phosphatase 2A as an interacting protein of leucine-rich repeat kinase 2. *Biol Chem* **397**, 541–554 (2016).

44. J. P. L. Sim, *et al.*, Identification of PP2A and S6 Kinase as Modifiers of Leucine-Rich Repeat Kinase-Induced Neurotoxicity. *Neuromolecular Med* **22**, 218–226 (2020).

45. L. Iannotta, *et al.*, PAK6-mediated phosphorylation of PPP2R2C regulates LRRK2-PP2A complex formation. *Front Mol Neurosci* **16** (2023).

46. Z. Berger, K. A. Smith, M. J. LaVoie, Membrane Localization of LRRK2 Is Associated with Increased Formation of the Highly Active LRRK2 Dimer and Changes in Its Phosphorylation. *Biochemistry* **49**, 5511–5523 (2010).

47. E. Greggio, *et al.*, The Parkinson Disease-associated Leucine-rich Repeat Kinase 2 (LRRK2) Is a Dimer That Undergoes Intramolecular Autophosphorylation. *Journal of Biological Chemistry* **283**, 16906–16914 (2008).

48. C. L. Klein, *et al.*, Homo- and heterodimerization of ROCO kinases: LRRK2 kinase inhibition by the LRRK2 ROCO fragment. *J Neurochem* **111**, 703–715 (2009).

49. S. Sen, P. J. Webber, A. B. West, Dependence of Leucine-rich Repeat Kinase 2 (LRRK2) Kinase Activity on Dimerization. *Journal of Biological Chemistry* **284**, 36346–36356 (2009).

50. W. W. Smith, *et al.*, Kinase activity of mutant LRRK2 mediates neuronal toxicity. *Nat Neurosci* **9**, 1231–1233 (2006).

51. E. Leandrou, *et al.*, Kinase activity of mutant LRRK2 manifests differently in hetero-dimeric vs. homo-dimeric complexes. *Biochemical Journal* **476**, 559–579 (2019).

52. L. G. Helton, *et al.*, Allosteric Inhibition of Parkinson's-Linked LRRK2 by Constrained Peptides. *ACS Chem Biol* **16**, 2326–2338 (2021).

53. Z. Liu, J. A. Mobley, L. J. DeLucas, R. A. Kahn, A. B. West, LRRK2 autophosphorylation enhances its GTPase activity. *The FASEB Journal* **30**, 336–347 (2016).

54. P. Pathak, *et al.*, Doubly Constrained C-terminal of Roc (COR) Domain-Derived Peptides Inhibit Leucine-Rich Repeat Kinase 2 (LRRK2) Dimerization. *ACS Chem Neurosci* **14**, 1971–1980 (2023).

55. R. K. Singh, *et al.*, Nanobodies as allosteric modulators of Parkinson's disease–associated LRRK2. *Proc Natl Acad Sci U S A* **119** (2022).

56. R. N. Fuji, *et al.*, Effect of selective LRRK2 kinase inhibition on nonhuman primate lung. *Sci Transl Med* **7** (2015).

57. S. Biskup, *et al.*, Dynamic and redundant regulation of LRRK2 and LRRK1 expression. *BMC Neurosci* **8**, 102 (2007).

58. M. C. Herzig, *et al.*, LRRK2 protein levels are determined by kinase function and are crucial for kidney and lung homeostasis in mice. *Hum Mol Genet* **20**, 4209–4223 (2011).

59. T. Kuwahara, *et al.*, Roles of lysosomotropic agents on LRRK2 activation and Rab10 phosphorylation. *Neurobiol Dis* **145**, 105081 (2020).

60. R. J. Nichols, *et al.*, Substrate specificity and inhibitors of LRRK2, a protein kinase mutated in Parkinson's disease. *Biochemical Journal* **424**, 47–60 (2009).

61. E. Ogris, D. M. Gibson, D. C. Pallas, Protein phosphatase 2A subunit assembly: the catalytic subunit carboxy terminus is important for binding cellular B subunit but not polyomavirus middle tumor antigen. *Oncogene* **15**, 911–917 (1997).

62. S. Longin, *et al.*, Selection of Protein Phosphatase 2A Regulatory Subunits Is Mediated by the C Terminus of the Catalytic Subunit. *Journal of Biological Chemistry* **282**, 26971–26980 (2007).

63. K. Hahn, *et al.*, PP2A Regulatory Subunit PP2A-B′ Counteracts S6K Phosphorylation. *Cell Metab* **11**, 438–444 (2010).

64. H. Park, *et al.*, Dysregulation of protein phosphatase 2A in parkinson disease and dementia with lewy bodies. *Ann Clin Transl Neurol* **3**, 769–780 (2016).

65. H. Tian, *et al.*, Leucine Carboxyl Methyltransferase Downregulation and Protein Phosphatase Methylesterase Upregulation Contribute Toward the Inhibition of Protein Phosphatase 2A by α-Synuclein. *Front Aging Neurosci* **10** (2018).

66. C. Fevga, *et al.*, *PTPA* variants and impaired PP2A activity in early-onset parkinsonism with intellectual disability. *Brain* **146**, 1496–1510 (2023).

67. J. Qu, *et al.*, The Molecular Mechanism of Alpha-Synuclein Dependent Regulation of Protein Phosphatase 2A Activity. *Cellular Physiology and Biochemistry* **47**, 2613–2625 (2018).

68. N. Ramalingam, *et al.*, Dynamic physiological α-synuclein S129 phosphorylation is driven by neuronal activity. *NPJ Parkinsons Dis* **9**, 4 (2023).

69. M. Kuszczyk, W. Gordon-Krajcer, J. W. Lazarewicz, Homocysteine-induced acute excitotoxicity in cerebellar granule cells in vitro is accompanied by PP2A-mediated dephosphorylation of tau. *Neurochem Int* **55**, 174–180 (2009).

70. L. Martin, A. Magnaudeix, F. Esclaire, C. Yardin, F. Terro, Inhibition of glycogen synthase kinase-3β downregulates total tau proteins in cultured neurons and its reversal by the blockade of protein phosphatase-2A. *Brain Res* **1252**, 66–75 (2009).

71. J.-M. Taymans, V. Baekelandt, Phosphatases of α-synuclein, LRRK2, and tau: important players in the phosphorylation-dependent pathology of Parkinsonism. *Front Genet* **5** (2014).

72. M. Bennecib, C.-X. Gong, I. Grundke-Iqbal, K. Iqbal, Role of protein phosphatase-2A and -1 in the regulation of GSK-3, cdk5 and cdc2 and the phosphorylation of tau in rat forebrain. *FEBS Lett* **485**, 87–93 (2000).

73. M. Sharma, *et al.*, Large-scale replication and heterogeneity in Parkinson disease genetic loci. *Neurology* **79**, 659–667 (2012).

74. M. Voronkov, S. P. Braithwaite, J. B. Stock, Phosphoprotein Phosphatase 2A: A Novel Druggable Target for Alzheimer's Disease. *Future Med Chem* **3**, 821–833 (2011).

75. L. Martin, *et al.*, Tau protein phosphatases in Alzheimer's disease: The leading role of PP2A. *Ageing Res Rev* **12**, 39–49 (2013).

76. T. Teruya, S. Simizu, N. Kanoh, H. Osada, Phoslactomycin targets cysteine-269 of the protein phosphatase 2A catalytic subunit in cells. *FEBS Lett* **579**, 2463–2468 (2005).

77. T. Eguchi, *et al.*, LRRK2 and its substrate Rab GTPases are sequentially targeted onto stressed lysosomes and maintain their homeostasis. *Proceedings of the National Academy of Sciences* **115**, E9115–E9124 (2018).

78. K. Melachroinou, *et al.*, Elevated In Vitro Kinase Activity in Peripheral Blood Mononuclear Cells of Leucine-Rich Repeat Kinase 2 G2019S Carriers: A Novel Enzyme-Linked Immunosorbent Assay-Based Method. *Movement Disorders* **35**, 2095–2100 (2020).

79. N. Antoniou, *et al.*, A motif within the armadillo repeat of Parkinson's-linked LRRK2 interacts with FADD to hijack the extrinsic death pathway. *Sci Rep* **8**, 3455 (2018).

80. L. Mao, *et al.*, Role of Protein Phosphatase 2A in mGluR5-regulated MEK/ERK Phosphorylation in Neurons. *Journal of Biological Chemistry* **280**, 12602–12610 (2005).

81. B. K. Gilsbach, *et al.*, Roco kinase structures give insights into the mechanism of Parkinson disease-related leucine-rich-repeat kinase 2 mutations. *Proceedings of the National Academy of Sciences* **109**, 10322–10327 (2012).

82. G. Guaitoli, *et al.*, Structural model of the dimeric Parkinson's protein LRRK2 reveals a compact architecture involving distant interdomain contacts. *Proc Natl Acad Sci U S A* **113**, E4357–E4366 (2016).

83. C. J. Gloeckner, *et al.*, Phosphopeptide Analysis Reveals Two Discrete Clusters of Phosphorylation in the N-Terminus and the Roc Domain of the Parkinson-Disease Associated Protein Kinase LRRK2. *J Proteome Res* **9**, 1738–1745 (2010).

84. C.-C. Tsou, *et al.*, DIA-Umpire: comprehensive computational framework for data-independent acquisition proteomics. *Nat Methods* **12**, 258–264 (2015).

85. L. K. Pino, *et al.*, The Skyline ecosystem: Informatics for quantitative mass spectrometry proteomics. *Mass Spectrom Rev* **39**, 229–244 (2020).

Figure 1: PPP2CA overexpression results in decreased LRRK2 dimerization and kinase activity. We examined the degree of dimer formation (A) and intrinsic kinase activity of isolated dimers (B) for wild-type (WT) LRRK2 co-expressed in HEK293T cells with WT-PPP2CA. Dimers were labeled with biotin *in situ* and captured on streptavidin-coated ELISA plates for ELISA and in-well activity measurements. Over-expression of PPP2CA reduces dimer formation as well as the intrinsic kinase activity of LRRK2. C) Co-expression of PPP2CA reduces cellular kinase activity of LRRK2 in HEK293T cells, as assessed by auto-phosphorylation at Ser1292 in both total cell extracts as well as isolated dimers (SA eluate, lower panel), and phosphorylation of Rab10 (T73). Additionally, note that PPP2CA co-elutes with dimeric LRRK2 purified on SA-coated resin. Data are means±SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 using unpaired T-test (two-tailed).

Figure 2: PP2A is able to dephosphorylate LRRK2 *in vitro* and *in* cells at T1503. A) Purified His-MBP-RocCOR was incubated together with GST-Roco4 kinase and ATP to ensure RocCOR phosphorylation. The reaction was stopped by addition of the H1152 Roco4 kinase inhibitor and commercially available GST-PPP2CA was added. Phosphorylation of LRRK2 T1503 was determined by Western blotting (n=3). B) Quantification of (A). Shown is the normalized ratio of LRRK2 pT1503 phosphorylation (pT1503 signal divided by the His-MBP-RocCOR signal) . C) LRRK2 pT1503 phosphorylation in A549 cells treated with DMSO or Okadaic acid for 60 minutes. In the presence of Okadaic acid the autophosphorylation of LRRK2 at T1503 is increased compared to cells treated only with DMSO (Okadaic acid solvent). As a control for the immunoprecipitation, no LRRK2 antibody was used for the immunoprecipitation (n=4). D) Quantification of (C). Shown is the ratio of the pT1503 and LRRK2 signal normalized to the DMSO condition. Data are means±SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, using unpaired T-test (two-tailed).

Figure 3: The phosphorylation status of the T1503 residue is essential for LRRK2 dimerization and kinase activity. To determine the impact of auto-phosphorylation at Thr1503-LRRK2 on dimerization and activity, we co-expressed proximity biotinylation construct pairs to isolate dimers of the following composition: WT/WT dimers, WT/T1503A hetero-dimers, or T1503A/T1503A homo-dimers. Dimer formation and intrinsic kinase activity were measured as above using an ELISA-based format. Loss of phosphorylation at T1503 on one of the dimer pairs does not alter dimer formation (A), but slightly reduces intrinsic activity of isolated dimers (B); whereas complete loss of phosphorylation at T1503 in LRRK2 dimers results in de-stabilization of LRRK2 dimers (C) and more substantial loss of kinase activity (D). Data are means±SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 using unpaired T-test (two-tailed).

Figure 4: *In vitro* and *in cellulo* identification data showing LRRK2 phosphorylating PPP2CA at the Thr304 residue. A) GFP-LRRK2-G2019S was co-expressed with NSF-tagged WT-PPP2CA or T340A-PPP2CA in HEK293 cells. The PP2A proteins were isolated by streptavidin pull-down and phosphorylation of Thr304 WT-PPP2CA was visualized by a pThrPro specific antibody B) Purified LRRK2 was incubated with commercially available purified WT-PPP2CA and ATP, with or without MLi-2 inhibitor. The Western-blot results show that purified LRRK2 is phosphorylating PPP2CA at T304, while in the presence of MLi-2, PPP2CA is not phosphorylated (n=3). C) The catalytic PPP2CA and PPP2CB were immunoprecipitated from parental or KO-LRRK2 RAW 264.7 macrophages using the PP2Ac antibody. Phosphorylation was determined by western blotting (n=3). D) Quantification of panel C. Shown is the ratio of the pT304 and PP2Ac signal normalized to the WT condition Data are means±SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, using unpaired T-test (two-tailed).

Figure 5: The regulation of the phosphorylation status of PPP2CA at T304 residue has an effect on the holoenzyme association as well as in its enzymatic activity. A) NSF tagged PPP2R2B or (B) NSF tagged PPP2R5C were co-expressed with the indicated myc-tagged PPP2CA constructs and the interaction was tested by immunoprecipitation. As a positive control, the endogenous catalytic PP2A subunit (anti PP2Ac staining, lower band, panel A), is interacting in the myc-PPP2CA-T304D sample pull down (n=3). C) Quantification of (B). D) PP2A activity of the indicated NSF tagged PPP2CA proteins was measured using an Ser/Thr Phosphatase assay kit (#17-127, Merck). The release of pmoles of phosphate is normalized to WT-PPP2CA . The PPP2CA-T304 mutants have significantly less phosphatase activity compared to WT-PPP2CA (n=2 for biological experiments with a triplicate per biological experiment). Data are means±SEM. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ using unpaired T-test (two-tailed).

Figure 6: The overexpression of WT-PPP2CA, but not T304A-PPP2CA, rescues cortical neurons from G2019S-induced cell death. Primary embryonic mouse cortical neurons were transiently transfected with WT or G2019S-LRRK2 together with empty vector, WT-PPP2CA, T304A-PPP2CA, or T304D-PPP2CA. Three days following transfection, the neurons were fixed and processed for immunofluorescence to label transfected cells (GFP reporter), activated caspase-3 (red, A), Flag (PPP2CA; red, B), and nuclear morphology (DAPI, blue). C) The percentage of transfected neurons with apoptotic features was determined in a blinded manner. The plot shown is representative of 2-3 independent cultures and transfections, and each bar is comprised of at least 3 technical replicates (individual coverslips, with a minimum of 100 neurons counted in each coverslip). Mutation of the T304 site abolishes the neuroprotective capacity of PPP2CA. ** $p < 0.01$; *** $p < 0.001$, using unpaired T-test (two-tailed).

Supplementary Fig 1: Evidence for LRRK2 pT1503. (A) Extracted PRM spectrum for pT1503 for LRRK2 either incubated without or with ROCO4. (B) Extracted PRM spectrum for a reference peptide corresponding to the LRRK2 Roc sequence.

Supplementary Fig. 2: Evidence for the PPP2CA T304 phosphorylation. Mascot-annotated MS2 spectra either for (A) the phosphorylated peptide or (B) unphosphorylated peptide are shown.

Supplementary Fig. 3: LRRK2 is able to dephosphorylate PPP2CA at T304 residue in A549 cells.
A549 lung carcinoma cells were treated with 10nM MLi-2 LRRK2 kinase inhibitor for 90 minutes, prior to cell lysis. Furthermore, the PP2A catalytic subunits were immunoprecipitated. In the absence of the MLi-2 inhibitor, T304 signal of PP2A catalytic subunits is detected, however when the LRRK2 kinase activity is inhibited pharmacologically (by MLI-2), the pT304 signal is diminished

Supplementary Fig. 4: The c-terminal part of PPP2CA is essential for interaction with PPP2R2B subunit. NSF-PPP2R2B or NSF-empty and the indicated myc-tagged PPP2CA constructs were co-transfected in HEK293 cells. Subsequently the cells were lysed and streptavidin beads were used to pull down NSF- PPP2R2B. In addition, we checked whether myc-PPP2CA constructs are in the same complex with NSF- PPP2R2B. While myc-WT-PPP2CA interacts with NSF- PPP2R2B the myc-1-300aa-PPP2CA and myc-1-293aa-PPP2CA proteins are showing a very low interaction affinity with the NSF- PPP2R2B