Planarian behavioral screening is a useful invertebrate model for evaluating
seizurogenic chemicals

Danielle Ireland[a], Evangeline Coffinas[a], Christina Rabeler[a], and Eva-Maria S. Collins[a,b,c,d,*]

[a]Department of Biology, Swarthmore College, 500 College Avenue, Swarthmore, PA 19081,
USA

[b]Department of Physics and Astronomy, Swarthmore College, 500 College Avenue,
Swarthmore, PA 19081, USA

[c]Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania,
3400 Civic Center Blvd, Philadelphia, PA 19104 USA

[d]Center of Excellence in Environmental Toxicology, University of Pennsylvania, 421 Curie
Boulevard Philadelphia, PA 19104 USA

[*]Corresponding author: Email: ecollin3@swarthmore.edu (E-MSC)

Abstract
Detecting adverse health effects of drugs and other chemicals early during chemical/drug
development saves significant time and resources. Freshwater planarians are an emerging
invertebrate model for rapid, cost-effective neurotoxicity screening. Because planarians
exhibit seizure-like behavior when exposed to chemicals that cause seizures in mammals,
such as N-methyl-D-aspartate (NMDA) and picrotoxin, they could be a useful first-tier model
for seizure screening and thus reduce the need for slow and expensive mammalian tests.
However, planarian seizure studies to date have been low-throughput and lacking the
necessary standardization and automated analysis to make this model a viable screening
solution. Here, we present results from medium-throughput behavioral testing conducted in
48-well plates using two popular models for planarian pharmacological and toxicological
studies: *Dugesia japonica* and *Girardia dorotocephala*. Planarian behavior was scored using
automated image analysis, measuring both translational behavior and body shape changes.
We found that known seizurogenic compounds in mammals (NMDA, nicotine, picrotoxin,
pilocarpine, and pentylenetetrazole (PTZ)) induced seizure-like behavior in both planarian

species within 30 minutes of exposure. We also tested three pesticides (parathion, carbaryl,
and permethrin). Parathion and carbaryl, but not permethrin, caused planarian seizure-like
activity. While the planarian species responded similarly to most compounds, some
compounds showed potency differences of 10-100-fold (pilocarpine and nicotine,
respectively). *G. dorotocephala* planarians were generally more sensitive, but *D. japonica*
planarians displayed more reproducible behaviors. By standardizing both experimental
approach and analysis methods and making them available, this work can serve as a
framework for future testing of chemicals for seizurogenic potential in planarians.

Keywords

Dugesia japonica; *Girardia dorotocephala*; seizures; parathion; PTZ; behavior

1. Introduction

Drugs that target the central nervous system (CNS) or non-CNS drugs that can cross the
blood brain barrier may cause adverse outcomes on brain function, such as the induction of
seizures. In fact, a recent study (Behl et al., 2025) reports that failure during seizure liability
testing accounts for 67% of failures of CNS-drugs during pre-clinical drug development,
based on survey results by the Safety Pharmacology Society in 2015 (Authier et al., 2016).
Seizures can be induced through different molecular initiation events (Behl et al., 2025; Jett,
2012; Tukker and Westerink, 2021). For example, the well-known seizurogenic chemicals
pentylenetetrazole (PTZ) and picrotoxin (PTX) cause seizures by inhibiting the $GABA_A$
receptor (Squires et al., 1984) while acute toxic exposure to organophosphorus nerve gases
and pesticides causes seizures due to inhibition of acetylcholinesterase (AChE)
(Kozhemyakin et al., 2010). Thus, seizure liability testing is an important step during pre-
clinical drug development for drugs that enter the CNS or possess structural similarity to
known seizurogenic compounds (Behl et al., 2025; Easter et al., 2009; Roberts et al., 2021).

Detection of potential seizurogenic compounds has traditionally relied on mice and rat *in vivo*
and *ex vivo* models during late stage preclinical studies (Easter et al., 2009; Tukker and
Westerink, 2021). However, there is increasing awareness that testing in rodents is too
expensive and slow to keep up with the demands for chemical screening and may be limited
in its predictive capabilities for human health (Behl et al., 2025; Smirnova et al., 2014; Van
Norman, 2019). Therefore, multiple *in silico* and *in vitro* assays have been developed in

recent years to fill the need to more quickly and efficiently identify seizure liability (Behl et
al., 2025; Belair et al., 2025; Easter et al., 2009; Grainger et al., 2018; Tukker and Westerink,
2021). Specifically, micro-electrode arrays (MEAs), particularly those utilizing human
induced pluripotent stem cell derived neuronal /astrocyte co-cultures, have emerged as a
promising new approach method (Belair et al., 2025; Grainger et al., 2018; Tukker and
Westerink, 2021). In these assays, whole network function can be quantified and specific
changes in neuronal activity have been shown to be indicative of *in vivo* seizures (Ishii et al.,
2017). However, cell systems, including 3D cultures (Pelkonen et al., 2020), lack the
complexity and metabolic capacity necessary to fully evaluate the effects of chemicals on
brain function in a living organism (Hogberg and Smirnova, 2022). Small organismal models
help overcome these challenges as they contain the entire complexity of the nervous system
in its original context and have metabolic capacity, albeit with species-dependent differences,
and thus can bridge between *in vitro* and mammalian models (Collins et al., 2024).
Developing zebrafish allow for rapid and cost-effective screening in an intact organism and
therefore have become a popular complementary test system, especially for rapid screening
of anti-seizure compounds (Baraban et al., 2005; McGraw and Poduri, 2025; Milder et al.,
2022; Whyte-Fagundes et al., 2025).
81
No model can 100% predict human *in vivo* effects and therefore modern toxicology aims to
leverage the evidence gained from multiple complementary models via integrated approaches
to hazard and risk assessment (Hogberg et al., 2022; Jaworska et al., 2016; Rockley et al.,
2023). Freshwater planarians are invertebrates that have been utilized for pharmacological
studies for over a century (Buttarelli et al., 2008; Ireland and Collins, 2022; Raffa, 2008).
Additionally, planarians have recently gained recognition for their amenability to rapid
behavioral testing of toxicants and neuroactive drugs (Ireland et al., 2025a; Ireland and
Collins, 2022, 2023). Planarians are sensitive to chemical exposure and respond with
stereotypical behavioral responses that can be measured (reviewed in (Buttarelli et al., 2008;
Hagstrom et al., 2016; Raffa, 2008)). Chemically-induced behaviors have been qualitatively
described in the planarian literature, but with the exception of "scrunching", a well-
characterized planarian behavior that shows oscillatory body shape changes (Cochet-Escartin
et al., 2015; Sabry et al., 2019), chemically-induced behaviors remain to be quantitatively
defined (Reho et al., 2022). Because planarians live in water, chemical exposure is simple
and behavioral experiments can be conducted easily and inexpensively (Ireland and Collins,
2022). Planarians share key neurotransmitter systems with humans, including GABA,

glutamate, serotonin, dopamine, and acetylcholine (Ross et al., 2017), thus making them a

relevant neurobiology model. Different species have been used for chemical exposure

experiments for different reasons, such as commercial availability, size, and more recently

genome availability and amenability to high-throughput studies in multi-well plates (Ireland

and Collins, 2023; Wu and Li, 2018).

Studies focused specifically on seizure-like behavior in planarians (Figure 1) have primarily

used the commercially available *Girardia* (formerly *Dugesia*) *dorotocephala* (GD) and

Girardia (formerly *Dugesia) tigrina* (GT) species and found that these species are sensitive to

known seizurogenic compounds such as N-methyl-D-aspartate (NMDA) (Rawls et al., 2009),

nicotine (Bach et al., 2016; Pagán et al., 2015; Rawls et al., 2011), and pilocarpine (Miller et

al., 2025). Because it has not yet been shown that the behavioral phenotypes induced by these

compounds are due to seizures through EEG or other means to visualize neuronal activity, we

will refer to them as 'planarian seizure-like activity' (pSLA), a term originally introduced by

(Rawls et al., 2009) and defined as "asynchronous paroxysms (C-shape, twitching behavior)".

Other researchers have also referred to pSLA as planarian seizure-like motility (pSLM)

(Pagán et al., 2013), defined as "vigorous writhing and bending of the body" (Bach et al.,

2016). A clear definition of what these terms encapsulate does not exist but they seem to

denote rapid shape changes – each lasting approximately 1 second (Rawls et al., 2009) - that

include C-shape, writhing, head flailing, dorsal oscillation, corkscrew, among others. Some of

these shapes are well-defined (e.g., corkscrew) and examples have previously been reviewed

(Hagstrom et al., 2016; Wu and Li, 2018), while others are ambiguous, e.g., "dorsal

oscillations" (Miller et al., 2025).

Figure 1. Overview of planarian studies on seizurogenic chemicals in the last 25 years (2000-2025). Tested seizurogenic compounds are listed. Some of the studies also tested other compounds, including therapeutics to treat seizures, which are not listed here. Abbreviations indicate the species tested: GD – *Girardia dorotocephala*; GT – Girardia *tigrina*.

While object tracking has been employed to characterize the spatial behavior of planarians exposed to seizurogenic chemicals (Miller et al., 2025), pSLA thus far has been scored manually, by counting the number of abnormal behavioral changes during a 5-10 minute recording period. Manual pSLA scoring relies on qualitative criteria about shape changes, which may differ between laboratories and even researchers. Moreover, behavioral data is intrinsically noisy due to the biological variability of organisms; thus, standardization of methods to minimize variability and automated pSLA analysis is indispensable to overcome these limitations and guarantee reproducibility and transferability of results.

Here, we present a medium-throughput method for screening chemicals for seizurogenic potential in two planarian species that are popular models in toxicology and pharmacology research: *Dugesia japonica* (DJ) and GD planarians. Screening is performed in 48-well plates and scoring of pSLA is achieved for all 48 planarians simultaneously via automated image analysis using a Motility Index (MI) metric that uses a standard image subtraction algorithm. The MI method is similar to methods routinely used to evaluate zebrafish behavior (Bruni et al., 2016), including specifically for seizures (e.g., (Afrikanova et al., 2013; Whyte-Fagundes et al., 2025)). The 48-well plate format and automated analysis greatly increases the throughput from previous planarian studies, which utilized exposure in petri dishes (e.g.,

(Bach et al., 2016; Rawls et al., 2009)) or 12-well plates (Miller et al., 2025) and used

primarily manual pSLA scoring of individual planarians. Manual scoring of 48 worms

recorded for 30 minutes would take 24 hours compared to ~15 minutes of run time of the

automated computational analysis we present here. Moreover, automated analysis allows for

a standardized, quantitative definition of pSLA that can be used across laboratories.

Using this automated analysis, we quantified the behavior of DJ and GD planarians to 9 test

compounds. Four of the compounds have previously been shown to induce pSLA in some

planarian species (NMDA, nicotine, picrotoxin (PTX), pilocarpine; Figure 1) and were

confirmed as pSLA inducers by our automated analysis, which quantitatively defined pSLA

based on species-specific thresholds in MI scores derived from qualitative scoring of NMDA-

induced pSLA. PTZ, which is used to create rodent and zebrafish seizure models (Baraban et

al., 2005; Dhir, 2012; Gawel et al., 2020; Milder et al., 2022; Shimada and Yamagata, 2018;

Yang et al., 2023), also induced pSLA in both planarian species. We also evaluated allyl

isothiocyanate (AITC) as a control compound that induces scrunching (Cochet-Escartin et al.,

2015; Sabry et al., 2019) and show that our analysis can distinguish it from pSLA. The other

test compounds were three pesticides with seizurogenic potential in mammals, namely two

acetylcholinesterase inhibitors (parathion and carbaryl) and the pyrethroid permethrin (Jett,

2012). Permethrin, tested to the solubility limit, did not cause effects but parathion and

carbaryl induced pSLA, thus demonstrating that pSLA can be triggered through different

pathways in planarians, similar to in mammals.

2. Methods

2.1. Planarian species and maintenance

The freshwater planarian species DJ and GD were used in this study. DJ planarians have been

maintained in the laboratory for > 15 years on organic liver (beef and chicken). GD

planarians were obtained commercially from Carolina Biologicals (Cat #132954) and then

maintained on frozen bloodworms (San Francisco Bay Brand, #88055) in the laboratory. All

species were maintained in planarian water (0.21 g/L Instant Ocean salts [Spectrum Brands,

Blacksburg, VA, USA], 0.83 mM $MgSO_4$, 0.9 mM $CaCl_2$, 0.04 mM $KHCO_3$, 0.9mM

$NaHCO_3$), fed 1-2x per week and cleaned on the day of feeding and 2 days following feeding.

Unless otherwise noted, all experiments used intact, adult planarians which had been fasted

4-9 days. Note that planarians can live for months without any food (McConnell, 1967) and

we did not observe differences in response to the chemicals depending on the fasting state

(Supplemental Figure S1). To determine if a brain is required for pSLA (section 3.6),

planarians of similar size to intact ones were amputated 4 hours prior to chemical exposure

with an ethanol-sterilized razor or scalpel blade between the auricles and pharynx.

2.2. Chemicals and preparation

Chemicals used in this study are listed in Table 1. For chemicals that had previously been

tested in planarian seizure studies, test concentrations were guided by previous literature

(Table 2), testing a range of concentrations around those shown to previously induce pSLA.

For chemicals which had not previously been tested acutely in planarians, initial test

concentration ranges were chosen based on either concentrations previously used in zebrafish

to induce seizures (Afrikanova et al., 2013; Baraban et al., 2005) or concentrations which

induced chronic effects in planarians (Ireland et al., 2022b; Zhang et al., 2019). The reported

concentrations were refined by preliminary experiments to identify a concentration range

spanning no effects to induction of pSLA or the solubility limit of the chemical, and thus

were often shifted for the two planarian species. PTX was only tested at one concentration as

this was the only concentration we found induced pSLA and higher concentrations were

limited by solubility. Depending on water solubility, chemical stock solutions were prepared

either in planarian water or in 100% dimethyl sulfoxide (DMSO) in a fume hood (Table 1).

For all chemicals prepared in DMSO, the final DMSO concentration in the 1x solution for all

concentrations was 0.5% (v/v) and 0.5% (v/v) DMSO was used as the solvent control for

these chemicals. Because DMSO alone can affect planarian behavior (Pagán et al., 2006), we

tested DMSO up to 5% for acute behavioral effects. No overt qualitative behavioral

phenotypes or pSLA were found for DMSO at concentrations <4% in DJ or <5% in GD

(Supplemental Figure S2). Chemicals prepared in planarian water were used the same day,

while stocks made in DMSO were aliquoted and stored at -20□ or -80□ until use. Frozen

stocks were thawed a maximum of 3 times (Kozikowski et al., 2003). Depending on chemical

solubility, chemicals were added as 1x, 2x or 10x solutions to the planarians in the wells

(Table 1).

pH measurements were taken with an Apera PH60-MS pH Tester kit (Apera Instruments,

Columbus, OH) after allowing the probe to stabilize >10 minutes. The pH of the highest

tested concentration of each chemical is listed in Supplemental Table S1. As the tested

concentrations of NMDA were found to have very low pH (3-3.6), we compared NMDA

exposure to planarian water at the same pH. To obtain planarian water at acidic pH, dilute

hydrochloric acid was added until the desired pH was obtained and pH was verified

immediately before use.

Table 1. Overview of chemicals and concentrations used

Chemical	CAS #	Tested concentrations[1]	Vendor	Purity (%)
allyl isothiocyanate (AITC)[2]	57-06-7	25 µM (GD); 100 µM (DJ)	Sigma-Aldrich	95%
carbaryl[3]	63-25-2	10-75 µM	Sigma-Aldrich	≥98.0 %
dimethyl sulfoxide (DMSO)[2]	67-68-5	0.1%-5% (GD); 0.5-5% (DJ)	Sigma-Aldrich	≥99.7%
N-methyl-D-aspartate (NMDA)[4]	6384-92-5	3 mM (GD); 15 mM (DJ)	Tocris Bioscience	≥99% (HPLC)
(–)-nicotine[2]	54-11-5	10-50 µM (GD); 50-1000 µM (DJ)	Sigma-Aldrich	≥99% (GC)
parathion[3]	56-38-2	10-50 µM (GD); 10-75 µM[6] (DJ)	Sigma-Aldrich; ChemService	≥98.0% (HPLC); 98.8% (GC/FID)
pentylenetetrazole (PTZ)[5]	54-95-5	5- 20 mM	Cayman Chemical	≥98%
permethrin (mixture of cis and trans isomers) [3]	52645-53-1	100 µM; 1 mM[6]	Sigma-Aldrich	>97% (29.8% cis, 67.8% trans isomers)
picrotoxin, *Anamirta cocculin* (PTX)[4]	124-87-8	5 mM[6]	Sigma-Aldrich	≥98% (mixture of picrotoxin and picrotin, HPLC)
(+)-pilocarpine hydrochloride[5]	54-71-7	2-6 mM (GD); 6-100 mM (DJ)	Cayman Chemical	≥98%

[1]if the same concentration or concentration range was used in both species, there is no species reference; else, it
is indicated in parentheses which species was tested at this concentration.
[2] indicates highest concentration stock was 10x of the nominal, prepared in planarian water.
[3] indicates highest concentration stock was 200x of the nominal, prepared in DMSO and diluted to an
intermediate 2x or 10x.
[4] indicates highest concentration stock was 1x of the nominal, prepared in planarian water.
[5] indicates highest concentration stock was 2x of the nominal, prepared in planarian water.
[6] indicates solubility issues with the 1x stock in planarian water.

Table 2. Chemicals and concentrations reported to induce pSLA in planarians.

Chemical	Species	Concentration range	References
Nicotine	GT; GD	10-2000 µM (GT) 1-5 mM (GD)	(Bach et al., 2016; Pagán et al., 2015; Ramakrishnan and DeSaer, 2011; Rawls et al., 2011)
Pilocarpine	GD	1,2,3,4,6 mM	(Miller et al., 2025)
PTX	GT	0.01-5 mM	(Ramakrishnan and DeSaer, 2011)

2.3. Plate setup and acute recording

Planarians were loaded into the wells of a tissue culture treated 48-well plate (Genesee
Scientific, El Cajon, CA), one planarian per well, using a p1000 pipettor with the pipette tip
slightly trimmed with a clean razor blade to avoid damage to the planarians. The planarians
were loaded in planarian water and an appropriately concentrated chemical stock was added
such that the total volume of 1x solution in each well equalled 200 µL. Chemical addition
was performed in the fume hood. The chemical stock was added using a multichannel
pipettor (CAPP Aero 12-Channel Pipette, 5-50 µL), the plate was sealed using thermal film
(Excel Scientific, Victorville, CA, USA), and behavior was recorded, with the recordings
beginning 2-6 minutes after chemical addition, including the transfer time between the fume
hood and the recording setup and starting the recording. Four conditions (3 test conditions
and 1 vehicle control) were tested per plate, with each condition comprising two columns of
wells, with n=12 planarians per condition (chemical concentration and worm type) (Figure
2A). Separate plates were tested for each planarian species. If any chemical in the plate was
prepared in DMSO, the vehicle control consisted of 0.5% (v/v) DMSO, else it contained
untreated planarian water. As needed due to loss of worms crawling out of the wells or for
cases with high data dispersion, independent replicates were performed to increase the
sample size. The sample sizes for all experiments are listed in Supplemental Table S3.

Figure 2. Schematic of 48-well plate and behavioral imaging setup. A) Testing was

performed in 48-well plates with one worm per well. Each chemical condition (different

colors) was tested across 2 columns of wells (n=12). Every plate contained one population of

vehicle controls (white wells). Separate plates were tested for each planarian species. Worm

drawing by Mackenzie Malia. B) Custom imaging setup allowing for recording of 2 plates

simultaneously. Details of the components used are provided in Supplemental Table S2.

As shown in Figure 2B, recordings were taken with a custom imaging setup inside a

NEEWER Photo Studio Shooting tent (Amazon.com, Seattle WA) that allows for recording

in a controlled lighting environment. All components used in this set up are described in

Supplemental Table S2. In brief, the plates were set on top of an LED light panel with

brightness set to 1350 to 1450 lux, as measured with a lux meter. A Fresnel lens was placed

on top of the plate to minimize imaging aberrations. While published studies primarily used

5-10 minutes of recording (e.g., (Bach et al., 2016; Pagán et al., 2015; Rawls et al., 2009) and

up to 1 hour (Miller et al., 2025), we chose to record for 30 minutes, except where indicated,

to allow for more sensitivity to detect pSLA that may not be induced immediately. Images

were collected at 5 frames per second (fps) using a Point Grey Flea3 camera (FLIR,

Wilsonville, OR) using Flycap software. This frame rate was chosen because we wanted to

compare pSLA to other hyperactive behaviors, such as scrunching, which requires a

minimum 5 fps frame rate for detection (Cochet-Escartin et al., 2015; Sabry et al., 2020), and

we found that higher frame rates did not improve signal detection (Supplemental Figure S3).

2.4. Automated image analysis

Analysis of acute behavioral recordings was performed in MATLAB (2025a, MathWorks,
Natick, MA) using custom scripts. Any worm that crawled out of the water during recording
was excluded from analysis. To create a standardized image of the background to be using for
masking the individual wells and removing background noise, the average projection of the
first 2 minutes of recording was used. For determination of overall activity and changes in
body shape, we calculated the absolute value of the difference of consecutive frames divided
by the absolute value of the sum of the two images. The difference image was then binarized
using a noise threshold determined from recordings of plates containing water only under the
same imaging conditions. The motility index (MI) was then calculated for each well and
frame as the sum of pixels that changed between the two images above the noise threshold.
The area of each worm was calculated via thresholding of the image following subtraction by
the average projection background image. As optical aberrations during imaging create some
"blind spots" in which the planarian can become lost, any frames where no worm was found
were excluded in the MI calculations. As individual bouts of pSLA have been previously
described as lasting approximately 1 second (Rawls et al., 2009), the median MI score for 1
second (5 frame) bins was calculated. These scores were then normalized by the median
worm area in the first 2 minutes of recording, where the worms are most likely to be moving
the most and thus have elongated shapes.

To correlate normalized MI scores with the presence of pSLA, we manually evaluated the
recording of 10-12 planarians exposed to NMDA (3 mM in GD, 15 mM in DJ) across 3
independent plates to identify a training set of manually determined pSLA events. For each
analyzed worm, 1 minute of recording (60 1-second bins) was chosen, which exhibited high
pSLA qualitatively by eye, and each 1 second bin containing pSLA was selected. For each
species, we then evaluated the distribution of normalized MI scores across all identified
pSLA events (Supplemental Figure S4). The 10^{th} percentile of each species' pSLA
distributions (GD: 0.11; DJ: 0.21) was then used as a threshold, where normalized MI scores
above this threshold were counted as pSLA events.

MI provides information about the general activity of a planarian but gives no context to the
spatial displacement of the worm. Thus, we also quantified the translational activity of each
planarian over 20 seconds (100 frame) bins. This bin size was chosen because a normal
gliding planarian can circle around the outside of a well of a 48-well plate in about this time.
For each bin, the minimum intensity projection was determined. The average projection of
the first 2 minutes of recording was subtracted from the minimum intensity projection to

remove the background. The resulting image was binarized using a pre-determined threshold.

The translational activity of each worm was calculated as the sum of pixels in this filtered

image (thus representing the motion of the worm) divided by the median area of the worm

during the first 2 minutes of recording. All image analysis codes have been deposited on

Zenodo (doi: 10.5281/zenodo.18686049).

2.5. Lethality analysis

To determine if the presence of acute pSLA is correlated with subsequent lethality, some

plates were removed after the initial acute recording and stored in a dark 20 °C incubator

until the following day. Plates were recorded for 5 minutes at 21-27 hours after plate setup.

From the recordings, each planarian was manually scored as alive, dead, "crawl-out" when

the worm crawls out of the solution in the well, or "sick" when the worm was alive/moving

but displayed abnormal morphology, indicated by head regression, pharynx extrusion,

lesions/disintegration or C-shape body postures.

2.6. Statistical analysis

The number of pSLA events, defined by each species threshold (see section 2.4), over the

entire 30 minute recording were counted and are provided in Supplemental File S1. Statistical

analysis was performed in R (4.5.0, (R Core Team, 2016)). For each compound, data were

compiled across all independent replicates and compared with the compiled in plate vehicle

controls (planarian water or 0.5% DMSO) across the same replicates. A negative binomial

generalized linear model was performed using the MASS package (Venables and Ripley,

2002) to determine an effect of condition on the number of pSLA events. Post hoc pairwise

comparisons of the estimated marginal means of the model was performed using the contrast

function of emmeans (Lenth, 2023) using the Benjamini & Hochberg (BH) p-value

adjustment method for multiple comparisons. The median, 25^{th} and 75^{th} percentiles for each

group, and final adjusted p-values for all pairwise comparisons are provided in Supplemental

File S1.

3. Results

3.1. Defining pSLA events using prototypical concentrations of NMDA

The term pSLA was first introduced in 2009 by Rawls and co-authors to describe planarian

behavior that looked convulsive and indicative of seizures in response to NMDA treatment

(Rawls et al., 2009). However, pSLA identification was qualitative and the manual counting

of pSLA is extremely time consuming and susceptible to bias. To overcome these limitations

and thus make rapid screening of potential seizurogenic compounds in freshwater planarians

possible, we developed a pipeline to robustly and quantitatively identify pSLA.

Because we needed a prototypical example of pSLA to build our analytical pipeline, we first

identified a NMDA concentration that qualitatively induced robust pSLA in both planarian

species within 30 minutes of exposure. Using the published data as a guide (Rawls et al.,

2009), we tested 3 mM NMDA and found that this concentration immediately induced

convulsive behavior in GD. However, robust convulsions were not observed in DJ until 15

mM NMDA was used. At these selected concentrations, NMDA-induced various paroxysms,

including head flailing, turning, rapid twitching and convulsions, over-extension, C-shapes,

corkscrews and spiral curling in both species (Supplemental Videos S1 and S2), as previously

described in the literature. Figure 3 shows example high magnification images of the different

shapes and behaviors that we observed in both species and Supplemental Figure S4 show the

progression of behaviors over 15 minutes of recording for a *D. japonica* planarian exposed to

NMDA.

362
Figure 3. High magnification examples of abnormal body shapes and behaviors

observed during NMDA-induced pSLA. Normal gliding behavior (left) as observed in

planarian water exhibits relatively fixed body lengths and straight motion. During NMDA

exposure (3 mM in GD and 15 mM in DJ), planarians change rapidly between different

abnormal body shapes. Scale bars: 0.5 mm.

Next, to allow for rapid behavioral screening of multiple conditions simultaneously, we

developed a medium-throughput screening set-up using 48-well plates as explained in

Methods Section 2.3 and shown in Figure 2. Each well contained one planarian, with 2
columns per treatment condition, for a total of 12 planarians per condition per plate, allowing
for simultaneous testing of 4 conditions in the same plate (Figure 2A). The use of commercial
components (provided in Supplemental Table S2) for the behavioral imaging box (Figure 2B)
allows for replication of this method in other laboratories. Using this set-up, chemically
treated planarians were imaged for 30 minutes immediately (within 2-6 minutes) following
exposure. To quantitatively identify pSLA, we measured the motility index (MI, see Methods
Section 2.4), normalized by worm area, for each planarian over the 30-minute recording
period. As individual bouts of pSLA have been previously described as lasting approximately
1 second (Rawls et al., 2009), we calculated the median MI per 1 second (5 frames) bins for
each worm. Species-specific pSLA thresholds were then determined based on the distribution
of normalized MI scores for a subset of representative NMDA-exposed planarians (see
Methods Section 2.4, Supplemental Figure S5), where normalized MI scores per 1 second bin
above this threshold were identified as potential pSLA events. We then applied this analysis
to all NMDA data.

We found that NMDA showed significantly more pSLA events than the respective controls in
each species (Figure 4A-B). Because these NMDA concentrations had low pH (~3-3.6,
Supplemental Table S1), we also tested the effect of pH alone using planarian water at
approximately the same pH as the respective NMDA concentration. Acidic planarian water
induced significantly more pSLA events than the respective non-adjusted controls, though not
to the same extent as with NMDA. However, in DJ, the number of pSLA events in the acidic
pH sample was not significantly different from NMDA. We conclude that pH is a
contributing factor to NMDA-induced pSLA in both species.

While the total number of pSLA events over the entire recording time allows classification of
a chemical as seizurogenic or not, there is additional information about the chemical's action
in the dynamics of the response, which can be evaluated from the normalized MI scores as a
function of time (Figure 4C-D). In both species, NMDA induced high activity within 0-2
minutes of recording time, which dissipated within ~5 (GD) or ~10 (DJ) minutes. After this
time, NMDA-exposed planarians became contracted and immobile and/or crawled out of the
NMDA solution. As 0 or nearly 0 MI scores are seen with both normal gliding behavior and
immobility, we also quantified the translational activity of the planarians (see Methods
Section 2.4). These translational activity plots show that the decrease in MI scores in NMDA-

exposed planarians is concomitant with decreased motion around the well (Figure 4E-F). This

decreased motility may indicate the toxic nature of NMDA as almost all worms in both

species were dead or had crawled out of their wells by ~24 hours (Supplemental Figure S6).

410
Figure 4. Validation of quantitative analysis of pSLA activity using NMDA. (A-B)

Boxplots of the number of pSLA events per 30 min recording in A) GD and B) DJ exposed to

NMDA and planarian water at the same pH as the respective NMDA concentration. Dots

represent individual planarians. Statistical significance was determined using pairwise

contrasts (with a Benjamini & Hochberg p-value correction) of the estimated marginal means

of a negative binomial generalized linear model of condition versus number of pSLA events.

Different lower-case letters indicate the groups are statistically significantly different (p<0.05). (C-D) Median normalized MI for all C) GD or D) DJ planarians exposed to their prototypical NMDA concentration and their respective in plate vehicle controls as a function of time. (E-F) Median normalized translational activity of all E) GD or F) DJ planarians exposed to their prototypical NMDA concentration and their respective in plate vehicle controls as a function of time. Sample sizes are listed in Supplemental Table S3.

3.2. MI analysis distinguishes pSLA from other hyperactive behaviors

To verify that our analysis method is specific to pSLA, we exposed GD and DJ planarians to AITC, which has been shown to activate the planarian Transient Receptor Potential Ankyrin channel (DjTRPAa) and induce forward scrunching, a periodic muscle-based form of locomotion (Sabry et al., 2019). AITC concentrations that induced scrunching within a few minutes of exposure (25 µM for GD and 100 µM for DJ) had increased normalized MI scores (Figure 5A-B) and showed differences in translational activity compared to controls (Figure 5C-D). However, the MI scores were not high enough to be detected as significant pSLA activity (Figure 5E-F). AITC behavior was only analyzed for 10 minutes following exposure as scrunching did not persist past this time.

Figure 5. Quantitative MI analysis can distinguish pSLA from scrunching behavior induced by AITC. (A-B) Median normalized MI for all A) GD or B) DJ planarians exposed to AITC and their respective in plate vehicle controls as a function of time. Note that in A the control curve is not visible as all scores were 0. (C-D) Median normalized translational activity of all C) GD or D) DJ planarians exposed to AITC and their respective in plate vehicle controls as a function of time. (E-F) Boxplots of the number of pSLA events per 10 min recording in E) GD and F) DJ exposed to AITC and the respective in plate vehicle control. Dots represent individual planarians. No significant differences were found between controls and AITC in each using pairwise contrasts (with a Benjamini & Hochberg p-value correction) of the estimated marginal means of a negative binomial generalized linear model of condition versus number of pSLA events. Note the y-axes are scaled differently in the various plots to best reflect each data set. Sample sizes are listed in Supplemental Table S3.

3.3. Compounds known to induce pSLA in planarians are detected via MI analysis

Next, we tested other compounds that have previously been reported to induce pSLA in planarians (Table 2).

3.3.1 Nicotine

Studies by different research groups have reported hyperactivity and pSLA in GD and GT planarians upon acute nicotine exposure in the µM to mM range (Figure 1 and Table 2). Nicotine binds to and activates nicotinic acetylcholine receptors. While there is supporting pharmacological evidence for nicotinic acetylcholine receptors in GD planarians (Rawls et al., 2011), the receptors have not yet been characterized. No significantly increased pSLA events were observed in GD planarians exposed to 10 – 50 µM nicotine (Figure 6A) although we qualitatively observed hyperactivity and pSLA-like events in GD planarians exposed to 10 µM nicotine immediately following exposure. This hyperactivity was slower and less intense than pSLA seen with 3 mM NMDA, as supported by lower amplitudes in the MI versus time plot for nicotine (Supplemental Figure S7) compared to NMDA (Figure 4C). In contrast to 3 mM NMDA, no lethality was observed at ~24 hours with 10 µM nicotine (Supplemental Figure S6).

Figure 6. MI analysis of known seizurogenic compounds in planarians. Boxplots of the

number of pSLA events per 30 min recording in (A-B) Nicotine, (C-D) Pilocarpine, and E-F)

PTX in GD (left column, A, C, E) and DJ (right column, B, D, F). Dots represent individual

planarians. Sample sizes are listed in Supplemental Table S3. Statistical significance was

determined using pairwise contrasts (with a Benjamini & Hochberg p-value correction) of the

estimated marginal means of a negative binomial generalized linear model of condition

versus number of pSLA events. Different lower-case letters indicate the groups are

statistically significantly different (p<0.05). Note the y-axes are scaled differently in the

various plots to best reflect each data set.

Nicotine-induced pSLA quickly dissipated within the first 2 minutes, followed by a return to

gliding behavior (Supplemental Figure S7 and S8). For these reasons, and because of the

noise in the controls, no statistical significance was found when considering the number of

pSLA events over the entire 30 minutes recording. When we focused on only the first 5 minutes of recording, we found significant pSLA for 10 µM nicotine (Supplemental Figure S9). Thus, our analysis shows that nicotine induces pSLA, in agreement with published data, which only scored 5 minutes of exposure (Ramakrishnan and DeSaer, 2011; Rawls et al., 2011).

Interestingly, pSLA was not observed in higher concentrations (15 and 25 µM) which largely displayed normal gliding. Exposure to 50 µM nicotine resulted in contraction and paralysis over the course of the 30 minutes recording (Supplemental Figure S10A). In contrast, in DJ planarians, significantly increased pSLA was observed starting at 100 µM nicotine, with the most pSLA events observed in 750 µM, which was highly hyperactive and demonstrated NMDA-like pSLA consisting of head flailing, convulsions and dynamic changes between S- and C-shapes (Figure 6B). In 750 µM nicotine, pSLA activity dissipated/slowed after 5 minutes of recording and was replaced by contraction with some sporadic, slow convulsive events (Supplemental Figure S11) and very little translational motion (Supplemental Figure S12). Contraction and further morphological abnormalities such as C-shapes and lesions were observed at ~24 hours of exposure to 750 µM nicotine, although little lethality was observed (Supplemental Figure S6).

3.3.2 Pilocarpine

It has recently been shown that a muscarinic receptor agonist, pilocarpine, can also trigger pSLA in GD . Using particle tracking to quantify the time spent in the center and the periphery of a well of a 12-well plate and manual scoring of different behaviors such as "dorsal oscillations", "head flick", "C-shape", and others, concentrations of 4-6 mM pilocarpine were found to be significantly different from controls (Miller et al., 2025). Thus, we first tested 2-6 mM in GD planarians. Using MI analysis, we found that all tested concentrations of pilocarpine induced statistically significant pSLA in GD (Figure 6C), in agreement with the study by Miller et al. (Miller et al., 2025). No lethality was observed at 24 hours (Supplemental Figure S6). However, DJ planarians only displayed significantly increased pSLA at $\geq$ 40 mM pilocarpine, with the most pSLA events observed in 100 mM pilocarpine (Figure 6D), which also caused 100% lethality at 24 hours (Supplemental Figure S6). Thus, the two species show clear differences in sensitivity to pilocarpine, which may be due to differences in uptake, molecular targets, or metabolism.

3.3.3 PTX

The GABA$_A$ receptor antagonist PTX has been shown to induce pSLA in GT planarians (Ramakrishnan and DeSaer, 2011) but no studies have yet been conducted in GD or DJ planarians. Because of limited water solubility, the highest concentration of PTX that could be tested was 5 mM. We found that 5 mM PTX induced hyperactivity and pSLA within 30 minutes of recording in both species (Figure 6E-F). However, in contrast to the other chemicals tested, which induced pSLA within the first few minutes, 5 mM PTX induced pSLA was only observed after 20-25 minutes of recording, following initial gliding behavior and sporadic bouts of head twitching and dorsal oscillations (Supplemental Figures S7 and S11), suggesting that this concentration of PTX was at the potency threshold. PTX did not induce lethality in either species at ~24 hours (Supplemental Figure S6), but the worms were elongated and were still exhibiting hyperactive behaviors.

3.4. PTZ can induce pSLA

Similar to PTX, PTZ also antagonizes the GABA$_A$ receptor (albeit through a different mechanism) and is frequently used to generate animal seizure models (Afrikanova et al., 2013; Chitolina et al., 2023; Dhir, 2012; Milder et al., 2022; Shimada and Yamagata, 2018; Yang et al., 2023). We therefore tested varying concentrations of PTZ for its ability to induce pSLA in both GD and DJ planarians. We found that concentrations exceeding 10 mM (GD) or 5 mM (DJ) PTZ significantly induced pSLA (Figure 7A-B). PTZ-induced pSLA was characterized by an initial phase of head wiggling and turning, followed by random twitching and convulsions that included longitudinal over-extension and C-shapes, corkscrews and spiral curling. In GD, these hyperactive behaviors slowed down over the 30 minutes of recording and some of the planarians moved into a contracted state, indicating muscle paralysis (Supplemental Figures S7-8). In contrast, in DJ planarians, hyperactive behavior persisted throughout the entire 30-minute recording (Supplemental Figure S11). Both species were impacted at 24 hours with high numbers of death, sickness and crawl-out behavior, though the exact mix of these three abnormal conditions differed across the species (Supplemental Figure S6)

Figure 7. PTZ and parathion induce pSLA in GD and DJ planarians. Boxplots of the number of pSLA events per 30-minute recording in (A-B) PTZ, (C-D) parathion, (E-F) carbaryl, and (G-H) permethrin in GD (left column) and DJ (right column). Dots represent individual planarians. Sample sizes are listed in Supplemental Table S3. Statistical significance was determined using pairwise contrasts (with a Benjamini & Hochberg p-value correction) of the estimated marginal means of a negative binomial generalized linear model

of condition versus number of pSLA events. Different lower-case letters indicate the groups

are statistically significantly different ($p < 0.05$). Note the y-axes are scaled differently in the

various plots to best reflect each data set.

3.5. Some pesticides induce pSLA

Pesticides with different molecular targets have been found to be seizurogenic in mammalian

models (Jett, 2012). Because both nicotine and pilocarpine, two cholinergic receptor

agonists, induced pSLA, we hypothesized that organophosphorus pesticides and carbamates

that inhibit acetylcholinesterase (AChE) in planarians as in humans (Hagstrom et al., 2017)

and can cause seizures in mammals at high concentrations (Jett, 2012), could trigger pSLA

upon acute high concentration exposure. To test this hypothesis, we tested one chemical from

each class, parathion and carbaryl, respectively. Concentrations of 10 µM parathion and

above caused significant pSLA in both planarian species (Figure 7C-D). While 50 µM caused

the most pSLA events in both species, 31.6 µM parathion was a better concentration to

induce pSLA in GD due to the high number of crawl-out events seen in this species at 50 µM.

Following ~24 hour exposure to these concentrations of parathion, both species showed

sickness and lethality (Supplemental Figure S6). Carbaryl induced much weaker pSLA events

in both species, which were characterized by small contractions and head twitches. While 25

µM showed significantly increased pSLA in DJ planarians compared to vehicle controls, 10-

75 µM carbaryl showed no significant effect on the number of pSLA in GD planarians

(Figure 7E-F), despite 25 µM showing increased median MI scores per bin (Supplemental

Figure S7). At 75 µM, some scrunching and head twitching was observed within the first 5

minutes of recording, following by decreased movement and contraction (Supplemental

Figure S10B), suggesting that this concentration exceeded the pSLA-inducing range of

concentrations. No significant lethality was observed in either species exposed to 25 µM

carbaryl at ~24 hours (Supplemental Figure S6).

Some pyrethroid insecticides, such as permethrin, have been shown to cause seizures in

rodents at lethal concentrations (Staatz and Hosko, 1985) or in environmentally relevant

concentrations in cats (Dymond and Swift, 2008). Pyrethoids inhibit voltage-gated sodium

channels, leading to neuronal hyper-excitability and at high doses can inhibit GABA

signalling (Jett, 2012). We tested permethrin up to its solubility limit (1 mM) and did not

observe significant pSLA in either planarian species (Figure 7G-H). Additionally, no obvious

differences in behavior could be seen in the MI or translational activity plots (Supplemental

Figures S7, S8, S11, S12).

3.6. The planarian brain is not required to induce pSLA in DJ planarians

It had previously been shown that pSLA in headless GT planarians was similar to that in

intact planarians when exposed to nicotine but not to cocaine, another pSLA inducer (Pagán

et al., 2012). Cocaine-induced pSLA was nearly completely lost in regenerating tails and was

gradually restored during regeneration (Pagán et al., 2013). These results suggest that cocaine

sensitive receptors are primarily found in the brain, consistent with dopamine receptors in

vertebrates. In contrast, nicotinic acetylcholine receptors must be distributed in both the head

and the body because the response in headless planarians was similar to that of intact

planarians. Therefore, comparing responses of intact and headless planarians to seizurogenic

chemicals may provide insights into where the molecular targets of these chemicals are

expressed in planarians. Thus, we tested whether the presence of the brain was required for

pSLA by exposing regenerating DJ tails to each seizurogenic compound at a concentration

shown to significantly induce pSLA in adult/intact DJs. Exposure began 4 h after amputation

to allow for wound closure and to mitigate the increased mucus production following

amputation (Peiris et al., 2014) that may hinder chemical uptake. All chemicals except for

carbaryl induced pSLA in regenerating DJ tails (Table 3).

Table 3. **Median number of pSLA in adult or regenerating tail DJ planarians exposed to**

representative concentrations of seizurogenic compounds. Sample sizes are listed in

Supplemental Table S3.

Condition	Adult Median # pSLA (25th, 75th percentile)	Regenerating Median # pSLA (25th, 75th percentile)
15 mM NMDA	42 (6, 63.5)	45 (33, 68)
pH 3	14 (5, 21)	69 (24, 132)*
750 µM nicotine	96 (83, 152)	29 (11, 55)*
100 mM pilocarpine	37 (15,88)	69 (43, 201)
5 mM PTX	12 (6, 15)	15 (10, 21)
15 mM PTZ	75 (49 , 100)	150 (93, 197)
50 µM parathion	394 (194, 470)	59 (37, 136)*
25 µM carbaryl	3 (2, 13)	0 (0,1)

607 * non-overlapping interquartile ranges between adult and regenerating planarians for the same chemical
608

Some compounds, including NMDA, pilocarpine, PTX, and PTZ induced similar amounts of

pSLA in the two developmental stages with overlapping interquartile ranges. Nicotine and

parathion showed significantly more active behaviors - evidenced by higher numbers of

pSLA - in adult DJ planarians than in tails, suggesting that the presence of the brain plays a

role for pSLA induction in DJ planarians. Interestingly, pH 3 showed significantly greater
activity in the regenerating tails than in adults. Thus, in DJ planarians, while the seizurogenic
compounds do not strictly require the planarian brain to induce pSLA, presence of the brain
can affect pSLA dynamics depending on the compound. Carbaryl was at the detection
threshold and thus a comparison may not be meaningful.
618

4. Discussion
4.1. MI analysis can classify seizurogenic compounds and distinguish pSLA from other
behaviors
To overcome the limitations associated with manual scoring of pSLA, namely scoring bias
and time effort, we created an automated MI analysis method that uses image subtraction to
detect changes in body shape. Using frame subtraction to quantify activity is a commonly
used method to quantify behaviors including seizures in other model systems such as
zebrafish (Afrikanova et al., 2013; Bruni et al., 2016; Whyte-Fagundes et al., 2025). By using
qualitative scoring of NMDA-induced pSLA as a framework, we derived a quantitative MI
threshold for pSLA in GD and DJ planarians. This analysis method was able to correctly
identify known seizurogenic chemicals in planarians (NMDA, nicotine, PTX, pilocarpine)
and distinguish pSLA from other abnormal behaviors, such as forward scrunching, which – in
contrast to pSLA – exhibits periodic changes in body shape (Cochet-Escartin et al., 2015).
Complementary analysis of the translational worm activity as a function of time helped us to
distinguish normal behavior from contraction and paralysis, which both show low MI scores.
634

We confirmed previous studies by other research groups that NMDA (Rawls et al., 2009),
nicotine (Rawls et al., 2011), and pilocarpine (Miller et al., 2025) induce pSLA in GD
planarians and showed that they also induce pSLA in DJ planarians. PTX had previously only
been studied in GT planarians (Ramakrishnan and DeSaer, 2011); here, we show that it also
induces pSLA in GD and DJ planarians. The concentrations of NMDA, pilocarpine, and PTX
that induced pSLA in GD planarians here were comparable to previous studies using GD or
GT planarians. However, we found that 3 mM NMDA lowered pH to a range where pH alone
caused pSLA while previous work found that 3 mM NMDA did not lower the pH as much
(pH 6.6), and thus pH alone did not induce pSLA (Rawls et al., 2009). The discrepancy in
these data may be due to differences in the exact water composition as the previous study
used Artificial Pond Water (APW; 6 mM NaCl; 0.1 mM NaHCO$_3$; 0.6 mM CaCl$_2$; pH 7.3)

(Rawls et al., 2009) which has slightly different salt composition compared to our planarian

water (see Methods Section 2.1) and thus may have different buffering capacity.

For nicotine, we found that 50 µM caused paralysis in GD planarians, comparable to

concentrations used in GT planarians (Ramakrishnan and DeSaer, 2011). In contrast, other

studies in GD planarians have used nicotine in the mM range (Bach et al., 2016; Pagán et al.,

2015, 2013; Rawls et al., 2011). The difference is likely be due to the product used. While we

and (Ramakrishnan and DeSaer, 2011) used pure (-) nicotine solution, the other studies (Bach

et al., 2016; Pagán et al., 2015, 2013; Rawls et al., 2011) used nicotine hydrogen tartrate salt.

The pesticides parathion and carbaryl are both cholinesterase inhibitors and also induced

pSLA in at least one planarian species whereas the pyrethroid permethrin, which inhibits

voltage-gated sodium channels, did not.

Finally, we showed that PTZ, which is used to generate seizure models in mice (Dhir, 2012;

Shimada and Yamagata, 2018) and zebrafish (Afrikanova et al., 2013; Baraban et al., 2005;

Chitolina et al., 2023; Milder et al., 2022), robustly induces pSLA in both planarian species.

Thus, PTZ could be used to generate a planarian seizure-like model to test the efficacy of

possible anti-seizure treatments. In fact, existing studies that used compounds other than PTZ

suggest that pSLA in planarians can be modulated by exposure to known anti-convulsant

therapeutics (e.g., (Pagán et al., 2015; Ramakrishnan and DeSaer, 2011; Rawls et al., 2009)).

Now that we have developed this automated platform, it can be used to validate existing

manual studies on therapeutic effectiveness.

4.2 GD and DJ planarians respond similarly to chemicals but with different sensitivities

GD and DJ planarians displayed very different sensitivities to NMDA, nicotine, and

pilocarpine (Table 3, Supplemental Tables S4-S5). GD planarians were more sensitive, with

the most extreme difference in sensitivity of nearly 100-fold for nicotine. While NMDA and

pilocarpine induced similar behavioral phenotypes in the two species, there were also

qualitative differences in the behaviors displayed by the two species in response to nicotine as

DJ planarians exhibited extensive pSLA events that were never observed in GD planarians.

Other studies have reported similar behavioral and sensitivity differences between planarian

species in response to certain chemicals (e.g., (Ireland et al., 2020; Sabry et al., 2019)).

Together, these data emphasize that extrapolation between planarian species is not

recommended due to possible differences in pharmacokinetics (chemical uptake, metabolism)

and/or pharmacodynamics (presence of targets/receptors). Further mechanistic studies will be

needed to better understand these species-specific differences.

We found that DJ planarians are more behaviorally heterogeneous than DJ planarians. These

differences may be due to the DJ planarians being bred under controlled laboratory

conditions with standardized food and culture conditions for years whereas GD planarians

had only been maintained under these conditions for a few weeks in our study. Changes and

differences in diet could also contribute to the behavioral differences (Pacis et al., 2025). The

variability in GD behavior was the primary reason for reduced statistical power in some of

the exposure conditions (Supplemental Table S6). However, the increased sensitivity of GD

planarians to some of the compounds tested is a strength for screening seizurogenic

chemicals. More compounds will need to be screened in both species to evaluate their

strengths and weaknesses for this application. A good starting point would be to test

chemicals that have been proposed to induce pSLA in planarians in recent manual studies,

such as sertraline (Thumé and Frizzo, 2017) and various cholinesterase inhibitors (Bezerra

Da Silva et al., 2016), which were not included here.

4.3 The brain is not required for pSLA

Most seizurogenic chemicals induced pSLA in headless DJ planarians to a similar extent as in

intact adults, indicating that the cephalic ganglion is not required for pSLA induction by the

compounds tested here. The planarian CNS is distributed and includes both the cephalic

ganglion and two ventral nerve cords (VNCs) that extend along the head–tail axis (Ross et

al., 2017). Thus, hyperexcitation of neural circuits within the VNCs may be sufficient to

generate pSLA. This parallels observations in mammals that spinal cord circuits can

independently support seizure-like activity. For example, GABA$_A$ receptor blockade with

PTX induces epileptiform activity in mouse spinal neurons *in vitro* (Barker and MacDonald,

1980). These findings suggest that neural circuits can generate hyperexcitability even without

brain involvement.

Given that planarians reproduce asexually through binary fission, autonomous motor control

of a headless body fragment is essential for survival of tail offspring (Le et al., 2021). The

persistence of pSLA following decapitation may therefore reflect functional redundancy

within this distributed neural architecture rather than the absence of a neural origin.

Consistent with this view, planarian behavioral responses are complex and many motor

behaviors — including turning, rectifying, and scrunching — can occur in the absence of the
head (Cochet-Escartin et al., 2015; Le et al., 2021), indicating that the VNCs can
independently coordinate motor output. However, our data do not allow us to distinguish
whether the observed effects arise from central versus peripheral nervous system
mechanisms, nor whether they reflect neuronal hyperactivity or direct activation of muscle.
Future mechanistic studies (see 4.5.3) will be needed to distinguish these scenarios.
720
Interestingly, parathion and nicotine both produced significantly more pSLA events in intact
adult *D. japonica* than in headless planarians, suggesting that the brain enhances or
modulates the behavioral response to these compounds. Although cholinesterases (Hagstrom
et al., 2018) and nicotinic receptors (Pagán et al., 2013) are distributed throughout the
planarian body, cholinesterase expression is highest in the brain, indicating that cholinergic
signalling may be more strongly regulated there. The cephalic ganglion may therefore
amplify or coordinate cholinergic network activity, leading to more robust pSLA when intact.
Consistent with a modulatory role for the brain, nicotine exposure previously produced fewer
pSLA events in decapitated GT planarians, although the effect was not statistically significant
(Pagán et al., 2013). Differences between that study and ours may reflect species differences,
nicotine formulation, or analytical methods. Notably, cocaine-induced pSLA has also been
reported to require the presence of the brain (Pagán et al., 2013), further supporting the idea
that at least some substances depend on cephalic neural integration to produce maximal
behavioral output. Because our analysis compared only the medians and interquartile ranges
of the total number of pSLA events over the 30-minute recording, differences in response
dynamics between intact and headless planarians would not have been detected and may
warrant future investigation.
738

4.4 Use of planarians for screening seizurogenic chemicals and therapeutics
Planarians offer practical advantages for early seizure liability screening. Although they are
simpler than zebrafish and lack a circulatory system and blood–brain barrier (Rawls et al.,
2010), they are inexpensive to maintain and, as invertebrates, do not require the animal
welfare protocols mandated for vertebrate models like zebrafish (Collins et al., 2024).
Zebrafish larvae at 5 days post fertilization (dpf) are primarily used for screening when
experiments are not classified as animal testing (Strähle et al., 2012) but their nervous system
and metabolic machinery are still maturing (Menezes et al., 2014; Saad et al., 2017; Tiedeken
and Ramsdell, 2010). Seizure models have also been developed for *Caenorhabditis elegans*,

but rely on specific genetic strains and have largely been used for identification of anti-
seizure therapeutics (Jones et al., 2021; Wong et al., 2018). We envision a drug discovery
pipeline where, following *in silico* chemical modelling and/or human *in vitro* MEA assays,
compounds can be screened in adult planarians with metabolic capacity (Ireland et al., 2022a)
and mature neuronal function, before subsequent testing in developing zebrafish and
preclinical rodent models. This tiered screening approach would only prioritize the best lead
candidates at each stage, ultimately reducing cost and vertebrate use in early seizure liability
testing. We have previously shown that organismal planarian and zebrafish behavioral
screening and MEA assays were the most informative to identify developmental
neurotoxicants (Ireland et al., 2025b) and thus may prove to be a promising combination for
identify seizurogenic chemicals and anti-seizure therapeutics.

All models have limitations, some of which are unique and some which are shared across
models. The strengths and limitations of planarians and developing zebrafish (<5 dpf) for
chemical screening have recently been reviewed (Collins et al., 2024). One limitation shared
by organismal models and cell culture models is the requirement for aqueous chemical
exposure, making poorly water-soluble or volatile compounds difficult to test (Collins et al.,
2024). As a result, sufficiently high concentrations to induce pSLA may not always be
achievable, as illustrated by our ability to test PTX only at its seizure induction threshold. *In*
silico modelling can help overcome some experimental limitations, as has been shown
recently for screening anti-seizure drugs in zebrafish larvae (Belair et al., 2025; Cassar et al.,
2017). Comparative testing of compounds known to be seizurogenic in humans across these
models can clarify their complementarity within a rapid seizure liability test battery and
identify data gaps and limitations, similar to ongoing efforts in developmental neurotoxicity
testing (Ireland et al., 2025b; Masjosthusmann et al., 2020).

4.5 Limitations and Opportunities
The work here can act as a starting point for future screens of seizurogenic chemicals and
anti-convulsants in planarians but further optimization and system validation are required to
decide whether planarian screening could be a valuable part of an early-stage seizure liability
testing battery. Three major tasks still remain: 1) optimized experimentation to improve
throughput, 2) optimized image analysis and 3) mechanistic validation of pSLA.

4.5.1 Optimize experimental throughput

The experimental approach described here provides a proof-of-concept design that should be
further optimized to fit within the needs of a preclinical drug discovery pipeline. The
recording period could be shortened to 15 minutes to achieve higher throughput as most
chemicals induced pSLA within the first 5-10 minutes. However, it should be recognized this
may decrease detection power for some chemicals. Our recordings began 2-6 minutes
following chemical exposure due to experimental constraints in starting recording
immediately, which was sufficient for the compounds we studied (Supplemental Figure S13).
This window could be decreased to increase the possibility of catching very short-lasting
pSLA events. Here, we used a minimum of n=12 specimen per condition, similar to or greater
than the sample sizes used in previous planarian seizure studies (Bach et al., 2016; Miller et
al., 2025; Rawls et al., 2009). While these sample sizes were generally sufficient to achieve
adequate statistical power, intrinsic variability in control behavior of GD planarians required
larger sample sizes in some cases (Supplemental Table S6). Therefore, while the column
setup that we used here, utilizing n=12 within a single experiment, was helpful for testing a
few selected compounds for this proof-of-concept study, for new screens, we recommend a
row setup, similar to what is used for rapid behavioral screening in other studies (Ireland et
al., 2025a; Zhang et al., 2019), because it allows for the simultaneous testing of 5
concentrations/conditions plus a vehicle control. By using 2 plates with alternating row setup,
one can control for potential edge effects (Zhang et al., 2019) and achieve n=16 per condition
while testing in independent experiments to ensure robustness. In addition to acute behavioral
recordings, we also evaluated lethality one day after chemical exposure to determine if
lethality was correlated with the presence of significant pSLA. This was not the case. Thus,
lethality likely results from other toxic effects besides pSLA and thus is not an informative
endpoint for this type of study. To accelerate system evaluation and have multiple research
groups participate in characterizing the value of planarian behavioral screening, the recording
and analysis methods developed here are made available to other laboratories. This will
ensure standardized conditions that allow cross-laboratory data comparisons and integration.
809

4.5.2 Optimize automated image analysis
The MI analysis pipeline described here provides an automated, quantitative way to detect
pSLA that improves on previous manual techniques that were not well defined and tedious.
One limitation of MI analysis is that it has difficulty to detect pSLA when the
convulsive/twitching behavior is small and/or primarily in the third dimension (e.g., in the
case of carbaryl), which is not well captured in the simple 2D imaging setup we used. While

these behaviors could still be detected as pSLA in DJ planarians, they were not found to be
significant for GD planarians because GD planarians showed more variable control behavior,
including swimming, scrunching, and turning over. Future work could incorporate more
sophisticated image analysis methods, incorporating machine learning to identify specific
body shape changes and use them to characterize pSLA, as has recently been done in
zebrafish (Whyte-Fagundes et al., 2025). Such a tailored approach would likely be more
robust than MI analysis and may reveal subtypes of pSLA, which could provide additional
insight into the underlying biological mechanisms. For example, the initial behavioral effects
of parathion are thought to be due to acetylcholinesterase inhibition while later effects are
from secondary neurotransmitter systems such as glutamate (Jett, 2012; Kozhemyakin et al.,
2010); thus, the type of convulsions may differ over time. Data analysis could thus also
consider the temporal dynamics of the behavioral response by evaluating effects over
different time bins.

4.5.3 Mechanistic validation of pSLA as a model for human seizure activity

For planarians to be used as a screening system for seizure liability, it is crucial to confirm
that pSLA is indicative of seizures using electrophysiological measurements of neuronal
activity. Some electrophysiological techniques have been developed in planarians (Aoki et
al., 2009; Benita et al., 2024; Freiberg et al., 2023), with varying levels of invasiveness.
However, existing techniques are relatively low throughput and not well suited to chemical
exposure. Alternatively, MEA has been successfully used to measure neuronal activity *in vivo*
in zebrafish larvae (Tomasello and Sive, 2020) and could be adapted for planarians. In
addition, RNA interference (RNAi) could be used to verify that the known seizurogenic
compounds tested here induce pSLA through the same pathways as in mammals. For
example, PTX and PTZ both target the $GABA_A$ receptor, thus making it a good starting point
for mechanistic studies. Both parathion and carbaryl induced pSLA in DJ planarians and
inhibit DJ cholinesterases (Hagstrom et al., 2017; Ireland et al., 2022b); thus, RNAi
knockdown of the planarian cholinesterases could be used to verify that the molecular
initiation event of pSLA in planarians is cholinesterase inhibition, as in humans (Jett, 2012).
These verification studies will be critical for adoption of the planarian model for screening
seizurogenic chemicals as they would show that results found in planarians are relevant and
translatable to mammals and humans.

In summary, this work demonstrates that pSLA can be triggered through various pathways in
GD and DJ planarians, similar to in mammals. By standardizing both experimental approach
and analysis methods and making the tools available to the research community, this work
can serve as a framework for future studies of chemicals for their seizurogenic potential in
planarians.

CRediT authorship contribution statement
Danielle Ireland: Conceptualization; Software; Visualization; Validation; Formal analysis;
Data curation; Writing – original draft; Writing - review and editing. **Evangeline Coffinas:**
Investigation; Writing – review and editing. **Christina Rabeler:** Investigation; Writing –
review and editing. **Eva-Maria S. Collins:** Conceptualization; Software; Methodology;
Visualization; Validation; Writing – original draft; Writing – review and editing; Supervision;
Resources; Project administration.

Funding
This work was supported by the National Institutes of Health CounterACT program [grant
number 5R21NS132528-02]. The content is solely the responsibility of the authors and does
not necessarily represent the official views of the National Institutes of Health.

Conflict of interest statement
EMSC is the founder of Inveritek, LLC, which offers planarian HTS commercially. EMSC
and DI are inventors on a patent for planarian rapid screening. The remaining authors declare
that the research was conducted in the absence of any commercial or financial relationships
that could be construed as a potential conflict of interest.

Data availability
The datasets supporting the conclusions of this article are included within the article and
available from the corresponding author upon reasonable request. The MATLAB code used
for image analysis is available on Zenodo (doi: 10.5281/zenodo.18686049).

Acknowledgements

The authors thank Kate Sun and Erin Szuromi for their help with experiments, Dr. Scott Rawls, Dr. Hilary McCarren, and Mathieu Conroy for discussions, Mackenzie Malia for the worm schematic in Figure 2A and Hannah Poon for comments on the manuscript. During the preparation of this work, the authors used ChatGPT (OpenAI) in order to help design R code for the power analysis presented in Supplemental Table S4. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article .

References

Afrikanova, T., Serruys, A.S.K., Buenafe, O.E.M., Clinckers, R., Smolders, I., de Witte, P.A.M., Crawford, A.D., Esguerra, C.V., 2013. Validation of the Zebrafish Pentylenetetrazol Seizure Model: Locomotor versus Electrographic Responses to Antiepileptic Drugs. PLoS ONE 8, 54166. https://doi.org/10.1371/journal.pone.0054166

Aoki, R., Wake, H., Sasaki, H., Agata, K., 2009. Recording and spectrum analysis of the planarian electroencephalogram. Neuroscience 159, 908–914. https://doi.org/10.1016/j.neuroscience.2008.11.011

Authier, S., Arezzo, J., Delatte, M.S., Kallman, M.-J., Markgraf, C., Paquette, D., Pugsley, M.K., Ratcliffe, S., Redfern, W.S., Stevens, J., Valentin, J.-P., Vargas, H.M., Curtis, M.J., 2016. Safety pharmacology investigations on the nervous system: An industry survey. J Pharmacol Toxicol Methods 81, 37–46. https://doi.org/10.1016/j.vascn.2016.06.001

Bach, D.J., Tenaglia, M., Baker, D.L., Deats, S., Montgomery, E., Pagán, O.R., 2016. Cotinine antagonizes the behavioral effects of nicotine exposure in the planarian Girardia tigrina. Neuroscience Letters 632, 204–208. https://doi.org/10.1016/j.neulet.2016.09.005

Baraban, S.C., Taylor, M.R., Castro, P.A., Baier, H., 2005. Pentylenetetrazole induced changes in zebrafish behavior, neural activity and c-fos expression. Neuroscience 131, 759–768. https://doi.org/10.1016/j.neuroscience.2004.11.031

Barker, J.L., MacDonald, J.F., 1980. Picrotoxin Convulsions Involve Synaptic and Nonsynaptic Mechanisms on Cultured Mouse Spinal Neurons. Science 208, 1054–1056. https://doi.org/10.1126/science.7375918

Behl, M., Karmaus, A., Rao, M., Lane, T., Harris, J., Sachs, C., Borrel, A., Oyetade, O., Unnikrishnan, A., Hamm, J., Hogberg, H.T., 2025. De-Risking Seizure Liability: Integrating Adverse Outcome Pathways (AOPs), New Approach Methodologies (NAMs) and In Silico Approaches while Highlighting Knowledge Gaps. Toxicological Sciences 207, 306–319. https://doi.org/10.1093/toxsci/kfaf109

Belair, D.G., Kohnken, R., McCloud, R.L., Sandoval, S., Green, J., Buck, W.R., Polakowski, J.S., 2025. Predicting seizure liability of small molecules using an in vitro multi-electrode array based assay coupled with modeling of brain disposition. Current Research in Toxicology 8, 100236. https://doi.org/10.1016/j.crtox.2025.100236

Benita, O., Nesher, N., Shomrat, T., 2024. Neurophysiological measurements of planarian brain activity: a unique model for neuroscience research. Biology Open 13, bio060480. https://doi.org/10.1242/bio.060480

Bezerra Da Silva, C., Pott, A., Elifio-Esposito, S., Dalarmi, L., Fialho Do Nascimento, K., Moura Burci, L., De Oliveira, M., De Fátima Gaspari Dias, J., Maria Warumby Zanin,

S., Gomes Miguel, O., Dallarmi Miguel, M., 2016. Effect of Donepezil, Tacrine,
Galantamine and Rivastigmine on Acetylcholinesterase Inhibition in Dugesia tigrina.
Molecules 21, 53. https://doi.org/10.3390/molecules21010053

Bruni, G., Rennekamp, A.J., Velenich, A., McCarroll, M., Gendelev, L., Fertsch, E., Taylor,
J., Lakhani, P., Lensen, D., Evron, T., Lorello, P.J., Huang, X.-P., Kolczewski, S.,
Carey, G., Caldarone, B.J., Prinssen, E., Roth, B.L., Keiser, M.J., Peterson, R.T.,
Kokel, D., 2016. Zebrafish behavioral profiling identifies multitarget antipsychotic-
like compounds. Nature Chemical Biology 12, 559–566.
https://doi.org/10.1038/nchembio.2097

Buttarelli, F.R., Pellicano, C., Pontieri, F.E., 2008. Neuropharmacology and behavior in
planarians: Translations to mammals. Comparative Biochemistry and Physiology Part
C: Toxicology & Pharmacology 147, 399–408.
https://doi.org/10.1016/j.cbpc.2008.01.009

Cassar, S., Breidenbach, L., Olson, A., Huang, X., Britton, H., Woody, C., Sancheti, P.,
Stolarik, D., Wicke, K., Hempel, K., LeRoy, B., 2017. Measuring drug absorption
improves interpretation of behavioral responses in a larval zebrafish locomotor assay
for predicting seizure liability. J Pharmacol Toxicol Methods 88, 56–63.
https://doi.org/10.1016/j.vascn.2017.07.002

Chitolina, R., Gallas-Lopes, M., Reis, C.G., Benvenutti, R., Stahlhofer-Buss, T., Calcagnotto,
M.E., Herrmann, A.P., Piato, A., 2023. Chemically-induced epileptic seizures in
zebrafish: A systematic review. Epilepsy Res 197, 107236.
https://doi.org/10.1016/j.eplepsyres.2023.107236

Cochet-Escartin, O., Mickolajczk, K.J., Collins, E.-M.S., 2015. Scrunching: a novel escape
gait in planarians. Physical Biology 12, 055001.
https://doi.org/doi:%2010.1088/1478-3975/12/5/056010

Collins, E.M.S., Hessel, E.V.S., Hughes, S., 2024. How neurobehavior and brain
development in alternative whole-organism models can contribute to prediction of
developmental neurotoxicity. NeuroToxicology 102, 48–57.
https://doi.org/10.1016/j.neuro.2024.03.005

Dhir, A., 2012. Pentylenetetrazol (PTZ) kindling model of epilepsy. Curr Protoc Neurosci
Chapter 9, Unit9.37. https://doi.org/10.1002/0471142301.ns0937s58

Dymond, N., Swift, I., 2008. Permethrin toxicity in cats: a retrospective study of 20 cases.
Aust Veterinary J 86, 219–223. https://doi.org/10.1111/j.1751-0813.2008.00298.x

Easter, A., Bell, M.E., Damewood, J.R., Redfern, W.S., Valentin, J.-P., Winter, M.J., Fonck,
C., Bialecki, R.A., 2009. Approaches to seizure risk assessment in preclinical drug
discovery. Drug Discovery Today 14, 876–884.
https://doi.org/10.1016/j.drudis.2009.06.003

Freiberg, J., Lang, L., Kaernbach, C., Keil, J., 2023. Characterization of the planarian surface
electroencephalogram. BMC Neurosci 24, 29. https://doi.org/10.1186/s12868-023-
00799-z

Gawel, K., Langlois, M., Martins, T., Van Der Ent, W., Tiraboschi, E., Jacmin, M., Crawford,
A.D., Esguerra, C.V., 2020. Seizing the moment: Zebrafish epilepsy models.
Neuroscience & Biobehavioral Reviews 116, 1–20.
https://doi.org/10.1016/j.neubiorev.2020.06.010

Grainger, A.I., King, M.C., Nagel, D.A., Parri, H.R., Coleman, M.D., Hill, E.J., 2018. In vitro
Models for Seizure-Liability Testing Using Induced Pluripotent Stem Cells. Front
Neurosci 12, 590. https://doi.org/10.3389/fnins.2018.00590

Hagstrom, D., Cochet-Escartin, O., Collins, E.-M.S., 2016. Planarian brain regeneration as a
model system for developmental neurotoxicology. Regeneration 3, 65–77.
https://doi.org/10.1002/reg2.52

Hagstrom, D., Hirokawa, H., Zhang, L., Radić, Z., Taylor, P., Collins, E.-M.S., 2017.
Planarian cholinesterase: in vitro characterization of an evolutionarily ancient enzyme
to study organophosphorus pesticide toxicity and reactivation. Archives of Toxicology
91, 2837–2847. https://doi.org/10.1007/s00204-016-1908-3
Hagstrom, D., Zhang, S., Ho, A., Tsai, E.S., Radić, Z., Jahromi, A., Kaj, K.J., He, Y., Taylor,
P., Collins, E.M.S., 2018. Planarian cholinesterase: molecular and functional
characterization of an evolutionarily ancient enzyme to study organophosphorus
pesticide toxicity. Archives of Toxicology 92, 1161–1176.
https://doi.org/10.1007/s00204-017-2130-7
Hogberg, H.T., Hsieh, J.-H., Chang, X., Sipes, N.S., Shafer, T., Behl, M., 2022. Case study on
the use of Integrated Approaches for Testing and Assessment for DNT to prioritize a
class of Organophosphorus flame retardants, Series on Testing and Assessment.
Organisation for the Economic Cooperation and Development Environment
Directorate, Paris.
Hogberg, H.T., Smirnova, L., 2022. The Future of 3D Brain Cultures in Developmental
Neurotoxicity Testing. Front Toxicol 4, 808620.
https://doi.org/10.3389/ftox.2022.808620
Ireland, D., Bochenek, V., Chaiken, D., Rabeler, C., Onoe, S., Soni, A., Collins, E.-M.S.,
2020. Dugesia japonica is the best suited of three planarian species for high-
throughput toxicology screening. Chemosphere 253, 126718.
https://doi.org/10.1016/j.chemosphere.2020.126718
Ireland, D., Collins, E.-M.S., 2022. New Worm on the Block: Planarians in
(Neuro)Toxicology. Current Protocols 2, e637. https://doi.org/10.1002/CPZ1.637
Ireland, D., Collins, E.S., 2023. Planarians as a model to study neurotoxic agents, in:
Alternative Methods in Neurotoxicology, Alternative Methods in Neurotoxicology.
Elsevier Inc., pp. 29–60. https://doi.org/10.1016/bs.ant.2023.01.002
Ireland, D., Rabeler, C., Gong, T., Collins, E.-M.S., 2022a. Bioactivation and detoxification
of organophosphorus pesticides in freshwater planarians shares similarities with
humans. Archives of Toxicology. https://doi.org/10.1007/s00204-022-03387-y
Ireland, D., Rabeler, C., Rao, S., Richardson, R.J., Collins, E.-M.S., 2025a. Distinguishing
classes of neuroactive drugs based on computational physicochemical properties and
experimental phenotypic profiling in planarians. PLoS ONE 20, e0315394.
https://doi.org/10.1371/journal.pone.0315394
Ireland, D., Word, L.J., Collins, E.-M.S., 2025b. Statistical analysis of multi-endpoint
phenotypic screening increases sensitivity of planarian neurotoxicity testing.
Toxicological Sciences 208, 104–121. https://doi.org/10.1093/toxsci/kfaf117
Ireland, D., Zhang, S., Bochenek, V., Hsieh, J.-H., Rabeler, C., Meyer, Z., Collins, E.-M.S.,
2022b. Differences in neurotoxic outcomes of organophosphorus pesticides revealed
via multi-dimensional screening in adult and regenerating planarians. Frontiers in
Toxicology 4, 948455. https://doi.org/10.3389/ftox.2022.948455
Ishii, M.N., Yamamoto, K., Shoji, M., Asami, A., Kawamata, Y., 2017. Human induced
pluripotent stem cell (hiPSC)-derived neurons respond to convulsant drugs when co-
cultured with hiPSC-derived astrocytes. Toxicology 389, 130–138.
https://doi.org/10.1016/j.tox.2017.06.010
Jaworska, J., Corsini, E., Basketter, D., 2016. Integrated Testing Strategies for Skin
Sensitization Hazard and Potency Assessment—State of the Art and Challenges.
Cosmetics 2016, Vol. 3, Page 16 3, 16. https://doi.org/10.3390/COSMETICS3020016
Jett, D.A., 2012. Chemical toxins that cause seizures. NeuroToxicology 33, 1473–1475.
https://doi.org/10.1016/j.neuro.2012.10.005

Jones, A., Marson, A.G., Cunliffe, V.T., Sills, G.J., Morgan, A., 2021. C. elegans as a
Potential Model for Acute Seizure-Like Activity, in: Vohora, D. (Ed.), Experimental
and Translational Methods to Screen Drugs Effective Against Seizures and Epilepsy,
Neuromethods. Springer US, New York, NY, pp. 233–248.
https://doi.org/10.1007/978-1-0716-1254-5_13
Kozhemyakin, M., Rajasekaran, K., Kapur, J., 2010. Central Cholinesterase Inhibition
Enhances Glutamatergic Synaptic Transmission. Journal of Neurophysiology 103,
1748–1757. https://doi.org/10.1152/jn.00949.2009
Kozikowski, B.A., Burt, T.M., Tirey, D.A., Williams, L.E., Kuzmak, B.R., Stanton, D.T.,
Morand, K.L., Nelson, S.L., 2003. The Effect of Freeze/Thaw Cycles on the Stability
of Compounds in DMSO. SLAS Discovery 8, 210–215.
https://doi.org/10.1177/1087057103252618
Le, D., Sabry, Z., Chandra, A., Kristan, W.B., Collins, E.M.S., 2021. Planarian fragments
behave as whole animals. Current Biology 31, 5111-5117.e4.
https://doi.org/10.1016/j.cub.2021.09.056
Lenth, R.V., 2023. emmeans: Estimated Marginal Means, aka Least-Squares Means.
Masjosthusmann, S., Blum, J., Bartmann, K., Dolde, X., Holzer, A., Stürzl, L., Keßel, E.H.,
Förster, N., Dönmez, A., Klose, J., Pahl, M., Waldmann, T., Bendt, F., Kisitu, J.,
Suciu, I., Hübenthal, U., Mosig, A., Leist, M., Fritsche, E., 2020. Establishment of an
a priori protocol for the implementation and interpretation of an in vitro testing
battery for the assessment of developmental neurotoxicity. EFSA Supporting
Publications 17. https://doi.org/10.2903/SP.EFSA.2020.EN-1938
McConnell, J.V., 1967. A manual of psychological experimentation on planarians. Planarian
Press, Ann Arbor, Michigan.
McGraw, C.M., Poduri, A., 2025. Machine learning enables high-throughput, low-replicate
screening for novel anti-seizure targets and compounds using combined movement
and calcium fluorescence in larval zebrafish. Eur J Pharmacol 991, 177327.
https://doi.org/10.1016/j.ejphar.2025.177327
Menezes, F.P., Rico, E.P., Da Silva, R.S., 2014. Tolerance to seizure induced by kainic acid is
produced in a specific period of zebrafish development. Prog Neuropsychopharmacol
Biol Psychiatry 55, 109–112. https://doi.org/10.1016/j.pnpbp.2014.04.004
Milder, P.C., Zybura, A.S., Cummins, T.R., Marrs, J.A., 2022. Neural Activity Correlates
With Behavior Effects of Anti-Seizure Drugs Efficacy Using the Zebrafish
Pentylenetetrazol Seizure Model. Front. Pharmacol. 13, 836573.
https://doi.org/10.3389/fphar.2022.836573
Miller, T., Vu, J., Tran, C.J., Ritvik, A., Cruz-Ham, K., Haq, K., Musto, B., Musto, A.E.,
2025. Planarian as an Animal Model for Experimental Acute Seizure. JoVE e67307,
67307. https://doi.org/10.3791/67307
Pagán, O.R., Baker, D., Deats, S., Montgomery, E., Tenaglia, M., Randolph, C., Kotturu, D.,
Tallarida, C., Bach, D., Wilk, G., Rawls, S., Raffa, R.B., 2012. Planarians in
pharmacology: parthenolide is a specific behavioral antagonist of cocaine in the
planarian Girardia tigrina. The International journal of developmental biology 56,
193–6.
Pagán, O.R., Deats, S., Baker, D., Montgomery, E., Wilk, G., Tenaglia, M., Semon, J., 2013.
Planarians require an intact brain to behaviorally react to cocaine, but not to react to
nicotine. Neuroscience 246, 265–270.
https://doi.org/10.1016/j.neuroscience.2013.05.010
Pagán, O.R., Montgomery, E., Deats, S., Bach, D., Baker, D., 2015. Evidence of nicotine-
induced, curare-insensitive, behavior in planarians. Neurochemical Research 40,
2087–2090. https://doi.org/10.1007/s11064-015-1512-6

Pagán, O.R., Rowlands, A.L., Urban, K.R., 2006. Toxicity and behavioral effects of
dimethylsulfoxide in planaria. Neuroscience Letters 407, 274–278.
Peiris, T.H., Hoyer, K.K., Oviedo, N.J., 2014. Innate immune system and tissue regeneration
in planarians: An area ripe for exploration. Seminars in Immunology 26, 295–302.
https://doi.org/10.1016/j.smim.2014.06.005
Pelkonen, A., Mzezewa, R., Sukki, L., Ryynänen, T., Kreutzer, J., Hyvärinen, T., Vinogradov,
A., Aarnos, L., Lekkala, J., Kallio, P., Narkilahti, S., 2020. A modular brain-on-a-chip
for modelling epileptic seizures with functionally connected human neuronal
networks. Biosens Bioelectron 168, 112553.
https://doi.org/10.1016/j.bios.2020.112553
R Core Team, 2016. R: A language environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.
Raffa, R.B., 2008. Planaria: A Model for Drug Action and Abuse, 0 ed. CRC Press.
https://doi.org/10.1201/9781498713597
Ramakrishnan, L., DeSaer, C., 2011. Carbamazepine inhibits distinct chemoconvulsant-
induced seizure-like activity in Dugesia tigrina. Pharmacology Biochemistry and
Behavior 99, 665–670. https://doi.org/10.1016/j.pbb.2011.06.003
Rawls, S.M., Patil, T., Tallarida, C.S., Baron, S., Kim, M., Song, K., Ward, S., Raffa, R.B.,
2011. Nicotine behavioral pharmacology: Clues from planarians. Drug and Alcohol
Dependence 118, 274–279. https://doi.org/10.1016/j.drugalcdep.2011.04.001
Rawls, S.M., Patil, T., Yuvasheva, E., Raffa, R.B., 2010. First evidence that drugs of abuse
produce behavioral sensitization and cross-sensitization in planarians. Behavioural
pharmacology 21, 301. https://doi.org/10.1097/FBP.0B013E32833B0098
Rawls, S.M., Thomas, T., Adeola, M., Patil, T., Raymondi, N., Poles, A., Loo, M., Raffa,
R.B., 2009. Topiramate antagonizes NMDA- and AMPA-induced seizure-like activity
in planarians. Pharmacology Biochemistry and Behavior 93, 363–367.
https://doi.org/10.1016/j.pbb.2009.05.005
Reho, G., Lelièvre, V., Cadiou, H., 2022. Planarian nociception: Lessons from a scrunching
flatworm. Frontiers in Molecular Neuroscience 0, 394.
https://doi.org/10.3389/FNMOL.2022.935918
Roberts, R., Authier, S., Mellon, R.D., Morton, M., Suzuki, I., Tjalkens, R.B., Valentin, J.-P.,
Pierson, J.B., 2021. Can We Panelize Seizure? Toxicol Sci 179, 3–13.
https://doi.org/10.1093/toxsci/kfaa167
Rockley, K., Roberts, R., Jennings, H., Jones, K., Davis, M., Levesque, P., Morton, M., 2023.
An integrated approach for early in vitro seizure prediction utilizing hiPSC neurons
and human ion channel assays. Toxicol Sci 196, 126–140.
https://doi.org/10.1093/toxsci/kfad087
Ross, K.G., Currie, K.W., Pearson, B.J., Zayas, R.M., 2017. Nervous system development
and regeneration in freshwater planarians. Wiley Interdisciplinary Reviews:
Developmental Biology 6, 1–26. https://doi.org/10.1002/wdev.266
Saad, M., Matheeussen, A., Bijttebier, S., Verbueken, E., Pype, C., Casteleyn, C., Van
Ginneken, C., Apers, S., Maes, L., Cos, P., Van Cruchten, S., 2017. In vitro CYP-
mediated drug metabolism in the zebrafish (embryo) using human reference
compounds. Toxicol In Vitro 42, 329–336. https://doi.org/10.1016/j.tiv.2017.05.009
Sabry, Z., Ho, A., Ireland, D., Rabeler, C., Cochet-Escartin, O., Collins, E.M.S., 2019.
Pharmacological or genetic targeting of Transient Receptor Potential (TRP) channels
can disrupt the planarian escape response. PLoS ONE 14, e0226104.
https://doi.org/10.1371/journal.pone.0226104

Sabry, Z., Rabeler, C., Ireland, D., Bayingana, K., Collins, E.M.S., 2020. Planarian scrunching as a quantitative behavioral readout for noxious stimuli sensing. Journal of Visualized Experiments 2020, 1–18. https://doi.org/10.3791/61549

Shimada, T., Yamagata, K., 2018. Pentylenetetrazole-Induced Kindling Mouse Model. J Vis Exp 56573. https://doi.org/10.3791/56573

Smirnova, L., Hogberg, H.T., Leist, M., Hartung, T., 2014. Food for Thought …: Developmental neurotoxicity – Challenges in the 21(st) century and in vitro opportunities. ALTEX 31, 129–156. https://doi.org/10.14573/altex.1403271

Squires, R.F., Saederup, E., Crawley, J.N., Skolnick, P., Paul, S.M., 1984. Convulsant potencies of tetrazoles are highly correlated with actions on GABA/benzodiazepine/picrotoxin receptor complexes in brain. Life Sciences 35, 1439–1444. https://doi.org/10.1016/0024-3205(84)90159-0

Staatz, C.G., Hosko, M.J., 1985. Effect of pyrethroid insecticides on EEG activity in conscious, immobilized rats. Pesticide Biochemistry and Physiology 24, 231–239. https://doi.org/10.1016/0048-3575(85)90133-6

Strähle, U., Scholz, S., Geisler, R., Greiner, P., Hollert, H., Rastegar, S., Schumacher, A., Selderslaghs, I., Weiss, C., Witters, H., Braunbeck, T., 2012. Zebrafish embryos as an alternative to animal experiments—A commentary on the definition of the onset of protected life stages in animal welfare regulations. Reproductive Toxicology 33, 128–132. https://doi.org/10.1016/J.REPROTOX.2011.06.121

Thumé, I.S., Frizzo, M.E., 2017. Sertraline Induces Toxicity and Behavioral Alterations in Planarians. BioMed Research International 2017, 1–8. https://doi.org/10.1155/2017/5792621

Tiedeken, J.A., Ramsdell, J.S., 2010. Zebrafish Seizure Model Identifies p , p' -DDE as the Dominant Contaminant of Fetal California Sea Lions That Accounts for Synergistic Activity with Domoic Acid. Environ Health Perspect 118, 545–551. https://doi.org/10.1289/ehp.0901301

Tomasello, D.L., Sive, H., 2020. Noninvasive Multielectrode Array for Brain and Spinal Cord Local Field Potential Recordings from Live Zebrafish Larvae. Zebrafish 17, 271–277. https://doi.org/10.1089/zeb.2020.1874

Tukker, A.M., Westerink, R.H.S., 2021. Novel test strategies for in vitro seizure liability assessment. Expert Opin Drug Metab Toxicol 17, 923–936. https://doi.org/10.1080/17425255.2021.1876026

Van Norman, G.A., 2019. Limitations of Animal Studies for Predicting Toxicity in Clinical Trials: Is it Time to Rethink Our Current Approach? JACC: Basic to Translational Science 4, 845. https://doi.org/10.1016/J.JACBTS.2019.10.008

Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S, Fourth. ed. Springer, New York.

Whyte-Fagundes, P., Efromson, J., Vance, A., Carpenter, S., Bègue, A., Carroll, A., Doman, T.J.J., Harfouche, M., Baraban, S.C., 2025. Automated detection of complex zebrafish seizure behavior at scale. Commun Biol 8, 872. https://doi.org/10.1038/s42003-025-08310-6

Wong, S.Q., Jones, A., Dodd, S., Grimes, D., Barclay, J.W., Marson, A.G., Cunliffe, V.T., Burgoyne, R.D., Sills, G.J., Morgan, A., 2018. A Caenorhabditis elegans assay of seizure-like activity optimised for identifying antiepileptic drugs and their mechanisms of action. J Neurosci Methods 309, 132–142. https://doi.org/10.1016/j.jneumeth.2018.09.004

Wu, J.P., Li, M.H., 2018. The use of freshwater planarians in environmental toxicology studies: Advantages and potential. Ecotoxicology and Environmental Safety 161, 45–56. https://doi.org/10.1016/j.ecoenv.2018.05.057

Yang, C.-S., Wu, M.-C., Lai, M.-C., Wu, S.-N., Huang, C.-W., 2023. Identification of New
Antiseizure Medication Candidates in Preclinical Animal Studies. Int J Mol Sci 24,
13143. https://doi.org/10.3390/ijms241713143
Zhang, S., Hagstrom, D., Hayes, P., Graham, A., Collins, E.-M.S., 2019. Multi-behavioral
endpoint testing of an 87-chemical compound library in freshwater planarians.
Toxicological Sciences 167, 26–44. https://doi.org/10.1093/toxsci/kfy145
1181