Planarian behavioral screening is a useful invertebrate model for evaluating seizurogenic

chemicals

Danielle Ireland[1], Evangeline Coffinas[1], Christina Rabeler[1], and Eva-Maria S. Collins[1,2,3,4]

[1]Swarthmore College, Biology, 500 College Avenue, Swarthmore, PA, 19081, USA

[2]Department of Physics and Astronomy, Swarthmore College, Swarthmore, Pennsylvania,

United States of America

[3]Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA, USA

[4]Center of Excellence in Environmental Toxicology, University of Pennsylvania, Philadelphia,

Pennsylvania, United States of America

Highlights

- Classic seizurogenic chemicals induce convulsions in planarians

- A 48-well exposure paradigm allows for rapid testing of seizurogenic compounds

- Automated analysis replaces manual scoring of seizure-like planarian behavior

- Acute parathion exposure causes seizure-like activity in planarians

Keywords

Dugesia japonica; *Dugesia dorotocephala*; seizures; parathion; PTZ; behavior

Abstract

Detecting adverse health effects of drugs and other chemicals early during chemical/drug

development saves significant time and resources. Freshwater planarians are an emerging

invertebrate model for rapid, cost-effective neurotoxicity screening. Because planarians exhibit

seizure-like behavior when exposed to chemicals that cause seizures in mammals, such as N-

methyl-D-aspartate (NMDA) and picrotoxin, they could be a useful first-tier model for seizure

screening and thus reduce the need for slow and expensive mammalian tests. However,

planarian seizure studies to date have been low-throughput and lacking the necessary

standardization and automation to make this model a viable screening solution. Here, we

present results from medium-throughput behavioral testing conducted in 48-well plates using

two popular models for planarian pharmacological and toxicological studies: *Dugesia japonica*

and *Dugesia dorotocephala*. Planarian behavior was scored using automated image analysis,

measuring both translational behavior and body shape changes. We found that known

seizurogenic compounds in mammals (NMDA, nicotine, picrotoxin, pilocarpine, and

pentylenetetrazole (PTZ)) induced seizure-like behavior in both planarian species within 30

min of exposure. We also tested three pesticides (parathion, carbaryl, and permethrin).

Parathion and carbaryl, but not permethrin, caused planarian seizure-like activity. While the

planarian species responded similarly to most compounds, some compounds showed potency

differences of 10-100-fold (pilocarpine and nicotine, respectively). *D. dorotocephala*

planarians were generally more sensitive, but *D. japonica* planarians displayed more

reproducible behaviors. By standardizing both experimental approach and analysis methods

and making them available, this work can serve as a framework for future testing of chemicals

for seizurogenic potential in planarians.

1. Introduction

Drugs that target the central nervous system (CNS) or non-CNS drugs that can cross the blood

brain barrier may cause adverse outcomes on brain function, such as the induction of seizures.

In fact, a recent study (Behl et al., 2025) reports that failure during seizure liability testing

accounts for 67% of failures of CNS-drugs during pre-clinical drug development, based on

survey results by the Safety Pharmacology Society in 2015 (Authier et al., 2016). Seizures can

be induced through different molecular initiation events (Behl et al., 2025; Jett, 2012; Tukker

and Westerink, 2021). For example, the well-known seizurogenic chemicals pentylenetetrazole

(PTZ) and picrotoxin (PTX) cause seizures by inhibiting the $GABA_A$ receptor (Squires et al.,

1984) while acute toxic exposure to organophosphorus nerve gases and pesticides causes

seizures due to inhibition of acetylcholinesterase (AChE) (Kozhemyakin et al., 2010). Thus,

seizure liability testing is an important step during pre-clinical drug development for drugs that

enter the CNS or possess structural similarity to known seizurogenic compounds (Behl et al.,

2025; Easter et al., 2009; Roberts et al., 2021).

Detection of potential seizurogenic compounds has traditionally relied on mice and rat *in vivo*
and *ex vivo* models in during late stage preclinical studies (Easter et al., 2009; Tukker and
Westerink, 2021). However, there is increasing awareness that testing in rodents is too
expensive and slow to keep up with the demands for chemical screening and may be limited in
its predictive capabilities for human health (Behl et al., 2025; Smirnova et al., 2014; Van
Norman, 2019). Therefore, multiple *in silico* and *in vitro* assays have been developed in recent
years to fill the need to more quickly and efficiently identify seizure liability (Behl et al., 2025;
Belair et al., 2025; Easter et al., 2009; Grainger et al., 2018; Tukker and Westerink, 2021).
Specifically, micro-electrode arrays (MEAs), particularly those utilizing human induced
pluripotent stem cell derived neuronal /astrocyte co-cultures, have emerged as a promising new
approach method (Belair et al., 2025; Grainger et al., 2018; Tukker and Westerink, 2021). In
these assays, whole network function can be quantified and specific changes in neuronal
activity have been shown to be indicative of *in vivo* seizures (Ishii et al., 2017). However, cell
systems, including 3D cultures (Pelkonen et al., 2020), lack the complexity and metabolic
capacity necessary to fully evaluate the effects of chemicals on brain function in a living
organism (Hogberg and Smirnova, 2022). Developing zebrafish allow for rapid and cost-
effective screening in an intact organism and therefore have become a popular complementary
test system, especially for rapid screening of anti-seizure compounds (McGraw and Poduri,
2025; Milder et al., 2022; Whyte-Fagundes et al., 2025).

No model can 100% predict human *in vivo* effects and therefore modern toxicology aims to
leverage the evidence gained from multiple complementary models via integrated approaches
to hazard and risk assessment (Hogberg et al., 2022; Jaworska et al., 2016; Rockley et al.,
2023). Freshwater planarians are invertebrates that have been utilized for pharmacological
studies for over a century (Buttarelli et al., 2008; Ireland and Collins, 2022; Raffa, 2008).
Planarians have recently gained recognition for their amenability to rapid behavioral testing of
toxicants and neuroactive drugs (Ireland et al., 2025a; Ireland and Collins, 2022, 2023).
Planarians are sensitive to chemical exposure and respond with stereotypical behavioral
responses that can be measured. Because planarians live in water, chemical exposure is facile
and behavioral experiments can be conducted easily and inexpensively (Ireland and Collins,
2022). Planarians share key neurotransmitter systems with humans, including GABA,
glutamate, serotonin, dopamine, and acetylcholine (Ross et al., 2017), thus making them a
relevant neurobiology model. Different species have been used for chemical exposure
experiments for different reasons, such as commercial availability, size, and more recently

genome availability and amenability to high-throughput studies in multi-well plates (Ireland
and Collins, 2023; Wu and Li, 2018).

Studies focused specifically on seizure-like behavior in planarians (**Figure 1**) have primarily
used the commercially available *Dugesia dorotocephala* (DD) and *Dugesia tigrina* (DT)
species and found that these species are sensitive to known seizurogenic compounds such as
N-methyl-D-aspartate (NMDA) (Rawls et al., 2009), nicotine (Bach et al., 2016; Pagán et al.,
2015; Rawls et al., 2011), and pilocarpine (Miller et al., 2025). When exposed to these
compounds, planarians display stereotypical behaviors. Because it has not yet been shown that
the behavioral phenotypes induced by these compounds are due to seizures through EEG or
other means to visualize neuronal activity, we will refer to them as 'planarian seizure-like
activity (pSLA)', a term originally introduced by (Rawls et al., 2009) and defined as
"asynchronous paroxysms (C-shape, twitching behavior)".

Figure 1. Overview of planarian studies on seizurogenic chemicals in the last 25 years
(2000-2025). Tested seizurogenic compounds are listed. Some of the studies also tested other
compounds, including therapeutics to treat seizures, which are not listed here. Abbreviations
indicate the species tested: DD – *Dugesia dorotocephala*; DT – *Dugesia tigrina*.

Other researchers have referred to pSLA as planarian seizure-like motility (pSLM) (Pagán et
al., 2013), defined as "vigorous writhing and bending of the body" (Bach et al., 2016). A clear
definition of what these terms encapsulate does not exist but they seem to denote rapid shape
changes – each lasting approximately 1 second (Rawls et al., 2009) - that include C-shape,

writing, head flailing, dorsal oscillation, corkscrew, among others. Some of these shapes are well-defined (e.g., corkscrew), while others are ambiguous, e.g., "dorsal oscillations" (Miller et al., 2025). While object tracking has been employed to characterize the spatial behavior of planarians exposed to seizurogenic chemicals (Miller et al., 2025), pSLA thus far has been scored manually, by counting the number of abnormal behavioral changes during a 5-10 min recording period. Manual pSLA scoring relies on qualitative criteria about shape changes, which may differ between laboratories and even researchers. Moreover, behavioral data is intrinsically noisy due to the biological variability of organisms; thus, standardization of methods to minimize variability and automated pSLA analysis is indispensable to overcome these limitations and guarantee reproducibility and transferability of results.

Here, we present a medium-throughput method for screening chemicals for seizurogenic potential in two planarian species that are popular models in toxicology and pharmacology research: *Dugesia japonica* (DJ) and DD planarians. Screening is performed in 48-well plates. Scoring of pSLA is obtained from automated image analysis using a Motility Index (MI) metric that uses a standard image subtraction algorithm, similar to methods routinely used to evaluate zebrafish behavior (Bruni et al., 2016), including specifically for seizures (e.g., (Afrikanova et al., 2013; Whyte-Fagundes et al., 2025)). Using this method, we quantified the behavior of DJ and DD planarians to 9 test compounds. Four of the compounds have previously been shown to induce pSLA in some planarian species (NMDA, nicotine, picrotoxin, pilocarpine; Figure 1) and were confirmed as pSLA inducers by our automated analysis. PTZ, which is used to create rodent and zebrafish seizure models (Dhir, 2012; Gawel et al., 2020; Milder et al., 2022; Shimada and Yamagata, 2018; Yang et al., 2023), also induced pSLA in both planarian species. We also evaluated allyl isothiocyanate (AITC) as a control compound that induces "scrunching", a well-characterized planarian behavior that shows oscillatory body shape changes (Cochet-Escartin et al., 2015; Sabry et al., 2019), and show that our analysis can distinguish it from pSLA. The other test compounds were three pesticides with seizurogenic potential in mammals, namely two acetylcholinesterase inhibitors (parathion and carbaryl) and the pyrethroid permethrin (Jett, 2012). Permethrin, tested to the solubility limit, did not cause effects but parathion and carbaryl induced pSLA, thus demonstrating that pSLA can be triggered through different pathways in planarians, similar to the situation in mammals.

2. Methods

2.1. Planarian species and maintenance

The freshwater planarian species DJ and DD were used in this study. DJ planarians have been maintained in the laboratory for > 15 years on organic liver (beef and chicken). DD planarians were obtained commercially from Carolina Biologicals (Cat #132954) and then maintained on frozen bloodworms (San Francisco Bay Brand, #88055) in the laboratory. All species were maintained in planarian water (0.21 g/L Instant Ocean salts [Spectrum Brands, Blacksburg, VA, USA], 0.83 mM $MgSO_4$, 0.9 mM $CaCl_2$, 0.04 mM $KHCO_3$, 0.9mM $NaHCO_3$), fed 1-2x per week and cleaned on the day of feeding and 2 days following feeding. Adult intact planarians which had been fasted 4-9 days were selected for experiments. If amputated planarians were used in an experiment alongside intact planarians, they were selected to be of similar size to the matched intact ones. Amputations were performed 4 hours prior to chemical exposure with an ethanol-sterilized razor or scalpel blade between the auricles and pharynx.

2.2. Chemicals and preparation

Chemicals used in this study are listed in Table 1. Depending on water solubility, chemical stock solutions were prepared either in planarian water or in 100 % dimethyl sulfoxide (DMSO) in a fume hood. For all chemicals prepared in DMSO, the final DMSO concentration in the 1x solution was 0.5%. Because DMSO alone can affect planarian behavior (Pagán et al., 2006), we tested DMSO up to 5% for acute behavioral effects. No overt qualitative behavioral phenotypes or pSLA were found for DMSO at concentrations <4% in DJ or <5% in DD (Supplemental Figure S1). Chemicals prepared in planarian water were used the same day, while stocks made in DMSO were aliquoted and stored at -20°C or -80°C until use. Frozen stocks were thawed a maximum of 3 times. Depending on chemical solubility, chemicals were added as 1x, 2x or 10x solutions to the planarians in the wells (Table 1).

pH measurements were taken with an Apera PH60-MS pH Tester kit (Apera Instruments, Columbus, OH) after allowing the probe to stabilize >10 minutes. The pH of the highest tested concentration of each chemical is listed in Supplemental Table S1. To obtain planarian water at acidic pH, dilute hydrochloric acid was added until the desired pH was obtained and pH was verified immediately before use.

Table 1. Overview of chemicals and concentrations used

Chemical	CAS #	Tested concentrations[1]	Vendor	Purity (%)

allyl isothiocyanate (AITC)[2]	57-06-7	25 µM (DD); 100 µM (DJ)	Sigma-Aldrich	95%
carbaryl[3]	63-25-2	10-75 µM	Sigma-Aldrich	≥98.0 %
dimethyl sulfoxide (DMSO)[2]	67-68-5	0.1%-5% (DD); 0.5-5% (DJ)	Sigma-Aldrich	≥99.7%
N-methyl-D-aspartate (NMDA)[4]	6384-92-5	3 mM (DD); 15 mM (DJ)	Tocris Bioscience	≥99% (HPLC)
(−)-nicotine[2]	54-11-5	10-50 µM (DD); 50-1000 µM (DJ)	Sigma-Aldrich	≥99% (GC)
parathion[3]	56-38-2	10-50 µM (DD); 10-75 µM[6] (DJ)	Sigma-Aldrich; ChemService	≥98.0% (HPLC); 98.8% (GC/FID)
pentylenetetrazole (PTZ)[5]	54-95-5	5- 20 mM	Cayman Chemical	≥98%
permethrin (mixture of cis and trans isomers) [3]	52645-53-1	100 µM; 1 mM[6]	Sigma-Aldrich	>97% (29.8% cis, 67.8% trans isomers)
picrotoxin, *Anamirta cocculin* (PTX)[4]	124-87-8	5 mM[6]	Sigma-Aldrich	≥98% (mixture of picrotoxin and picrotin, HPLC)
(+)-pilocarpine hydrochloride[5]	54-71-7	2-6 mM (DD); 6-100 mM (DJ)	Cayman Chemical	≥98%

[1]if the same concentration or concentration range was used in both species, there is no species reference; else, it is indicated in parentheses which species was tested at this concentration.
[2] indicates highest concentration stock was 10x of the nominal, prepared in planarian water.
[3] indicates highest concentration stock was 200x of the nominal, prepared in DMSO and diluted to an intermediate 2x or 10x.
[4] indicates highest concentration stock was 1x of the nominal, prepared in planarian water.
[5] indicates highest concentration stock was 2x of the nominal, prepared in planarian water.
[6] indicates solubility issues with the 1x stock in planarian water.

2.3. Plate setup and acute recording

Planarians were loaded into the wells of a tissue culture treated 48-well plate (Genesee Scientific, El Cajon, CA), one planarian per well, using a p1000 pipettor with the pipette tip slightly trimmed with a clean razor blade to avoid damage to the planarians. The planarians were loaded in planarian water and an appropriately concentrated chemical stock was added such that the total volume of 1x solution in each well equalled 200 µL. Chemical addition was performed in the fume hood. The chemical stock was added using a multichannel pipettor (CAPP Aero 12-Channel Pipette, 5-50 µL), the plate was sealed using thermal film (Excel

Scientific, Victorville, CA, USA), and behavior was recorded, with the recordings beginning
2-6 minutes after chemical addition. Four conditions (3 test conditions and 1 vehicle control)
were tested per plate, with each condition comprising two columns of wells, with n=12
planarians per condition (Figure 2A). If any chemical in the plate was prepared in DMSO, the
vehicle control consisted of 0.5% DMSO, else it contained untreated planarian water. As
needed, independent replicates were performed to increase the sample size.

Figure 2. Schematic of 48-well plate and behavioral imaging setup. A) Testing was
performed in 48-well plates with one worm per well. Each chemical condition (different colors)
was tested across 2 columns of wells (n=12). Every plate contained one population of vehicle
controls (white wells). B) Custom imaging setup allowing for recording of 2 plates
simultaneously. Details of the components used are provided in Supplemental Table S2.

As shown in Figure 2B, recordings were taken with a custom imaging setup inside a NEEWER
Photo Studio Shooting tent (Amazon.com, Seattle WA) that allows for recording in a controlled
lighting environment. All components used in this set up are described in Supplemental Table
S2. In brief, the plates were set on top of an LED light panel with brightness set to 1350 to
1450 lux, as measured with a lux meter. A Fresnel lens was placed on top of the plate to
minimize imaging aberrations. Recordings were 30 minutes in length, except where indicated,
and taken at 5 frames per second using a Point Grey Flea3 camera (FLIR, Wilsonville, OR)
using Flycap software. This frame rate was chosen because we wanted to compare pSLA to
other hyperactive behaviors, such as scrunching, which requires a minimum 5 fps frame rate
for detection (Cochet-Escartin et al., 2015; Sabry et al., 2020), and we found that higher frame
rates did not improve signal detection (Supplemental Figure S2).

2.4. Automated image analysis

Analysis of acute behavioral recordings was performed in MATLAB (2025a, MathWorks, Natick, MA) using custom scripts. Any worm that crawled out of the water during recording was excluded from analysis. To create a standardized image of the background to be using for masking the individual wells and removing background noise, the average projection of the first 2 min of recording was used. For determination of overall activity and changes in body shape, we calculated the absolute value of the difference of consecutive frames divided by the absolute value of the sum of the two images. The difference image was then binarized using a noise threshold determined from recordings of plates containing water only under the same imaging conditions. The motility index (MI) was then calculated for each well and frame as the sum of pixels that changed between the two images above the noise threshold. The area of each worm was calculated via thresholding of the image following subtraction by the average projection background image. As optical aberrations during imaging create some "blind spots" in which the planarian can become lost, any frames where no worm was found were excluded in the MI calculations. As individual bouts of pSLA have been previously described as lasting approximately 1 second (Rawls et al., 2009), the median MI score for 1 sec (5 frame) bins was calculated. These scores were then normalized by the median worm area in the first 2 min of recording, where the worms are most likely to be moving the most and thus have elongated shapes.

To correlate normalized MI scores with the presence of pSLA, we manually evaluated the recording of 10-12 planarians exposed to NMDA (3 mM in DD, 15 mM in DJ) across 3 independent plates to identify pSLA events. For each analyzed worm, 1 minute of recording (300 1 sec bins) was chosen, which exhibited high pSLA activity qualitatively by eye and 1 sec bins containing pSLA were marked. For each species, we then evaluated the distribution of normalized MI scores across all identified pSLA events (Supplemental Figure S3). The 10^{th} percentile of each species' pSLA distributions (DD: 0.11; DJ: 0.21) was then used as a threshold, where normalized MI scores above this threshold were counted as pSLA events.

MI provides information about the general activity of a planarian but gives no context to the spatial displacement of the worm. Thus, we also quantified the translational activity of each planarian over 20 sec (100 frame) bins. This bin size was chosen because a normal gliding planarian can circle around the outside of a well of a 48-well plate in about this time. For each bin, the minimum intensity projection was determined. The average projection of the first 2

min of recording was subtracted from the minimum intensity projection to remove the background. The resulting image was binarized using a pre-determined threshold. The translational activity of each worm was calculated as the sum of pixels in this filtered image (thus representing the motion of the worm) divided by the median area of the worm during the first 2 min of recording.

2.5. Lethality analysis

To determine if the presence of acute pSLA is correlated with subsequent lethality, some plates were removed after the initial acute recording and stored in a dark 20 °C incubator until the following day. Plates were recorded for 5 min at 21-27 hours after plate setup. From the recordings, each planarian was manually scored as alive, dead, "crawl-out" when the worm crawls out of the solution in the well, or "sick" when the worm was alive/moving but displayed abnormal morphology, indicated by head regression, pharynx extrusion, lesions/disintegration or C-shape body postures.

2.6. Statistical analysis

The number of pSLA events, defined by each species threshold (see section 2.4), over the entire 30 min recording were counted and are provided in Supplemental File S1. Statistical analysis was performed in R (4.5.0, (R Core Team, 2016)). For each compound, data were compiled across all independent replicates and compared with the compiled in plate vehicle controls (planarian water or 0.5% DMSO) across the same replicates. A negative binomial generalized linear model was performed using the MASS package (Venables and Ripley, 2002) to determine an effect of condition on the number of pSLA events. Post hoc pairwise comparisons of the estimated marginal means of the model was performed using the contrast function of emmeans ((Lenth, 2023) using the Benjamini & Hochberg (BH) p-value adjustment method for multiple comparisons.

3. Results

3.1. Defining pSLA events using prototypical concentrations of NMDA

The term pSLA was first introduced in 2009 by Rawls and co-authors to describe planarian behavior that looked convulsive and indicative of seizures in response to NMDA treatment (Rawls et al., 2009). However, pSLA identification was qualitative and the manual counting of pSLA is extremely time consuming and susceptible to bias. To overcome these limitations and

thus make rapid screening of potential seizurogenic compounds in freshwater planarians

possible, we developed a pipeline to robustly and quantitatively identify pSLA.

Because we needed a prototypical example of pSLA to build our analytical pipeline, we first

identified a NMDA concentration that qualitatively induced robust pSLA in both planarian

species within 30 minutes of exposure. Using the published data as a guide (Rawls et al., 2009),

we tested 3 mM NMDA and found that this concentration immediately induced convulsive

behavior in DD. However, robust convulsions were not observed in DJ until 15 mM NMDA

was used. At these selected concentrations, NMDA-induced various paroxysms, including head

flailing, turning, rapid twitching and convulsions, over-extension and C-shapes, corkscrews

and spiral curling in both species (Supplemental Videos S1 and S2), as previously described in

the literature. Figure 3 shows example high magnification images of the different shapes and

behaviors that we observed in both species.

Figure 3. High magnification examples of abnormal body shapes and behaviors observed

during NMDA-induced pSLA. Normal gliding behavior (left) as observed in planarian water

exhibits relatively fixed body lengths and straight motion. During NMDA exposure (3 mM in

DD and 15 mM in DJ), planarians change rapidly between different abnormal body shapes.

Scale bars: 0.5 mm.

Next, to allow for rapid behavioral screening of multiple conditions simultaneously, we

developed a medium-throughput screening set-up using 48-well plates as explained in Methods

Section 2.3 and shown in Figure 2. Each well contained one planarian, with 2 columns per

treatment condition, for a total of 12 planarians per condition per plate, allowing for

simultaneous testing of 4 conditions in the same plate (Figure 2A). The use of commercial components (provided in Supplemental Table S2) for the behavioral imaging box (Figure 2B) allows for replication of this method in other laboratories. Using this set-up, chemically treated planarians were imaged for 30 minutes immediately (within 2-6 min) following exposure. To quantitatively identify pSLA, we measured the motility index (MI, see Methods Section 2.4), normalized by worm area, for each planarian over the 30 min recording period. As individual bouts of pSLA have been previously described as lasting approximately 1 sec (Rawls et al., 2009), we calculated the median MI per 1 sec (5 frames) bins for each worm. Species-specific pSLA thresholds were then determined based on the distribution of normalized MI scores for a subset of representative NMDA-exposed planarians (see Methods Section 2.4, Supplemental Figure S3), where normalized MI scores per 1 sec bin above this threshold were identified as potential pSLA events. We then applied this analysis to all NMDA data.

We found that NMDA showed significantly more pSLA events than the respective controls in each species (Figure 4A-B). Because these NMDA concentrations had low pH (~3, Supplemental Table S1), we also tested the effect of pH alone using planarian water at approximately the same pH as NMDA. Acidic planarian water induced significantly more pSLA events than the respective non-adjusted controls, though not to the same extent as with NMDA. However, in DJ, the number of pSLA events in the acidic pH sample was not significantly different from NMDA. We conclude that pH is a contributing factor to NMDA-induced pSLA in both species.

While the total number of pSLA events over the entire recording time allows classification of a chemical as seizurogenic or not, there is additional information about the chemical's action in the dynamics of the response, which can be evaluated from the normalized MI scores as a function of time (Figure 4C-D). In both species, NMDA induced high activity within 0-2 min of recording time, which dissipated within ~5 (DD) or ~10 (DJ) min. After this time, NMDA-exposed planarians became contracted and immobile and/or crawled out of the NMDA solution. As 0 or nearly 0 MI scores are seen with both normal gliding behavior and immobility, we also quantified the translational activity of the planarians (see Methods Section 2.4). These translational activity plots show that the decrease in MI scores in NMDA-exposed planarians is concomitant with decreased motion around the well (Figure 4E-F). This decreased motility may indicate the toxic nature of NMDA as almost all worms in both species were dead or had crawled out of their wells by ~24 h (Supplemental Figure S4).

Figure 4. Validation of quantitative analysis of pSLA activity using NMDA. (A-B) Boxplots of the number of pSLA events per 30 min recording in A) DD and B) DJ exposed to NMDA and planarian water at the same pH as the respective NMDA concentration. Dots represent individual planarians. Statistical significance was determined using pairwise contrasts (with a Benjamini & Hochberg p-value correction) of the estimated marginal means of a negative binomial generalized linear model of condition versus number of pSLA events. Different lower-case letters indicate the groups are statistically significantly different (p<0.05). (C-D) Median normalized MI for all C) DD or D) DJ planarians exposed to their prototypical NMDA concentration and their respective in plate vehicle controls as a function of time. (E-F)

Median normalized translational activity of all E) DD or F) DJ planarians exposed to their

prototypical NMDA concentration and their respective in plate vehicle controls as a function

of time. Sample sizes are listed in Supplemental Table S3.

3.2. MI analysis distinguishes pSLA from other hyperactive behaviors

To verify that our analysis method is specific to pSLA, we exposed DD and DJ planarians to

AITC, which has been shown to activate the planarian Transient Receptor Potential Ankyrin

channel (*Dj*TRPAa) and induce forward scrunching, a periodic muscle-based form of

locomotion (Sabry et al., 2019). AITC concentrations that induced scrunching within a few

minutes of exposure (25 µM for DD and 100 µM for DJ) had increased normalized MI scores

(Figure 5A-B) and showed differences in translational activity compared to controls (Figure

5C-D). However, the MI scores were not high enough to be detected as significant pSLA

activity (Figure 5E-F). AITC behavior was only analyzed for 10 minutes following exposure

as scrunching did not persist past this time.

Figure 5. Quantitative MI analysis can distinguish pSLA from scrunching behavior induced by AITC. (A-B) Median normalized MI for all A) DD or B) DJ planarians exposed to AITC and their respective in plate vehicle controls as a function of time. Note that in A the control curve is not visible as all scores were 0. (C-D) Median normalized translational activity of all C) DD or D) DJ planarians exposed to AITC and their respective in plate vehicle controls as a function of time. (E-F) Boxplots of the number of pSLA events per 10 min recording in E) DD and F) DJ exposed to AITC and the respective in plate vehicle control. Dots represent individual planarians. No significant differences were found between controls and AITC in each using pairwise contrasts (with a Benjamini & Hochberg p-value correction) of the estimated marginal means of a negative binomial generalized linear model of condition versus

number of pSLA events. Note the y-axes are scaled differently in the various plots to best reflect
each data set. Sample sizes are listed in Supplemental Table S3.

3.3. Compounds known to induce pSLA in planarians are detected via MI analysis
Next, we tested other compounds that have previously been reported to induce pSLA in
planarians (Table 2).

Table 2. Chemicals and concentrations reported to induce pSLA in planarians.

Chemical	Species	Concentration range	References
Nicotine	DT; DD	10-2000 µM (DT) 1-5 mM (DD)	(Bach et al., 2016; Pagán et al., 2015; Ramakrishnan and DeSaer, 2011; Rawls et al., 2011)
Pilocarpine	DD	1,2,3,4,6 mM	(Miller et al., 2025)
PTX	DT	0.01-5 mM	(Ramakrishnan and DeSaer, 2011)

3.3.1 Nicotine
Studies by different research groups have reported hyperactivity and pSLA in DD and DT
planarians upon acute nicotine exposure in the µM to mM range (Figure 1 and Table 2).
Nicotine binds to and activates nicotinic acetylcholine receptors. While there is supporting
pharmacological evidence for nicotinic acetylcholine receptors in DD planarians (Rawls et al.,
2011), the receptors have not yet been characterized. No significantly increased pSLA events
were observed in DD planarians exposed to 10 – 50 µM nicotine (Figure 6A) although we
qualitatively observed hyperactivity and pSLA-like events in DD planarians exposed to 10 µM
nicotine immediately following exposure. This hyperactivity was slower and less intense than
pSLA seen with 3 mM NMDA, as supported by lower amplitudes in the MI versus time plot
for nicotine (Supplemental Figure S5) compared to NMDA (Figure 4C). In contrast to 3 mM
NMDA, no lethality was observed at ~24 h with 10 µM nicotine (Supplemental Figure S4).

Figure 6. MI analysis of known seizurogenic compounds in planarians. Boxplots of the number of pSLA events per 30 min recording in (A-B) Nicotine, (C-D) Pilocarpine, and E-F) PTX in DD (left column, A, C, E) and DJ (right column, B, D, F). Dots represent individual planarians. Sample sizes are listed in Supplemental Table S3. Statistical significance was determined using pairwise contrasts (with a Benjamini & Hochberg p-value correction) of the estimated marginal means of a negative binomial generalized linear model of condition versus number of pSLA events. Different lower-case letters indicate the groups are statistically significantly different (p<0.05). Note the y-axes are scaled differently in the various plots to best reflect each data set.

Nicotine-induced pSLA quickly dissipated within the first 2 minutes, followed by a return to gliding behavior (Supplemental Figure S5 and S6). For these reasons, and because of the noise in the controls, no statistical significance was found when considering the number of pSLA

events over the entire 30 min recording. When we focused on only the first 5 min of recording, we found significant pSLA for 10 µM nicotine (Supplemental Figure S7). Thus, our analysis shows that nicotine induces pSLA, in agreement with published data, which only scored 5 min of exposure (Ramakrishnan and DeSaer, 2011; Rawls et al., 2011).

Interestingly, pSLA was not observed in higher concentrations (15 and 25 µM) which largely displayed normal gliding. Exposure to 50 µM nicotine resulted in contraction and paralysis over the course of the 30 min recording (Supplemental Figure S8A). In contrast, in DJ planarians, significantly increased pSLA was observed starting at 100 µM nicotine, with the most pSLA events observed in 750 µM, which was highly hyperactive and demonstrated NMDA-like pSLA consisting of head flailing, convulsions and dynamic changes between S- and C-shapes (Figure 6B). In 750 µM nicotine, pSLA activity dissipated/slowed after 5 min of recording and was replaced by contraction with some sporadic, slow convulsive events (Supplemental Figure S9) and very little translational motion (Supplemental Figure S10). Contraction and further morphological abnormalities such as C-shapes and lesions were observed at ~24 h of exposure to 750 µM nicotine, although little lethality was observed (Supplemental Figure S4).

3.3.2 Pilocarpine

It has recently been shown that a muscarinic receptor agonist, pilocarpine, can also trigger pSLA in DD. Using particle tracking to quantify the time spent in the center and the periphery of a well of a 12-well plate and manual scoring of different behaviors such as "dorsal oscillations", "head flick", "C-shape", and others, concentrations of 4-6 mM pilocarpine were found to be significantly different from controls (Miller et al., 2025). Thus, we first tested 2-6 mM in DD planarians. Using MI analysis, we found that all tested concentrations of pilocarpine induced statistically significant pSLA in DD (Figure 6C), in agreement with the study by Miller et al. (Miller et al., 2025). No lethality was observed at 24 h (Supplemental Figure S4). However, DJ planarians only displayed significantly increased pSLA at ≥ 40 mM pilocarpine, with the most pSLA events observed in 100 mM pilocarpine (Figure 6D), which also caused 100% lethality at 24 h (Supplemental Figure S4). Thus, the two species show clear differences in sensitivity to pilocarpine, which may be due to differences in uptake, molecular targets, or metabolism.

3.3.3 PTX

The GABA$_A$ receptor antagonist PTX has been shown to induce pSLA in DT planarians (Ramakrishnan and DeSaer, 2011) but no studies have yet been conducted in DD or DJ planarians. Because of limited water solubility, the highest concentration of PTX that could be tested was 5 mM. We found that 5 mM PTX induced hyperactivity and pSLA within 30 min of recording in both species (Figure 6E-F). However, in contrast to the other chemicals tested, which induced pSLA within the first few minutes, 5 mM PTX induced pSLA was only observed after 20-25 min of recording, following initial gliding behavior and sporadic bouts of head twitching and dorsal oscillations (Supplemental Figures S4 and S8), suggesting that this concentration of PTX was at the potency threshold. PTX did not induce lethality in either species at ~24 hours (Supplemental Figure S4), but the worms were elongated and were still exhibiting hyperactive behaviors.

3.4. PTZ can induce pSLA

Similar to PTX, PTZ also antagonizes the GABA$_A$ receptor (albeit through a different mechanism) and is frequently used to generate animal seizure models (Afrikanova et al., 2013; Chitolina et al., 2023; Dhir, 2012; Milder et al., 2022; Shimada and Yamagata, 2018; Yang et al., 2023). We therefore tested varying concentrations of PTZ for its ability to induce pSLA in both DD and DJ planarians. We found that concentrations exceeding 10 mM (DD) or 5 mM (DJ) PTZ significantly induced pSLA (Figure 7A-B). PTZ-induced pSLA was characterized by an initial phase of head wiggling and turning, followed by random twitching and convulsions that included longitudinal over-extension and C-shapes, corkscrews and spiral curling. In DD, these hyperactive behaviors slowed down over the 30 min of recording and some of the planarians moved into a contracted state, indicating muscle paralysis (Supplemental Figures S5-6). In contrast, in DJ planarians, hyperactive behavior persisted throughout the entire 30 min recording (Supplemental Figure S9). Both species were impacted at 24 hours with high numbers of death, sickness and crawl-out behavior, though the exact mix of these three abnormal conditions differed across the species (Supplemental Figure S4)

Figure 7. PTZ and parathion induce pSLA in DD and DJ planarians. Boxplots of the

number of pSLA events per 30 min recording in (A-B) PTZ, (C-D) parathion, (E-F) carbaryl,

and (G-H) permethrin in DD (left column) and DJ (right column). Dots represent individual

planarians. Sample sizes are listed in Supplemental Table S3. Statistical significance was

determined using pairwise contrasts (with a Benjamini & Hochberg p-value correction) of the

estimated marginal means of a negative binomial generalized linear model of condition versus

number of pSLA events. Different lower-case letters indicate the groups are statistically
significantly different (p<0.05). Note the y-axes are scaled differently in the various plots to
best reflect each data set.

3.5. Some pesticides induce pSLA

Pesticides with different molecular targets have been found to be seizurogenic in mammalian
models (Jett, 2012). Because both nicotine and pilocarpine, two cholinergic receptor agonists,
induced pSLA, we hypothesized that organophosphorus pesticides and carbamates that inhibit
acetylcholinesterase (AChE) in planarians as in humans (Hagstrom et al., 2017) and can cause
seizures in mammals at high concentrations (Jett, 2012), could trigger pSLA upon acute high
concentration exposure. To test this hypothesis, we tested one chemical from each class,
parathion and carbaryl, respectively. Concentrations of 10 µM parathion and above caused
significant pSLA in both planarian species (Figure 7C-D). While 50 µM caused the most pSLA
events in both species, 31.6 µM parathion was a better concentration to induce pSLA in DD
due to the high number of crawl-out events seen in this species at 50 µM. Following ~24 hour
exposure to these concentrations of parathion, both species showed sickness and lethality
(Supplemental Figure S4). Carbaryl induced much weaker pSLA events in both species, which
were characterized by small contractions and head twitches. While 25 µM showed significantly
increased pSLA in DJ planarians compared to vehicle controls, 10-75 µM carbaryl showed no
significant effect on the number of pSLA in DD planarians (Figure 7 E-F), despite 25 µM
showing increased median MI scores per bin (Supplemental Figure S5). At 75 µM, some
scrunching and head twitching was observed within the first 5 minutes of recording, following
by decreased movement and contraction (Supplemental Figure S8B), suggesting that this
concentration exceeded the pSLA-inducing range of concentrations. No significant lethality
was observed in either species exposed to 25 µM carbaryl at ~24 h (Supplemental Figure S4).

Some pyrethroid insecticides, such as permethrin, have been shown to cause seizures in rodents
at lethal concentrations (Staatz and Hosko, 1985) or in environmentally relevant concentrations
in cats (Dymond and Swift, 2008). Pyrethoids inhibit voltage-gated sodium channels, leading
to neuronal hyper-excitability and at high doses can inhibit GABA signalling (Jett, 2012). We
tested permethrin up to its solubility limit (1 mM) and did not observe significant pSLA in
either planarian species (Figure 7G-H). Additionally, no obvious differences in behavior could
be seen in the MI or translational activity plots (Supplemental Figures S5, S6, S9, S10).

3.6. The planarian brain is not required to induce pSLA in DJ planarians

It had previously been shown that pSLA in headless DT planarians was similar to that in intact

planarians when exposed to nicotine but not to cocaine, another pSLA inducer (Pagán et al.,

2012). Cocaine-induced pSLA was nearly completely lost in regenerating tails and was

gradually restored during regeneration (Pagán et al., 2013). These results suggest that cocaine

sensitive receptors are primarily found in the brain, consistent with dopamine receptors in

vertebrates. In contrast, nicotinic acetylcholine receptors must be distributed in both the head

and the body because the response in headless planarians was similar to that of intact

planarians. Therefore, comparing responses of intact and headless planarians to seizurogenic

chemicals may provide insights into where the molecular targets of these chemicals are

expressed in planarians. Thus, we tested whether the presence of the brain was required for

pSLA by exposing regenerating DJ tails to each seizurogenic compound at a concentration

shown to significantly induce pSLA in adult/intact DJs. Exposure began 4 h after amputation

to allow for wound closure and to mitigate the increased mucus production following

amputation (Peiris et al., 2014) that may hinder chemical uptake. All chemicals except for

carbaryl induced pSLA in regenerating DJ tails (Table 3).

Table 3. **Median number of pSLA in adult or regenerating tail DJ planarians exposed to**

representative concentrations of seizurogenic compounds. Sample sizes are listed in

Supplemental Table S3.

Condition	Adult Median # pSLA (25th, 75th percentile)	Regenerating Median # pSLA (25th, 75th percentile)
15 mM NMDA	42 (6, 63.5)	45 (33, 68)
pH 3	14 (5, 21)	69 (24, 132)*
750 μM nicotine	96 (83, 152)	29 (11, 55)*
100 mM pilocarpine	37 (15,88)	69 (43,201)
5 mM PTX	12 (6, 15)	15 (10, 21)
15 mM PTZ	75 (49 , 100)	150 (93, 197)
50 μM parathion	394 (194,470)	59 (37, 136)*
25 μM carbaryl	3 (2, 13)	0 (0,1)

* significantly different from controls ($p < 0.05$) using pairwise contrasts (with a Benjamini & Hochberg p-value
correction) of the estimated marginal means of a negative binomial generalized linear model of condition versus
number of pSLA events

Some compounds, including NMDA, pilocarpine, PTX, and PTZ induced similar amounts of

pSLA in the two developmental stages with overlapping interquartile ranges. Nicotine and

parathion showed significantly more active behaviors - evidenced by higher numbers of pSLA

- in adult DJ planarians than in tails, suggesting that the presence of the brain plays a role for

pSLA induction in DJ planarians. Interestingly, pH 3 showed significantly greater activity in the regenerating tails than in adults. Thus, in DJ planarians, while the seizurogenic compounds do not strictly require the planarian brain to induce pSLA, presence of the brain can affect pSLA dynamics depending on the compound. Carbaryl was at the detection threshold and thus a comparison may not be meaningful.

4. Discussion

4.1. MI analysis can classify seizurogenic compounds and distinguish pSLA from other behaviors

To overcome the limitations associated with manual scoring of pSLA, namely scoring bias and time effort, we created an automated MI analysis method that uses image subtraction to detect changes in body shape. Using frame subtraction to quantify activity is a commonly used method to quantify behaviors including seizures in other model systems such as zebrafish (Afrikanova et al., 2013; Bruni et al., 2016; Whyte-Fagundes et al., 2025). By using qualitative scoring of NMDA-induced pSLA as a framework, we derived a quantitative MI threshold for pSLA in DD and DJ planarians. This analysis method was able to correctly identify known seizurogenic chemicals in planarians (NMDA, nicotine, PTX, pilocarpine) and distinguish pSLA from other abnormal behaviors, such as forward scrunching, which – in contrast to pSLA – exhibits periodic changes in body shape (Cochet-Escartin et al., 2015). Complementary analysis of the translational worm activity as a function of time helped us to distinguish normal behavior from contraction and paralysis, which both show low MI scores.

With the aim of providing a simple metric for easy identification of potential seizurogenic compounds, we quantified the number of pSLA events over the 30 minutes of recording. This recording length is longer than many previous planarian seizure studies, which primarily evaluated the first 5-10 minutes after chemical exposure. We increased this testing window to provide a better opportunity to identify seizurogenic compounds without the need for precise concentration tuning, while still being short enough to allow for medium-throughput screening. However, this longer recording time also potentially decreased our detection power for transient, short-lived behaviors that did not have time to accumulate many pSLA events, as was the case for DD planarians exposed to nicotine. Thus, in some cases, looking at the number of pSLA as a function of time may be necessary to identify weaker effects.

One limitation of MI analysis is that it has difficulty to detect pSLA when the
convulsive/twitching behavior is small and/or primarily in the third dimension (e.g., in the case
of carbaryl), which is not well captured in the simple 2D imaging setup we used. While these
behaviors could still be detected as pSLA in DJ planarians, they were not found to be significant
for DD planarians because DD planarians showed more variable control behavior, including
swimming, scrunching, and turning over. This variability in DD behavior is likely due to them
having been obtained commercially and not having been raised for generations under
controlled laboratory conditions.
603

4.2. MI analysis confirms that known seizurogenic chemicals identified by qualitative studies
induce pSLA in planarians

We confirmed previous studies by other research groups that NMDA (Rawls et al., 2009),
nicotine (Rawls et al., 2011), and pilocarpine (Miller et al., 2025) induce pSLA in DD
planarians and showed that they also induce pSLA in DJ planarians. PTX had previously only
been studied in DT planarians (Ramakrishnan and DeSaer, 2011); here, we show that it also
induces pSLA in DD and DJ planarians. The concentrations of NMDA, pilocarpine, and PTX
that induced pSLA in DD planarians here were comparable to previous studies using DD or
DT planarians.

However, we found that 3 mM NMDA lowered pH to a range where pH alone caused pSLA
while previous work found that 3 mM NMDA did not lower the pH as much (pH 6.6), and thus
pH alone did not induce pSLA (Rawls et al., 2009). The discrepancy in these data may be due
to differences in the exact water composition as the previous study used Artificial Pond Water
(APW; 6 mM NaCl; 0.1 mM NaHCO$_3$; 0.6 mM CaCl$_2$; pH 7.3) (Rawls et al., 2009) which has
slightly different salt composition compared to our planarian water (see Methods Section 2.1)
and thus may have different buffering capacity.
620

For nicotine, we found that 50 μM caused paralysis in DD planarians, comparable to
concentrations used in DT planarians (Ramakrishnan and DeSaer, 2011). In contrast, other
studies in DD planarians have used nicotine in the mM range (Bach et al., 2016; Pagán et al.,
2015, 2013; Rawls et al., 2011). The difference is likely be due to the product used. While we
and (Ramakrishnan and DeSaer, 2011) used pure (-) nicotine solution, the other studies (Bach
et al., 2016; Pagán et al., 2015, 2013; Rawls et al., 2011) used nicotine hydrogen tartrate salt.
627

4.3 PTZ and cholinesterase inhibitors as pSLA inducers

We showed that PTZ, which is used to generate mice (Dhir, 2012; Shimada and Yamagata, 2018) and zebrafish (Afrikanova et al., 2013; Baraban et al., 2005; Chitolina et al., 2023; Milder et al., 2022) seizure models, robustly induces pSLA in both planarian species. Thus, PTZ could be used to generate a planarian seizure-like model to test the efficacy of possible anti-seizure treatments. In fact, existing studies that used other compounds than PTZ suggest that pSLA in planarians can be modulated by exposure to known anti-convulsant therapeutics (e.g., (Pagán et al., 2015; Ramakrishnan and DeSaer, 2011; Rawls et al., 2009)). Now that we have developed this automated platform, it can be used to validate existing manual studies on therapeutic effectiveness. In addition, RNA interference (RNAi) could be used to verify that these known seizurogenic chemicals induce pSLA through the same pathways as in mammals. For example, PTX and PTZ both target the $GABA_A$ receptor, thus making it a good starting point for mechanistic studies. Parathion, an organophosphorus pesticide, induced pSLA in both planarian species, and carbaryl, a carbamate, caused pSLA in DJ planarians. Both compounds inhibit planarian cholinesterases (Hagstrom et al., 2017; Ireland et al., 2022b); thus, RNAi knockdown of the planarian cholinesterases could be used to verify that the molecular initiation event of pSLA in planarians is cholinesterase inhibition, as in humans (Jett, 2012).

Interestingly, parathion and nicotine both showed significantly more pSLA events in adult (intact) DJ planarians than in decapitated ones. Thus, the brain is not required for pSLA induced by these compounds, as expected because cholinesterases (Hagstrom et al., 2018) and nicotinic receptors (Pagán et al., 2013) are found throughout the planarian body, but is important for inducing and potentially modulating pSLA. It was previously shown in DT planarians that decapitation decreased the number of pSLA events in nicotine, though the effect was not significant (Pagán et al., 2013). The difference between this and our study may be due to species differences, different nicotine formulations as mentioned above, or analysis methods.

Finally, it is also still necessary to confirm that pSLA is indicative of seizures using measures of neuronal activity. A recent study has shown that using surface wire electrodes one can measure EEG activity non-invasively in planarians (Freiberg et al., 2023). Alternatively, microelectrode arrays have been successfully used to measure neuronal activity *in vivo* in zebrafish larvae (Tomasello and Sive, 2020) and could be adapted for planarians. These verification studies will be critical for adoption of the planarian model for screening

seizurogenic chemicals as they would show that results found in planarians are relevant and
translatable to mammals and humans.

4.4 DD and DJ planarians respond similarly to chemicals but with different sensitivities

DD and DJ planarians displayed very different sensitivities to NMDA, nicotine, and
pilocarpine. DD planarians were more sensitive, with the most extreme difference in sensitivity
of nearly 100-fold for nicotine. While NMDA and pilocarpine induced similar behavioral
phenotypes in the two species, there were also qualitative differences in the behaviors displayed
by the two species in response to nicotine as DJ planarians exhibited extensive pSLA events
that were never observed in DD planarians. Other studies have reported similar behavioral and
sensitivity differences between planarian species in response to certain chemicals (e.g., (Ireland
et al., 2020, p. 20; Sabry et al., 2019)). Together, these data emphasize that extrapolation
between planarian species is not recommended due to possible differences in chemical uptake,
metabolism and/or presence of targets. We also showed that DJ planarians are more
behaviorally homogeneous than DD planarians, which may be due to having been bred under
controlled laboratory conditions with standardized food for years whereas DD planarians had
only been maintained under these conditions for a few weeks in our study. However, the
increased sensitivity of DD planarians to some of the compounds tested may make them a
better model for screening seizurogenic chemicals. More compounds will need to be screened
in both species to evaluate their strengths and weaknesses for this application. A good starting
point would be to test chemicals that have been proposed to induce pSLA in planarians in recent
manual studies, such as sertraline (Thumé and Frizzo, 2017) and various cholinesterase
inhibitors (Bezerra Da Silva et al., 2016), which were not included here.

4.5 Use of planarians for screening seizurogenic chemicals and therapeutics

Our results demonstrate that these two planarian species are promising models for rapid
screening of chemicals for seizurogenic potential. While planarians are simpler than zebrafish
and lack a circulatory system and blood-brain barrier (Rawls et al., 2010), they have multiple
advantages for early screening. They are less expensive than zebrafish, which as vertebrates
require animal welfare protocols for husbandry that are not necessary for invertebrates such as
planarians (Collins et al., 2024). While zebrafish larvae at 5 days post fertilization (dpf) are
primarily used for screening because at this stage experiments are not considered animal testing
(Strähle et al., 2012), their nervous system and metabolic machinery are still maturing at this
time (Menezes et al., 2014; Saad et al., 2017; Tiedeken and Ramsdell, 2010). Thus, by

screening compounds in adult planarians with metabolic capacity (Ireland et al., 2022a) and mature neuronal function first, in conjunction with *in silico* chemical modelling and/or human *in vitro* MEA assays, we can reduce cost and vertebrate use in early seizure liability testing.. We have previously shown that organismal planarian and zebrafish behavioral screening and MEA assays were the most informative to identify developmental neurotoxicants (Ireland et al., 2025b) and thus may prove to be a promising combination for identify seizurogenic chemicals and anti-seizure therapeutics.

As discussed in the Introduction, all models have limitations, some of which are unique and some which are shared across models. The strengths and limitations of planarians and developing zebrafish (<5 dpf) for chemical screening have recently been reviewed (Collins et al., 2024). One limitation shared between these organismal models and cell culture models is the requirement for aqueous chemical exposure. Thus, it is challenging to test poorly water soluble or volatile compounds in these systems (Collins et al., 2024) and sufficiently high concentrations to induce pSLA may not be possible, as evidenced by our ability to only test PTX at its seizure induction threshold. *In silico* modelling can help overcome some experimental limitations, as has been shown recently for screening anti-seizure drugs in zebrafish larvae (Belair et al., 2025; Cassar et al., 2017). By performing comparative studies of compounds known to be seizurogenic in humans, we can determine the complementarity of these rapid screening seizure models for a potential seizure liability test battery and identify current data gaps and limitations, similar to current efforts in developmental neurotoxicity testing (Ireland et al., 2025b; Masjosthusmann et al., 2020).

The work here can act as a starting point for future screens of seizurogenic chemicals and anti-convulsants in planarians but further optimization and system validation are required to decide whether planarian screening could be a valuable part of an early stage seizure liability testing battery. In addition to the mechanistic studies mentioned earlier that are needed to confirm molecular targets and distinguish seizures from convulsions, more compounds that have been studied in cell culture and zebrafish need to be evaluated in planarians to compare system sensitivity and specificity (Belair et al., 2025). To accelerate system evaluation and have multiple research groups participate in characterizing the value of planarian behavioral screening, the recording and analysis methods developed here are made available to other laboratories. This will ensure standardized conditions that allow cross-laboratory data comparisons and integration.

While the column setup that we used here was helpful for testing a few selected compounds,

for new screens, we recommend a row setup, similar to what is used for rapid behavioral

screening in other studies (Ireland et al., 2025a; Zhang et al., 2019), because it allows for the

simultaneous testing of 5 concentrations/conditions plus a vehicle control. By using 2 plates

with alternating row setup, one can control for potential edge effects (Zhang et al., 2019) and

achieve n=16 per condition. There is a trade-off between screening throughput and reliability

of hit calls: fewer wells per condition increase screening speed while more wells increase

screen robustness and statistical power. In zebrafish, n=6 wells/condition have been used to

screen chemicals for potential anti-seizure drugs (Dinday and Baraban, 2015). Here, we used

a minimum of n=12 specimen per condition because of the observed variability in DD

planarians, which requires larger sample sizes to have more statistical power. In terms of

timing, we used 30 min to allow for throughput while simultaneously being able to detect late-

onset pSLA (as in PTX). The recording period could be shortened to 5-10 min to achieve higher

throughput as most chemicals induced pSLA within the first 5 min. It's also possible that effects

may wear off, as for example in the case of nicotine for DD planarians, and thus shorter

incubation would be helpful here. But we also showed that additional analysis incorporating

the temporal dynamics can reveal such short-term effects. Our recordings began 2-6 min

following chemical exposure, which was sufficient for the compounds we studied

(Supplemental Figure S11), but this window could be decreased to increase the possibility of

catching very short-lasting pSLA events. In addition to acute behavioral recordings, we also

evaluated lethality one day after chemical exposure to determine if lethality was correlated with

the presence of significant pSLA. This was not the case. Thus, lethality likely results from other

toxic effects besides pSLA and thus is not an informative endpoint for this type of study.

Future work could also incorporate more sophisticated image analysis methods, incorporating

machine learning, to identify specific body shape changes and use them to characterize pSLA,

as has recently been done in zebrafish (Whyte-Fagundes et al., 2025). Such a tailored approach

would likely be more robust than MI analysis and may reveal subtypes of pSLA, which could

provide additional insight into the underlying biological mechanisms. For example, the initial

behavioral effects of parathion are thought to be due to acetylcholinesterase inhibition while

later effects are from secondary neurotransmitter systems such as glutamate (Jett, 2012;

Kozhemyakin et al., 2010); thus, the type of convulsions may differ over time.

In summary, this work demonstrates that pSLA can be triggered through various pathways in DD and DJ planarians, similar to in mammals. By standardizing both experimental approach and analysis methods and making the tools available to the research community, this work can serve as a framework for future studies of chemicals for their seizurogenic potential in planarians.

CRediT authorship contribution statement

Danielle Ireland: Conceptualization; Software; Visualization; Validation; Formal analysis; Data curation; Writing – original draft; Writing - review and editing. **Evangeline Coffinas:** Investigation; Writing – review and editing. **Christina Rabeler:** Investigation; Writing – review and editing. **Eva-Maria S. Collins:** Conceptualization; Software; Methodology; Visualization; Validation; Writing – original draft; Writing – review and editing; Supervision; Resources; Project administration.

Funding

This work was supported by the National Institutes of Health CounterACT program [grant number 5R21NS132528-02]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Acknowledgements

The authors thank Kate Sun and Erin Szuromi for their help with experiments, Dr. Scott Rawls, Dr. Hilary McCarren, and Mathieu Conroy for discussions, and Hannah Poon for comments on the manuscript.

Conflict of interest statement

EMSC is the founder of Inveritek, LLC, which offers planarian HTS commercially. EMSC and DI are inventors on a patent for planarian rapid screening. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at XXX.

Data availability

The datasets supporting the conclusions of this article are included within the article and available from the corresponding author upon reasonable request.

References

Afrikanova, T., Serruys, A.S.K., Buenafe, O.E.M., Clinckers, R., Smolders, I., de Witte, P.A.M., Crawford, A.D., Esguerra, C.V., 2013. Validation of the Zebrafish Pentylenetetrazol Seizure Model: Locomotor versus Electrographic Responses to Antiepileptic Drugs. PLoS ONE 8, 54166. https://doi.org/10.1371/journal.pone.0054166

Authier, S., Arezzo, J., Delatte, M.S., Kallman, M.-J., Markgraf, C., Paquette, D., Pugsley, M.K., Ratcliffe, S., Redfern, W.S., Stevens, J., Valentin, J.-P., Vargas, H.M., Curtis, M.J., 2016. Safety pharmacology investigations on the nervous system: An industry survey. J Pharmacol Toxicol Methods 81, 37–46. https://doi.org/10.1016/j.vascn.2016.06.001

Bach, D.J., Tenaglia, M., Baker, D.L., Deats, S., Montgomery, E., Pagán, O.R., 2016. Cotinine antagonizes the behavioral effects of nicotine exposure in the planarian Girardia tigrina. Neuroscience Letters 632, 204–208. https://doi.org/10.1016/j.neulet.2016.09.005

Baraban, S.C., Taylor, M.R., Castro, P.A., Baier, H., 2005. Pentylenetetrazole induced changes in zebrafish behavior, neural activity and c-fos expression. Neuroscience 131, 759–768. https://doi.org/10.1016/j.neuroscience.2004.11.031

Behl, M., Karmaus, A., Rao, M., Lane, T., Harris, J., Sachs, C., Borrel, A., Oyetade, O., Unnikrishnan, A., Hamm, J., Hogberg, H.T., 2025. De-Risking Seizure Liability: Integrating Adverse Outcome Pathways (AOPs), New Approach Methodologies (NAMs) and In Silico Approaches while Highlighting Knowledge Gaps. Toxicological Sciences. https://doi.org/10.1093/toxsci/kfaf109

Belair, D.G., Kohnken, R., McCloud, R.L., Sandoval, S., Green, J., Buck, W.R., Polakowski, J.S., 2025. Predicting seizure liability of small molecules using an in vitro multi-electrode array based assay coupled with modeling of brain disposition. Current Research in Toxicology 8, 100236. https://doi.org/10.1016/j.crtox.2025.100236

Bezerra Da Silva, C., Pott, A., Elifio-Esposito, S., Dalarmi, L., Fialho Do Nascimento, K., Moura Burci, L., De Oliveira, M., De Fátima Gaspari Dias, J., Maria Warumby Zanin, S., Gomes Miguel, O., Dallarmi Miguel, M., 2016. Effect of Donepezil, Tacrine, Galantamine and Rivastigmine on Acetylcholinesterase Inhibition in Dugesia tigrina. Molecules 21, 53. https://doi.org/10.3390/molecules21010053

Bruni, G., Rennekamp, A.J., Velenich, A., McCarroll, M., Gendelev, L., Fertsch, E., Taylor, J., Lakhani, P., Lensen, D., Evron, T., Lorello, P.J., Huang, X.-P., Kolczewski, S., Carey, G., Caldarone, B.J., Prinssen, E., Roth, B.L., Keiser, M.J., Peterson, R.T., Kokel, D., 2016. Zebrafish behavioral profiling identifies multitarget antipsychotic-like compounds. Nature Chemical Biology 12, 559–566. https://doi.org/10.1038/nchembio.2097

Buttarelli, F.R., Pellicano, C., Pontieri, F.E., 2008. Neuropharmacology and behavior in planarians: Translations to mammals. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology 147, 399–408. https://doi.org/10.1016/j.cbpc.2008.01.009

Cassar, S., Breidenbach, L., Olson, A., Huang, X., Britton, H., Woody, C., Sancheti, P., Stolarik, D., Wicke, K., Hempel, K., LeRoy, B., 2017. Measuring drug absorption improves

interpretation of behavioral responses in a larval zebrafish locomotor assay for
predicting seizure liability. J Pharmacol Toxicol Methods 88, 56–63.
https://doi.org/10.1016/j.vascn.2017.07.002
Chitolina, R., Gallas-Lopes, M., Reis, C.G., Benvenutti, R., Stahlhofer-Buss, T., Calcagnotto,
M.E., Herrmann, A.P., Piato, A., 2023. Chemically-induced epileptic seizures in
zebrafish: A systematic review. Epilepsy Res 197, 107236.
https://doi.org/10.1016/j.eplepsyres.2023.107236
Cochet-Escartin, O., Mickolajczk, K.J., Collins, E.-M.S., 2015. Scrunching: a novel escape gait
in planarians. Physical Biology 12, 055001. https://doi.org/doi: 10.1088/1478-
3975/12/5/056010
Collins, E.M.S., Hessel, E.V.S., Hughes, S., 2024. How neurobehavior and brain development
in alternative whole-organism models can contribute to prediction of developmental
neurotoxicity. NeuroToxicology 102, 48–57.
https://doi.org/10.1016/j.neuro.2024.03.005
Dhir, A., 2012. Pentylenetetrazol (PTZ) kindling model of epilepsy. Curr Protoc Neurosci
Chapter 9, Unit9.37. https://doi.org/10.1002/0471142301.ns0937s58
Dinday, M.T., Baraban, S.C., 2015. Large-Scale Phenotype-Based Antiepileptic Drug Screening
in a Zebrafish Model of Dravet Syndrome. eNeuro 2, ENEURO.0068-15.2015.
https://doi.org/10.1523/ENEURO.0068-15.2015
Dymond, N., Swift, I., 2008. Permethrin toxicity in cats: a retrospective study of 20 cases.
Aust Veterinary J 86, 219–223. https://doi.org/10.1111/j.1751-0813.2008.00298.x
Easter, A., Bell, M.E., Damewood, J.R., Redfern, W.S., Valentin, J.-P., Winter, M.J., Fonck, C.,
Bialecki, R.A., 2009. Approaches to seizure risk assessment in preclinical drug
discovery. Drug Discovery Today 14, 876–884.
https://doi.org/10.1016/j.drudis.2009.06.003
Freiberg, J., Lang, L., Kaernbach, C., Keil, J., 2023. Characterization of the planarian surface
electroencephalogram. BMC Neurosci 24, 29. https://doi.org/10.1186/s12868-023-
00799-z
Gawel, K., Langlois, M., Martins, T., Van Der Ent, W., Tiraboschi, E., Jacmin, M., Crawford,
A.D., Esguerra, C.V., 2020. Seizing the moment: Zebrafish epilepsy models.
Neuroscience & Biobehavioral Reviews 116, 1–20.
https://doi.org/10.1016/j.neubiorev.2020.06.010
Grainger, A.I., King, M.C., Nagel, D.A., Parri, H.R., Coleman, M.D., Hill, E.J., 2018. In vitro
Models for Seizure-Liability Testing Using Induced Pluripotent Stem Cells. Front
Neurosci 12, 590. https://doi.org/10.3389/fnins.2018.00590
Hagstrom, D., Hirokawa, H., Zhang, L., Radić, Z., Taylor, P., Collins, E.-M.S., 2017. Planarian
cholinesterase: in vitro characterization of an evolutionarily ancient enzyme to study
organophosphorus pesticide toxicity and reactivation. Archives of Toxicology 91,
2837–2847. https://doi.org/10.1007/s00204-016-1908-3
Hagstrom, D., Zhang, S., Ho, A., Tsai, E.S., Radić, Z., Jahromi, A., Kaj, K.J., He, Y., Taylor, P.,
Collins, E.M.S., 2018. Planarian cholinesterase: molecular and functional
characterization of an evolutionarily ancient enzyme to study organophosphorus
pesticide toxicity. Archives of Toxicology 92, 1161–1176.
https://doi.org/10.1007/s00204-017-2130-7
Hogberg, H.T., Hsieh, J.-H., Chang, X., Sipes, N.S., Shafer, T., Behl, M., 2022. Case study on the
use of Integrated Approaches for Testing and Assessment for DNT to prioritize a class
of Organophosphorus flame retardants, Series on Testing and Assessment.

891   Organisation for the Economic Cooperation and Development Environment
892   Directorate, Paris.

893   Hogberg, H.T., Smirnova, L., 2022. The Future of 3D Brain Cultures in Developmental
894   Neurotoxicity Testing. Front Toxicol 4, 808620.
895   https://doi.org/10.3389/ftox.2022.808620

896   Ireland, D., Bochenek, V., Chaiken, D., Rabeler, C., Onoe, S., Soni, A., Collins, E.-M.S., 2020.
897   Dugesia japonica is the best suited of three planarian species for high-throughput
898   toxicology screening. Chemosphere 253, 126718.
899   https://doi.org/10.1016/j.chemosphere.2020.126718

900   Ireland, D., Collins, E.-M.S., 2022. New Worm on the Block: Planarians in (Neuro)Toxicology.
901   Current Protocols 2, e637. https://doi.org/10.1002/CPZ1.637

902   Ireland, D., Collins, E.S., 2023. Planarians as a model to study neurotoxic agents, in: Advances
903   in Neurotoxicology, Alternative Methods in Neurotoxicology. Elsevier Inc., pp. 29–60.
904   https://doi.org/10.1016/bs.ant.2023.01.002

905   Ireland, D., Rabeler, C., Gong, T., Collins, E.-M.S., 2022a. Bioactivation and detoxification of
906   organophosphorus pesticides in freshwater planarians shares similarities with
907   humans. Archives of Toxicology. https://doi.org/10.1007/s00204-022-03387-y

908   Ireland, D., Rabeler, C., Rao, S., Richardson, R.J., Collins, E.-M.S., 2025a. Distinguishing classes
909   of neuroactive drugs based on computational physicochemical properties and
910   experimental phenotypic profiling in planarians. PLoS ONE 20, e0315394.
911   https://doi.org/10.1371/journal.pone.0315394

912   Ireland, D., Word, L.J., Collins, E.-M.S., 2025b. Statistical analysis of multi-endpoint
913   phenotypic screening increases sensitivity of planarian neurotoxicity testing.
914   Toxicological Sciences kfaf117. https://doi.org/10.1093/toxsci/kfaf117

915   Ireland, D., Zhang, S., Bochenek, V., Hsieh, J.-H., Rabeler, C., Meyer, Z., Collins, E.-M.S., 2022b.
916   Differences in neurotoxic outcomes of organophosphorus pesticides revealed via
917   multi-dimensional screening in adult and regenerating planarians. Frontiers in
918   Toxicology 4, 948455. https://doi.org/10.3389/ftox.2022.948455

919   Ishii, M.N., Yamamoto, K., Shoji, M., Asami, A., Kawamata, Y., 2017. Human induced
920   pluripotent stem cell (hiPSC)-derived neurons respond to convulsant drugs when co-
921   cultured with hiPSC-derived astrocytes. Toxicology 389, 130–138.
922   https://doi.org/10.1016/j.tox.2017.06.010

923   Jaworska, J., Corsini, E., Basketter, D., 2016. Integrated Testing Strategies for Skin
924   Sensitization Hazard and Potency Assessment—State of the Art and Challenges.
925   Cosmetics 2016, Vol. 3, Page 16 3, 16. https://doi.org/10.3390/COSMETICS3020016

926   Jett, D.A., 2012. Chemical toxins that cause seizures. NeuroToxicology 33, 1473–1475.
927   https://doi.org/10.1016/j.neuro.2012.10.005

928   Kozhemyakin, M., Rajasekaran, K., Kapur, J., 2010. Central Cholinesterase Inhibition
929   Enhances Glutamatergic Synaptic Transmission. Journal of Neurophysiology 103,
930   1748–1757. https://doi.org/10.1152/jn.00949.2009

931   Lenth, R.V., 2023. emmeans: Estimated Marginal Means, aka Least-Squares Means.

932   Masjosthusmann, S., Blum, J., Bartmann, K., Dolde, X., Holzer, A., Stürzl, L., Keßel, E.H.,
933   Förster, N., Dönmez, A., Klose, J., Pahl, M., Waldmann, T., Bendt, F., Kisitu, J., Suciu, I.,
934   Hübenthal, U., Mosig, A., Leist, M., Fritsche, E., 2020. Establishment of an a priori
935   protocol for the implementation and interpretation of an in-vitro testing battery for
936   the assessment of developmental neurotoxicity. EFSA Supporting Publications 17.
937   https://doi.org/10.2903/SP.EFSA.2020.EN-1938

McGraw, C.M., Poduri, A., 2025. Machine learning enables high-throughput, low-replicate
screening for novel anti-seizure targets and compounds using combined movement
and calcium fluorescence in larval zebrafish. Eur J Pharmacol 991, 177327.
https://doi.org/10.1016/j.ejphar.2025.177327

Menezes, F.P., Rico, E.P., Da Silva, R.S., 2014. Tolerance to seizure induced by kainic acid is
produced in a specific period of zebrafish development. Prog Neuropsychopharmacol
Biol Psychiatry 55, 109–112. https://doi.org/10.1016/j.pnpbp.2014.04.004

Milder, P.C., Zybura, A.S., Cummins, T.R., Marrs, J.A., 2022. Neural Activity Correlates With
Behavior Effects of Anti-Seizure Drugs Efficacy Using the Zebrafish Pentylenetetrazol
Seizure Model. Front. Pharmacol. 13, 836573.
https://doi.org/10.3389/fphar.2022.836573

Miller, T., Vu, J., Tran, C.J., Ritvik, A., Cruz-Ham, K., Haq, K., Musto, B., Musto, A.E., 2025.
Planarian as an Animal Model for Experimental Acute Seizure. JoVE 67307.
https://doi.org/10.3791/67307

Pagán, O.R., Baker, D., Deats, S., Montgomery, E., Tenaglia, M., Randolph, C., Kotturu, D.,
Tallarida, C., Bach, D., Wilk, G., Rawls, S., Raffa, R.B., 2012. Planarians in
pharmacology: parthenolide is a specific behavioral antagonist of cocaine in the
planarian Girardia tigrina. The International journal of developmental biology 56,
193–6.

Pagán, O.R., Deats, S., Baker, D., Montgomery, E., Wilk, G., Tenaglia, M., Semon, J., 2013.
Planarians require an intact brain to behaviorally react to cocaine, but not to react to
nicotine. Neuroscience 246, 265–270.
https://doi.org/10.1016/j.neuroscience.2013.05.010

Pagán, O.R., Montgomery, E., Deats, S., Bach, D., Baker, D., 2015. Evidence of nicotine-
induced, curare-insensitive, behavior in planarians. Neurochemical Research 40,
2087–2090. https://doi.org/10.1007/s11064-015-1512-6

Pagán, O.R., Rowlands, A.L., Urban, K.R., 2006. Toxicity and behavioral effects of
dimethylsulfoxide in planaria. Neuroscience Letters 407, 274–278.

Peiris, T.H., Hoyer, K.K., Oviedo, N.J., 2014. Innate immune system and tissue regeneration in
planarians: An area ripe for exploration. Seminars in Immunology 26, 295–302.
https://doi.org/10.1016/j.smim.2014.06.005

Pelkonen, A., Mzezewa, R., Sukki, L., Ryynänen, T., Kreutzer, J., Hyvärinen, T., Vinogradov, A.,
Aarnos, L., Lekkala, J., Kallio, P., Narkilahti, S., 2020. A modular brain-on-a-chip for
modelling epileptic seizures with functionally connected human neuronal networks.
Biosens Bioelectron 168, 112553. https://doi.org/10.1016/j.bios.2020.112553

R Core Team, 2016. R: A language environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Raffa, R.B., 2008. Planaria: A Model for Drug Action and Abuse, 0 ed. CRC Press.
https://doi.org/10.1201/9781498713597

Ramakrishnan, L., DeSaer, C., 2011. Carbamazepine inhibits distinct chemoconvulsant-
induced seizure-like activity in Dugesia tigrina. Pharmacology Biochemistry and
Behavior 99, 665–670. https://doi.org/10.1016/j.pbb.2011.06.003

Rawls, S.M., Patil, T., Tallarida, C.S., Baron, S., Kim, M., Song, K., Ward, S., Raffa, R.B., 2011.
Nicotine behavioral pharmacology: Clues from planarians. Drug and Alcohol
Dependence 118, 274–279. https://doi.org/10.1016/j.drugalcdep.2011.04.001

Rawls, S.M., Patil, T., Yuvasheva, E., Raffa, R.B., 2010. First evidence that drugs of abuse
produce behavioral sensitization and cross-sensitization in planarians. Behavioural
pharmacology 21, 301. https://doi.org/10.1097/FBP.0B013E32833B0098

Rawls, S.M., Thomas, T., Adeola, M., Patil, T., Raymondi, N., Poles, A., Loo, M., Raffa, R.B.,
2009. Topiramate antagonizes NMDA- and AMPA-induced seizure-like activity in
planarians. Pharmacology Biochemistry and Behavior 93, 363–367.
https://doi.org/10.1016/j.pbb.2009.05.005

Roberts, R., Authier, S., Mellon, R.D., Morton, M., Suzuki, I., Tjalkens, R.B., Valentin, J.-P.,
Pierson, J.B., 2021. Can We Panelize Seizure? Toxicol Sci 179, 3–13.
https://doi.org/10.1093/toxsci/kfaa167

Rockley, K., Roberts, R., Jennings, H., Jones, K., Davis, M., Levesque, P., Morton, M., 2023. An
integrated approach for early in vitro seizure prediction utilizing hiPSC neurons and
human ion channel assays. Toxicol Sci 196, 126–140.
https://doi.org/10.1093/toxsci/kfad087

Ross, K.G., Currie, K.W., Pearson, B.J., Zayas, R.M., 2017. Nervous system development and
regeneration in freshwater planarians. Wiley Interdisciplinary Reviews:
Developmental Biology 6, 1–26. https://doi.org/10.1002/wdev.266

Saad, M., Matheeussen, A., Bijttebier, S., Verbueken, E., Pype, C., Casteleyn, C., Van
Ginneken, C., Apers, S., Maes, L., Cos, P., Van Cruchten, S., 2017. In vitro CYP-
mediated drug metabolism in the zebrafish (embryo) using human reference
compounds. Toxicol In Vitro 42, 329–336. https://doi.org/10.1016/j.tiv.2017.05.009

Sabry, Z., Ho, A., Ireland, D., Rabeler, C., Cochet-Escartin, O., Collins, E.M.S., 2019.
Pharmacological or genetic targeting of Transient Receptor Potential (TRP) channels
can disrupt the planarian escape response. PLoS ONE 14, e0226104.
https://doi.org/10.1371/journal.pone.0226104

Sabry, Z., Rabeler, C., Ireland, D., Bayingana, K., Collins, E.M.S., 2020. Planarian scrunching as
a quantitative behavioral readout for noxious stimuli sensing. Journal of Visualized
Experiments 2020, 1–18. https://doi.org/10.3791/61549

Shimada, T., Yamagata, K., 2018. Pentylenetetrazole-Induced Kindling Mouse Model. J Vis
Exp 56573. https://doi.org/10.3791/56573

Smirnova, L., Hogberg, H.T., Leist, M., Hartung, T., 2014. Food for Thought …: Developmental
neurotoxicity – Challenges in the 21(st) century and in vitro opportunities. ALTEX 31,
129–156. https://doi.org/10.14573/altex.1403271

Squires, R.F., Saederup, E., Crawley, J.N., Skolnick, P., Paul, S.M., 1984. Convulsant potencies
of tetrazoles are highly correlated with actions on GABA/benzodiazepine/picrotoxin
receptor complexes in brain. Life Sciences 35, 1439–1444.
https://doi.org/10.1016/0024-3205(84)90159-0

Staatz, C.G., Hosko, M.J., 1985. Effect of pyrethroid insecticides on EEG activity in conscious,
immobilized rats. Pesticide Biochemistry and Physiology 24, 231–239.
https://doi.org/10.1016/0048-3575(85)90133-6

Strähle, U., Scholz, S., Geisler, R., Greiner, P., Hollert, H., Rastegar, S., Schumacher, A.,
Selderslaghs, I., Weiss, C., Witters, H., Braunbeck, T., 2012. Zebrafish embryos as an
alternative to animal experiments—A commentary on the definition of the onset of
protected life stages in animal welfare regulations. Reproductive Toxicology 33, 128–
132. https://doi.org/10.1016/J.REPROTOX.2011.06.121

Thumé, I.S., Frizzo, M.E., 2017. Sertraline Induces Toxicity and Behavioral Alterations in
Planarians. BioMed Research International 2017, 1–8.
https://doi.org/10.1155/2017/5792621
Tiedeken, J.A., Ramsdell, J.S., 2010. Zebrafish Seizure Model Identifies p , p' -DDE as the
Dominant Contaminant of Fetal California Sea Lions That Accounts for Synergistic
Activity with Domoic Acid. Environ Health Perspect 118, 545–551.
https://doi.org/10.1289/ehp.0901301
Tomasello, D.L., Sive, H., 2020. Noninvasive Multielectrode Array for Brain and Spinal Cord
Local Field Potential Recordings from Live Zebrafish Larvae. Zebrafish 17, 271–277.
https://doi.org/10.1089/zeb.2020.1874
Tukker, A.M., Westerink, R.H.S., 2021. Novel test strategies for in vitro seizure liability
assessment. Expert Opin Drug Metab Toxicol 17, 923–936.
https://doi.org/10.1080/17425255.2021.1876026
Van Norman, G.A., 2019. Limitations of Animal Studies for Predicting Toxicity in Clinical
Trials: Is it Time to Rethink Our Current Approach? JACC: Basic to Translational
Science 4, 845. https://doi.org/10.1016/J.JACBTS.2019.10.008
Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S, Fourth. ed. Springer,
New York.
Whyte-Fagundes, P., Efromson, J., Vance, A., Carpenter, S., Bègue, A., Carroll, A., Doman,
T.J.J., Harfouche, M., Baraban, S.C., 2025. Automated detection of complex zebrafish
seizure behavior at scale. Commun Biol 8, 872. https://doi.org/10.1038/s42003-025-
08310-6
Wu, J.P., Li, M.H., 2018. The use of freshwater planarians in environmental toxicology
studies: Advantages and potential. Ecotoxicology and Environmental Safety 161, 45–
56. https://doi.org/10.1016/j.ecoenv.2018.05.057
Yang, C.-S., Wu, M.-C., Lai, M.-C., Wu, S.-N., Huang, C.-W., 2023. Identification of New
Antiseizure Medication Candidates in Preclinical Animal Studies. Int J Mol Sci 24,
13143. https://doi.org/10.3390/ijms241713143
Zhang, S., Hagstrom, D., Hayes, P., Graham, A., Collins, E.-M.S., 2019. Multi-behavioral
endpoint testing of an 87-chemical compound library in freshwater planarians.
Toxicological Sciences 167, 26–44. https://doi.org/10.1093/toxsci/kfy145
1059