Title

Sex Differences in Low-dose Ethanol Effects on Motivated Behavior and Limbic Corticostriatal Activity

Authors

Christina M Curran-Alfaro, MS[1,2], Kathleen G Bryant, PhD[1,2], Toni-Shae Ledgister[1], MS, Sana Amin[1],

BS, Jacqueline M Barker, PhD[1*]

Affiliation

1. Department of Pharmacology and Physiology

2. Graduate Program in Neuroscience

Drexel University College of Medicine

Philadelphia, PA, 19102

*Address correspondence to:

Jacqueline M Barker, PhD

Department of Pharmacology and Physiology

Drexel University College of Medicine

245 N 15[th] Street

Philadelphia, PA 19102

215-762-8794

Jmb893@drexel.edu

Funding sources

NIH: F31AA031439 (CMCA); R21AA027629 (JMB); T32AA007462 (CC; trainee KGB)

Structured abstract and key words:

Background: In 2023, the World Health Organization declared that there is no safe amount of alcohol

consumption. It is becoming increasingly clear that even at lower doses, ethanol exposure impacts both

the brain and behavior. Emerging work has shown that chronic exposure to lower doses of ethanol may

lead to inflexible behaviors and promote aberrant reward seeking. This study aimed to determine the

impact of chronic, low-dose ethanol exposure on neural substrates of reward and associated alterations

in behavioral strategy in response to change in reward value.

Methods: Female and male C57BL/6J mice underwent operant training to self-administer 10% sucrose

under a fixed ratio schedule. After acquisition, response criteria were graduated to a random interval

schedule to promote automated behaviors. Throughout training, mice were given an i.p. injection of

low-dose ethanol (0.5g/kg) or saline, 1 hour after each training session. Mice did not receive low-dose

ethanol or saline i.p. injections after completion of training. Mice were then tested in a PR task in which

the reward magnitude of reinforcer was reduced (small or large) or increased (small or large). A subset

of mice expressed a retrograde tracer in the nucleus accumbens (NAc), and cFos expression within

NAc circuits was analyzed following a sucrose self-administration session.

Results: Chronic low-dose ethanol exposure altered behavioral responding in female mice following

small changes in reward magnitude. Female mice showed divergent response patterns when there was

a small reduction in reward magnitude, with greater proportions of ethanol-exposed female mice either

increasing or decreasing responding versus controls. Following a small increase, low-dose ethanol

female mice significantly increased responding versus controls. Female mice exposed to chronic low-

dose ethanol shifted behavioral strategy with a reduction in 'checking' behavior (magazine entry after a

lever press) in response to changes in reward magnitude. This effect was not observed in male mice.

Low-dose ethanol exposure altered cFos expression within the prelimbic cortex and its projections to

the NAc during reward seeking.

Conclusions: Chronic, low-dose ethanol altered behavioral responding and strategy in female mice in

response to changes in reward value. During reward seeking, low-dose ethanol exposure impacted

prelimbic cortex and its projections to the nucleus accumbens activity in both female and male mice.

Future studies should investigate the consequences of chronic low-dose ethanol on both the brain and

behavior to further understand what underlying processes drive aberrant reward-seeking behaviors.

Keywords: Motivation; ethanol; reward; sex differences; nucleus accumbens

Introduction

It is increasingly appreciated that even low doses of ethanol can impact health. This includes recognition from the World Health Organization that there is no safe amount of ethanol consumption as any level of ethanol consumption increases risk of negative health outcomes (Anderson et al., 2023). Despite this, consumption of ethanol is very common (WHO, 2024), necessitating a greater characterization of its impacts on brain and behavior. Low-dose ethanol exposure has distinct physiological effects from higher dose ethanol exposure. At lower doses, ethanol elicits a more stimulatory response, promoting disinhibition and relaxation, whereas at higher doses ethanol exposure induces sedative effects resulting in impaired motor function, coma and at severe levels, potentially death (Cui and Koob, 2017). Chronic exposure to lower doses of ethanol impacts both brain and behavior (Cui and Koob, 2017; Davidson et al., 1997; Daviet et al., 2022; Weafer and Fillmore, 2016). These neurobehavioral changes arising from chronic low-dose ethanol exposure contribute to increased risk for the development of alcohol use disorder (AUD) or other disorders associated with aberrant motivated behavior. Our previous work has shown that chronic, low-dose ethanol exposure enhanced motivation for a reward in a sex-specific manner (Bryant et al., 2022). However, it is not clear which aspects of motivated behavior and associated neurocircuitry are impacted by chronic low-dose ethanol exposure. Corticolimbic brain regions such as the prefrontal cortex (PFC) and basolateral amygdala (BLA) play a large role in cognitive processes necessary for flexible behavior and are known to be sensitive to higher doses of ethanol (Abernathy et al., 2010; Gass and Chandler, 2013; Klenowski et al., 2021; Koob and Volkow, 2016; Nippert et al., 2024; Pleil et al., 2015; Shields et al., 2022; Trantham-Davidson and Chandler, 2015). To expand on these findings, the impact of chronic low-dose ethanol exposure on behavioral sensitivity to changes in nondrug reward value was investigated using a progressive ratio task in which reward magnitude was either decreased or increased. The effect of chronic, low-dose ethanol on cFos expression within projections to the nucleus accumbens (NAc) was assessed following a sucrose self-administration session to determine the impact of low-dose ethanol

on NAc circuit engagement. These findings give insight to chronic low-dose ethanol exposure effects on

aberrant reward motivation and seeking.

Materials and Methods

Subjects

Adult male and female C57BL/6J mice from Jackson Laboratory (delivered at 9 weeks in age; $N = 71$

(35 females, 36 males; 1 female excluded due to failure to acquire the task) were used in the following

studies. All studies were performed in accordance with approved protocols from the Drexel University

Institutional Animal Care and Use Committee. All mice were housed in a vivarium with a standard

12h:12h light-dark cycle. Mice were acclimated to the facilities for at least one week before beginning

experiments. Prior to the start of experiments, mice were food restricted to approximately 90% of their

ad libitum weight, which lasted for the duration of the behavioral studies.

Surgeries

In order to investigate the effects of low-dose ethanol on NAc circuitry, 24 adult C57BL/6J mice ($N = 12$

males and 12 females) underwent stereotaxic surgery for microinjections of a retrograde GFP tagged

viral tracer (pAAV-hSyn-EGFP; Addgene, Watertown, MA; #50465) into the NAc. Surgeries were

performed using isoflurane anesthesia. Virus was injected (0.2 uL infusion at 0.1 uL/minute) targeting

1.5 mm anterior/posterior (A/P), + 0.6 mm medial/lateral (M/L) and -4.7 mm dorsal/ventral (D/V) to

bregma (0 mm). The syringe was left in place for 5 minutes for diffusion.

Behavior training

All behavioral experiments were conducted using standard Med-Associates (St. Albans, VT) operant

boxes. On one wall of each box there were two retractable levers on either side of a reward magazine

containing a slot for pellet or liquid reinforcer delivery. Cue lights were directly above each lever but

were not used in these experiments. On the opposite wall of each box, there were five nose pokes with

lights which were not used for any of the experiments. The back wall, ceiling and door of the box were

comprised of Plexiglas. All operant boxes were housed within sound attenuating chambers which had

ventilation by fan and white noise.

Mice underwent operant training to self-administer 10% sucrose (20 uL per reinforcer delivery) on a

fixed ratio 1 (FR1) schedule on two levers, where each lever press resulted in one reinforcer delivery.

Levers were presented consecutively, and mice had access to each lever for 15 minutes, with the order

of lever presentation counterbalanced across days for each animal. After acquisition of stable

responding on the FR1 schedule (> 15 presses on each lever for 3 consecutive days), the

reinforcement schedules were transitioned. One lever was assigned to be reinforced on a random

interval (RI) schedule, which has been shown to promote automated behaviors, and the other on a

variable ratio (VR) schedule, which promotes maintenance of goal-directed responding (Barker et al.,

2018; Bryant et al., 2022; Dickinson et al., 2002; Gremel and Costa, 2013). The RI lever was reinforced

at RI30 (the first lever press after, on average, 30 seconds is reinforced) for 3 days, followed by RI60

for 3 days. The VR lever was reinforced at VR5 (on average, the 5th lever press was reinforced) for 3

days followed by 3 days on a VR8 schedule. All progressive ratio testing was conducted on the lever

trained using the RI schedule as behavioral impacts of training on this schedule on reward motivation

have been observed following chronic, low-dose ethanol exposure (Bryant et al., 2022).

Low-dose ethanol exposure

Mice were matched by sex to receive low-dose ethanol (0.5 g/kg, i.p.) or saline vehicle injections 1 hour

after the initiation of each training session (with a mean of 22 days of training/exposure per animal).

The 1-hour time point was chosen to match previous literature looking at the effects of low-dose ethanol

exposure (Bryant et al., 2023, 2022). After training was completed, mice no longer received ethanol or

saline injections and thus went into progressive ratio (PR) testing with a history of low-dose ethanol

exposure.

Progressive ratio testing

In order to assess the impact of chronic low-dose ethanol exposure on motivated behavior under

changing reward conditions, 47 mice (24 males, 23 females) with a history of low-dose ethanol

exposure (or vehicle controls) underwent PR4 testing. In these sessions, only the lever trained on the

RI schedule was protracted. During the PR test, reinforcer delivery requirements progressively

increased, with the number of lever presses required for each successive reward progressing in

increments of 4 (1, 5, 9, 13, 17...). During training, reinforcer magnitude was 20 uL of 10% sucrose. To

assess the effects of a history of low-dose ethanol exposure on sensitivity to changes in reward value -

including increases vs. decreases in magnitude, and small vs. large changes - reinforcer magnitude

was altered across PR testing sessions. PR testing was counterbalanced to assess small changes

(from 20 uL to 15 or 25 uL) or large changes (from 20 uL to 5 or 30 uL) from the baseline reinforcer

volume. All testing was performed on the PR4 schedule and conducted across an 8-hour session until

animals reached a "breakpoint" where they discontinued pressing (no lever presses for 5 min) (Bryant

et al., 2022; Gourley et al., 2007).

Immunofluorescence

To probe the effects of chronic low-dose ethanol exposure on NAc circuits involved with reward

motivation and seeking, tissue generated from animals from a previously published behavioral data set

(Bryant et al., 2022, 2024) were utilized. Following the completion of PR4 testing, animals with a

retrograde tracer in the NAc underwent a 15-minute RI60 session (N = 12 males and 12 females). To

detect cFos protein expression, mice underwent transcardial perfusion 90 minutes from the midpoint of

the 15-minute session. Brains were postfixed in 4% paraformaldehyde solution then transferred to 30%

sucrose for cryoprotecting. Brain tissue was then cut at 40 microns in preparation for

immunofluorescence staining. Tissue was incubated in cFos rabbit (Abcam, Cambridge, UK;

#ab222699, 1:500 dilution) and GFP goat (Abcam, Cambridge, UK; #ab5450, 1:1000 dilution) primaries

in 4°C overnight. Tissue was then incubated in Alexa Fluor 594 anti-rabbit (Invitrogen, Carlsbad, CA;

#A32754, 1:250 dilution) and Alexa Fluor 488 anti-goat (Jackson ImmunoResearch Labs, West Grove,

PA; #705-545-147, 1:250 dilution) secondary antibodies for 2 hours at room temperature. Slices were

then sorted and mounted onto slides for immunofluorescence microscopy imaging using a Nikon

motorized microscope with 10x objective. Quantification of cFos, GFP and colocalization puncta

analysis was conducted by a scientist blind to all groups/conditions using ImageJ and normalized by

area of the region of interest. Labeled cells were quantified from slices at bregma -.95, -1.06, -1.23 mm

for BLA, +1.97, + 1.69, +1.53 mm for PFC, and +1.7, +1.4, and +1.1 mm for NAc (9 slices/mouse; 3

slices/region) using the Franklin & Paxinos 'The Mouse Brain Atlas', 3rd edition as reference.

Statistical analysis

All statistical analyses were conducted using GraphPad Prism. Differences in group means were

determined using unpaired t-tests or, for repeated behavioral testing, rmANOVA. To characterize

population differences in behavior, responding during the PR4 test sessions was calculated as percent

change. Using a quartile split, the mice that were in the highest 25% were identified as 'increasers', the

bottom 25% as 'decreasers', while the middle 50% were considered not to change their behavior. A chi

square was performed to analyze distribution of changes in breakpoint for each PR4 test (5uL, 15uL,

25uL and 35uL). 'Checking' behavior analysis was calculated for the complete PR session as well as

for each block within the PR session as effort requirement for reinforcer delivery increased for each PR

block using MATLAB (version R2022a; Barker Lab Github; https://github.com/bryantkg/Barker-lab-

code).

Results

Low-dose ethanol does not impact sucrose self-administration acquisition

To assess the effects of chronic low-dose ethanol exposure on sucrose self-administration acquisition

(FR1) [See behavioral timeline, **Fig 1A**], the number of days required to meet response criteria during

training was compared in saline versus low-dose ethanol-exposed mice. Ethanol did not impact days

required to reach acquisition criteria in female [unpaired t-test, t = 1.524, df = 21; p = 0.1424; **Fig 1B**] or

male mice [unpaired t-test: t = 0.4683, df = 22, p = 0.6442; **Fig 1C**]. Two-way ANOVA indicated no

effect of low-dose ethanol on lever responding across day in female [F(8,168) = 0.3668, p = 0.3933; **Fig**

1D] or male mice [F(8,176) = 1.030, p = 0.4154**; Fig 1E**]. A main effect of training days was observed in

females [F(3.147, 66.08) = 5.189, p = 0.0024, Greenhouse-Geisser corrected], with elevated

responding compared to day 1 on days 7, 8, and 9 [p's < 0.05 Sidak's]. A similar main effect of day was

observed in males [F(5.645, 124.2) = 7.607, p < 0.0001, Greenhouse-Geisser corrected], with elevated

responding vs day 1 on days 4, 6, 7, 8, and 9 [all p's < 0.05, Sidak's].

Low-dose ethanol alters reward-circuit engagement following reward seeking

To investigate sex-specific effects of low-dose ethanol on neural substrates involved with reward

motivation, we analyzed colocalization of a retrograde tracer injected within the NAc and cFos activity in

NAc-projecting regions following sucrose reward seeking (15-minute RI60 session) in male and female

mice who had a history of exposure to either saline or ethanol (**Fig 2A**). The percent of NAc-projecting

neurons from the BLA, prelimbic cortex (PrL) or infralimbic cortex (IL) which expressed cFos (% cFos+

and GFP+/total GFP+), the percent of all cFos expressing neurons that projected to the NAc from the

BLA, PrL or IL (% cFos+ and GFP+/total cFos+), and overall BLA, PrL and IL cFos+ cells were

quantified.

Within the BLA, there was no effect of sex or treatment on percent of BLA projections to the NAc that

expressed cFos [two-way ANOVA; *Sex:* F(1,21) = 0.3363, p = 0.5681; *EtOH*: F(1,21) = 0.7320, p =

0.4019; *Sex x EtOH*: F(1,21) = 0.2044, p = 0.6558; **Fig 2B**]. However, females exhibited a lower

percentage of cFos expressing cells that projected from the BLA to NAc than males [two-way ANOVA;

Sex: F(1,21) = 6.561, p = 0.0182]. There was no effect of ethanol treatment or interaction [*EtOH*:

F(1,21) = 1.268, p = 0.2728; *Sex x EtOH*: F(1,21) = 1.500, p = 0.2342; **Fig 2C**]. This stands in contrast

to greater total cFos expression in the BLA in females than males [two-way ANOVA: *Sex*: F(1,21) =

11.97, p=0.0023], potentially indicating activity in another BLA circuit in females [*EtOH*: F(1,21) =

0.3957, p=0.5361; *Sex x EtOH*: F(1,21) = 1.694, p = 0.2072; **Fig 2D**].

Within the PrL, there was a main effect of sex such that a greater percentage of PrL projections to NAc

expressed cFos in female mice as compared to male mice [two-way ANOVA: *Sex*: $F_{(1,21)}$ = 5.521, p =

0.0286; **Fig 2E**]. There was no effect of ethanol treatment or interaction [*EtOH*: $F_{(1,21)}$=0.9713,

p=0.3356; *Sex x EtOH*: $F_{(1,21)}$ = 0.1868, p = 0.6700; **Fig 2E**]. In contrast, for both female and males

there was a main effect of ethanol such that ethanol-exposed mice exhibited a lower percentage of

cFos expressing cells that projected from the PrL to the NAc [two-way ANOVA: *EtOH*: $F_{(1,21)}$ = 5.636,

p = 0.0277; *Sex x EtOH*: $F_{(1,21)}$ = 0.6829, p = 0.4184; **Fig 2F**] but a greater total of PrL cFos

expressing cells when compared to saline controls [two-way ANOVA: *EtOH*: $F_{(1,21)}$ = 6.622, p =

0.0177; *Sex x EtOH*: $F_{(1,21)}$ = 0.7450, p = 0.7450; **Fig 2G**]. There was no effect of sex on cFos

expressing cells projecting from the PrL to the NAc [two-way ANOVA: *Sex*: $F_{(1,21)}$ = 0.5879, p =

0.4522; **Fig 2F**] or total cFos expression within the PrL [two-way ANOVA: *Sex*: $F_{(1,21)}$ = 0.5569, p =

0.4638; **Fig 2G**].

Within the IL, there was no effect of sex or ethanol treatment on circuit-specific or total cFos expression

[percent of IL projections to the NAc projections that expressed cFos: two-way ANOVA: *Sex*: $F_{(1,21)}$ =

0.7489, p = 0.3976; *EtOH*: $F_{(1,21)}$ = 0.2997, p = 0.5905; *Sex x EtOH*: $F_{(1,21)}$ = 0.02830, p = 0.8682;

Fig 2H; percent total cFos that project to NAc: two-way ANOVA: *Sex*: $F_{(1,21)}$ = 2.794, p =

0.1110; *EtOH*: $F_{(1,21)}$ = 1.573, p = 0.2250; *Sex x EtOH*: $F_{(1,21)}$ = 0.1807, p = 0.6755; **Fig 2I**; total

cFos: two-way ANOVA: *Sex*: $F_{(1,21)}$ = 0.3470, p = 0.5624; *EtOH*: $F_{(1,21)}$=3.785, p=0.0659; *Sex x*

EtOH: $F_{(1,21)}$ =1.412, p = 0.2487; **Fig 2J**]. Overall, these findings show that the effects of low-dose

ethanol on NAc-projecting circuit engagement during reward seeking depend on sex and exposure and

differ across cortical and subcortical regions.

Chronic low-dose ethanol exposure alters sucrose reward motivation in female – but not male - mice

when changes in reward magnitude occur during progressive ratio testing

To investigate the effects of chronic, low-dose ethanol on reward motivation in response to changes in

reward magnitude, changes in breakpoints during PR testing were assessed. Control and low-dose

ethanol-exposed mice exhibited similar overall mean breakpoints in both males and females following

either a small reduction [*females:* t = 0.7273, df = 21, p = 0.4750; **Fig 3A**; *males:* t = 1.807, df = 22, p =

0.0845; **Fig 3B**] or a large reduction [*females:* t = 0.7309, df = 21, p = 0.4729; **Fig 3C**; *males*: t =

0.6052, df = 22, p = 0.5513; **Fig 3D**]. Investigation of individual differences showed that low-dose

ethanol exposure shifted behavioral response patterns in female mice. Specifically, an observed shift in

the distribution of female mice exhibiting changes in behavior was observed, where a greater proportion

of ethanol-exposed female mice changed their response patterns following a small reduction in reward

magnitude [X^2(2, 23) = 6.378, p = 0.0412; **Fig 3A**]. This was not observed in males [X^2 (2, 24) = 2.762,

p = 0.2514; **Fig 3B**]. Following a large reduction in reward magnitude, no differences in breakpoints

between control and low-dose ethanol-exposed females [X^2 (2, 23) = 2.071, p = 0.3551, **Fig 3C**] or low-

dose ethanol-exposed male mice were observed [X^2 (2, 24) = 0.2338, p = 0.8897; **Fig 3D**].

When there was a small increase in reward magnitude, low-dose ethanol-exposed female mice

exhibited significantly higher overall mean breakpoints compared to control female mice [unpaired t-

test, t = 2.103, df = 21, p = 0.0477; **Fig 4A**] this effect was not observed in male mice [t = 1.060, df =

22, p = 0.3006; **Fig 4B**]. When there was a large increase in reward magnitude, both female and male

mice responded similarly to controls [females: t = 0.4723, df = 21, p = 0.6416; **Fig 4C**; males; t =

0.2374, df = 22, p = 0.8145; **Fig 4D**]. When there was small increase in reward magnitude, both female

[(X^2 (2, 23) = 2.644, p = 0.2666, **Fig 4A**] and male mice [X^2 (2, 24) = 1.333, p = 0.5134, **Fig 4B**] did not

show significant differences in distribution in responding. Similarly, when there was a large increase in

reward magnitude, both females [(X^2 (2, 23) = 3.448, p = 0.1784**, Fig 4C**] and male [X^2 (2, 24) = 1.333,

p = 0.5134, **Fig 4D**] mice did not exhibit significant differences in population distribution versus controls.

These findings indicted that the effects of low-dose ethanol on sensitivity to changes in reward were

restricted to small changes (reduction or increase) as no effects were observed when there were large

changes (reduction or increase).

Low-dose ethanol exposure alters behavioral strategy in a sex-specific manner

To probe the effects of chronic low-dose ethanol exposure on underlying behavioral strategies during

PR testing, we quantified "checking" behavior as defined by percent of lever presses followed by a

magazine entry for the entire PR session as well as within session as each PR block progressed.

When there was a small reduction in reward magnitude, we observed a significant decrease in overall

checking behavior during the PR session [unpaired t-test: $t = 2.168$, df = 21, $p = 0.0418$; **Fig 5A (top)**]

as well as within session in ethanol-exposed female mice in comparison to controls [two-way

rmANOVA: *PR block*: $F_{(2.383,26.21)} = 1.323$, $p = 0.2863$, Greenhouse-Geisser corrected; *EtOH*:

$F_{(1,21)} = 5.438$, $p=0.0297$; *PR block x EtOH*: $F_{(10,110)} = 0.5659$, $p = 0.8385$; **Fig 5A (bottom)**]. This

main effect of treatment was not observed in males when there was a small reduction in reward

magnitude [$t = 0.08621$, df = 22, $p = 0.9321$, **Fig 5B (top)**; *PR block*: $F_{(2.041,29.13)} = 13.89$,

$p<0.0001$, Greenhouse-Geisser corrected; *EtOH*: $F_{(1,22)} = 0.2465$, $p = 0.6245$; *PR block x EtOH*:

$F_{(11,157)} = 0.3515$, $p = 0.9720$; **Fig 5B (bottom)**]. There was an effect of PR block in checking

behavior as the session progressed with a significant decrease in checking behavior during blocks with

lever requirement 9 to 45 in comparison to the 1[st] PR block [Dunnett's multiple comparison: all p's <

0.0259; **Fig 5B (bottom)**].

Chronic low-dose female mice had lower checking behavior than controls when there was a large

reduction in reward magnitude. This was observed both in overall [$t = 2.121$, df = 21, $p = 0.0460$; **Fig

5C (top)**] and within session analysis [*PR block*: $F_{(2.900, 36.25)} = 1.765$, $p = 0.1726$, Greenhouse-

Geisser corrected; *EtOH*: $F_{(1, 21)} = 8.279$, $p = 0.0090$; *PR block x EtOH*: $F_{(8, 100)} = 1.162$, $p = 0.3299$;

Fig 5C (bottom)]. There was no effect of ethanol exposure on overall percent magazine checking [$t =$

0.5698, df = 22, $p=0.5746$; **Fig 5D (top)**] or within PR session analysis in mice male when there was a

large reduction in reward magnitude [*PR block*: $F_{(2.624, 38.19)}$ = 4.873, p = 0.0077, Greenhouse-Geisser corrected; *EtOH*: $F_{(1, 22)}$ = 0.1189, p = 0.7335; *PR block x EtOH*: $F_{(9, 131)}$ = 0.3501, p = 0.9560; **Fig 5D (bottom)**]. There was a significant decrease in checking behavior, regardless of treatment, during the block in which the lever requirement was 25 in comparison to the 1st PR block of the session [p = 0.0290; **Fig 5D (bottom)**].

When there was a small increase in reward magnitude, there was a significant reduction in checking behavior by ethanol-exposed female mice in overall PR session [t = 2.471, df = 21, p = 0.0221; **Fig 6A (top)**] but not when within session checking behavior was analyzed to that of control female mice checking behavior [*PR block*: $F_{(2.109, 31.19)}$ = 14.41, p<0.0001, Greenhouse-Geisser corrected; ETOH: $F_{(1, 21)}$ = 3.453, p = 0.0772; *PR block x EtOH*: $F_{(14, 207)}$ = 1.405, p = 0.1528; **Fig 6A (bottom)**]. We did observe an effect of PR block with a decrease in checking behavior in blocks with a lever requirement of 13 to 57 in comparison to the 1st block regardless of treatment [all p < 0.0088; **Fig 6A (bottom)**]. This was no effect of low-dose ethanol on male mice overall checking behavior [t = 0.05214, df = 22, p = 0.9589; **Fig 6B (top)**] or within PR session analysis when there was a small increase in reward magnitude [*PR block*: $F_{(5.024, 74.70)}$ = 16.21, p<0.001, Greenhouse-Geisser corrected; *EtOH*: $F_{(1, 22)}$ = 1.540, p = 0.2277; *PR block x EtOH*: $F_{(15, 223)}$ = 0.5917, p = 0.8799; **Fig 6B (bottom)**]. We did observe an effect of PR block with a decrease in checking behavior during blocks with a lever requirement of 9 to 61 in comparison to the 1st PR block [all p's < 0.0029, **Fig 6B (bottom)**].

When there was a large increase in reward magnitude, there was a significant decrease in checking behavior in both overall session analysis [t = 2.476, df = 21, p = 0.0219; **Fig 6C (top)**] as well as within PR session analysis in ethanol-exposed female mice [*PR block*: $F_{(3.456, 52.99)}$ = 13.61, p < 0.0001, Greenhouse-Geisser corrected; *EtOH*: $F_{(1, 21)}$ = 7.804, p = 0.0109; *PR block x EtOH*: 12, 184) = 1.681, p = 0.0738; **Fig 6C (bottom)**]. There was an effect of PR block with reduced checking behavior in blocks with a lever requirement of 25,33,37,41,45 and 49 vs the 1st PR block of the session in female mice [all p's < 0.047; **Fig 6C (bottom)**]. In response to a large increase in reward magnitude, there was

no difference in percent 'checking' during overall session [t = 0.06169, df = 22, p = 0.9514, **Fig 6D**

(top)] or within PR session analysis in ethanol male mice in comparison to controls [*PR block*: 4.003,

59.76) = 17.76, p < 0.0001, Greenhouse-Geisser corrected; *EtOH*: $F_{(1, 22)}$ = 0.05818, p = 0.8116; *PR*

block x EtOH: $F_{(14, 209)}$ = 1.106, p = 0.3542; **Fig 6D (bottom)**]. There was a significant decrease in

checking behavior in male mice during blocks with a lever requirement of 9 to 57 vs the 1[st] PR block of

the session [all p's < 0.0380; **Fig 6D (bottom)**].

Discussion

There is increasing evidence of neurobiological and behavioral consequences of repeated or chronic

exposure to lower doses of ethanol, though this remains understudied. Previous findings have identified

alterations in motivated reward seeking and limbic circuitry (Bryant et al., 2022, 2023, 2024; Corbit et

al., 2012; Cui and Koob, 2017; Fillmore and Rush, 2001). To further characterize low-dose ethanol

effects on reward seeking and neural correlates, the current experiments investigated the

consequences of chronic, low-dose ethanol exposure on motivated reward seeking following changes

in reward magnitude. While no effects of low-dose ethanol exposure on acquisition of sucrose self-

administration were observed, low-dose ethanol had sex-specific effects on behavioral response to

change in reward magnitude, such that ethanol-exposed females – but not males – exhibited greater

behavioral shifts following changes in reward magnitude compared to ethanol-naive mice. This was

accompanied by a change in response strategy, with reduced reward magazine 'checking behavior' in

ethanol-exposed females, but not males, during self-administration.

The absence of any effects of low-dose ethanol exposure on the acquisition of sucrose self-

administration replicates our previous findings using a similar ethanol exposure procedure (Bryant et

al., 2022, 2023, 2024) and further aligns with findings that higher ethanol exposure using vapor

inhalation also does not impact overall sucrose self-administration (Barker et al., 2020; Renteria et al.,

2018). Despite the similar acquisition curves and overall response rates between ethanol-exposed and

control mice, the neural circuits engaged by reward seeking – as indicated by cFos expression – were

impacted by sex and/or a history of ethanol exposure, with differences observed in the basolateral

amygdala and prelimbic cortex. Consistent with our findings, cortico-limbic structures are particularly

vulnerable to chronic ethanol exposure (Abernathy et al., 2010; Gass and Chandler, 2013; Klenowski et

al., 2021; Koob and Volkow, 2016; Pleil et al., 2015; Shields et al., 2022; Stuber et al., 2011), with sex-

specific effects of higher dose ethanol exposure on cFos expression within NAc circuitry (Chan et al.,

2025).

The BLA, a key structure within the mammalian reward circuit, is thought to encode reward value and

contribute to flexible actions (Ambroggi et al., 2008; Keefer et al., 2022; Mahler and Berridge, 2012;

O'Doherty, 2003; Wassum, 2022). Activity within the BLA during reward seeking has been associated

with 'updating' changes in reward value with disruptions in BLA activity corresponding to impaired ability

to update behavior, potentially producing maladaptive reward seeking (Hinz et al., 2025; Jezzini and

Padoa-Schioppa, 2020; Keefer et al., 2022; Wassum, 2022; Wassum et al., 2012). BLA's function

associated with the ability to update behavior consistent with reward value may be mediated by BLA

projections to the NAc (Mahler and Berridge, 2012; Stuber et al., 2011; Wassum, 2022; Wassum and

Izquierdo, 2015; Wellman et al., 2005). In both non-human primates and rodents, disconnection of the

BLA and NAc impaired sensitivity to outcome devaluation, implicating this circuit in the ability to use

reward value to guide behavior (Ambroggi et al., 2008; Wellman et al., 2005).

The current findings identified sex differences in cFos expression following self-administration in the

BLA, independent of ethanol exposure. Despite overall higher cFos counts in the BLA of female mice

compared to males, cFos expression was lower in BLA projections to the NAc in females. These

findings not only identify greater activity of BLA neurons following self-administration in females, it also

may suggest that BLA circuits other than the projection to the NAc are recruited during self-

administration in female mice. This may be consistent with other findings of sex differences in BLA and

NAc excitability and plasticity, with females exhibiting greater excitability in BLA (Price and McCool,

2022, 2021). This may impact circuit recruitment as well as behavioral strategies in response to

changes in reward magnitude in a sex-specific manner (Copenhaver and LeGates, 2024; Price and

McCool, 2022, 2021). While chronic low-dose ethanol effects on BLA cFos were not observed, sex

differences in BLA cFos activity following self-administration may reflect differential sensitivity of female

mice to changes in reward value. However, the current studies only assessed cFos expression

following a standard self-administration session. It is possible that probing cFos expression following a

session in which reward magnitude was altered would have revealed effects of chronic low-dose

ethanol.

In contrast to findings in the BLA, where sex but not ethanol exposure history impacted cFos

expression, cFos counts in the PrL were mediated by both ethanol exposure and sex. Following self-

administration, male and female mice had similar total cFos expression within the PrL, however, female

mice exhibited greater cFos expression than males within PrL to NAc projections, regardless of ethanol

treatment. Total cFos expression was increased in the PrL of mice both male and female mice with a

history of chronic, low-dose ethanol exposure. Despite this, the percentage of PrL to NAc projections

expressing cFos was reduced in ethanol-exposed mice, suggesting that the increase overall PrL

increase in ethanol-exposed mice was driven by PrL projections to other targets. One function of the

PrL is evaluation of motivational conflicts or contexts that influence whether goal-directed actions

should be suppressed or initiated (Green and Bouton, 2021; Shipman et al., 2018) thus, the PrL is

essential for mediating flexible behavior and reward-driven decision making (Killcross and Coutureau,

2003; Wassum et al., 2012; Woon et al., 2020). While the IL is also implicated in the regulation of

reward seeking, no changes in IL cFos were observed in this paradigm, suggesting that these findings

to not generalize across the medial PFC. Though not quantified in the current manuscript, the PrL also

projects to the BLA. These projections are also involved in the regulation of reward seeking and may

contribute to low-dose ethanol effects on behavior (Cardinal et al., 2002; Land et al., 2014; Song et al.,

2020; Tavares et al., 2023). These changes in PrL activity following chronic low-dose ethanol exposure

may independently impact the ability to flexibly regulate behavior or may drive deficits through

interactions with downstream structures like the BLA. Mechanistic studies selectively modulating these

structures and circuits will provide a more complete causal perspective on the contribution of these

circuits to low-dose ethanol-induced behavioral alterations.

Reward motivation is determined by the value of the reward and the effort (cost) in seeking and

obtaining the reward (Albert-Lyons et al., 2024; Bissonette and Roesch, 2016; Roesch and Olson,

2007). The current findings suggest that even at lower doses, chronic ethanol may impact the flexible

relationship between effort and reward magnitude. Following either a large increase or decrease in

reward magnitude, no effect of chronic, low-dose ethanol on reward motivation in either male or female

mice in comparison to controls was observed. However, when changes in reward magnitude in either

direction (decreases or increases) were smaller, ethanol-exposed female mice exhibited greater

changes in behavior compared to controls. Following a small reduction in reward magnitude, low-dose

ethanol female mice exhibited divergent behavioral responses. One population of female mice

increased responding for the smaller reward, while another population reduced responding, suggesting

that while ethanol exposure increased behavioral sensitivity to small reward reductions in female mice,

this resulted in distinct behavioral phenotypes. In clinical studies, women have been reported to work

harder for less reward, displaying higher motivation in the pursuit of goals or reward to that of men

(Lewis et al., 2023). The current data suggest that at lower doses, ethanol may promote one similar

phenotype in which a subset of female mice were willing to work harder for less reward. These

divergent behavioral phenotypes observed in female mice may reflect vulnerabilities induced by

chronic, low-dose ethanol exposure. For example, failure to persist in seeking goals when reward value

is reduced may indicate risk for depression-related behaviors (Admon and Pizzagalli, 2015; Vrieze et

al., 2013). Future work should consider whether these divergent populations reflect risk for

neuropsychiatric illness, such as depression, or subsequent development of AUD.

In contrast to findings following reward reductions, low-dose ethanol-exposed female mice showed a

greater increase in reward motivation following a small increase in reward magnitude in comparison to

control female mice. This effect was not observed in chronic, low-dose ethanol-exposed male mice.

Notably, previous work has found an overall increase in reward motivation as measured on the PR in

low-dose ethanol-exposed male, but not female, mice (Bryant et al., 2023). Together, this suggests that

chronic, low-dose ethanol exposure may alter sensitivity to changes in reward value in females, while

having more general effects on reward motivation in males as observed in other higher dose ethanol

exposure studies (Radke et al., 2017; Risher et al., 2013; Somkuwar et al., 2018). Though effects of

ethanol on behavioral response were only observed following small changes in reward magnitude,

ethanol-exposed female mice exhibited overall reduced 'checking' behavior – defined as percent of

entries to the reward magazine that follow a lever press - across all PR test sessions. This may be

consistent with deficits in the use of action (lever press) outcome (reward delivery) contingencies to

guide behavior in low-dose ethanol-exposed female mice. However, analysis of within-session checking

behavior showed that chronic, low-dose ethanol female mice shifted checking behavior as the PR test

sessions progressed, suggesting monitoring of reward contingencies. This would be consistent with

previous work which found that female mice exposed to chronic, low-dose ethanol use checking

behavior to guide their behavior in response to changes in contingency (Barker et al., 2024) and greater

sensitivity to changes in outcome value in higher dose ethanol-exposed females (Barker et al., 2020;

Pickens et al., 2020). Thus, the shift in behavioral strategy observed here during changes of reward

magnitude may suggest increased reward sensitivity driven by increased goal-directed behavior in

female mice. This is potentially consistent with findings that male, but not female, adult rats exhibit

facilitated habit formation following higher doses of ethanol exposure (Barker et al., 2017), and when

self-administering lower doses of ethanol (Barker et al., 2010). Together, this may suggest divergent

effects of ethanol on value sensitivity and use of contingencies in males and females.

Caveats and future considerations

The current study characterized the impacts of low-dose ethanol on non-drug reward seeking. It is

possible that low-dose ethanol exposure would have differing effects on ethanol reward seeking as is

observed following higher doses of ethanol. For example, previous work has shown that female mice

subjected to chronic intermittent ethanol vapor inhalation increase reward sensitivity to ethanol (in the

absence of conflict), promoting enhanced reward seeking in a conditioned place preference paradigm

(Xie et al., 2019). It is also not clear how chronic, low-dose ethanol exposure would impact other effort

or motivation assays other than the PR test used in this study. An important consideration to note in our

current study is that ethanol exposure was administered by the experimenter in a bolus injection dose.

While this procedure enabled controlled administration of this targeted dose of ethanol to determine

effects on the brain and behavior in both males and females, and targeted timing of administration

relative to behavior, it is likely that self-administered ethanol would engage different circuitry. This

intraperitoneal injection method of delivery was selected to minimize stress of oral gavage, but this

yields distinct metabolic impacts. In addition, while this dosing procedure yields similar BECs in male

and female mice (Nothem et al., 2023), it is possible that the behavioral effects observed in females

might emerge in males following an alternative dose of ethanol.

These studies assessed correlates of neural activation following reward seeking, using cFos expression

was used as a marker of putative cellular activity during reward seeking. This allowed investigation of

NAc circuitry engagement following sucrose self-administration sessions, however, these experiments

did not provide fine temporal resolution on changes in neural activity or characterize activity following

changes in reward magnitude. It is important in future studies to probe changes in cellular activity

across training to capture as these changes occur prior to testing. In this study sex and low-dose

ethanol effects on PrL and BLA regional activity and their projections to NAc were identified. Future

work should incorporate other circuitry such as PrL to BLA as well as BLA to PrL to help further identify

the effects of low-dose ethanol on cortico-limbic structures involved with reward seeking and

motivation. Incorporation of mechanistic focused studies would be beneficial to determine if the

observed changes in substrate engagement and behavioral outputs were causally related.

Conclusions

These findings indicate that repeated exposure to low-dose ethanol induced sex-specific effects, with

ethanol-exposed female mice exhibiting alterations in motivated responding for sucrose and underlying

behavioral strategies. This was accompanied by alterations in cortico-limbic circuit engagement

following self-administration reward seeking, including low-dose ethanol effects on prelimbic activity in

both male and female mice. This work highlights the importance of studying the effects of lower dose

ethanol exposure on the brain and behavior to further understand how sex plays a role in the

development of ethanol-induced aberrant reward seeking which may lead to increased risk of negative

outcomes.

References

1. Abernathy, K., Chandler, L.J., Woodward, J.J., 2010. Alcohol and the Prefrontal Cortex. Int Rev Neurobiol 91, 289–320. https://doi.org/10.1016/S0074-7742(10)91009-X

2. Admon, R., Pizzagalli, D.A., 2015. Dysfunctional Reward Processing in Depression. Curr Opin Psychol 4, 114–118. https://doi.org/10.1016/J.COPSYC.2014.12.011

3. Albert-Lyons, R., Capan, S., Ng, K.H., Nautiyal, K.M., 2024. Reward value and internal state differentially drive impulsivity and motivation. Behavioural brain research 471, 115073. https://doi.org/10.1016/J.BBR.2024.115073

4. Alcohol [WWW Document], URL https://www.who.int/news-room/fact-sheets/detail/alcohol (accessed 5.14.25).

5. Ambroggi, F., Ishikawa, A., Fields, H.L., Nicola, S.M., 2008. Basolateral Amygdala Neurons Facilitate Reward-Seeking Behavior by Exciting Nucleus Accumbens Neurons. Neuron 59, 648–661. https://doi.org/10.1016/J.NEURON.2008.07.004

6. Anderson, B.O., Berdzuli, N., Ilbawi, A., Kestel, D., Kluge, H.P., Krech, R., Mikkelsen, B., Neufeld, M., Poznyak, V., Rekve, D., Slama, S., Tello, J., Ferreira-Borges, C., 2023. Health and cancer risks associated with low levels of alcohol consumption. Lancet Public Health 8, e6–e7. https://doi.org/10.1016/S2468-2667(22)00317-6

7. Barker, J.M., Bryant, K.G., Montiel-Ramos, A., Goldwasser, B., Chandler, L.J., 2020. Selective Deficits in Contingency-Driven Ethanol Seeking Following Chronic Ethanol Exposure in Male Mice. Alcohol Clin Exp Res 44, 1896–1905. https://doi.org/10.1111/ACER.14418

8. Barker, J.M., Bryant, K.G., Osborne, J.I., Chandler, L.J., 2017. Age and sex interact to mediate the effects of intermittent, high-dose ethanol exposure on behavioral flexibility. Front Pharmacol 8. https://doi.org/10.3389/FPHAR.2017.00450

9. Barker, J.M., Glen, W.B., Linsenbardt, D.N., Lapish, C.C., Chandler, L.J., 2018. Habitual Behavior

Is Mediated by a Shift in Response-Outcome Encoding by Infralimbic Cortex. eNeuro 4.

https://doi.org/10.1523/ENEURO.0337-17.2017

10. Barker, J.M., Torregrossa, M.M., Arnold, A.P., Taylor, J.R., 2010. Dissociation of Genetic and

Hormonal Influences on Sex Differences in Alcoholism-Related Behaviors. Journal of

Neuroscience 30, 9140–9144. https://doi.org/10.1523/JNEUROSCI.0548-10.2010

11. Bissonette, G.B., Roesch, M.R., 2016. Editorial: Neural Circuitry of Behavioral Flexibility:

Dopamine and Related Systems. Front Behav Neurosci 10.

https://doi.org/10.3389/FNBEH.2016.00006

12. Bryant, K.G., Nothem, M.A., Buck, L.A., Singh, B., Amin, S., Curran-Alfaro, C.M., Barker, J.M.,

2023. A History of Low-Dose Ethanol Shifts the Role of Ventral Hippocampus during Reward

Seeking in Male Mice. eNeuro 10. https://doi.org/10.1523/ENEURO.0087-23.2023

13. Bryant, Kathleen G, Singh, B., Barker, J.M., 2024. Sex and individual differences in the effect of

chronic low-dose ethanol on behavioral strategy selection. Alcohol: Clinical and Experimental

Research 48, 132–141. https://doi.org/10.1111/ACER.15218

14. Bryant, K.G., Singh, B., Barker, J.M., 2022. Reinforcement History Dependent Effects of Low Dose

Ethanol on Reward Motivation in Male and Female Mice. Front Behav Neurosci 16.

https://doi.org/10.3389/FNBEH.2022.875890

15. Cardinal, R.N., Parkinson, J.A., Hall, J., Everitt, B.J., 2002. Emotion and motivation: The role of the

amygdala, ventral striatum, and prefrontal cortex. Neurosci Biobehav Rev 26, 321–352.

https://doi.org/10.1016/S0149-7634(02)00007-6

16. Chan, A.E., Driscoll, G.S., Usmani, Z., Ozburn, A.R., 2025. Nucleus accumbens core

chemogenetic excitation in male mice and chemogenetic inhibition in female mice reduced ethanol

reward. Biol Sex Differ 16, 1–16. https://doi.org/10.1186/S13293-025-00745-0/FIGURES/5

17. Copenhaver, A.E., LeGates, T.A., 2024. Sex-specific mechanisms underlie long-term potentiation

at hippocampus-nucleus accumbens synapses. bioRxiv.

https://doi.org/10.1101/2024.01.15.575709

18. Corbit, L.H., Nie, H., Janak, P.H., 2012. Habitual alcohol seeking: time course and the contribution

of subregions of the dorsal striatum. Biol Psychiatry 72, 389–395.

https://doi.org/10.1016/J.BIOPSYCH.2012.02.024

19. Cui, C., Koob, G.F., 2017. Titrating Tipsy Targets: The Neurobiology of Low-Dose Alcohol. Trends

Pharmacol Sci 38, 556–568. https://doi.org/10.1016/J.TIPS.2017.03.002

20. Davidson, D., Camara, P., Swift, R., 1997. Behavioral Effects and Pharmacokinetics of Low-Dose

Intravenous Alcohol in Humans. Alcohol Clin Exp Res 21, 1294–1299.

https://doi.org/10.1111/J.1530-0277.1997.TB04451.X

21. Daviet, R., Aydogan, G., Jagannathan, K., Spilka, N., Koellinger, P.D., Kranzler, H.R., Nave, G.,

Wetherill, R.R., 2022. Associations between alcohol consumption and gray and white matter

volumes in the UK Biobank. Nat Commun 13. https://doi.org/10.1038/S41467-022-28735-5

22. Dickinson, A., Wood, N., Smith, J.W., 2002. Alcohol seeking by rats: action or habit? Q J Exp

Psychol B 55, 331–348. https://doi.org/10.1080/0272499024400016

23. Fillmore, M.T., Rush, C.R., 2001. Alcohol effects on inhibitory and activational response strategies

in the acquisition of alcohol and other reinforcers: Priming the motivation to drink. J Stud Alcohol

62, 646–656. https://doi.org/10.15288/JSA.2001.62.646

24. Gass, J.T., Chandler, L.J., 2013. The Plasticity of Extinction: Contribution of the Prefrontal Cortex

in Treating Addiction through Inhibitory Learning. Front Psychiatry 4.

https://doi.org/10.3389/FPSYT.2013.00046

25. Gourley, S.L., Wu, F.J., Kiraly, D.D., Ploski, J.E., Kedves, A.T., Duman, R.S., Taylor, J.R., 2007.

REGIONALLY-SPECIFIC REGULATION OF pERK1/2 MAP KINASE IN A MODEL OF

ANTIDEPRESSANT-SENSITIVE CHRONIC DEPRESSION. Biol Psychiatry 63, 353.

https://doi.org/10.1016/J.BIOPSYCH.2007.07.016

26. Green, J.T., Bouton, M.E., 2021. New functions of the rodent prelimbic and infralimbic cortex in

instrumental behavior. Neurobiol Learn Mem 185. https://doi.org/10.1016/j.nlm.2021.107533

27. Gremel, C.M., Costa, R.M., 2013. Orbitofrontal and striatal circuits dynamically encode the shift

between goal-directed and habitual actions. Nat Commun 4.

https://doi.org/10.1038/NCOMMS3264

28. Hinz, J., Mahn, M., Müller, S., Szőnyi, A., Eichlisberger, T., Lüthi, A., 2025. Stimulus-specific and

adaptive value representations in the basolateral amygdala in male mice. Nat Commun 16.

https://doi.org/10.1038/S41467-025-60414-Z

29. Jezzini, A., Padoa-Schioppa, C., 2020. Neuronal Activity in the Primate Amygdala during Economic

Choice. J Neurosci 40, 1286–1301. https://doi.org/10.1523/JNEUROSCI.0961-19.2019

30. Keefer, S.E., Kochli, D.E., Calu, D.J., 2022. Inactivation of the Basolateral Amygdala to Insular

Cortex Pathway Makes Sign-Tracking Sensitive to Outcome Devaluation. eNeuro 9,

ENEURO.0156-22.2022. https://doi.org/10.1523/ENEURO.0156-22.2022

31. Killcross, S., Coutureau, E., 2003. Coordination of Actions and Habits in the Medial Prefrontal

Cortex of Rats. Cerebral Cortex 13, 400–408. https://doi.org/10.1093/CERCOR/13.4.400

32. Klenowski, P.M., Fogarty, M.J., Drieberg-Thompson, J.R., Bellingham, M.C., Bartlett, S.E., 2021.

Reduced Inhibitory Inputs On Basolateral Amygdala Principal Neurons Following Long-Term

Alcohol Consumption. Neuroscience 452, 219–227.

https://doi.org/10.1016/J.NEUROSCIENCE.2020.10.039

33. Koob, G.F., Volkow, N.D., 2016. Neurobiology of addiction: a neurocircuitry analysis. Lancet

Psychiatry 3, 760–773. https://doi.org/10.1016/S2215-0366(16)00104-8

34. Land, B.B., Narayanan, N.S., Liu, R.J., Gianessi, C.A., Brayton, C.E., M Grimaldi, D., Sarhan, M.,

Guarnieri, D.J., Deisseroth, K., Aghajanian, G.K., Dileone, R.J., 2014. Medial prefrontal D1

dopamine neurons control food intake. Nature Neuroscience 2014 17:2 17, 248–253.

https://doi.org/10.1038/nn.3625

35. Lewis, C.A., Grahlow, M., Kühnel, A., Derntl, B., Kroemer, N.B., 2023. Women compared with men
work harder for small rewards. Sci Rep 13, 1–10. https://doi.org/10.1038/S41598-023-32391-
0;SUBJMETA=2811,477,631;KWRD=HUMAN+BEHAVIOUR,PSYCHOLOGY

36. Mahler, S. V., Berridge, K.C., 2012a. What and when to "want"? Amygdala-based focusing of
incentive salience upon sugar and sex. Psychopharmacology (Berl) 221, 407–426.
https://doi.org/10.1007/s00213-011-2588-6

37. Nippert, K.E., Rowland, C.P., Vazey, E.M., Moorman, D.E., 2024. Alcohol, flexible behavior, and
the prefrontal cortex: Functional changes underlying impaired cognitive flexibility.
Neuropharmacology 260. https://doi.org/10.1016/J.NEUROPHARM.2024.110114

38. Nothem, M.A., Wickman, J.R., Giacometti, L.L., Barker, J.M., 2023. Effects of ethanol on
mechanical allodynia and dynamic weight bearing in male and female mice with spared nerve
injury. Alcohol: Clinical and Experimental Research 47, 382–394.
https://doi.org/10.1111/ACER.14997

39. O'Doherty, J., 2003. Can't learn without you: Predictive value coding in orbitofrontal cortex requires
the basolateral amygdala. Neuron 39, 731–733. https://doi.org/10.1016/S0896-6273(03)00525-7

40. Pickens, C.L., Cook, A., Gaeddert, B., 2020. Dose-dependent effects of alcohol injections on
omission-contingency learning have an inverted-U pattern. Behavioural brain research 392,
112736. https://doi.org/10.1016/J.BBR.2020.112736

41. Pleil, K.E., Lowery-Gionta, E.G., Crowley, N.A., Li, C., Marcinkiewcz, C.A., Rose, J.H., McCall,
N.M., Maldonado-Devincci, A.M., Morrow, A.L., Jones, S.R., Kash, T.L., 2015. Effects of chronic
ethanol exposure on neuronal function in the prefrontal cortex and extended amygdala.
Neuropharmacology 99, 735. https://doi.org/10.1016/J.NEUROPHARM.2015.06.017

42. Price, M.E., McCool, B.A., 2022. Chronic Alcohol Dysregulates Glutamatergic Function in the
Basolateral Amygdala in a Projection-and Sex-Specific Manner. Front Cell Neurosci 16, 857550.
https://doi.org/10.3389/FNCEL.2022.857550/BIBTEX

43. Price, M.E., McCool, B.A., 2021. Structural, Functional, and Behavioral Significance of Sex and

Gonadal Hormones in the Basolateral Amygdala: A Review of Preclinical Literature. Alcohol 98, 25.

https://doi.org/10.1016/J.ALCOHOL.2021.08.001

44. Radke, A.K., Jury, N.J., Kocharian, A., Marcinkiewcz, C.A., Lowery-Gionta, E.G., Pleil, K.E.,

McElligott, Z.A., McKlveen, J.M., Kash, T.L., Holmes, A., 2017. Chronic EtOH effects on putative

measures of compulsive behavior in mice. Addiction biology 22, 423–434.

https://doi.org/10.1111/ADB.12342

45. Renteria, R., Baltz, E.T., Gremel, C.M., 2018. Chronic alcohol exposure disrupts top-down control

over basal ganglia action selection to produce habits. Nature Communications 2018 9:1 9, 1–11.

https://doi.org/10.1038/s41467-017-02615-9

46. Risher, M.L., Fleming, R.L., Boutros, N., Semenova, S., Wilson, W.A., Levin, E.D., Markou, A.,

Swartzwelder, H.S., Acheson, S.K., 2013. Long-Term Effects of Chronic Intermittent Ethanol

Exposure in Adolescent and Adult Rats: Radial-Arm Maze Performance and Operant Food

Reinforced Responding. PLoS One 8, e62940. https://doi.org/10.1371/JOURNAL.PONE.0062940

47. Roesch, M.R., Olson, C.R., 2007. Neuronal activity related to anticipated reward in frontal cortex:

does it represent value or reflect motivation? Ann N Y Acad Sci 1121, 431–446.

https://doi.org/10.1196/ANNALS.1401.004

48. Shields, C.N., Baltz, E.T., Valencia, M.L., Gremel, C.M., 2022. Effects of chronic alcohol exposure

on motivation-based value updating. Alcohol 101, 53–64.

https://doi.org/10.1016/J.ALCOHOL.2022.03.004

49. Shipman, M.L., Trask, S., Bouton, M.E., Green, J.T., 2018. Inactivation of prelimbic and infralimbic

cortex respectively affects minimally-trained and extensively-trained goal-directed actions.

Neurobiol Learn Mem 155, 164. https://doi.org/10.1016/J.NLM.2018.07.010

50. Somkuwar, S.S., Quach, L.W., Quigley, J.A., Purohit, D.C., Fannon, M.J., Koob, G.F., Mandyam,

C.D., 2018. Ethanol Reinforcement Elicits Novel Response Inhibition Behavior in a Rat Model of

Ethanol Dependence. Brain Sciences 2018, Vol. 8, Page 119 8, 119.

https://doi.org/10.3390/BRAINSCI8070119

51. Song, Y., Wang, J., Lukasiewicz, T., Xu, Z., Zhang, S., Wojcicki, A., Xu, M., 2020. Mega-reward:

Achieving human-level play without extrinsic rewards. AAAI 2020 - 34th AAAI Conference on

Artificial Intelligence 5826–5833. https://doi.org/10.1609/AAAI.V34I04.6040

52. Stuber, G.D., Sparta, D.R., Stamatakis, A.M., Van Leeuwen, W.A., Hardjoprajitno, J.E., Cho, S.,

Tye, K.M., Kempadoo, K.A., Zhang, F., Deisseroth, K., Bonci, A., 2011. Excitatory transmission

from the amygdala to nucleus accumbens facilitates reward seeking. Nature 2011 475:7356 475,

377–380. https://doi.org/10.1038/nature10194

53. Tavares, G.E.B., Bianchi, P.C., Yokoyama, T.S., Palombo, P., Cruz, F.C., 2023. Involvement of

cortical projections to basolateral amygdala in context-induced reinstatement of ethanol-seeking in

rats. Behavioural Brain Research 448, 114435. https://doi.org/10.1016/J.BBR.2023.114435

54. Trantham-Davidson, H., Chandler, L.J., 2015. Alcohol-induced alterations in dopamine modulation

of prefrontal activity. Alcohol 49, 773–779. https://doi.org/10.1016/J.ALCOHOL.2015.09.001

55. Vrieze, E., Pizzagalli, D.A., Demyttenaere, K., Hompes, T., Sienaert, P., De Boer, P., Schmidt, M.,

Claes, S., 2013. Reduced reward learning predicts outcome in major depressive disorder. Biol

Psychiatry 73, 639–645. https://doi.org/10.1016/J.BIOPSYCH.2012.10.014

56. Wassum, K.M., 2022. Amygdala-cortical collaboration in reward learning and decision making.

Elife 11. https://doi.org/10.7554/ELIFE.80926

57. Wassum, K.M., Izquierdo, A., 2015. The basolateral amygdala in reward learning and addiction.

Neurosci Biobehav Rev 57, 271. https://doi.org/10.1016/J.NEUBIOREV.2015.08.017

58. Wassum, K.M., Tolosa, V.M., Tseng, T.C., Balleine, B.W., Monbouquette, H.G., Maidment, N.T.,

2012. Transient Extracellular Glutamate Events in the Basolateral Amygdala Track Reward-

Seeking Actions. Journal of Neuroscience 32, 2734–2746.

https://doi.org/10.1523/JNEUROSCI.5780-11.2012

59. Weafer, J., Fillmore, M.T., 2016. Low-Dose Alcohol Effects on Measures of Inhibitory Control,

Delay Discounting, and Risk-Taking. Curr Addict Rep 3, 75–84. https://doi.org/10.1007/S40429-

016-0086-Y

60. Wellman, L.L., Gale, K., Malkova, L., 2005. GABAA-Mediated Inhibition of Basolateral Amygdala

Blocks Reward Devaluation in Macaques. Journal of Neuroscience 25, 4577–4586.

https://doi.org/10.1523/JNEUROSCI.2257-04.2005

61. Woon, E.P., Sequeira, M.K., Barbee, B.R., Gourley, S.L., 2020. Involvement of the rodent prelimbic

and medial orbitofrontal cortices in goal-directed action: A brief review. J Neurosci Res 98, 1020–

1030. https://doi.org/10.1002/JNR.24567

62. Xie, Q., Buck, L.A., Bryant, K.G., Barker, J.M., 2019. Sex Differences in Ethanol Reward Seeking

Under Conflict in Mice. Alcohol Clin Exp Res 43, 1556–1566. https://doi.org/10.1111/ACER.14070

Figure Legends

Figure 1. Low-dose ethanol did not impact acquisition. (**A**) Timeline of behavioral experiment. (**B,C**)

Low-dose ethanol exposure did not alter the number of days required to achieve stable responding on

an FR1 schedule in either female (**B**) or male (**C**) mice. (**D, E**) Low-dose ethanol exposure did not alter

number of lever presses across training schedules in female (**D**) or male (**E**) mice. Females- saline: n =

12, ethanol: m = 11; males- saline: n = 12, ethanol: n = 12. Data represent mean +/- SEM.

Figure 2: Low-dose ethanol had brain-region specific effects on neural activity. (**A**) Timeline of

stereotaxic surgery, chronic low-dose ethanol exposure during behavioral training and sacrifice post

reward seeking session for cFos immunofluorescence staining. (**B**) Neither a history of low-dose

ethanol exposure nor sex altered percent of BLA projections to the NAC expressing cFos. (**C**) The

proportion of cFos expressing cells projecting to the NAc within the BLA was greater in males than

females, with no effect of ethanol exposure. (**D**) Female mice had a greater total of cells within the BLA

expressing cFos than male mice, with no effect of ethanol exposure. (**E**) Female mice showed greater

percent of PrL cells projecting to the NAc expressing cFos than males. (**F**) Ethanol-exposed mice

showed reduction in PrL cells expressing cFos which projected to the NAc to that of controls. (**G**)

Ethanol-exposed mice had a greater number of cells within the PrL expressing cFos than controls.

Neither a history of low-dose ethanol exposure nor sex altered percent of IL projecting cells to the NAc

expressing cFos (**H**) percent of cFos expressing cells which project to the NAc within the IL (**I**) or total

cells within the IL expressing cFos (**J**). **$p < 0.01$, *$p < 0.05$. Females- saline: n = 6, ethanol: n = 6;

males- saline: n = 6, ethanol: n = 6, Data represent +/- SEM.

Figure 3: Low-dose ethanol altered sensitivity to small reductions in reward magnitude in female

mice. (**A**) Female mice exposed to low-dose ethanol exhibited divergent responding when a small

reduction in reward magnitude occurred. (**B**) Low-dose ethanol exposure had no effect on responding

in male mice when there was a small reduction in reward magnitude. (**C**, **D**) When there was a large

reduction in reward magnitude, there was no effect of ethanol in either female or male mice. X^2, *p <

0.05. Females- saline: n = 12, ethanol: n = 11; males- saline n = 12, ethanol: n = 12, dashed line

represent upper and lower quartiles with solid line representing mean.

Figure 4: Low-dose ethanol altered responding in female mice when there was a small increase in

reward magnitude. (**A**) Ethanol-exposed female mice increased responding when there was a small

increase in reward magnitude in comparison to saline female mice. (**B**) Low-dose ethanol did not alter

male mice responding when there was a small increase in reward magnitude. (**C**, **D**) Low-dose ethanol

did not alter responding in female or male mice when there was a large increase in reward magnitude.

*p < 0.05. Females- saline: n = 12, ethanol: n = 11; males- saline n = 12, ethanol: n = 12, dashed line

represent upper and lower quartiles with solid line representing mean.

Figure 5: Analysis of 'checking' behavior during progressive ratio (PR) 4 testing when there was a

reduction in reward magnitude. (**A**)Ethanol-exposed female mice reduced percent checking behavior

during total (top) and within (bottom) PR testing session when there was a small reduction in reward

magnitude. (**B**) Low-dose ethanol exposure did not affect percent checking behavior in male mice for

total session or within session analysis when there was a small reduction in reward magnitude. (**C**)

Ethanol-exposed female mice reduced percent checking behavior in total session (top) and within

(bottom) PR testing session analysis when there was a large reduction in reward magnitude. (**D**) There

was no effect of low-dose ethanol exposure on checking behavior in male mice, for the total or within

PR4 testing session analysis, when there was a large reduction in reward magnitude. **p < 0.01, *p <

0.05. Females- saline: n = 12, ethanol: m = 11; males- saline: n = 12, ethanol: n = 12. Data represent

mean +/- SEM.

Figure 6: Analysis of 'checking' behavior during progressive ratio (PR) 4 testing when an increase in

reward magnitude occurred. (**A**) Ethanol-exposed female mice reduced percent checking behavior in

total (top) but not within PR testing session analysis when there was a small increase in reward

magnitude. (**B**) Low-dose ethanol did not impact percent checking behavior in male mice in total

session or within PR4 testing session analysis when there was a small increase in reward magnitude.

(**C**) Ethanol-exposed female mice reduced percent checking behavior for total (top) and within (bottom)

PR testing session analysis when there was a large increase in reward magnitude. (**D**) Low-dose

ethanol exposure did not impact checking behavior in male mice, in total or within PR4 session

analysis, when there was a large reduction in reward magnitude. ***$p < 0.001$, *$p < 0.01$. Females-

saline: n = 12, ethanol: m = 11; males- saline: n = 12, ethanol: n = 12. Data represent mean +/- SEM.

Females   *Males*

Sm. reduction

A — % mag. followed by lever press; Saline vs EtOH; *

B — % mag. followed by lever press; Saline vs EtOH

%checking vs Response requirement (PR blocks); Saline, EtOH; *

Lg. reduction

C — % mag. followed by lever press; Saline vs EtOH; *

D — % mag. followed by lever press; Saline vs EtOH

%checking vs Response requirement (PR blocks); Saline, EtOH; **