# A human lysosomal storage disorder toolkit for decoding proteome landscapes in cortical and dopaminergic-like induced neurons

Felix Kraus[1], Yuchen He[1], Yizhi Jiang[1,2], Delong Li[2,3], Yohannes A. Ambaw[4], Federico M. Gasparoli[1], Joao A. Paulo[1], Tobias C. Walther[4,5], Robert V. Farese, Jr.[4], Steven P. Gygi[1], Florian Wilfling[2,3] and J. Wade Harper[1,2]*

[1]Department of Cell Biology, Harvard Medical School, Boston MA, USA

[2]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815, USA

[3]Mechanisms of Cellular Quality Control, Max Planck Institute of Biophysics, Frankfurt, Germany.

[4]Cell Biology Program, Sloan Kettering Institute, New York, NY, USA.

[5]Howard Hughes Medical Institute, New York, NY, USA

*J. Wade Harper

**Email:** wade_harper@hms.harvard.edu

PNAS requires the corresponding author to provide an ORCID identifier at submission and strongly encourages all authors to use an ORCID ID. Do not include ORCIDs in the manuscript file; individual authors must link their ORCID account to their PNAS profile at www.pnascentral.org. For proper authentication, authors are not permitted to add ORCIDs on proofs. Learn more or register for ORCID.

**Author Contributions:** Conceptualization: F.K., J.W.H.; Investigation: F.K., Y.H., D.L., J.A.P., Y.J., Y.A.A; Analysis: F.K., D.L., F.M.G., J.W.H.; Visualization: F.K. Supervision: R.V.F., T.C.W., S.P.G., F.W., J.W.H.; Writing (original draft): F.K., J.W.H.; Writing (reviewing and editing): F.K., J.W.H., with input from all authors.

**Competing Interest Statement:** J.W.H. is a co-founder of Caraway Therapeutics, a subsidiary of Merck & Co., Inc., Rahway, NJ, USA (Caraway) and is a scientific advisory board member for Lyterian Therapeutics. This publication is unrelated to these competing interests. All other authors have no competing interests to declare.

**Keywords:** Proteomics, Organelle quality control, iNeuron, iDA, Lysosomal Storage Disorder

**Abstract**

Lysosomes maintain cellular homeostasis by degrading proteins delivered via endocytosis and autophagy, and by recycling building blocks for organelle biogenesis. Lysosomal Storage Disorders (LSDs) comprise a broad group of diseases affecting diverse lysosomal functions. To facilitate molecular phenotyping across diverse LSD gene classes, we are developing a library of human embryonic stem cells engineered to lack individual LSD genes. Here, we report the generation and initial analysis of a first-generation toolkit of ES cells lacking one of 23 LSD genes, including the majority of genes associated with sphingolipidoses and neuronal ceroid lipofucinoses. Global proteomic analysis of induced cortical-like and midbrain dopaminergic-like neurons coupled with in-depth abundance and correlation profiling across organelles and sub-organelle components revealed potential vulnerabilities that reflect distinct patterns of proteome alterations across both genotypes and neuronal cell types. We characterize alterations in the mitochondrial proteome associated with GBA1 and ASAH1 deficiency, and identify synaptic defects in $ASAH1^{-/-}$ induced neurons that correlate with defects in neuronal firing rates. Moreover, we developed an informatic pipeline for proteome-wide identification of individual protein interactions and protein complexes that may be disrupted as a result of LSD gene deficiency. Finally, we visualized structural alterations of *ASAH1*-deficient endolysosomes *in situ* using cryo-electron tomography, revealing swollen organelles that were largely devoid of dense internal membranes characteristic of wild-type cells, but containing numerous intralumenal vesicle compartments. This toolkit and associated proteomic landscapes provide a resource for defining molecular signatures associated with LSD gene dysfunction and organelle vulnerability.

**Introduction**

Lysosomes are a primary degradative membrane-bound compartment within cells, and contain hydrolases that digest glycoproteins and lipids delivered via endocytosis, as well as proteins and organelles delivered via autophagy, thereby facilitating the recycling of molecular building blocks(1, 2). Additionally, lysosomes contain transporters responsible for trafficking of recycled building blocks to other cellular compartments. As such, lysosomes function as important signaling hubs for nutrient availability via the MTOR pathway(1). Genetic and clinical studies have identified ~70 distinct types of lysosomal storage disorders (LSDs) that map to at least 54 genes directly or indirectly linked with lumenal catabolic functions. LSDs are primarily classified as sphingolipidoses, mucopolysaccharidoses, glycoproteinoses, and neuronal ceroid lipofuscinoses (NCL), based on the type of storage material that accumulates within lysosomes (e.g. glycosphingolipids, or cholesterol) (3). Additional storage material can also accumulate as a secondary response. However, how diverse primary and secondary storage material accumulation interfaces with distinct multisystem disease phenotypes is poorly understood. One possibility is that various LSD risk alleles create molecular vulnerabilities that can be mapped onto organelles and pathways in a cell-type dependent manner, potentially providing a deeper understanding of LSD pathogenesis mechanisms.

As an initial approach to addressing this knowledge gap, we previously reported a library of more than 2 dozen HeLa cell lines lacking genes linked with LSDs, facilitating global lipidomic and proteomic profiling and revealing patterns of alterations in autophagy and cholesterol trafficking systems with specific LSD mutations (4). Here, we extend this approach through the development of a versatile toolkit of human embryonic stem (ES) cell lines lacking one of 23 LSD genes, facilitating the generation of cortical-like and midbrain dopaminergic-like induced neuron (iN and iDA) models for global longitudinal molecular phenotyping. The current toolkit is focused on genes linked with sphingolipidoses, NCL, and integral membrane protein disorder classes. The focus on sphingolipidoses and NCL genes reflects: 1) an emerging framework for understanding the mechanisms and phenotypes associated with defects in sialic-acid-containing glycosphingolipid breakdown on intraluminal vesicles (ILVs) formed within endolysosomes, including the role of bis(monoacylglycerol)phosphate (BMP) lipids in this process (5-8), 2) the proposed genetic linkage between glycosphingolipid breakdown enzymes GBA1, SMPD1, ASAH1 and Parkinson's disease (PD) (9-18), and 3) the finding that specific NCL genes, including CLN11 (also called GRN), may regulate the abundance of BMP within lysosomes, with loss of BMP resulting in the accumulation of GM2 and GM3 gangliosides (7). Using deep global proteome analysis of day 50 iN and iDA cells coupled with informatic pipelines for decoding organelle protein abundance landscapes and underlying protein-protein interaction (PPI) networks, we identify patterns of proteome alterations that may define functional vulnerabilities associated with specific LSD genes. Candidate vulnerabilities include alterations in mitochondrial proteomes particularly in $GBA1^{-/-}$ iDA cells, as previously observed (19, 20), and synaptic proteomes in $ASAH1^{-/-}$ cells, with $ASAH1^{-/-}$ iDA cells displaying reduced neuronal firing. Finally, to understand the impact of ASAH1 deficiency on lysosomal ultrastructure, we performed cryo-electron tomography (cryo-ET) on lysosomal structures in the HeLa cell system, revealing swollen structures largely devoid of lamellar membrane material typically found in lysosomal structures in wild-type HeLa cells. This toolkit and proteomic data provides a resource for the field.

**Results**

**A Stem Cell Toolkit for Systematic Analysis of LSD Genes**
To systematically examine potential alterations in the properties of organellar systems in the context of LSDs, we developed a versatile stem cell loss of function toolkit. We employed an H9 human embryonic stem cell (hESC)-based system (4, 21) containing an inducible AAVS1-NGN2 cassette for efficient neuronal differentiation and endogenously tagged TMEM192-3xHA (heterozygous) for LysoIP (H9^NGN2;TMEM192-HA). These cells, referred to hereafter as "Control" cells,

can be efficiently convert to cortical-like iNeurons (iN) or midbrain-like dopaminergic neurons (iDA)
using established methods (22, 23), as accessed by immunofluorescence with $\alpha$-MAP2 and $\alpha$-TH
(Tyrosine Hydroxylase) (**Fig. S1A**). The initial version of this toolkit includes inactivating indels in
23 LSD genes, including genes implicated in sphingolipidoses (*GBA1*, *ASAH1*, *HEXA*, *HEXB*,
*PSAP*, *SMPD1*), NCL (*CLN1*, *CLN2*, *CLN3*, *DNAJC5*, *CLN5*, *CLN6*, *CLN7*, *CLN8*, *ATP13A2*, *GRN*,
*CTSD*, *CTSF*), Mucolipidosis type IV (*MCOLN1* (encoding TRPML1), as well as Niemann-Pick
Type-C (*NPC1*, *NPC2*), *LIPA* (Wolman's disease) and *GAA* (Pompe's disease). CRISPR RNPs
were transfected into Control cells and homozygous clones with frameshift indels were identified
by locus sequencing (see **Dataset S1** for a detailed analysis of indels and clone identification).
Proteomic data described below broadly confirms loss of function mutations, and points to the utility
of this system for our ongoing effort to create a complete LSD toolkit.

**Proteomic Fingerprinting of LSD Mutant iN and iDA Neurons**
Our previous work has indicated the utility of total proteome analysis to reveal alterations in
organelle and pathway abundance in response to loss of specific pathways (e.g. autophagy) or
during changes in cell state (24-26). To profile alterations in the proteome of our LSD toolkit, we
prioritized generation of day 50 iN and iDA cells in biological triplicate for 23 individual mutant lines
along with Control cells (**Fig. 1A and Dataset S2**), which is a sufficient period for creation of
functional synapses (22, 23). In parallel, a focused set of cell lines lacking genes directly (*ASAH1*,
*GBA1*, *SMPD1*) or indirectly (*GRN*) (7) linked with sphingolipidoses were differentiated for 30 days,
in biological triplicates (**Fig. 1A and Dataset S3**). Total cell proteins and interspersed quality control
samples were subjected to analysis using an Astral mass spectrometer with nDIA-hrMS2 (narrow-
window data-independent acquisition-high resolution MS$^2$) acquisition (**Fig. 1A**), allowing detection
of ~10,000 proteins across cells of each genotype (**Fig. S1C,D**).

We validated the robustness of the dataset as a resource by performing multiple types of quality
control analyses, allowing demonstration of: 1) consistent instrument performance over long data
collection periods (standard deviation <10%) (**Fig. S1E and Dataset S2,3**), 2) Pearson coefficients
of pairwise replicates typically >0.98 (**Fig. S1F**), 3) separation of neuronal subtype and length of
differentiation in principal component analysis (**Fig. S1G**), and 4) the expected loss of specific LSD
proteins, when detected, based on proteomics (**Fig. S1H**). Additionally, we demonstrated the
expected induction of several neuronal markers by proteomics or imaging in Control iN and iDA
cells (**Fig. S1I-K**) as well as consistent induction of a set of neuronal markers via proteomics in
both types of neurons, and TH protein in iDA cells, across all of the LSD mutant cell lines during
differentiation (**Fig. S1L,M**). Taken together, this data supports the utility of this resource for further
elucidation of proteomic phenotyping across LSD mutants.

**LSD Mutant iNeuron Organelle Landscapes**
We initially examined alterations in protein abundance, finding that the number of proteins
significantly changed compared to Control (q-value < 0.05, FDR-corrected) varied with genotype
and cell type (**Fig. S2A**). This was reflected distinct patterns of alterations in the abundance of
specific organelles or protein complexes/pathways (e.g. autophagy) (**Fig. 1B**). Alterations in the
abundance of proteins assigned to individual organelles provides a means by which to explore
possible organelle vulnerability across genotype and disease class, with, for example, a subset of
iDA and iN cells of sphingolipidoses and NCL cells displaying the largest alterations of changes in
protein abundance for lysosome and endosome compartments (**Fig. 1C**). The ability to identify
organelle vulnerability signatures within the proteomic data depends upon the quality of the
underlying annotations. For example, although various synaptic protein categories in the GO (gene
ontology) database are contained within SYNGO, there are varying degrees of overlap within the
various GO groups (**Fig. S2B**). We therefore used these more refined compartment annotations
along with an analogous treatment of the endolysosomal compartment (**Fig. S2B and Dataset S1**)
to perform hierarchical clustering of protein log$_2$FC alterations across organelle and sub-cellular
annotations (**Fig. S2C**). This analysis revealed shared patterns of increased or reduced protein
abundance especially for synaptic vesicle, lysosomal, and mitochondrial annotations, with iDA and

iN cell types sometimes demonstrating distinct patterns (**Fig. S2C**). F- ratio analysis of genotype
variance by annotation (between vs. within) demonstrated that the most impacted organelle
annotations reflected biological signals rather than technical noise (F-ratio>1.0) (**Fig. S2D**).
162
To capture the global impact of each LSD gene deletion, we generated an organelle correlation
map by computing all pairwise correlations of annotation-specific mean log$_2$fold-changes across
genotypes and cell types (**Fig. 1D,E**). This composite map integrates the direction and strength of
these relationships (as a correlation factor) to reveal how each knockout reshapes proteome
organization across subcellular compartments. For example, both $CLN5^{-/-}$ and $GRN^{-/-}$ iDA share
high similarity in synaptic annotation categories (SynGO, SynapaseSVs), whereas iN of $SMPD1^{-/-}$
and $ASAH1^{-/-}$ clustered together with iDA of $SMPD1^{-/-}$ and $CTSD^{-/-}$ (but interestingly not ASAH1$^{-/-}$
iDA) (**Fig. 1E**). The highest variability in correlations across disease classes was found for groups
of proteins linked with mitochondria, endolysosome, and synaptic vesicles/SynGO, for both iN and
iDA neurons (**Fig. 1F**). For example, the SynapseSV category of proteins displays high variance
across genotypes, particularly in iDA neurons, while in iN cells, the sphingolipidoses group of
proteins displayed the most extensive variability (**Fig. 1F**). Distinct patterns of variation between
organelles for the two cell types can be visualized in correlation plots for particular LSD mutations
(**Fig. 2A**). For example, synaptic proteins are among the most variable (most negative correlations)
in both iDA and iN $GBA1^{-/-}$ (Sphingolipidoses) cells, while mitochondria are selectively affected
(negatively correlated) in iDA cells (**Fig. 2A**). In contrast, synaptic proteins were positively
correlated in iDA $GRN^{-/-}$ (NCL) or $MCOLN1^{-/-}$ cells, but were negatively correlated in iN cells (**Fig.
2A**). As an orthogonal approach to comparing the effects of LSD mutations, we implemented a
ranked impact score that assesses the extent of variability in subcellular compartments in either
iDA or iN cells of individual disease classes, with the lowest score reflecting the greatest impact
(**Fig. 2B**) (see **MATERIALS and METHODS**). Taken together, this data suggests a complex
interplay between genotype and cell type in defining organelle landscapes across LSDs, and led
us to focus further on mitochondrial and synaptic compartments.
186
**Sphingolipidoses-related Mitochondrial Proteome Signatures**
Previous linkage of GBA1 mutants with mitochondrial dysfunction (19, 20) together with
mitochondria being among the most impacted organelles with Sphingolipidoses-related mutants
(**Fig. 2A**) led us to examine this relationship in detail. Correlation plots for iN and iDA cells for the
mitochondrial proteome revealed extensive variation for Sphingolipidoses mutants in iDA cells but
much less so in iN cells (**Fig. 2C**). Interestingly, while both $GBA1^{-/-}$ and $ASAH1^{-/-}$ iDA cells displayed
negative correlations, $HEXA^{-/-}$, $HEXB^{-/-}$ and $PSAP^{-/-}$ positive correlations (**Fig. 2C**), indicative of
distinct behavior, which was further visualized through unsupervised cluster analysis of individual
organelle annotations (**Fig. 2D**). We separated specific mitochondrial proteins based on functional
categories and abundance, including proteins involved in oxidative phosphorylation (OXPHOS) and
mitochondrial matrix (mtMatrix), and examined overlap between the various Sphingolipidoses
genotypes in iN and iDA cells (**Fig. 2E**). While $GBA1^{-/-}$ and $ASAH1^{-/-}$ mutants displayed either
complete (OXPHOS) or partial (mito- and mtMatrix) overlap in proteins with altered abundance in
iDA cells, no overlap was observed in the context of iN cells (**Fig. 2E**). Consistent with OXPHOS
components underlying the dependency on neuron subtype, negative correlation connectivity
(links) with other LSD gene deletions were prominent for $ASAH1^{-/-}$ iDA cells in the context of
OXPHOS proteins, but were much less prominent in the context of total mitochondrial proteins (**Fig.
2F**). Interestingly, the only positive link observable in the dataset for $ASAH1^{-/-}$ cells was with
$SMPD1^{-/-}$, and this was specific to iN cells (**Fig. 2F**). Together, these data provide insight into how
loss of specific Sphingolipidoses genes alters the abundance of selective classes of proteins within
the mitochondrial compartment in a neuronal type-dependent manner.
208
209  **Sphingolipidoses-related Synaptic Proteome Signatures**
210  Impact scores for synaptic and endocytic components among Sphingolipidoses mutants (**Fig. 2B**)
211  led us to focus on alterations in these compartments. Cluster analysis of log$_2$FC data for individual
212  organelles revealed distinct patterns across the LSD genotypes (**Fig. 3A,B**). For example, some

genotypes, including all Sphingolipidoses mutants, displayed increased levels of lysosomal proteins, which was frequently mirrored by reduced levels of synaptic proteins. In both cases, $ASAH1^{-/-}$ mutants displayed a small negative correlation for SynapseSV and SV fusion annotations in iDA cells but a positive correlation in iN cells, suggesting a distinct behavior (**Fig. 3C,D**). Most Sphingolipidoses cells displayed reductions in SV endocytosis and SV exocytosis protein abundance, but unlike other genotypes, $GBA1^{-/-}$ and $ASAH1^{-/-}$ cells also showed reductions in the abundance of SV fusion proteins selectively in iDA neurons (**Fig. 3B**). Proteins within the SV fusion category displayed numerous negatively correlated links with specific LSD mutants in either iDA or iN cells, and as expected, the number of such links within the SynapseSV category was substantially reduced relative to SV fusion (**Fig. 3E**). These data suggest potential synaptic vulnerability in $ASAH^{-/-}$ iNeurons, which is address below.

**An Approach for Decoding Protein Complex-level Vulnerabilities**

The experiments described above provide an atlas for evaluating the effects of LSD gene loss at an organelle level based on protein abundance. In order to extend this analysis to the level of individual PPIs and protein complexes, we developed an analysis pipeline that encompasses three major types of interaction data in a master PPI database: 1) well-validated protein complexes as are available in CORUM, Complex Portal (CP) or protein structure databases (PDB) (27-29), 2) high throughput interaction data, as represented in BioGrid and STRING databases (30, 31), and 3) predicted pairwise PPIs based on large-scale Alphafold Multimer or coevolutionary analyses (32, 33) (see **Fig. 4A,** *see* **Materials and Methods, Dataset S4**). Combining these resources with supporting structural evidence allowed us to generate a baseline" neuronal cell line PPI network based on Tier 1 confidence scores (**Fig. 4A**), ensuring that only high-quality predictions would be used for the downstream biological analysis. Over half (55.9%) of all edges in the neuronal baseline dataset were matched to structures in PDB, while nearly a quarter were also found in either CORUM or CP (**Fig. S3A**). Of the ~7100 PPIs found in the baseline database only 245 (~3%) had no additional structural or biological validation beyond the very high protein structure prediction metrics. To assess the combinatorial overlap of supporting evidence in a numerical fashion, we developed a cumulative score capturing the structural and biological evidence for each PPI (**Fig. S3B,C**). Not surprisingly, the highest counts per annotation were measured in mitochondrial, synaptic and nuclear annotations, thus scaling with both the number of proteins within a given annotation and the experimental evidence available in databases (**Fig. S3D,E,F**). To normalize for the varying size of annotations we expressed the coverage of PPI edges in support databases as a fraction of the size of all edges found in the annotation. Compartments with extensive structural and biological characterization (e.g. mitochondria) displayed complete coverage across all sets, whereas structurally less-well defined compartments (e.g. ER) scored more poorly (**Fig. S3D,** *bottom*). To expand this network further, we used the approach of Zhang et al (33) to first build 2571 and 1695 additional trimers and tetramers, respectively, from pairwise PPIs, with AF-Mprob >0.995, and >74% of these exhibiting structural support in the PDB (**Fig. 4A-C, Fig. S3G,H,I**). Fusion of high-confidence PPIs to at least 2 nodes of pre-existing trimers and tetramers seeds resulted in 408 complexes with a median support score of 4.06 across all databases (see **Materials and Methods** for details of confidence limits). While 246 (60.3%) of these matched one or more high confidence complexes, 138 represent candidate new assemblies (**Fig. 4D-F, Fig. S3J**). In the identified complexes, between 45-60% of edges were matched to pre-existing structural database evidence, suggesting robust empirical support for these structures (**Fig.4E,F**).

Having developed this protein interaction database, we first identified individual pairwise PPIs where the abundance of one or both proteins was significantly reduced (q < 0.05) compared with controls across LSD genotypes (**Fig. S4A**). The majority of pairwise interactions were unaffected by LSD gene mutation, although genotypes displaying the largest number of proteome alterations (e.g. CLN3, CLN5, CLN6 and MCOLN1) displayed the highest fractional PPI reduction, as expected. Reduced pairwise interactions were enriched in annotations for specific subcellular compartments in a genotype-dependent and in some cases, disease class-dependent manner in both iN and iDA cells, suggesting selective vulnerability (**Fig. 4G, Fig. S4B**). To map proteome

alterations at the protein complex level (i.e. beyond pairwise interactions), we assigned PPI nodes
within complexes as "disrupted" when the abundance of at least one interaction (edge) was
significantly reduced and as "lost" when *all* detected interactions with that node displayed reduced
abundance (**Fig. 4*A*, Fig. S4*C* and Dataset S4**) The number of lost or disrupted nodes varied
across LSD genotypes and was generally more prevalent in iN cells (**Fig. S4*C* and Dataset S4**).
Additionally, individual LSD genotypes displayed enrichment of specific organelles for disrupted or
lost protein complexes, including trimers built using the method of Zhang et al (33) (**Fig. 4*G*, Fig.
S4*D* and Dataset S4**).
275
Examples of disrupted complexes are shown in **Fig. 4*H* and Fig. S4*E***. One robust case is Complex
I from the mitochondrial electron transport chain, which displayed 7 or 10 lost or disrupted nodes,
respectively, in $GBA1^{-/-}$ iN or iDA cells, consistent with enrichment for OXPHOS localization terms
being affected at the organelle level (**Fig. 2*D***). Similarly, the BLOC-1 (Biogenesis of lysosome-
related organelles) complex known to regulate lysosome position in cells (34) displayed loss or
disruption of 4 or 5 nodes in $MCOLN1^{-/-}$ iN cells, and similarly, the core Commander Complex
involved in sorting of proteins into recycling vesicles (35) displays 9 disrupted nodes in $NPC1^{-/-}$ iN
cells. Additional examples of altered complexes include: 1) glutamate receptor assemblies (36) in
$GBA1^{-/-}$ and $MCOLN1^{-/-}$ cells, 2) CAMK2 in $GBA1^{-/-}$, $ASAH1^{-/-}$ and $MCOLN1^{-/-}$ cells, 3) the GATOR
complex involved in amino acid sensing (37) in $ASAH1^{-/-}$ cells, 4) the electron transport chain
complex cytochrome C oxidase in $ASAH1^{-/-}$ cells, and 5) the RAB7 guanine nucleotide exchange
factor CCZ1-MON1-RMC1 on lysosomes (38-40) in $ASAH1^{-/-}$ cells (**Fig. S4*E***). The entire collection
of protein complexes with alterations in specific subunits is available in **Dataset S4**. Taken together,
these examples demonstrate how the interaction analysis pipeline allows interrogation of altered
proteomes beyond the level of organelles in a systematic manner.
291
**Synaptic Function in $ASAH1^{-/-}$ iDA cells**
As described above, we detected alterations in synaptic protein abundance in $ASAH1^{-/-}$ iNeurons,
which led us to examine synaptic structure and function in these cells. We initially performed thin-
section electron microscopy on day 50 Control and $ASAH1^{-/-}$ iDA and iN cells (**Fig. 5*A*, Fig. S5*A***).
In Control iDA cells, we observed pre-synaptic structures containing numerous synaptic vesicles
with uniform vesicle sizes (**Fig. 5*A***). In contrast, apparent synaptic structures in $ASAH1^{-/-}$ iDA cells
appeared more disorganized with often fewer synaptic vesicles and with less uniform size
distributions (**Fig. 5*A***). Similar findings were made in day 50 iN cells (**Fig. S5*A***). Immunostaining
of day 38 iDA cultures with the pre-synaptic markers SYP (synaptophysin) and BSN (Bassoon)
revealed largely uniform pre-synaptic with the majority of SYP puncta co-staining with Bassoon in
Control cells (**Fig. 5*B-D*, Fig. S5*B***). In contrast, $ASAH1^{-/-}$ iDA cells frequently displayed enlarged
$\alpha$-tubulin positive neuronal structures that contained $\alpha$-SYP puncta that was more frequently not
co-stained with $\alpha$-Bassoon (**Fig. 5*B-D*, Fig. S5*A***), suggesting defects in pre-synaptic compartment
organization.
306
To examine the functional impact of $ASAH1$ deficiency, we performed Fluo-4-based $Ca^{2+}$ imaging
of Control and $ASAH1^{-/-}$ iDA and iN cells over increasing time periods of differentiation using a
robust evaluation machine-learning-based pipeline (41, 42) (**Fig. S5*C-E***) (see Materials and
Methods). As expected, stimulation with KCL resulted in an increase in relative firing frequency,
which could be reduced by pre-treatment with CQNX/D-AP5 (**Fig. S5*D***). We found that neuronal
firing, as measured by either the percent of total active regions of interest (ROI) over time or the
mean percentage of active versus non-active frames, were significantly reduced in day 50 iDA
$ASAH1^{-/-}$ vs Control cells, but this reduced activity could be partially rescued, albeit not to Control-
levels, by the addition of KCl (**Fig. 5*E*, Fig. S5*E***). In contrast, firing rates were only minorly affected
in iN cells over the selected timepoints (**Fig. S5*F***). Thus, alterations in the synaptic proteome in
day 50 iDA cells correlates with reduced firing rates.
318
**Visualization of $ASAH1^{-/-}$ endolysosomal compartments by cryo-ET**

ASAH1 plays a central role in glycosphingolipid catabolism within the lysosome through the
conversion of ceramide to sphingosine and fatty acids and functions downstream of GBA1 and
upstream of SMPD1 (**Fig. S6A**). The finding that *ASAH1*[-/-] iDA and iN cells display altered
lysosomal proteome abundance (**Fig. 3A,B**) prompted a biochemical and ultrastructural analysis
of ASAH1-deficient lysosomes in the experimentally tractable HeLa cells system.  Initially, we
performed LysoIP from Control and *ASAH1*[-/-] HeLa cells containing an endogenously tagged
TMEM192[HA] locus (4) along with parental HeLa cells as a negative control. Proteomic analysis of
isolated lysosomes displayed the expected increase in endolysosomal proteins, which were absent
in cells lacking the LysoIP tag (**Fig. S6B,C and Dataset S5**), with a subset of lumenal hydrolytic
proteins displaying increased abundance (**Fig. S6D**). As expected, lipidomics revealed a ~3-fold
increase in ceramide abundance in lysosomes from *ASAH1*[-/-] cells, consistent with *ASAH1*
deficiency (**Fig. S6E and Dataset S6**). Interestingly, several BMP species were increased in
abundance in lysosomes in the absence of *ASAH1*, which may underlie the modest (~2-fold or
less) reduction in glycosphingolipids (GSL) GM1, GM2, GM3 in lysosomes (7) (**Fig. S6F**). Taken
together, this analysis validates the *ASAH1*[-/-] cell line and is consistent with its role as an acid
ceramidase.
336
We next examined endolysosomal structures *in situ* in *ASAH1*[-/-] HeLa cells using cryo-focused ion
beam (cryo-FIB) milling paired with cryo-ET (**Fig. 5F-H**). Endolysosomal structures in Control cells
have typical ultrastructural features of auto-/endolysosomes, including frequent single or double
membrane structures within the limiting membrane and dense material indicative of degradative
cargo (**Fig. 5F,G**). Endolysosomal structures in *ASAH1*[-/-] cells appeared swollen compared to those
in Control cells, which showed dense packing of cellular material (**Fig. 5F,G**). Notably, membranous
material was largely absent in endolysosomal structures in *ASAH*[-/-] cells. In one tomogram of
*ASAH1*[-/-] cells, we observed an inward-budding vesicle within an endolysosome, along with
numerous intraluminal vesicles (ILVs) that appeared to be studded with glycosylated proteins (**Fig
5F-H** and additional examples in **Fig. S7A,B**). ILVs are thought to be the sites of breakdown of
glycosphingolipids (5, 8). Segmentation of tomograms highlights the difference in size and
membrane structure packing within the endolysosomes of Control or *ASAH1*[-/-] cells, and reveals
the close proximity of ER to the limiting membrane (**Fig. 5G, Movie S1, S2**). Taken together, these
results suggest endolysosomal dysfunction in the context of *ASAH1* deficiency.
351
352  **Discussion**
353
The LSD toolkit reported here provides a means by which to examine functional consequences of
individual LSD gene loss during changes in cell states while also allowing comparisons across
various classes of storage disorders. Through extensive analysis of total proteome data for both iN
and iDA cells, we provided a landscape of alterations reflecting the activities of individual LSD
genes. To our knowledge, this is the first comparison between iN and iDA cells in culture at this
level of proteome depth, and the first comparison across divergent LSD mutants. Additionally, we
developed an informatic framework to globally map alterations in specific protein complexes based
on alterations in protein abundance. This PPI approach allows us to go beyond protein abundance
changes at the organelle level to map alterations in the context of protein complexes, including
potentially new interactions based on proteome-wide Alphafold prediction. Although deployed here
on neuronal proteomes across LSD mutations, the approach should be adaptable for mapping
protein complex alterations across virtually any large-scale data types, including CRISPR, Perturb-
seq or degrader screens. Additionally, our analysis focused on Tier 1 PPIs, which represents the
most stringent evidence for validated interactions. Further analysis, including Tier 2 complexes
(**Fig. 4A**), may reveal additional complexes of interest whose abundance may be altered in specific
genetic contexts. Our ongoing studies seek to expand the toolkit to ultimately encompass all genes
associated with LSDs.
371
372  Our analysis focused on genes linked with sphingolipidoses, in part due to potential links with PD
373  risk(18). We found that *GBA1* and *ASAH1* mutants sometimes displayed distinct proteome

alterations compared with other sphingolipidoses genes. In particular, we observed alternations in
OXPHOS and pre-synaptic proteins in *GBA1* and *ASAH1* mutant iDA cells. The selectivity observed
for alterations in the mitochondrial OXPHOS system in iDA cells is interesting in light of the known
linkage between OXPHOS pathways and PD(43, 44). Conditional deletion of murine mtComplex I
subunit *Ndufs4* results in a deficit in nigrostriatal axons, and is associated with a PD-like loss of
motor function, suggesting that loss of mtComplex I component of OXPHOS is sufficient to produce
motor dysfunction(44). Likewise, the mtComplex I inhibitors MPTP and rotenone promote a
Parkinson's-like phenotype in mice (45, 46). Similarly, synaptic dysfunction has also been linked
with PD, including synaptotagmin-11 (*SYT11*) and synaptojanin (*SYNJ1*) variants that have been
associated with familial PD (47, 48). It is conceivable that alterations in OXPHOS and pre-synaptic
protein pools observed in *ASAH1*[-/-] and *GBA1*[-/-] iDA cells could represent vulnerabilities that
increase the risk of PD in patients with reduced enzymatic activity for these key sphingolipid
catabolism enzymes. Loss of *ASAH1* results in increased levels of ceramide, as confirmed here by
lipidomics in HeLa cells. Ceramides are known to affect the electron transport chain in
mitochondria(49), suggesting additional potential mechanisms independent of alterations in
OXPHOS protein abundance. Interestingly, we also observed increased levels of BMP species in
*ASAH1*[-/-] HeLa cells. We speculate that increased levels of BMP may increase the efficiency of
catabolism of sphingolipids in the context of *ASAH1* mutants, as indicated by reduced levels of
GM2 and GM3 species. The toolkit developed here, which includes a gene edited epitope for
LysoIP, will allow a detailed analysis of lysosomal lipidomes and proteomes in iN and iDA cells
lacking a broad set of LSD genes in future studies.

**Materials and Methods**
394
Detailed procedures related to cell culture, gene editing, biochemical procedures, MS, microscopy,
and electron tomography are described in *Materials and Methods* in the Supplemental Information
document. Statistical analyses were performed using MSstats and R. All error bars represent SEM,
and statistical significance was determined by *t* tests, as specified in the corresponding figure
legends.

**Data, Materials, and Software Availability**

Proteomic data (.RAW files) for nDIA of iN/iDA day 30, 50 and 70 iNeurons is deposited in the
MASSIVE Database (https://massive.ucsd.edu/ProteoSAFe/static/massive.jsp): MSV000099237
(reviewer login: MSV000099237_reviewer; password: lsd). HeLa LysoIP TMT data was deposited
into ProteomeXchange (50): PXD067219. Lipidomics data is deposited on Metabolomics
Workbench (51): Study ID ST004217; http://dx.doi.org/10.21228/M8Z556. Raw Cryo-ET
tomograms have been deposited at EMDB under accession numbers EMD-55210 (Control) and
EMD-55211 (*ASAH1*[-/-]). A Key Resource Table containing reagents and materials, source data,
segmentation models and datasets associated with this publication are available from Zenodo.org:
10.5281/zenodo.17296003. Scripts affiliated with this work is available on Github under the
repository https://github.com/sauerkrausi/neuroLSD.

**Acknowledgments**
412
We thank members of the Harper lab for feedback. We thank the Central Electron Microscopy
Facility at the Max Planck Institute of Biophysics for support. This work was funded by the Warren
Alpert Foundation (J.W.H.), the Bluefield Project (J.W.H., R.V.F, T.C.W.), NIH R01NS110395
(J.W.H.), NIH R01 GM132129 (J.A.P.), the Max Planck Society (F.W.), a Fred and Joan Goldberg
Post-doctoral Fellowship (F.K.). D. L. was supported by a Boehringer Ingelheim Fonds PhD
fellowship. This research was also funded in part by Aligning Science Across Parkinson's [ASAP-
000282 and 024268 to J.W.H. and F.W.] through the Michael J. Fox Foundation for Parkinson's
Research (MJFF). For the purpose of open access, the author has applied a CC BY public copyright
license to all Author Accepted Manuscripts arising from this submission. We acknowledge the Core

424
**References**

1.    R. E. Lawrence, R. Zoncu, The lysosome as a cellular centre for signalling, metabolism
and quality control. *Nat Cell Biol* **21**, 133-142 (2019).
2.    A. P. Mutvei, M. J. Nagiec, J. Blenis, Balancing lysosome abundance in health and
disease. *Nat Cell Biol* **25**, 1254-1264 (2023).
3.    J. A. Martina, N. Raben, R. Puertollano, SnapShot: Lysosomal Storage Diseases. *Cell*
**180**, 602-602 e601 (2020).
4.    F. Kraus *et al.*, Global cellular proteo-lipidomic profiling of diverse lysosomal storage
disease mutants using nMOST. *Sci Adv* **11**, eadu5787 (2025).
5.    G. Wilkening, T. Linke, K. Sandhoff, Lysosomal degradation on vesicular membrane
surfaces. Enhanced glucosylceramide degradation by lysosomal anionic lipids and
activators. *J Biol Chem* **273**, 30271-30278 (1998).
6.    H. D. Gallala, K. Sandhoff, Biological function of the cellular lipid BMP-BMP as a key
activator for cholesterol sorting and membrane digestion. *Neurochem Res* **36**, 1594-1600
(2011).
7.    S. Boland *et al.*, Deficiency of the frontotemporal dementia gene GRN results in
gangliosidosis. *Nat Commun* **13**, 5924 (2022).
8.    H. Schulze, K. Sandhoff, Sphingolipids and lysosomal pathologies. *Biochim Biophys Acta*
**1841**, 799-810 (2014).
9.    C. Blauwendraat *et al.*, Genetic modifiers of risk and age at onset in GBA associated
Parkinson's disease and Lewy body dementia. *Brain* **143**, 234-248 (2020).
10.   E. Sidransky *et al.*, Multicenter analysis of glucocerebrosidase mutations in Parkinson's
disease. *N Engl J Med* **361**, 1651-1661 (2009).
11.   O. Goker-Alpan *et al.*, Parkinsonism among Gaucher disease carriers. *J Med Genet* **41**,
937-940 (2004).
12.   J. Aharon-Peretz, H. Rosenbaum, R. Gershoni-Baruch, Mutations in the
glucocerebrosidase gene and Parkinson's disease in Ashkenazi Jews. *N Engl J Med* **351**,
1972-1977 (2004).
13.   D. Chatterjee, D. Krainc, Mechanisms of Glucocerebrosidase Dysfunction in Parkinson's
Disease. *J Mol Biol* **435**, 168023 (2023).
14.   R. N. Alcalay *et al.*, SMPD1 mutations, activity, and alpha-synuclein accumulation in
Parkinson's disease. *Mov Disord* **34**, 526-535 (2019).
15.   Z. Gan-Or *et al.*, The emerging role of SMPD1 mutations in Parkinson's disease:
Implications for future studies. *Parkinsonism Relat Disord* **21**, 1294-1295 (2015).
16.   Z. Gan-Or *et al.*, The p.L302P mutation in the lysosomal enzyme gene SMPD1 is a risk
factor for Parkinson disease. *Neurology* **80**, 1606-1610 (2013).
17.   J. N. Foo *et al.*, Rare lysosomal enzyme gene SMPD1 variant (p.R591C) associates with
Parkinson's disease. *Neurobiol Aging* **34**, 2890 e2813-2895 (2013).
18.   L. A. Robak *et al.*, Excessive burden of lysosomal storage disorder gene variants in
Parkinson's disease. *Brain* **140**, 3191-3203 (2017).
19.   S. Kim, Y. C. Wong, F. Gao, D. Krainc, Dysregulation of mitochondria-lysosome contacts
by GBA1 dysfunction in dopaminergic neuronal models of Parkinson's disease. *Nat*
*Commun* **12**, 1807 (2021).
20.   C. Moren *et al.*, GBA mutation promotes early mitochondrial dysfunction in 3D
neurosphere models. *Aging (Albany NY)* **11**, 10338-10355 (2019).
21.   F. V. Hundley *et al.*, Endo-IP and lyso-IP toolkit for endolysosomal profiling of human-
induced neurons. *Proc Natl Acad Sci U S A* **121**, e2419079121 (2024).
22.   Y. Zhang *et al.*, Rapid single-step induction of functional neurons from human pluripotent
stem cells. *Neuron* **78**, 785-798 (2013).

23.  J. E. Heebner *et al.*, Deep Learning-Based Segmentation of Cryo-Electron Tomograms. *J
Vis Exp* 10.3791/64435 (2022).
24.  A. Ordureau *et al.*, Temporal proteomics during neurogenesis reveals large-scale
proteome and organelle remodeling via selective autophagy. *Mol Cell* **81**, 5082-5098
e5011 (2021).
25.  K. L. Hickey *et al.*, Proteome census upon nutrient stress reveals Golgiphagy membrane
receptors. *Nature* **623**, 167-174 (2023).
26.  M. J. Hoyer *et al.*, Combinatorial selective ER-phagy remodels the ER during
neurogenesis. *Nat Cell Biol* **26**, 378-392 (2024).
27.  G. Tsitsiridis *et al.*, CORUM: the comprehensive resource of mammalian protein
complexes-2022. *Nucleic Acids Res* **51**, D539-D545 (2023).
28.  H. M. Berman *et al.*, The Protein Data Bank. *Nucleic Acids Res* **28**, 235-242 (2000).
29.  S. Balu *et al.*, Complex portal 2025: predicted human complexes and enhanced
visualisation tools for the comparison of orthologous and paralogous complexes. *Nucleic
Acids Res* **53**, D644-D650 (2025).
30.  R. Oughtred *et al.*, The BioGRID database: A comprehensive biomedical resource of
curated protein, genetic, and chemical interactions. *Protein Sci* **30**, 187-200 (2021).
31.  D. Szklarczyk *et al.*, The STRING database in 2023: protein-protein association networks
and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids
Res* **51**, D638-D646 (2023).
32.  E. W. Schmid *et al.*, Proteome-wide in silico screening for human protein-protein
interactions. *BioRxiv* doi.org/10.1101/2025.11.10.687652 (2025).
33.  J. Zhang *et al.*, Predicting protein-protein interactions in the human proteome. *Science*
**390**, eadt1630 (2025).
34.  R. De Pace, S. Ghosh, C. D. Williamson, J. S. Bonifacino, BLOC-1 and BORC: Complex
regulators of endolysosomal dynamics. *Cell Chem Biol* **32**, 1106-1124 (2025).
35.  K. E. McNally, P. J. Cullen, Endosomal Retrieval of Cargo: Retromer Is Not Alone.
*Trends Cell Biol* **28**, 807-822 (2018).
36.  I. H. Greger, M. L. Mayer, Structural biology of glutamate receptor ion channels: towards
an understanding of mechanism. *Curr Opin Struct Biol* **57**, 185-195 (2019).
37.  Z. Cui, A. M. N. Joiner, R. M. Jansen, J. H. Hurley, Amino acid sensing and lysosomal
signaling complexes. *Curr Opin Struct Biol* **79**, 102544 (2023).
38.  L. Dehnen *et al.*, A trimeric metazoan Rab7 GEF complex is crucial for endocytosis and
scavenger function. *J Cell Sci* **133** (2020).
39.  L. P. Vaites, J. A. Paulo, E. L. Huttlin, J. W. Harper, Systematic Analysis of Human Cells
Lacking ATG8 Proteins Uncovers Roles for GABARAPs and the CCZ1/MON1 Regulator
C18orf8/RMC1 in Macroautophagic and Selective Autophagic Flux. *Mol Cell Biol* **38**
(2018).
40.  M. Nordmann *et al.*, The Mon1-Ccz1 complex is the GEF of the late endosomal Rab7
homolog Ypt7. *Curr Biol* **20**, 1654-1659 (2010).
41.  M. Pachitariu, M. Rariden, C. Stringer, Cellpose-SAM: superhuman generalization for
cellular segmentation. *BioRxiv* https://doi.org/10.1101/2025.04.28.651001 (2025).
42.  S. Lenci, D. Sieger, CalciumNetExploreR: An R Package for Network Analysis of Calcium
Imaging Data. *BioRxiv* https://doi.org/10.1101/2025.01.24.634485 (2025).
43.  W. S. Choi *et al.*, Conditional deletion of Ndufs4 in dopaminergic neurons promotes
Parkinson's disease-like non-motor symptoms without loss of dopamine neurons. *Sci Rep*
**7**, 44989 (2017).
44.  P. Gonzalez-Rodriguez *et al.*, Disruption of mitochondrial complex I induces progressive
parkinsonism. *Nature* **599**, 650-656 (2021).
45.  A. D. Van Laar *et al.*, Transient exposure to rotenone causes degeneration and
progressive parkinsonian motor deficits, neuroinflammation, and synucleinopathy. *NPJ
Parkinsons Dis* **9**, 121 (2023).
46.  J. W. Tetrud, J. W. Langston, MPTP-induced parkinsonism as a model for Parkinson's
disease. *Acta Neurol Scand Suppl* **126**, 35-40 (1989).

47.    M. A. Nalls *et al.*, Large-scale meta-analysis of genome-wide association data identifies
529           six new risk loci for Parkinson's disease. *Nat Genet* **46**, 989-993 (2014).
530    48.    M. Cao *et al.*, Parkinson Sac Domain Mutation in Synaptojanin 1 Impairs Clathrin
531           Uncoating at Synapses and Triggers Dystrophic Changes in Dopaminergic Axons.
532           *Neuron* **93**, 882-896 e885 (2017).
533    49.    M. Vos, C. Klein, A. A. Hicks, Role of Ceramides and Sphingolipids in Parkinson's
534           Disease. *J Mol Biol* **435**, 168000 (2023).
535    50.    Y. Perez-Riverol *et al.*, The PRIDE database resources in 2022: a hub for mass
536           spectrometry-based proteomics evidences. *Nucleic Acids Res* **50**, D543-D552 (2022).
537    51.    M. Sud *et al.*, Metabolomics Workbench: An international repository for metabolomics
538           data and metadata, metabolite standards, protocols, tutorials and training, and analysis
539           tools. *Nucleic Acids Res* **44**, D463-470 (2016).

**Figures and Tables**

**544** Fig. 1. Proteome atlas of lysosomal Storage Disorder neuronal cell models. (*A*) Schematic of
**545** experimental workflow. Control and 23 LSD knockout human ES cell lines were differentiated to iN
**546** and iDA in two cohorts (day 30, day 50, each in biological triplicate) and analyzed using nDIA LFQ.
**547** (*B*) Half-circos plot of log$_2$FC mean protein abundance per annotation (q < 0.05, FDR-corrected) of
**548** each LSD knockout/Control for iN (left) and iDA (right) neurons for the indicated annotations or
**549** organelles. (*C*) Ternary plot of log$_2$FC [KO/Control] for endosome, lysosome and OXPHOS
**550** annotations across iN and iDA neurons. Triangles, iDA cells; circles, iN cells. (*D*) Example of
**551** pairwise, proteome-wide cross-correlation of lysosome (top), SynapseSV (middle) and recycling
**552** endosome (bottom) annotated proteins. (*E*) Composite correlation across 20 organelle
**553** annotations. (*F*) Mean organelle correlation based on by disease classes. ± SD for iN (top) and iDA
**554** (bottom). Disease class is indicated by shading.

**Fig. 2. Differential mitochondrial organelle vulnerability across Sphingolipidoses mutant iDA neurons. (A)** iN vs. iDA organelle and protein complex/compartment correlation plots for cells deficient in *GBA1*, *ASAH1*, *GRN*, or *MCOLN1* (TRPML1). **(B)** Impact score reflecting the extent to which genotypes within Sphingolipidoses, Integral Membrane Protein Disorders, and Neuronal Ceroid Lipofuscinoses categories display alterations across various organelles and protein complex/compartment proteomes in either iN or iDA cells. The numbers in each square represent the rank in each cell lineage. The lower the number indicates a more severe disruption across the genotypes examined. **(C)** iN vs. iDA correlation plot for mitochondria across 23 LSD mutants. **(D)** Hierarchical cluster correlation heatmap of indicated organelle annotations across Sphingolipidoses mutants for iN (left) and iDA (right) cells. **(E)** Euler plots depicting overlap and unions of mitochondrial (top), OXPHOS (middle) and mito-matrix (bottom) protein IDs log$_2$FC > -0.75 across sphingolipidoses mutants for iN and iDA cells. **(F)** Circos plots showing correlations between genotypes for mitochondria (left) and OXPHOS (right). Center panel depicts genotypic links originating from only *GBA1$^{-/-}$* and *ASAH1$^{-/-}$*. Only links with Pearson scores (r) smaller than -0.2 and larger than 0.7 are shown. Blue labels indicate genotypes belonging to Sphingolipidoses disease class. Bold labels are PD-risk genes.

584
**Fig 3. Systematic analysis of synaptic and endolysosomal compartments across LSD iN and**
**iDA cells. (A,B)** Log$_2$FC [KO vs Control] heatmap of vesicular annotations for iN (panel **A**) and iDA
(panel **B**) cells at day 50 of differentiation. The annotations are sorted from high to low from left to
right based on iN cell data. Sphingolipidoses related genotypes are in blue and LSD-PD risk genes
are labeled in bold. **(C,D)** iN vs. iDA organelle correlation of SynapseSV (panel **C**) and SV fusion
(panel **D**) proteins. Genotypes are color-coded by disease class. **(E)** Circos plots showing
correlations between genotypes for SynapseSV (left) and SV fusion (right) proteins. Center plots
depict correlation links originating from only *GBA1*$^{-/-}$ and *ASAH1*$^{-/-}$. Only links with Pearson (r)
smaller than -0.2 and larger than 0.7 are shown. Blue labels indicate genotypes belonging to
Sphingolipidoses-risk genes. Bold labels are PD-risk genes.

**Fig. 4. Protein-Protein interaction atlas of neuronal lysosomal dysfunction. (A)** Schematic depiction of data evaluation pipeline for neuronal protein-protein interactions (PPIs) in LSD mutants. **(B)** Bar graph of trimers found in neuronal baseline PPIs matched to pre-existing complex and structural databases (left). Heatmap representing the fraction of trimers (in %) found in databases per annotations or organelle (right). **(C)** Bar graph showing number of PPIs, trimers and tetramers (from left to right) with matching structural validation in PDB database. **(D)** Like panel **B**, but for complexes detected in neuronal baseline PPI dataset. **(E)** Bar graph of mean percent of PPI edges found in ComplexPortal (CP), CORUM and PDB. **(F)** Boxplot of mean support score of complexes belonging to novel, relaxed match and strict match classes. **(G)** Heatmap of fraction of PPIs (q < 0.05) disrupted by organelle across LSD KOs in iN (left) and iDA (right). **(H)** Examples representation of protein complex vulnerability. (top) Mitochondrial complex I is depicted as example with loss or disruption of several nodes in *GBA1*-/- iN (middle) or *GBA1*-/- iDA (right) compared with Control cells (left). (bottom left) BLOC-1 complex as an example of lost or disrupted interactions in *MCOLN1*-/- iN cells. (bottom right) Commander complex as an example of a complex disrupted in *NPC1*-/- iN cells. The legend summarizes color codes for edge evidence and PPI state.

**Fig. 5**

**Fig. 5: Synaptic morphology defects correlate with reduced basal Ca$^{2+}$ firing activity in ASAH1$^{-/-}$ iDA cells. (A)** Representative EM images depicting synapses and neuronal ultrastructure of Control and *ASAH1$^{-/-}$* iDA neurons at day 50 of *in vitro* differentiation. Scale bars are indicated. Pre-synaptic structures with synaptic vesicles are indicated by yellow arrowheads. Panels on the right display enlarged vesicular structures within *ASAH1$^{-/-}$* synapses. **(B)**, Confocal images of Control and *ASAH1$^{-/-}$* iN and iDA cells at day 38 of *in vitro* differentiation, labeled for SYP, Bassoon and Tubulin. Scale bar: 20 μm and 2 μm. Images depicting a wider field view are provided in **Fig. S5B**. **(C)** Number of colocalized SYP-Bassoon object pairs in Control iN and iDA and iDA *ASAH1$^{-/-}$* neurons, based on confocal images in panel *B* (n(Control iN): 20 stacks; n(Control iDA): 19 stacks; n(*ASAH1$^{-/-}$* iDA): 19 stacks). iN: **P(0.001); iDA: ****P< 0.0001 ; iDA Ctrl vs *ASAH1$^{-/-}$:* ****P< 0.0001. **(D)** Quantification of SYP object counts (from left to right), based on confocal images in panel *B*. (n(Control iN): 40 stacks; n(Control iDA): 40 stacks; n(*ASAH1$^{-/-}$* iN): 20 stacks; n(*ASAH1$^{-/-}$* iDA): 40 stacks). Object count: Control iN vs iDA: ***P(0.0002), iN Control vs *ASAH1$^{-/-}$*: **P(0.003), iDA Control vs *ASAH1$^{-/-}$*: ****P<0.0001. **(E)** Stacked bar graph for Control (left) and *ASAH1$^{-/-}$* (right) iDA cell Ca$^{2+}$ firing depicting the cumulative % of active ROIs over the time course. d33 Control vs *ASAH1$^{-/-}$*: *P(0.02); d50 Control vs *ASAH1$^{-/-}$*: *P(0.04). Data from day 33 and 50 of *in vitro* differentiation with or without stimulation by KCl. **(F)** Example tomogram of Control and *ASAH1$^{-/-}$*

635    endo-lysosomal structures. Scale bar 100 nm. **(G)** 3D-renderings of tomogram segmentations from
636    panel **F**. **(H)** Higher magnification of the white boxes in panel **F** showing inward budding membrane
637    events (purple arrowheads) and ILVs studded with glycosylated proteins (yellow arrowheads).
638    Scale bar 100 nm.
639