A *Plasmodium falciparum* redox survival mechanism licenses killing by artemisinins

Smritikana Dutta,[1], Mohammad Faaiz[1], Saikat Bhattacharjee[2], Kasturi Haldar[3,*] and Souvik

Bhattacharjee[1, *]

[1] Special Centre for Molecular Medicine, Jawaharlal Nehru University, New Delhi 110067. India.

[2] Regional Center for Biotechnology, 3rd Milestone, Faridabad-Gurgaon Expressway, Faridabad,

Haryana 121001. India.

[3] Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, IN 46556, USA.

[*] Corresponding authors

Corresponding authors address:

souvik@jnu.ac.in . Special Centre for Molecular Medicine, Jawaharlal Nehru University, New Delhi
110067. India.

khaldar@nd.edu. Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana,
IN 46556, USA.

Abstract

Mutations in *Plasmodium falciparum* Kelch13 (K13) confer artemisinin resistance (ART-R) which threatens global malaria control, but known K13 functions fail to explain clinical ART-R. We reported that K13 binds the oxidant heme *in vitro*, however, its functions in redox-stress, cell survival and death remained unknown. Since taut control of free heme is not feasible in infected erythrocytes, we utilized a non-erythroid cell model to show that K13 directly binds and is stabilized by nanomolar heme levels. K13 also binds and regulates a major redox transcription factor, which is displaced by heme into the nucleus, to raise redox-stress responses that become suppressed during artemisinin-induced death (ART-death). K13's evolutionarily conserved kelch domain confers heme-binding and ART-death characteristics to its mammalian orthologue KEAP1. Chemical or genetic elevation of K13, fuels ART-death proportionate to K13 levels even in vast excess of heme, suggesting a novel plasmodial redox-survival mechanism licenses ART-death in clinical ART-R.

Introduction

Artemisinins (ARTs) are highly effective in curing blood-stage infection of *Plasmodium falciparum,* the deadliest of human malaria parasites[1]. These drugs are critical components of artemisinin-based combination therapies (ACT) that have played a key role in reducing the global burden of malaria and remain the cornerstone of its control worldwide[2]. Despite the prevalent model that ART kills through indiscriminate alkylation and damage of parasite lipids, proteins and DNA[3–5], robust clinical resistance has been ascribed to one major gene *Pfkelch13* (K13; PlasmoDB ID: PF3D7_1343700)[6], which has been shown to be causal for resistance[7]. Since ART-resistance (ART-R) poses a significant threat to malaria control[8,9], K13 has been intensely studied over the last decade[10–14]. The consensus data suggest that it possess an evolutionarily conserved 'kelch domain' is cytoplasmically synthesized and associated with vesicular structures, including the parasite's digestive endosomes, apicoplast, mitochondria, endoplasmic reticulum as well as vesicles enriched in Rabs, ATG18 and phosphatidylinositol 3-phosphate (PI3P)[9–11]. At the parasite's trophozoite stage, when haemoglobin uptake and digestion are most active, K13 regulates endocytosis *via* structures called cytostomes, and resistance mutations suppress this process[15,16]. But notably, these very trophozoite stages harbouring K13 resistance mutations remain highly susceptible to ART suggesting that while hemoglobin uptake regulated by K13, it may not be the major target of resistance, which may arise from changes in additional but still unknown functions of K13.

ARTs are 'pro-drugs' in that they must all be converted to dihydroartemisinin (DHA) and which in turn is subsequently activated to release their killing action. High (micromolar) concentration of heme have been shown to increase ART-potency because it cleaves the endoperoxide bridge yielding highly reactive oxy-centered and C4-centered radicals[17] that alkylates a wide range of cellular components causing oxidative damage, which results in parasite cell death[18,19]. Malaria parasites digestion of host hemoglobin actively produces heme in the parasite's food vacuole (FV). Heme is presumed to confer the sensitivity of ARTs in killing malaria parasites[20,21]. However, it is poorly understood whether this can be achieved by physiological heme concentrations and how killing is effected in 'ring' parasites early in blood-stage infection that lack a food vacuole, yet are critical clinical targets of ART-R[22,23].

We have recently reported that recombinant and parasite-derived K13 directly bind molecular heme

with a dissociation constant (K_d) of 61.5 nM[24]. Heme binds to many proteins[25,26]. This includes

albumins with a K_d range of 2-40 μM[27], explaining their ready capacity to transfer heme to wide range

of acceptors[28]. In contrast heme-oxygenases which degrade heme, bind it with nanomolar (3-300

nM) affinity[29]. This suggests that with a K_d of 61.5 nM, K13 engages in at least moderately tight heme-

binding. Heme binding is known to effect profound structural changes in proteins[30] and heme is

implicated in oxidative stress metabolism[31]. However, the consequences of heme binding for K13's

stability and turnover, and whether heme-binding induces K13-mediated redox-controlled responses

remain entirely unknown. Furthermore, K13's engagement of antioxidant and detoxification systems,

as regulated by its closest mammalian orthologue Kelch-like ECH-associated protein 1 (KEAP1)[6,32,33]

has not been investigated. Finally, the implications of dislodging or neutralizing heme bound to K13,

for mechanisms of ART-based killing remain to be discovered.

Since *in vitro* studies suggest that K13 binds heme with a K_d of 61.5 nM[24], regulation of associated

cellular mechanisms may be expected to be initiated at nanomolar to low micromolar concentrations.

However, carefully controlling heme levels in this range remains unfeasible in blood-stage parasites.

This is because the host erythrocyte contains high levels of hemoglobin (3.2-3.6 mM)[34], whose

digestion is a major source of parasite heme. Erythrocytes also contain high levels (21 ± 2 μM) of free

heme[35] which may translocate through passive and facilitated pathways[26,36]. Since mature human red

cells are enucleated and terminally differentiated, these host sources of heme cannot be genetically

'turned off' post infection. Moreover, their reduction compromises parasite health[37]. Genetic or

chemical targeting of parasite's *de novo* heme biosynthesis in infected erythrocytes appears to have

no effect on parasite growth[38,39], suggesting that heme provided by exogenous pathways can be

supplemented from host-derived sources. Notably, free heme levels in most non-erythroid cells are in

the nanomolar range (25-300 nM)[36], but in *P. falciparum* they are reported to be considerably higher

(at 1.6 μM or more)[40,41], suggesting that throughout blood-stage infection the parasite may reside in a

high intracellular heme-rich environment, but the consequences for redox stress remain obscure.

We desired a cell system to study the molecular functions of K13 and their regulations by nanomolar

concentrations of heme. We therefore used a model non-erythroid mammalian cell line Du145[42] with

reduced levels of KEAP1, adapted to grow in heme-depleted media, and whose free heme levels

were tightly regulable by blocking endogenous heme biosynthetic enzymes. These Du145 cells were

otherwise capable of an active redox response[43]. Our goal was to query whether K13 and/or heme

contribute to redox-stress responses well as obtain mechanistic understanding of how either or both

trigger sensitization of cells to killing by ARTs and validate the model system as a rapid, functional

screen to identify redox cell survival and death mechanisms of *Plasmodium falciparum* malaria

Results

K13 cellular levels are increased by heme-binding and -stabilization

Human Du145 cells (HTB-81, ATCC, *www.atcc.org*) contain a full length wild-type *Keap1* gene, but

CpG dinucleotide methylation of its core promoter results in low *keap1* mRNA expression[44], making

them suitable for ectopically expressing kelch domain proteins with minimal interference from the

native KEAP1 protein. Sequence comparison clearly revealed that K13 is a homologue of KEAP1

(**Supplementary Fig. 1a**). We generated three stable transgenic lines expressing: (i) K13 as a C-

terminus fusion to GFP; K13-GFP, (ii) GFP alone, as well as (iii) a C-terminus fusion of flag-strep tag

KEAP1 (see *Materials and Methods*). Indirect immunofluorescence assays (IFA) with quantitative

analyses, suggested cellular distribution of K13-GFP, GFP and KEAP1, at comparable levels in

transgenic Du145 cells (**Supplementary Fig. 1b**). K13-GFP, GFP and flag-strep tag KEAP1

respectively appeared as fusion proteins at expected sizes of 105-kDa, 27-kDa and 80-kDa

(**Supplementary Fig. 1c, d**).

We recently showed that *in vitro* heme bound recombinant K13 protein (TrK13-WT) with high

affinity[24], but action of heme in cells was not known. Since the K_d for heme binding to K13 protein was

61.5 nM, we wanted to measure the effects at low nanomolar concentrations of heme. This required

growing Du145 cells under conditions where available heme was substantially depleted from the

extracellular sources, and the production of biosynthetic heme was blocked by inhibitors. Fetal bovine

serum (FBS) is the largest extracellular donor of heme and contributed median concentration of 3.2 ±

0.06 µM heme to culture media. We reduced FBS concentrations of heme to <850 nM

(**Supplementary Fig. 2a, b**) to create heme-depleted media (HDM; *Material and Methods*). Since the

K_d of heme for albumins is >1 µM, this was expected to dramatically reduce the transfer of

extracellular heme to cells. We also added succinylacetone [SA; an inhibitor of the second enzyme of

heme biosynthesis pathway, δ-aminolevulinic acid (ALA) dehydratase] to the culture media, which

restricted intracellular *de novo* heme synthesis. Our overall goal was to achieve culture conditions that

would be responsive to nanomolar concentrations of exogenously added heme. Accordingly, under

HDM conditions, the parental Du145, transgenic GFP and K13-GFP cells all showed intracellular

concentrations of ~60 nM heme. Addition of extracellular heme at 5 μM, raised intracellular heme to

400-450 nM. When extracellular concentrations were raised to 50 μM, the intracellular levels

plateaued at ~750 nM (**Supplementary Fig. 2c**). Together these data suggested that in the course of

all experiments intracellular heme concentrations ranged from ~60-750 nM.

K13-GFP cells cultured in HDM (K13-GFP^{HDM}) were subsequently exposed to different concentrations

of exogenously added heme (0-5 μM) in the media (referred to as heme-replenished media). As

shown in **Fig. 1a**, concentrations of 10 nM heme induced detectable increase in K13-GFP levels.

Progressive increase of K13-GFP levels were seen at 50 nM, 100 nM, 1 μM and 5 μM heme. At 5 μM

heme, K13-GFP levels were restored to those seen in K13-GFP 'control' cultures (**Fig. 1b**). There was

no effect on the cellular viability of K13-GFP (or GFP) expressing cells at 5 μM heme;

(**Supplementary Fig. 2d:** the IC$_{50}$ for heme was ~25-fold higher for all cells). Notably, there was no

detectable change in the relative intensity of GFP or KEAP1 proteins when cells were subjected to

heme depletion or when supplemented with 5 μM heme (**Supplementary Fig. 2e, f**).

We next investigated the specificity of heme action. As shown in **Fig. 1c,** PPIX alone or when

complexed with Zn, had no effect even at 50 μM. However, increasing heme levels from 5 to 50 μM

continued to steadily increase levels of K13-GFP. Together, the data suggest that heme at

extracellular concentrations of 10 nanomolar or higher (and up to 50 μM) increase cellular levels of

K13-GFP (but not GFP or KEAP1), suggesting it is specific for K13. Increases in K13 were only

mediated by heme (not PPIX or Zn-PPIX) and likely due to heme's directly binding to K13 (since we

have previously shown that heme binds K13 with nanomolar affinity *in vitro*[24]).

Amongst heme binding proteins, hemopexin, a serum glycoprotein (which displays highest affinity for

heme at K$_d$ ~ 1 pM[45,46]) has been reported to show heme-dependent resistance to protease

digestion[45]. We therefore investigated if the protease sensitivity of K13 was altered in its heme-bound

state. For *in vitro* assay, we used the affinity purified 61-kDa recombinant TrK13-WT protein as K13

representative, described in our previous studies[24,47]. Recombinant TrK13-WT protein (10 μM) was

incubated with or without equimolar concentrations of heme and desalted to remove excess unbound

heme (see *Materials and methods*). Heme-bound TrK13-WT protein was subsequently treated with

trypsin under optimal buffer and pH conditions for varied incubation times. For trypsin (a serine

protease that cleaves between the carboxyl group of arginine or lysine and the amino group of the

adjacent amino acid), we also included commercially procured BSA, a confirmed heme-binding

protein[48] (but with substantially lower affinity) as a control. As shown in **Fig. 1d**, Coomassie-stained

SDS-PAGE revealed full-length (FL) TrK13-WT protein at 61-kDa. Treatment with trypsin yielded

additional low molecular weight fragments (doublets at 30-kDa and at 12-kDa; blue arrowheads) in

the absence of heme, whose appearance was blocked when the TrK13-WT protein was pre-

incubated with heme. Heme-binding only conferred partial resistance of BSA to trypsin

(**Supplementary Fig. 2g**). In absence of heme, multiple small protein fragments were detected (from

22-40 kDa; green arrowheads), while in its presence, larger fragments were observed at 45-55-kDa

(red arrowheads). The effects of heme on trypsin digestion of both TrK13-WT protein and BSA were

likely because of changes in the 3-dimensional conformation of these proteins (which is consistent

with previous reports in the literature[48] and observed in our earlier CD studies[24]). As shown in **Fig. 1e**,

resistance to trypsin digestion was not seen when the TrK13-WT protein was incubated with PPIX or

Zn-PPIX or iron salts (ferrous ammonium sulfate, FAS or ferric ammonium sulfate, FES, although

these salts are known to bind K13)[24]. Together, these data suggest that heme binding to K13 shows

chemical and molecular specificity, may induce confirmational change and thereby induce

stabilization of K13 in cells.

K13-regulated oxidative stress responses are triggered by heme

Nuclear factor erythroid 2-related factor 2 (Nrf2) is a major redox-regulated transcription factor

controlled by KEAP1[49], the closest mammalian orthologue of K13. We therefore assessed whether

Nrf2 was regulated by K13 and the impact of heme, a known, major redox regulator[50]. First, we

investigated changes in Nrf2 levels in GFP or KEAP1 expressing cells under HDM or heme-

replenished culture conditions *i.e.,* GFP[HDM] *versus* GFP[HEME] or KEAP1[HDM] *versus* KEAP1[HEME] cells.

Relative intensities of both GFP and KEAP1 proteins remained unaffected by heme (**Supplementary**

Fig. 3a, b). Consequently, we investigated only K13-GFP cells, cultured in the absence or presence

of increasing concentrations of heme, and followed by immunoprecipitations from cellular lysates

using anti-GFP antibodies. Western blot (**Fig. 2a**) and quantitative densitometric analysis (**Fig. 2b**) of

band intensities revealed increasing amounts of immunoprecipitated K13-GFP with increased heme

concentrations (**2a i** and **2b i**). In contrast, levels of co-immunopurified Nrf2 decreased with increasing

heme (**Fig. 2a i**), although there was increase in the overall cellular Nrf2 levels (**Fig. 2a ii** and **2b ii-**

iii). These results strongly implied that interactions between K13 and Nrf2 were negatively affected by

heme. To investigate whether these reflected direct protein-protein interactions, we co-incubated

recombinant Nrf2-6×his protein with either heme-free or heme-bound recombinant TrK13-WT (that

was desalted to remove excess unbound heme). As shown in **Fig. 2c**, after incubation with heme for

1 h, there was ~26% reduction in Nrf2-6×his binding to TrK13-WT, suggesting heme may directly

prevent/disrupt Nrf2 binding to K13.

We next performed immunolocalization studies to determine subcellular distribution of Nrf2 in K13-

GFP[HDM] and K13-GFP[HEME] cells (**Fig. 2d**). Here, data revealed predominantly cytoplasmic localization

of Nrf2 with K13-GFP under HDM conditions (left panel). In contrast, the K13-GFP[HEME] cells displayed

prominent nuclear localized Nrf2 protein (right panel). Moreover, in the presence of heme, levels of

K13 were higher than under HDM conditions. Subcellular fractionation confirmed that under

conditions of HDM, a vast majority (~83%) of Nrf2 protein was detected in the cytoplasmic fraction

(**Fig. 2e**; second blot). However, in the K13-GFP[HEME] cells, ~87% of the Nrf2 protein was detected in

the nuclear fraction. As expected, the K13-GFP protein was present exclusively in the cytoplasmic

fraction independent of the absence or presence of heme. Furthermore, the intensity of K13-GFP was

much higher under heme-replenished condition (top blot). The GFP expressing cells revealed no

changes in the relative nuclear/cytoplasmic fractionation of the Nrf2 protein or intensities of GFP

(**Supplementary Fig. 3c**). As Nrf2 is a redox-controlled transcription factor, we also compared the

expression of stress-response genes *heme oxygenase* (*ho-1*), *NAD(P)H quinone oxidoreductase 1*

(*nqo1*) and *thioredoxin reductase 1* (*txnrd1*) in K13-GFP[HDM] and K13-GFP[HEME] cells. As shown in **Fig.**

2f, RT-qPCR data confirmed increased expression of these stress-response genes as a consequence

of increased Nrf2 translocation to the cell nucleus.

Chaperone mediate autophagy is a major pathway of K13 degradation

201   A recent study implicated chaperone-mediated autophagy (CMA) in the lysosomal degradation

202   KEAP1 in mouse midbrain progenitor cell line SN4741[51]. Since K13 shares structural and sequence

identity with KEAP1, we investigated if similar degradation pathway was involved in the proteolysis of K13-GFP under HDM conditions. Our analyses of the primary sequence of K13, suggested it contained 'KFERQ-like' motifs as recognized by an online algorithm KFERQ finder v0.8 (*https://rshine.einsteinmed.edu*)[52]. These motifs are targeting signals of CMA. In CMA the chaperone heat shock 70-kDa (cognate) protein 8 (also known as HSPA8) binds 'substrate' proteins bearing 'KFERQ-like' motifs[53,54]. LAMP2A acts as a receptor on the lysosomal membrane. HSPA8 binds the cytosolic tail of LAMP2A[55]. This in turn unfolds the substrate and enables its translocation into lysosomes where it is subsequently degraded[56]. As shown in **Fig. 3a**, K13 contains eighteen 'KFERQ-like' motif, strongly suggesting it may be targeted to lysosomes. Immunolocalization studies using indirect IFA suggested moderate association between K13 and LAMP2A with Pearson's correlation coefficient (PCC) of 0.59. Notably, heme raised the PCC to 0.79, suggesting it induced a high level of co-association of K13-GFP with LAMP2A (**Fig. 3b**). We therefore concluded that although heme induces conformational change in K13, this does not obscure targeting of K13-GFP to LAMP2A labeled lysosomes. But since heme increases levels of K13, it is not likely to promote degradation of K13-GFP within the lysosome. We therefore propose (as schematic in **Fig. 3b**) that the heme-induced conformational stability may prevent unfolding of the K13-GFP substrate and its translocation into the lysosome, thereby limiting its degradation in cells.

Since LAMP2A is an exclusive CMA marker, we investigated the effects of a previously reported CMA-inhibitor, *i.e.*, the RARα (retinoic acid receptor alpha) activator ATRA (*all-trans* retinoic acid), known to be a potent inhibitor of LAMP2A expression[57]. As expected, under HDM conditions, there was moderate co-association of K13 and LAMP2A, with a PCC of 0.58 (**Fig. 4a i**). Treatment with ATRA elevated levels of K13-GFP as well as its co-association with LAMP2A (raising the PCC to 0.90; **Fig. 4a ii**). As expected, heme also increased K13-GFP levels and the PCC with LAMP2A rose to 0.77 (**Fig. 4a iii**). Notably, ATRA markedly increased K13-GFP in HDM to levels far higher than those seen with heme replenishment (**Fig. 4b i** and **4b iii**). ATRA also significantly reduced LAMP2A levels compared to controls, both in K13-GFP[HDM] and K13-GFP[HEME] cells (**Fig. 4b i** and **4b ii**). Together these data strongly suggested ATRA-mediated reduction of LAMP2A prevented the degradation of K13, explaining why it increased K13 levels more than heme.

To further confirm the role of LAMP2A-mediated degradation of K13-GFP, we obtained stable

transgenic cell lines expressing K13-GFP or GFP in the HeLa *lamp2a* knockout (*Δlamp2a*; Abcam;

ab255402) and the corresponding parental HeLa cells (Abcam, ab255928). HeLa cells have

previously been reported to inherently express very low levels of native KEAP1 protein and

correspondingly have high basal Nrf2 activity[58]. We designated them as *Δlamp2a*[K13-GFP], HeLa[K13-GFP],

Δlamp2a[GFP] and HeLa[GFP], cells, respectively. As shown in **Supplementary Fig. 4a**, LAMP2A protein

was undetectable in *Δlamp2a* cells both by fluorescence microscopy and in western blots. Moreover,

there was no difference in levels of K13-GFP in the transgenic *Δlamp2a*[K13-GFP] cells regardless of

whether they were cultured under HDM or 5 µM heme-supplemented conditions (**Fig. 4c**). In contrast,

significant reduction in the K13-GFP level was evident in the parental HeLa[K13-GFP] cells in HDM

(compare lanes 3 and lane 4). In *Δlamp2a*[GFP] cells, we observed no changes in the level of GFP

(**Supplementary Fig. 4b**) when compared to the HeLa[GFP] cells irrespective of HDM or heme-

replenished conditions. Thus, the absence of LAMP2A adversely affected the proteolysis of K13-GFP,

but not GFP, which further substantiated the involvement of CMA pathway in proteolysis of K13-GFP

under HDM conditions.

K13's kelch domain transfers heme-dependent stability and heme binding property to KEAP1

We next investigated if the heme-regulated stability of K13 was due to its kelch repeat (KRP) domain

and could be transferred to KEAP1. As shown in **Fig. 5a**, we made chimeras of the N-terminus of

KEAP1 and the kelch domain of K13 fused to GFP (KEAP1[N]K13[KRP]-GFP) as well N-terminus of K13

and the kelch domain of KEAP1 fused to GFP (K13[N]KEAP1[KRP]-GFP). Stable expression of

KEAP1[N]K13[KRP]-GFP and K13[N]KEAP1[KRP]-GFP proteins in Du145 cells were confirmed by western

blots (**Fig. 5b**). As shown in **Fig. 5c**, immunolabeling assays revealed significant (72%) reduction in

KEAP1[N]K13[KRP]-GFP cells under HDM conditions when compared to respective counterparts in 5 µM

heme-supplemented media. In contrast K13[N]KEAP1[KRP]-GFP cells were reduced by only 30%.

Supplementing HDM with the heme precursor ALA had no significant effect, but heme largely restored

levels of KEAP1[N]K13[KRP]-GFP expression (**Fig. 5d**). K13[N]KEAP1[KRP]-GFP showed much less heme-

responsive expression (**Fig. 5e**). Together, these results suggested that the kelch domain of K13

conferred robust heme-stimulated expression of the KEAP1-K13 chimera which was closely

259 comparable to the wild-type K13-GFP. The NTR, CCD and BTB-POZ domains of K13 also made a

260 minor contribution in transferring heme-dependent phenotype to the KEAP1 chimera.

261 In addition, we investigated the relative affinity of recombinant K13 (also known as TrK13-WT),

262 KEAP1, KEAP1^{N}K13^{KRP} and K13^{N}KEAP1^{KRP} proteins for hemin-agarose. Recombinant proteins

263 expressed and purified from *E. coli* were incubated with hemin-agarose beads and percentage

264 binding was determined (see *Materials and Methods*). As shown in **Fig. 5f**, K13/TrK13-WT, KEAP1,

265 KEAP1^{N}K13^{KRP} K13^{N}KEAP1^{KRP} proteins respectively showed 30.1%, 5.9%, 32.5% and 10.3% binding

266 to hemin-agarose, respectively. Again, this suggested that the kelch domain of K13 could confer the

267 same efficiency of heme binding to KEAP1 as seen associated with K13/TrK13-WT protein. When

268 soluble cellular extracts from stable transgenic Du145 cells ectopically expressing the K13-GFP,

269 KEAP1 and their chimera proteins were used, we again found that K13-GFP, KEAP1,

270 KEAP1^{N}K13^{KRP}-GFP and K13^{N}KEAP1^{KRP}-GFP bound hemin-agarose beads with efficiencies of

271 58.9%, 3.1%, 64.5% and 29.3%, respectively (**Fig. 5g**). Cumulatively, the results in **Fig. 5** strongly

272 support that the kelch domain of K13 transfers robust heme binding capacity and heme-based

273 stabilization to KEAP1. The N-terminal domains of K13 may also contribute a minor role consistent

274 with our prior studies suggesting a subset of heme binding motifs are also detected outside of K13's

275 Kelch domain[24].

K13 and its chimeras confer sensitivity to heme-dependent ART-killing of Du145 cells

277 Since heme is implicated in the activation of ART, we next assessed the ART sensitivity of Du145

278 cells expressing K13-GFP and its chimeras (as well as GFP alone) in a standard killing MTT [3-(4,5-

279 dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay (see *Materials and Methods*) comparing

280 outcomes in HDM or media replenished with 5 µM heme. Du145 cells expressing control GFP

281 showed an IC$_{50}$ of 117.7 ± 0.9 µM in the absence of added heme (**Fig. 6a i**), thereby indicating that

282 they are intrinsically highly resistant to ART. The addition of 5 µM heme minimally reduced the IC$_{50}$ by

283 2.6%. Expression of KEAP1 also showed a small effect inducing approximately 6.1% reduction.

284 However, K13-GFP and the KEAP1^{N}K13^{KRP} cells had a greater effect, which translated into increasing

285 heme-dependent sensitivity to ART by 18.2% and 23.7%, respectively (**Fig. 6a i-ii**). K13^{N}KEAP1^{KRP}-

286 GFP had lesser effect showing ART sensitivity increase of 13.5%. The relative order of conferring

287 increased ART sensitivity was KEAP1^{N}K13^{KRP}-GFP > K13-GFP > K13^{N}KEAP1^{KRP}-GFP > KEAP1 >

GFP, which matched the heme binding properties reported for these transgenes in **Fig. 5**. Together, these data suggested that the presence of K13, particularly its kelch domain, conferred sensitivity to ART in the presence of 5 µM heme.

Since the K13 also binds Nrf2 (which is antagonized by heme), we examined the effects of ART on the dynamics of K13 and Nrf2 in HDM as well as after heme supplementation. As shown in **Fig. 6b i-ii,** ART had no sizeable effect on K13-GFP levels in the presence or absence of heme. However, Nrf2 levels were reduced by 50% in the presence of both ART and heme. This reduction was due to the depletion of Nrf2 in the cytoplasm (**Fig. 6c i-ii**), suggesting it was through regulation by K13. These data support a model shown in **Fig. 6d**, where heme plays a major role in binding K13 to displace Nrf2, which in turn is targeted to the nucleus to induce a protective redox (or pro-survival) response. Heme bound to K13 activates ART to generate toxic heme-ART adducts. K13's renewed ability to regulate Nrf2 suggests that these toxic heme-ART adducts do not inactivate K13. Rather, the adducts dislodge from K13 and disperse to mediate damage which occurs in absence of a protective redox response, resulting in cell death (or pro-death) responses. Thus, ART-induced killing reconstituted in Du145 cells may arise from the dual effect of inducing toxic ART-heme adducts as well as preventing a protective redox response to render an overall 'pro-death response (**Fig. 6d**).

ART sensitivity is increased proportional to the K13 intensity

Our findings in **Fig. 6** strongly supported that expression of K13 or chimeras, containing its kelch domain in presence of 5 µM heme, increased sensitivity to ART. We wanted to gain better understanding on whether increasing K13 at sub-micromolar heme concentration also played a role. Since our studies in **Fig. 4** established that ATRA increased levels of K13, we separately evaluated the effects of heme and ATRA. As shown in **Fig. 7a, b**, in cells expressing GFP, 5 µM heme increased sensitivity to ART by ~ 6%, while ATRA had no effect. In K13-GFP cells, heme alone stimulated ART-sensitivity by 18%, ATRA by 25%, and the combination of two stimulated ART-sensitivity by 36% (**Fig. 7a, b**). These data strongly suggested that elevation in levels of K13-GFP by ATRA even under HDM conditions increased sensitivity to ART. But heme at 5 µM provided additional sensitization effect possibly by (i) increasing K13 levels (ii) providing substrate to form toxic heme-ART adducts, whose disassociation from K13 suppression of pro-survival responses (as hypothesized in **Fig. 6d**).

We were therefore interested in determining whether ART sensitivity was proportional to K13

intensity. To do so, we compared the sensitivity to killing by ART to levels of K13 (or its kelch chimera)

normalized to GAPDH (a control cytoplasmic protein). As shown in **Fig. 7c**, we observed an overall

linear relationship between increase in ART sensitivity and the K13/GAPDH (R^2 value = 0.97).

Notably, cell lines designated as 1, 2 and 4 reflected measurements generated in HDM, and all

showed lower degree of ART sensitivity and K13/GAPDH levels than counterpart measurements

generated in the presence of 5 µM heme. Also, the presence of ATRA increased K13 levels, thereby

boosting K13/GAPDH ratio and correspondingly showed increased ART sensitivity of K13-GFP cells

under HDM conditions (as cell line #3 in **Fig. 7c**). Moreover, combination of heme and ATRA in these

cells (designated as 3H) further elevated K13/GAPDH ratio and showed highest sensitivity to ART

(38%). Comparisons in HeLa cells revealed that knockout of LAMP2A, which increased K13/GAPDH

by three-fold raised ART sensitivity by one-third. However, 5 µM heme further raised the K13/GAPDH

ratio four-fold and doubled ART-sensitivity (compared to parental HeLa in HDM).

When we separately compared the effects of extracellular 5 µM *versus* 50 µM heme (**Fig. 7d**), the

higher concentration of heme further augmented killing. In GFP cells, this could be easily interpretable

as increasing intracellular heme alone from ~400 to 750 nM directly sensitized ART-killing, increasing

it from ~2 to ~6%. However, as we have previously shown for K13 kelch domain proteins, heme may

also act by raising levels of the transgene alone. We therefore tracked transgene expression at 50 µM

heme (**Fig. 7d**, black bars). This revealed that even at 50 µM, increasing K13-GFP and K13 chimeras

stimulated ART-killing to much higher levels. Taken together, these data suggest that heme alone (at

intracellular concentrations of 400 nM to 1 µM), is comparatively weak stimulator of ART's killing

activity but can be potentiated by expression of K13 and its appropriate chimeras.

Discussion

K13 was identified as marker of ART-R more than a decade ago[6]. Our studies provide the first

evidence that it acts as redox sensor regulated by the oxidant heme (summarized in **Fig. 8a**). This is

also the first report that a small molecule like heme can bind a member of an evolutionary conserved

kelch family (whilst multiple protein partners have been defined for kelch proteins[15,59–61]). Notably,

K13-heme activates ARTs, suggesting the potent killing action of these drugs exploits a parasite

protective redox response (**Fig. 8a, cycle 1**) Whether *Plasmodium* encodes for functional orthologue

of Nrf2 (the transcription factor that induces the stress-induced protective response in Du145 cells)

remains unknown. However, since heme is a strong oxidant[31], upon its binding to K13, a protective

redox response is likely elicited as a compensatory effect (expected for **cycles 2 and 3** shown in **Fig.**

8a). A redox stress response may be needed to protect the parasite against high levels of heme

reported at all stages of parasite growth[62].

The models in **Fig. 8a, b** maybe particularly informative for clinically reported ART-induced ring-stage

death[63,64] based on the following considerations. Young rings (0-3 h) lack a digestive food vacuole

and therefore depend on heme from biosynthetic sources, analogous to Du145 cells. Similarly in *P.*

falciparum rings, K13-heme triggers a pro-survival response exploited by ART to spring a pro-death

response of oxidizing ART-heme adducts that are no longer redox active but can kill the ring parasite

(**Fig. 8b i**), and this function is compromised in ART-R (**Fig. 8b ii**). To date, K13 function has been

best understood in context of regulation of uptake of hemoglobin into the food vacuole at the

trophozoite stage. However, it is well documented that K13 is expressed in daughter merozoites that

assemble when trophozoites mature to the segmented schizont stage. K13 is also detected from the

earliest stages of ring infection (mode in **Fig. 8b ii**)[59,65,66]. However, K13's precise location is not well

mapped in *P. falciparum* merozoites and rings. But in the related apicomplexan *Toxoplasma gondii*, a

K13 orthologue was localized to endosomal structures called Type I micropores and shown to

regulate plasma membrane endocytosis[67,68]. Since all apicomplexans (including *Plasmodium*) contain

a 'Type I' micropore in their 'apical' stages[69,70], by virtue of evolutionary conservation of mechanisms

of apical organelles, we have proposed that in merozoites and rings K13 localizes to *P. falciparum*

Type I micropores and regulates essential plasma membrane endocytic functions of these parasite

stages [71] (**Fig. 8b i**). Indeed the *P. falciparum* food vacuole where K13 is abundant develops from a

Type II micropore, which is also consistent with a conserved micropore function for K13.

The Du145 cell model enabled assessment of the effects of K13 binding to heme from nanomolar to

sub-micromolar range. The intrinsic concentration of heme in Du145 cells is 60 nM. Supplementing

with extracellular heme at 5 µM and 50 µM resulted in maximal intracellular concentrations of 0.45 µM

and 0.75 µM, which approach but are nonetheless lower than the published physiological range (1-5

µM) reported for *Plasmodium*[40,41]. Whether kelch proteins of other apicomplexans bind heme has yet

to be investigated. Moreover, although the kelch domain of K13 is necessary and sufficient to confer

heme binding property to its closest mammalian orthologue KEAP1, whether and how heme is released from K13 under physiological conditions is not known, since it cannot be ubiquitinylated and degraded, which is a hallmark of protein substrates and their cellular regulation by kelch proteins. But our studies suggest that when K13-bound heme activates ART, the kelch structure may protect against self-oxidation by heme-ART adducts[24]. This is consistent with prior studies showing that alkyne-ART analog despite being highly active against a wide range of malarial cellular targets failed to alkylate K13[72,73]. Heme-ART adducts show low affinity for recombinant K13[24], suggesting a mechanism for their release and a new model where a single molecule of K13 may catalyze multiple cycles of heme-ART toxic adduct formation and oxidative killing (**Fig. 8a**; K13 cycle 1).

That we find that intracellular heme concentrations of 400- 750 nM alone yield low sensitization to ART, counters prevailing models that suggest that heme alone is sufficient to activate a low level of ART at the trophozoite stages (**Fig. 8c**). Notably, our data show that even at the lowest intensities, K13 (expected not to exceed nanomolar levels) was more effective than heme alone and in addition, the extent of sensitization was directly proportional to the level of K13. This is consistent with our findings that raising K13 (by ATRA treatment) in HDM increased ART-sensitivity more than an eight-fold in free heme concentrations of 60 nM. These data strongly support the idea that K13 acts as 'catalyst' to facilitate assembly of activated ART-heme complexes, and once the activated toxic adducts are released, K13 may facilitate successive rounds of ART-based killing. We therefore propose that even at the trophozoite stage, rather than heme alone, it is the K13-bound heme which effects sensitization to ART (**Fig. 8c**). These stages are known to be far more sensitive to ART than rings, and trophozoites expressing K13 mutations of ART-R are highly susceptible to killing by ART, which is consistent with their higher levels of K13.

K13-heme concentration at late endosomes in Du145 cells suggests that in addition to triggering a pro-survival response, heme binding to K13 may stabilize endosomal location in both rings and trophozoites and possibly create concentrated cellular foci of K13 function as well as efficiently activate ART at all stages of parasite growth. That heme binding, endosomal concentration and ART-based killing are not properties of mammalian KEAP1, suggests they are parasite evolutionary adaptations conferred by the K13 kelch domain. Yet, orthologues of LAMP2A (a receptor for K13 in Du145 and HeLa cells that controls its autophagic degradation), have not been reported in *P.*

falciparum or *T. gondii.* Nonetheless, autophagic systems exist in both parasites[74,75] and additional

studies are needed to determine their influence on heme-dependent K13 stability, as well as

endosomal degradation and turnover of K13 and its orthologues across apicomplexans. Another key

priority is identification of K13-coupled parasite redox factors that remain unidentified because a lack

of conservation at the level of primary sequences prevents easy prediction from nucleotide sequence

comparisons. While our studies establish that K13 is an important determinant in ART sensitization,

other redox and oxidative factors may also play a role. Our data suggest Du145 may continue to

provide a powerful model system for future functional screens of endosomal effectors as well as novel

plasmodial pro-oxidant responses, regulated by both heme and K13 and targeted in killing by ART.

One generalized lesson already learned may be that suppression of redox sensing may sensitize a

wider range of infectious agents as well as cancers to killing by ARTs.

Materials and methods

Generation of plasmid constructs

The pPR-IBA101 (TrK13-WT) plasmid was generated as described in earlier studies[24,47]. For the

generation of pPR-IBA101 (KEAP1), the full length (1875 base pairs) transcript of *keap1* was

amplified from cDNA library prepared from human cell line (HEK293T) using the primer pairs (Hs-

Keap1-BsaI-Fwd (5'-CAAATGGGAGACCTTATGCAGCCAGATCCCAGGC-3') and Hs-Keap1-BsaI-

Rev (5'-CCAAGCGCTGAGACCAGCAGCACAGGTACAGTTCTGCTGGTCAATCTG-3'). The PCR

product was digested with BsaI and cloned at corresponding sites in similarly digested pPR-IBA101.

For the generation of pPR-IBA101 (KEAP1^{N}K13^{KRP}), nucleotide sequences coding for the N-terminus

region of KEAP1 was amplified from pPR-IBA101 (KEAP1) plasmid using (Hs-Keap1-BsaI-Fwd and

KEAP1toK13-KRP-Rev   (5'-AAATCCACCGGCAGCAGCGCGGCCCACCTTGGG-3')   primers.

Nucleotide sequence encoding for the KRP region of K13 was amplified using the pPR-IBA101

(TrK13-WT)   plasmid   as   template   and   KEAP1toK13-KRP-Fwd   (5'-

GTGGGCCGCGCTGCTGCCGGTGGATTTGATGGTGTAGAATATTTAAATTCG-3')   and   PfKelch13-

BsaI-Rev   (5'-GGAGGACAAAATGGCAATGTTCTAAATTCATGTGGATTCTTTTCACCAGAT

ACAAATGAATGGC-3') primer pairs. The two PCR products were used as templates to reconstitute

KEAP1^{N}K13^{KRP} by overlapping PCR, which was then digested with BsaI and cloned into pPR-IBA101.

Similarly, to generate pPR-IBA101 (K13^{N}KEAP1^{KRP}), nucleotide sequences encoding for the N-

terminus of K13 was amplified from pPR-IBA101 (TrK13-WT) plasmid as template and PfKelch13-BsaI-Fwd (5'-CAAATGGGAGACCTTATGGAAGGAGAAAAGTAAAAACAAAAGCAAATAGTATCTCG-3') and K13toKEAP1-KRP-Rev (5'-GTAGATCAG GGCAGCAGCTATACAAAATACTAATGGGAATGGTAAAAATTTAATACC-3') primers. Overlapping PCR, using the above two PCR products as templates, generated K13^{N}KEAP1^{KRP} which then introduced to pPR-IBA101 after digestion with BsaI.

For generating recombinant Nrf2-6×his, the full length (1626 bp) transcript was amplified from cDNA library prepared from human cell line (Du145) using the primer pairs HsNrf2-XbaIFwd (5'-CACGTGTCTAGAATGATGGACTTGGAGCTGCCG-3') and HsNrf2ns-XhoIRev (5'-CCTAGGCTCGAGGTTTTTCTTAACATCTGGCTTCTTACTTTTGGG-3') and the PCR product was then digested with XbaI-XhoI and clone into similarly digested pET303-ct-his plasmid (ThermoFisher, USA) for in-frame fusion with C-terminus 6×his tag. To generate GFP expressing constructs for mammalian cells, pAcGFP1-C1 (Addgene, USA) was used and AgeI and NheI sites considered for inserting the gene of interest. To generate pAcGFP1-C1 (K13-GFP), pAcGFP1-C1 (KEAP1^{N}K13^{KRP}) and pAcGFP1-C1 (K13^{N}KEAP1^{KRP}) plasmids, their pPR-IBA101 plasmids were used as corresponding templates. For K13-GFP, the primers pairs were PfKelch13-NheI-Fwd (5'-AAGCTTGCTAGCATGAACAGCGAGGTGTGTTCGCG-3') and PfKelch13-AgeI-Rev (5'-AGCGCTACCGGTAGCAGCCAAATTTGCCAAGAGAATGCCGTG-3'), for generating KEAP1^{N}K13^{KRP}GFP, the primer pairs were KEAP1-NheI-Fwd (5'-AGATCTGCTAGCATGCAGCCAGATCCCAGGC-3') and PfKelch13-AgeI-Rev (5'-AGCGCTACCGGTAGCAGCCAAATTTGCCAAGAGAATGCCGTG-3'); and for generating K13^{N}KEAP1^{KRP}, PfKelch13-NheI-Fwd (5'-AAGCTTGCTAGCATGAACAGCGAGGTGTGTTCGCG-3') and KEAP1-AAA-AgeIRev (5'-GGATCCACCGGTAGCTGCTGCACAGGTACAGTTCTGC TGGTCAATC-3') primers were used. To generate the KEAP1 construct, pPR-IBA101 (KEAP1) plasmid was used as the template to amplify nucleotides encoding for KEAP1-strep using KEAP1-NheI-Fwd and Strep-XhoI-Rev (5'-GGATCCCTCGAGCTTTCTCGAACTGCGGGTGGCTCCAA-3'). The PCR amplified product was subjected to NheI-XhoI digestion and inserted into similarly digested pAcGFP1-C1.

462   All constructs generated for this study were confirmed by Sanger's sequencing prior to further

463   experiments.

Cell culture, generation of transgenic lines and their treatments

The Du145 cell line was the kind gift from Prof. Rana Pratap Singh, SLS, JNU, India. The parental

HeLa (ab255928) and HeLa-Δ*lamp2a* (ab255402) were procured from Abcam, USA and cultured

according to the manufacturer's instructions in complete media RPMI (Gibco, USA). Cells were

transfected using Lipofectamine 3000 (Invitrogen, USA) following the manufacturer protocol. Stable

transgenic lines were generated by G418 (0.5 mg/ml, Sigma, USA) selection. For the heme depleted

(HDM) conditions, the cells were grown in RPMI media supplemented with heme-depleted FBS and

succinyl acetone (0.5 mM, Caymanchem, USA) for 48 h. For heme-replenished media 5 µM hemin

chloride (Merck, USA) was added to the HDM. To assess *in vivo* stability of K13 protein upon heme

binding, the cells were treated with heme analogs PPIX (50 µM, Merck, USA) or Zn-PPIX (50 µM,

Merck, USA) for 48 h. Cells were also subjected to treatment either with for 48 h with ART (50 µM,

Caymanchem, USA) in presence or absence of heme.

Preparation of heme-depleted FBS

Commercially purchased FBS (Gibco, USA) was incubated with ascorbic acid (10 mM, SRL, India) at

37°C in a shaking incubator at 200 rpm for overnight. Spectral scans of the pre- and post-treated FBS

measured to check the presence or absence of soret peak at 410 nm, respectively using JASCO

UV660 spectrophotometer. Finally, the ascorbate-treated FBS was dialyzed against PBS, pH 7.4, with

at least three periodic changes for 24 h at 4°C. Dialyzed FBS was filter-sterilized using 0.2 µm

polyether sulfone filter (Sartorius, USA).

Assessment of cellular viability

Cell viability was assessed using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT,

Merck, USA)-based colorimetric assay. Cells at 50-60% confluency and within 72 h incubation in the

96-well cell culture plate were only considered. Following treatment, the cells incubated with 20 µl of 5

mg/ml MTT solution for 2h at 37° in the dark. Finally, the culture media was discarded, and formazan

crystal was dissolved with 200 µl of DMSO, and absorbance was recorded at 570 nm using a

microplate reader (Thermo Fisher Scientific, USA). Cell viability was calculated and expressed as

percentage in comparison to the untreated controls. For MTT assay, the viability was calculated for

six different concentrations (0, 40, 80, 120, 160 and 200 µM) in three replicates to generate the IC_{50}

values for ART.

Estimation of total cellular heme content

Cells were grown up to 90-95% confluency and dislodged from the plate by treatment with trypsin and

were washed twice with saline PBS. Cell count was recorded prior to lysis using hemocytometer.

Cells were then solubilized in five volumes of RIPA buffer for 2 h at 4ºC. After a brief centrifugation,

the supernatant was diluted ten times and subjected for spectrophotometric scanning at 410 nm. The

absorbance was plotted against the standard curve of hemin chloride to calculate the total cellular

heme content. The standard curve has been generated using gradually increasing concentration (0,

0.05, 0.1, 0.5, 1 and 10 µM) of hemin chloride.

Nuclear and cytoplasmic fractionation of transgenic cells

Cellular fractionation into nuclear and cytoplasmic fractions was performed following previously

described protocol[76]. Briefly, the cells from the 60 mm culture plates were scrapped and washed twice

with ice-cold phosphate buffered saline (PBS) pH 7.4. The cell pellets were washed, resuspended in

250 µl of ice-cold 0.2% (v/v) NP40 (HiMedia Labs, India), minced with micro-pestle, vortexed and kept

on rocker shaker for 15 mins. The resuspended pellets were then centrifuged at 10,000 rpm for 1 min,

and the supernatant was collected as the cytoplasmic fraction. The pellet was then resuspended in 1

ml ice-cold 0.2% (v/v) NP40 to remove remnant cytoplasmic debris and centrifuged at same speed.

Finally, the pellet was resuspended in 200 µl ice-cold PBS and designated as the nuclear fraction.

The nuclear fraction was also briefly sonicated using micro pulser (Sonics, USA).

Indirect immunofluorescence assay (IFA)

IFA was performed using parental or transfected cell lines. Briefly, cells were initially grown on poly-L-

lysine coated coverslips and fixed using 4% (w/v) paraformaldehyde/ 0.0075% (v/v) glutaraldehyde

followed by permeabilization with 0.1% (v/v) Triton X-100 and neutralized with 50 mM NH_4Cl. Blocking

was done using 3% (w/v) BSA. The coverslips were incubated with respective primary antibodies (10

µg/ml) at 4°C for overnight followed by incubation in FITC- or TRITC/ Rhodamine-conjugated

secondary IgG antibodies (1:200; MP biomedicals, USA). Cellular nuclei were stained with 5 µg/ml

Hoechst 33342 (Molecular Probes) and slides were mounted with DABCO. The slides were finally

visualized by SP8 confocal laser scanning microscope (Leica, Germany). Fluorescence intensity was

calculated using Image-J software (*https://imagej.net/ij/*) software by considering 3 biological

replicates for each set and each replicate containing 30 individual cells. The RGB intensity plot and

Person's correlation coefficient (PCC) was generated from the Image-J graphics and JaCop,

respectively.

SDS-PAGE and western blotting

The induced bacterial cell pellet was directly solubilized in Laemmli's sample buffer and heat

denatured for 15 min at 95°C. For mammalian cells, following scrapping from plates, cells were

washed twice with saline PBS and solubilized in five volumes of RIPA buffer for 2 h at 4°C. After a

brief centrifugation, the supernatant was considered as whole protein pool and boiled for 15 min at

95°C using Laemmli's sample buffer and resolved by 8-10% SDS-PAGE followed by Coomassie or

silver staining. For subsequent western blots, resolved proteins were transferred into nitrocellulose

membrane (MDI, USA) followed by blocking with ice cold 5% (w/v) fat-free skim milk (SRL, India) for 2

h at RT. The membranes were incubated with specific primary antibodies (1:3000 to 1:5000 dilution)

for overnight at 4°C on a platform shaker followed by subsequent washing and incubation with

respective HRP-conjugated secondary antibodies (1:7000 to 1:10000 dilution). Blots were finally

developed by ECL assay and detected using gel imager (Bio-Rad, USA). The primary antibodies are

used in this study were: anti-GFP (Invitrogen, USA; Cloud clone, China), anti-UBQ, anti-Lamp2 and

anti-GAPDH (Cloud clone, China), anti-k13 (Custom generated, Biobharati Life Sciences[47]), anti-

Histone 3 (kind gift from Prof. Suman Kumar Dhar, SCMM, JNU, India), anti-Nrf2, anti-KEAP1, anti-

HO1, anti-NQO1, anti-MCM3, anti-DPP3 and anti-TRAF2 (ABclonal, Woburn), anti-His and

streptavidin-HRP (BioBharati Life Sciences, India), and anti-HA. The subsequent HRP conjugated

secondary antibodies used in this study were goat anti-rabbit (Bio-Rad, USA), goat anti-mouse, goat

anti-rat (MP biomedicals, USA). Band intensity was calculated from Image-J (*https://imagej.net/ij/*)

software by considering 3 individual blots for each data set.

Purification of recombinant protein and *in vitro* trypsin digestion

Recombinant Trk13-WT, KEAP1, K13^{N}KEAP1^{KRP}and KEAP1^{N}K13^{KRP} proteins as C-terminus fusion

with strep tag and Nrf2-6×his were purified from isopropyl-1-thio-β-D-galactopyranoside (IPTG, SRL,

India) and arabinose (0.2%, SRL, India) induced bacterial pellet. Briefly, the induced bacterial pellet

was lysed using lysozyme (SRL, India), sonicated and washed thrice with 2% (w/v) sodium

deoxycholate (SRL, India) to purify inclusion bodies (IBs). The IBs were dissolved into 2 M urea, 50

mM Tris-HCl buffer pH 12.5 and then dialyzed against buffer W (50 mM Tris-HCl, 150 mM NaCl, pH

8). Dialyzed solution was kept for binding to StrepTactin XT Superflow resin (IBA, Germany) for

Trk13-WT, KEAP1, K13^{N}KEAP1^{KRP}and KEAP1^{N}K13^{KRP} and Ni-NTA resin (Thermo, USA) for Nrf2-

6×his. Purified proteins were dialyzed overnight at 4°C against buffer W with at least three periodic

changes. Purity of each protein was checked by SDS-PAGE and Coomassie staining and further

confirmed by western blotting using commercial antibodies against the 6×his tag, strep tag (Biobharati

Life Sciences) or custom-generated anti-K13.

For *in vitro* trypsin digestion, 10 µg of recombinant TrK13-WT or commercially procured BSA (A3294,

Sigma, USA) was incubated with or without equimolar concentrations of heme in buffer W for 1 h.

Samples were desalted as described in our earlier study[24], and treated with 0.5 µg trypsin (T6567,

Sigma-Aldrich) for different incubation times. Following digestions, Laemmli's sample buffer was

immediately added to each sample, boiled for 15 min at 95°C and processed for SDS-PAGE and

Coomassie staining.

***In vitro* protein-protein interactions**

The purified recombinant Nrf2-6×his (10 µM) was incubated with Ni-NTA beads overnight and washed

thrice with buffer W containing 30 mM imidazole. The Ni-NTA bound recombinant Nrf2-6×his was

then incubated with purified recombinant TrK13-WT (10 µM) in buffer W for 4 h at RT. The beads

were thoroughly washed with buffer W, and Ni-NTA-bound proteins were resolved by SDS-PAGE

followed by Coomassie staining or western blot using antibodies against the 6×his tag, strep-tag

(Biobharati Life Sciences) or custom-generated anti-K13. For experiments involving the interactions

between heme-bound or heme-free recombinant TrK13-WT and Nrf2-6×his, the TrK13-WT was

preincubated with heme (0 or 10 µM) at RT for 1 h at room temperature (RT) and desalted using PD

Spintrap™ G-25 columns (Cytiva) to remove excess unbound heme.

Hemin agarose binding

Affinity-purified recombinant Trk13-WT, KEAP1, K13^{N}KEAP1^{KRP}and KEAP1^{N}K13^{KRP} proteins (10 µM)

were incubated with 50 µl suspensions of 50% v/v hemin-agarose (Merck, USA) beads for 1 h at RT

in buffer W, pH 8, under rotating conditions, followed by five washes with ten volumes of buffer W to

remove unbound proteins. The hemin-agarose beads were then solubilized in Laemmli's sample

buffer, resolved by SDS-PAGE, and visualized by Coomassie staining. For assessing hemin-agarose

binding from transgenic cellular lysates, the total cell pellet was resuspended in 5 volumes of ice-cold

RIPA buffer complemented with 1X protease inhibitor cocktail (cOmplete, Roche) and incubated at

rotating wheel for 1 h. After centrifugation (15 mins, 21,100 $\times$ g, 4°C), the supernatant containing

soluble proteins fractions was incubated with hemin agarose beads for 4 h. Unbound proteins in the

lysates were washed with ten volumes of RIPA buffer and bound fractions were then solubilized in

Laemmli's sample buffer and subjected to western blotting using anti-strep tag antibody (Biobharati

Life Sciences).

Co-immunoprecipitation

Protein A beads (IBA Life Sciences, Germany) were incubated with 10 µg of respective primary

antibodies for 4 h at 4 °C under rotating conditions and unbound antibodies were removed by

extensive washes with pre-cooled PBS. Mammalian cells were harvested, lysed with five volumes of

RIPA buffer supplemented with 1 mM PMSF (Thermo, USA) and protease inhibitor cocktail (Roche)

and centrifuged at 10,000 $\times$ g for 15 min at 4°C to remove the cell debris. Clarified lysates were then

incubated with antibody-coated protein A beads (30-50 µl, 50% v/v resuspension) for overnight at 4°C

under rotating conditions. Following binding, unbound lysates were removed and the beads washed

twice with RIPA buffer and twice with ice-cold saline PBS. Beads were boiled with Laemmli's sample

buffer for 15 min at 95°C and processed for western blotting.

Total RNA, cDNA synthesis, and real time RT qPCR

The total RNA was isolated by Trizol reagent (Invitrogen, USA). Briefly, cell pellets were washed twice

with PBS and homogenized using 500 µl of Trizol reagent. The supernatants were isolated after a

brief centrifugation, mixed with 300 µl chloroform and incubated at RT. After centrifugation at 13,000 $\times$

g for 15 min at 4°C, the upper aqueous phase was removed and 500 µl isopropanol added. Following

incubation for 30 min at RT, the tubes were centrifuged at 13,000 $\times$ g for 15 min at 4°C. The pellet

was washed twice with 70% ethanol and finally dissolved in TE. Samples were treated with DNase I

enzyme (Thermo scientific, USA) to remove genomic DNA contamination. The quantity and quality of

RNA was estimated using Bio spectrophotometer (Eppendorf, Germany) and agarose-formamide gel

electrophoresis. Approximately 1 µg of total RNA was used for cDNA synthesis using PrimeScript 1st

strand cDNA Synthesis Kit (Takara-Bio, Japan) following manufacturer's protocol. For real time RT-

qPCR analyses, 2 µl of ten-fold diluted stock solution of cDNA samples was used. To perform RT-

qPCR analyses, primers were designed based on the coding sequence of specific genes using Oligo

Analyzer software (*https://eurofinsgenomics.eu/en/ecom/tools/oligo-analysis/*). Primers pairs were

HsHO1_RT_Fwd (5'-ATGACACCAAGGACCAGAGC-3') and HsHO1_RT_Rev (5'-

GCATAAAGCCCTACAGCAACT-3'); HsNQO1_RT_Fwd (5'-CCTTCCGGAGTAAGAAGGCAG-3') and

HsNQO1_RT_Rev (5'-TCCAGGCGTTTCTTCCATCC-3'); HsTXNRD1_RT_Fwd (5'-

GGAACTAGATGGGGTCTCGG-3') and HsTXNRD1_RT_Rev (5'-TCTTGCAGGGCTTGTCCTAA-

3'). RT-qPCR analyses were performed by using HOTFIRE Pol EvaGreen qPCR Mix Plus (ROX)

(Solis BioDyne, USA) and Step one plus system (Applied biosystem, USA). To confirm the absence of

any primer dimers in the amplified products, a standard melting curve analysis was conducted.

Primers for *gapdh* included HsGAPDH_RT_Fwd (5'-CAGGAGGCATTGCTGATGAT-3') and

HsGAPDH_RT_Rev (5'-GAAGGCTGGGGCTCATTT-3') as reference. Relative fold change was

calculated by $2^{-\Delta\Delta Ct}$ method.

Statistical validation

One-way ANOVA was carried out to test significance in difference considering at least three biological

replicates using GraphPad prism 10. The adjusted p values (p.adj ≤ 0.001) were obtained via Tukey's

HSD.

Author contributions

Conceptualization Ideas: S.D., K.H., S.B.[1] and S.B.[2].

Methodology: S.D., M.F.

Validation: S.D.

Investigation: S.D.

Resources: S.B.[1], S.B.[2], K.H.

Writing: S.D., K.H., S.B.[1]

Supervision: K.H., S.B.[1]

Project Administration: S.B.[1]

Funding acquisition: S.B.[1]

634 **Acknowledgements**

635 This work was supported by research grants from the Indian Council of Medical Research (ICMR),

636 (58/25/2020/PHA/BMS) and Indian Ministry of Human Resource Development (MHRD) Scheme for

637 Translational and Advanced Research in Sciences (STARS) (MoE-STARS/STARS-2/2023-0111) from

638 the Government of India (SB). SD is a recipient of Department of Biotechnology Research Associate

639 fellowship (DBTRA/2023 24/N/JNU/118) from the Government of India.

640 **Funding sources and disclosure of conflicts of interest**

641 The authors declare that the funding source(s) had no role in the study design, data collection,

642 analyses, and interpretation, and in the decision to submit the article for publication. The authors also

643 declare that they have no competing interests related to this manuscript.

644 **Data availability**

645 The authors declare that the source data supporting the findings of this study are available within the

646 paper and the associated supplementary information files (Supplementary Figures 1-4).

647 **Materials & Correspondence**

648 Correspondence and material requests should be addressed to Souvik Bhattacharjee

649 (*souvik@jnu.ac.in*)

650 **Figure and figure legends**

651

Fig. 1

Fig. 1. Heme stabilizes K13-GFP and confers *in vitro* resistance to trypsin digestion. **a.** Western blots (top) and densitometric analysis (bottom) for K13-GFP under HDM condition or when supplemented with increasing concentrations of heme. GAPDH is the loading control. **b.** IFA images

(left) and mean fluorescence intensity plot (MFI, right) of K13-GFP cells under control, HDM or 5 μM

heme-supplemented conditions. Scale bar, 5 μm. **c.** Western blot (left) and quantitative densitometric

analysis (right) of K13-GFP band intensity under HDM condition (lane 1), or upon supplementation

with heme analogs PPIX (lane 2), Zn-PPIX (lane 3) or increasing concentrations (μM) of heme (lanes

4-7). Values in parenthesis indicate concentration (in μM). **d.** Coomassie-stained SDS-PAGE showing

trypsin-digested fragments (blue arrowheads) derived from the full length (FL, black arrowhead)

recombinant K13 (TrK13-WT) in heme-free (lanes 2-4) and heme-bound TrK13-WT (lanes 6-8)

protein. The heme-free and heme-bound TrK13-WT proteins without trypsin treatment are shown in

lanes 1 and 5, respectively. **e.** Coomassie-stained SDS-PAGE showing trypsin-digested fragments

(blue arrowheads) derived from the recombinant FL TrK13-WT protein (black arrowhead) when

preincubated with heme analogs PPIX, Zn-PPIX, FAS or FES (lanes 4, 6, 8 and 10, respectively) and

not by the heme-bound TrK13-WT (lane 2) protein. Lanes 1, 3, 5, 7 and 9 represent samples with no

trypsin treatment. Molecular weight standards (M) for all SDS-PAGE and western blots (in kDa) are as

indicated. Graphs represent mean from three biological replicates ± SE. Statistical significance

calculated by one-way Anova. Distinct alphabets represent significant difference at p.adj ≤ 0.05.

Fig. 2

Fig. 2. Heme alters the level and dynamics of the redox reporter Nrf2 in K13-GFP-expressing

cells. a. i. Western blots showing immunoprecipitated K13-GFP (top) and the co-associated Nrf2

(bottom) protein from K13-GFP cells cultured in HDM or in the presence of different concentrations of

heme. **ii.** Total Nrf2 (top) and GAPDH (bottom, as loading control) in the cellular lysates for conditions

in a. **b.** Graphical densitometric analyses of K13-GFP, total Nrf2 and co-purified Nrf2 for data shown

in a. **c.** Western blots (left) and graphical densitometric analysis (right) showing reduced co-

purification of the heme-bound (lane 4) recombinant K13 (TrK13-WT) by the recombinant Nrf2-6×his

protein (pre-bound to Ni-NTA beads) as compared to the heme-free TrK13-WT protein (lane 3) .

Lanes 1 and 2 indicate TrK13-WT input and the Ni-NTA bound Nrf2-6×his protein, respectively. Lanes

5 and 6 show no nonspecific binding of heme-free or heme-bound TrK13-WT protein to the Ni-NTA

beads, respectively. **d.** IFA images showing increased levels of K13-GFP (green) fluorescence and

nuclear localization of Nrf2 (red; right panel) in the K13-GFP[HEME] cells (right). In K13-GFP[HDM] cells

(right), the Nrf2 fluorescence distribution was predominantly cytoplasmic. Scale bar, 5 μm. **e.**

Western blots and graphical densitometric quantitation for cytoplasmic and nuclear fractionation of

Nrf2 in K13-GFP[HDM] and K13-GFP[HEME] cells. **f.** RT-qPCR data showing relative expression of stress-

response genes in K13-GFP[HDM] cells and K13-GFP[HEME] cells. Molecular weights (in kDa) are as

indicated. Error bars represent mean of three technical replicates ± SE. Statistical significance was

calculated by one-way Anova. Distinct alphabets represent significant difference at p.adj ≤ 0.05.

Fig. 3

Fig. 3. Signature motifs of autophagy in K13 and heme-dependent co-association with
LAMP2A. a. Schematic showing the distribution of 18 putative KFERQ-like motifs in different domains
of K13, as identified by KFERQ finder v0.8 (*https://rshine.einsteinmed.edu*). NTR, N-terminus domain;
CCD, coiled-coil domain; BTB-POZ, Broad-complex, tramtrack and bric-a-brac (BTB)-poxvirus and
zinc-finger (POZ) domain. b-c. IFA images (left) and RGB intensity plots (right) showing the extent of
colocalization between K13-GFP (green) and LAMP2A (red) in K13-GFP[HDM] (b) and K13-GFP[HEME]

697   cells (c). Pearson's correlation coefficient (PCC) values for the selected region (white bar) are

698   indicated in red text. Magnified images of the white-dotted boxed region for respective cells are also

699   shown. Schematic in b represents proposed association of K13 with lysosomes based on the inset

700   magnification. Scale bar, 5 µm is shown at the bottom of each image. Cell nucleus (blue) stained with

701   Hoechst 33342.

Fig. 4

Fig. 4. Reduction or knockout of LAMP2A decreases degradation of K13-GFP under HDM conditions. a. In Du145 cells, IFA images and RGB intensity plots (with PCC values) showing relative colocalization between K13-GFP (green) and LAMP2A (red) in K13-GFPHDM cells in the absence (i) or presence (ii) of ATRA (inhibitor of LAMP2A expression) compared to K13-GFPHEME cells (iii). Corresponding fluorescence and merged images were acquired under identical optics and exposure

conditions. **b.** Western blots (i) and fold change (ii-iii) for LAMP2A (top), K13-GFP (middle) in the absence or presence of ATRA in cells expressing K13-GFP^{HDM} (lanes 1-2) or K13-GFP^{HEME} (lanes 3-4). GAPDH (bottom) was the loading control. **c.** In HeLa cells, IFA images (left), and graphical MFI (right) of K13-GFP fluorescence (green) in transgenic $\varDelta$lamp2a$^{K13-GFP}$ and parental HeLa$^{K13-GFP}$ cells under HDM (-) or heme-supplemented (+) conditions. **d.** Western blots (top) and graphical fold change of K13-GFP (top) in transgenic $\varDelta$lamp2a$^{K13-GFP}$ (lanes 1-2) and parental HeLa (lanes 3-4) cells under HDM (lanes 2 and 4) or heme-supplemented (lanes 1 and 3) conditions. GAPDH was the loading control. Scale bar for IFA images, 5 μm. Boxed (white dotted) images are magnified in sub-panel. For all statistical analyses, each bar denotes the mean from three technical replicates ± SE. Statistical significance was calculated by one-way Anova. Distinct alphabets represent difference at p.adj ≤ 0.05. Molecular weights (in kDa) for all blots are as indicated.

Fig. 5

Fig. 5. The K13 KRP domain confers heme-dependent stability and heme-binding characteristics to KEAP1. a. Schematics for domain organization in K13, KEAP1, K13^{N}KEAP1^{KRP} and KEAP1^{N}K13^{KRP} chimeric proteins. NTR, N-terminus domain; CCD, coiled-coil domain; BTB-POZ, Broad-complex, tramtrack and bric-a-brac (BTB)-poxvirus and zinc-finger (POZ) domain; IVR, intervening region. **b.** Western blot detecting K13^{N}KEAP1^{KRP}-GFP and KEAP1^{N}K13^{KRP}-GFP (arrowhead) chimeras in transgenic Du145 cells. Loading control is GAPDH. **c.** IFA images and mean

fluorescence intensity (MFI) of K13^{N}KEAP1^{KRP}-GFP and KEAP1^{N}K13^{KRP}-GFP cells cultured under

HDM or heme-repleted conditions. Nuclei stained with Hoechst 33342 (blue), scale bar, 5 µm.

Respective boxed (white dotted) regions are magnified and shown below. **d-e.** Western blots (left)

and densitometric fold change (right) in KEAP1^{N}K13^{KRP}-GFP (d) and K13^{N}KEAP1^{KRP} (e) protein band

intensities under HDM, heme-repleted or heme precursor aminolevulinic acid (ALA)-containing HDM

conditions. **f-g.** Coomassie-stained SDS-PAGE (f) and western blots (g) showing hemin agarose

binding by indicated KEAP1-K13 recombinant proteins or chimeras (f) or from respective transgenic

Du145 cellular lysates (g). Percentage binding (as compared to the total input) is shown graphically.

Molecular weights (in kDa) for SDS-PAGE and western blots are as indicated. Error bars in the

graphs represent mean of three technical replicates ± SE. Statistical significance was calculated by

one-way Anova. Distinct alphabets represent significant difference at p.adj ≤ 0.05.

Fig. 6

a i.

Transgenic Du145 cell line	IC$_{50}$ (µM)		% change in IC$_{50}$
	HDM	HEME (5 µM)	
GFP	117.7 ± 0.9	114.7 ± 0.5	2.6 ± 0.7 %
K13-GFP	109.7 ± 0.2	93.8 ± 0.4	18.2 ± 0.6 %
KEAP1	87.8 ± 0.2	83.9 ± 0.8	6.1 ± 1.4 %
KEAP1^{N}K13^{KRP}-GFP	102.3 ± 0.3	84.8 ± 0.9	23.7 ± 1.4 %
K13^{N}KEAP1^{KRP}-GFP	80.8 ± 0.1	73.9 ± 0.9	13.5 ± 1.8 %

Fig. 6. Dynamics of K13, heme and Nrf2 in a redox survival mechanism that promotes ART-induced death. a. IC$_{50}$ values (µM, i) and change in percentage sensitivity to ART (right, ii) in respective transgenic Du145 cell lines on supplementation with 5 µM heme. **b.** Western blots (i) and quantitative densitometric analysis (versus GAPDH, ii) for K13-GFP (top) and Nrf2 (middle) in K13-

GFP^{HDM} (lanes 1 and 2) and K13-GFP^{HEME} (lanes 3 and 4) cells in the absence (lanes 1 and 3) or

presence (lanes 2 and 4) of ART. GAPDH (bottom) is the loading control. **c.** Western blots (i) and

graphical densitometric quantitation (ii) for cytoplasmic and nuclear fractionation of K13-GFP and Nrf2

in K13-GFP^{HDM} and K13-GFP^{HEME} cells in the presence of ART. Molecular weight (in kDa) are as

indicated. Error bars in a-c represent mean of three technical replicates ± SE. Statistical significance

was calculated by one-way Anova. Distinct alphabets represent significant difference at p.adj ≤ 0.05.

d. Model for a redox survival mechanism that promotes artemisinin-induced death**.** Dynamics of Nrf2

binding by free K13, its displacement by heme inducing (i) a redox survival response and (ii) K13-

heme mediated activation of ART inducing oxidative pro-death response in Du145 cells.

Fig. 7

Fig. 7. ART sensitivity is increased proportional to K13 intensity. a. IC_{50} values (µM) for GFP and K13-GFP cells in presence or absence of ATRA or heme (5 µM). **b.** Effect of heme (5 µM) and ATRA on percentage change in ART sensitivity in cells expressing GFP (grey bars) or K13-GFP (black bars). **c.** Graph showing the percentage change in ART sensitivity as a function of K13/GAPDH intensity ratio under HDM (solid black circles) or 5 µM heme-repleted (solid red circles) conditions in transgenic Du145 and HeLa cells. Black dotted trendline and R-squared values are indicated. Description of cells is shown at the bottom. Numbers only and numbers followed by initial 'H' indicate cells cultured under HDM or 5 µM heme-repleted conditions, respectively. **d.** Graph showing change in percentage sensitivity to ART at extracellular 5 µM (grey) and 50 µM (black) heme in the repleted

765   media. Error bars in a, b and d represent mean of three technical replicates ± SE. Statistical

766   significance was calculated by one-way Anova. Distinct alphabets represent significant difference at

767   p.adj ≤ 0.05.

Fig. 8

Fig. 8. A new model for clinical ART-R, implicated by K13-heme redox survival responses and ART-induced death. **a-b.** ART induced pro-death mechanisms and their modulation in rings. Our primary conclusion is for **K13 in cycle 1** (bold lines, **a**), where K13 (solid yellow circle) binds heme (solid red circles, **a**) and becomes stabilized at cytoplasmic face of endosomal Type I micropore membranes of merozoites and rings (shown in panel **b**). For merozoites and rings, heme may come from mitochondrial and extracellular sources. K13-heme activates ART (solid black stars) to heme-ART adducts (dark brown hybrid star-circles), which dissociate from K13 (resulting in suppression of K13 redox sensing). Widespread toxicity of heme-ART adducts to cellular targets, triggers a pro-death response in the parasite. Heme-free K13 may re-cycle back to resume heme binding and re-initiate cycle 1 and reinforce death outcomes. **K13 cycle 2** may be initiated if a (yet hypothetical) parasite redox factor (PRF; solid blue rectangle) binds K13 and becomes suppressed until heme-binding dislodges the PRF to activate ART and pro-death effects, acting additively with cycle 1. **K13 cycle 3** may be initiated upon K13 displacement of PRF to trigger pro-survival responses and reduce pro-death effects of cycles 1 and 2. **b i**. Model proposing K13-heme complexes (solid yellow circles with

solid red circles) assemble at Type I micropores of *P. falciparum* merozoites as they develop in schizonts. Released merozoites invade new red blood cells to form rings associated with Type I micropores and K13-heme complexes[71]. **b ii.** In wild type rings, K13-heme trigger ART (solid black stars)-induced oxidative damage radicals that destroy essential Type I micropores, resulting in ring-death. K13 ART-R mutants (solid purple circles) show reduced heme binding, oxidative damage enabling ring survival and ART-R. **c.** K13-heme is also catalytic for ART activation in trophozoites. K13 (solid yellow circles) concentrates at cytostome structures that enable hemoglobin digestion and liberation of free heme (solid red circles) in the food vacuole. Existing models propose that free heme activates ART (solid black stars) to toxic heme-ART adducts (dark brown hybrid star-circle) in the food vacuole (see red box). We propose that heme from the food vacuole and other sources (like the mitochondria) transported to the parasite bind K13 concentrated at the cytostome, which greatly stimulates formation of activated ART, and toxic heme-ART adducts (dark brown hybrid star-circle in bold red box) compared to encounters of ART with free heme in the food vacuole, thus explaining why K13 mutants do not confer ART-R at these stages.

References

1. Nosten, F. & White, N. J. Artemisinin-based combination treatment of falciparum malaria. *Am. J. Trop. Med. Hyg.* **77**, 181–192 (2007).

2. Eastman, R. T. & Fidock, D. A. Artemisinin-based combination therapies: A vital tool in efforts to eliminate malaria. *Nature Reviews Microbiology* vol. 7 (2009).

3. Pires, C. V. *et al.* Oxidative stress changes the effectiveness of artemisinin in Plasmodium falciparum. *MBio* **15**, (2024).

4. Zhu, P. & Zhou, B. The Antagonizing Role of Heme in the Antimalarial Function of Artemisinin: Elevating Intracellular Free Heme Negatively Impacts Artemisinin Activity in Plasmodium falciparum. *Molecules* **27**, (2022).

5. Rocamora, F. *et al.* Oxidative stress and protein damage responses mediate artemisinin resistance in malaria parasites. *PLoS Pathog* **14**, e1006930 (2018).

6.   Ariey, F. *et al.* A molecular marker of artemisinin-resistant Plasmodium falciparum malaria.

Nature **505**, 50–55 (2014).

7.   Straimer, J. *et al.* Drug resistance. K13-propeller mutations confer artemisinin resistance in

Plasmodium falciparum clinical isolates. *Science (80-.).* **347**, 428–431 (2015).

8.   WHO. World malaria report 2024. https://www.who.int/teams/global-malaria-

programme/reports/world-malaria-report-2024 (2024).

9.   Cui, L., Mharakurwa, S., Ndiaye, D., Rathod, P. K. & Rosenthal, P. J. Antimalarial drug

resistance: Literature review and activities and findings of the ICEMR network. *American*

Journal of Tropical Medicine and Hygiene vol. 93 (2015).

10.   Menard, D. *et al.* A Worldwide Map of Plasmodium falciparum K13-Propeller Polymorphisms.

N Engl J Med **374**, 2453–2464 (2016).

11.   Xie, S. C., Ralph, S. A. & Tilley, L. K13, the Cytostome, and Artemisinin Resistance. *Trends in*

Parasitology vol. 36 (2020).

12.   Blasco, B., Leroy, Di. & Fidock, D. A. Antimalarial drug resistance: Linking Plasmodium

falciparum parasite biology to the clinic. *Nature Medicine* vol. 23 (2017).

13.   Fairhurst, R. M. & Dondorp, A. M. Artemisinin-Resistant Plasmodium falciparum Malaria.

Microbiol Spectr **4**, (2016).

14.   Bhattacharjee, S. *et al.* Remodeling of the malaria parasite and host human red cell by vesicle

amplification that induces artemisinin resistance. *Blood* **131**, 1234–1247 (2018).

15.   Birnbaum, J. *et al.* A Kelch13-defined endocytosis pathway mediates artemisinin resistance in

malaria parasites. *Science (80-.).* **367**, 51–59 (2020).

16.   Tutor, M. V. *et al.* The Plasmodium falciparum artemisinin resistance-associated protein Kelch

13 is required for formation of normal cytostomes. *bioRxiv* (2023).

17.   Meshnick, S. R. Artemisinin: Mechanisms of action, resistance and toxicity. in *International*

Journal for Parasitology vol. 32 (2002).

18.   Sutherland, C. J., Henrici, R. C. & Artavanis-Tsakonas, K. Artemisinin susceptibility in the

838   malaria parasite Plasmodium falciparum: propellers, adaptor proteins and the need for cellular

839   healing. *FEMS Microbiol. Rev.* **45**, (2021).

840   19.   Ribbiso, K. A. *et al.* Artemisinin-based drugs target the plasmodium falciparum heme

841   detoxification pathway. *Antimicrob. Agents Chemother.* **65**, (2021).

842   20.   Zhang, S. & Gerhard, G. S. Heme activates artemisinin more efficiently than hemin, inorganic

843   iron, or hemoglobin. *Bioorganic Med. Chem.* **16**, (2008).

844   21.   Rosenthal, M. R. & Ng, C. L. Plasmodium falciparum Artemisinin Resistance: The Effect of

845   Heme, Protein Damage, and Parasite Cell Stress Response. *ACS Infectious Diseases* vol. 6

846   (2020).

847   22.   Yu, X. *et al.* Ring-stage growth arrest: Metabolic basis of artemisinin tolerance in Plasmodium

848   falciparum. *iScience* **26**, (2023).

849   23.   Platon, L. & Ménard, D. Plasmodium falciparum ring-stage plasticity and drug resistance.

850   *Trends in Parasitology* vol. 40 (2024).

851   24.   Rahman Abdur *et al.* Artemisinin-resistant Plasmodium falciparum Kelch13 mutant proteins

852   display reduced heme-binding affinity and decreased artemisinin activation. *Commun Biol.* **13**,

853   1499 (2024).

854   25.   Severance, S. & Hamza, I. Trafficking of heme and porphyrins in metazoa. *Chem. Rev.* **109**,

855   (2009).

856   26.   Schultz, I. J., Chen, C., Paw, B. H. & Hamza, I. Iron and porphyrin trafficking in heme

857   biogenesis. *Journal of Biological Chemistry* vol. 285 (2010).

858   27.   Ascenzi, P. *et al.* Hemoglobin and heme scavenging. *IUBMB Life* vol. 57 (2005).

859   28.   Morgan, W. T., Heng Liem, H., Sutor, R. P. & Muller-Eberhard, U. Transfer of heme from

860   heme-albumin to hemopexin. *BBA - Gen. Subj.* **444**, (1976).

861   29.   Yi, L. & Ragsdale, S. W. Evidence that the heme regulatory motifs in heme oxygenase-2 serve

862 as a thiol/disulfide redox switch regulating heme binding. *J. Biol. Chem.* **282**, (2007).

863 30. Gamiz-Arco, G. *et al.* Heme-binding enables allosteric modulation in an ancient TIM-barrel
864  glycosidase. *Nat. Commun.* **12**, (2021).

865 31. Chiabrando, D., Vinchi, F., Fiorito, V., Mercurio, S. & Tolosano, E. Heme in pathophysiology: A
866  matter of scavenging, metabolism and trafficking across cell membranes. *Front. Pharmacol.* **5**,
867  (2014).

868 32. Jiang Tempel W. Loppnau P. Graslund S. He H. Ravichandran M. Seitova A. Arrowsmith C.H.
869  Edwards A.M. Bountra C. El Bakkouri M. Senisterra G. Osman K.T. Lovato D.V. Hui R.
870  Hutchinson A. Lin Y.H., D. Q. Crystal Structure Analysis of Kelch protein from Plasmodium
871  falciparum. *Struct. Genomics Consort.* (2015).

872 33. Li, X., Zhang, D., Hannink, M. & Beamer, L. J. Crystal structure of the Kelch domain of human
873  Keap1. *J. Biol. Chem.* **279**, 54750–54758 (2004).

874 34. Billett, H. H. *Hemoglobin and Hematocrit. Clinical Methods: The History, Physical, and
875  Laboratory Examinations* (Butterworths, 1990).

876 35. Aich, A., Freundlich, M. & Vekilov, P. G. The free heme concentration in healthy human
877  erythrocytes. *Blood Cells, Mol. Dis.* **55**, (2015).

878 36. Donegan, R. K., Moore, C. M., Hanna, D. A. & Reddi, A. R. Handling heme: The mechanisms
879  underlying the movement of heme within and between cells. *Free Radical Biology and
880  Medicine* vol. 133 (2019).

881 37. White, N. J. Anaemia and malaria. *Malaria Journal* vol. 17 (2018).

882 38. Nagaraj, V. A. *et al.* Malaria Parasite-Synthesized Heme Is Essential in the Mosquito and Liver
883  Stages and Complements Host Heme in the Blood Stages of Infection. *PLoS Pathog.* **9**,
884  (2013).

885 39. Ke, H. *et al.* The heme biosynthesis pathway is essential for Plasmodium falciparum
886  development in mosquito stage but not in blood stages. *J. Biol. Chem.* **289**, (2014).

887   40.   Abshire, J. R., Rowlands, C. J., Ganesan, S. M., So, P. T. C. & Niles, J. C. Quantification of

888   labile heme in live malaria parasites using a genetically encoded biosensor. *Proc. Natl. Acad.*

889   *Sci. U. S. A.* **114**, (2017).

890   41.   Heller, L. E. & Roepe, P. D. Quantification of Free Ferriprotoporphyrin IX Heme and Hemozoin

891   for Artemisinin Sensitive versus Delayed Clearance Phenotype Plasmodium falciparum

892   Malarial Parasites. *Biochemistry* **57**, (2018).

893   42.   Stone, K. R., Mickey, D. D., Wunderli, H., Mickey, G. H. & Paulson, D. F. Isolation of a human

894   prostate carcinoma cell line (DU 145). *Int. J. Cancer* **21**, (1978).

895   43.   Jayakumar, S., Kunwar, A., Sandur, S. K., Pandey, B. N. & Chaubey, R. C. Differential

896   response of DU145 and PC3 prostate cancer cells to ionizing radiation: Role of reactive

897   oxygen species, GSH and Nrf2 in radiosensitivity. *Biochim. Biophys. Acta - Gen. Subj.* **1840**,

898   (2014).

899   44.   Zhang, P. *et al.* Loss of Kelch-like ECH-associated protein 1 function in prostate cancer cells

900   causes chemoresistance and radioresistance and promotes tumor growth. *Mol Cancer Ther* **9**,

901   336–346 (2010).

902   45.   Morgan, W. T. & Smith, A. Domain structure of rabbit hemopexin. Isolation and

903   characterization of a heme-binding glycopeptide. *J. Biol. Chem.* **259**, (1984).

904   46.   Tolosano, E. & Altruda, F. Hemopexin: Structure, function, and regulation. *DNA and Cell*

905   *Biology* vol. 21 (2002).

906   47.   Goel, N. *et al.* Plasmodium falciparum Kelch13 and its artemisinin-resistant mutants assemble

907   as hexamers in solution: a SAXS data-driven modelling study. *FEBS J.* **289**, (2022).

908   48.   Makarska-Bialokoz, M. Interactions of hemin with bovine serum albumin and human

909   hemoglobin: A fluorescence quenching study. *Spectrochim. Acta - Part A Mol. Biomol.*

910   *Spectrosc.* **193**, (2018).

911   49.   Kensler, T. W., Wakabayashi, N. & Biswal, S. Cell survival responses to environmental

912   stresses via the Keap1-Nrf2-ARE pathway. *Annual Review of Pharmacology and Toxicology*

913   vol. 47 (2007).

914  50. Mense, S. M. & Zhang, L. Heme: A versatile signaling molecule controlling the activities of

915   diverse regulators ranging from transcription factors to MAP kinases. *Cell Research* vol. 16

916   (2006).

917  51. Taguchi, K. *et al.* Keap1 degradation by autophagy for the maintenance of redox homeostasis.

918   *Proc. Natl. Acad. Sci. U. S. A.* **109**, (2012).

919  52. Kirchner, P. *et al.* Proteome-wide analysis of chaperone-mediated autophagy targeting motifs.

920   *PLoS Biol.* **17**, (2019).

921  53. Agarraberes, F. A. & Dice, J. F. A molecular chaperone complex at the lysosomal membrane

922   is required for protein translocation. *J. Cell Sci.* **114**, (2001).

923  54. Stricher, F., Macri, C., Ruff, M. & Muller, S. HSPA8/HSC70 chaperone protein. *Autophagy* **9**,

924   (2013).

925  55. Ikami, Y., Terasawa, K., Watabe, T., Yokoyama, S. & Hara-Yokoyama, M. The two-domain

926   architecture of LAMP2A within the lysosomal lumen regulates its interaction with

927   HSPA8/Hsc70. *Autophagy Reports* vol. 1 (2022).

928  56. Zhu, L. *et al.* Chaperone-mediated autophagy degrades Keap1 and promotes Nrf2-mediated

929   antioxidative response. *Aging Cell* **21**, (2022).

930  57. Anguiano, J. *et al.* Chemical modulation of chaperone-mediated autophagy by retinoic acid

931   derivatives. *Nat. Chem. Biol.* **9**, (2013).

932  58. Probst, B. L., McCauley, L., Trevino, I., Wigley, W. C. & Ferguson, D. A. Cancer cell growth is

933   differentially affected by constitutive activation of NRF2 by KEAP1 deletion and

934   pharmacological activation of NRF2 by the synthetic triterpenoid, RTA 405. *PLoS One* **10**,

935   (2015).

936  59. Gnadig, N. F. *et al.* Insights into the intracellular localization, protein associations and

937   artemisinin resistance properties of Plasmodium falciparum K13. *PLoS Pathog* **16**, e1008482

(2020).

60. Siddiqui, G., Srivastava, A., Russell, A. S. & Creek, D. J. Multi-omics Based Identification of

Specific Biochemical Changes Associated With PfKelch13-Mutant Artemisinin-Resistant

Plasmodium falciparum. *J Infect Dis* **215**, 1435–1444 (2017).

61. Atul *et al.* Artemisinin resistance in P. falciparum: Probing the interacting partners of Kelch13

protein in parasite. *J. Glob. Antimicrob. Resist.* (2023) doi:10.1016/j.jgar.2023.08.012.

62. Müller, S. Redox and antioxidant systems of the malaria parasite Plasmodium falciparum.

Molecular Microbiology vol. 53 (2004).

63. Hott, A. *et al.* Artemisinin-resistant Plasmodium falciparum parasites exhibit altered patterns of

development in infected erythrocytes. *Antimicrob Agents Chemother* **59**, 3156–3167 (2015).

64. Witkowski, B. *et al.* Novel phenotypic assays for the detection of artemisinin-resistant

Plasmodium falciparum malaria in Cambodia: in-vitro and ex-vivo drug-response studies.

Lancet Infect Dis **13**, 1043–1049 (2013).

65. Birnbaum, J. *et al.* A genetic system to study Plasmodium falciparum protein function. *Nat*

Methods **14**, 450–456 (2017).

66. Mbengue, A. *et al.* A molecular mechanism of artemisinin resistance in Plasmodium falciparum

malaria. *Nature* **520**, 683–687 (2015).

67. Wan, W. *et al.* The Toxoplasma micropore mediates endocytosis for selective nutrient salvage

from host cell compartments. *Nat. Commun.* **14**, (2023).

68. Koreny, L. *et al.* Stable endocytic structures navigate the complex pellicle of apicomplexan

parasites. *Nat. Commun.* **14**, (2023).

69. Garnham, P. C. C., Bird, R. G. & Baker, J. R. Electron microscope studies of motile stages of

malaria parasites: III. The ookinetes of Haemamoeba and Plasmodium. *Trans. R. Soc. Trop.*

Med. Hyg. **56**, 116–120 (1962).

70. Jiong Yang , Shaojun Long , Geoff Hide , Zhao-Rong Lun, D.-H. L. Apicomplexa micropore:

history, function, and formation. *Trends Parasitol.* **40**, 416–426 (2024).

71.   Haldar, K. & Bhattacharjee, S. Vesicular mechanisms of drug resistance in apicomplexan

parasites. *Microbiol. Mol. Biol. Rev.* **89**, 1–17 (2025).

72.   Wang, J. *et al.* Haem-activated promiscuous targeting of artemisinin in Plasmodium

falciparum. *Nat Commun* **6**, 10111 (2015).

73.   Gao, P. *et al.* Profiling the Antimalarial Mechanism of Artemisinin by Identifying Crucial Target

Proteins. *Engineering* (2023) doi:10.1016/j.eng.2023.06.001.

74.   Sinai, A. P. & Roepe, P. D. Autophagy in Apicomplexa: a life sustaining death mechanism?

Trends Parasitol **28**, 358–364 (2012).

75.   Ghosh, D., Walton, J. L., Roepe, P. D. & Sinai, A. P. Autophagy is a cell death mechanism in

Toxoplasma gondii. *Cell. Microbiol.* **14**, (2012).

76.   Suzuki, K., Bose, P., Leong-Quong, R. Y., Fujita, D. J. & Riabowol, K. REAP: A two minute cell

fractionation method. *BMC Res. Notes* **3**, (2010).