Home range is not constrained by numbers of hippocampal neurons

Mia C. Elbon, Greg Casalino and Suzana Herculano-Houzel

Dept. of Psychology, Vanderbilt Brain Institute, Vanderbilt University, Nashville TN

Correspondence to: suzana.herculano@vanderbilt.edu

Abstract

Numbers of hippocampal neurons vary by over three orders of magnitude across mammalian species. What evolutionary pressures shape this diversity? Given the role of the hippocampus in spatial mapping, the greater spatial navigation demands of larger home ranges may drive selection for more hippocampal neuron. Using data from 379 species, we crossed home range and population density data with cortical and hippocampal neuron counts predicted from clade-specific brain scaling laws to examine whether home range scales universally with estimated hippocampal neuron numbers across mammals. We confirm that home range scales universally with the inverse of population density across species and increases with body mass and metabolic rate. However, home range scaling with hippocampal or cortical neuron numbers differs by clade, such that bats, carnivorans and cetartiodactyls traverse home ranges over 1,000-fold larger than other mammals with equivalent hippocampal neuron numbers. These findings persist across data subsets controlling for study method, duration, and temporal scope. Numbers of hippocampal neurons are thus not limiting to spatial navigation in the wild, calling into question adaptationist explanations for the evolution of more hippocampal neurons based on a supposed need for increased spatial processing capacity. We propose that home range is determined primarily by population density, mediated by field metabolic rate and diet. The diversity in hippocampal neuron numbers across mammals, in turn, arises as a byproduct of clade-specific scaling of numbers of cortical neurons which we suggest is contingent on energetic opportunity, not on navigational or other cognitive demands.

Significance Statement

Why do some animals have so many more brain neurons than others? One explanation is evolution through selection for more neurons as needed for instance to navigate their environments to find food. However, this study finds that bats, carnivorans and whales traverse distances more than 1,000-fold larger than other mammals with similar numbers of neurons in the hippocampus, the brain structure required for spatial navigation. The authors question the traditional need-based explanation for brain diversity in favor of an opportunity-based account of the evolution of numbers of brain neurons.

Introduction

There is enormous diversity in numbers of cortical neurons across mammalian species. What are the drivers of that diversity, that is, what explains why some species have very few and others have up to a thousand times as many cortical neurons? Ever since Darwin, need-based natural selection is as a rule invoked as the main driver of diversity in evolution. By that logic, and assuming that more neurons increase the information-processing capacity of brain structures, different parts of the brain with different functions should be found to have more neurons as species rely more heavily on those functions.

In what concerns the hippocampus, a cortical structure responsible for the associative learning underlying concept building and spatial navigation[1-6], those species with increasing needs for spatial navigation should be under selective pressure for increasing numbers of hippocampal neurons. Across mammalian species, the need for spatial navigation capacity should increase with home range, which is the area that an animal traverses over a set period of time in its normal activities of food gathering, mating, and caring for young[7]. Mammals create spatial maps using their hippocampus[1,8,9]It has been suggested that hippocampus size varies with relative selection pressures on cognitive mapping abilities and spatial memory[10-12]. Larger animals have increasingly larger home ranges[13-20], which is considered to be a response to the larger metabolic needs of species with increasing body mass[7,13,14,18,19,20-24]. Increased metabolic needs would also explain the larger home ranges of carnivores compared to herbivores of similar body mass[13,19,24,25]. Accordingly, larger animals with larger home ranges should be under selective pressure for increased hippocampal volume. Indeed, animals who rely more on complex spatial behaviors (e.g., animals with larger home ranges) have larger hippocampi[10,26-30].

Similarly, increasing the spatial complexity of the environment has been shown to increase

hippocampal volume[31-35].

The underlying rationale behind the argument that animals need larger hippocampi to

navigate larger home ranges is that larger hippocampi should consist of proportionately more

neurons; absent direct data on numbers of hippocampal neurons, hippocampal volume has been

expected to serve as a proxy for the number of neurons in the hippocampal network serving

spatial navigation. So far, however, very little data exist on actual numbers of neurons in the

hippocampus. Models shows that increasing numbers of neurons in the hippocampal network

should afford a linear increase in the capacity of this network to represent the area of the

environment that an individual traverses with a given spatial resolution[9,36-38]. If covering the

increased surface areas of larger home ranges requires increased numbers of hippocampal

neurons to the point that it imposes a selective pressure, then one should find a universal

correlation of both hippocampal size and numbers of hippocampal neurons with home range

across a wide variety of mammalian species.

We report in an accompanying paper that the total number of neurons in the mammalian

hippocampus measured with the isotropic fractionator, a non-stereological method that provides

estimates that are completely independent of structure mass, scales universally across species not

with hippocampal or brain mass, but with the total number of neurons in the cerebral cortex[39].

Importantly, our finding of clade-specific scaling of numbers of hippocampal neurons with brain

mass contradicts stereological estimates in a recent report that simply reflect brain size across a

wide range of mammalian species[40], which could be due to undersampling that makes

stereological estimates inadvertently sensitive to structure size[41]. Our new finding that numbers

of hippocampal neurons scale approximately with the square root of the total number of neurons

in the cerebral cortex, both of which can be estimated from brain mass in a clade-specific

manner, puts us in position to finally examine directly whether home range scales universally

with the estimated number of hippocampal neurons across species covering the entire spectrum

of mammalian body sizes. Here we use the home range dataset compiled by Broekman et al.[42] to

which we add estimates of population density for those same species compiled by Santini et al.[43];

data on body mass and basal metabolic rate compiled in the AnAge database by de Magalhães

and Costa[44]; and data on field metabolic rates from three different sources[45-47].We then cross

these data both with numbers of neurons in the cerebral cortex predicted from brain mass in each

species according to the clade-specific relationships that have been found to apply separately to

primates, glires, certartiodactyls, afrotherians, marsupials, chiropterans, and carnivorans[48]; and

with the numbers of neurons in the hippocampus of each species predicted from their number of

cerebral cortical neurons using the scaling relationships that apply to their clade[39]. The latter was

used as an added check because the universal scaling of numbers of hippocampal neurons with

the square root of total numbers of cerebral cortical neurons[39] means that any scaling

relationships that apply to total numbers of cerebral cortical neurons also apply to hippocampal

neurons with the exponent divided by 2. We could then test whether the expected universal

correlation between home range and numbers of hippocampal neurons applied across

mammalian species.

Results

Across 350 mammalian species in our dataset, we find that average home range scales as power functions of adult body mass (Fig. 1A) and basal metabolic rate (Fig. 1B) with exponents 1.06 ± 0.04 and 1.36 ± 0.09, respectively. The residual variation in $\log_{10}$ average home range after accounting for $\log_{10}$ body mass or $\log_{10}$ basal metabolic rate shows that bats, cetaceans and most carnivorans, especially the semi-aquatic species and some bears (asterisks in Fig. 1), consistently have much larger home ranges than predicted for their body mass (Fig. 1C) and basal metabolic rate (Fig. 1D; ANOVA $p<0.0001$). However, the apparent effect of clade is due to diet type (Fig. 1E, F, both ANOVA $p<0.0001$), as carnivorous species have consistently larger home ranges than herbivorous and omnivorous species. Within each diet type, average home range scales similarly across clades (all ANOVA $p>0.01$, with the exclusion of bats, which are always outliers as shown in Fig .1A). Importantly, across herbivorous and omnivorous species (not including bats), average home range scales across clades with an exponent of 1.006 ± 0.044 ($r^2=0.682$, $p<0.0001$), identical to unity; in contrast, across carnivorous species (not including bats), average home range scales across species of all clades with a larger scaling exponent of 1.308 ± 0.056 ($r^2=0.870$, $p<0.0001$).

Figure 1. Average home range scales universally with body mass and basal metabolic rate across a wide range of mammalian species, with the exception of bats, semi-aquatic carnivorans and some bears. All plotted functions include all species and clades in the dataset. Allometric exponents and r^2 values are shown in the graphs. **A**, average home range scales with adult body mass ($p < 0.0001$, $n = 350$) in a manner that leaves bats, semi-aquatic carnivorans and some bear species (asterisks) outside the 95% prediction interval (dotted lines). **B**, average home range scales with basal metabolic rate ($p < 0.0001$, $n = 157$) in an apparently universal fashion, although bats remain outliers; notice that data for aquatic carnivores, several bats and some bears are not available in the dataset. **C,E**, residual $\log_{10}$ home range for each species after accounting for $\log_{10}$ adult body mass, grouped by clade (**C**) or by diet type (**E**). **D,F**, residual $\log_{10}$ home range for each species after accounting for $\log_{10}$ basal metabolic rate, grouped by clade (**D**) or by diet type (**F**). Diet types are carnivory (C), herbivory (H) or omnivory (O).

In contrast, we find that the scaling relationships between home range and numbers of

neurons in the cerebral cortex (Fig. 2A) and hippocampus (Fig. 2B) do not apply universally

across all mammals or clades. Using the functions fitted across all species, primates consistently

have smaller home ranges and cetaceans and chiropterans as well as semi-aquatic carnivorans

consistently have larger home ranges than expected for their numbers of neurons, as the residuals

show (Fig. 2C, D; ANOVA for all clades, p<0.0001; ANOVA for carnivorans, chiropterans, and

primates, p<0.0001). Importantly, the distinct scaling of home range with numbers of cortical

and hippocampal neurons in primates is not due to diet type (Fig. 2E, F), because primate home

ranges remain significantly different (smaller than expected) after accounting for diet type

(ANOVA for carnivorans, glires and primates, p<0.0001 for each diet type). Semi-aquatic

carnivores (all of which are carnivoran species) and bats again are the species with the largest

residual home range, now joined by several cetacean species (Fig. 2A, B). The systematic

disparities between bats, carnivorans, cetaceans and primates suggest that the scaling of home

range with numbers of cortical and hippocampal neurons is clade-specific.

Figure 2: Home range does not scale universally with numbers of cortical or hippocampal neurons across
mammalian species. All plotted functions include all species and clades in the dataset. Allometric
exponents and r^2 values are shown in the graphs. A, B, average home range scales with estimated total
numbers of cerebral cortical neurons (A; p < 0.0001, n = 303) or of hippocampal neurons (B; p < 0.0001,
n = 292) in a manner that leaves semi-aquatic carnivorans and some bear species (asterisks) as well as
bats outside the 95% prediction interval (dotted lines), and additionally leaves all primate species below
the fitted line, although mostly still inside the prediction interval. C, residual $\log_{10}$ home range for each
species after accounting for $\log_{10}$ number of cerebral cortical neurons, grouped by clade. D, residual $\log_{10}$
home range for each species after accounting for $\log_{10}$ number of hippocampal, grouped by clade. Note
that primate species consistently have smaller home ranges and carnivorans, especially the semi-aquatic
species, consistently have larger home ranges than predicted for their numbers of neurons.

The clade-specificity of the scaling of home range with numbers of cortical and hippocampal

neurons is evidenced in Figure 3. Glires, eulipotyphlans, marsupials, afrotherians and primates

share scaling exponents below 3.0 or 6.0 (for numbers of cortical or hippocampal neurons,

respectively), whereas carnivorans and artiodactyls (including cetaceans) share exponents above

those values (exponents indicated in Fig. 3A, B; the scaling function for chiropterans is not

significant, with p=0.7757). Interestingly, while bats are always outliers with much larger home

ranges for their numbers of neurons, as with body mass (Fig. 1), the data for the remaining clades

can be grouped in two different ways that make either primates or large carnivorans and

cetartiodactyls into outliers. When carnivorans and cetartiodactyls (in addition to bats) are

excluded from the analysis, yielding joint power functions with an r^2 of 0.641, primates are well

aligned with glires, marsupials and eulipotyphlans in the scaling of home range with numbers of

neurons (Fig. 3A, B). In this case, all large carnivorans (not only the semi-aquatic species) and

cetartiodactyls would be described as having far larger home ranges than predicted for their

numbers of cortical and hippocampal neurons. Alternatively, if only primates (and bats) are

excluded while carnivorans and cetartiodactyls are included in the analysis together with all

other clades (Fig. 3C, D), we find that average home range scales with numbers of all cortical or

hippocampal neurons as power functions of a higher r^2 of 0.788 and 0.655 that however still

exclude the aquatic carnivorans and some bears in the dataset, as well as bats. Using this latter

analysis, which provides a better description of the variation in the dataset, primates and bats are

the outlier clades, and in different directions: primates consistently occupy much smaller home

ranges, and most bats consistently occupy much larger home ranges, than other mammals of

similar numbers of cortical or hippocampal neurons. Either way, two facts remain: that home

range and numbers of cortical and hippocampal neurons do not vary in a universally correlated

manner across mammalian species; and that semi-aquatic carnivorans, many bats and some bears

occupy much larger home ranges than can be predicted by their body size or numbers of neurons.

Figure 3. Home range scales with numbers of cortical or hippocampal neurons in a clade-specific
manner. Scaling exponents ± standard error that apply to each clade (not plotted) are indicated in the
graphs. Plotted functions include all species and clades in the dataset (excluding chiropterans) minus
carnivorans and cetartiodactyls (A and B) or minus primates (C and D). Allometric exponents and r^2
values for the plotted functions are shown in black in the top left of each graph. **A and B:** When
carnivorans and cetartiodactyls are excluded from analysis, average home range scales with number of
cortical neurons and number of hippocampal neurons ($p < 0.0001$, $n = 216$) in a manner that includes
primates and leaves most carnivorans, most cetaceans and some artiodactyls outside the 95% prediction
interval (dotted lines). **C and D**: When primates are excluded from analysis, average home range scales as
power functions of numbers of cortical and hippocampal neurons ($p < 0.0001$, $n = 216$) with a higher r^2
and a 95% prediction interval that either excludes primates (C) or leaves them systematically below the
prediction line (D) and still excludes the semi-aquatic carnivorans and some bears in the dataset. Notice
that most bats in the dataset are always outliers.

Effect of tracking methods, sample size, and study duration

The average home range estimates used so far in the analysis combine various tracking methods,
sample size, and study duration, as has been standard practice in the field[19,42,50,51]. The lack of

consistence in the time span (daily, annual, other) and level (individual, group, population) that apply to the values of home range analyzed raises the concern that the finding that primates are outliers in the scaling of home range with numbers of neurons is a measurement artifact. To address this possibility, we determined whether primates are still outliers when the scaling between numbers of cortical or hippocampal neurons and home range is calculated exclusively for subsets of data restricted to the same combination of time span (home range covered in a day or in a year) and level (home range of single individuals or of a population). Figure 4 shows that even in these much more restricted datasets (which lack data for bats, semi-aquatic carnivorans and cetaaceans), and despite the inclusion of primates in the fitted data, species in this clade still consistently have smaller home ranges than predicted for their numbers of cortical or hippocampal neurons, well below the 95% prediction interval.

Figure 4. Primates have smaller home ranges than predicted for their numbers of cortical and
hippocampal neurons even in data sets restricted to similar time span and population level. Average home
range is plotted as a function of numbers of cortical neurons (A-D) or estimated numbers of hippocampal
neurons (E-H) using data restricted to daily home range measured in individuals (A, E); annual home
range measured in individuals (B, F); daily home range measured in a population (C, G); or annual home
range measured in a population (D, H). The plotted functions apply to all species in each data set,
including primates, which still fall below the 95% prediction interval (dotted lines) in every case.
Allometric exponents (all $p<0.0001$), r^2 values and n are shown in the graphs.
221

Population density is a universal predictor of home range, and also does not scale universally with numbers of cortical neurons

We find that average home range scales with the near inverse of population density, with an

exponent indistinguishable from -1 (Fig. 5A), in a manner that applies equally across carnivores,

herbivores and omnivores (Fig. 5B, ANOVA $p=0.3789$) and, importantly, also includes bats

(Fig. 5C). Our dataset thus reproduces the previous observations that home range is a negative

inverse function of population density[15,18,52-54]. While an ANOVA across all clades (Fig. 5C)

indicates a significant effect of clade ($p<0.0001$), residual home ranges after accounting for

population density do not differ across the three clades that share large numbers of hippocampal

neurons but differ markedly in average home range (ANOVA for carnivorans, cetartiodactyls,

and primates, $p=0.1429$), nor across bats and rodents that share small numbers of hippocampal

neurons but also differ strikingly in home range (ANOVA, $p=0.3347$). Interestingly, while the

scaling of home range with the inverse of population density includes bats, primates and

cetaceans, most semi-aquatic carnivore species remain outliers, with larger home ranges than

predicted for their population density (Fig. 5A).

A

B

C

238 **Figure 5.** Population density is a better universal predictor of home range than any other variable
239 examined. A, home range scales as a power function of population density (A, p < 0.0001, n = 262). B,
240 residual $\log_{10}$ home range by population density for each species grouped by diet. C residual $\log_{10}$ home
241 range by population density for each species grouped by clade.

In line with the lack of a universal correlation between home range and numbers of hippocampal

neurons, we find that the power functions that relate population density to numbers of cortical

(Fig. 6A) or hippocampal neurons (Fig. 6B) are not universal across all mammals in the dataset

(both ANOVA p<0.0001), and also do not apply equally across species of different diet types

(Fig. 6C,D, ANOVA p<0.0001). Across clades primates and glires have systematically higher

and carnivorans, cetaceans and the one bat species with estimates available have systematically

lower population densities than predicted for their numbers of neurons (Fig. 6E,F), even when

analysis is restricted to herbivorous and omnivorous diet types (ANOVA, p<0.0001), which

argues for clade-specific relationships between population density and numbers of hippocampal

and cortical neurons.

Figure 6. Population density is not a universal predictor of number of hippocampal neurons. Population density scales as a power function of number of cerebral cortical (A; p<0.0001. n=XXX) or hippocampal neurons (B; p < 0.0001, n = 216) across species in a manner that includes primates but still leaves cetaceans and one semi-aquatic carnivoran (asterisks) below the 95% prediction interval (dotted lines). C, D, residual $\log_{10}$ population density after accounting for number of cerebral cortical (C) or hippocampal (D) neurons for each species, grouped by diet type. E, F, residual $\log_{10}$ population density after accounting for numbers of cortical (E) or hippocampal (F) neurons for each species, grouped by clade. Note that there is a single species of chiropterans with data available for both population density and estimated numbers of neurons.

Finally, we sought to determine the source of the larger home ranges and lower population densities of carnivorans and carnivorous species as a whole compared to primates of similar numbers of hippocampal neurons by examining how these three variables scale with the field metabolic rate that indicates how much energy different species must be able to procure per day in order to survive. Figure 7 shows that field metabolic rate is a good but not universal predictor of home range (Fig. 7A), population density (Fig. 7B) and number of hippocampal neurons (Fig. 7C). Carnivorous species (Fig. 7D) and also bats and carnivoran species (Fig. 7G, the vast majority of which also have carnivorous diets) consistently have larger home ranges, whereas primates and glires have consistently smaller home ranges (Fig. 7G), than predicted for their field metabolic rate. Carnivorous species, which include carnivorans and cetaceans and the one available bat species, also have lower than predicted population densities, while primates have higher than predicted population densities, for their field metabolic rates (Figs. 7E, H). Diet, on the other hand, does not differentiate the relationship between numbers of hippocampal neurons and field metabolic rate (Fig. 7I; ANOVA, p=0.0343 versus p<0.001 for all other comparisons); the distinguishing factor here is clade, as primates have systematically more hippocampal neurons than predicted for their field metabolic rates (Fig. 7F), in line with their systematically larger numbers of cortical neurons compared to non-primate mammals of similar body mass[50], and the two bats with data available have fewer hippocampal neurons than predicted for their field metabolic rates. We thus find that while the home range occupied by mammalian species is a universal function of their population density, with both variables affected coordinately by diet,

299 the number of neurons in the hippocampus of mammalian species is not universally related to the

300 home range, population density, or field metabolic needs of a species.

Figure 7. Field metabolic rate mediates population density. Field metabolic rate scales as a power
function of average home range (A), population density (B), and estimated number of hippocampal
neurons (C) across mammalian species. All plotted functions include all species and clades in the dataset.
Allometric exponents and r^2 values are shown in the graphs. A-C, average home range scales with field
metabolic rate (A; p < 0.0001, n = 67), population density (B; p < 0.0001, n = 50), and number of
hippocampal neurons (C; p < 0.0001, n = 60) in a manner that includes semi-aquatic carnivorans and
some bear species (asterisks) and primates inside the 95% prediction interval (dotted lines). D-I, residual
$\log_{10}$ home range, population density, and number of hippocampal neurons after accounting for field
metabolic rate for each species, grouped by clade (D-F) or by diet (G-I). Note that while primates are
included in the 95% prediction interval, they deviate systematically from the predicted values.

Discussion

Obtaining sufficient energy to make it repeatedly through the days, together with

maintaining physical integrity, is the one non-negotiable condition for animal survival. Our

findings are consistent with a scenario in which, given an environmental reality of limited food

availability per surface area, the higher rates of energy use per day of larger animals cause them

to self-organize, through food-seeking exploitative behaviors, into predictably lower population

densities, which in turn necessitate exploratory behaviors and dispersal into inversely larger

home ranges[15]. Importantly, our finding that home range scales linearly with body mass across

herbivorous and omnivorous species but with a larger exponent of 1.3 across carnivorous species

indicates that the inconsistent scaling exponents reported in the literature were due to the

inclusion of varying numbers of carnivorous species ([14,17-19,22,25,49]). We propose that diet type

affects feeding efficacy and efficiency and thus modulates the relationship between field

metabolic rate and population density (Fig. 7), which, given the necessarily inverse relationship

between population density and home range[20], explains the dependence on diet of the scaling of

home range with body mass[19]. Specifically, carnivory exposes animals to the relative scarcity of

prey compared to vegetable foodstuffs per surface area in the environment[18,19,54]. As a result, for

a given body size and field metabolic rate, carnivorous animals exist in lower population

densities and explore larger home ranges than omnivorous animals, and herbivorous species exist

in larger population densities and explores smaller home ranges, all the while home range and

population density continue to scale universally across all diet types.

While we find that microchiropterans have larger home ranges than expected for their

body mass, micro- and megachiropteran bats alike have the home ranges predicted for their

population densities, which is explained by unusually low population densities for the body mass

found in microchiropterans. Low population densities could be expected for insect-eating bats

due to food scarcity combined with their very high specific metabolic rates, even if two of these

bat species with very large home ranges in the dataset also consume fruits (*Carollia perspicillata*

and *Glossophaga soricina*). Similarly, we propose that the relative scarcity of prey combined

with the increased metabolic cost of living in cold water and the three-dimensional nature of

hunting in an aquatic environment explain the finding that semi-aquatic carnivorans and

cetaceans are consistently outliers in our dataset, exploring home ranges that are larger than

expected not only for their body mass and field metabolic rate but also population density. The

resting metabolic costs of aquatic mammals are about twice as high as those of terrestrial

mammals, with a scaling relationship for resting metabolic rate with body mass of 26.9 $M_{bd}^{0.69}$ in

aquatic mammals[56] in contrast to McNab's allometric prediction for the basal metabolic rate of

similarly sized terrestrial mammals[57] of 10.55 $M_{bd}^{0.724}$ (after converting for M_{bd} in kg), probably

due to heat lost to cold water. The effect of prey scarcity to drive decreased population density

and larger home range is expected to be especially severe in both aerial and aquatic

environments, where there is an added dimension to prey movement. Interestingly, given the

difference between stroke vs. stride energetic costs, fully submerged mammals are able to avoid

hydrodynamic drag and exhibit lower mean stroke costs compared to surface swimming

carnivores[58]. The higher efficiency of fully submerged carnivores, such as cetaceans, might

explain how come their home ranges match the predicted values for their very low population

densities whereas the aquatic carnivorans, as inefficient surface swimmers, explore much larger

home ranges than predicted for their population densities (Fig. 5A).

To the heart of our central question, we find that the enormous range of coordinated

variation in home range and population density occurs without the expected universal correlation

with numbers of hippocampal neurons typically presumed to be necessary to support spatial

exploration[10,26-35]. To the contrary, we show that carnivoran species are capable of exploring

home ranges of around 1,000 km^2 on land and as much as 100,000 km^2 over water with the same

estimated ca. 30 million hippocampal neurons found in primates who remain in an area of just

about 1 km^2; likewise, bats explore areas between 1 and 10 km^2 with fewer than the 2-3 million

hippocampal neurons found in small rodents and marsupials who forage in under 0.01 km^2 (Fig.

2B). Conversely, bats, carnivorans and primates that explore home ranges of about 10 km^2 do so

with numbers of hippocampal neurons that are 10 times as large in primates as in carnivorans

and ca. 30 times as large as in bats. If bats, carnivorans and primates alike obtain the energy that

they need by occupying the home range that is predicted for their population density (Fig. 5A),

the number of hippocampal neurons available to represent the surface area covered by a species

is clearly not limiting to the point that it would be under selective pressure to increase together

with home range as animals became larger in evolution. Known mechanisms of multiplexing
hippocampal representations of environments on different scales[38] and dynamically scaling place
field sizes to context[59] would suffice to account for how a given number of hippocampal neurons
suffices to allow exploration of home ranges varying by a factor of over 1,000-fold that we
uncover here.

The finding that bats cover inordinately large home ranges for their diminutive body
mass using the smallest numbers of hippocampal neurons in our dataset should suffice to dismiss
the notion that a need to navigate large environments imposes a selective pressure for more
neurons, or even larger brains for that matter. Those determined to find adaptationist
explanations to the discrepancy in scaling of home range with numbers of hippocampal neurons
between primates and other mammals might turn to Fig. 3D and argue that non-primate species
with larger home ranges do have the correspondingly larger numbers of hippocampal neurons
predicted by a supposed need for more hippocampal spatial representational power. In that case,
however, a new argument would have to be invoked to explain how come primates evolved
much larger numbers of hippocampal neurons than "necessary" to represent their comparatively
small environments – for instance, that primates benefit from the increased complexity in
representing their environment afforded by more neurons (see below). In our view, a much more
parsimonious explanation lies in acknowledging that the scaling relationship between numbers of
neurons and body size is not universal[55,60]. If there is any correlation between home range and
numbers of hippocampal (and cerebral cortical) neurons, we argue that it is simply because
larger body mass, which necessitates universally larger home ranges as predicted by their
population densities (according to the combination of field metabolic rate and diet), come with
more cortical neurons[61] – but following scaling rules that differ between primate and non-
primate clades[55]. In this case, the clade-specific correlation between home range and numbers of
hippocampal neurons is a trivial mathematical contingency, not a biological necessity.

This is not to say that larger numbers of hippocampal neurons do not bring a cognitive
advantage. According to theory, the more the neurons in the hippocampal network, the bigger
should be the capacity of the network to represent an area with a certain spatial resolution[38],
which can be presumed to extend to cognitive representations. Given the combination of the
predictably small home ranges of primate species and their large numbers of hippocampal
neurons (which are also predictable for their cerebral cortices[39]), we propose that primates do

benefit from both increased complexity in cognitive representations and increased spatial

resolution in their hippocampal maps of the environment, as the radius of their place cells is

expected to be reduced[38]. This latter prediction can be tested with systematic comparisons of

place cell radius across primate and non-primate species of either similar home ranges or

numbers of hippocampal neurons. Evidently, the comparatively lower spatial resolution of non-

primates and presumably simpler cognitive representations are perfectly sufficient for their

successful navigation in exploring and exploiting whatever size home range is required for their

survival. Thus, the increased spatial resolution we predict for primates cannot be considered a

necessity, and therefore also not the result of selective pressures – unless one resorts to a

convoluted argument that only primates were subject to such pressures for increased spatial

resolution.

Rather, we propose that it is more parsimonious that in exploring their environment,

mammals go as far as they have to go in order to meet their metabolic demands (given how their

diet impacts population density), but their numbers of hippocampal neurons are just not limiting

to how far they can go. The diversity of numbers of hippocampal neurons, which cannot be said

to have been driven by selective pressure for increased spatial navigation capacity, is tied to the

square root of the variation in total numbers of cortical neurons[39]. Thus, what needs to be

explained is what leads to the enormous variation in total numbers of cortical neurons in

mammalian evolution. We favor the simpler alternative explanation that numbers of cortical (and

so also hippocampal) neurons increase in evolution as consistently increased energetic

opportunities present themselves to a population across generations[62], with body size increasing

in tandem as still affordable[61].

Materials and Methods

The home range and population densities of 342 mammalian species were compiled using

two databases: "HomeRange: A global database of mammalian home ranges"[42] and

"TetraDENSITY 2.0 – A database of population density estimates in Tetrapods"[43]. Home range

data collection and statistical techniques for each species differed across studies, with often

differing temporal measurements and differing observation techniques employed for any given

species. To combine home range and population density data into a single database, it was

necessary to calculate average home range values for each species. The combined database was

then integrated with data on body mass, basal metabolic rate and diet type compiled by de

Magalhães and Costa[44] in the AnAge database and with data on numbers of cortical neurons

predicted from brain mass[48]. From the numbers of cortical neurons, the number of hippocampal

neurons was estimated for each species according to clade-specific formulas: $e^{7.673} N_{cx}^{0.815}$ for

cetartiodactyls, $e^{3.181} N_{cx}^{0.658}$ for carnivorans, $e^{-4.792} N_{cx}^{1.183}$ for chiropterans, $e^{4.322} N_{cx}^{0.640}$ for

marsupials, $e^{9.841} N_{cx}^{0.346}$ for primates, $e^{3.384} N_{cx}^{0.693}$ for glires, and $e^{7.767} N_{cx}^{0.450}$ for afrotherians.

Note that these clade-specific calculations provide an alternate set of estimates for N_{hp}, which

can also be more simply obtained as roughly $N_{cx}^{0.5}$ [39]. Finally, field metabolic rate data were

added to the combined database using values gathered from Nagy[45], de Castro[46] and Pontzer[47].

From these data, average home range (measured in km^2) for each species was plotted in

log-transformed bivariate-fit graphs comparing home range against adult body mass (measured

in grams), basal metabolic rate (in W), field metabolic rate (in kJ/day), number of cortical

neurons (N_{cx}), estimated number of hippocampal neurons (N_{hp}), and population density

(individuals/km^2) in JMP Pro 18. Mammalian data points were separated by clade for

comparison, as indicated in the graphs. We purposefully do not use any phylogenetic correction

methods because clade effects are addressed directly in our analysis.

Acknowledgments

Thanks to Vanderbilt University for supporting Mia C. Elbon. This work was not supported by
any external sources of funding.

References

1. J. O'Keefe, J. Dostrovsky, The hippocampus as a spatial map: Preliminary evidence from unit activity in the freely-moving rat. Brain Res. 34, 171–175 (1971).

2. J. O'Keefe, L. Nadel. "The Hippocampus as a Cognitive Map." Oxford University Press (1978).

3.   M. A. Wilson, B. L. McNaughton, Dynamics of the hippocampal ensemble code for space. Science 261, 1055–1058 (1993). https://doi.org/10.1126/science.8351520

4.   S.M. Weisberg, V.R. Schinazi, N.S. Newcombe, T.F. Shipley, R.A. Epstein. Variations in cognitive maps: Understanding individual differences in navigation. Journal of Experimental Psychology: Learning Memory and Cognition, 40(3), 669-682 (2014). https://doi.org/10.1037/a0035261.

5.   M. Geva Sagiv, L. Las, Y. Yovel, N. Ulanovsky, Spatial cognition in bats and rats: from sensory acquisition to multiscale maps and navigation. Nat. Rev. Neurosci. 16, 94–108 (2015). https://doi.org/10.1038/nrn3888

6. S.M. Weisberg, A.D. Ekstrom. Hippocampal volume and navigational ability: The map(ping) is not to scale. Neuroscience and biobehavioral reviews, 126, 102–112 (20211). https://doi.org/10.1016/j.neubiorev.2021.03.012.

7. W.H. Burt. Territoriality and Home Range Concepts as Applied to Mammals, Journal of Mammalogy, Volume 24, Issue 3,, Pages 346–352 (1943), https://doi.org/10.2307/1374834

8.   M. Fyhn, S. Molden, M. P. Witter, E. I. Moser, M. B. Moser, Spatial representation in the entorhinal cortex. Science 305, 1258–1264 (2004).

9.   K. B. Kjelstrup, et al., Finite scale of spatial representation in the hippocampus. Science 321, 140–143 (2008).

10. J. R. Krebs, D. F. Sherry, S. D. Healy, V. H. Perry, A. L. Vaccarino, Hippocampal specialization of food-storing birds. Proc. Natl. Acad. Sci. U.S.A. 86, 1388–1392 (1989).

11. N. S. Clayton, J. C. Reboreda, A. Kacelnik, Seasonal changes of hippocampus volume in parasitic cowbirds. Behav. Process. 41, 237–243 (1997).

12. R. A. Powell, M. S. Mitchell, What is a home range? J. Mammal. 93, 948–958 (2012).

13. B. K. McNab, Bioenergetics and the determination of home range size. Am. Nat. 97, 133–140 (1963).

14. A. S. Harestad, F. L. Bunnel, Home range and body weight—A reevaluation. Ecology 60, 389–402 (1979). https://doi.org/10.2307/1937667

15. J. Damuth, Population density and body size in mammals. Nature 290, 699–700 (1981). https://doi.org/10.1038/290699a0

16. P.H. Harvey, G.M. Mace. Comparisons between taxa and adaptive trends: problems of methodology. Current Problems in Sociobiology, p. 343-361. University Press, Cambridge (1982).

17. S. L. Lindstedt, et al., Home range, time, and body size in mammals. Ecology 67, 413–418 (1986). https://doi.org/10.2307/1938584

18. W. Jetz, et al., The scaling of animal space use. Science 306, 266–268 (2004). https://doi.org/10.1126/science.1102138

19. M. A. Tucker, T. J. Ord, T. L. Rogers, Home range-body mass patterns: are all mammals equal? Glob. Ecol. Biogeogr.23, 1105–1114 (2014). https://doi.org/10.1111/geb.12194

20. J. Damuth, Interspecific allometry of population density in mammals and other animals: the independence of body mass and population energy use. Biol. J. Linn. Soc. 31, 193–246 (1987). https://doi.org/10.1111/j.1095-8312.1987.tb01990.x

21. M. E. Gompper, J. L. Gittleman, Home range scaling: intraspecific and comparative trends. Oecologia 87, 343–348 (1991). https://doi.org/10.1007/BF00634589

22. D. A. Kelt, D. H. Van Vuren, The ecology and macroecology of mammalian home range area. Am. Nat. 157, 637–645 (2001). https://doi.org/10.1086/320621

23. K. A. Nagy, Field metabolic rate and body size. J. Exp. Biol. 208, 1621–1625 (2005). https://doi.org/10.1242/jeb.01553

24. Z. Boratyński, Energetic constraints on mammalian home range size. Funct. Ecol. 34, 468–474 (2020). https://doi.org/10.1111/1365-2435.13480

25. T. W. Schoener, Sizes of feeding territories among birds. Ecology 49, 123–141 (1968). https://doi.org/10.2307/1933567

26. D. F. Sherry, A. L. Vaccarino, Hippocampus and memory for food caches in black capped chickadees. Behav. Neurosci.103, 308–318 (1989). https://doi.org/10.1037/0735-7044.103.2.308

27. L. F. Jacobs, W. D. Spencer, Natural space use patterns and hippocampal size in kangaroo rats. Brain Behav. Evol. 44, 125–132 (1994). https://doi.org/10.1159/000113584

28. S. D. Healy, J. R. Krebs, Food storing and the hippocampus in Paridae. Brain Behav. Evol. 47, 195–199 (1996).

29. O. S. Todorov, V. Weisbecker, E. Gilissen, K. Zilles, A. A. de Sousa, Primate hippocampus size and organization are predicted by sociality but not diet. Proc. R. Soc. B 286, 20191712 (2019). https://doi.org/10.1098/rspb.2019.1712

30. B. M. Schilder, H. M. Petry, P. R. Hof, Evolutionary shifts dramatically reorganized the human hippocampal complex. J. Comp. Neurol. 528, 3143–3170 (2020). https://doi.org/10.1002/cne.24822

31. H. Van Praag, G. Kempermann, F. H. Gage, Neural consequences of environmental enrichment. Nat. Rev. Neurosci. 1, 191–198 (2000).

32. M. L. Holding, J. A. Frazier, E. N. Taylor, C. R. Strand. Experimentally altered navigational demands induce changes in the cortical forebrain of free-ranging Northern Pacific Rattlesnakes (*Crotalus o. oreganus*). Brain Behav. Evol. 79, 144–154 (2012).

33. D. F. Pfau, Plasticity in Adult Male Sceloporus occidentalis. (California Polytechnic State University, San Luis Obispo, 2012).

34. T. C. Roth, L. D. LaDage, C. A. Freas, V. V. Pravosudov, Variation in memory and the hippocampus across populations from different climates: a common garden approach. Proc. R. Soc. B 279, 402–410 (2012).

35. L. D. LaDage, T. C. Roth II, B. Sinervo, V. V. Pravosudov, Environmental experiences influence cortical volume in territorial and nonterritorial side blotched lizards, Uta stansburiana. Anim. Behav. 115, 11–18 (2016).

36. J. J. Hopfield, Neurons with graded response have collective computational properties like those of two state neurons. Proc. Natl. Acad. Sci. U.S.A. 81, 3088–3092 (1984). https://doi.org/10.1073/pnas.81.10.3088

37. Y. Roudi, P. E. Latham, A balanced memory network. PLoS Comput. Biol. 3, e141 (2007). https://doi.org/10.1371/journal.pcbi.0030141

38. K. R. Hedrick, K. Zhang, Megamap: flexible representation of a large space embedded with nonspatial information by a hippocampal attractor network. J. Neurophysiol. 116, 868–891 (2016). https://doi.org/10.1152/jn.00856.2015

39. S. Herculano-Houzel, B. Torres, F. Cunha, D. J. Messeder, P. R. Manger, J. H. Kaas. The number of hippocampal neurons is a linear predictor of the pace of life across mammalian species. In preparation.

40. J. Maliković, J. L. Cantalapiedra, L. Vinciguerra, K. Schönbächler, A. L. F. Destro, J. Rodger, M. Jörimann, L. Las, S. G. Hörpel, D. P. Wolfer, L. Slomianka & I. Amrein, Traces of phylogeny and ecology in hippocampal neuron numbers. *PNAS Nexus* 4, pgaf261 (2025). https://doi.org/10.1093/pnasnexus/pgaf261

41. K. Avelino-de-Souza, N. Patzke, K. A. Karlsson, P. R. Manger, Cellular composition of the brain of a Northern minke whale. *J. Comp. Neurol.* 533: e70089 (2025).

42. M. J. E. Broekman, S. Hoeks, R. Freriks, M. M. Langendoen, K. M. Runge, E. Savenco, R. ter Harmsel, M. A. J. Huijbregts, M. A. Tucker, HomeRange: A global database of mammalian home ranges. Glob. Ecol. Biogeogr. 32, 198–205 (2023). https://doi.org/10.1111/geb.13625

43. L. Santini, V. Y. Mendez Angarita, C. Karoulis, D. Fundarò, N. Pranzini, C. Vivaldi, T. Zhang, A. Zampetti, S. J. Gargano, D. Mirante, L. Paltrinieri, TetraDENSITY 2.0—A database of population density estimates in tetrapods. Glob. Ecol. Biogeogr. 33, e13929 (2024). https://doi.org/10.1111/geb.13929

44. J. P. de Magalhães, J. A. Costa, A database of vertebrate longevity records and their relation to other life history traits. J. *Evol. Bio*l. 22, 1770–1774 (2009). https://doi.org/10.1111/j.1420-9101.2009.01783.x

45. K.A. Nagy. Field Bioenergetics of Mammals—What Determines Field Metabolic Rates. *Aust J Zool* 42 (1): 43-53 (1994). https://doi.org/10.1071/ZO9940043

46. F. de Castro. FmrBT: A comprehensive data set of field metabolic rates, body mass and ambient temperature. *Scientific Data* 12: 1589 (2025).

47. H. Pontzer, D. A. Raichlen, A. D. Gordon et al. Primate energy expenditure and life history. *Proc. Natl. Acad. Sci. USA* 111:1433-1437 (2014).

48. S. Herculano-Houzel, Longevity and sexual maturity vary across species with number of cortical neurons, and humans are no exception. *J. Comp. Neurol*. 527, 1689-1705 (2019). https://doi.org/10.1002/cne.24564.

49. G. M. Mace, P. H. Harvey, Energetic constraints on home range size. Am. Nat. 121, 120–132 (1983).

50. K. E. Jones, J. Bielby, M. Cardillo, S. A. Fritz, J. O'Dell, C. D. L. Orme, K. Safi, W. Sechrest, E. H. Boakes, C. Carbone, C. Connolly, M. J. Cutts, J. K. Foster, R. Grenyer, M. Habib, C. A. Plaster, S. A. Price, E. A. Rigby, J. Rist, A. Teacher, O. R. P. Bininda Emonds, J. L. Gittleman, G. M. Mace, A. Purvis, PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. Ecology 90, 2648 (2009).

51. J. Seigle-Ferrand, K. Atmeh, J.-M. Gaillard, V. Ronget, N. Morellet, M. Garel, A. Loison, G. Yannic, A systematic review of within population variation in the size of home range across ungulates: What do we know after 50 years of telemetry studies? Front. Ecol. Evol. 8, 555429 (2021). https://doi.org/10.3389/fevo.2020.555429

52. M. Aronsson, M. Low, J. V. López Bao, J. Persson, J. Odden, J. D. C. Linnell, H. Andrén, Intensity of space use reveals conditional sex specific effects of prey and conspecific density on home range size. Ecol. Evol. 6, 2957–2967 (2016).

53. J. Balluffi Fry, Y. N. Majchrzak, M. J. L. Peers, E. K. Studd, A. K. Menzies, L. G. Horne, E. Monk, et al., Why does animal home range size decrease with population density? Ecology 106, e70054 (2025).

54. G. C. Sanderson, The study of mammal movements: a review. J. Wildl. Manage. 30, 215–235 (1966).

55. S. Herculano-Houzel, K. Catania, P. R. Manger, J. H. Kaas, Mammalian brains are made of these: a dataset of the numbers and densities of neuronal and nonneuronal cells in the brain of Glires, primates, Scandentia, Eulipotyphlans, Afrotherians and Artiodactyls, and their

relationship with body mass. Brain Behav. Evol. 86, 145–163
(2015). https://doi.org/10.1159/000437413

56. T. M. Williams, Racing time: physiological rates and metabolic scaling in marine
mammals. Integr. Comp. Biol. (2022). https://doi.org/10.1093/icb/icac054

57. B.K. McNab. An analysis of the factors that influence the level and scaling of
mammalian BMR. Comparative biochemistry and physiology. Part A, Molecular &
integrative physiology, 151(1), 5–28 (2008). https://doi.org/10.1016/j.cbpa.2008.05.008.

58. T. M. Williams, M. P.-H. Jørgensen, A. M. Pagano, C. M. Bryce, Hunters versus hunted:
new perspectives on the energetic costs of survival at the top of the food chain. Funct.
Ecol. 34, 2015–2029 (2020). https://doi.org/10.1111/1365-2435.13649

59. T. Eliav et al., Science 372, eabg4020 (2021). DOI: 10.1126/science.abg4020

60. S. Herculano-Houzel, The neuron number manifesto: Let's drop body mass from the
search for biological correlates of intelligent, flexible cognition in evolution. In Kaas JH,
Herculano-Houzel S (eds), Evolution of the Nervous System, 3rd edition, Volume II
(Krubitzer L, ed.). Elsevier, London, in press.

61. S. Herculano-Houzel, A supply-limited framework of brain function accounts for the
evolution of endothermic brains. In Kaas JH, Herculano-Houzel S (eds), Evolution of the
Nervous System, 3rd edition, Volume I (Herculano-Houzel S, ed.). Elsevier, London, in press.

62. M.C. Teles, G. Melo, S. Herculano-Houzel, R.F. Oliveira, Larger fish have larger brains
with more neurons across but not within cohorts raised in different growth conditions. J.
Comp. Neurol. 533, e70090 (2025).