Title: Small- and large-scale patches shape benthic microbial
community structure and function in streams at the subcontinental
scale

Running head: Patches shape stream microorganisms

Authors: Richard LaBrie[1,*], Rebecca L. Maher[2], Kelly S. Aho[3], Brittni L. Bertolet[4], Nicholas E Ray[5],
Paula C.J. Reis[6], Andrew L. Robison[7], Shannon L. Speir[8]

[1]Department of Earth & Planetary Sciences, McGill University, 3450 rue University,
Montréal, Québec, Canada, H3A 0E8
[2]Friday Harbor Labs, University of Washington, Friday Harbor, WA, USA
[3]Department of Biology, Boston University, Boston, MA 02215, USA
[4]Department of Earth and Environmental Sciences, Rutgers University Newark, Newark, NJ
07102, USA
[5]School of Marine Science and Policy, University of Delaware, Lewes DE 19958 USA
[6]Department of Biology, University of Waterloo, Waterloo, ON, N2L3G1, Canada
[7]Department of Natural Resources and the Environment, University of New Hampshire,
Durham, NH, USA 03824
[8]Department of Crop, Soil, and Environmental Sciences, University of Arkansas,
Fayetteville, AR, USA 72701
22
23 *Corresponding author(s): richard.labrie90@gmail.com
24

ORCID
RL 0000-0003-1681-4888
RLM 0000-0003-3108-8503
KSA 0000-0002-1992-6378
BLB 0000-0002-0581-1158
NER 0000-0002-1959-3120
PCJR 0000-0001-6940-0545
ALR 0000-0003-1628-6237
SLS 0000-0002-3624-0327

Keywords

Stream microbial ecology, biogeochemistry, nitrogen cycle, NEON, open science

AUTHOR CONTRIBUTION STATEMENT
RL wrote the first draft, all others contributed; All authors edited the manuscript; RLM
curated metagenomic data, PCJR and BB curated 16S data, ALR, SLS and KSA curated
environmental data; RL, RLM, PCJR, BB and ALR analyzed the data; SLS, ALR and RL

managed the project at different stages; ALR conceived the study as part of the Eco-DAS
program, all authors contributed to the final study design.
44

SCIENTIFIC SIGNIFICANCE STATEMENT (125 words, max is 125)
Microbial communities in streams process materials as water flows toward the ocean. As
microbial communities are influenced by environmental conditions, it remains challenging to
predict stream microbial metabolism at large spatial scales. In this study, we used the
National Ecological Observatory Network (NEON) public database to investigate the drivers
of microbial community structure and functions in stream sediments across the USA. This
unique dataset revealed high variability in microbial communities among streams that were
best explained by local conditions, then by water quality and streambed habitat type.
However, stream physico-chemical characteristics emerged as strong predictors of microbial
communities when field replicates were averaged and considered as single, large microbial
communities. These findings indicate multiple spatial scales must be considered to fully
understand benthic microbial communities.
57
58

ABSTRACT
Streams and rivers process dissolved and particulate matter as water moves along the land-to-
ocean continuum, making important contributions to global biogeochemical cycles. Yet,
predicting stream and river microbial metabolism associated with biogeochemical
transformations at broad spatial scales remains challenging. Here, we used data from the
National Ecological Observatory Network program to investigate whether ecological
relationships among microbial community structure and function and environmental
conditions observed at small scales hold at the subcontinental scale. We found that microbial
communities were best explained by site-specific conditions. However, when field replicates
were averaged, stream physico-chemical characteristics such as pH and temperature emerged
as driving factors. This indicates that water quality acted as an environmental filter on
microbial communities at the subcontinental scale, but was masked by small-scale patches
that created high spatial heterogeneity. Our findings underscore the importance of
considering multiple spatial scales to fully understand benthic microbial communities' role in
stream biogeochemistry.

74
75

INTRODUCTION

Streams and rivers are critical components of global elemental cycles, transporting and processing dissolved and particulate matter along the land-to-ocean aquatic continuum (Cole et al. 2007; Maranger et al. 2018; Regnier et al. 2022). Within these ecosystems, metabolism and biogeochemical conditions are driven by microbial communities (Battin et al. 2016). Simultaneously, stream biogeochemical conditions—such as pH, oxygen concentration and organic matter inputs—can also regulate microbial community structure, and there have been substantial research efforts to link microbial communities to specific biogeochemical functions (Bier et al. 2015; Zeglin 2015). Understanding the bidirectional relationship between microbes and their environment is essential for predicting how aquatic ecosystems respond to environmental and climate change. Yet, this remains challenging because of the scale-dependency between microbial communities and environmental variables (Wu and Loucks 1995; LaBrie et al. 2020).

Benthic microbial communities in lotic ecosystems are structured by environmental conditions acting at varying scales. For instance, the immediate overlaying water column controls many physical and chemical conditions such as temperature, organic matter, and nutrient delivery (Battin et al. 2003, 2016; Akinwole et al. 2021). Benthic microbial communities are also influenced by the physical structure of their local habitat, including substrate type and stream depth (Hullar et al. 2006; Cruaud et al. 2020; Gao et al. 2024). These environmental conditions can change at the centimeter to meter-scale, and together, favor high heterogeneity in benthic microbial communities (Wu and Loucks 1995; Clapcott and Barmuta 2010; Briggs et al. 2015). As ecological scale increases, drivers of microbial community structure and function change from being dominated by local stochasticity to larger scale processes such as geological formations and geomorphological and climatological changes (Wu and Loucks 1995; Besemer et al. 2009; Ezzat et al. 2022). Yet, we lack an understanding of the key controls on stream benthic microbial community composition and function at continental scales as most studies have been largely limited to local and regional scales (Zeglin 2015; Hosen et al. 2017; Ruiz-González et al. 2019). In turn, this limits our ability to predict microbial responses to broader climatic changes. A first step in addressing this challenge is to determine whether universal biogeochemical drivers of lotic microbial communities can be identified from existing datasets approaching the continental-scale.

Here, we used data from 23 National Ecological Observatory Network (NEON) streams sampled in 2019 to relate patterns in microbial diversity and function to biogeochemistry across the United States, from Alaska to Puerto Rico. NEON sites can be classified in six different groups based on characteristic water chemistry signatures (Edmonds et al. 2022). As such, we aimed to understand if these water quality groups translated to patterns in microbial communities, as water chemistry has been shown to be an important driver of microbial community structure up to regional scales (Battin et al. 2016; Crowther et al. 2019). Specifically, we sought to answer the following research questions: 1) how do benthic microbial community structure and functional gene diversity vary at the subcontinental scale? 2) is this variation, if present, correlated with patterns in stream

120 biogeochemistry? By using a subcontinental-scale dataset, we can assess the relative
121 contribution of local and regional drivers (e.g., water temperature, biogeochemical
122 conditions, sediment properties) to large-scale patterns in microbial community structure and
123 function.
124
METHODS
We leveraged data collected by NEON from 24 wadeable streams (**Table S1**) across the
United States, from Puerto Rico to Alaska (**Figure S1**). These streams span 16 of the 20
ecoclimatic domains used by NEON when determining site placement (Hargrove and
Hoffman 2004). We targeted sampling efforts between 2019/01/01 and 2019/12/31. This
timeframe was selected for even sample coverage across sites; one site (OKSR) was excluded
from analysis due to low sequencing yield in 16S rRNA gene and metagenomic data. Land
cover varies by site, but the majority of NEON sites are located on smaller streams not highly
impacted by anthropogenic activity. Site characteristics such as latitude and longitude,
watershed area, and elevation were obtained from the NEON site descriptions
(neonscience.org/field-sites/explore-field-sites). Streambed slope was collected from NEON
thalweg surveys (DP4.00131.001) and land cover was taken from the NEON Aquatic
Watershed shapefile (neon.maps.arcgis.com). Benthic quality-controlled 16S rRNA gene
sequences (DP1.20280.001, 2019/01-2019/11), shotgun metagenomic sequences
(DP1.20279.001, 2019/06-2019/08), and environmental data (2019/01-2019/11) were
downloaded from the NEON data portal using the R package NeonUtilities (Lunch et al.
2024).
142
Sampling
The detailed strategy for benthic microbial sampling is available on NEON website
(NEON.DOC.003044). Briefly, benthic microbes were sampled three times per year (roughly
spring, summer and fall). Only benthic samples taken during the summer sampling event
were analyzed for metagenomics. Scrub sampling was performed on rocks (epilithon) and
wood (epixylon) to collect the attached biofilm. Biofilms were then resuspended in 125 to
140 mL of 0.2 μm filtered deionized water and then at least 50 mL (or until filter got clogged)
were filtered on 0.22 μm filters (Sterivex). Grab samples were collected for sand
(epipsammon), silt (epipelon) and plant surfaces (epiphyton) samples. Filters and whirl-pak
bags with sample materials were flash-frozen on dry ice in the field and stored at -80 °C until
analysis. DNA extraction and sequencing protocols can be found on NEON website
(BMI_dnaExtractionSOP, BMI_16Sv4v5_SOP, and BMI_metagenomicsSequencingSOP)
and in the SI.
Bioinformatics
16S rRNA gene
Benthic 16S rRNA gene sequences samples encompass 23 benthic NEON sites (i.e., all but
OKSR), totaling 459 samples analyzed in this study. Sequences were analyzed using the
dada2 pipeline v1.22.0 (Callahan et al. 2016) in R v4.1.2 (R Core Team 2022). Read quality
of each sequence run was visually inspected using quality profiles, and the forward and
reverse reads were trimmed to 275 and 250 nucleotides, respectively. Primers were removed

using the trimLeft() argument in the filterAndTrim() function. After filtering for quality, the
average number of reads was 22,324 per sample. Amplicon Sequence Variant (ASV) tables
were produced following the dada2 pipeline including the removal of chimera sequences with
an additional step to collapse together sequences that are only different in length
(collapseNoMismatch(); minOverlap = 300) to minimize false singletons. Due to the low
quality of samples produced by a specific sequence run, we used a threshold of a minimum
2,000 final reads per sample, resulting in the exclusion of 30 samples. Taxonomy assignment
was performed at the Genus level with SILVA v138 using dada2's 'assignTaxonomy'
function (Wang et al. 2007). Sequences that were unclassified (i.e., no Kingdom assigned;
2567 ASVs) or assigned to eukaryotes or chloroplasts were removed, resulting in 63,813
prokaryotic ASVs. Although not used in this study, surface samples were also curated (see
SI) and the final ASV table is publicly available (https://doi.org/10.5281/zenodo.15490674).

<u>Metagenomic data</u>
Benthic shotgun metagenomic samples underwent the following bioinformatic pipeline.
Paired-end reads were filtered using BBDuk v37.25 (sourceforge.net/projects/bbmap) to
remove adapters (k = 12, mink = 11) using normal mode with recommended flags "tpe" and
"tbo". Read pairs for 'dnaSamples' corresponding to a single field sample were co-assembled
using SPAdes v3.13.0 (Bankevich et al. 2012) without error correction. Filtered reads were
aligned to the metagenome assemblies using Bowtie2 v2.4.4 (Langmead et al. 2019) and
SAMtools v1.9 (Li et al. 2009) to generate coverage bam files. Contigs greater than 500
basepairs were processed in Anvi'o v7.1 (Eren et al. 2015) for each metagenome assembly.
Briefly, anvi-gen-contigs-database was used to generate contig databases and call genes with
Prodigal v2.6.3 (Hyatt et al. 2010). Metagenome assemblies were functionally annotated with
KOfam, a customized HMM database of KEGG Orthologs (Aramaki et al. 2020), Pfam
(Mistry et al. 2021), and NCBI's database of Clusters of Orthologous Groups of proteins
(Tatusov et al. 2003). Next, anvi-profile-blitz was used to export gene-level coverage and
detection statistics from the contig databases and coverage bam files. Lastly, functional
annotations from all annotation sources were exported from Anvi'o and combined with gene-
level statistics into gene-by-sample tables using the custom script process_blitz_neon.R
(https://zenodo.org/records/17154747). Gene coverages were normalized based on reads per
kilobase per million mapped (RPKM) which corrects differences in both sample sequencing
depth and gene length (Konwar et al. 2015). Prodigal gene calls were also annotated using
DIAMOND v2.0.15 against NCycDB, a curated database of nitrogen cycling genes (Tu et al.
2019).

197
Environmental data
Environmental data includes grab samples of chemical properties of surface water
(DP1.20093.001), continuous discharge (DP4.00130.001), photosynthetically active radiation
at the water surface (DP1.20042.001), sediment chemical and physical properties
(DP1.20194.001), water temperature (DP1.20053.001), and sensor-based measurements of
dissolved oxygen, pH, conductivity, and turbidity (DP1.20288.001). For grab samples,
concentrations below the published detection limit were assigned a value of half the detection
limit.

206
Environmental data was aligned with microbial sampling dates by finding the nearest
sampling period in time. Some variables were then either combined or removed from the
dataset due to significant autocorrelation with other variables (see SI for details). The final
dataset contains 35 variables, including 11 watershed and sampling location characteristics,
19 water column variables, and five sediment variables. To assess the influence of water
quality using lower dimensionality, we classified sites into one of six water quality groups as
determined by Edmonds et al. (2022).
214
Statistical analysis
As sequencing depths and the number of replicates varied among sites for the 16S rRNA
gene data, we pooled all field replicates together, computed 1000 different rarefied ASV
tables per sample (rrarefy function, vegan, Oksanen et al. 2010) and then averaged them. This
enabled us to calculate total richness and Shannon Diversity Index on comparable sites and
avoided pseudoreplication when looking at relationships with environmental drivers.
Functional diversity metrics were also calculated from gene-by-sample tables (*diversity*
function, *vegan*). To ordinate samples based on their compositional dissimilarity, we
performed non-metric multidimensional scaling (NMDS) with Bray-Curtis distances
(*metaMDS* function*, vegan*) on both metagenomic and 16S rRNA gene data using *phyloseq*
v1.50.0 (McMurdie and Holmes 2013). We then projected environmental variables that were
significant at p-value < 0.01 on 16S rRNA gene data (*envfit* function, *vegan*). Lastly,
statistical differences in taxonomic and functional composition among groups (substrate type,
water quality group, NEON site) were tested using PERMANOVA (*adonis2* function, *vegan*)
with 999 permutations (Anderson 2001). PERMDISP (*betadisper* and *permutest* functions,
vegan) was used to test for uneven dispersion of groups (Anderson 2006).
231
RESULTS
Stream benthic communities
Microbial community structure and function composition in benthic streams were best
described by local conditions, not water chemistry. Indeed, community structure (**Figure 1A**
and B) was best explained by site (PERMANOVA, $R^2 = 0.31$, p < 0.001), followed by water
quality group ($R^2 = 0.08$, p < 0.001) and substrate type ($R^2 = 0.06$, p < 0.001). Alpha diversity
metrics were generally not well predicted by sites, water quality groups and substrate types,
except for the Shannon index which was well constrained by substrate types (**Table S2;**
Figure S2). Although there were fewer samples for functional diversity (**Figure 1C and D**),
microbial functions were still best explained by site (PERMANOVA, $R^2 = 0.37$, p < 0.001),
though substrate type ($R^2 = 0.23$, p < 0.001) and water quality group explained slightly more
variation ($R^2 = 0.14$, p < 0.001). Site, substrate type, and water quality group were all
significant predictors of the number and diversity of microbial functions (**Table S2; Figure**
S3). A general summary of microbial community composition (**Figure S4 to S8**) and
metagenomic samples (**Figure S9 and S10**) can be found in the supplementary information.
247

Figure 1. Non-metric multidimensional scaling (NMDS) ordinations showing variation in microbial community composition and function using the 16S rRNA gene (A and B) and KOfam annotations (C and D). Panels A and C are color-coded as a function of site name, B and D, as a function of water quality groups. For all panels, shapes represent substrate types.

Interrelationships between pH and water quality, and benthic communities

Beyond local conditions, we investigated which environmental variables correlated with community structure across substrate types and within epilithon samples by averaging field replicates. Using NMDS, we displayed samples using either water quality groups (**Figure 2**) or sites (**Figure S11**) and projected environmental variables that were statistically significant. For all substrate types combined, and in epilithon only, the top environmental variables were sediment pH, water quality groups, and water column dissolved inorganic carbon (**Table S3**). Latitude and water temperature were also important for epilithon samples and had opposite effects to each other (**Figure 2B**). Overall, these results show that water quality groups and pH are significantly related to benthic community structure.

268
269 **Figure 2**. Non-metric multidimensional scaling (NMDS) plot showing variation in microbial community
270 structure (16S rRNA gene) for all substrate types (A) and epilithon only (B) after averaging field replicates.
271 Environmental variables that were statistically significant (p-value < 0.01) were passively projected on the
272 ordination plots using the envfit function. Colors represent water quality groups and shapes, substrate type.
273
274

Abundance of genes related to nitrogen and methane cycling genes vary by substrate type

Using epilithon and epipsammon metagenomic samples, we found that nitrogen- and
methane-related genes varied among substrate types (**Figure 3**). Specifically, the average
RPKM related to nitrogen was lower in epilithon than in epipsammon, both overall and in
site-specific pairwise comparisons (**Figure 3A**), but many nitrogen cycling pathways were
ubiquitous across sites and substrates. Anaerobic ammonium oxidation (i.e., anammox),
however, was found at only one site (REDB epipsammon) in low abundance (not visible in
figure 3A). Across streams, genes associated with heterotrophic processes (e.g.,
denitrification and dissimilatory nitrate reduction to ammonium [DNRA]) dominated. Taken

together, genes associated with nitrogen recycling were more abundant than nitrogen
removal. Genes related to methanogenesis were always more abundant than genes related to
methane oxidation (**Figure 3B**). While methanogenesis and methane oxidation-related genes
covaried in epipsammon samples, they did not in epilithon samples, where methane
oxidation-related genes were almost absent.
290
291

292
293 **Figure 3**. Abundance of nitrogen-related genes as a function of sites and substrate types where colors represent
294 nitrogen cycling pathways (A). Methanogenesis-related and methane oxidation-related genes for epilithon
295 (black) and epipsammon (blue) substrate types (B). In epipsammon, the two methane pathways covaried, but not
296 in epilithon. Anammox: anaerobic ammonium oxidation; DNRA: dissimilatory nitrate reduction to ammonium;
297 RPKM: reads per kilobase per million mapped
298
299

DISCUSSION
Using publicly available NEON microbial sequencing data and biogeochemistry data,
we present the first subcontinental patterns in microbial taxonomic and functional diversity in
streams located across the USA. When considering field replicates individually, we found
that water quality was not the main driver of community structure. However, when field
replicates were averaged, water quality groups as an integrative variable became a strong
predictor of community structure and function. More specifically, pH and average

temperature were strong predictors of community structure as observed in soil and other
stream studies (Sutton and Findlay 2003; Lauber et al. 2009; Niño□García et al. 2016;
Philippot et al. 2024). Previous studies observing ecologically significant relationships
between geochemical parameters and benthic microbial communities have generally been
limited to highly impacted streams (Zhou et al. 2021) or extreme environments (Ezzat et al.
2022). Thus, although the water quality groups used in this study are coarse, our findings
suggest that broad-scale patterns in water quality act as environmental filters for benthic
microbial communities at subcontinental scales even in relatively pristine systems, with
implications for understanding stream microbial processes at the continental scale.
316
Substrate type influences genes related to biogeochemical transformations
Across all sites, community structure and function were best explained by location,
but could also be explained by substrate type. This supports the idea that habitat structure
influences microbial communities in aquatic ecosystems, as observed at smaller scales
(Besemer et al. 2009; Singer et al. 2010; Cruaud et al. 2020; Sun et al. 2023). For instance,
streambed composition controls the life-cycle of biofilms (Chen et al. 2022) and influences
oxygen exchange between running water and hyporheic flow (Jones and Holmes 1996;
Mueller et al. 2013). Oxygen availability likely contributed to the contrasting microbial
communities, and specifically genes related to nitrogen metabolic pathways, between
epipsammon and epilithon substrates. Observed differences in nitrogen cycling microbial
communities were likely related to redox gradients and although anoxia can develop in
epilithic biofilms (Anderson-Glenna et al. 2008; Flemming et al. 2016), it appeared to be less
common in our dataset, potentially as a result of the sampling strategy—grab samples were
used for epipsammon in which oxic-anoxic gradients can be present at the centimeter scale
(Briggs et al. 2015). In epipsammon samples, nitrification genes coexisted with
denitrification and DNRA, as did methanogenesis and methane oxidation-related genes. As
these metabolic pathways are dependent on oxic-anoxic coexistence, it suggests that spatial
complexity (i.e., patches) of oxygen availability had a stronger influence on microbial
communities than nutrient availability. Patches of oxygen availability were likely more
abundant and in closer proximity in epipsammon due to finer substrates than in epilithon
samples.
338
Patches across spatial scales
Averaging community structure on a site–substrate type basis revealed the strong
influence of water quality groups, pH and temperature on microbial communities. This
implies high spatial heterogeneity in which small-scale patches within stream reaches
obscured the influence of biogeochemical conditions (Akinwole et al. 2021; Bier et al. 2023).
Indeed, the choice of scale strongly influences our ability to observe patterns (Wu and
Loucks 1995) where small-scale processes may appear as stochastic variation at larger scales,
and likewise, large-scale relationships may dissipate at smaller scales. Patches, however, are
not limited to small-scale features; they can also appear at larger scales, creating nested
processes (Wu and Loucks 1995). For instance, Puerto Rican CUPE and GUIL sites did not
differ much from other sites in terms of water and sediments biogeochemical conditions
(**Figure S12**), but had unique epilithon communities, displayed the highest level of unique

taxa and among the highest number of metabolic functions. This suggests that local
conditions specific to these sites were not well-captured despite the broad NEON scope.
Local enrichments to benthic microbial communities likely created patches of high diversity
in both CUPE and GUIL sites—Puerto Rico may thus represent a large-scale patch up to
~10,000 km² that is both nesting smaller scale patches and is nested within the larger system
(**Figure 4**). Revealing the ecological importance of patches at the subcontinental scale was
possible given the high spatial sampling effort—both within reach and across ecoregions—of
the NEON program.

359
Figure 4. Representation of nested patches across spatial scales, including at the subcontinental scale with water
quality groups (A, < ~10,000,000 km²), Puerto Rico CUPE and GUIL sites (B, < ~10,000 km²), among substrate
type (C, < ~1,250 m²), and within a grab sample (D, < 0.1 m²). Symbols in A and B represent water quality
groups as defined by Edmonds et al. (2022), and represent microbial communities in C and D. The inset in B
uses the same NMDS as figure 1A but only displays sites for water quality group 3; colors are shared between
the map and the NMDS.

366
CONCLUSION
Open science provides a segue from spatially-limited field observation programs to
large-scale studies. By expanding spatial extents to subcontinental scales, environmental
gradients and ecological niches are more diverse, allowing robust assessments of our
fundamental understanding of aquatic ecosystems and ecological theories. While
biogeochemical conditions can be the main driver of community structure in streams

(Peipoch et al. 2019), our subcontinental approach highlighted the greater influence of
patches, from small (within reach) to large (Puerto Rico and larger) scales. Although within-
reach patches did not relate broadly to water and sediment biogeochemical conditions,
presumably because of the scale-dependency of environmental variables (LaBrie et al. 2020),
patches are critical for stream ecosystem processes by maximizing microbial biomass and
functions through niche partitioning (Cardinale 2011; Battin et al. 2016), ensuring high
functional redundancy and resilience (Comte et al. 2013; Pelletier et al. 2020). Our findings
suggest that multiple spatial scales—from within a stream reach to continental scale—must
be considered to fully understand stream benthic microbial communities and how they impact
biogeochemical cycles.
383
384

ACKNOWLEDGEMENTS
This material is based in part upon work supported by the National Ecological Observatory
Network (NEON), a program sponsored by the U.S. National Science Foundation (NSF) and
operated under cooperative agreement by Battelle. We thank the Ecological Dissertations in
the Aquatic Sciences (Eco-DAS)-XIV organized by Paul Kemp and Krissy Remple. We
thank Andrew H. Morris for discussions and advice on analyzing metagenomic data. We
thank the National Science Foundation (NSF, award OCE-1925796) and Association for the
Sciences of Limnology and Oceanography (ASLO) for funding and fostering this
collaboration. Part of this work was funded by the Wares and Trottier Space Institute
postdoctoral fellowships to RL; the AMTD Waterloo Global Talent postdoctoral fellowship
to PCJR and the Friday Harbor Laboratories William Calvin Postdoctoral Fellowship to
RLM.

REFERENCES
Akinwole, P. O., J. Kan, L. A. Kaplan, and R. H. Findlay. 2021. Spatial Variability in

Streambed Microbial Community Structure across Two Watersheds. Microbiol.

Spectr. **9**: e01972-21. doi:10.1128/Spectrum.01972-21

Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance.

Austral Ecol. **26**: 32–46. doi:10.1111/j.1442-9993.2001.01070.pp.x

Anderson, M. J. 2006. Distance-based tests for homogeneity of multivariate dispersions.

Biometrics **62**: 245–253. doi:10.1111/j.1541-0420.2005.00440.x

Anderson-Glenna, M. J., V. Bakkestuen, and N. J. W. Clipson. 2008. Spatial and temporal

variability in epilithic biofilm bacterial communities along an upland river gradient.

FEMS Microbiol. Ecol. **64**: 407–418. doi:10.1111/j.1574-6941.2008.00480.x

409 Aramaki, T., R. Blanc-Mathieu, H. Endo, K. Ohkubo, M. Kanehisa, S. Goto, and H. Ogata.

410   2020. KofamKOALA: KEGG Ortholog assignment based on profile HMM and

411   adaptive score threshold. Bioinformatics **36**: 2251–2252.

412   doi:10.1093/bioinformatics/btz859

413 Bankevich, A. and others. 2012. SPAdes: A New Genome Assembly Algorithm and Its

414   Applications to Single-Cell Sequencing. J. Comput. Biol. **19**: 455–477.

415   doi:10.1089/cmb.2012.0021

416 Battin, T. J., K. Besemer, M. M. Bengtsson, A. M. Romani, and A. I. Packmann. 2016. The

417   ecology and biogeochemistry of stream biofilms. Nat. Rev. Microbiol. **14**: 251–263.

418   doi:10.1038/nrmicro.2016.15

419 Battin, T. J., L. A. Kaplan, J. D. Newbold, and S. P. Hendricks. 2003. A mixing model

420   analysis of stream solute dynamics and the contribution of a hyporheic zone to

421   ecosystem function. Freshw. Biol. **48**: 995–1014. doi:10.1046/j.1365-

422   2427.2003.01062.x

423 Besemer, K., G. Singer, I. Hödl, and T. J. Battin. 2009. Bacterial Community Composition of

424   Stream Biofilms in Spatially Variable-Flow Environments. Appl. Environ. Microbiol.

425   **75**: 7189–7195. doi:10.1128/AEM.01284-09

426 Bier, R. L. and others. 2015. Linking microbial community structure and microbial processes:

427   an empirical and conceptual overview. FEMS Microbiol. Ecol. **91**: fiv113.

428 Bier, R. L., J. J. Mosher, L. A. Kaplan, and J. Kan. 2023. Spatial scale impacts microbial

429   community composition and distribution within and across stream ecosystems in

430   North and Central America. Environ. Microbiol. **25**: 1860–1874. doi:10.1111/1462-

431   2920.16396

432 Briggs, M. A., F. D. Day-Lewis, J. P. Zarnetske, and J. W. Harvey. 2015. A physical

433   explanation for the development of redox microzones in hyporheic flow. Geophys.

434   Res. Lett. **42**: 4402–4410. doi:10.1002/2015GL064200

Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P.

Holmes. 2016. DADA2: high-resolution sample inference from Illumina amplicon

data. Nat. Methods **13**: 581. doi:10.1038/nmeth.3869

Cardinale, B. J. 2011. Biodiversity improves water quality through niche partitioning. Nature

472: 86–89. doi:10.1038/nature09904

Chen, X., C. Zhang, I. H. Townend, Z. Gong, Q. Feng, and X. Yu. 2022. The Resilience of

Biofilm-Bound Sandy Systems to Cyclic Changes in Shear Stress. Water Resour.

Res. **58**: e2021WR031098. doi:10.1029/2021WR031098

Clapcott, J. E., and L. A. Barmuta. 2010. Metabolic patch dynamics in small headwater

streams: exploring spatial and temporal variability in benthic processes. Freshw. Biol.

55: 806–824. doi:10.1111/j.1365-2427.2009.02324.x

Cole, J. J. and others. 2007. Plumbing the Global Carbon Cycle: Integrating Inland Waters

into the Terrestrial Carbon Budget. Ecosystems **10**: 172–185. doi:10.1007/s10021-

006-9013-8

Comte, J., L. Fauteux, and P. A. del Giorgio. 2013. Links between metabolic plasticity and

functional redundancy in freshwater bacterioplankton communities. Front. Microbiol.

4. doi:10.3389/fmicb.2013.00112

Crowther, T. W., J. van den Hoogen, J. Wan, M. A. Mayes, A. D. Keiser, L. Mo, C. Averill,

and D. S. Maynard. 2019. The global soil community and its influence on

biogeochemistry. Science **365**: eaav0550. doi:10.1126/science.aav0550

Cruaud, P., A. Vigneron, C. C. Dorea, M. J. Rodriguez, and S. J. Charette. 2020. Rapid

Changes in Microbial Community Structures along a Meandering River.

Microorganisms **8**: 1631. doi:10.3390/microorganisms8111631

Edmonds, J. W., K. B. S. King, M. B. Neely, R. T. Hensley, K. J. Goodman, and K. M.

Cawley. 2022. Using large, open datasets to understand spatial and temporal

patterns in lotic ecosystems: NEON case studies. Ecosphere **13**: e4102.

doi:10.1002/ecs2.4102

Eren, A. M., Ö. C. Esen, C. Quince, J. H. Vineis, H. G. Morrison, M. L. Sogin, and T. O.

Delmont. 2015. Anvi'o: an advanced analysis and visualization platform for 'omics

data. PeerJ **3**: e1319. doi:10.7717/peerj.1319

Ezzat, L. and others. 2022. Benthic Biofilms in Glacier-Fed Streams from Scandinavia to the

Himalayas Host Distinct Bacterial Communities Compared with the Streamwater.

Appl. Environ. Microbiol. **88**: e00421-22. doi:10.1128/aem.00421-22

Flemming, H.-C., J. Wingender, U. Szewzyk, P. Steinberg, S. A. Rice, and S. Kjelleberg.

2016. Biofilms: an emergent form of bacterial life. Nat. Rev. Microbiol. **14**: 563–575.

doi:10.1038/nrmicro.2016.94

Gao, Y. and others. 2024. Structural changes and assembly mechanisms of microbial

communities during rapid sedimentation of Yellow River sediments. Environ.

Technol. Innov. **35**: 103702. doi:10.1016/j.eti.2024.103702

Hargrove, W. W., and F. M. Hoffman. 2004. Potential of Multivariate Quantitative Methods

for Delineation and Visualization of Ecoregions. Environ. Manage. **34**: S39–S60.

doi:10.1007/s00267-003-1084-0

Hosen, J. D., C. M. Febria, B. C. Crump, and M. A. Palmer. 2017. Watershed Urbanization

Linked to Differences in Stream Bacterial Community Composition. Front. Microbiol.

8. doi:10.3389/fmicb.2017.01452

Hullar, M. A. J., L. A. Kaplan, and D. A. Stahl. 2006. Recurring Seasonal Dynamics of

Microbial Communities in Stream Habitats. Appl. Environ. Microbiol. **72**: 713–722.

doi:10.1128/AEM.72.1.713-722.2006

Hyatt, D., G.-L. Chen, P. F. LoCascio, M. L. Land, F. W. Larimer, and L. J. Hauser. 2010.

Prodigal: prokaryotic gene recognition and translation initiation site identification.

BMC Bioinformatics **11**: 119. doi:10.1186/1471-2105-11-119

Jones, J. B., and R. M. Holmes. 1996. Surface-subsurfaceinteractions in stream

ecosystems. Trends Ecol. Evol. **11**: 239–242. doi:10.1016/0169-5347(96)10013-6

Konwar, K. M. and others. 2015. MetaPathways v2.5: quantitative functional, taxonomic and

usability improvements. Bioinformatics **31**: 3345–3347.

doi:10.1093/bioinformatics/btv361

LaBrie, R., S. Bélanger, R. Benner, and R. Maranger. 2020. Spatial abundance distribution

of prokaryotes is associated with dissolved organic matter composition and

ecosystem function. Limnol. Oceanogr. **66**: 575–587. doi:10.1002/lno.11624

LaBrie, R., R. Maher, K. Aho, B. Bertolet, N. Ray, P. C. Reis, A. Robison, and S. Speir.

2025. 16S surface water samples for paper titled "Small- and large-scale patches

shape benthic microbial community structure and function in streams at the

subcontinental scale."doi:10.5281/zenodo.15490674

Langmead, B., C. Wilks, V. Antonescu, and R. Charles. 2019. Scaling read aligners to

hundreds of threads on general-purpose processors. Bioinformatics **35**: 421–432.

doi:10.1093/bioinformatics/bty648

Lauber, C. L., M. Hamady, R. Knight, and N. Fierer. 2009. Pyrosequencing-Based

Assessment of Soil pH as a Predictor of Soil Bacterial Community Structure at the

Continental Scale. Appl. Environ. Microbiol. **75**: 5111–5120. doi:10.1128/AEM.00335-

09

Li, H. and others. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics

25: 2078–2079. doi:10.1093/bioinformatics/btp352

Lunch, C., C. Laney, N. Mietkiewicz, E. Sokol, K. Cawley, and NEON (National Ecological

Observatory Network). 2024. neonUtilities: Utilities for Working with NEON Data,.

Maranger, R., S. E. Jones, and J. B. Cotner. 2018. Stoichiometry of carbon, nitrogen, and

phosphorus through the freshwater pipe. Limnol. Oceanogr. Lett. **3**: 89–101.

doi:https://doi.org/10.1002/lol2.10080

McMurdie, P. J., and S. Holmes. 2013. phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. PloS One **8**: e61217.

doi:10.1371/journal.pone.0061217

Mistry, J. and others. 2021. Pfam: The protein families database in 2021. Nucleic Acids Res.

49: D412–D419. doi:10.1093/nar/gkaa913

Mueller, M., J. Pander, R. Wild, T. Lueders, and J. Geist. 2013. The effects of stream

substratum texture on interstitial conditions and bacterial biofilms: Methodological

strategies. Limnologica **43**: 106–113. doi:10.1016/j.limno.2012.08.002

Niño-García, J. P., C. Ruiz-González, and P. A. Del Giorgio. 2016. Landscape-scale spatial

abundance distributions discriminate core from random components of boreal lake

bacterioplankton. Ecol. Lett. **19**: 1506–1515. doi:10.1111/ele.12704

Oksanen, J. and others. 2010. Vegan: community ecology package. R package version

1.17-4. HttpCRANR-Proj.

Peipoch, M., S. R. Miller, T. R. Antao, and H. M. Valett. 2019. Niche partitioning of microbial

communities in riverine floodplains. Sci. Rep. **9**: 16384. doi:10.1038/s41598-019-

52865-4

Pelletier, M. C. and others. 2020. Resilience of aquatic systems: Review and management

implications. Aquat. Sci. **82**: 1–44. doi:10.1007/s00027-020-00717-z

Philippot, L., C. Chenu, A. Kappler, M. C. Rillig, and N. Fierer. 2024. The interplay between

microbial communities and soil properties. Nat. Rev. Microbiol. **22**: 226–239.

doi:10.1038/s41579-023-00980-5

R Core Team. 2022. R: A Language and Environment for Statistical Computing, R

Foundation for Statistical Computing.

Regnier, P., L. Resplandy, R. G. Najjar, and P. Ciais. 2022. The land-to-ocean loops of the

global carbon cycle. Nature **603**: 401–410. doi:10.1038/s41586-021-04339-9

Ruiz-González, C. and others. 2019. Higher contribution of globally rare bacterial taxa

reflects environmental transitions across the surface ocean. Mol. Ecol. **28**: 1930–

1945. doi:10.1111/mec.15026

Singer, G., K. Besemer, P. Schmitt-Kopplin, I. Hödl, and T. J. Battin. 2010. Physical

Heterogeneity Increases Biofilm Resource Use and Its Molecular Diversity in Stream

Mesocosms. PLOS ONE **5**: e9988. doi:10.1371/journal.pone.0009988

Sun, Y., F. Ye, Q. Huang, F. Du, T. Song, H. Yuan, X. Liu, and D. Yao. 2023. Linking

ecological niches to bacterial community structure and assembly in polluted urban

aquatic ecosystems. Front. Microbiol. **14**: 1288304. doi:10.3389/fmicb.2023.1288304

Sutton, S. D., and R. H. Findlay. 2003. Sedimentary microbial community dynamics in a

regulated stream: East Fork of the Little Miami River, Ohio. Environ. Microbiol. **5**:

256–266. doi:10.1046/j.1462-2920.2003.00396.x

Tatusov, R. L. and others. 2003. The COG database: an updated version includes

eukaryotes. BMC Bioinformatics **4**: 41. doi:10.1186/1471-2105-4-41

Tu, Q., L. Lin, L. Cheng, Y. Deng, and Z. He. 2019. NCycDB: a curated integrative database

for fast and accurate metagenomic profiling of nitrogen cycling genes. Bioinformatics

35: 1040–1048. doi:10.1093/bioinformatics/bty741

Wang, Q., G. M. Garrity, J. M. Tiedje, and J. R. Cole. 2007. Naïve Bayesian Classifier for

Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy. Appl.

Environ. Microbiol. **73**: 5261–5267. doi:10.1128/AEM.00062-07

Wu, J., and O. L. Loucks. 1995. From Balance of Nature to Hierarchical Patch Dynamics: A

Paradigm Shift in Ecology. Q. Rev. Biol. **70**: 439–466. doi:10.1086/419172

Zeglin, L. H. 2015. Stream microbial diversity in response to environmental changes: review

and synthesis of existing research. Front. Microbiol. **6**: 454.

doi:10.3389/fmicb.2015.00454

Zhou, L., P. Wang, S. Huang, Z. Li, H. Gong, W. Huang, Z. Zhao, and Z. Yu. 2021.

Environmental filtering dominates bacterioplankton community assembly in a highly

urbanized estuarine ecosystem. Environ. Res. **196**: 110934.

doi:10.1016/j.envres.2021.110934