P. aeruginosa disrupts CDC-42/ARP-2/3 mediated F-actin organization to

induce intestinal deformation in *C. elegans*

Authors: AN Divyashree, Tanushree Sinha and Anup Padmanabhan[1]*

Affiliations:

[1]Department of Biology, Trivedi School of Biosciences, Ashoka University, Sonipat,

Haryana India -131029

*Correspondence to: anup.padmanabhan@ashoka.edu.in

ORCiD ids:

0000-0003-3127-2132 (AN Divyashree)

0009-0000-7405-283X (Tanushree Sinha)

0000-0001-6007-6929 (Anup Padmanabhan)

SUMMARY

Apico-basal polarity maintains the integrity of cortical actin cytoskeleton and regulates

secretory functions in intestinal epithelium essential for barrier defense against

infections. The opportunistic pathogen, *Pseudomonas aeruginosa*, exploits the

interplay between cell polarity and cytoskeletal machineries to subvert these defenses.

Mechanistic insights into such pathogen-induced cytoskeletal alterations have largely

been derived from *in-vitro* epithelial monolayers, which may not capture the influences

of multicellular physiology such as tissue mechanics, and innate immunity. Here, we

show that the extracellular *P. aeruginosa* disrupts apical polarity in *C. elegans*

enterocytes, leading to fragmentation of ARP-2/3 clusters and disorganization of apical

F-actin. This disruption causes shedding of actin-rich vesicles into lumen and a

pronounced apical deformation. Inhibition of CDC-42-mediated actin polymerization or

PI3K-AKT signalling attenuated enterocyte deformation and extended the lifespan of

C. elegans upon *P. aeruginosa* exposure*.* Our findings reveal a conserved strategy

by which *P. aeruginosa* exploits cellular polarity machinery to disrupt host actin

organization during extracellular infection of the intestinal epithelium.

Keywords : F-actin / ARP-2/3 / CDC-42 / Polarity / *P. aeruginosa*/ Host-pathogen

INTRODUCTION

Symbiotic interactions with the gut microbiota are essential for maintaining health and metabolism in metazoans. Constantly exposed to a diverse micro-organisms, the intestinal epithelium acts as a frontline sentinel for immune surveillance, detecting perturbations caused by dysbiosis, distinguishing harmless commensals from pathogens, and mounting context-appropriate immune response. During infection, bacterial pathogens elicit a wide range of responses from intestinal epithelial cells - some orchestrated by the pathogens to promote their survival and dissemination, and others initiated by the host to limit damage and trigger innate immunity. Unlike professional immune cells, enterocytes lack canonical pathogen recognition receptors (PRRs) that detect microbe-associated molecular patterns (MAMPs). Instead, they respond to pathogen effector activity and the disruptions they cause to physiological processes and cellular homeostasis, collectively termed "patterns of pathogenesis". These include damage associated molecular patterns (DAMPs) triggered by pathogen effectors. Studies in model organisms such as *Drosophila melanogaster* and *Caenorhabditis elegans (C. elegans)* have highlighted the role of effector-triggered, cell autonomous surveillance in epithelial immunity (Stuart *et al*, 2013; Tse-Kang *et al*, 2025; Vance *et al*, 2009).

C. elegans, a bacterivorous soil nematode, inhabits microbe-rich environments and encounters a broad spectrum of microorganisms, including pathogens (Berg *et al*, 2016; Schulenburg & Félix, 2017). While a protective cuticle (Gravato-Nobre *et al*, 2005) and sophisticated pathogen-avoidance behaviours (Lei *et al*, 2024; Melo & Ruvkun, 2012; Pereira *et al*, 2020; Prakash *et al*, 2021; Singh & Aballay, 2019b; Zhang *et al*, 2005) limit exposure, the intestinal epithelium remains the primary site of infection due to the worm's bacterial diet. Ingested bacteria are mechanically ground in the pharynx before entering the lumen, yet some pathogens survive, proliferate, and colonize the intestine. The *C. elegans* intestine is composed of 20 polarized epithelial cells arranged longitudinally, with apical surfaces facing the central lumen and basolateral surface engaged in cell-cell and cell-matrix adhesions (Dimov & Maduro, 2019; McGhee, 2007). The apical surface is enriched in microvilli, providing the primary site for host-pathogen interaction (MacQueen *et al,* 2005a) (Fig. 1B). Polarity is maintained by structural and signalling complexes: the Crumbs complex (CRB-1,

CRB-3, MAGUK-2) and the PAR complex (comprising PAR-3, PAR-6, PKC-3 kinase and CDC-42 GTPase) localize apically, while the Scribble complex (LET-413, LGL-1, and DLG-1) and PAR-1 kinase localize basolaterally. Lipids are also polarized, with phosphatidylinositol 4,5-bisphosphate [$PI(4,5)P_2$ or PIP_2] enriched apically and phosphatidylinositol 3,4,5-trisphosphate [$PI(3,4,5)P_3$ or PIP_3] basolaterally. This polarity is reinforced by polarity complexes, cytoskeletal organization, membrane trafficking, and junctional assemblies. (Armenti & Nance, 2012; Bossinger *et al*, 2004; Legouis *et al*, 2000; Winter *et al*, 2012). Membrane polarity is also essential for the spatial localization and activation of Rho family GTPases (Jaffe & Hall, 2005). In particular, CDC-42 activates the ARP-2/3 complex to promote branched actin polymerization, essential for the assembly and maintenance of apical microvilli (Bernadskaya *et al*, 2011; Ma *et al*, 1998; Martin-Belmonte *et al*, 2007; Shafaq-Zadah *et al*, 2012; Thuenauer *et al*, 2022). During infection many pathogens employ diverse virulence strategies to subvert the host polarity and actin cytoskeletal machinery to alter cellular shape and mechanics (Ruch & Engel, 2017). This facilitates intracellular entry and spread, or extracellular effector-induced inflammation and tissue damage (Bastounis *et al*, 2022; Colonne *et al*, 2016). Conversely, epithelial cells may actively reorganize actin as part of their defense response. Thus, plasma membrane polarity-actin cytoskeleton cross talk is vital for epithelial integrity during infection (Fleiszig *et al*, 1997; Tran *et al*, 2014).

The opportunistic pathogen, *Pseudomonas aeruginosa* a major cause of nosocomial infections and infects a wide range of hosts, including *C. elegans* (Balla & Troemel, 2013; Mahajan-Miklos *et al*, 1999; Tan *et al*, 1999a). In worms, *P. aeruginosa* infection induces diverse stress responses, mitotic quiescence, and apoptosis (Kuss-Duerkop & Keestra-Gounder, 2020; Pellegrino *et al*, 2014; Dunbar *et al*, 2012; McEwan *et al*, 2012; Bollen *et al*, 2024). In the absence of professional immune cells, *C. elegans* upon detecting infection activate innate immune signalling such as the TIR-1-p38 MAP kinase and the DAF-2/DAF-16 insulin/IGF-1 pathways (Peterson *et al*, 2023). DAF-2 dependent phosphorylation of PI3K/AGE-1 recruits AKT-1/AKT-2 kinases that in turn phosphorylate DAF-16. This kinase cascade results in cytoplasmic retention of DAF-16, thereby regulating gene expression (Ewbank, 2006). DAF-16 is known to induce expression of antimicrobial genes, detoxifying enzymes and bacterial avoidance (Balla

& Troemel, 2013; Cohen & Troemel, 2015; Irazoqui *et al*, 2010; Peterson *et al*, 2022;

Pukkila-Worley & Ausubel, 2012). Avoidance behaviour is also triggered by

mechanical cues, such as intestinal distension ("bloating"), mediated by TRPM

channels (Filipowicz *et al*, 2021; Kumar *et al*, 2019; Singh & Aballay, 2019b, 2019a).

Here, we investigated the mechanism underlying cytoskeletal remodelling and polarity

disruption following *P. aeruginosa* infection. We find that *P. aeruginosa* exploits PI3K-

AKT pathway to perturb CDC-42-ARP-2/3-mediated actin organization in C. elegans

enterocytes. The resultant disruption in epithelial polarity and loss of actin rich vesicles

from the apical surface locally alters the surface mechanics resulting in deforms

intestinal lumen surface contributing to host lethality.

RESULTS

***Pseudomonas aeruginosa* infection results in *C. elegans* intestinal deformation**

Consistent with previous studies, exposure to *P. aeruginosa* strain PA14 resulted in

reduced *C. elegans* survival (lifespan^{PA14} ~8 days; TD$_{50}$^{PA14} ~3 days) compared to *E.*

coli (OP50) (lifespan^{OP50} ~18 days; TD$_{50}$^{OP50} ~8 days) (Fig 1A). This lethality has been

partly attributed to the colonization of the intestinal lumen by *P. aeruginosa* (Tan *et al*,

1999a). To examine the impact *P. aeruginosa* infection on host cytoskeletal

organization, we imaged *C. elegans* intestines expressing mCherry fused to the N-

terminus of ACT-5, the intestine specific actin isoform (MacQueen *et al*, 2005b;

Szumowski *et al*, 2016). In *E. coli* fed worms, ACT-5, appeared as smooth parallel

lining along the intestinal lumen (Fig. 1B and 1C; blue arrowhead, OP50). Despite age

related intestinal distension (Egge *et al*, 2019; Irazoqui *et al*, 2010; Singh & Aballay,

2019a) the ACT-5 signal remained smooth even after 3 days on OP50 (Fig.1C; red

double-headed arrows in OP50/day-3, and S1A). Although lumen was distended

following *P. aeruginosa* exposure (Fig. 1C, red double-headed arrows in PA14 day-3,

and Fig S1A), apical surface characteristics were markedly distinct. In contrast to the

smooth surface seen in *E. coli*-fed animals, *P. aeruginosa*-fed worms displayed

extensive wrinkles and surface deformations (Fig. 1C; PA14,filled red arrowheads),

with vesicle-like structures budding outward from the deformed apical surface (Fig.

1C; PA14, red arrows in enlarged ROI). Whereas only ~4% (2/54) of *E. coli*-fed worms

showed such deformations after 1-day, ~48%(27/56) of *P. aeruginosa* -exposed

worms did (Fig. 1C). By day-3, the incidence rose to ~83% (43/52) in *P. aeruginosa-*

fed worms compared to ~19% (10/52) on *E. coli.*

Magnitude of intestinal deformation depends on the duration *of P. aeruginosa*

infection

To quantify changes in intestinal morphology, we defined a dimensionless "apical

deformation index" (δ_{apical}), as the ratio of apical lumen length to total worm length (Fig.

1E). *E. coli*-fed worms consistently maintained 'δ_{apical}^{OP50}=1', whereas P. aeruginosa

exposure resulted in 'δ_{apical}^{PA14}>1', with values increasing over time, indicating

progressive deformation (Fig. 1F). Tracking individual worms revealed that worms fed

with *E. coli* , the value of δ_{apical}^{OP50}~1, but in worms exposed to *P. aeruginosa*

deformation was evident by day 1 and worsened over subsequent days (Fig. 1G). In

acute infection assays, 12-hour exposure did not cause significant deformation,

whereas 24-hour exposure induced noticeable apical distortion even after transfer

back to *E.coli* (Fig. S1B). To test whether deformation extended beyond the apical

surface, we examined transgenic worms co-expressing LET-413::mCherry and DLG-

1::GFP, the Scribble and Discs Large orthologs in *C. elegans,* as markers for

basolateral surface and AJ, respectively (Fig. 5A) (Bossinger *et al*, 2004; Legouis *et*

al, 2000; McMahon *et al*, 2001; Riga *et al*, 2021). Neither AJ positioning nor basal

deformation index (δ_{basal}) was significantly altered by *P. aeruginosa* exposure (Fig. 5A

and S1C). However, lateral surfaces showed LET-413 enrichment and altered

morphology (Fig. 5A and 5C), indicating that deformation was restricted to apical and

lateral domains. Next we confirmed that the intestinal deformation was independent of

germline or uterine expansion by examining males, which lack these tissues (Fig. S1D

and S1E). F-actin staining using Phalloidin-AlexaFluor647 and LifeAct::mRuby

expression showed visible apical deformation specifically upon *P. aeruginosa*

exposure, eliminating the possibility that the infection phenotype might be an artifact

of mCherry::ACT-5 transgene expression. (Fig. S1F and S1H, red filled arrowheads).

Finally, we validated the phenotype using additional apical/sub-apical markers: PGP-

1::GFP, ERM-1::GFP, PAR-6::GFP, and IFB-2::GFP, that localized to apical

membrane, microvillar structure or subapical positions, respectively (Fig. 1H) (Geisler
et al, 2020; Göbel *et al*, 2004; Sepers *et al*, 2022). In all cases, *P. aeruginosa* exposure
resulted in clear deformation of the apical surface topology. 'Slow killing' assay,
typically uses nematode growth media (NGM) containing 0.35% peptone(Tan *et al*,
1999a). To minimize any confounding effects from elevated peptone concentrations
on intestinal epithelial response, we conducted our infections on standard NGM
containing 0.25% peptone, optimised for *C. elegans* growth (Stiernagle, 2006). We
observed intestinal deformation in worms infected on 0.35% peptone media
indistinguishable from 0.25% peptone (Fig. S1I and S1J). Our lifespan assays mirrored
previously reported kinetics using 0.35% peptone-containing NGM (Figure 1A) (Tan
et al, 1999a). Thus we confirmed that our infection conditions faithfully recapitulate
chronic *P. aeruginosa* infection, leading to progressive apical and lateral intestinal
deformation.

CDC-42 mediates *Pseudomonas*-induced intestinal deformation

RHO family GTPases are the key upstream regulators of F-actin assembly and
organization, that co-ordinate actin dynamics in a spatiotemporally controlled manner
(Ann Mack & Georgiou, 2014; Jaffe & Hall, 2005). *P. aeruginosa* infecting MDCK cells
disrupt RHO signalling through effector proteins that interfere with GTPase-activating
proteins (GAPs), leading to the disorganization of actin architecture (Cowell *et al*,
2005; Kazmierczak & Engel, 2002; Krall *et al*, 2000; Pederson *et al*, 1999; Sun &
Barbieri, 2004). To test whether *Pseudomonas* induced surface deformation in *C.*
elegans enterocytes might similarly depend on RHO GTPase-mediated actin re-
organization, we individually depleted RHO-1, CED-10, and CDC-42 in adult worms
expressing intestinal mCherry::ACT-5, and subsequently exposed them to *P.*
aeruginosa or *E. coli*. As in controls, exposure to *P. aeruginosa* resulted in apical
deformation in *rho-1(RNAi)* or *ced-10(RNAi)* animals (Fig. 2A, S2A and S2B). In
contrast, *cdc-42(RNAi)* worms exposed to *P. aeruginosa* displayed no deformation
(Fig. 2A and 2B). This lack of deformation was not due to impaired pathogen sensing
in *cdc-42(RNAi)* worms (Fig. S2D and S2E) (Estes *et al*, 2010). Consistent with its role
in supressing intestinal deformation, *cdc-42(RNAi)* worms exhibited significantly

extended total lifespan (8 days) compared to RNAi controls (4 days) upon *P.*
aeruginosa exposure (Fig. 2C).

205 *P. aeruginosa* infection leads to cytoplasmic accumulation of G-actin

The absence of intestinal deformation in *cdc-42(RNAi)* worms prompted us to
investigate the intracellular localization of CDC-42 and its downstream effector, the
actin nucleating ARP-2/3 complex, during *P. aeruginosa* infection. We used worms co-
expressing ARX-2::TagRFP (an essential subunit of the ARP-2/3 complex) at its
endogenous locus, and CDC-42::GFP, expressed exogenously under the native *cdc-*
42 promoter (Neukomm *et al*, 2014; Wu *et al*, 2017). In control animals maintained on
E. coli, both CDC-42 and ARX-2 localized predominantly to the apical membrane of
the enterocytes, with little accumulation in the cytoplasm over 2 days of exposure (Fig.
2D and 2G; OP50 panel and Fig. S2C). Upon exposure to *P. aeruginosa*, both
cytoplasmic and apical CDC-42 levels increased markedly (Fig. 2D enlarged ROI, 2E
and 2F). Cytoplasmic levels of ARX-2 also increased (~2.6 fold) (Fig. 2H) though
apical ARX-2 remained unchanged relative to *E. coli* (Fig. 2I). Intracellular actin
mirrored that of ARX-2, with ~2.5 fold increase in cytoplasmic levels but none in apical
levels (Fig. 2J- 2L). Surprisingly, Phalloidin-AlexaFluor647 staining and
LifeACT::mRuby did not show any cytoplasmic staining, suggesting that the enhanced
cytoplasmic actin pool following *P. aeruginosa* infection might represent filamentous
actin (Fig. S1F-S1H). Thus, the increased cytoplasmic levels following *P. aeruginosa*
infection reflects accumulation of G-actin rather than loss of F-actin from the apical
surface.

226 *P. aeruginosa* induced disruption of cortical actin results in apical surface
227 deformation

It seemed counterintuitive that apical ARX-2 and ACT-5 intensities did not increase
despite CDC-42 enrichment. To clarify this conundrum, we measured apical levels of
ACT-5 and ARX-2 in *cdc-42(RNAi)* worms, and found them to be significantly reduced
(see OP50 panels in Figs. 3A, S3A, S3B and S3D). Thus apical actin and ARP-2/3

complex are sensitive to CDC-42-depletion, but not to *P. aeruginosa* induced CDC-
42-enrichment. The absence of deformation in *cdc-42(RNAi)* animals could therefore
reflect a direct effect of CDC-42 depletion or an indirect effect via reduced ARP-2/3-
mediated F-actin polymerization. To distinguish these, we depleted ACT-5 or ARX-2
in worms co-expressing CDC-42::GFP and ARX-2::TagRFP. Due to *act-5(RNAi)-*
associated developmental defects, analyses were restricted to day 1 post-infection.
Strikingly, *act-5(RNAi)* and *arx-2(RNAi)* worms, like *cdc-42(RNAi)* showed rescue of
deformation phenotype following infection despite elevated CDC-42 levels (red open
arrowheads in Fig.3C-E,3G and S3E).

To probe the function of ARP-2/3 dependent actin polymerization in *P. aeruginosa*
induced intestinal deformation, we analysed worms co-expressing mCherry::ACT5
and an apical membrane maker, PGP-1::GFP. In control worms fed with *E. coli*, ACT-
5 and PGP-1 were uniformly co-localized at apical surface (Fig. 3F; control panel). In
arx-2(RNAi) intestines, apical actin was absent and instead appeared as cytoplasmic
aggregates, even when fed with *E. coli* (Fig. 3F; *arx-2(RNAi)* red filled arrowheads).
This ruled out cytoplasmic actin accumulation as a possible cause of *P. aeruginosa*-
induced deformation. ARP-2/3 disruption has been implicated in cytoplasmic
mislocalization of membrane bound components such as DLG-1 and ERM-1
(Bernadskaya *et al*, 2011). PGP-1 displayed diffused apical localization in *arx-2(RNAi)*
intestines (Fig. 3F; *arx-2(RNAi)* red hollow arrowheads). These results suggested the
central role of ARP-2/3 polymerized actin in maintenance of apical membrane
organization. Importantly, CDC-42 and ARX-2 maintained apical localization even in
the absence of actin (Fig. 3F and 3G), confirming their upstream role in the actin
polymerization pathway. Collectively, these results demonstrate that rescue of apical
deformation in *cdc-42(RNAi)* and *arx-2(RNAi)* intestines following *P. aeruginosa*
exposure was due to disruption of ARP-2/3 mediated actin polymerization.

Fragmented ARP-2/3 clustering and F-actin disorganization drives release of
membrane vesicles into the intestinal lumen following *P. aeruginosa* exposure.

Although overall intensities of apical ARX-2 and ACT-5 were comparable between
worms exposed to *P. aeruginosa* and *E. coli* (Fig. 2I and 2L), their surface localization

patterns were markedly distinct. In *E. coli*-fed worms, ARX-2 was evenly distributed at

the apical surface (Fig. 4A; OP50, cyan arrowhead) essential for contiguous actin

polymerization, supporting brush border integrity (Bidaud-Meynard *et al*, 2021;

MacQueen *et al*, 2005b; Stutz *et al*, 2015). In contrast, *P. aeruginosa*-infected animals

exhibited fragmented ARX-2 clusters, interspersed with ARX-2-depleted regions (Fig.

4A; PA14, white arrows). We sought to correlate the cortical actin distribution with this

fragmented ARP-2/3 localization using worms co-expressing mCherry::ACT-5 and

ARX-2::GFP. Medial confocal sections of intestines from this strain when exposed to

P. aeruginosa resulted in apical actin to protrude as a 'bud-like' structure into the

lumen, emerging from ARX-2-free apical surface (Fig 4B; actin protrusions - yellow

arrowheads, fragmented ARP-2/3 – cyan arrows and Fig. 1C; red arrows in enlarged

ROI). By day-2, these protrusions pinched off as vesicles, coinciding with apical

surface deformation (Fig. 1C; day-2 PA14 panel - red arrows in enlarged ROI and Fig.

4D). Lattice light sheet microscopy of the intestinal lumen surface of animals

expressing mCherry::ACT-5 exposed to *P. aeruginosa* expressing GFP revealed

extracellular pinching of actin rich vesicles from the apical surface into the lumen (Fig.

4C; enlarged ROI, cyan arrows).

To determine whether these extracellular actin structures were membrane-enclosed,

we examined worms co-expressing mCherry::ACT-5 and PGP-1::GFP. Medial

confocal images on day-2 post-infection revealed apical buds at various stages: initial

protrusion (Fig. 4D; ROIs 1a & 2a), membrane pinching (1b & 2b), and free vesicles

in the lumen (cyan arrows, 1c & 2c). Intensity line-scan analysis confirmed that these

vesicles consisted of cortical actin cores surrounded by PGP-1::GFP-labelled apical

membrane. LifeACT::mRuby labeled these vesicles, indicating filamentous actin

content (Fig. 4E). Diameter of actin containing vesicles was $\sim 0.59\,\mu$m (Fig. 4E). These

results support the view that *P. aeruginosa* exposure led to disordered ARP-2/3

localization at the apical lumen surface destabilising cortical actin organization,

resulting in the release of F-actin rich vesicles from the apical surface into lumen.

P. aeruginosa infection disrupts apical polarity of enterocyte epithelium

Fragmented ARP-2/3 clustering and vesicle shedding suggested a loss of apical

membrane identity. *P. aeruginosa* attachment to MDCK cells triggers a transient

PI(3,4,5)P$_3$ accumulation at the apical surface resulting in the loss of apical membrane

identity (Kierbel *et al*, 2007, 2005; Thuenauer *et al*, 2022). To check if *P. aeruginosa*

might cause a similar disruption to the polarity in *C. elegans* intestinal epithelium, we

analysed the localization of the *C. elegans* Scribble homolog, LET-413 and the cell

junction marker, DLG-1. The LET-413 complex and DLG-1 have been reported to be

essential for maintenance of basolateral domains in *C. elegans* by regulating the

localization of apical polarity components (Bossinger *et al*, 2004; McMahon *et al*,

2001). In *E. coli*-fed worms, the localization of LET-413 was limited to the basolateral

membrane with the apical surface largely devoid of any LET-413 (Fig. 5A; OP50 panel,

basolateral surface -yellow filled arrowhead, apical surface -yellow asterisks) (Legouis

et al, 2000). Upon exposure to *P. aeruginosa,* we observed a significant increase in

LET-413 levels along the apical and lateral membranes (Fig. 5A; PA14 panel, apical

surface - yellow arrows, lateral membrane- yellow open arrowhead, 5B and 5C),

indicating disruption in cell polarity. *P. aeruginosa* infection did not affect the position

of cell junctions (AJ)(Fig. 5A; cyan arrows).

The absence of intestinal deformation in *cdc-42(RNAi)* animals prompted us to

investigate if CDC-42 might also regulate apico-lateral polarity following *P. aeruginosa*

exposure. To this end, we checked the localization of LET-413 in control (L4440) and

cdc-42(RNAi) worms. The enhanced apical localization of LET-413 in the intestines of

control (L4440) worms was significantly diminished in *cdc-42(RNAi)* animals following

P. aeruginosa exposure (Fig. 5D). Conversely, we did not detect any disturbance to

the apical localization of CDC-42 and ARP-2/3 in *let-413(RNAi)* intestines, in

agreement with previous reports indicating that LET-413 depletion has minimal effect

on epithelial polarity (Fig S5A) (Riga *et al*, 2021). Moreover, similar to control animals,

P. aeruginosa exposure to *let-413(RNAi)* worms resulted in apical surface deformation

as well as elevated cytoplasmic levels of CDC-42 (S5A and S5B). Taken together,

these results suggest that upon exposure to *P. aeruginosa,* the apical surface *C.*

elegans intestine acquires basolateral characteristics in a CDC-42-dependent

mechanism.

PI3K-AKT pathway depletion abrogates *P. aeruginosa* induced CDC-42 mis-localization and intestinal deformation

Disruption of the PI3K-AKT pathway impedes *P. aeruginosa* infection in MDCK cells (Kierbel *et al*, 2005). Similarly, long-lived *C. elegans* mutants such as *akt-1*, *akt-2* and *age-1*/PI3K exhibit resistance to *P. aeruginosa* (Evans *et al*, 2008; Garsin *et al*, 2003). Our observation that *P. aeruginosa* disrupted intestinal polarity in *C. elegans* suggested that host response to *P. aeruginosa* infection could be conserved between mammalian epithelium and *C. elegans* enterocytes. To examine the conservation in cellular response to *P. aeruginosa* between mammmals and *C. elegans*, we diminished PI3K-AKT signalling by a simultaneous depletion of AKT-1 and AKT-2 using RNAi and checked the localizations of LET-413, CDC-42 and ARX-2 in the intestines of *C elegans*. Depletion of AKT-1/AKT-2, like in *cdc-42(RNAi)* worms, prevented the apical localization of LET-413 following *P. aeruginosa* infection (Fig. 5D). Evidence for crosstalk between CDC-42 dependent actin assembly and PI3K-AKT in mammalian cells prompted us to measure CDC-42 and ARX-2 levels in the intestines of *akt-1(RNAi)+akt-2(RNAi)* and *age-1(RNAi)* animals following *P. aeruginosa* exposure (McCormick *et al*, 2019; Yang *et al*, 2012). Intestinal epithelium of animals depleted of AKT or AGE-1 displayed diminished apical and cytosolic ARX-2 localization compared to control animals (Fig. 5E, S5E - S5H), and also did not display apical deformation following *P. aeruginosa* exposure (Fig. 5D). Taken together our results suggest that PI3K activation induced apical enrichment of PI(3,4,5)P$_3$ following exposure to *P. aeruginosa*, disrupts apical membrane polarity and de-localizes ARP-2/3 dependent actin assembly from the apical surface resulting in intestinal deformation. These results re-enforces the importance of polarity disruption as a key event following *P. aeruginosa* infection and prior to intestinal deformation.

Extracellular localization of *Pseudomonas* is sufficient to induce intestinal deformation

Pseudomonas aeruginosa functions both as an extracellular and intracellular pathogen infecting wide range of hosts (Resko *et al*, 2024). Transient activation of

PI3K pathway resulting in apical membrane acquiring basolateral characteristics was
resulted in internalization of *P. aeruginosa* by MDCK cells (Kierbel *et al*, 2005).
Electron microscopy studies of *P. aeruginosa* -infected *C. elegans* intestine have also
reported *Pseudomonas* invasion into enterocytes (Irazoqui *et al*, 2010; Xue *et al*,
2024).   To determine whether intestinal deformation in *C. elegans* results from *P.
aeruginosa* breaching the epithelial barrier, we performed live confocal imaging of
infection using GFP-expressing *E. coli* or *P. aeruginosa*. Unlike E. coli (OP50-GFP),
which were in very few numbers even after two days of exposure, *P. aeruginosa*
(PA14-GFP) cells formed visible aggregates throughout the intestinal lumen as early
as one-day post infection (Fig. 6A). Interestingly, apical surfaces adjacent to these
aggregates displayed mild contortions, which progressed into pronounced
deformation by day 2 (Fig. 6A; red triangles in bottom row-PA14). Orthogonal
reconstruction (YZ plane) of confocal (XY) planes revealed that PA14-GFP localization
remained restricted to the lumen with no detectable signal in the cytoplasm of the
intestinal cells (Fig.6A; PA14 panel). These observations were confirmed using lattice
light-sheet microscopy (Fig. 4C and Supplementary Movie S1). To test whether apical
surface deformation is a general enterocyte response to any pathogen, we conducted
parallel infection assays using *Salmonella enterica* serovar *typhimurium* (Stm), a
Gram-negative bacterium known to modulate CDC-42 activity during mammalian
infections (Bandyopadhyay *et al*, 2024) and cause lethality in *C. elegans (*Fig. S6A)
(Aballay *et al*, 2000; Desai *et al*, 2019). Stm-GFP infection in worms expressing
mCherry::ACT-5, resulted in efficient intestinal colonization by day 2 similar to *P.
aeruginosa* (Fig. S5B). However, despite this robust lumen colonization, in contrast to
P. aeruginosa , Stm-exposed intestines did not display any surface deformation (Fig.
6A; middle row, yellow open arrowheads). Taken together these results indicate that
the intestinal deformation is a pathogen specific response by the *C. elegans*
enterocyte to extracellular *Pseudomonas* infection.

Physical association between live *P .aeruginosa* and the intestinal surface is not essential for apical deformation

Given the key role for physical association with host cells in *P. aeruginosa*

pathogenesis (via various adhesins, including flagella, type IV pili, type III secretion

system (T3SS), and lectins (Bucior *et al*, 2012; Comolli *et al*, 1999; Thuenauer *et al*,

2022), we asked whether a direct attachment of *P. aeruginosa* to the apical membrane

of *C. elegans* intestine was necessary to trigger tissue deformation. First, we tested if

exposure to live microbes is essential for inducing deformation of apical surface, by

exposing animals to heat-killed *E. coli* or *P. aeruginosa* instead of live bacteria. Since

worms displayed no noticeable defects in growth, maturation or egg laying, we

concluded that heat-killed bacteria as food source was not nutritionally limiting.

However, unlike live *P. aeruginosa*, heat killed *P. aeruginosa* failed to induce intestinal

deformation (Fig. 6B; open triangle). Our observation agrees with previous reports

showing heat-killed *P. aeruginosa* does not elicit immune response in *C.* elegans

(Irazoqui *et al*, 2010). Since extracellular *P. aeruginosa* induces intestinal deformation,

we hypothesized that secreted factors may mediate this effect. To test this, we

exposed worms to cell-free filter sterilized supernatant collected from *P. aeruginosa* or

E. coli cultures. Remarkably, a 12-hour exposure to *P. aeruginosa* supernatant alone

was sufficient to trigger apical surface deformation in the *C. elegans* intestine (Fig.

6C).

The global response regulator, GacA governs the expression of numerous virulence

factors secreted by *P. aeruginosa,* including toxins and components in biofilm

regulation (Barta *et al*, 1992; Kitten *et al*, 1998; Parkins *et al*, 2001; Reimmann *et al*,

1997) and has also been implicated in inducing developmental defects and activating

the unfolded protein response (UPR) in *C. elegans* intestinal epithelium (Pellegrino *et*

al, 2014). To assess the role of GacA in apical surface deformation, we exposed

mCherry::ACT-5-expressing worms to *E. coli*, *P. aeruginosa* or a *gacA* deletion mutant

(PA14*ΔgacA*) strain. The intestines of worms exposed to PA14*ΔgacA* exhibited an

intermediate level of intestinal deformation compared to those infected with wild-type

PA14 (Fig. 6D and 6E). Importantly, a *gacA* deletion mutant (PA14*ΔgacA*) retains

partial pathogenicity, causing ~50% lethality in infected worms (Fig. 6F) (Feinbaum *et*

al, 2012; Tan *et al*, 1999b). These results suggest that while GacA partially contributes

to *P. aeruginosa* pathogenesis in *C. elegans*, a GacA-independent mechanism also

exists, through which the pathogen elicits tissue surface deformation. Taken together,

these results support the view that while intestinal distension is a general tissue

response to microbial colonization of the lumen, mechanical deformation of apical

surface is a specific response to extracellular heat labile component from *P.*

aeruginosa.

DISCUSSION

Here we report that *P. aeruginosa* exposure induces two distinct morphological

alterations in the *C. elegans* intestine; (a) lumenal distension, reflected by increased

intestinal width, and (b) deformation of the apical surface, seen as wrinkling of the

lumen. While lumen distension has been previously been described as a general

response to microbial colonisation, we show that apical surface deformation is specific

to *P. aeruginosa* infection.

P. aeruginosa infection in MDCK cells recruits PI3K to the attachment sites, causing

PIP_3 enrichment, apical mislocalization of basolateral proteins, and disruption of

epithelial polarity through perturbation of the Rho GTPases - RHO-1, RAC-1 and CDC-

42 (Aktories, 2011; Kazmierczak & Engel, 2002; Kierbel *et al*, 2007; Krall *et al*, 2000).

Consistent with this, we find that inhibition of either PI3K-AKT signalling or CDC-42-

dependent actin assembly completely abolished both intestinal deformation and

pathogen-induced lethality. *P. aeruginosa* infection also led to apical mis-localization

of the basolateral marker LET-413, indicating a loss of apical membrane identity,

though no apical markers were mislocalized baso-laterally. The rescue observed in

cdc-42(RNAi) animals further implicated CDC-42 in pathogen-induced actin

remodelling. However, depletion of the downstream polarity facrot, LET-413 failed to

rescue deformation, suggesting that CDC-42 contributes primarily via actin assembly

rather than polarity regulation. Supporting this, depletion of ACT-5 or disruption of

ARP-2/3 phenocopied *cdc-42(RNAi)*. In line with previous reports (Kierbel *et al*, 2005;

Paradis & Ruvkun, 1998), we noted that attenuation of PI3K-AKT signalling rescued

the intestinal deformation by limiting disruption in CDC-42 dependent ARP-2/3 mis-

localization. These results highlights the central role of CDC-42-ARP-2/3-mediated

actin polymerization in *P. aeruginosa* induced deformation.

In uninfected animals, ARP-2/3, together with actin-binding proteins such as PLST-1
and FLN-2, forms evenly distributed apical clusters (Bidaud-Meynard *et al*, 2021).
ARP-2/3 polymerizes contiguous F-actin to support microvillar structures (Fig. 4A).
Infection by *P. aeruginosa* fragmented these clusters, disrupting local actin
organization, destabilizing the apical surface, which promoted vesiculation of actin-
rich membrane into the lumen. Because actin structures in enterocytes show minimal
mobility, their disruption cannot be rapidly replenished, destabilizing the surface
(Bernadskaya *et al*, 2011; Bidaud-Meynard *et al*, 2021) (Fig. 6G). This agrees with
electron micrographs showing actin protrusions in MDCK cells (Kierbel *et al*, 2005) as
well as microvillar loss and accumulation of host debris in the *C. elegans* lumen during
infection (Irazoqui *et al*, 2010; Xue *et al*, 2024). Intriguingly, depletion of ARX-2 or
ACT-5 prevented vesicle release and surface deformation. We speculate that a
uniform depletion of ARP-2/3 or actin reduces actin availability for vesiculation. Similar
protection has been reported in MDCK cells, where Latrunculin A mediated actin
depolymerization attenuated *P. aeruginosa* infection by upto 60-fold (Kazmierczak *et*
al, 2004). CDC-42 upregulation and activation have been reported in polarized MDCK
cells (Kazmierczak *et al*, 2004). Consistent with these findings, *P. aeruginosa* infection
in *C. elegans* enterocytes led to cytoplasmic accumulation of CDC-42, ARP-2/3 and
actin. While RHO-mediated actin regulation and homeostasis are monitored in
mammals by pyrin- and NOD-dependent pathways, no equivalent surveillance been
reported in *C. elegans* (Mostowy & Shenoy, 2015). Although cytoplasmic actin
accumulation correlated with infection, it did not directly cause deformation. Whether
it may be sensed as a pattern of pathogenesis is remains to be addressed. Moreover,
MAPK signalling remained intact in *cdc-42(RNAi)* animals following infection,
indicating that CDC-42 depletion does not impair microbial sensing.

Similar to polarized MDCK cells*, P. aeruginosa* does not breach the cell membrane
of *C. elegans* epithelium. Additionally, while the surface deformation phenotype could
be recapitulated by exposure to cell-free supernatant from *P . aeruginosa* growth
culture but not by heat-killed bacteria suggested the role of a heat-labile factor. Given
the extracellular lifestyle of *P. aeruginosa* in polarized epithelia, extracellular
membrane vesicles (OMVs) rich in surface components, quorum-sensing molecules ,

toxins and other virulence factors, are plausible mediators of deformation signal
(Bauman & Kuehn, 2009, 2006; Kuehn & Kesty, 2005; Mashburn & Whiteley, 2005).

Pseudomonas is generally considered as an extracellular pathogen and its
intracellular lifestyle is usually associated with tissue injury. While *P. aeruginosa* has
been shown to infiltrate the cytoplasmic space many host systems, we have not
observed epithelial breach in our experiments in *C. elegans* intestine. We demonstrate
that extracellular *P. aeruginosa* exploits PI3K-AKT activation to perturb CDC-42-ARP-
2/3-mediated actin organization leading to loss of apical polarity and cytoskeletal
integrity in the *C. elegans* enterocytes. Our findings reveal an evolutionarily conserved
strategy by which *P. aeruginosa* subverts actin regulation during extracellular infection.
We propose that the pathogen mediated disruption of polarity impairs apically directed
transport and secretion of antimicrobial peptides, and surface binding lectins. An
attenuated antimicrobial response could allow bacterial proliferation in the lumen and
a rapid spreading of the bacteria in the population.

ACKNOWLEDGEMENTS

This work was supported by DBT-Wellcome India Alliance Fellowship
(IA/I/18/1/503624) and ANRF Core research grant (CRG/2023/004638) to A.P,
Department of Biotechnology Junior research Fellowship (DBT/2024-
25/AshokaUni/2486) to T.S., and core funding support from the Trivedi School of
Biosciences, Ashoka University. We acknowledge the infrastructure support from the
Central Bio-imaging facility and Ashoka-Zeiss Core Imaging Facility at Ashoka
University. We thank Ms. Megha Rai for help with vesicle size quantification, Ms.
Shraddha Nirmal for assisting with generating strains for some of the experiments and
Ms. Purna Pardeshi for some of the schematic images in this manuscript. We are
grateful to the labs of Arnab Mukhopadhyay, Anindya Ghosh-Roy, Emily Troemel,
Grégoire Michaux, Jogender Singh, Mike Boxem, Mahak Sharma and Varsha Singh,
for sharing *C. elegans* and bacterial strains. Some strains were provided by the CGC,
which is funded by NIH Office of Research Infrastructure Programs (P40OD010440).
We thank Jogender Singh for discussions.

AUTHOR CONTRIBUTIONS**:** D.A.N and A.P conceived the project. D.A.N, T.S and A.P designed the
experiments. D.A.N and T.S performed experiments. D.A.N and T.S analysed the data with input from
A.P. A.P wrote the manuscript with input from D.A.N and T.S.

References:

Aballay A, Yorgey P & Ausubel FM (2000) Salmonella typhimurium proliferates and
517   establishes a persistent infection in the intestine of Caenorhabditis elegans.
518   *Curr Biol* 10: 1539–1542

519   Aktories K (2011) Bacterial protein toxins that modify host regulatory GTPases. *Nat Rev*
520   *Microbiol* 9: 487–498

521   Ann Mack N & Georgiou M (2014) The interdependence of the Rho GTPases and
522   apicobasal cell polarity. *Small GTPases* 5: 37–41

523   Armenti ST & Nance J (2012) Adherens junctions in C. elegans embryonic
524   morphogenesis. *Subcell Biochem* 60: 279–299

525   Balla KM & Troemel ER (2013) Caenorhabditis elegans as a model for intracellular
526   pathogen infection. *Cell Microbiol* 15: 1313–1322

527   Bandyopadhyay S, Zhang X, Ascura A, Edelblum KL, Bonder EM & Gao N (2024)
528   Salmonella engages CDC42 effector protein 1 for intracellular invasion. *J Cell*
529   *Physiol* 239

530   Barta TM, Kinscherf TG & Willis DK (1992) Regulation of tabtoxin production by the lemA
531   gene in Pseudomonas syringae. *J Bacteriol* 174: 3021–3029

532   Bastounis EE, Radhakrishnan P, Prinz CK & Theriot JA (2022) Mechanical Forces Govern
533   Interactions of Host Cells with Intracellular Bacterial Pathogens. *Microbiol Mol*
534   *Biol Rev* 86: e00094-20

535   Bauman SJ & Kuehn MJ (2006) Purification of outer membrane vesicles from
536   Pseudomonas aeruginosa and their activation of an IL-8 response. *Microbes*
537   *Infect* 8: 2400–2408

538   Bauman SJ & Kuehn MJ (2009) Pseudomonas aeruginosa vesicles associate with and
539   are internalized by human lung epithelial cells. *BMC Microbiol* 9: 26

540   Berg M, Stenuit B, Ho J, Wang A, Parke C, Knight M, Alvarez-Cohen L & Shapira M (2016)
541   Assembly of the *Caenorhabditis elegans* gut microbiota from diverse soil
542   microbial environments. *ISME J* 10: 1998–2009

543   Bernadskaya YY, Patel FB, Hsu HT & Soto MC (2011) Arp2/3 promotes junction
544   formation and maintenance in the Caenorhabditis elegans intestine by
545   regulating membrane association of apical proteins. *Mol Biol Cell* 22: 2886–2899

546   Bidaud-Meynard A, Demouchy F, Nicolle O, Pacquelet A, Suman SK, Plancke CN, Robin
547   FB & Michaux G (2021) High-resolution dynamic mapping of the *C. elegans*
548   intestinal brush border. *Development* 148

Bollen DP, Reddy KC, Lascarez-Lagunas LI, Kim DH & Colaiácovo MP (2024) Germline
mitotic quiescence and cell death are induced in *Caenorhabditis elegans* by
exposure to pathogenic *Pseudomonas aeruginosa*. *GENETICS* 226: iyad197

Bossinger O, Fukushige T, Claeys M, Borgonie G & McGhee JD (2004) The apical
disposition of the Caenorhabditis elegans intestinal terminal web is maintained
by LET-413. *Dev Biol* 268: 448–456

Bucior I, Pielage JF & Engel JN (2012) Pseudomonas aeruginosa Pili and Flagella Mediate
Distinct Binding and Signaling Events at the Apical and Basolateral Surface of
Airway Epithelium. *PLoS Pathog* 8: e1002616

Cohen LB & Troemel ER (2015) Microbial pathogenesis and host defense in the
nematode C. elegans. *Curr Opin Microbiol* 23: 94–101

Colonne PM, Winchell CG & Voth DE (2016) Hijacking Host Cell Highways: Manipulation
of the Host Actin Cytoskeleton by Obligate Intracellular Bacterial Pathogens.
Front Cell Infect Microbiol 6

Comolli JC, Waite LL, Mostov KE & Engel JN (1999) Pili Binding to Asialo-GM1 on
Epithelial Cells Can Mediate Cytotoxicity or Bacterial Internalization by
Pseudomonas aeruginosa. *Infect Immun* 67: 3207–3214

Cowell BA, Evans DJ & Fleiszig SMJ (2005) Actin cytoskeleton disruption by ExoY and its
effects on *Pseudomonas aeruginosa* invasion. *FEMS Microbiol Lett* 250: 71–76

Desai SK, Padmanabhan A, Harshe S, Zaidel-Bar R & Kenney LJ (2019) Salmonella
biofilms program innate immunity for persistence in Caenorhabditis elegans.
Proc Natl Acad Sci U S A 116: 12462–12467

Dimov I & Maduro MF (2019) The C. elegans intestine: organogenesis, digestion, and
physiology. *Cell Tissue Res* 377: 383–396

Dunbar TL, Yan Z, Balla KM, Smelkinson MG & Troemel ER (2012) C. elegans detects
pathogen-induced translational inhibition to activate immune signaling. *Cell
Host Microbe* 11: 375–386

Egge N, Arneaud SLB, Wales P, Mihelakis M, McClendon J, Fonseca RS, Savelle C,
Gonzalez I, Ghorashi A, Yadavalli S, *et al* (2019) Age-Onset Phosphorylation of a
Minor Actin Variant Promotes Intestinal Barrier Dysfunction. *Dev Cell* 51: 587-
601.e7

Estes KA, Dunbar TL, Powell JR, Ausubel FM & Troemel ER (2010) bZIP transcription
factor zip-2 mediates an early response to Pseudomonas aeruginosa infection in
Caenorhabditis elegans. *Proc Natl Acad Sci U S A* 107: 2153–2158

Evans EA, Chen WC & Tan M (2008) The DAF-2 insulin-like signaling pathway
independently regulates aging and immunity in *C. elegans*. *Aging Cell* 7: 879–893

Ewbank J (2006) Signaling in the immune response. *WormBook*

Feinbaum RL, Urbach JM, Liberati NT, Djonovic S, Adonizio A, Carvunis A-R & Ausubel FM (2012) Genome-Wide Identification of Pseudomonas aeruginosa Virulence-Related Genes Using a Caenorhabditis elegans Infection Model. *PLoS Pathog* 8: e1002813

Filipowicz A, Lalsiamthara J & Aballay A (2021) TRPM channels mediate learned pathogen avoidance following intestinal distention. *eLife* 10: e65935

Fleiszig SM, Evans DJ, Do N, Vallas V, Shin S & Mostov KE (1997) Epithelial cell polarity affects susceptibility to Pseudomonas aeruginosa invasion and cytotoxicity. *Infect Immun* 65: 2861–2867

Garsin DA, Villanueva JM, Begun J, Kim DH, Sifri CD, Calderwood SB, Ruvkun G & Ausubel FM (2003) Long-Lived *C. elegans daf-2* Mutants Are Resistant to Bacterial Pathogens. *Science* 300: 1921–1921

Geisler F, Coch RA, Richardson C, Goldberg M, Bevilacqua C, Prevedel R & Leube RE (2020) Intestinal intermediate filament polypeptides in C. elegans: Common and isotype-specific contributions to intestinal ultrastructure and function. *Sci Rep* 10: 3142

Göbel V, Barrett PL, Hall DH & Fleming JT (2004) Lumen morphogenesis in C. elegans requires the membrane-cytoskeleton linker erm-1. *Dev Cell* 6: 865–873

Gravato-Nobre MJ, Nicholas HR, Nijland R, O'Rourke D, Whittington DE, Yook KJ & Hodgkin J (2005) Multiple genes affect sensitivity of Caenorhabditis elegans to the bacterial pathogen Microbacterium nematophilum. *Genetics* 171: 1033–1045

Irazoqui JE, Troemel ER, Feinbaum RL, Luhachack LG, Cezairliyan BO & Ausubel FM (2010) Distinct Pathogenesis and Host Responses during Infection of C. elegans by P. aeruginosa and S. aureus. *PLoS Pathog* 6: e1000982

Jaffe AB & Hall A (2005) Rho GTPases: Biochemistry and biology. *Annu Rev Cell Dev Biol* 21: 247–269

Kazmierczak BI & Engel JN (2002) *Pseudomonas aeruginosa* ExoT Acts In Vivo as a GTPase-Activating Protein for RhoA, Rac1, and Cdc42. *Infect Immun* 70: 2198–2205

Kazmierczak BI, Mostov K & Engel JN (2004) Epithelial Cell Polarity Alters Rho-GTPase Responses to *Pseudomonas aeruginosa*. *Mol Biol Cell* 15: 411–419

Kierbel A, Gassama-Diagne A, Mostov K & Engel JN (2005) The Phosphoinositol-3-Kinase–Protein Kinase B/Akt Pathway Is Critical for *Pseudomonas aeruginosa* Strain PAK Internalization. *Mol Biol Cell* 16: 2577–2585

Kierbel A, Gassama-Diagne A, Rocha C, Radoshevich L, Olson J, Mostov K & Engel J (2007) *Pseudomonas aeruginosa* exploits a PIP3-dependent pathway to transform apical into basolateral membrane. *J Cell Biol* 177: 21–27

Kitten T, Kinscherf TG, McEvoy JL & Willis DK (1998) A newly identified regulator is
required for virulence and toxin production in Pseudomonas syringae. *Mol*
Microbiol 28: 917–929

Krall R, Schmidt G, Aktories K & Barbieri JT (2000) Pseudomonas aeruginosa ExoT is a
Rho GTPase-activating protein. *Infect Immun* 68: 6066–6068

Kuehn MJ & Kesty NC (2005) Bacterial outer membrane vesicles and the host–pathogen
interaction. *Genes Dev* 19: 2645–2655

Kumar S, Egan BM, Kocsisova Z, Schneider DL, Murphy JT, Diwan A & Kornfeld K (2019)
Lifespan Extension in C. elegans Caused by Bacterial Colonization of the
Intestine and Subsequent Activation of an Innate Immune Response. *Dev Cell*
49: 100-117.e6

Kuss-Duerkop SK & Keestra-Gounder AM (2020) NOD1 and NOD2 Activation by Diverse
Stimuli: a Possible Role for Sensing Pathogen-Induced Endoplasmic Reticulum
Stress. *Infect Immun* 88: e00898-19

Legouis R, Gansmuller A, Sookhareea S, Bosher JM, Baillie DL & Labouesse M (2000)
LET-413 is a basolateral protein required for the assembly of adherens junctions
in Caenorhabditis elegans. *Nat Cell Biol* 2: 415–422

Lei M, Tan Y, Tu H & Tan W (2024) Neuronal basis and diverse mechanisms of pathogen
avoidance in Caenorhabditis elegans. *Front Immunol* 15: 1353747

Ma L, Rohatgi R & Kirschner MW (1998) The Arp2/3 complex mediates actin
polymerization induced by the small GTP-binding protein Cdc42. *Proc Natl Acad*
Sci U S A 95: 15362–15367

MacQueen AJ, Baggett JJ, Perumov N, Bauer RA, Januszewski T, Schriefer L & Waddle JA
(2005a) ACT-5 Is an Essential *Caenorhabditis elegans* Actin Required for
Intestinal Microvilli Formation. *Mol Biol Cell* 16: 3247–3259

MacQueen AJ, Baggett JJ, Perumov N, Bauer RA, Januszewski T, Schriefer L & Waddle JA
(2005b) ACT-5 Is an Essential *Caenorhabditis elegans* Actin Required for
Intestinal Microvilli Formation. *Mol Biol Cell* 16: 3247–3259

Mahajan-Miklos S, Tan M-W, Rahme LG & Ausubel FM (1999) Molecular Mechanisms of
Bacterial Virulence Elucidated Using a Pseudomonas aeruginosa–
Caenorhabditis elegans Pathogenesis Model. *Cell* 96: 47–56

Martin-Belmonte F, Gassama A, Datta A, Yu W, Rescher U, Gerke V & Mostov K (2007)
PTEN-Mediated Apical Segregation of Phosphoinositides Controls Epithelial
Morphogenesis through Cdc42. *Cell* 128: 383–397

Mashburn LM & Whiteley M (2005) Membrane vesicles traffic signals and facilitate group
activities in a prokaryote. *Nature* 437: 422–425

McCormick B, Chu JY & Vermeren S (2019) Cross-talk between Rho GTPases and PI3K in
the neutrophil. *Small GTPases* 10: 187–195

McEwan DL, Kirienko NV & Ausubel FM (2012) Host translational inhibition by
Pseudomonas aeruginosa exotoxin A triggers an immune response in
Caenorhabditis elegans. *Cell Host Microbe* 11: 364–374

McGhee J (2007) The C. elegans intestine. *WormBook*

McMahon L, Legouis R, Vonesch JL & Labouesse M (2001) Assembly of C. elegans apical
junctions involves positioning and compaction by LET-413 and protein
aggregation by the MAGUK protein DLG-1. *J Cell Sci* 114: 2265–2277

Melo JA & Ruvkun G (2012) Inactivation of conserved C. elegans genes engages
pathogen- and xenobiotic-associated defenses. *Cell* 149: 452–466

Mostowy S & Shenoy AR (2015) The cytoskeleton in cell-autonomous immunity:
structural determinants of host defence. *Nat Rev Immunol* 15: 559–573

Neukomm LJ, Zeng S, Frei AP, Huegli PA & Hengartner MO (2014) Small GTPase CDC-42
promotes apoptotic cell corpse clearance in response to PAT-2 and CED-1 in C.
elegans. *Cell Death Differ* 21: 845–853

Paradis S & Ruvkun G (1998) *Caenorhabditis elegans* Akt/PKB transduces insulin
receptor-like signals from AGE-1 PI3 kinase to the DAF-16 transcription factor.
Genes Dev 12: 2488–2498

Parkins MD, Ceri H & Storey DG (2001) Pseudomonas aeruginosa GacA, a factor in
multihost virulence, is also essential for biofilm formation. *Mol Microbiol* 40:
1215–1226

Pederson KJ, Vallis AJ, Aktories K, Frank DW & Barbieri JT (1999) The amino-terminal
domain of *Pseudomonas aeruginosa* ExoS disrupts actin filaments via small-
molecular-weight GTP-binding proteins. *Mol Microbiol* 32: 393–401

Pellegrino MW, Nargund AM, Kirienko NV, Gillis R, Fiorese CJ & Haynes CM (2014)
Mitochondrial UPR-regulated innate immunity provides resistance to pathogen
infection. *Nature* 516: 414–417

Pereira AG, Gracida X, Kagias K & Zhang Y (2020) *C. elegans* aversive olfactory learning
generates diverse intergenerational effects. *J Neurogenet* 34: 378–388

Peterson ND, Icso JD, Salisbury JE, Rodríguez T, Thompson PR & Pukkila-Worley R (2022)
Pathogen infection and cholesterol deficiency activate the C. elegans p38
immune pathway through a TIR-1/SARM1 phase transition. *eLife* 11: e74206

Peterson ND, Tse SY, Huang QJ, Wani KA, Schiffer CA & Pukkila-Worley R (2023) Non-
canonical pattern recognition of a pathogen-derived metabolite by a nuclear
hormone receptor identifies virulent bacteria in C. elegans. *Immunity* 56: 768-
782.e9

Prakash D, Ms A, Radhika B, Venkatesan R, Chalasani SH & Singh V (2021) 1-Undecene
from *Pseudomonas aeruginosa* is an olfactory signal for flight-or-fight response
in *Caenorhabditis elegans*. *EMBO J* 40: e106938

Pukkila-Worley R & Ausubel FM (2012) Immune defense mechanisms in the
Caenorhabditis elegans intestinal epithelium. *Curr Opin Immunol* 24: 3–9

Reimmann C, Beyeler M, Latifi A, Winteler H, Foglino M, Lazdunski A & Haas D (1997)
The global activator GacA of Pseudomonas aeruginosa PAO positively controls
the production of the autoinducer N-butyryl-homoserine lactone and the
formation of the virulence factors pyocyanin, cyanide, and lipase. *Mol Microbiol*
24: 309–319

Resko ZJ, Suhi RF, Thota AV & Kroken AR (2024) Evidence for intracellular
Pseudomonas aeruginosa. *J Bacteriol* 206

Riga A, Cravo J, Schmidt R, Pires HR, Castiglioni VG, Van Den Heuvel S & Boxem M
(2021) Caenorhabditis elegans LET-413 Scribble is essential in the epidermis for
growth, viability, and directional outgrowth of epithelial seam cells. *PLOS Genet*
17: e1009856

Ruch TR & Engel JN (2017) Targeting the Mucosal Barrier: How Pathogens Modulate the
Cellular Polarity Network. *Cold Spring Harb Perspect Biol* 9: a027953

Sauvanet C, Wayt J, Pelaseyed T & Bretscher A (2015) Structure, Regulation, and
Functional Diversity of Microvilli on the Apical Domain of Epithelial Cells. *Annu*
Rev Cell Dev Biol 31: 593–621

Schulenburg H & Félix M-A (2017) The Natural Biotic Environment of *Caenorhabditis*
elegans. *Genetics* 206: 55–86

Sepers JJ, Ramalho JJ, Kroll JR, Schmidt R & Boxem M (2022) ERM-1 Phosphorylation and
NRFL-1 Redundantly Control Lumen Formation in the C. elegans Intestine. *Front*
Cell Dev Biol 10

Shafaq-Zadah M, Brocard L, Solari F & Michaux G (2012) AP-1 is required for the
maintenance of apico-basal polarity in the *C. elegans* intestine. *Development*
139: 2061–2070

Singh J & Aballay A (2019a) Microbial Colonization Activates an Immune Fight-and-Flight
Response via Neuroendocrine Signaling. *Dev Cell* 49: 89-99.e4

Singh J & Aballay A (2019b) Intestinal infection regulates behavior and learning via
neuroendocrine signaling. *eLife* 8: e50033

Stiernagle T (2006) Maintenance of C. elegans. *WormBook*

Stuart LM, Paquette N & Boyer L (2013) Effector-triggered versus pattern-triggered
immunity: how animals sense pathogens. *Nat Rev Immunol* 13: 199–206

Stutz K, Kaech A, Aebi M, Künzler M & Hengartner MO (2015) Disruption of the C.
elegans intestinal brush border by the fungal lectin CCL2 phenocopies dietary
lectin toxicity in mammals. *PLoS ONE* 10: 1–23

Sun J & Barbieri JT (2004) ExoS Rho GTPase-activating Protein Activity Stimulates
Reorganization of the Actin Cytoskeleton through Rho GTPase Guanine
Nucleotide Disassociation Inhibitor. *J Biol Chem* 279: 42936–42944

Szumowski SC, Estes KA, Popovich JJ, Botts MR, Sek G & Troemel ER (2016) Small
GTPases promote actin coat formation on microsporidian pathogens traversing
the apical membrane of*Caenorhabditis elegans*intestinal cells: Actin coats on
microsporidia-containing vesicles. *Cell Microbiol* 18: 30–45

Tan MW, Mahajan-Miklos S & Ausubel FM (1999a) Killing of Caenorhabditis elegans by
Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis.
Proc Natl Acad Sci U S A 96: 715–720

Tan M-W, Rahme LG, Sternberg JA, Tompkins RG & Ausubel FM (1999b) *Pseudomonas*
aeruginosa killing of *Caenorhabditis elegans* used to identify *P. aeruginosa*
virulence factors. *Proc Natl Acad Sci* 96: 2408–2413

Thuenauer R, Kühn K, Guo Y, Kotsis F, Xu M, Trefzer A, Altmann S, Wehrum S,
Heshmatpour N, Faust B, *et al* (2022) The Lectin LecB Induces Patches with
Basolateral Characteristics at the Apical Membrane to Promote Pseudomonas
aeruginosa Host Cell Invasion. *mBio* 13: e00819-22

Tran CS, Eran Y, Ruch TR, Bryant DM, Datta A, Brakeman P, Kierbel A, Wittmann T,
Metzger RJ, Mostov KE, *et al* (2014) Host Cell Polarity Proteins Participate in
Innate Immunity to Pseudomonas aeruginosa Infection. *Cell Host Microbe* 15:
636–643

Tse-Kang S, Wani KA & Pukkila-Worley R (2025) Patterns of pathogenesis in innate
immunity: insights from C. elegans. *Nat Rev Immunol*

Vance RE, Isberg RR & Portnoy DA (2009) Patterns of Pathogenesis: Discrimination of
Pathogenic and Nonpathogenic Microbes by the Innate Immune System. *Cell*
Host Microbe 6: 10–21

Winter JF, Höpfner S, Korn K, Farnung BO, Bradshaw CR, Marsico G, Volkmer M,
Habermann B & Zerial M (2012) Caenorhabditis elegans screen reveals role of
PAR-5 in RAB-11-recycling endosome positioning and apicobasal cell polarity.
Nat Cell Biol 14: 666–676

Wu D, Chai Y, Zhu Z, Li W, Ou G & Li W (2017) CED-10-WASP-Arp2/3 signaling axis
regulates apoptotic cell corpse engulfment in C. elegans. *Dev Biol* 428: 215–223

Xue F, Ragno M, Blackburn SA, Fasseas M, Maitra S, Liang M, Rai S, Mastroianni G,
Tholozan F, Thompson R, *et al* (2024) New tools to monitor Pseudomonas
aeruginosa infection and biofilms in vivo in C. elegans. *Front Cell Infect Microbiol*
14

Yang HW, Shin M-G, Lee S, Kim J-R, Park WS, Cho K-H, Meyer T & Do Heo W (2012)
Cooperative Activation of PI3K by Ras and Rho Family Small GTPases. *Mol Cell*
47: 281–290

Zhang Y, Lu H & Bargmann CI (2005) Pathogenic bacteria induce aversive olfactory
learning in Caenorhabditis elegans. *Nature* 438: 179–184

Figure 1

A

% Surviving vs Exposure period (Days)
- OP50
- PA14

B

Cuticle, Enterocyte, Lumen, Apical surface, Microvilli, Cell cortex, Adherens Junctions, Terminal web (Endotube), Baso-lateral region, Nucleus, Enterocyte

D

Exposure period (Days) 1 2 3
No. of worms showing intestinal deformation (%)
- n = 2/54 (OP50), n = 27/56 (PA14)
- n = 3/43 (OP50), n = 28/43 (PA14)
- n = 10/52 (OP50), n = 43/52 (PA14)

C

OP50 | PA14

Exposure period (Days): 1, 2, 3

ACT-5

E

$$\text{Deformation Index } (\delta) = \frac{\ell}{L}$$

$\delta \sim 1$ (control)

$\delta > 1$ (deformed intestine)

F

Exposure period (Days) 1 2 3
Deformation Index (δ^{apical})
- OP50, PA14
- ns, ***, ***

G

Deformation Index (δ^{apical}) vs Exposure period (Days)
- OP50
- PA14

H

2 Days post exposure

OP50 | PA14

- ERM-1 — Microvilli
- PAR-6 — Sub-apical
- IFB-2 — Endotube
- PGP-1 — Apical Membrane

FIGURE LEGENDS

Figure 1: *P. aeruginosa* infection deforms the apical surface of intestinal
epithelium in *C. elegans*

(A) Representative Kaplan-Meier survival plot of N2 worms infected with *P. aeruginosa* (PA14-
magenta) or *E. coli* (OP50-black) at 25^0 C. (n = 100 worms, N=2)

(B) Schematic illustration of *C. elegans* intestinal epithelium depicting different regions of the
enterocyte.

(C) Medial confocal sections of C. elegans intestines expressing mCherry::ACT-5 localized to
apical surface. While lumen distension increases with age in animals exposed to with *P.*
aeruginosa or *E. coli* (red double headed arrows), intestinal surface deformation is seen only
upon exposure to with *P. aeruginosa* (red filled arrowheads), but not *E. coli* (blue ROI; blue
filled arrowheads). with *P. aeruginosa* infection also led to actin vesicles budding of apical
surface (Red ROI; red arrows).

(D) Bar graphs showing increased percentage of intestinal deformation with increase in the
duration of exposure to with *P. aeruginosa*.

(E) Schematic showing the estimation of deformation index (δ^{apical}) of apical surface

(F) Representative violin plot showing quantification of deformation index of apical surface
shown in Fig.1C. (N=3; n= 10 worms per treatment per set).

(G) Plot showing progressive deformation of apical surface of intestinal lumen from individual
worms measured over 3 days following *P. aeruginosa* or *E. coli* exposure. (n=5)

(H) Medial confocal images of *C. elegans* expressing fluorescently labelled apical markers
upon exposure to *P. aeruginosa* or *E. coli*. ERM-1::eGFP (apical regions containing microvilli),
PAR-6::mNeonGreen (apical PAR polarity complex), IFB-2::wScarlet (intermediate filaments),
PGP-1::GFP (apical plasma membrane).

Scale bar is 20 μm. P-values were calculated by Mann-Whitney U test. ****p<0.0001, ***p
<0.001, and * p <0.05

Figure 2: Exposure to *P. aeruginosa* results in cytoplasmic mis-localization of
CDC-42, ARX-2 and ACT-5

(A) RNAi mediated depletion of *cdc-42*, but not *rho-1 or ced-10* rescues the *PA-14* induced
intestinal deformation. Red filled arrowheads arrow heads show deformed regions of the
lumen, while red open arrow heads denote the smooth intestinal lumen surface in *cdc-*
42(RNAi) worms.

(B) Quantification showing reduced deformation index of intestines in *cdc-42(RNAi)* worms
compared to L4440 (control) upon *P. aeruginosa* exposure. Measurements are plotted as
violin plot indicating median and interquartile of each data set. (N = 3; n≥12 worms per
condition and per replicate)

(C) Representative Kaplan-Meier survival plot of *cdc-42(RNAi)* or L4440 (control) worms
subjected to *P. aeruginosa* or *E. coli* infection. CDC-42 depletion partially rescues *P.*
aeruginosa induced shortening of lifespan (TD50$^{cdc42(RNAi)}$ = 8 days, TD50^{control} = 4 days) (N=2;
n= 50 worms per set and per condition)

(D, G and J) Medial cross-sections of intestinal lumen of *C. elegans* expressing (D) CDC-
42::GFP, (G) ARX-2::TagRFP, and (J) mCherry::ACT-5 following exposure to *E. coli* or *P.*
aeruginosa. Images shown are pseudo-colored to highlight changes in the intensity. Warm
colors represent higher intensity and lower intensity by the dimmer colors. Cyan arrows
indicate the cytoplasmic enrichment of these proteins in animals infected with *P. aeruginosa*.
Where visible, nuclei of the intestinal cells are marked with letter 'n'. Scale bar -20 *µ*m.

(E and F) Violin plots showing cytoplasmic (E) and apical (F) intensities of CDC-42::GFP upon
P. aeruginosa or *E. coli* infection. Measurements are mean ±95% C.I. Each dot represents
one animal and the data shown is the cumulation of three independent experiments (N=3;
n=33).

(H and I) Violin plots indicating cytoplasmic (E) and apical (F) intensities of ARX-2::TagRFP
upon *P. aeruginosa* or *E. coli* infection. Measurements are mean ±95% C.I. Each dot
represents individual worm (N=2; n≥21)

(K and L) Violin plots of cytoplasmic (E) and apical (F) intensity measurements of
mCherry::ACT-5 upon *P. aeruginosa* or *E. coli* infection. Measurements are mean ±95% C.I.
Each dot represents individual worm (N=2; n≥14)

Scale bar is 20 *µ*m. P-values were calculated by Mann-Whitney U test. ****p<0.0001, ***p
<0.001, **p <0.01and * p <0.05.

Figure 3: Inhibition of ARP-2/3 dependent actin polymerization prevents *P.*

***aeruginosa* -induced apical surface deformation**

(A) Intestinal lumen marked by mCherry::ACT-5 from control and *cdc-42(RNAi)* animals

exposed to *P. aeruginosa* or *E. coli*. Red arrowheads indicate rescue of intestinal deformation

in *cdc-42(RNAi)* animals.

(B) Cytoplasmic ACT-5 intensity quantified from intestinal epithelium of control and

cdc42(RNAi) worms, day 2 post infection. Quantifications are mean ±95% C.I. (n ≤ 14).

(C) Confocal images of intestinal epithelium from control and *arx-2(RNAi)* animals co-

expressing CDC-42::GFP and ARX-2::tagRFP, and subjected to *E. coli* and *P. aeruginosa*

infection. All images are from 2- days post infection. Open red arrows indicate rescue of the

intestinal deformation.

(D) Quantification of cytoplasmic CDC-42::GFP in control and *arx-2(RNAi)* animals following

2-day exposure to *P. aeruginosa* or *E. coli*. The violin plot shows the distribution of data with

median and interquartile for each data set. p-values n≥8 for all the sets. expect n= 5 for PA14

arx-2 (RNAi).

(E) Violin plot showing apical deformation index quantification for *arx-2(RNAi)* animals

exposed to *E. coli* or *P. aeruginosa*. Plot shows the median and interquartile of each data set.

(n ≤ 11).

(F) Medial plane image of intestinal lumen co-expressing PGP1::GFP and mCherry::ACT5

subjected to control (L4440) and *arx-2* RNAi followed by *E. coli* and *Pseudomonas* exposure.

All the images are from one day post *E. coli* and *Pseudomonas* exposure.

(G) Medial plane image of intestinal lumen co-expressing CDC42::GFP and ARX2::TagRFP

subjected to control (L4440, empty vector) and *act-5* RNAi followed by *E. coli* and

Pseudomonas exposure. All the images are from one day post *E. coli* and *Pseudomonas*

exposure. Open red arrow marks represent rescue of the intestinal deformation.

unless otherwise specified, Scale bar is 20 μm. P-values were calculated by Mann-Whitney U

test. ****$p<0.0001$, ***$p <0.001$, and * $p <0.05$.

Figure 4: Fragmented ARP-2/3 disrupts actin assembly leading to extracellular

vesicle release from apical surface

(A) Intensity visualization of medial confocal images of *C. elegans* intestines expressing ARX-

2::TagRFP showing a fragmented distribution of ARX-2 (cyan arrows) on day-2 post *P.*

aeruginosa exposure. ARX-2 shows a contiguous localization in worms exposed to *E. coli*

(cyan arrowheads). Surface intensity of plots of enlarged ROIs highlights the fragmentation.

Scale bar -10 μm.

(B) *C. elegans* co-expressing ARX-2::GFP and mCherry::ACT-5 showing actin structures

(yellow arrowheads) permeating into the lumen from ARX-2-free regions (cyan arrows). Scale

bar -10 μm.

(C) Lattice light sheet images acquired 2 days post exposure of PA14-GFP or OP50-GFP -fed

C. elegans expressing intestinal mCherry::ACT5. Images were acquired in 13.3X objective

and represented as maximum intensity projection in volume mode. Cyan arrows in enlarged

ROI indicate the vesicles emerging from apical surface into the lumen.

(D) Membrane bound actin rich vesicles bud off from intestinal apical surface following 2-day

P. aeruginosa exposure (cyan arrows in ROI). Enlarged ROI shows vesicles at various stages

of budding – (a) initial bulging, (b) bud formation, and (c) released vesicle. An intensity linescan

drawn across the extracellular vesicle indicates actin (magenta) surrounded by the apical

plasma membrane marked by PGP-1::GFP (green). Scale bar - 5 μm.

(E) Line scan-intensity plot of mCherry::ACT5 and PGP1::GFP vesicles 2 days *P. aeruginosa*

exposure. (n=20)

(F) LifeACT::mRuby localizes to F-actin in the apical vesicles (cyan arrow) of *C. elegans*

intestines exposed to *Pseudomonas*. Scale bar - 10 μm.

P-values were calculated by Mann -Whitney U test. ****$p<0.0001$, ***$p <0.001$, and * $p <0.05$.

A

LET-413   DLG-1   LET-413 / DLG-1

DPI 2   OP50 / PA14

B

Exposure period (Days)

1   2

LET-413^apical (A.U)

OP50   PA14   OP50   PA14

****   ****

C

Exposure period (Days)

1   2

LET-413^lateral (A.U)

OP50   PA14   OP50   PA14

****   ****

D

DLG-1 / LET-413

OP50   PA14

DPI 2   L4440 / cdc-42 (RNAi) / akt-1/ akt-2 (RNAi)

E

CDC-42 / ARX-2

OP50   PA14

L4440 / akt-1/ akt-2 (RNAi) / age-1 (RNAi)

Figure 5 : *P. aeruginosa* induced polarity disruption is mediated via PI3K-AKT

kinase pathway

(A)   LET-413 shows enhanced accumulation in the apical (yellow arrow) and lateral (yellow

arrowhead) surfaces *C. elegans* enterocytes upon *P. aeruginosa* infection. DLG-1 marks

cell junctions (cyan arrows). Double asterisks shows absence of LET-413 at the apical

surface in intestines of *C. elegans* exposed to *E. coli*.

(B and C) Quantification of (B) apical, and (C) cytoplasmic LET413 levels in enterocyte of

worms following *E. coli* (OP50) or *P. aeruginosa* (PA14) infection. Each data point

represents individual worm. Measurements are plotted as violin plot indicating median and

interquartile of each data set. (n=12). p-value was calculated using Mann-Whitney U test.

(n=12)

(A) Depletion of AKT-1 and AKT-2, or CDC-42 rescues *P. aeruginosa* induced loss of polarity.

In contrast to control (L4440; yellow asterisks), apical recruitment of LET-413 in response

to *P. aeruginosa* infection is absent in *akt-1/akt-2(RNAi)* and *cdc-42(RNAi)* (yellow

asterisks).

(B) Disruption of PI3K-AKT pathway abrogates *P. aeruginosa* induced mis-localization of CDC-

42, ARX-2 and actin. Medial confocal plane of intestinal lumen co-expressing CDC-

42::GFP and ARX-2::TagRFP subjected to *akt-1/akt-2 and age-1* RNAi followed by *E. coli*

and *Pseudomonas* exposure for two days. Closed yellow arrow indicate intestinal

deformation. Yellow asterisks denote rescue of deformation.

Scale bar in 20 μm. P-values were calculated by Mann -Whitney U test. ****$p < 0.0001$, ***p

< 0.001, and * $p < 0.05$

Figure 6

Divyashree et. al (2025)

Figure 6:   Extracellular *P. aeruginosa* can induce intestinal deformation in *C.*

elegans

(A) Confocal medial images of intestinal epithelium from *C. elegans* expressing ACT-

::mCherry and exposed to GFP expressing *E. coli* (top panel), *S. typhymurium* (middle panel)

and *P. aeruginosa* (bottom panel) for 3-days post infection. Dashed lines indicate cross-

sectional plane whose orthogonal view is presented adjacent to the image. Red hollow (*E. coli*

) and yellow (*S. typhimurium)* arrowheads indicate intestinal distension (bloating), but smooth

lumen surface. Red filled arrowheads indicate the deformation of the intestine in worms

infected with *P. aeruginosa*. Scale bar - 20 *µ*m.

(B) Medial confocal images of intestines from *C. elegans* expressing mCherry::ACT-5 animals

grown on heat killed *E. coli* (top panel) and *P. aeruginosa* (bottom panel). Hollow red arrow

indicates the absence of intestinal deformation upon exposure to heat killed *P. aeruginosa*.

Scale bar- 20 *µ*m.

(C) Medial confocal images of intestines from *C.* elegans expressing mCherry::ACT-5

exposed to *E. coli*(OP50) (top panel), *P. aeruginosa* (PA14) (middle panel) and *P. aeruginosa*

(PA14ΔgacA) (bottom panel). Filled red arrows indicates intestinal deformation. Scale bar -

20 *µ*m

(D) Quantification of deformation index (δ^{apical}) of the apical intestinal surface in C. elegans

expressing intestinal mCherry::ACT-5 exposed to *E. coli*(OP50) ,*P. aeruginosa* (PA14) and *P.*

aeruginosa. Each data point represents individual animal and measurements are plotted as

violin plot indicating median and interquartile of each data set. (n =10)

(E) Kaplan- Meier survival plot of worms described in Fig. 6A upon exposure to *E. coli*(OP50)

, *P. aeruginosa* (PA14) and *P. aeruginosa* . Worms were grown on *E. coli* till L4 stage and

transferred on to the respective infection plates. (n = 50 worms, N=2)

(F) Schematic representation of the proposed mechanism for *P. aeruginosa* intestinal apical

surface deformation in *C. elegans*

Scale bar in 20 *µ*m. P-values were calculated by Mann -Whitney U test. ****p<0.0001, ***p

<0.001, and * p <0.05