Fair-weather friends: Unequal partnerships between *Parastagonospora nodorum* and *Pyrenophora tritici-repentis* define disease dynamics in wheat

Running title: Unequal partners in wheat co-infection

Leon Lenzo [1], Evan John [2], Jason Bradley [3], Geoff Thomas [3], Dion Bennett [4,5] and Kar-Chun Tan [1*]

[1]Centre for Crop and Disease Management, School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, 6102, Australia

[2]Department of Plant Pathology, Institute of Crop Science and Resource Conservation (INRES), University of Bonn, Nussallee 9, Bonn, 53115 Germany

[3]Department of Primary Industries and Regional Development, 6151, Perth, Western Australia, Australia

[4]Australian Grain Technologies, Northam, Western Australia, 6401, Australia

[5]School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, 6102, Australia

*Corresponding author; Kar-Chun.Tan@curtin.edu.au

Keywords: *Parastagonospora nodorum*, *Pyrenophora tritici-repentis*, necrotrophic, fungal, pathogens, co-infection, septoria nodorum blotch, wheat, tan spot, yellow leaf spot, priority effects, digital PCR

Abstract

Parastagonospora nodorum and *Pyrenophora tritici-repentis* are the causal agents of septoria nodorum blotch and tan spot of wheat, respectively. Though these fungal phytopathogens have been found to frequently cohabitate the same leaf, their interaction dynamics in the manifestation of disease remain poorly understood due to limitations in species-specific detection methods. We developed a digital PCR based model targeting conserved regions of the α-tubulin gene, enabling biomass quantification of both pathogens during infection. Field surveys revealed up to two in three symptomatic infections involved both pathogens, with co-infected plants showing significantly higher individual pathogen biomass than single-species infections. Host plants in the field with moderate resistance to both pathogens were found to be significantly more necrotic under co-infection, with individual pathogen biomass up to twice that observed value for single infections. However, like fair-weather friends the partnership between these two pathogens seems to be conditional. When *P. tritici-repentis* established first, secondary *P. nodorum* colonisation led to a breakdown of host resistance. Conversely, when *P. nodorum* established on the host first, it suppressed *P. tritici-repentis* colonisation regardless of host resistance. To our knowledge this is the first description of asymmetric priority effects overcoming host resistance in a plant pathosystem. Resistance breeding strategies evaluating single pathogen challenges may inadvertently select for cultivars vulnerable to sequential co-infection, necessitating integrated disease complex approaches for durable resistance development.

Introduction

The order in which microorganisms colonise habitats fundamentally shapes community assembly and species interactions, a phenomenon known as priority effects [1]. In the colonisation of plant tissues, early arriving species modify environmental conditions and resource availability, altering biotic conditions to determine the trajectory of community development [2]. Agricultural phyllospheres represent a synthetic, simplified ecological context, where vast expanses of genetically homogeneous hosts grown at high density create conditions that accelerate pathogen evolution and promote rapid population growth [3]. Previous work

has demonstrated priority effects in plant-associated microbial communities, with pathogen arrival order

determines infection outcomes and host responses, yet their operation in agricultural pathosystems

remains poorly understood despite their potential to fundamentally alter disease outcomes [4,5]. Recent

surveys using molecular analysis across European wheat fields revealed that over two thirds of

symptomatic leaf samples harboured at least two pathogenic fungal species [6]. Environmental monitoring

studies using spore trapping showed propagules from fungal pathogens circulate simultaneously during

growing seasons, creating opportunities for co-infection [7]. While priority effects have been extensively

documented in bacterial pathosystems, with pathogens employing diverse strategies including

competition, cooperation, and temporal separation to succeed in polymicrobial environments, their

operation in fungal plant pathosystems remains poorly understood despite their potential to fundamentally

alter disease outcomes [8]. Polymicrobial infections in plant systems may result in resource competition, host

immune modulation or direct pathogen-pathogen interactions that chemically or mechanically modulate

the colonization, reproduction, and transmission success of coinfecting species [9]. These interactions can

be either facilitative, where one pathogen creates conditions that enhance proliferation of another [10,11], or

inhibitory, where pathogens actively suppress competitors through antimicrobial compounds or resource

monopolization [12,13].

Parastagonospora nodorum and *Pyrenophora tritici-repentis*, the causal agents of septoria nodorum blotch

(SNB) and tan spot (TS) of wheat, represent an ideal system for investigating priority effects in necrotrophic

plant pathosystems [14,15], with previous work documenting near ubiquitous co-occurrence across wheat

growing regions [16]. Empirical studies of interactions between species have been limited however, largely due

to the morphological similarity of their constituent diseases and lack of robust quantitative methods. Both

pathogens are members of the order Pleosporales, which also includes pathogens such as *Leptosphaeria*

and *Alternaria* spp., and share similar infection strategies [17]. They each employ necrotrophic effectors that

interact with dominant host susceptibility genes to trigger an uncontrolled hypersensitive response,

resulting in necrotic lesions that support pathogen proliferation [18]. Notably, both species share the

necrotrophic effector ToxA, with evidence suggesting a horizontal gene transfer event, indicating stable

physical proximity between the two species [19]. Breeding efforts have made significant progress in improving

wheat resistance to both diseases through the elimination of dominant susceptibility genes such as *TaTsn1*,

which interacts with ToxA [20], and through the stacking of resistance alleles for both SNB and TS [21,22]. Despite

these advances, both diseases continue to persist in wheat-growing regions worldwide and remain priority

targets for resistance breeding [23], suggesting that current single-pathogen evaluation approaches may not

capture the complexity of field disease dynamics.

Previous efforts to study co-infection dynamics between these pathogens have faced significant

methodological constraints. Due to the morphological similarity of lesions produced by both pathogens,

and with coalescence obscuring distinct characteristics, visual assessment of disease symptoms requires

microscopic analysis even by trained pathologists [24,25]. Neural-network based disease identification

algorithms, which have achieved superior classification accuracy for other wheat diseases, have not yet

addressed the challenging differentiation between morphologically similar leaf spot diseases such as SNB

and TS [26]. The development of Real-Time Polymerase Chain Reaction (RT-PCR) assays to quantify both

pathogens in host tissue as well as the closely related hemi-biotroph *Zymoseptoria tritici* has provided

pioneering insight into the dynamics of foliar wheat pathogens during co infection [6,27,28]. However, these

studies rely on DNA as a direct analogue for pathogen biomass well as the utilization of targets with unstable

copy number in the genome including ribosomal and transposon associated sequences. For example, a

multicopy target for *P. tritici-repentis* [29] which was found to be variable across strains and regions (Figure S1)

[28]. Although this increases its sensitivity, it renders the assay unsuitable for field application where

uncharacterised strains may be present. Additionally, RT-PCR suffers from a lack of reproducibility at low

pathogens titre, especially in complex environmental samples where PCR inhibitors are known to be

present [30]. Digital PCR (dPCR) addresses limitations of RT-PCR by providing absolute quantification of target

nucleic acids without standard curves, offering greater precision and reproducibility while being more

tolerant of inhibitors and capable of reliably detecting low abundance targets [31].

Despite significant advances in breeding for resistance against SNB and TS, both *P. nodorum* and *P. tritici-*

repentis continue to persist in wheat-growing regions globally [6,16,23]. They are most commonly found under

co-infection conditions, suggesting that interactions between the two organisms may facilitate their

sustained proliferation. Consequently, SNB-TS wheat pathosystem has been a focal point and model of

research into polymicrobial infection of plants in recent years, where a range of RT-PCR-based targets have

been developed for pathogen monitoring and demonstrated a high field co-infection incidence [6,16,28]. It is not

known how wheat varieties that differ in resistance to TS and SNB dictate the outcome of single versus SNB-

TS co-infection, and so we sought to clarify whether *P. nodorum* and *P. tritici-repentis* form cooperative or

competitive associations during co-infection in wheat. We hypothesise that pathogen proliferation will be

determined by: (1) genetic resistance present in the host wheat leaves, (2) co-infecting pathogen presence

or absence and (3) pathogen arrival order. Here, we develop a duplex digital PCR assay alongside automated

image analysis enabling simultaneous absolute quantification of both pathogens from infected tissue and

apply these tools to investigate polymicrobial disease dynamics in a necrotrophic plant pathosystem.

Materials and Methods

Fungal Manipulations

Reference isolates for *P. nodorum* SN15 [32] and *P. tritici-repentis* M4 [33] were used for all *in vitro* and defined

in planta inoculation in this study. All isolates were maintained on 90 mm V8 PDA plates [34], at ambient

temperature under a 12h light/dark photoperiod. To generate mycelial tissue, isolates were grown in Fries 3

broth [35] before lyophilisation . To generate inoculum, *P. nodorum* conidiospores were harvested directly from

V8PDA plates and resuspended in 0.02% w/v Tween . Inoculum for *P. tritici-repentis* isolates was generated

as previously described [36].

Visual Disease Analysis

Harvested leaves were photographed *in situ* using a Google Pixel 6 Pro (USA). Individual leaf images were

manually extracted and catalogued using ImageJ [37]. Differential hue value masks were then generated to

distinguish between healthy, chlorotic and necrotic tissue, with cutoffs at 90, 43 and 30 respectively utilizing

the opencv library for Python. For full HSV values, refer to Table S1. The Python tools "LeafScan" for image

processing and analysis were developed and are available at https://github.com/LeonLenzo/leaf-scan.

Normalised pixel values to total leaf areas were used for downstream analyses.

Nucleic Acid Purification

All fungal and plant tissues were snap frozen in liquid nitrogen and homogenized using a Tissue Lyser II

(Retsch, Duesseldorf, Germany), and DNA was purified using the Biosprint DNA Plant Kit in a Biosprint15

(both, Qiagen, Hilden, Germany) DNA concentrations were determined by fluorometric analysis using a

Qubit Flex system employing the dsBRDNA Working Solution (Thermofisher, Massachusetts, USA).

Digital PCR Assay

Primers aT_F and aT_R were designed to simultaneously amplify a 187-bp region in *P. nodorum* and a 185-

bp region in *P. tritici-repentis*. These sequences were then used to query 179 *P. nodorum* [38,39] and 67 *P. tritici-*

repentis [14,40] genome assemblies to ensure within-species conservation and single copy status. Internal from

the primer-binding-site sequences, species-specific fluorescent probes Pn_FAM and Ptr_HEX were

designed to differentially amplify the target organisms (Table 1). Purified gDNA from *P. nodorum* SN15 and

P. tritici-repentis M4 were used to optimize the assay. dPCR reactions were carried out in 22-µl volumes,

where each well contained 11µl ddPCR Supermix for Probes (No dUTP) (Bio-Rad, California, USA), 5µl DNA

template, and 0.25µM of primer/probe mix at equimolar concentration. A QX200 Automatic Droplet

Generator (Bio-Rad, California, USA) was used to fractionate samples according to manufacturer's

instructions, before undergoing thermal cycling at 95°C for 10 min, then 40 cycles of 94°C for 30 s, 60°C for

60 s, then a final denaturation of 95°C for 10 min. The QX200 Droplet Reader (Bio-Rad, California, USA) was

used to quantify positive droplets, with data analysis performed using QuantaSoft Analysis Pro v1.0.596

(Bio-Rad, California) to determine absolute copy numbers following Poisson distribution correction. For

downstream analysis, copy per microlitre outputs were only counted for values >1, and all input DNA values

were normalised to 1ng/µl. In silico off-target analysis was performed by querying all primer and probe

sequences against the NCBI RefSeq fungi database using BLAST nucleotide search. BLAST results were

filtered using minimum sequence identity thresholds of 60% for primers and 85% for probes. Cross-reactive

organisms were identified as those showing significant hits to both primers (aT_F and aT_R) and at least one

species-specific probe (Pn_FAM or Ptr_HEX), representing potential targets for non-specific dPCR

amplification (Figure 1D). Primers and probe specificity were tested *in vitro* using the ddPCR assay (Figure

2C) against the purified gDNA of closely related fungal pathogens and wheat (Table S2).

Biomass Modelling

To approximate a relationship between dPCR copy number and fungal biomass, we developed a two-step

calibration approach. First, we quantified the DNA yield per unit fungal tissue by extracting DNA from known

quantities of freeze-dried mycelium. Three biological replicates of *P. nodorum* SN15 and *P. tritici-repentis*

M4 mycelia were weighed out in technical triplicates between 0.15 and 15 mg and DNA was extracted as

described above. Total DNA content was plotted as a function of tissue mass and analysed using linear

regression to derive the species-specific DNA-to-tissue ratio (β) for each pathogen. Second, we established

the relationship between DNA input and dPCR target copy number by analysing serial dilutions of gDNA

from both species. Triplicate samples at concentrations ranging from 0.05 to 500 pg/µl were analysed using

the dPCR assay described above. The integrated biomass model combines these relationships through the

formula below where C represents dPCR-detected alpha-tubulin copies/µl, α is the target copies per

nanogram of DNA, and β is the micrograms of DNA per milligram of tissue. After substituting the

experimentally derived values, we obtained simplified predictive equations for each species that enable

direct conversion of dPCR-measured target copies into estimated fungal biomass.

$$Tissue_{(ug)} = \frac{C}{\alpha\beta}$$

Field Sites and Experimental Design

Symptomatic leaf tissue was collected between May and October of 2022 from field sites sown in early April

in Western Australia. The first, a disease nursery unattenuated by fungicide for more than ten years

(Northam, 31°32'23.4"S 116°42'17.3"E). The second plot (South Perth, -31.991046, 115.888172), contained

three adjacent plots, established by incorporating infected wheat stubble containing single or mixed

pathogen species into the soil prior to sowing, with cross-contamination between adjacent plots expected

over the growing season to test competitive dynamics under varying establishment priorities (Figure S2).

Cultivars were selected for resistance through Department of Primary Industries and Regional Development

of Western Australia wheat variety disease ratings to select lines with differential susceptibility to SNB and

TS (Table 2). SVS (susceptible-very susceptible) indicates wheat cultivars with minimal disease resistance,

while MRMS (Moderately Resistant-Moderately Susceptible) denotes cultivars with intermediate resistance

levels. Samples were photographed on site, trimmed with a scalpel to 50 mm in length, then stored on ice

for ≤ 24h before snap freezing and storage at -80°C until processing.

Attached Leaf Assay

Wheat seedlings were grown under white light on a 12hr photoperiod until 12 days old. Leaves were

mounted adaxial side up onto 45mm XPS foam boards using micropore tape inside a seedling tray. Attached

leaves were inoculated with 15µl of either 1×10^7 spores.ml^{-1} of *P. nodorum* SN15 spores, 1×10^5 spores.ml^{-1} of

P. tritici-repentis M4 spores suspended in 0.02% Tween 20 or a mixture of the two maintaining the original

concentration. Following inoculation spore suspension was spread with a paintbrush (WestArt, 4mm

Synthetic), leaves allowed to dry, then sealed inside humidity chambers and maintained at 22C ± 1C, ≥ 95%

humidity under white light on a 12 hr photoperiod. Staggered infections were carried out 72 hrs post-initial

inoculation following the same procedure. Leaves were photographed seven days after secondary infection,

before excising a 50mm section with a sterile scalpel.

Statistical analysis

Statistical analyses were performed using R version 4.5.0 [41]. Linear regression analysis was conducted using

the lm() function to fit a linear model. Pathogen biomass data were log10-transformed to improve normality

and stabilize variance. Normality was assessed using Shapiro-Wilk tests, which confirmed that log10

transformation substantially improved the distribution of biomass data. Student's t-tests were used to

compare pathogen biomass between co-infection and single infection treatments within each site and

wheat line. For sequential inoculation data, one-way ANOVA was conducted separately for each wheat line

with treatment as the factor. Post-hoc comparisons used Tukey's HSD test when significant effects were

detected ($p < 0.05$). ANOVA assumptions were verified through residual analysis and Shapiro-Wilk tests for

normality.

Results

Biomass modelling and absolute pathogen DNA quantification using dPCR

To identify a suitable locus for marker development, gene annotations from SN15 and M4 were compared

using reciprocal BLAST analysis to identify highly conserved orthologues. Candidates were filtered manually

to identify regions ≤200 bp with conserved 5' and 3' termini for primer binding sites adjacent to a divergent

central region for a species-specific probe, before selecting a region spanning the final intron of the alpha-

tubulin ortholog in both species (Pn_SN15_01762, Ptr_M4_98590). Suitable loci within the orthologous

alpha-tubulin coding sequences were identified so that a single primer set could target binding sites

conserved across both species, while species-specific internal probes (Pn_FAM and Ptr_HEX respectively)

permitted species discrimination (Figure 1A & B). Single copy primer-probe sites were conserved at a single

copy across pan-genomes of the respective species, with in silico cross-reactivity analysis confirming

minimal risk of in species off target effects (Figure 1C). No cross reactivity to known wheat associated fungi

was observed when tested in silico, with only seven primer + probe matches ≤ 98% found excluding target

organisms (Figure 1D). Species-specific DNA yield analysis revealed *P. tritici-repentis* produces 0.160 μg

DNA per mg dry tissue compared to *P. nodorum* at 0.059 μg DNA per mg dry tissue (Figure 2A). Calibration

curves for both probes demonstrated near-perfect linearity ($R^2 > 0.99$) across a broad range of fungal DNA

(0.5 pg.µl^{-1} to 0.5 ng.µl^{-1}) for both pathogens (Figure 2B). When tested against gDNA of the wheat host or

related fungal species and alternative wheat pathogens, the assay showed no off-target effects (Figure 2C).

The integrated biomass model yielded conversion factors of 1.3 µg biomass per target copy for *P. tritici-*

repentis and 3.4 µg per target copy for *P. nodorum*, providing the foundation for all subsequent comparative

analyses (Figure S3).

Incidence of host tissue necrosis (but not chlorosis) correlates with fungal biomass accumulation

Total fungal biomass measured through dPCR correlated with necrosis measured with automatic pixel

thresholding (Figure 3A). Controlled infection of wheat cvs Axe and Scepter with single or mixed *P.*

nodorum/*P. tritici-repentis* inoculum were monitored through photography and tissue sampling over a 10-

day post-infection period. Linear regression models were constructed for each of three distinct disease

metrics: chlorosis, necrosis and total disease (necrosis plus chlorosis) (Figure 3B). Of the three metrics

evaluated, necrosis demonstrated the strongest positive correlation ($R^2 = 0.705$). Total disease coverage

(chlorosis plus necrosis) exhibited a moderate positive correlation with pathogen biomass ($R^2 = 0.445$).

Interestingly, chlorosis showed a weak negative correlation with pathogen biomass ($R^2 = 0.214$), suggesting

that chlorotic symptoms are not indicative of high pathogen loads. We then extended this analysis to

naturally infected wheat plants sampled weekly over a 16-week period beginning at tillering stage at a field

site. Field data revealed substantial temporal variation in disease-biomass correlations (Figure 3C). Strong

correlations similar to controlled conditions were observed in week 1, but collapsed and recovered across

the season, likely reflecting temporal confounding as infections at varying stages are sampled

simultaneously. This temporal pattern highlighted the importance of assessment timing for the correlation

between visual symptom assessment and fungal biomass.

Field co-infection facilitates biomass accumulation and the breakdown of host resistance

Pathogen biomass responses to genetic resistance in host plants were examined in four wheat cultivars with

combinatorial differences in TS and SNB disease resistance ratings (Table 1), sampled in a naturally wheat-

disease nursery endemic for both *P. nodorum* and *P. tritici-repentis*. Co-infected samples were the most

common type observed (63%) validating previous observations [6,16], with single infections representing 21%

and 16% of samples for *P. tritici-repentis* and *P. nodorum,* respectively (Figure 4). For Axe (SVS to both TS and

SNB), no significant differences in necrosis severity were observed between co-infected and single-infected

samples, with significantly higher *P. tritici-repentis* but not *P. nodorum* biomass observed under co-

infection. With Emu Rock and Yitpi, the only significant difference observed was a higher proportion of *P.*

nodorum biomass in Emu Rock, with no single infections of a given pathogen observed in hosts with a

resistance to said pathogen.

Wheat cv. Scepter, which is rated MRMS to both TS and SNB, demonstrated greater necrosis when both

pathogens were present (Figure 4A). Both *P. tritici-repentis* and *P. nodorum* were detected at very low single-

infection frequencies and significantly less biomass than co-infected samples. In general, both pathogens

exhibited significantly higher biomass accumulation under co-infection conditions compared to single-

pathogen infections (Figures 4B & C). Notably, the only wheat cultivar to exhibit higher necrosis and biomass

during co-infection was Scepter, the host with the highest resistance to both pathogens.

Fair weather friends: P. tritici-repentis facilitates co-infection while P. nodorum suppresses

We sought to isolate priority effects in the field by using a defined primary inoculum in a field trial involving

three adjacent plots, capable of spreading airborne inoculum for secondary infections. Exterior plots

contained either stubble infected with *P. tritici-repentis* or *P. nodorum* and the central plot contained stubble

with an equal mixture of both (Figure S2). For the *P. tritici-repentis* and mixed stubble site, co-infection was

again ubiquitous throughout the growing season, suggesting the presence of secondary airborne inoculum

[7]. Disease incidences measured by necrosis and biomass for both species were significantly higher under

co-infection than for single infection, with the exception of *P. nodorum* at the mixed site, broadly reflecting

the patterns seen under endemic conditions (Figures 5, S4 and S5).

The *P. nodorum* inoculum site revealed dramatically different interaction dynamics. *P. tritici-repentis* failed

to establish single-pathogen infection and was only observed at a low frequency in co-infected samples

(Figures 5 and S6). This suggests that *P. nodorum* may antagonise secondary *P. tritici-repentis* host-infection

indicating priority effects in the tripartite *P. nodorum-P. tritici-repentis*-wheat.

Controlled sequential infection reveals an asymmetric partnership

To confirm this antagonism, we conducted sequential co-infection assays to investigate the role of temporal

dynamics. *P. nodorum* and *P. tritici-repentis* were inoculated onto wheat cvs. Axe and Scepter seedlings in

a controlled environmental growth chamber, where secondary infection was carried out 72 hrs after primary

infection using defined pathogen isolates. Analysis of wheat cv. Axe revealed consistent biomass

accumulation and necrosis development across treatments as well as secondary-infection timings (Figure

6A). Simultaneous inoculation resulted in *P. tritici-repentis* dominance, while *P. tritici-repentis* priority led to

balanced pathogen loads with no significant competitive advantage for either species pertaining to biomass

accumulation. When *P. nodorum* was applied as a primary inoculum ahead of *P. tritici-repentis*, it again

inhibited subsequent *P. tritici-repentis* biomass accumulation. This was the strongest competitive

asymmetry observed across all treatments on wheat cv. Axe and provides additional evidence of the

inhibitory dynamics observed under field conditions.

Sequential co-infection on wheat cv. Scepter displayed markedly different infection dynamics. Most

treatments resulted in substantially reduced pathogen biomass and necrosis compared to wheat cv Axe,

likely due to background resistance (Figure 6B). *P. tritici-repentis* followed by *P. nodorum* inoculation was

the sole exception, representing a breakdown of seedling resistance where both pathogens achieved

biomass levels comparable to those in susceptible plants. This treatment was also the only one to induce

significant necrosis in resistant plants, suggesting that *P. tritici-repentis* establishment compromises

subsequent resistance responses to *P. nodorum,* allow both pathogens to proliferate.

Collectively, this work confirms priority effects influence pathogen proliferation in the SNB-TS/wheat

pathosystem. Under natural field conditions, co-infection is ubiquitous and facilitates biomass

accumulation for both pathogens, synergistically break down host resistance. When the inoculum source

is controlled, *P. nodorum* suppresses *P. tritici-repentis* establishment, while *P. tritici-repentis* facilitates

balanced co-infection. In controlled sequential infections, *P. tritici-repentis* primacy enables both

pathogens to overcome genetic resistance, while *P. nodorum* primary infection consistently suppresses *P.*

tritici-repentis infection regardless of host background.

Discussion

Co-infection dynamics play crucial roles in wheat pathosystems, affecting pathogen virulence and

epidemiology. Priority effects have been shown to impact disease development and competitive advantage

[43], immune suppression enabling persistence of normally avirulent strains [44], and effector complementation

between strains enhancing disease severity [45]. The challenge of accurately measuring pathogen abundance

in polymicrobial leaf spot diseases has been well established prior to this study [24,25]. Previous studies using

RT-PCR were able to highlight the high frequency of wheat co-infection by *P. nodorum* and *P. tritici-repentis*,

however the work was limited by available technology to measuring relative pathogen abundance [6,27,28].

Our methods represent a significant advancement in the study of polymicrobial plant diseases. Through

species-specific DNA yield normalization, we showed *P. tritici-repentis* yielded almost three times the DNA

per milligram of tissue compared to *P. nodorum*. Using a conserved single copy gene eliminates variability

between strains while the utilization of dPCR retains sufficient sensitivity for field applications. This provided

a strong foundation to investigate the relationship pathogen abundance and visual symptoms, wheat

cultivar disease resistance ratings, and the role of priority effects in the outcome of disease. Our findings

reveal two interactions. The first is co-operative, where *P. tritici-repentis* priority establishment facilitates

subsequent *P. nodorum* colonisation and enhances disease severity and biomass for both pathogens in a

resistant host . The second is competitive, where *P. nodorum* priority establishment suppresses *P. tritici-*

repentis colonisation regardless of host resistance background. To our knowledge, this is the first study to

observe asymmetric facilitation overcoming host resistance in a fungal plant pathosystem.

Field trials where *P. tritici-repentis* was introduced first reproduced facilitation patterns observed in naturally

infected fields. Priority establishment by *P. tritici-repentis* may drive the co-infection observed in wheat

production systems, where *P. tritici-repentis* ascospores have been shown to mature up to three months

before those of *P. nodorum* in south-west Western Australia [46,47]. In addition, *P. tritici-repentis* may benefit

from superior establishment characteristics, including larger resource-rich and temperature-resistant

conidia [48,49]. When *P. tritici-repentis* establishes on resistant host leaves it facilitates *P. nodorum* colonisation

leading to greater disease through an unknown mechanism that leads positive interference where temporal

sequence determines resistance breakdown. *Zymoseptoria tritici* followed by *Pseudomonas syringae*

infection leads to systemically induced bacterial susceptibility in wheat, where fungal infection facilitates

colonization through suppression of immune-related metabolites [50]. Priority effects may exploit depleted

immune responses, where initial infection activates energetically costly defence signalling pathways

creating windows of vulnerability for secondary infections [51]. Necrotrophic pathogens can subvert plant

defence responses that evolved primarily against biotrophic pathogens including the salicylic acid pathway,

which while providing resistance against biotrophs, can enhance susceptibility to necrotrophs [52].

Necrotrophic pathogens create favourable conditions for subsequent colonisation by generating necrotic

lesions with effectors that cause necrosis and provide nutrient-rich substrates from lysed plant cells,

support necrotrophic growth [53]. Our observations are distinguished by the asymmetric nature of this

facilitation despite both pathogens producing necrotrophic effectors infection on wheat [14,39]. This suggests

that mechanistic differences beyond necrotrophic virulence determine the facilitative outcomes we

observed. Previously, *P. nodorum* strains producing high levels of necrotrophic effectors rescued reduced

virulence strains on susceptible wheat, maintaining overall virulence of the pathogen population[45]. Similarly

the introduction reduced virulence mutants of *Magnaporthe oryzae* into populations of wild-type strains

resulted in enhanced overall population fitness and increased damage to rice hosts[11]. These examples

illustrate that cooperation between pathogen strains with different virulence capabilities can lead to

enhanced disease outcomes, representing a distinct mechanism from host immune suppression-

mediated facilitation. The mechanism underlying the facilitation observed in our data remains unclear, but

could involve direct pathogen-pathogen interactions, host immune suppression, or both processes working

in combination[9].

P. nodorum priority establishment antagonises *P. tritici-repentis* colonisation across all host genotypes

observed in the field and laboratory, suggesting a physiological mechanism possessed by *P. nodorum* to

suppress the growth of competitors. While *P. nodorum* and *P. tritici-repentis* share a near-identical ToxA[19],

P. nodorum possesses a diverse collection of characterised necrotrophic effectors including SnTox1,

SnTox3, SnTox5[54] and SnTox267[55]. *Verticillium dahliae* has co-opted antimicrobial proteins specifically for

manipulating fungal competitors during host colonization[56], suggesting exclusion through biochemical

interference may be an evolved strategy rather than an incidental consequence of infection. It is not known

if characterised *P. nodorum* NEs possess antimicrobial functions however, SnTox1 possess dual functions;

facilitating necrosis in wheat and binding chitin, possibly to evade host PAMP-triggered recognition[57]. Recent

pan-genomic analysis of *P. nodorum* identified an abundance of candidate secreted effector proteins,

including numerous homologs to known antimicrobial effectors from other fungal species[38]. In addition, *P.

nodorum* possesses an extensive arsenal of secondary metabolite biosynthesis genes[58]. Previous studies

revealed that *P. nodorum* was capable of producing volatile organic compounds that exhibit phytotoxic and

antimicrobial activities[49,59,60]. It remains to be seen if these secondary metabolites possess bioactivity that

antagonises *P. tritici-repentis*[61].

Molecular quantification revealed competitive exclusion patterns masked by visual assessment. While

necrosis provides a reliable estimate of total pathogen biomass under controlled conditions, visual

symptoms cannot reliably distinguish between single and co-infection states. Additionally, quantification of

disease can be hindered by simultaneous biotic and abiotic stresses, emphasising the need for molecular

approaches that quantify pathogens biomass [62]. Even with automated image analysis tools like LeafScan,

which can accurately quantify necrotic tissue area, morphologically indistinguishable lesions produced by

P. nodorum and *P. tritici-repentis* provide no visual indication of their microbial composition. The correlation

observed under controlled infection suggests that both pathogens primarily colonise necrotic tissue

regions, likely secreting necrotrophic effectors into surrounding tissue to induce chlorosis ahead of

colonisation, supporting previous mechanistic studies [63].

These sequence-dependent interactions represent a form of historical contingency where early colonisers

fundamentally alter the trajectory of microbial community development within host tissues [1]. The facilitative

and competitive mechanisms we observed demonstrate that pathogen communities are not simply

assembled through independent colonisation events, but rather through complex ecological networks

where arrival order determines interaction outcomes [5]. This challenges traditional views of agricultural

pathosystems as simplified host-pathogen dyads and instead positions them as dynamic microbial

ecosystems governed by community assembly rules similar to those operating in natural systems [2], but with

accelerated temporal dynamics due to the intensive selection pressures of agricultural environments [3].

Agricultural monocultures may thus serve as model systems for understanding rapid microbial community

assembly processes under simplified environmental conditions, offering insights into fundamental

ecological principles that operate across diverse ecosystems while providing practical applications for

sustainable disease management strategies.

Several priorities emerge from this work. We recommend that future disease resistance trials of commercial

wheat cultivars consider rating for the SNB-TS complex over the current single disease resistance rating

system in wheat-growing regions where *P. nodorum* and *P. tritici-repentis* are endemic. Furthermore,

additional analysis of the effect that co-infection may have on grain yields is needed. Field validation studies

across diverse environments and wheat genetic backgrounds are needed to determine the generalisability

of our sequence-dependent findings, particularly investigating whether similar priority effects occur in

cultivars with different resistance backgrounds.

Acknowledgements

The authors would like to thank; Noel Knight (Curtin University) and Lincoln Harper (Curtin University) for

assistance with dPCR assay design; Julie Lawrence (Curtin University) and Manisha Shankar (DPIRD) for

providing off-target isolates for dPCR validation. We extend our thanks to Dr Araz Abdullah (Curtin University)

for insightful discussions and for establishing the precursor groundwork for this study. This study was

conducted by the Centre for Crop and Disease Management, a co-investment between the Grains

Research and Development Corporation (GRDC) and Curtin University - grant CUR1403-002BLX.

Data Accessibility Statement

All data supporting the results of this study are available as follows: Raw dPCR data, image analysis outputs,

and statistical analysis code are archived in the Dryad Digital Repository at

[https://doi.org/10.5061/dryad.mcvdnckch]. The LeafScan image processing and analysis tools are freely

available at https://github.com/LeonLenzo/leaf-scan. Reference genome assemblies for

Parastagonospora nodorum SN15 and *Pyrenophora tritici-repentis* M4 are publicly available through NCBI

GenBank. Primer and probe sequences for the duplex dPCR assay are provided in the Materials and

Methods section. Field sampling coordinates and experimental metadata are included in the

supplementary materials. All materials and protocols are available upon reasonable request to the

corresponding author.

References

1. Fukami T. Historical Contingency in Community Assembly: Integrating Niches, Species Pools, and Priority Effects. *Annu. Rev. Ecol. Evol. Syst.* 2015 **46**:1–23. https://doi.org/10.1146/annurev-ecolsys-110411-160340.

2. Vacher C, A Hampe, AJ Porté, et al. The Phyllosphere: Microbial Jungle at the Plant-Climate Interface. *Annu. Rev. Ecol. Evol. Syst.* 2016 **47**:1–24. https://doi.org/10.1146/annurev-ecolsys-121415-032238.

3. McDonald BA, EH Stukenbrock. Rapid emergence of pathogens in agro-ecosystems: global threats to agricultural sustainability and food security. *Philos. Trans. R. Soc. B Biol. Sci.* 2016 **371**. https://doi.org/10.1098/rstb.2016.0026.

4. Halliday FW, RM Penczykowski, B Barrès, et al. Facilitative priority effects drive parasite assembly under coinfection. *Nat. Ecol. Evol.* 2020 **4**:1510–21. https://doi.org/10.1038/s41559-020-01289-9.

5. Debray R, RA Herbert, AL Jaffe, et al. Priority effects in microbiome assembly. *Nat. Rev. Microbiol.* 2022 **20**:109–21. https://doi.org/10.1038/s41579-021-00604-w.

6. Fejer Justesen A, B Corsi, A Ficke, et al. Hidden in plain sight: a molecular field survey of three wheat leaf blotch fungal diseases in North-Western Europe shows co-infection is widespread. *Eur. J. Plant Pathol.* 2021 **160**:949–62. https://doi.org/10.1007/S10658-021-02298-5.

7. Van der Heyden H, P Dutilleul, J-B Charron, et al. Monitoring airborne inoculum for improved plant disease management. A review. *Agron. Sustain. Dev.* 2021 **41**:40–40. https://doi.org/10.1007/s13593-021-00694-z.

8. Ferreira RBR, LCM Antunes, N Sal-Man. Pathogen-pathogen interactions during co-infections. *ISME J.* 2025 **19**:wraf104. https://doi.org/10.1093/ismejo/wraf104.

9. Dutt A, D Andrivon, C Le May. Multi-infections, competitive interactions, and pathogen coexistence. *Plant Pathol.* 2022 **71**:5–22. https://doi.org/10.1111/ppa.13469.

10. Harrison F, LE Browning, M Vos, et al. Cooperation and virulence in acute Pseudomonas aeruginosa infections. *BMC Biol.* 2006 **4**:21. https://doi.org/10.1186/1741-7007-4-21.

11. Lindsay RJ, MJ Kershaw, BJ Pawlowska, et al. Harbouring public good mutants within a pathogen population can increase both fitness and virulence. *eLife* 2016 **5**:e18678. https://doi.org/10.7554/eLife.18678.

12. Mousa WK, A Schwan, J Davidson, et al. An endophytic fungus isolated from finger millet (Eleusine coracana) produces anti-fungal natural products. *Front. Microbiol.* 2015 **6**:1157. https://doi.org/10.3389/fmicb.2015.01157.

13. Orton ES, JKM Brown. Reduction of growth and reproduction of the biotrophic fungus Blumeria graminis in the presence of a necrotrophic pathogen. *Front. Plant Sci.* 2016 **7**:742–742. https://doi.org/10.3389/FPLS.2016.00742/BIBTEX.

14. Gourlie R, M McDonald, M Hafez, et al. The pangenome of the wheat pathogen Pyrenophora tritici-repentis reveals novel transposons associated with necrotrophic effectors ToxA and ToxB. *BMC Biol.* 2022 **20**. https://doi.org/10.1186/s12915-022-01433-w.

15. McDonald BA. How knowledge of pathogen population biology informs management of Septoria nodorum blotch on wheat. *Eur. J. Plant Pathol.* 2025 **171**:531–45. https://doi.org/10.1007/s10658-024-02996-w.

16. Abdullah AS, MR Gibberd, J Hamblin. Co-infection of wheat by Pyrenophora tritici-repentis and Parastagonospora nodorum in the wheatbelt of Western Australia. *Crop Pasture Sci.* 2020 **71**:119–27. https://doi.org/10.1071/CP19412.

17. Möller M, EH Stukenbrock. Evolution and genome architecture in fungal plant pathogens. *Nat. Rev. Microbiol.* 2017 **15**:756–71. https://doi.org/10.1038/nrmicro.2017.76.

18. McDonald MC, PS Solomon. Just the surface: advances in the discovery and characterization of necrotrophic wheat effectors. *Curr. Opin. Microbiol.* 2018 **46**:14–8. https://doi.org/10.1016/J.MIB.2018.01.019.

19. Friesen TL, EH Stukenbrock, Z Liu, et al. Emergence of a new disease as a result of interspecific virulence gene transfer. *Nat. Genet. 2006 388* 2006 **38**:953–6. https://doi.org/10.1038/ng1839.

20. Vleeshouwers VGAA, RP Oliver. Effectors as tools in disease resistance breeding against biotrophic, hemibiotrophic, and necrotrophic plant pathogens. *Mol. Plant. Microbe Interact.* 2014 **27**:196–206. https://doi.org/10.1094/MPMI-10-13-0313-IA.

21. Francki MG, E Walker, CJ McMullan, et al. Multi-Location Evaluation of Global Wheat Lines Reveal Multiple QTL for Adult Plant Resistance to Septoria Nodorum Blotch (SNB) Detected in Specific Environments and in Response to Different Isolates. *Front. Plant Sci.* 2020 **11**. https://doi.org/10.3389/fpls.2020.00771.

22. Shankar M, D Mather, H Golzar, et al. *Stacking yellow spot resistance genes into fixed lines results in genetic gain*. 2015. https://www.researchgate.net/publication/305748820.

23. Peters Haugrud AR, G Shi, S Seneviratne, et al. Genome-wide association mapping of resistance to the foliar diseases septoria nodorum blotch and tan spot in a global winter wheat collection. *Mol. Breed. New Strateg. Plant Improv.* 2023 **43**:54. https://doi.org/10.1007/s11032-023-01400-5.

24. Beard C, G Thomas, A Hills, et al. *Yellow spot and septoria nodorum blotch and their management in wheat*. 2024. https://library.dpird.wa.gov.au/fc_factsheets/70.

25. Friskop A, Z Liu. *Fungal Leaf Spot Diseases of Wheat: Tan spot, Septoria/Stagonospora nodorum blotch and Septoria tritici blotch (PP1249)*. 2021. www.ag.ndsu.edu/extension.

26. Long M, M Hartley, RJ Morris, et al. Classification of wheat diseases using deep learning networks with field and glasshouse images. *Plant Pathol.* 2023 **72**:536–47. https://doi.org/10.1111/ppa.13684.

27. Abdullah AS, C Turo, CS Moffat, et al. Real-time PCR for diagnosing and quantifying co-infection by two globally distributed fungal pathogens of wheat. *Front. Plant Sci.* 2018 **9**:1086–1086. https://doi.org/10.3389/FPLS.2018.01086/BIBTEX.

28. Fu H, J Jiang, MW Harding, et al. Development of a triplex qPCR system for the detection of Parastagonospora nodorum, Zymoseptoria tritici and Pyrenophora tritici-repentis from wheat. *Plant Dis.* 2025. https://apsjournals.apsnet.org/doi/10.1094/PDIS-01-25-0003-SR (18 August 2025, date last accessed).

29. Moffat CS, PT See, RP Oliver. Leaf yellowing of the wheat cultivar Mace in the absence of yellow spot disease. *Australas. Plant Pathol.* 2015 **44**:161–6. https://doi.org/10.1007/s13313-014-0335-2.

30. Hall AT, AM Zovanyi, DR Christensen, et al. Evaluation of Inhibitor-Resistant Real-Time PCR Methods for Diagnostics in Clinical and Environmental Samples. *PLOS ONE* 2013 **8**:e73845. https://doi.org/10.1371/journal.pone.0073845.

31. Huggett JF, CA Foy, V Benes, et al. The digital MIQE guidelines: Minimum information for publication of quantitative digital PCR experiments. *Clin. Chem.* 2013 **59**:892–902. https://doi.org/10.1373/clinchem.2013.206375.

32. Solomon PS, SW Thomas, P Spanu, et al. The utilisation of di/tripeptides by Stagonospora nodorum is dispensable for wheat infection. *Physiol. Mol. Plant Pathol.* 2003 **63**:191–9. https://doi.org/10.1016/j.pmpp.2003.12.003.

33. Moolhuijzen P, PT See, CS Moffat. A new PacBio genome sequence of an Australian Pyrenophora tritici-repentis race 1 isolate. *BMC Res. Notes* 2019 **12**:642–642. https://doi.org/10.1186/s13104-019-4681-6.

34. Rybak K, PT See, HTT Phan, et al. A functionally conserved Zn2Cys6 binuclear cluster transcription factor class regulates necrotrophic effector gene expression and host-specific virulence of two major Pleosporales fungal pathogens of wheat. *Mol. Plant Pathol.* 2017 **18**:420–34. https://doi.org/10.1111/mpp.12511.

35. Liu ZH, JD Faris, SW Meinhardt, et al. Genetic and physical mapping of a gene conditioning sensitivity in wheat to a partially purified host-selective toxin produced by Stagonospora nodorum. *Phytopathology* 2004 **94**:1056–60.

36. Jacques S, L Lenzo, K Stevens, et al. An optimized sporulation method for the wheat fungal pathogen Pyrenophora tritici-repentis. *Plant Methods* 2021 **17**:52–52. https://doi.org/10.1186/s13007-021-00751-4.

37. Abràmoff MD, PJ Magalhães, SJ Ram. Image processing with ImageJ. *Biophotonics Int.* 2004 **11**:36–42.

38. Jones DAB, K Rybak, M Hossain, et al. Repeat-induced point mutations driving Parastagonospora nodorum genomic diversity are balanced by selection against non-synonymous mutations. *Commun. Biol.* 2024 **7**:1614. https://doi.org/10.1038/s42003-024-07327-7.

39. Syme RA, K-C Tan, K Rybak, et al. Pan-Parastagonospora comparative genome analysis—effector
prediction and genome evolution. *Genome Biol. Evol.* 2018 **10**:2443–57.

40. Moolhuijzen PM, PT See, G Shi, et al. A global pangenome for the wheat fungal pathogen Pyrenophora
tritici-repentis and prediction of effector protein structural homology. *Microb. Genomics* 2022
8:mgen000872. https://doi.org/10.1099/mgen.0.000872.

41. R Core Team. R: A Language and Environment for Statistical Computing. 2025.

42. Keene ON. The log transformation is special. *Stat. Med.* 1995 **14**:811–9.
https://doi.org/10.1002/sim.4780140810.

43. Suffert F, G Delestre, F Carpentier, et al. Fashionably late partners have more fruitful encounters: Impact
of the timing of co-infection and pathogenicity on sexual reproduction in Zymoseptoria tritici. *Fungal Genet.*
Biol. 2016 **92**:40–9. https://doi.org/10.1016/J.FGB.2016.05.004.

44. Barrett LG, M Zala, A Mikaberidze, et al. Mixed infections alter transmission potential in a fungal plant
pathogen. *Environ. Microbiol.* 2021 **23**:2315–30. https://doi.org/10.1111/1462-2920.15417.

45. John E, S Jacques, HTT Phan, et al. Variability in an effector gene promoter of a necrotrophic fungal
pathogen dictates epistasis and effector-triggered susceptibility in wheat. *PLOS Pathog.* 2022 **18**:e1010149.

46. Bathgate JA, R Loughman. Ascospores are a source of inoculum of Phaeosphaeria nodorum, P. Avenaria
f. sp. Avenaria and Mycosphaerella graminicola in Western Australia. *Australas. Plant Pathol.* 2001 **30**:317–
22. https://doi.org/10.1071/AP01043.

47. Galloway J, M Salam, P Payne. *Less yellow spot in wheat: progress towards a decision support tool that*
will predict when yellow spot spores are released from the previous season stubble. 2016.

48. Benslimane H, S Aouali, A Khalfi, et al. In vitro morphological characteristics of Pyrenophora tritici-
repentis isolates from several Algerian agro-ecological zones. *Plant Pathol. J.* 2017 **33**:109–17.
https://doi.org/10.5423/PPJ.OA.09.2015.0189.

49. Muria-Gonzalez MJ, Y Yeng, S Breen, et al. Volatile Molecules Secreted by the Wheat Pathogen
Parastagonospora nodorum Are Involved in Development and Phytotoxicity. *Front. Microbiol.* 2020 **11**.
https://doi.org/10.3389/fmicb.2020.00466.

50. Seybold H, TJ Demetrowitsch, MA Hassani, et al. A fungal pathogen induces systemic susceptibility and
systemic shifts in wheat metabolome and microbiome composition. *Nat. Commun.* 2020 **11**.
https://pubmed.ncbi.nlm.nih.gov/32313046/.

51. Balint-Kurti P. The plant hypersensitive response: concepts, control and consequences. *Mol. Plant*
Pathol. 2019 **20**:1163–78. https://doi.org/10.1111/MPP.12821.

52. Glazebrook J. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens.
Annu. Rev. Phytopathol. 2005 **43**:205–27. https://doi.org/10.1146/annurev.phyto.43.040204.135923.

53. Tan KC, RP Oliver. Regulation of proteinaceous effector expression in phytopathogenic fungi. *PLoS*
Pathog. 2017 **13**. https://pubmed.ncbi.nlm.nih.gov/28426760/.

54. Kariyawasam GK, JK Richards, NA Wyatt, et al. The Parastagonospora nodorum necrotrophic effector
SnTox5 targets the wheat gene Snn5 and facilitates entry into the leaf mesophyll. *New Phytol.* 2022 **233**:409–
26. https://doi.org/10.1111/nph.17602.

55. Richards JK, GK Kariyawasam, S Seneviratne, et al. A triple threat: the Parastagonospora nodorum
SnTox267 effector exploits three distinct host genetic factors to cause disease in wheat. *New Phytol.* 2022
233:427–42. https://doi.org/10.1111/nph.17601.

56. Snelders NC, GC Petti, GCM van den Berg, et al. An ancient antimicrobial protein co-opted by a fungal
plant pathogen for in planta mycobiome manipulation. *Proc. Natl. Acad. Sci.* 2021 **118**:e2110968118–
e2110968118. https://doi.org/10.1073/pnas.2110968118.

57. Liu Z, Y Gao, YM Kim, et al. SnTox1, a Parastagonospora nodorum necrotrophic effector, is a dual-
function protein that facilitates infection while protecting from wheat-produced chitinases. *New Phytol.*
2016 **211**:1052–64. https://doi.org/10.1111/nph.13959.

58. Chooi YH, MJ Muria-Gonzalez, PS Solomon. A genome-wide survey of the secondary metabolite
biosynthesis genes in the wheat pathogen Parastagonospora nodorum. *Mycology* 2014 **5**:192–206.
https://doi.org/10.1080/21501203.2014.928386.

59. Chooi YH, C Krill, RA Barrow, et al. An In planta-expressed polyketide synthase produces (R)-mellein in
the wheat pathogen Parastagonospora nodorum. *Appl. Environ. Microbiol.* 2015 **81**:177–86.
https://doi.org/10.1128/AEM.02745-14.
60. Chooi YH, G Zhang, J Hu, et al. Functional genomics-guided discovery of a light-activated phytotoxin in
the wheat pathogen Parastagonospora nodorum via pathway activation. *Environ. Microbiol.* 2017 **19**:1975–
86. https://doi.org/10.1111/1462-2920.13711.
61. Robey MT, LK Caesar, MT Drott, et al. An interpreted atlas of biosynthetic gene clusters from 1,000 fungal
genomes. *Proc. Natl. Acad. Sci.* 2021 **118**:e2020230118. https://doi.org/10.1073/pnas.2020230118.
62. Bock CH, JGA Barbedo, EM Del Ponte, et al. From visual estimates to fully automated sensor-based
measurements of plant disease severity: status and challenges for improving accuracy. *Phytopathol. Res.*
2020 **2**. https://doi.org/10.1186/s42483-020-00049-8.
63. Faris JD, Z Zhang, H Lu, et al. A unique wheat disease resistance-like gene governs effector-triggered
susceptibility to necrotrophic pathogens. *Proc. Natl. Acad. Sci.* 2010 **107**:13544–9.
https://doi.org/10.1073/pnas.1004090107.
64. Shackley B, C Zaicou-Kunesch, H Dhammu, et al. *Wheat Variety Guide for WA Wheat variety guide for*
WA 2014. 2014.
65. Shackley B, D Nicol, J Curry, et al. *Department of Primary Industries and Regional Development Crop*
Sowing Guide 2022. 2022.
66. Bertazzoni S, DAB Jones, HT Phan, et al. Chromosome-level genome assembly and manually-curated
proteome of model necrotroph Parastagonospora nodorum Sn15 reveals a genome-wide trove of
candidate effector homologs, and redundancy of virulence-related functions within an accessory
chromosome. *BMC Genomics* 2021 **22**:382–382. https://doi.org/10.1186/s12864-021-07699-8.
67. McDonald MC, AP Taranto, E Hill, et al. Transposon-mediated horizontal transfer of the host-specific
virulence protein ToxA between three fungal wheat pathogens. *mBio* 2019 **10**.
https://doi.org/10.1128/mBio.01515-19.

Figure 6: Sequential Seedling Infection

(A) Axe and **(B)** Scepter. Bars represent mean (±SE) $\log_{10}$ biomass (µg) for *Pn* (orange) and *Ptr* (blue), with

Necrosis % (brown) plotted on the secondary y-axis. From left to right, Single pathogen treatments: Ptr → Ø

Ptr → Ptr, Pn → Ø, Pn → Pn, Co-infection treatments: Pn + Ptr, Pn → Ptr, Ptr → Pn. Letters inside bars indicate

statistical groupings within each series (Tukey HSD, α = 0.05). Asterisks above bars indicate significant

differences between *Pn* and *Ptr* biomass within co-infection treatments (paired t-test: ns = p ≥ 0.05, * = p <

0.05, ** = p < 0.01,). n = 3 per treatment.

Figure 5: Natural vs defined field inoculum

(A) necrosis severity (necrosis %, brown), **(B)** *P. nodorum* biomass (orange), and **(C)** *P. tritici-repentis*

biomass (blue) across four field sites. Natural (naturally occurring infection in an endemic field Figure 4),

Mixed (equal mix of *P. nodorum* and *P. tritici-repentis* infected stubble), *Pn (P. nodorum* infected stubble)

and *Ptr (P. tritici-repentis* infected stubble). Points represent wheat leaf samples with shapes indicating

wheat lines (circles = Axe, squares = Emu, diamonds = Scepter, triangles = Yitpi). Statistical significance of

differences between co-infection and single infection within each site indicated above brackets (ns = not

significant, * p < 0.05, ** p < 0.01, *** p < 0.001).

Figure 4: Host genetics in natural field disease

(A) disease severity (necrosis %, brown), **(B)** *P. nodorum* biomass (orange), and **(C)** *P. tritici-repentis* (blue)

biomass across four wheat lines: Axe (SVS to SNB and TS), Emu Rock (SVS to SNB, MRMS to TS), Scepter

(MRMS to SNB and TS), and Yitpi (MRMS to SNB SVS to TS). Points represent individual wheat leaves with

shapes indicating cultivar (circles = Axe, squares = Emu, diamonds = Scepter, triangles = Yitpi). Statistical

significance of differences between co-infection and single infection within each wheat line indicated above

brackets (ns = not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

A

51%	26%	24%
90	43	30	0

B

Total.Disease
(R^2 = 0.445)

Chlorosis
(R^2 = 0.214)

Necrosis
(R^2 = 0.705)

Log_{10} Biomass (µg)

LAA (%)

○ Axe   ◇ Scepter   100% Pn   100% Ptr

C

R^2 Value

Jul   Aug   Sep   Oct

● Chlorosis
● Necrosis

Figure 3: Disease biomass correlation

(A) Automated leaf segmentation showing original image, processed output with disease classification

(green = healthy, yellow = chlorotic, brown = necrotic), and HSV hue thresholds. **(B)** Correlations between

pathogen biomass and leaf area affected (LAA) by different visual disease metrics. Points represent Axe

(susceptible, circles) and Scepter (resistant, diamonds) varieties with single or mixed infections. Colour

gradient indicates *P. nodorum* (orange) to *P. tritici-repentis* (blue) proportion. R^2 shows correlation strength

for total disease, r shows correlation co-efficient. **(C)** Temporal variation in visual disease-biomass

correlations under field conditions over a 16-week sampling period. Lines show R^2 values for chlorosis

(yellow), necrosis (brown) metrics.

A

B

C

627 ***Figure 2: dPCR Validation***

628 **(A)** Standard curves showing linear relationship (β) between DNA yield for *P. tritici-repentis* (blue) and *P.*

629 *nodorum* (orange). **(B)** Standard curves for Ptr_HEX (blue) and Pn_FAM (orange) probes showing linear

630 relationship (α) between DNA concentration and copy number. **(C)** Specificity testing of the dPCR assay

631 against gDNA from wheat host and related fungal species, showing probe-specific amplification patterns.

A

B

C

D

Figure 1: dPCR Design

(A) Pairwise alignment of PnSN15_01762 (Orange) and PtrM4_098590 (Blue), the orthologues for alpha

tubulin. Locus aT is highlighted in red. **(B)** Primer and probe design showing aT_F and aT_R amplification

primers (green arrows) with species-specific probes Ptr_HEX (blue) and Pn_FAM (orange). Coloured bars

indicate sequence polymorphisms. **(C)** Percentage Identity of the aT region across pan-genome

assemblies. **(D)** Heatmap showing sequence identity between primer/probe sequences and organisms in

the Refseq fungi database.

Table 2

Australian wheat cultivars used in this study and their disease resistance ratings [64,65].
SVS, susceptible-very susceptible; MRMS, moderately resistant-moderately
susceptible

Wheat cultivar	SNB Rating	TS Rating
Axe	SVS	SVS
Emu Rock	SVS	MRMS
Scepter	MRMS	MRMS
Yitpi	MRMS	SVS

Table 1

Oligonucleotides used in this study.

Oligonucleotide	Sequence (5' – 3')
aT_F	AGGAACCTCGGCATTGAGC
aT_R	ACCAAGTTGGTCTGGAACTCG
Pn_FAM	CGGATCCCTGAACGTCGATCTC
Ptr_HEX	ATGTAGCTCCATTCTTCTCTCACTTGC