A *de novo* reference genome of the golden jackal, *Canis aureus*

Sven Winter[1,2,3,*], René Meißner[1], Stefan Prost[4,5], Carola Greve[6], Charlotte Gerheim[6],

Marine Arakelyan[7], Sargis Aghayan[8], Astghik Ghazaryan[7], Jennifer Hatlauf[9], Pamela A.

Burger[1,*]

[1]Research Institute of Wildlife Ecology, University of Veterinary Medicine Vienna, Vienna, Austria

[2]Faculty of Science and Technology, University of the Faroe Islands, Tórshavn, The Faroe Islands

[3]Senckenberg Biodiversity and Climate Research Centre, Frankfurt am Main, Germany

[4]Ecology and Genetics Research Unit, University of Oulu, Oulu, Finland

[5]Natural History Museum Vienna, Central Research Laboratories, Vienna, Austria

[6]Senckenberg Research Institute and Museum, Frankfurt am Main, Germany

[7]Institute of Zoology, Yerevan State University, Yerevan, Armenia

[8]Scientific Center of Zoology and Hydroecology, National Academy of Sciences of the Armenian

State

[9]Institute of Wildlife Biology and Game Management, BOKU University, Vienna, Austria

Contact for correspondence*:

Sven Winter, sven.winter@vetmeduni.ac.at

Pamela Burger, pamela.burger@vetmeduni.ac.at

Abstract

The golden jackal (*Canis aureus*) is rapidly expanding its range in Europe, driven by

climate and habitat changes, human influence, and changes in competition with

wolves. Its ecological flexibility enables it to thrive in various habitats, including urban

areas, raising concerns about its potential role in spreading zoonotic diseases. Jackals

may act as reservoirs for pathogens such as Lyme disease and babesiosis, affecting

wildlife, humans, and pets. Their close genetic relationship with domestic dogs also

increases the risk of hybridization and host-jumping, complicating disease dynamics.

To better understand their dispersal ability and host-pathogen dynamics, we present

the first chromosome-level genome assembly of the golden jackal, generated using

PacBio HiFi sequencing and reference-based scaffolding. The final assembly has a total

length of 2.53 Gb in 325 scaffolds, with 98.41% of the sequence anchored to the

expected 38+XY chromosomes. The assembly shows high contiguity, with scaffold and
contig N50 values of 68.03 Mb and 56.64 Mb, respectively. Annotation revealed 26,084
protein-coding genes, and repetitive elements account for 40.58% of the total
assembly.

This high-quality reference genome provides an essential resource for studying the
genetic basis of the golden jackal's adaptation, ecological interactions, and potential as
a zoonotic reservoir. It also supports efforts to monitor population expansion and its
effects on ecosystems. By advancing our understanding of golden jackal genetics, this
work enables future research on evolution, host-pathogen dynamics, and the broader
consequences of wildlife dispersal in a rapidly changing environment.

Keywords

Canidae, Zoonoses, Range expansion, Climate change, Pathogen transmission,
Wildlife–human interaction

Introduction

Canids (*Canidae*) are a diverse group of mostly carnivorous mammals that exhibit a
wide range of ecological adaptations and social behaviors (Allen, 2017). The
approximately 35 species are distributed across all continents except Antarctica,
highlighting their ability to thrive in vastly different environments (Krofel et al., 2022).
One such species, the golden jackal (*Canis aureus* LINNAEUS, 1758), is a mid-sized
canid that has been experiencing an accelerated natural dispersal, particularly within
Western Asia and into Europe (Hatlauf et al., 2021). This expansion is thought to be
driven by a combination of several factors, including climate change, shifts in habitat,
land-use changes, and other human activities, all of which have facilitated its range
expansion (Krofel et al., 2017; Spassov & Acosta-Pankov, 2019).

The golden jackal's notable adaptability allows it to occupy a wide array of
habitats, ranging from forests and grasslands to urban environments (Salek et al., 2014;
Torretta et al., 2020). Its opportunistic feeding habits, which include a broad diet of

small mammals, birds, insects, fruits, and vegetables, make it highly resilient and capable of colonizing new areas (Lanszki et al., 2022). While the presence of golden jackals in new ecosystems can provide benefits, such as natural pest control by regulating populations of rodents and other small vertebrates (Lanszki et al., 2016), the species' expansion also presents potential challenges (Gherman & Mihalca, 2017). As jackals move into new areas, they may compete with native species for resources and potentially disrupt local ecosystems (Krofel et al., 2017; Serva et al., 2023). Additionally, the golden jackal's capacity to survive in urban environments brings it into closer proximity with humans, which increases the likelihood of interaction with domestic animals, raising concerns about zoonotic disease transmission (Hatlauf, 2024; Lapid et al., 2023).

Figure 1. Golden jackal (*Canis aureus*) ©Jennifer Hatlauf

Recent studies have focused on the potential public health risks associated with the movement of golden jackals, especially in light of the discovery of jackals in Denmark carrying tick vectors and zoonotic pathogens (Klitgaard et al., 2017). Pathogens, such as those transmitted by ticks, can pose significant risks to both wildlife

and human health, which underscores the importance of monitoring the spread of this

species. The potential for golden jackals to carry and transmit diseases like Lyme

disease and babesiosis demonstrates how their expanding range could influence the

epidemiology of zoonoses, particularly in regions where the jackal may interact with

domestic animals and humans (Lymbery et al., 2014). Furthermore, the close

phylogenetic relationship between golden jackals and domestic dogs (*Canis lupus*

familiaris) raises the possibility of hybridization and host jumping, which could

exacerbate the spread of zoonotic diseases or introduce novel pathogens into local

ecosystems (Galov et al., 2015; Gherman & Mihalca, 2017; Stefanovic et al., 2024).

Given the important ecological role golden jackals play in their current and future

habitats, they serve as important models for understanding climate-mediated dispersal

of zoonotic diseases. Their expansion is a good example of how environmental changes

and human-induced factors influence the movement of wildlife and the pathogens they

carry. Studying jackals offers valuable insights into the broader dynamics of host-

pathogen coevolution and the transmission of diseases across species boundaries. To

support these efforts, a comprehensive genomic resource is essential.

In this study, we present the first chromosome-level genome assembly of the

golden jackal, generated using PacBio HiFi long-read sequencing, supplemented by

reference-based scaffolding. This high-quality reference genome provides an invaluable

tool for investigating the genetic basis of the species' adaptation to new environments

and its interactions with both its ecosystem and pathogens. It also enables more

effective monitoring of the species and the diseases it may carry, facilitating a better

understanding of how climate change and landscape fragmentation contribute to the

spread of zoonotic diseases. By advancing our genetic knowledge of golden jackals, we

provide a foundation for future studies on the evolution, population dynamics, and

ecological impacts of this increasingly prominent species.

Materials and Methods

Sampling and Sequencing

Tissue samples from a subadult male golden jackal (sample ID: CA_001; PathoNo.:

SK/123/23) shot by a hunter in Villach, Carinthia, Austria, were collected during post-

mortem examination at the University of Veterinary Medicine Vienna, Austria, on

06.02.2023 and stored frozen at -80°C. High molecular weight DNA was extracted from

spleen tissue using the innuPrep SE HMW kit (IST Innuscreen GmbH, Germany)

according to the manufacturer's recommendations. The quantity and quality of the DNA

were evaluated using the QuantiFluor ONE dsDNA System on the Quantus Fluorometer

(Promega Corporation, USA) and the Genomic DNA Screen Tape on the Agilent 4150

TapeStation system (Agilent Technologies, Inc., USA), respectively.

Library preparation for Pacific Biosciences of California, Inc (PacBio) circular consensus

sequencing (CCS) was prepared using the SMRTbell Express Prep Kit 3.0 (PacBio, USA)

and sequenced on the PacBio Revio system.

Assembly and QC

The PacBio-generated raw data BAM file was converted into FASTQ format using

BAM2fastx   v.1.3.0,   a   PacBio   Secondary   Analysis   Tool

(https://github.com/PacificBiosciences/pbbioconda). The initial assemblies, including

two, which resemble the two potential parental haplotypes, were generated with

hifiasm v.0.24.0 (RRID:SCR_021069) (Cheng et al., 2021) using standard parameters. As

sample quality was insufficient for a proximity-ligation library, we did not aim for a

haplotype-resolved assembly but instead continued with the pseudohaploid assembly

for the next steps. As we were unable to use Hi-C data for the scaffolding, due to the

fact that the sample was stored in ethanol, we instead used the Mexican gray wolf

assembly (Accession No. GCF_048164855.1) for a reference-based scaffolding with

RagTag v.2.1.0 (Alonge et al., 2022). Subsequently, gaps in the scaffolded assembly

were breached using the original PacBio long-read data with TGS_gapcloser v.1.2.1

(RRID:SCR_017633)(Xu et al., 2020). The mitochondrial genome was assembled using

the   MitoHiFi   workflow   (RRID:SCR_026369)   in   Galaxy   v.3.2.3

(RRID:SCR_006281)(Blankenberg et al., 2014; Uliano-Silva et al., 2023).

Assembly Quality Assessment

Assembly statistics to evaluate contiguity of the assembly were generated with Quast

v.5.0.2 (RRID:SCR_001228)(Gurevich et al., 2013). The completeness of the assembly

was   assessed   by   gene   set   completeness   analyses   using   BUSCO   v.5.8.3

(RRID:SCR_015008)   (Manni   et   al.,   2021)   and   compleasm   v.0.2.7

(RRID:SCR_026370)(Huang & Li, 2023) against the carnivora_odb12 orthologs dataset,

as   well   as   based   on   k-mers   (k=21)   using   Merqury   v.1.3   with   Meryl   v.   1.3

(RRID:SCR_022964)(Rhie et al., 2021). Minimap2 v.2.28 (RRID:SCR_018550)(Li, 2018)

and samtools v.1.21 (RRID:SCR_002105)(Danecek et al., 2021) were used to map the

PacBio reads to the final assembly and sort the mapping file to check the mapping rate

with   Qualimap   bamqc   v.2.3   (RRID:SCR_001209)(Okonechnikov   et   al.,   2016).   The

mapping   file   was   also   used   together   with   the   results   from   a   Diamond   v.2.1.10

(RRID:SCR_016071)(Buchfink et al., 2021) search against Swiss-Prot (release 2025-01)

(The UniProt Consortium, 2019) in Blobtools v. 1.1 (RRID:SCR_017618)(Laetsch &

Blaxter, 2017) to check for potential contamination.

Annotation

Repeats in the assembly were annotated with RepeatMasker v.4.1.0

(RRID:SCR_012954) (Smit et al., 2015) using a custom *de novo* repeat library generated

from the presented assembly with RepeatModeler v.2.0.1 (RRID:SCR_015027) (Flynn et

al., 2020) combined with the Canidae specific repeats from Dfam 3.1 and RepBase

(release 20181026) (Bao et al., 2015; Storer et al., 2021). Interspersed repeats were

hardmasked (Ns) while simple repeats were softmasked (lowercase) in preparation for

the gene annotation.

Genes were predicted using the homology-based approach implemented in the

GeMoMa pipeline v.1.7.1 (RRID:SCR_017646) (Keilwagen et al., 2016, 2018) using

MMseqs2 v. 13.45111 (RRID:SCR_022962) (Steinegger & Söding, 2017) as an alignment

tool. The following six mammalian genomes and corresponding annotations were used

as references: *Homo sapiens* (GCF_000001405.40), *Mus musculus*

(GCF_000001635.27), *Canis lupus familiaris* (German Shepherd, GCF_011100685.1;

Wang et al., 2021), *Vulpes vulpes* (GCF_048418805.1), *Vulpes lagopus*

(GCF_018345385.1;, *Canis lupus baileyi* (GCF_048164855.1; Rhie et al., 2021).

Subsequently, the proteins predicted by GeMoMa were functionally annotated by a

BLASTP v.2.11.0+ (RRID:SCR_001010) (Camacho et al., 2009) search against the Swiss-

Prot database (RRID:SCR_002380; release 2025-01) (The UniProt Consortium, 2019)

with an e-value cutoff of 10^{-6}. Gene ontology (GO) terms, domains, and motifs were

further annotated using InterProScan v.5.59.91 (RRID:SCR_005829) (Jones et al., 2014;

Quevillon et al., 2005). BUSCO v.5.8.3 (RRID:SCR_015008) (Manni et al., 2021) and

compleasm v.0.2.7 (RRID:SCR_026370) (Huang & Li, 2023) were used to evaluate gene

completeness in reference to the carnivora_odb12 dataset.

Table 1. Software and versions used to generate the *Canis aureus* assembly.

Pipeline Step	Software	Version
PacBio BAM to FASTQ	BAM2fastx	v.1.3.0
assembly & long-read polishing	hifiasm	v.0.24.0
Scaffolding	RagTag	v.2.1.0
Gap-closing	TGS-GapCloser	v.1.2.1
Mitochondrial genome assembly	MitoHiFi in Galaxy	v.3.2.3
Assembly statistics	Quast	v.5.0.2
Gene set completeness	BUSCO	v.5.8.3
	compelasm	v.0.2.7
Assembly completeness	Merqury	v.1.3
Mapping of long reads	minimap2	v.2.28
Sort & index BAM	samtools	v.1.21
Mapping statistics	QualiMap	v.2.3
Contamination analyses	Diamond	v.2.1.10
	Blobtools	v.1.1
Repeat library	RepeatModeler	v.2.0.1
Repeat masking	RepeatMasker	v.4.1.0
Homology-based gene prediction	GeMoMa pipeline	v.1.7.1
	MMseqs2	v.13.45111
Functional annotation	BLASTP	v.2.11.0+
	InterProScan	v.5.59.91

Results and Discussion

Sequencing of the two prepared CSS libraries on the PacBio Revio platform resulted in

6,520,091 HiFi reads (86.95 Gb) with a mean read length of 13.34 kb and a mean quality of

29.7 (Supplementary Material S1). The initial assembly with Hifiasm resulted in a draft

assembly of 2.54 Gb with 502 contigs and a contig N50 of 55.91 Mb (Table 2A). Due to the

sample quality, we were unable to generate proximity ligation libraries for scaffolding and

opted for a reference-based scaffolding approach. After removing some of the smallest

contigs, due to potential bacterial contamination (taxonomic assignments, variation in

sequencing depth and GC-content) identified by BlobTools (Fig. 2C; Supplementary Material

S2), the final highly contiguous chromosome-level assembly has a total length of 2.53 Gb

with 325 scaffolds and a scaffold and contig N50 of 68.03 MB and 56.64 Mb, respectively

(Table 2A). Of the 325 scaffolds (incl. the mitochondrial genome), the largest 40 (> 7 Mb)

corresponding to the expected 38 autosomes and the two allosomes (Tanomtong et al.,

2015) span 98.41% of the total assembly length, resulting in a L50 of 15 scaffolds. Of the

13,727 BUSCO genes in the carnivora_odb12 dataset, BUSCO identified 13,392 (97.6%) as

complete single copy and 158 (1.2%) as duplicated BUSCOs, while compleasm found 13,528

(98.55%) single copy and 126 (0.16%) duplicated BUSCOs (Fig. 2A). The high BUSCO and

compleasm scores, as well as a high mapping rate of 99.88% and a mapping quality of 39.7,

highlight the quality of the presented assembly. Furthermore, Merqury estimated a k-mer

completeness score of 96.93% with a QV score of 63.57, corresponding to an error rate of

4.39e-07 (see also Fig. 2 B).

The mitochondrial genome assembled from the HiFi reads was circular with a length of

16,729 bp, following the standard composition and structure of vertebrate mitochondrial

genome sequences with 13 protein-coding genes, 2 rRNAs, and 22 tRNAs (Fig. 2D).

Annotation

Repeat annotation

Repeat annotation identified a repeat content in the final assembly *VMU_Caureus_v1.0* of

40.58% or 1.02 Gb (Table 2B). With 24.19% of the assembly Long Interspersed Nuclear

Elements (LINEs) were the most common repeats, followed by LTR Elements (5.48%) and

Short Interspersed Nuclear Elements (SINEs, 4.95%). Non-Retroelements such as DNA

Transposons, small RNA, and simple repeats, each accounted for less than 4% of the

assembly.

Gene annotation

Homology-based gene prediction with GeMoMA identified 26,084 genes in

VMU_Caureus_v.1.0. The median gene length was 6,370 bp, spanning 469.26 Mb of the

assembly. BUSCO identified 41.2% (n = 5,653) of the 13,727 Carnivora orthologous genes

from the carnivora_odb12 BUSCO dataset in the assembly as single copy and 46.4% (n =

6,372) as duplicated, with 1,089 BUSCO genes (7.9%) missing (Fig. 2A). Compleasm identified

37.42% (n = 5,137) BUSCOs as single-copy and 50.11% (n = 6,879) as duplicated, with 7.71%

(n = 1,058) missing (Fig. 2A). InterProScan functionally annotated 61,027 of the 61,813

predicted proteins (98.73%) and assigned 53,147 proteins (85.98%) to the Reactome and

assigned at least one Gene Ontology (GO) term to 45,071 proteins (72.92%). Furthermore,

96.73 % (n=59,792) of the predicted proteins were matched to entries in the Swiss-Prot

database.

Conclusions

High-quality long-read-based genome assemblies are an essential resource for a large variety

of evolutionary, ecological, and phylogeographic analyses, as well as for biodiversity

monitoring. The presented long-read-based chromosome-level genome assembly of the

golden jackal is the first genome assembly for this species and thus provides a solid basis for

in-depth genomic studies on the species' range expansion, population dynamics, adaptability

to new environments, and potential as zoonotic disease reservoirs.

Table 2. Assembly statistics of *VMU_Caureus_v.1.0* (A) and its repeat content (B)

A

	Scaffold-level	Contig-level	
	VMU_Caureus_v.1.0	*VMU_Caureus_v.1.0**	*VMU_Caureus_draft_asm***
No. of scaffolds/contigs	325	384	502
No. of scaffolds/contigs (> 10 kbp)	325	384	502
L50	15	18	18
N50 (bp)	68,033,816	56,644,425	55,907,118
Max. scaffold/contig length(bp)	124,444,185	123,390,866	123,390,866
Total length (bp)	2,530,537,450	2,530,531,550	2,537,222,849
GC (%)	41.58	41.58	41.61
No. of N's	5,900	0	0
No. of N's per 100 kbp	0.23	0.00	0.00

*broken into contigs at gaps with a length of ≥ 10 N's. Statistics for these columns are based on contigs, the remaining columns are based on scaffolds.
**hifiasm draft assembly before reference-based scaffolding and gapclosing

B

Type of element	Number of elements	Percentage of assembly
SINEs	915,079	4.95%
LINEs:	1,980,627	24.19%
LTR elements	434,890	5.48%
DNA transposons	340,764	2.68%
Rolling-circles	1,562	0.02%
Unclassified	24,159	0.40%
Satellites	2	0.00%
Small RNA	632,699	3.31%

Simple repeats	835,984	2.25%
Low complexity	102,275	0.23%
Total		40.22%

Figure 2. Assembly quality assessment and mitochondrial genome of

_VMU_Caureus_v1.0._

(A) BUSCO and compleasm assessment results for the final assembly and the

annotation using the carnivora_obd12 dataset. (B) Merqury k-mer copy number

spectrum plot.(C) BlobPlot analysis comparing GC content (x-axis) and sequence

coverage of the PacBio data (y-axis) after removal of potentially contaminated scaffolds,

see Supplementary Material S2 for BlobPlot before removal. The color of the "blobs"

represents taxonomic assignment of the scaffolds based on the Swiss-Prot database.

(D) Annotation of the circular mitochondrial genome of CA01.

Data Accessibility

All underlying read data and the assembly are available under NCBI GenBank BioProject

PRJNA1259920. In addition, the chromosome-level and draft assembly, as well as the

annotation results, are available on Zenodo

(https://doi.org/10.5281/zenodo.16779415).

Competing Interests

The authors declare no competing interests.

Acknowledgements

We are grateful for funding from the Austrian Partnership Program for Higher Education

and Research (APPEAR) project 278 by Austria's Agency for Education and

Internationalisation (OEAD). S.W. acknowledges funding from the Austrian Science

Fund (FWF) project I508-B (to P.B.).We thank Anna Kübber-Heiss and Helmut Dier from

the Pathology Department of the University of Veterinary Medicine Vienna for access to

the frozen tissue collection and the Bioscientia Institut für Medizinische Diagnostik

GmbH (Ingelheim, Germany) for providing the PacBio SMRT sequencing service on the

PacBio Revio platform.

References

Allen, B. (2017). Challenges and opportunities for the study and conservation of large
carnivores. *Food Webs*, *12*, 1–2. https://doi.org/10.1016/j.fooweb.2017.09.001

Alonge, M., Lebeigle, L., Kirsche, M., Jenike, K., Ou, S., Aganezov, S., Wang, X., Lippman,
Z. B., Schatz, M. C., & Soyk, S. (2022). Automated assembly scaffolding using RagTag
elevates a new tomato system for high-throughput genome editing. *Genome Biology*,
23(1), 258. https://doi.org/10.1186/s13059-022-02823-7

Bao, W., Kojima, K. K., & Kohany, O. (2015). Repbase Update, a database of repetitive
elements in eukaryotic genomes. *Mobile DNA*, 6(1), 11. https://doi.org/10.1186/s13100-
015-0041-9

Blankenberg, D., Von Kuster, G., Bouvier, E., Baker, D., Afgan, E., Stoler, N., Taylor, J.,
Nekrutenko, A., & Galaxy Team. (2014). Dissemination of scientific software with Galaxy
ToolShed. *Genome Biology*, *15*(2), 403. https://doi.org/10.1186/gb4161

Buchfink, B., Reuter, K., & Drost, H.-G. (2021). Sensitive protein alignments at tree-of-
life scale using DIAMOND. *Nature Methods*, *18*(4), 366–368.
https://doi.org/10.1038/s41592-021-01101-x

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., &
Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1),
421. https://doi.org/10.1186/1471-2105-10-421

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de
novo assembly using phased assembly graphs with hifiasm. *Nature Methods*, *18*(2),
170–175. https://doi.org/10.1038/s41592-020-01056-5

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham,
A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools
and BCFtools. *GigaScience*, *10*(2), giab008.
https://doi.org/10.1093/gigascience/giab008

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.
(2020). RepeatModeler2 for automated genomic discovery of transposable element
families. *Proceedings of the National Academy of Sciences*, *117*(17), 9451–9457.
https://doi.org/10.1073/pnas.1921046117

Galov, A., Fabbri, E., Caniglia, R., Arbanasić, H., Lapalombella, S., Florijančić, T.,
Bošković, I., Galaverni, M., & Randi, E. (2015). First evidence of hybridization between
golden jackal (Canis aureus) and domestic dog (Canis familiaris) as revealed by genetic
markers. *Royal Society Open Science*, *2*(12), 150450.
https://doi.org/10.1098/rsos.150450

Gherman, C. M., & Mihalca, A. D. (2017). A synoptic overview of golden jackal parasites
reveals high diversity of species. *Parasites & Vectors*, *10*(1), 419.
https://doi.org/10.1186/s13071-017-2329-8

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: Quality assessment
tool for genome assemblies. *Bioinformatics*, *29*(8), 1072–1075.
https://doi.org/10.1093/bioinformatics/btt086

Hatlauf, J. (2024). Urban encounter: First confirmed record of a golden jackal (Canis
aureus) in Vienna, Austria. *Die Bodenkultur: Journal of Land Management, Food and
Environment, 75*(1), 1–7. https://doi.org/10.2478/boku-2024-0001

Hatlauf, J., Bayer, K., Trouwborst, A., & Hackländer, K. (2021). New rules or old
concepts? The golden jackal (Canis aureus) and its legal status in Central Europe.
European Journal of Wildlife Research, *67*(2), 25. https://doi.org/10.1007/s10344-020-
01454-2

Huang, N., & Li, H. (2023). compleasm: A faster and more accurate reimplementation of
BUSCO. *Bioinformatics*, *39*(10), btad595.
https://doi.org/10.1093/bioinformatics/btad595

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,
Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A.,
Scheremetjew, M., Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: Genome-
scale protein function classification. *Bioinformatics*, *30*(9), 1236–1240.
https://doi.org/10.1093/bioinformatics/btu031

Keilwagen, J., Hartung, F., Paulini, M., Twardziok, S. O., & Grau, J. (2018). Combining
RNA-seq data and homology-based gene prediction for plants, animals and fungi. *BMC
Bioinformatics*, *19*(1). https://doi.org/10.1186/s12859-018-2203-5

Keilwagen, J., Wenk, M., Erickson, J. L., Schattat, M. H., Grau, J., & Hartung, F. (2016).
Using intron position conservation for homology-based gene prediction. *Nucleic Acids
Research*, *44*(9), e89–e89. https://doi.org/10.1093/nar/gkw092

Klitgaard, K., Chriél, M., Isbrand, A., Jensen, T. K., & Bødker, R. (2017). Identification of
Dermacentor reticulatus Ticks Carrying Rickettsia raoultii on Migrating Jackal, Denmark.
Emerging Infectious Diseases, *23*(12), 2072–2074.
https://doi.org/10.3201/eid2312.170919

Krofel, M., Giannatos, G., Cirovic, D., Stoyanov, S., & Newsome, T. (2017). *Golden jackal
expansion in Europe: A case of mesopredator release triggered by continent-wide wolf
persecution?* https://doi.org/10.4404/hystrix-28.1-11819'%255D

Krofel, M., Hatlauf, J., Bogdanowicz, W., Campbell, L. a. D., Godinho, R., Jhala, Y. V.,
Kitchener, A. C., Koepfli, K.-P., Moehlman, P., Senn, H., Sillero-Zubiri, C., Viranta, S.,
Werhahn, G., & Alvares, F. (2022). Towards resolving taxonomic uncertainties in wolf,
dog and jackal lineages of Africa, Eurasia and Australasia. *Journal of Zoology*, *316*(3),
155–168. https://doi.org/10.1111/jzo.12946

Laetsch, D. R., & Blaxter, M. L. (2017). BlobTools: Interrogation of genome assemblies.
F1000Research, 6, 1287. https://doi.org/10.12688/f1000research.12232.1

Lanszki, J., Hayward, M. W., Ranc, N., & Zalewski, A. (2022). Dietary flexibility promotes
range expansion: The case of golden jackals in Eurasia. *Journal of Biogeography*, 49(6),
993–1005. https://doi.org/10.1111/jbi.14372

Lanszki, J., Kurys, A., Szabó, L., Nagyapáti, N., Porter, L. B., & Heltai, M. (2016). Diet
composition of the golden jackal and the sympatric red fox in an agricultural area
(Hungary). *Folia Zoologica*, 65(4), 310–322.
https://doi.org/10.25225/fozo.v65.i4.a3.2016

Lapid, R., Motro, Y., Craddock, H., Khalfin, B., King, R., Bar-Gal, G. K., & Moran-Gilad, J.
(2023). Fecal microbiota of the synanthropic golden jackal (Canis aureus). *Animal*
Microbiome, 5(1), 37. https://doi.org/10.1186/s42523-023-00259-3

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*,
34(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO
Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology*
and Evolution, 38(10), 4647–4654. https://doi.org/10.1093/molbev/msab199

Okonechnikov, K., Conesa, A., & García-Alcalde, F. (2016). Qualimap 2: Advanced
multi-sample quality control for high-throughput sequencing data. *Bioinformatics*,
32(2), 292–294. https://doi.org/10.1093/bioinformatics/btv566

Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R., & Lopez, R.
(2005). InterProScan: Protein domains identifier. *Nucleic Acids Research*, 33(suppl_2),
W116–W120. https://doi.org/10.1093/nar/gki442

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva,
M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G.
L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., … Jarvis, E. D.
(2021). Towards complete and error-free genome assemblies of all vertebrate species.
Nature, 592(7856), 737–746. https://doi.org/10.1038/s41586-021-03451-0

Salek, M., Červinka, J., Banea, O. C., Krofel, M., Ćirović, D., Selanec, I., Penezić, A., Grill,
S., & Riegert, J. (2014). Population densities and habitat use of the golden jackal (Canis
aureus) in farmlands across the Balkan Peninsula. *European Journal of Wildlife*
Research, 60(2), 193–200. https://doi.org/10.1007/s10344-013-0765-0

Serva, D., Iannella, M., Cittadino, V., & Biondi, M. (2023). A shifting carnivore's
community: Habitat modeling suggests increased overlap between the golden jackal
and the Eurasian lynx in Europe. *Frontiers in Ecology and Evolution*, 11.
https://doi.org/10.3389/fevo.2023.1165968

Smit, A., Hubley, R., & Green, P. (2015). RepeatModeler Open-1.0. 2008–2015. *Seattle,*
USA: Institute for Systems Biology.

Spassov, N., & Acosta-Pankov, I. (2019). Dispersal history of the golden jackal (Canis
aureus moreoticus Geoffroy, 1835) in Europe and possible causes of its recent
population explosion. *Biodiversity Data Journal, 7*, e34825.
https://doi.org/10.3897/BDJ.7.e34825

Stefanovic, M., Bogdanowicz, W., Adavoudi, R., Martínez-Sosa, F., Doan, K., Flores-
Manzanero, A., Srinivas, Y., Banea, O. C., Ćirović, D., D'Amico, G., Djan, M., Giannatos,
G., Hatlauf, J., Hayrapetyan, V., Heltai, M., Homel, K., Hulva, P., Ionică, A. M., Jhala, Y.
V., … Pilot, M. (2024). Range-wide phylogeography of the golden jackals (*Canis aureus*)
reveals multiple sources of recent spatial expansion and admixture with dogs at the
expansion front. *Biological Conservation, 290*, 110448.
https://doi.org/10.1016/j.biocon.2024.110448

Steinegger, M., & Soeding, J. (2017). MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets. *Nature Biotechnology, 35*(11), 1026–
1028. https://doi.org/10.1038/nbt.3988

Storer, J., Hubley, R., Rosen, J., Wheeler, T. J., & Smit, A. F. (2021). The Dfam community
resource of transposable element families, sequence models, and genome
annotations. *Mobile DNA, 12*(1), 2. https://doi.org/10.1186/s13100-020-00230-y

Tanomtong, A., Chaiyaphan, P., Supanuam, P., Puramongkol, T., Maneechot, N., &
Jangsuwan, N. (2015). The Standard Karyotype of the Asiatic Jackal, Canis aureus
(Carnivora, Canidae) from Thailand. *Cytologia, 80*(1), 3–13.
https://doi.org/10.1508/cytologia.80.3

The UniProt Consortium. (2019). UniProt: A worldwide hub of protein knowledge.
Nucleic Acids Research, 47(D1), D506–D515. https://doi.org/10.1093/nar/gky1049

Torretta, E., Dondina, O., Delfoco, C., Riboldi, L., Orioli, V., Lapini, L., & Meriggi, A.
(2020). First assessment of habitat suitability and connectivity for the golden jackal in
north-eastern Italy. *Mammalian Biology, 100*(6), 631–643.
https://doi.org/10.1007/s42991-020-00069-z

Uliano-Silva, M., Ferreira, J. G. R. N., Krasheninnikova, K., Blaxter, M., Mieszkowska, N.,
Hall, N., Holland, P., Durbin, R., Richards, T., Kersey, P., Hollingsworth, P., Wilson, W.,
Twyford, A., Gaya, E., Lawniczak, M., Lewis, O., Broad, G., Martin, F., Hart, M., … Darwin
Tree of Life Consortium. (2023). MitoHiFi: A python pipeline for mitochondrial genome
assembly from PacBio high fidelity reads. *BMC Bioinformatics, 24*(1), 288.
https://doi.org/10.1186/s12859-023-05385-y

Wang, C., Wallerman, O., Arendt, M.-L., Sundström, E., Karlsson, Å., Nordin, J.,
Mäkeläinen, S., Pielberg, G. R., Hanson, J., Ohlsson, Å., Saellström, S., Rönnberg, H.,

Ljungvall, I., Häggström, J., Bergström, T. F., Hedhammar, Å., Meadows, J. R. S., &

Lindblad-Toh, K. (2021). A novel canine reference genome resolves genomic

architecture and uncovers transcript complexity. *Communications Biology*, *4*(1), 185.

https://doi.org/10.1038/s42003-021-01698-x

Xu, M., Guo, L., Gu, S., Wang, O., Zhang, R., Peters, B. A., Fan, G., Liu, X., Xu, X., Deng,

L., & Zhang, Y. (2020). TGS-GapCloser: A fast and accurate gap closer for large genomes

with low coverage of error-prone long reads. *GigaScience*, *9*(giaa094).

https://doi.org/10.1093/gigascience/giaa094