A *de novo* reference genome of the golden jackal, *Canis aureus*

Sven Winter[1,2,3,*], René Meißner[1], Stefan Prost[4,5], Carola Greve[6], Charlotte Gerheim[6],

Marine Arakelyan[7], Sargis Aghayan[8], Jennifer Hatlauf[9], Pamela A. Burger[1,*]

[1]Research Institute of Wildlife Ecology, University of Veterinary Medicine Vienna, Vienna, Austria

[2]Faculty of Science and Technology, University of the Faroe Islands, Tórshavn, The Faroe Islands

[3]Senckenberg Biodiversity and Climate Research Centre, Frankfurt am Main, Germany

[4]Ecology and Genetics Research Unit, University of Oulu, Oulu, Finland

[5]Natural History Museum Vienna, Central Research Laboratories, Vienna, Austria

[6]Senckenberg Research Institute and Museum, Frankfurt am Main, Germany

[7]Institute of Zoology, Yerevan State University, Yerevan, Armenia

[8]Scientific Center of Zoology and Hydroecology, National Academy of Sciences of the Armenian

State

[9]Institute of Wildlife Biology and Game Management, BOKU University, Vienna, Austria

Contact for correspondence*:

Sven Winter, sven.winter@vetmeduni.ac.at

Pamela Burger, pamela.burger@vetmeduni.ac.at

Abstract

The golden jackal (*Canis aureus*) is rapidly expanding its range in Europe, driven by

climate and habitat changes, human influence, and changes in competition with wolves.

Its ecological flexibility enables it to thrive in various habitats, including urban areas,

raising concerns about its potential role in spreading zoonotic diseases. Jackals may act

as reservoirs for pathogens such as Lyme disease and babesiosis, affecting wildlife,

humans, and pets. Their close genetic relationship with domestic dogs also increases the

risk of hybridization and host-jumping, complicating disease dynamics.

To better understand their dispersal ability and host-pathogen dynamics, we present the

first chromosome-level genome assembly of the golden jackal, generated using PacBio

HiFi sequencing and reference-based scaffolding. The final assembly has a total length

of 2.53 Gb in 325 scaffolds, with 98.41% of the sequence anchored to the expected 38+XY

chromosomes. The assembly shows high contiguity, with scaffold and contig N50 values

of 68.03 Mb and 56.64 Mb, respectively. Annotation revealed 26,084 protein-coding
genes, and repetitive elements account for 40.58% of the total assembly.

This high-quality reference genome provides an essential resource for studying the
genetic basis of the golden jackal's adaptation, ecological interactions, and potential as
a zoonotic reservoir. It also supports efforts to monitor population expansion and its
effects on ecosystems. By advancing our understanding of golden jackal genetics, this
work enables future research on evolution, host-pathogen dynamics, and the broader
consequences of wildlife dispersal in a rapidly changing environment.

Keywords

Canidae, Zoonoses, Range expansion, Climate change, Pathogen transmission, Wildlife–
human interaction

Introduction

Canids (*Canidae*) are a diverse group of mostly carnivorous mammals that exhibit a wide
range of ecological adaptations and social behaviors (Allen, 2017). The approximately 35
species are distributed across all continents except Antarctica, highlighting their ability
to thrive in vastly different environments (Krofel et al., 2022). One such species, the
golden jackal (*Canis aureus* LINNAEUS, 1758), is a mid-sized canid that has been
experiencing an accelerated natural dispersal, particularly within Western Asia and into
Europe (Hatlauf et al., 2021). This expansion is thought to be driven by a combination of
several factors, including climate change, shifts in habitat, land-use changes, and other
human activities, all of which have facilitated its range expansion (Krofel et al., 2017;
Spassov & Acosta-Pankov, 2019).

The golden jackal's notable adaptability allows it to occupy a wide array of
habitats, ranging from forests and grasslands to urban environments (Salek et al., 2014;
Torretta et al., 2020). Its opportunistic feeding habits, which include a broad diet of small
mammals, birds, insects, fruits, and vegetables, make it highly resilient and capable of
colonizing new areas (Lanszki et al., 2022). While the presence of golden jackals in new

ecosystems can provide benefits, such as natural pest control by regulating populations

of rodents and other small vertebrates (Lanszki et al., 2016), the species' expansion also

presents potential challenges (Gherman & Mihalca, 2017). As jackals move into new

areas, they may compete with native species for resources and potentially disrupt local

ecosystems (Krofel et al., 2017; Serva et al., 2023). Additionally, the golden jackal's

capacity to survive in urban environments brings it into closer proximity with humans,

which increases the likelihood of interaction with domestic animals, raising concerns

about zoonotic disease transmission (Hatlauf, 2024; Lapid et al., 2023).

Figure 1. Golden jackal (*Canis aureus*) ©Jennifer Hatlauf

Recent studies have focused on the potential public health risks associated with

the movement of golden jackals, especially in light of the discovery of jackals in Denmark

carrying tick vectors and zoonotic pathogens (Klitgaard et al., 2017). Pathogens, such as

those transmitted by ticks, can pose significant risks to both wildlife and human health,

which underscores the importance of monitoring the spread of this species. The potential

for golden jackals to carry and transmit diseases like Lyme disease and babesiosis

demonstrates how their expanding range could influence the epidemiology of zoonoses,

particularly in regions where the jackal may interact with domestic animals and humans

(Lymbery et al., 2014). Furthermore, the close phylogenetic relationship between golden

jackals and domestic dogs (*Canis lupus familiaris*) raises the possibility of hybridization

and host jumping, which could exacerbate the spread of zoonotic diseases or introduce

novel pathogens into local ecosystems (Galov et al., 2015; Gherman & Mihalca, 2017;

Stefanovic et al., 2024).

Given the important ecological role golden jackals play in their current and future

habitats, they serve as important models for understanding climate-mediated dispersal

of zoonotic diseases. Their expansion is a good example of how environmental changes

and human-induced factors influence the movement of wildlife and the pathogens they

carry. Studying jackals offers valuable insights into the broader dynamics of host-

pathogen coevolution and the transmission of diseases across species boundaries. To

support these efforts, a comprehensive genomic resource is essential.

In this study, we present the first chromosome-level genome assembly of the

golden jackal, generated using PacBio HiFi long-read sequencing, supplemented by

reference-based scaffolding. This high-quality reference genome provides an invaluable

tool for investigating the genetic basis of the species' adaptation to new environments

and its interactions with both its ecosystem and pathogens. It also enables more effective

monitoring of the species and the diseases it may carry, facilitating a better

understanding of how climate change and landscape fragmentation contribute to the

spread of zoonotic diseases. By advancing our genetic knowledge of golden jackals, we

provide a foundation for future studies on the evolution, population dynamics, and

ecological impacts of this increasingly prominent species.

Materials and Methods

Sampling and Sequencing

Tissue samples from a subadult male golden jackal (sample ID: CA_001; PathoNo.:

SK/123/23) shot by a hunter in Villach, Carinthia, Austria, were collected during post-

mortem examination at the University of Veterinary Medicine Vienna, Austria, on

06.02.2023 and stored frozen at -80°C. High molecular weight DNA was extracted from

spleen tissue using the innuPrep SE HMW kit (IST Innuscreen GmbH, Germany) according

to the manufacturer's recommendations. The quantity and quality of the DNA were

evaluated using the QuantiFluor ONE dsDNA System on the Quantus Fluorometer

(Promega Corporation, USA) and the Genomic DNA Screen Tape on the Agilent 4150

TapeStation system (Agilent Technologies, Inc., USA), respectively.

Library preparation for Pacific Biosciences of California, Inc (PacBio) circular consensus

sequencing (CCS) was prepared using the SMRTbell Express Prep Kit 3.0 (PacBio, USA)

and sequenced on the PacBio Revio system.

Assembly and QC

The PacBio-generated raw data BAM file was converted into FASTQ format using

BAM2fastx v.1.3.0, a PacBio Secondary Analysis Tool

(https://github.com/PacificBiosciences/pbbioconda). The initial assemblies, including

two, which resemble the two potential parental haplotypes, were generated with hifiasm

v.0.24.0 (RRID:SCR_021069) (Cheng et al., 2021) using standard parameters. As sample

quality was insufficient for a proximity-ligation library, we did not aim for a haplotype-

resolved assembly but instead continued with the pseudohaploid assembly for the next

steps. As we were unable to use Hi-C data for the scaffolding, due to the fact that the

sample was stored in ethanol, we instead used the Mexican gray wolf assembly

(Accession No. GCF_048164855.1) for a reference-based scaffolding with RagTag v.2.1.0

(Alonge et al., 2022). Subsequently, gaps in the scaffolded assembly were breached using

the original PacBio long-read data with TGS_gapcloser v.1.2.1 (RRID:SCR_017633)(Xu et

al., 2020). The mitochondrial genome was assembled using the MitoHiFi workflow

(RRID:SCR_026369) in Galaxy v.3.2.3 (RRID:SCR_006281)(Blankenberg et al., 2014;

Uliano-Silva et al., 2023).

Assembly Quality Assessment

Assembly statistics to evaluate contiguity of the assembly were generated with Quast

v.5.0.2 (RRID:SCR_001228)(Gurevich et al., 2013). The completeness of the assembly

was assessed by gene set completeness analyses using BUSCO v.5.8.3

(RRID:SCR_015008) (Manni et al., 2021) and compleasm v.0.2.7

(RRID:SCR_026370)(Huang & Li, 2023) against the carnivora_odb12 orthologs dataset, as

well as based on k-mers (k=21) using Merqury v.1.3 with Meryl v. 1.3

(RRID:SCR_022964)(Rhie et al., 2021). Minimap2 v.2.28 (RRID:SCR_018550)(Li, 2018) and

samtools v.1.21 (RRID:SCR_002105)(Danecek et al., 2021) were used to map the PacBio

reads to the final assembly and sort the mapping file to check the mapping rate with

Qualimap bamqc v.2.3 (RRID:SCR_001209)(Okonechnikov et al., 2016). The mapping file

was also used together with the results from a Diamond v.2.1.10

(RRID:SCR_016071)(Buchfink et al., 2021) search against Swiss-Prot (release 2025-01)

(The UniProt Consortium, 2019) in Blobtools v. 1.1 (RRID:SCR_017618)(Laetsch & Blaxter,

2017) to check for potential contamination.

Annotation

Repeats in the assembly were annotated with RepeatMasker v.4.1.0 (RRID:SCR_012954)

(Smit et al., 2015) using a custom *de novo* repeat library generated from the presented

assembly with RepeatModeler v.2.0.1 ([RRID:SCR_015027](#)) (Flynn et al., 2020) combined

with the Canidae specific repeats from Dfam 3.1 and RepBase (release 20181026) (Bao

et al., 2015; Storer et al., 2021). Interspersed repeats were hardmasked (Ns) while simple

repeats were softmasked (lowercase) in preparation for the gene annotation.

Genes were predicted using the homology-based approach implemented in the GeMoMa

pipeline v.1.7.1 ([RRID:SCR_017646](#)) (Keilwagen et al., 2016, 2018) using MMseqs2 v.

13.45111 ([RRID:SCR_022962](#)) (Steinegger & Söding, 2017) as an alignment tool. The

following six mammalian genomes and corresponding annotations were used as

references: *Homo sapiens* (GCF_000001405.40), *Mus musculus* (GCF_000001635.27),

Canis lupus familiaris (German Shepherd, GCF_011100685.1; Wang et al., 2021), *Vulpes*

vulpes (GCF_048418805.1), *Vulpes lagopus* (GCF_018345385.1;, *Canis lupus baileyi*

(GCF_048164855.1; Rhie et al., 2021).

Subsequently, the proteins predicted by GeMoMa were functionally annotated by a

BLASTP v.2.11.0+ ([RRID:SCR_001010](#)) (Camacho et al., 2009) search against the Swiss-

Prot database ([RRID:SCR_002380](#); release 2025-01) (The UniProt Consortium, 2019) with

an e-value cutoff of 10^{-6}. Gene ontology (GO) terms, domains, and motifs were further

annotated using InterProScan v.5.59.91 ([RRID:SCR_005829](#)) (Jones et al., 2014;

Quevillon et al., 2005). BUSCO v.5.8.3 ([RRID:SCR_015008](#)) (Manni et al., 2021) and

compleasm v.0.2.7 ([RRID:SCR_026370](#)) (Huang & Li, 2023) were used to evaluate gene

completeness in reference to the carnivora_odb12 dataset.

Table 1. Software and versions used to generate the *Canis aureus* assembly.

Pipeline Step	Software	Version

PacBio BAM to FASTQ	BAM2fastx	v.1.3.0
assembly & long-read polishing	hifiasm	v.0.24.0
Scaffolding	RagTag	v.2.1.0
Gap-closing	TGS-GapCloser	v.1.2.1
Mitochondrial genome assembly	MitoHiFi in Galaxy	v.3.2.3
Assembly statistics	Quast	v.5.0.2
Gene set completeness	BUSCO	v.5.8.3
	compelasm	v.0.2.7
Assembly completeness	Merqury	v.1.3
Mapping of long reads	minimap2	v.2.28
Sort & index BAM	samtools	v.1.21
Mapping statistics	QualiMap	v.2.3
Contamination analyses	Diamond	v.2.1.10
	Blobtools	v.1.1
Repeat library	RepeatModeler	v.2.0.1
Repeat masking	RepeatMasker	v.4.1.0
Homology-based gene prediction	GeMoMa pipeline	v.1.7.1
	MMseqs2	v.13.45111
Functional annotation	BLASTP	v.2.11.0+
	InterProScan	v.5.59.91

Results and Discussion

Sequencing of the two prepared CSS libraries on the PacBio Revio platform resulted in

6,520,091 HiFi reads (86.95 Gb) with a mean read length of 13.34 kb and a mean quality of

29.7 (Supplementary Material S1). The initial assembly with Hifiasm resulted in a draft

assembly of 2.54 Gb with 502 contigs and a contig N50 of 55.91 Mb (Table 2A). Due to the

sample quality, we were unable to generate proximity ligation libraries for scaffolding and

opted for a reference-based scaffolding approach. After removing some of the smallest

contigs, due to potential bacterial contamination (taxonomic assignments, variation in

sequencing depth and GC-content) identified by BlobTools (Fig. 2C; Supplementary Material

S2), the final highly contiguous chromosome-level assembly has a total length of 2.53 Gb with

325 scaffolds and a scaffold and contig N50 of 68.03 MB and 56.64 Mb, respectively (Table

2A). Of the 325 scaffolds (incl. the mitochondrial genome), the largest 40 (> 7 Mb)

corresponding to the expected 38 autosomes and the two allosomes (Tanomtong et al., 2015)

span 98.41% of the total assembly length, resulting in a L50 of 15 scaffolds. Of the 13,727

BUSCO genes in the carnivora_odb12 dataset, BUSCO identified 13,392 (97.6%) as complete

single copy and 158 (1.2%) as duplicated BUSCOs, while compleasm found 13,528 (98.55%)

single copy and 126 (0.16%) duplicated BUSCOs (Fig. 2A). The high BUSCO and compleasm

scores, as well as a high mapping rate of 99.88% and a mapping quality of 39.7, highlight the

quality of the presented assembly. Furthermore, Merqury estimated a k-mer completeness

score of 96.93% with a QV score of 63.57, corresponding to an error rate of 4.39e-07 (see also

Fig. 2 B).

The mitochondrial genome assembled from the HiFi reads was circular with a length of 16,729

bp, following the standard composition and structure of vertebrate mitochondrial genome

sequences with 13 protein-coding genes, 2 rRNAs, and 22 tRNAs (Fig. 2D).

Annotation

Repeat annotation

Repeat annotation identified a repeat content in the final assembly *VMU_Caureus_v1.0* of

40.58% or 1.02 Gb (Table 2B). With 24.19% of the assembly Long Interspersed Nuclear

Elements (LINEs) were the most common repeats, followed by LTR Elements (5.48%) and Short

Interspersed Nuclear Elements (SINEs, 4.95%). Non-Retroelements such as DNA Transposons,

small RNA, and simple repeats, each accounted for less than 4% of the assembly.

Gene annotation

Homology-based gene prediction with GeMoMA identified 26,084 genes in

VMU_Caureus_v.1.0. The median gene length was 6,370 bp, spanning 469.26 Mb of the

assembly. BUSCO identified 41.2% (n = 5,653) of the 13,727 Carnivora orthologous genes from

the carnivora_odb12 BUSCO dataset in the assembly as single copy and 46.4% (n = 6,372) as

duplicated, with 1,089 BUSCO genes (7.9%) missing (Fig. 2A). Compleasm identified 37.42% (n

= 5,137) BUSCOs as single-copy and 50.11% (n = 6,879) as duplicated, with 7.71% (n = 1,058)

missing (Fig. 2A). InterProScan functionally annotated 61,027 of the 61,813 predicted proteins

(98.73%) and assigned 53,147 proteins (85.98%) to the Reactome and assigned at least one

Gene Ontology (GO) term to 45,071 proteins (72.92%). Furthermore, 96.73 % (n=59,792) of

the predicted proteins were matched to entries in the Swiss-Prot database.

Conclusions

High-quality long-read-based genome assemblies are an essential resource for a large variety

of evolutionary, ecological, and phylogeographic analyses, as well as for biodiversity

monitoring. The presented long-read-based chromosome-level genome assembly of the

golden jackal is the first genome assembly for this species and thus provides a solid basis for

in-depth genomic studies on the species' range expansion, population dynamics, adaptability

to new environments, and potential as zoonotic disease reservoirs.

Table 2. Assembly statistics of *VMU_Caureus_v.1.0* (A) and its repeat content (B)

A

	Scaffold-level	Contig-level	
	VMU_Caureus_v.1.0	*VMU_Caureus_v.1.0**	*VMU_Caureus_draft_asm***
No. of scaffolds/contigs	325	384	502
No. of scaffolds/contigs (> 10 kbp)	325	384	502
L50	15	18	18
N50 (bp)	68,033,816	56,644,425	55,907,118
Max. scaffold/contig length(bp)	124,444,185	123,390,866	123,390,866
Total length (bp)	2,530,537,450	2,530,531,550	2,537,222,849
GC (%)	41.58	41.58	41.61
No. of N's	5,900	0	0
No. of N's per 100 kbp	0.23	0.00	0.00

*broken into contigs at gaps with a length of ≥ 10 N's. Statistics for these columns are based on contigs, the remaining columns are based on scaffolds.
**hifiasm draft assembly before reference-based scaffolding and gapclosing

B

Type of element	Number of elements	Percentage of assembly
SINEs	915,079	4.95%
LINEs:	1,980,627	24.19%
LTR elements	434,890	5.48%
DNA transposons	340,764	2.68%
Rolling-circles	1,562	0.02%
Unclassified	24,159	0.40%
Satellites	2	0.00%
Small RNA	632,699	3.31%
Simple repeats	835,984	2.25%
Low complexity	102,275	0.23%
Total		40.22%

Figure 2. Assembly quality assessment and mitochondrial genome of VMU_Caureus_v1.0.

(A) BUSCO and compleasm assessment results for the final assembly and the annotation using the carnivora_obd12 dataset. (B) Merqury k-mer copy number spectrum plot.(C) BlobPlot analysis comparing GC content (x-axis) and sequence coverage of the PacBio data (y-axis) after removal of potentially contaminated scaffolds, see Supplementary Material S2 for BlobPlot before removal. The color of the "blobs" represents taxonomic

assignment of the scaffolds based on the Swiss-Prot database. (D) Annotation of the

circular mitochondrial genome of CA01.

Data Accessibility

Competing Interests

The authors declare no competing interests.

Acknowledgements

We are grateful for funding from the Austrian Partnership Program for Higher Education

and Research (APPEAR) project 278 by Austria's Agency for Education and

Internationalisation (OEAD). S.W. acknowledges funding from the Austrian Science Fund

(FWF) project I508-B (to P.B.).We thank Anna Kübber-Heiss and Helmut Dier from the

Pathology Department of the University of Veterinary Medicine Vienna for access to the

frozen tissue collection and the Bioscientia Institut für Medizinische Diagnostik GmbH

(Ingelheim, Germany) for providing the PacBio SMRT sequencing service on the PacBio

Revio platform.

References

Allen, B. (2017). Challenges and opportunities for the study and conservation of large
carnivores. *Food Webs*, *12*, 1–2. https://doi.org/10.1016/j.fooweb.2017.09.001

Alonge, M., Lebeigle, L., Kirsche, M., Jenike, K., Ou, S., Aganezov, S., Wang, X., Lippman,
Z. B., Schatz, M. C., & Soyk, S. (2022). Automated assembly scaffolding using RagTag

elevates a new tomato system for high-throughput genome editing. *Genome Biology*,

23(1), 258. https://doi.org/10.1186/s13059-022-02823-7

Bao, W., Kojima, K. K., & Kohany, O. (2015). Repbase Update, a database of repetitive

elements in eukaryotic genomes. *Mobile DNA*, *6*(1), 11. https://doi.org/10.1186/s13100-

015-0041-9

Blankenberg, D., Von Kuster, G., Bouvier, E., Baker, D., Afgan, E., Stoler, N., Taylor, J.,

Nekrutenko, A., & Galaxy Team. (2014). Dissemination of scientific software with Galaxy

ToolShed. *Genome Biology*, *15*(2), 403. https://doi.org/10.1186/gb4161

Buchfink, B., Reuter, K., & Drost, H.-G. (2021). Sensitive protein alignments at tree-of-

life scale using DIAMOND. *Nature Methods*, *18*(4), 366–368.

https://doi.org/10.1038/s41592-021-01101-x

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., &

Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1),

421. https://doi.org/10.1186/1471-2105-10-421

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de

novo assembly using phased assembly graphs with hifiasm. *Nature Methods*, *18*(2),

170–175. https://doi.org/10.1038/s41592-020-01056-5

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham,

A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools

and BCFtools. *GigaScience*, *10*(2), giab008.

https://doi.org/10.1093/gigascience/giab008

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.

(2020). RepeatModeler2 for automated genomic discovery of transposable element

families. *Proceedings of the National Academy of Sciences*, *117*(17), 9451–9457.

https://doi.org/10.1073/pnas.1921046117

Galov, A., Fabbri, E., Caniglia, R., Arbanasić, H., Lapalombella, S., Florijančić, T.,

Bošković, I., Galaverni, M., & Randi, E. (2015). First evidence of hybridization between

golden jackal (Canis aureus) and domestic dog (Canis familiaris) as revealed by genetic

markers. *Royal Society Open Science*, *2*(12), 150450.

https://doi.org/10.1098/rsos.150450

Gherman, C. M., & Mihalca, A. D. (2017). A synoptic overview of golden jackal parasites

reveals high diversity of species. *Parasites & Vectors*, *10*(1), 419.

https://doi.org/10.1186/s13071-017-2329-8

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: Quality assessment

tool for genome assemblies. *Bioinformatics*, *29*(8), 1072–1075.

https://doi.org/10.1093/bioinformatics/btt086

Hatlauf, J. (2024). Urban encounter: First confirmed record of a golden jackal (Canis
aureus) in Vienna, Austria. *Die Bodenkultur: Journal of Land Management, Food and*
Environment, 75(1), 1–7. https://doi.org/10.2478/boku-2024-0001

Hatlauf, J., Bayer, K., Trouwborst, A., & Hackländer, K. (2021). New rules or old
concepts? The golden jackal (Canis aureus) and its legal status in Central Europe.
European Journal of Wildlife Research, 67(2), 25. https://doi.org/10.1007/s10344-020-
01454-2

Huang, N., & Li, H. (2023). compleasm: A faster and more accurate reimplementation of
BUSCO. *Bioinformatics, 39*(10), btad595.
https://doi.org/10.1093/bioinformatics/btad595

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,
Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A.,
Scheremetjew, M., Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: Genome-
scale protein function classification. *Bioinformatics, 30*(9), 1236–1240.
https://doi.org/10.1093/bioinformatics/btu031

Keilwagen, J., Hartung, F., Paulini, M., Twardziok, S. O., & Grau, J. (2018). Combining
RNA-seq data and homology-based gene prediction for plants, animals and fungi. *BMC*
Bioinformatics, 19(1). https://doi.org/10.1186/s12859-018-2203-5

Keilwagen, J., Wenk, M., Erickson, J. L., Schattat, M. H., Grau, J., & Hartung, F. (2016).
Using intron position conservation for homology-based gene prediction. *Nucleic Acids*
Research, 44(9), e89–e89. https://doi.org/10.1093/nar/gkw092

Klitgaard, K., Chriél, M., Isbrand, A., Jensen, T. K., & Bødker, R. (2017). Identification of
Dermacentor reticulatus Ticks Carrying Rickettsia raoultii on Migrating Jackal, Denmark.
Emerging Infectious Diseases, 23(12), 2072–2074.
https://doi.org/10.3201/eid2312.170919

Krofel, M., Giannatos, G., Cirovic, D., Stoyanov, S., & Newsome, T. (2017). *Golden jackal*
expansion in Europe: A case of mesopredator release triggered by continent-wide wolf
persecution? https://doi.org/10.4404/hystrix-28.1-11819'%255D

Krofel, M., Hatlauf, J., Bogdanowicz, W., Campbell, L. a. D., Godinho, R., Jhala, Y. V.,
Kitchener, A. C., Koepfli, K.-P., Moehlman, P., Senn, H., Sillero-Zubiri, C., Viranta, S.,
Werhahn, G., & Alvares, F. (2022). Towards resolving taxonomic uncertainties in wolf,
dog and jackal lineages of Africa, Eurasia and Australasia. *Journal of Zoology, 316*(3),
155–168. https://doi.org/10.1111/jzo.12946

Laetsch, D. R., & Blaxter, M. L. (2017). BlobTools: Interrogation of genome assemblies.
F1000Research, 6, 1287. https://doi.org/10.12688/f1000research.12232.1

Lanszki, J., Hayward, M. W., Ranc, N., & Zalewski, A. (2022). Dietary flexibility promotes
range expansion: The case of golden jackals in Eurasia. *Journal of Biogeography*, *49*(6),
993–1005. https://doi.org/10.1111/jbi.14372

Lanszki, J., Kurys, A., Szabó, L., Nagyapáti, N., Porter, L. B., & Heltai, M. (2016). Diet
composition of the golden jackal and the sympatric red fox in an agricultural area
(Hungary). *Folia Zoologica*, *65*(4), 310–322.
https://doi.org/10.25225/fozo.v65.i4.a3.2016

Lapid, R., Motro, Y., Craddock, H., Khalfin, B., King, R., Bar-Gal, G. K., & Moran-Gilad, J.
(2023). Fecal microbiota of the synanthropic golden jackal (Canis aureus). *Animal*
Microbiome, *5*(1), 37. https://doi.org/10.1186/s42523-023-00259-3

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*,
34(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO
Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology*
and Evolution, *38*(10), 4647–4654. https://doi.org/10.1093/molbev/msab199

Okonechnikov, K., Conesa, A., & García-Alcalde, F. (2016). Qualimap 2: Advanced
multi-sample quality control for high-throughput sequencing data. *Bioinformatics*,
32(2), 292–294. https://doi.org/10.1093/bioinformatics/btv566

Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R., & Lopez, R.
(2005). InterProScan: Protein domains identifier. *Nucleic Acids Research*, *33*(suppl_2),
W116–W120. https://doi.org/10.1093/nar/gki442

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva,
M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G.
L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., … Jarvis, E. D.
(2021). Towards complete and error-free genome assemblies of all vertebrate species.
Nature, *592*(7856), 737–746. https://doi.org/10.1038/s41586-021-03451-0

Salek, M., Červinka, J., Banea, O. C., Krofel, M., Ćirović, D., Selanec, I., Penezić, A., Grill,
S., & Riegert, J. (2014). Population densities and habitat use of the golden jackal (Canis
aureus) in farmlands across the Balkan Peninsula. *European Journal of Wildlife*
Research, *60*(2), 193–200. https://doi.org/10.1007/s10344-013-0765-0

Serva, D., Iannella, M., Cittadino, V., & Biondi, M. (2023). A shifting carnivore's
community: Habitat modeling suggests increased overlap between the golden jackal
and the Eurasian lynx in Europe. *Frontiers in Ecology and Evolution*, *11*.
https://doi.org/10.3389/fevo.2023.1165968

Smit, A., Hubley, R., & Green, P. (2015). RepeatModeler Open-1.0. 2008–2015. *Seattle,*
USA: Institute for Systems Biology.

Spassov, N., & Acosta-Pankov, I. (2019). Dispersal history of the golden jackal (Canis
aureus moreoticus Geoffroy, 1835) in Europe and possible causes of its recent
population explosion. *Biodiversity Data Journal*, *7*, e34825.
https://doi.org/10.3897/BDJ.7.e34825

Stefanovic, M., Bogdanowicz, W., Adavoudi, R., Martínez-Sosa, F., Doan, K., Flores-
Manzanero, A., Srinivas, Y., Banea, O. C., Ćirović, D., D'Amico, G., Djan, M., Giannatos,
G., Hatlauf, J., Hayrapetyan, V., Heltai, M., Homel, K., Hulva, P., Ionică, A. M., Jhala, Y.
V., ... Pilot, M. (2024). Range-wide phylogeography of the golden jackals (*Canis aureus*)
reveals multiple sources of recent spatial expansion and admixture with dogs at the
expansion front. *Biological Conservation*, *290*, 110448.
https://doi.org/10.1016/j.biocon.2024.110448

Steinegger, M., & Soeding, J. (2017). MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets. *Nature Biotechnology*, *35*(11), 1026–
1028. https://doi.org/10.1038/nbt.3988

Storer, J., Hubley, R., Rosen, J., Wheeler, T. J., & Smit, A. F. (2021). The Dfam community
resource of transposable element families, sequence models, and genome
annotations. *Mobile DNA*, *12*(1), 2. https://doi.org/10.1186/s13100-020-00230-y

Tanomtong, A., Chaiyaphan, P., Supanuam, P., Puramongkol, T., Maneechot, N., &
Jangsuwan, N. (2015). The Standard Karyotype of the Asiatic Jackal, Canis aureus
(Carnivora, Canidae) from Thailand. *Cytologia*, *80*(1), 3–13.
https://doi.org/10.1508/cytologia.80.3

The UniProt Consortium. (2019). UniProt: A worldwide hub of protein knowledge.
Nucleic Acids Research, *47*(D1), D506–D515. https://doi.org/10.1093/nar/gky1049

Torretta, E., Dondina, O., Delfoco, C., Riboldi, L., Orioli, V., Lapini, L., & Meriggi, A.
(2020). First assessment of habitat suitability and connectivity for the golden jackal in
north-eastern Italy. *Mammalian Biology*, *100*(6), 631–643.
https://doi.org/10.1007/s42991-020-00069-z

Uliano-Silva, M., Ferreira, J. G. R. N., Krasheninnikova, K., Blaxter, M., Mieszkowska, N.,
Hall, N., Holland, P., Durbin, R., Richards, T., Kersey, P., Hollingsworth, P., Wilson, W.,
Twyford, A., Gaya, E., Lawniczak, M., Lewis, O., Broad, G., Martin, F., Hart, M., ... Darwin
Tree of Life Consortium. (2023). MitoHiFi: A python pipeline for mitochondrial genome
assembly from PacBio high fidelity reads. *BMC Bioinformatics*, *24*(1), 288.
https://doi.org/10.1186/s12859-023-05385-y

Wang, C., Wallerman, O., Arendt, M.-L., Sundström, E., Karlsson, Å., Nordin, J.,
Mäkeläinen, S., Pielberg, G. R., Hanson, J., Ohlsson, Å., Saellström, S., Rönnberg, H.,
Ljungvall, I., Häggström, J., Bergström, T. F., Hedhammar, Å., Meadows, J. R. S., &
Lindblad-Toh, K. (2021). A novel canine reference genome resolves genomic

architecture and uncovers transcript complexity. *Communications Biology*, *4*(1), 185.
https://doi.org/10.1038/s42003-021-01698-x

Xu, M., Guo, L., Gu, S., Wang, O., Zhang, R., Peters, B. A., Fan, G., Liu, X., Xu, X., Deng,
L., & Zhang, Y. (2020). TGS-GapCloser: A fast and accurate gap closer for large genomes
with low coverage of error-prone long reads. *GigaScience*, *9*(giaa094).
https://doi.org/10.1093/gigascience/giaa094

414