Kingdom-wide evolutionary characterization of RNA editing factors across Archaeplastida

Ming Chen[1, 2, 3], Qingming Qu[4], Zhang Zhang[1, 2, 3, *]

[1]National Genomics Data Center, China National Center for Bioinformation, Beijing, China

[2]Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing, China

[3]University of Chinese Academy of Sciences, Beijing, China

[4]State Key Laboratory of Cellular Stress Biology, School of Life Sciences, Xiamen University, China

*Corresponding author: Zhang Zhang (zhangzhang@big.ac.cn)

Abstract

RNA editing, a post-transcriptional modification in plant mitochondria and plastids, is essential for environmental adaptation and diverse physiological processes. Despite extensive identification of RNA editing factors, primarily including pentatricopeptide repeat (PPR), multiple organelle RNA editing factor (MORF), organelle RNA recognition motif-containing (ORRM), and organelle zinc finger (OZ) proteins, their evolutionary history remains poorly understood. Here, we perform kingdom-wide evolutionary analyses across 364 high-quality Archaeplastida genomes and find massive PPR gene expansions in early-diverging land plants, predominantly driven by dispersed duplication associated with retroposition. Furthermore, integrative analyses indicate that DYW subgroup PPR genes have been horizontally transferred from plants to bdelloid rotifers, potentially enhancing the recipients' adaptation to ultraviolet radiation. MORF proteins, accessory partners of PPRs, possess MORF hallmark domains structurally similar to protein-folding peptidase S8 propeptide/proteinase inhibitor I9 domains, suggesting a role in protein folding during RNA editing. Considering diverse domain compositions, we reclassify MORF, ORRM, and OZ proteins and uncover prevalent hallmark domain fusions. Together, these findings illuminate the kingdom-wide evolution of plant RNA editing machinery.

Key words: RNA editing; PPR; MORF; ORRM; OZ; evolution

Introduction

RNA editing, a post-transcriptional modification primarily involving cytidine-to-uridine (C-to-U) and uridine-to-cytidine (U-to-C) alterations in mitochondria and plastids, serves as a pivotal layer of genetic regulation in plants[1,2]. This process is essential for adaptation to diverse environments[3-6] and various physiological processes[7], including development[8], growth[9], flowering[10], fruit ripening[11], and resistance to stresses[12,13]. Editing factors, a group of nuclear-encoded proteins, are assembled into an editing complex to mediate RNA editing[6]. Given the remarkable diversity of editing factors in plants, elucidating deciphering their evolution is essential for understanding how the RNA editing machinery emerged and diversified[3,14-16].

Decades of work have been made to identify RNA editing factors, primarily including Pentatricopeptide Repeat (PPR), Multiple Organelle RNA Editing Factor (MORF) (also known as RNA Editing Factor Interacting Protein, RIP), Organelle RNA Recognition Motif-containing (ORRM), and Organelle Zinc (OZ) finger proteins[14], each characterized by hallmark domains or motifs that contribute distinctly to regulating RNA editing[17]. Among them, PPR proteins are key editing factors, specifically recognizing RNA sequences through P/PLS motif tracts and catalyzing C-to-U and U-to-C editing via carboxyl-terminal (C-terminal) DYW and DW:KP domains, respectively[18,19]. Furthermore, PPR proteins are classified into two subfamilies, P and PLS, with the latter further divided into PLS, E, E+, and DYW subgroups[17]. PPRs' accessory proteins, including MORF, ORRM, and OZ, which are named after their MORF, RRM, Ran Binding Protein 2 (RanBP2) type Zinc finger (Znf) hallmark domains, respectively, coordinate RNA editing through interactions both with PPR proteins and among themselves[6,10,20-27]. Beyond their hallmark domains, these

proteins also incorporate diverse additional domains, such as Glycine-Rich (GR)

domain in MORF proteins[28-30] and PPR/MORF domains in ORRM proteins[31].

Building on these advances, studies have increasingly focused on the

evolutionary histories of PPR, MORF, ORRM, and OZ genes[6,14,15,32,33]. The

PPR gene family is one of the largest gene families in plants[19], ranging from

tens of members in early-diverging chlorophytes to hundreds or thousands in

land plants[4,6] and thereby reflecting massive expansions[4,34]. PPR genes also

occur beyond plants, with P subfamily members widespread across eukaryotes

and occasionally detected in prokaryotes[35,36]. Notably, DYW subgroup PPR

genes are observed in multiple non-plant eukaryote lineages[37-40], including

Amoebozoa, Discoba, Fungi, Metazoa, and Malawimonadidae, presumably

resulting from horizontal gene transfer (HGT)[41]. In contrast, MORF genes are

present exclusively in angiosperms and gymnosperms[34]; ORRM genes are

broadly distributed across Viridiplantae lineages including chlorophytes,

mosses, liverworts, and angiosperms[31]; and OZ genes are sporadically

distributed in gymnosperms, mosses, and lycophytes, but are absent from

chlorophytes[25]. Despite these insights, a comprehensive evolutionary

characterization of these RNA editing factors across Archaeplastida remains

unexplored.

Toward this end, we collect a comprehensive list of genomes across

Archaeplastida, systematically identify PPR, MORF, ORRM, and OZ genes,

and reconstruct an evolutionary landscape of plant RNA editing factors. Building

on the landscape, we perform phylogenomic analyses to elucidate the

mechanisms underlying PPR expansions and investigate the extent and

evolutionary significance of HGT in DYW subgroup PPR genes. Through

sequence and structural alignments, we explore the conservation and diversity

93 of hallmark domains of MORF proteins. Finally, we reclassify MORF, ORRM,

94 and OZ proteins based on their domain composition.

Results

Evolutionary landscape of PPR, MORF, ORRM, and OZ genes across Archaeplastida

98 To systematically investigate the evolution of PPR, MORF, ORRM, and OZ

99 genes, we obtain a total of 364 high-quality genomes across Archaeplastida

100 that span early-diverging aquatic algae to highly derived land plants (for details

101 see Materials and Methods), including glaucophytes (*n*=1), rhodophytes (*n*=7),

102 prasinodermophytes (*n*=1), chlorophytes (*n*=33), charophytes (*n*=7), hornworts

103 (*n*=22), mosses (*n*=13), liverworts (*n*=4), lycophytes (*n*=6), ferns (*n*=5),

104 gymnosperms (*n*=13), and angiosperms (*n*=262) (Supplementary Table 1).

105 Based on these genomes, we first identify 138 single-copy orthologous gene

106 families (Supplementary Table 2) and then build a plant phylogenomic tree (Fig.

107 1a and Supplementary Fig. 1) that is topologically congruent with previous

108 studies[42,43]. Next, we systematically identify 212,779 PPR, 2,759 MORF, 7,760

109 ORRM, and 829 OZ genes (Supplementary Tables 3 and 4), and evolutionarily

110 characterize these genes in the phylogenomic tree as built above. Noticeably,

111 we find that PPR genes are scarcely present in glaucophytes, rhodophytes,

112 prasinodermophytes, chlorophytes, and charophytes, ranging only from 8 to

113 182 (Fig. 1b and Supplementary Table 4). In contrast, hundreds to thousands

114 of PPR genes are detected in land plants, particularly some hornworts, mosses,

115 and lycophytes, followed by gymnosperms, ferns, and angiosperms.

116 Remarkably, two lycophytes, namely, *Selaginella moellendorffii* and *Isoetes*

117 *sinensis*, have the highest number of PPR genes (*n*=3,861 and 3,256,

118 respectively) among the 364 Archaeplastida species in our study (Fig. 1b and

119 Supplementary Table 4). On the contrary, liverworts are short of PPR genes,

less than 100 in count (Supplementary Table 4).

Compared to PPR genes, the other three types of genes present distinct
patterns. Specifically, MORF genes are found only in seed plants, including
angiosperms and gymnosperms (Fig. 1c), in agreement with a previous study[34].
Additionally, seed plants generally harbor about 10 MORF genes, whereas
some species, such as *Brassica napus*, *Pogostemon cablin*, and *Cissus*
rotundifolia, have more MORF genes (*n*=40, 39, and 37, respectively;
Supplementary Table 4). Conversely, ORRM genes are ubiquitous across
Archaeplastida and exhibit a gradual increase from glaucophytes to
angiosperms, indicating their ancient origin (Fig. 1d). Albeit much fewer in
number than the rest three types of genes, OZ genes are enriched in
angiosperms, with a minor presence in chlorophytes, hornworts, mosses,
liverworts, lycophytes, ferns, and gymnosperms (Fig. 1e). Besides, many
angiosperms that have undergone diploidization or polyploidization, such as *P.*
cablin[44], *Crocus sativus*[45], and *Triticum aestivum*[46], exhibit concurrent
expansions of PPR, MORF, ORRM, and OZ genes (Fig. 1 and Supplementary
Table 4). In contrast, among these four types of genes, MORF genes are the
only ones markedly expanded in drought-resistant *C. rotundifolia* relative to
other angiosperms, whereas OZ genes are the only ones significantly
expanded in *Crataegus pinnatifida* and *Malus fusca*. Overall, PPR genes exhibit
dramatic expansions in land plants, in contrast to MORF, ORRM, and OZ genes.

PPR gene expansions in early-diverging land plants

Gene expansion can be achieved by gene duplication with four different modes,
viz., whole genome duplication (WGD), tandem duplication (TD), proximal
duplication (PD), and dispersed duplication (DD), which are classified
according to genomic loci of duplicated gene pairs[47-49]. To explore the

underlying mechanisms of PPR gene expansions in early-diverging land plants,

we select high-quality chromosome-level genomes, including four hornworts

(*Anthoceros agrestis*, 2,308 PPRs; *Notothylas orbicularis*, 1,772 PPRs;

Paraphymatoceros pearsonii, 1,583 PPRs; and *Phymatoceros phymatodes*,

1,627 PPRs), one moss (*Takakia lepidozioides*, 2,016 PPRs), and two

lycophytes (*I. sinensis*, 3,256 PPRs; and *Huperzia asiatica*, 1,583 PPRs)

(Supplementary Table 4), with the charophyte *Zygnema circumcarinatum* (67

PPRs) serving as the phylogenetic outgroup. Furthermore, we select well-

annotated PPR genes, accounting for 93.40~100% of our identified PPRs, and

detect that 90.04~95.89% of them form duplicated gene pairs based on

sequence similarity analysis (Supplementary Table 5). Meanwhile, we infer

WGDs in these selected plants through comprehensive analyses of gene

synteny and synonymous substitution rates (Ks) of homologous gene pairs (for

details see Materials and Methods). Accordingly, we find that *Z.*

circumcarinatum and four hornworts lack evidence of WGD, whereas *T.*

lepidozioides and *I. sinensis* each experience one WGD, and *H. asiatica*

undergoes two WGDs (Supplementary Fig. 2-9). These results are consistent

with previous studies[5,50-54], except that *I. sinensis* has been reported to

experience two WGDs based solely on Ks distributions[55]. Consequently,

integration of inferred WGD events with comparison of genomic loci of

duplicated PPR gene pairs (for details see Materials and Methods) indicates

that DD is primarily responsible for PPR gene expansions in the selected

species, followed by WGD in *I. sinensis* and *H. asiatica* (Fig. 2a and

Supplementary Table 5).

Furthermore, it has been reported that DD is related to retroposition[47,49], a

process that the complementary DNA transcript of a gene is inserted back into

the genome, thus producing a retrogene[56]. Generally, the retrogene of a

duplicated pair features intron loss and a polyadenylation (polyA) tail, and the

pair is flanked by short direct repeats (e.g., transposable element, TE)[56,57]. Thus,

we examine these features of DD PPR gene pairs as indicators of retroposition.

Specifically, we compare intron counts of PPR genes in each dispersed

duplicate pair, and find that 74.78~93.21% of DD-derived PPR genes in

hornworts and lycophytes have undergone intron loss, followed by the moss *T.*

lepidozioides with 51.98% (Supplementary Table 6 and Fig. 2b; for details see

Materials and Methods). Furthermore, we investigate the presence of polyA

tails downstream of the 3' untranslated regions (3'UTRs) of intronless DD-

derived PPR genes and find that 97.99~99.26% possess polyA tails in

lycophytes, 70.37~75.61% in hornworts, and 81.55% in the moss *T.*

lepidozioides (Supplementary Fig. 10). Meanwhile, we identify TEs within 20

kilobases (kb) upstream and downstream of intronless DD PPR gene pairs,

showing that 70.52~99.93% of them are flanked by TEs in lycophytes,

26.56~68.31% in hornworts, and 80.11% in *T. lepidozioides* (Supplementary

Fig. 11). Notably, most of these TEs are long terminal repeats (LTRs), including

LTR/Copia, LTR/Gypsy, and LTR/unknown (Supplementary Table 7).

Additionally, we present two DD PPR gene pairs with intron loss that are flanked

by the same type of LTR with high sequence similarity, viz., E2 subgroup PPR

genes, *HaA01G1373* and *HaA43G0090* in *H. asiatica* flanked by LTR/Copia

(96.43% upstream and 100% downstream) (Fig. 2c), and DYW subgroup PPR

genes, *Tle3c03864.1* and *Tle3c03878.1* in *T. lepidozioides* by LTR/Gypsy

(99.86% upstream and 100% downstream) (Fig. 2d). Furthermore, co-

occurrence of these three retroposition features is detected in 69.93~97.92%

of DD PPR gene pairs in lycophytes, 17.75~51.86% in hornworts, and 64.97%

in *T. lepidozioides* (Fig. 2e), suggesting lineage-specific degeneration of

retroposition signals. Collectively, these results indicate that PPR gene

expansions in early-diverging land plants are primarily driven by DD associated

with retroposition.

Horizontal transfer of DYW subgroup PPR genes from plants to bdelloid rotifers

As PPR genes consist of the P and PLS subfamilies, we explore their
phylogenetic patterns by reconstructing an evolutionary landscape across
Archaeplastida (Supplementary Fig. 12). We find that the P subfamily PPR
genes are ubiquitously distributed in Archaeplastida, with their numbers
progressively increasing from early-diverging glaucophytes to highly derived
angiosperms. Conversely, the PLS subfamily, including PLS, E, E+, DYW, and
DYW:KP subgroups, is confined to land plants, with rare occurrence in
liverworts and some mosses. In addition, the PLS and E subgroups are
numerous in lycophytes, while the E+ subgroup is exclusively enriched in
hornworts. The DYW subgroup is broadly distributed in land plants, with a
maximum of 750 members observed in the moss *T. lepidozioides*. In contrast,
its variant, the DYW:KP subgroup, is restricted to hornworts, lycophytes, and
ferns (Supplementary Table 3 and Supplementary Fig. 12), conforming with a
previous study[34].

Inspired by the previous findings that DYW subgroup PPR genes have also
been detected in non-plants[37,41], we next investigate the evolutionary
relationships of the DYW and DYW:KP subgroups between plants and non-
plants. Given that repetitive and variable PLS motif triplets may hinder reliable
phylogenetic inference and DYW domain catalyzes C-to-U editing via its PG
box (covering the first two β-strands), the zinc ion binding signature
$HxE(x)_nCxxC$, and C-terminal DYW tripeptide[58-60], we extract 6,646 DYW and
3,761 DYW:KP domains of 68 land plants as representatives of their respective
subgroup PPR genes (Supplementary Table 8). Based on these plant domains,

we further survey non-plant DYW and DYW:KP domains across 40 lineages

spanning eukaryotes, prokaryotes, and viruses (Supplementary Table 9; for

details see Materials and Methods). In total, we detect 1,666 DYW and 6

DYW:KP domains in Metazoa, 31 DYW in Rhizaria, 22 DYW and 1 DYW:KP in

Discoba, 15 DYW in Fungi, 9 DYW in Alveolata, 3 DYW in Haptista, and 1 DYW

in Amoebozoa (Supplementary Table 10), exhibiting 21.21~80% sequence

similarity to their plant counterparts (Supplementary Table 11). To reduce

domain redundancy, we further select 1,151 DYW and 356 DYW:KP domains

from six representative species spanning major land plant lineages, including

P. pearsonii (hornwort), *T. lepidozioides* (moss), *I. sinensis* (lycophyte),

Alsophila spinulosa (fern), *Gnetum montanum* (gymnosperm), and *Amborella*

trichopoda (angiosperm), together with all non-plant DYW and DYW:KP

domains for phylogenetic reconstruction (Supplementary Table 10). Notably,

three major clusters of non-plant DYW domains are embedded within plant

lineages, suggesting multiple waves of HGT from plants to non-plants (Fig. 3).

To explore the potential role of DYW subgroup PPR genes in non-plants, we

select a representative DYW domain from each lineage and compare its

sequence and structure with those of plant donors. On the one hand, most of

these non-plant DYW domains contain the conserved HxE(x)nCxxC motif,

except for the HxE(x)nGxxN in Fungi, whereas the PG-box (approximately 15

amino acid residues (aa) beginning with proline and glycine[61]) and DYW

tripeptide are variable (Fig. 4a and 4b; Supplementary Fig. 13). On the other

hand, these non-plant DYW domains exhibit significant structural similarity to

those of their plant donors (Root Mean Square Deviation, RMSD ≤ 2.5Å[62]), such

as the DYW domain in the bdelloid rotifer *Adineta vaga* (Metazoa) and its

counterpart in the moss *T. lepidozioides* (Fig. 4b and Supplementary Fig. 13).

Surprisingly, Metazoa acquire the largest number of DYW domains from plants

(Fig. 3), all of which occur in bdelloid rotifers that are resistant to ultraviolet (UV)

radiation and inhabit diverse environments, such as mosses, lichens, soils, and

deserts[63,64]. Notably, these bdelloid rotifers comprise *A. vaga*, *A. ricciae*, *A.*

steineri, *Didymodactylos carnosus*, *Philodina roseola*, *Rotaria magnacalcarata*,

R. socialis, *R. sordida*, *R.* sp. 'Silwood1', and *R.* sp. 'Silwood2'. Among them,

we find that *A. vaga* harbors lots of PPR genes (*n*=191), nearly one third of

which (*n*=60) are members of the DYW subgroup (Supplementary Table 12).

The majority of these DYW subgroup PPR genes (58/60) are expressed (Fig.

4c), suggesting their potential roles in biological processes. Intriguingly, the

moss *T. lepidozioides*, a potential donor of DYW subgroup PPR genes to *A.*

vaga, is also resistant to UV radiation[5]. Consistent with this resistance, its DYW

genes are upregulated during recovery from UV irradiation (Fig. 4d and

Supplementary Table 13), highlighting their potential roles in response to UV

radiation. Taken together, these results suggest that bdelloid rotifers living in

mosses may have acquired the DYW subgroup PPR genes as an adaptation

to high UV radiation (Fig. 4e).

Evolutionary conservation and diversity of MORF domains

MORF proteins, exclusive to seed plants (Fig. 1c) and comprising nine

members (MORF1~MORF9) in *A. thaliana*, are characterized by their MORF

hallmark domains and are essential for PPR-mediated RNA editing[28-30]. To

explore the evolutionary conservation of MORF domains, we extract 334

instances from 33 representative seed plants (Supplementary Table 14) and

identify their homologs across 45 lineages spanning eukaryotes, prokaryotes,

and viruses (Supplementary Table 9). We find that MORF domains are

homologous to peptidase S8 propeptide/proteinase inhibitor I9 domains (also

known as PA domain[28]) in metazoa, fungi, bryophytes, lycophytes, and ferns

(Fig. 5a and Supplementary Table 15). Furthermore, we compare the structures

of the MORF domain and its homologous PA domain with the highest sequence

similarity in each of these five lineages. We find that although they have

relatively low sequence similarity (33.33~41.94%), their structures are

significantly similar (RMSD ≤ 2.5Å), with RMSD values varying from

0.705~1.039Å (Fig. 5b and 5c; Supplementary Fig. 14). Notably, MORF

domains from MORF3 and MORF8 proteins share the same α-helix and β-

strand architecture as PA domains (Fig. 5c and Supplementary Fig. 14). As PA

domain is important in assisting fold zymogens and mature peptidases[65,66],

these results imply that MORF domains from MORF3 and MORF8 proteins may

help fold PPR proteins to regulate RNA editing.

To investigate the evolutionary diversity of MORF domains, we classify them

into MORF1 to MORF9 as identified in *A. thaliana* and survey PA domains

across our collected 364 high-quality Archaeplastida genomes. Accordingly, PA

domains are ubiquitous across major lineages, including glaucophytes,

charophytes, mosses, liverworts, hornworts, lycophytes, ferns, gymnosperms,

and angiosperms (Supplementary Fig. 15). Furthermore, we reconstruct a

phylogenetic tree of the 334 MORF domains, using five non-seed plant PA

domains with the highest sequence similarity as the outgroup. Consequently,

MORF2, MORF5, and MORF6 domains cluster into early-diverging clades in

the phylogenetic tree, indicating their earlier origins relative to other MORF

domains in seed plants (Fig. 5d). Moreover, most MORF domains (MORF3~6,

8, and 9) are broadly distributed across angiosperms and gymnosperms,

suggesting that MORF diversification predates the major divergences among

seed plant lineages. At the gene level, we analyze the expression of

MORF1~MORF9 genes, each harboring a single MORF hallmark domain,

across multiple tissues of the gymnosperm *Torreya grandis* and the angiosperm

A. thaliana. Notably, among these MORF genes, MORF8 and MORF9 display

higher expression across tissues in both species, consistent with observations

in *Apium graveolens* and *Populus trichocarpa*[67,68], indicating their conservation

in expression (Supplementary Fig. 16).

Hallmark domain fusions among MORF, ORRM, OZ, and PPR proteins

While MORF proteins contain the MORF hallmark domain, the presence of

additional domains, such as the GR domain in MORF1, MORF4, and MORF8

proteins[29], suggests the novel domain composition and underscores the need

for an updated classification of MORF proteins. Thus, in terms of domain

composition, we classify our identified 2,759 MORF proteins into four groups:

(1) Single-Pure MORF (SP-MORF, *n*=1,916), containing a single MORF domain;

(2) Single-Mixed MORF (SM-MORF, *n*=544), containing a single MORF domain

plus one or more non-MORF domains; (3) Multiple-Pure MORF (MP-MORF,

n=42), containing multiple MORF domains; and (4) Multiple-Mixed MORF (MM-

MORF, *n*=257), containing multiple MORF domains plus one or more non-

MORF domains (Fig. 6a and Supplementary Table 16).

Furthermore, we reconstruct the evolutionary landscape of these reclassified

MORF proteins across 262 angiosperms and 13 gymnosperms (Fig. 6b).

Specifically, SP-MORF proteins are widespread in gymnosperms and

angiosperms with larger numbers, whereas SM-MORF proteins are abundant

in angiosperms but rare in gymnosperms. Conversely, MP-MORF proteins

occur sparsely in both lineages and MM-MORF proteins are only observed in

angiosperms. Importantly, these reclassified MORF proteins exhibit a wide

range of domain compositions. Among them, 492 SM-MORF and 9 MM-MORF

proteins harbor a GR domain alongside MORF1~6, 8, and 9 domains in

angiosperms, whereas in the gymnosperm *T. grandis*, a single SM-MORF

protein contains a GR domain co-occurring with MORF9 domain, extending the

previously observed MORF-GR domain architectures in MORF1, MORF4, and

MORF8 proteins[29]. Surprisingly, we find that MORF domains and P motifs are

fused in two SM-MORF proteins, viz., AY_017782_RA in the angiosperm

Aquilaria yunnanensis and KAH9320946.1 in the gymnosperm *Taxus chinensis*

(Supplementary Table 16). Moreover, MORF domain and the RRM hallmark

domain of ORRM are fused in SM- and MM-MORF proteins in angiosperms.

Specifically, these domain fusions fall into four types: 1-MORF & 1-RRM (n=23),

1-MORF & 3-RRM (n=1), 2-MORF & 1-RRM (n=243), 3-MORF & 1-RRM (n=3)

(Supplementary Fig. 17 and Supplementary Table 16), where the leading

number indicates the copy number of each corresponding domain. Notably, the

most abundant fused type, 2-MORF & 1-RRM, is represented by the RNA

editing factor ORRM1[69].

ORRM proteins, specifically, ORRM1 to ORRM6, have been identified in

several angiosperms as mediators of RNA editing[23,70]. Likewise, we reclassify

our identified 7,760 ORRM proteins into SP-ORRM (n=2,689), SM-ORRM

(n=1,751), MP-ORRM (n=2,948), and MM-ORRM (n=372) (Supplementary

Table 17). Furthermore, we reconstruct their evolutionary landscape across

Archaeplastida, and find that SP-, SM-, and MP-ORRM proteins exhibit a

gradual increase from glaucophytes to angiosperms with larger number, while

MM-ORRM proteins show a sparse distribution (Supplementary Fig. 18). In

addition to previously identified ORRM1 to ORRM6 with a single RRM domain,

MP-ORRM proteins, such as CP29A, CP31A, and CP31B, each containing two

RRM domains, also affect RNA editing[23]. Intriguingly, numerous RRM and PPR

domains are fused in SM- and MM-ORRM proteins across Archaeplastida

(Supplementary Fig. 19). Specifically, SM-ORRM proteins contain the highest

number of RRM-P fusions in angiosperms (n=171), followed by hornworts (n=6),

mosses (*n*=4), liverworts (*n*=4), charophytes (*n*=3), gymnosperms (*n*=3),

lycophytes (*n*=1), and ferns (*n*=1). Moreover, SM-ORRM proteins exhibit one

RRM-DYW fusion in hornworts and angiosperms, and three RRM-E fusions in

angiosperms (Supplementary Table 17). Conversely, MM-ORRM proteins have

only five RRM-P and one RRM-E fusions in angiosperms. In addition, fusions

of RRM and the RanBP2 Znf hallmark domain of OZ protein are detected in

proteins that are not targeted to mitochondria or plastids (Supplementary Table

18).

Similarly, we reclassify OZ proteins and explore their evolution using the same

approach applied to MORF and ORRM proteins. Consequently, we identify 761

MP-OZ, 61 MM-OZ, 6 SP-OZ, and 1 SM-OZ proteins (Supplementary Table 19).

Based on their evolutionary landscape across Archaeplastida, we find that MP-

OZ proteins are enriched in angiosperms (*n*=725), followed by mosses (*n*=14),

lycophytes (*n*=6), ferns (*n*=6), gymnosperms (*n*=4), chlorophytes (*n*=3) and

liverworts (*n*=3) (Supplementary Fig. 20). Additionally, 60 MM-OZ, 6 SP-OZ,

and 1 SM-OZ proteins are found in angiosperms, while a single MM-OZ occurs

in hornworts. Notably, we detect that RanBP2 Znf and PPR domains (P and E)

are fused in four angiosperm proteins, despite their absence of targeting to

mitochondria or plastids (Supplementary Table 20). Collectively, our results

indicate that hallmark domains of PPR, MORF, ORRM, and OZ proteins can be

fused with more diverse patterns than previously thought (Fig. 6c).

Discussion

Our kingdom-wide characterization of RNA editing factors across

Archaeplastida reveals a complex evolutionary trajectory defined by massive

gene expansions, HGT, and the diversity and fusion of hallmark domains.

PPR genes, the primary mediators of plant RNA editing, exhibit dramatic

expansions in hornworts, mosses, and lycophytes. We demonstrate that these

expansions are predominantly driven by dispersed duplication associated with

retroposition, evidenced by signatures such as intron loss, polyA tails, and

flanking TEs. In contrast, WGD plays a subsidiary role in PPR expansion in

lycophytes and contributes little to that in the moss *T. lepidozioides* (Fig. 2a).

Notably, *T. lepidozioides* exhibits predominant dispersed duplication but limited

retroposition features, suggesting large-scale gene fractionation after an

ancient WGD event[71]. These results imply that retroposition, as well as WGD,

provided the genomic plasticity necessary for early land plants to diversify their

RNA editing machinery during the colonization of harsh terrestrial

environments[4].

The evolutionary reach of plant RNA editing extends far beyond Archaeplastida.

The discovery of DYW subgroup PPR genes in diverse non-plant lineages—

including Metazoa, Amoebozoa, Fungi, Haptista, and Rhizaria—indicates that

DYW-mediated C-to-U editing is a broadly distributed eukaryotic mechanism,

which is more common than previously recognized[41,72,73]. Noticeably, the high

expression of these genes in both recipient bdelloid rotifers (Metazoa) and their

putative plant donors (e.g., *T. lepidozioides*), coupled with shared resistance to

UV radiation, suggests a convergent adaptation where RNA editing may

facilitate the repair of UV-induced damage[74,75]. This highlights the functional

redeployment of editing factors following cross-kingdom transfer to meet

specific environmental pressures.

MORF proteins, restricted to seed plants, display significant structural similarity

between their hallmark MORF domains and PA domains of other proteins.

Specifically, MORF3 and MORF8 domains more closely resemble PA domains

associated with protein folding, whereas MORF1 domain has been reported to

be more structurally similar to an N-terminal ferredoxin-like domain that confers

RNA substrate positioning in bacterial 4-thio-uracil tRNA synthetases[28]. Our

results suggest that the structural diversity of MORF domains underpins their

functional versatility in RNA editing regulation.

Prompted by the structural diversity observed among MORF domains, we

extended our analysis to MORF, ORRM, and OZ proteins, systematically

reclassifying them based on domain composition. This reclassification revealed

not only a broader repertoire of potential RNA editing factors, but also extensive

hallmark domain fusions among PPR, MORF, ORRM, and OZ proteins (except

between MORF and OZ), mirroring their reported interactions[17,24,26,27,29] and

exemplifying the general link between domain fusion and protein-protein

networking[76-78]. Based on these findings, we propose an evolutionary model

where the composition of the editing machinery has progressively diversified

from hornworts to angiosperms, likely contributing to the maintenance of

efficient RNA editing (Extended Data Fig. 1). Collectively, these insights provide

a refined view of the genomic innovations that have shaped the plant RNA

machinery.

Materials and Methods

Collection and quality control of Archaeplastida genomes

To estimate the numbers of PPR, MORF, ORRM, and OZ genes and

reconstruct their evolutionary landscapes, 364 Archaeplastida high-quality

genomes were retrieved from public databases (Supplementary Table 1). To

ensure accurate gene counts, only the longest protein isoform of each gene

was retained. The completeness of annotated gene sets was estimated by

Benchmarking Universal Single-Copy Orthologue (BUSCO) (version 5.8.2) with

three   datasets   chlorophyta_odb10,   viridiplantae_odb10,   and

eukaryota_odb10[79]. As these datasets contain abundant sequences of

chlorophytes and angiosperms, genomes in these two lineages with BUSCO

scores ("complete") greater than 90% were retained. For other lineages,

chromosome-level genomes with BUSCO scores ("complete") greater than

85%; otherwise, the top 20% genomes with the highest BUSCO scores were

retained within each lineage.

Phylogenomic tree reconstruction of Archaeplastida

To reconstruct the phylogenomic tree of 364 Archaeplastida species, single-

copy orthologous gene families were inferred by OrthoFinder (version

3.0.1b1)[80], and 138 single-copy orthologous gene families covering at least 75%

species were retained. These single-copy orthologous gene families were

aligned by MAFFT (version 7.453)[81] with the parameter "--auto", and poorly

aligned regions were identified and trimmed by BMGE (version 1.12) with

BLOSUM30[82]. Gene trees were constructed by IQ-TREE (version 2.1.2)[83], with

the best-fit substitution model for each gene automatically selected according

to Bayesian Information Criterion, and using parameters "-B 1000 -bnni -m MFP

-T AUTO -ntmax 20". The resulting trees were subsequently merged, and the

species tree was inferred by ASTRAL (version 5.7.8)[84] using *Cyanophora*

paradoxa as the outgroup.

Identification of PPR, MORF, ORRM, and OZ proteins

The hallmark domains and motifs of PPR (P, P1, P2, S1, S2, SS, L1, L2, E1,

E2, E+, DYW, and DYW:KP), MORF (MORF), ORRM (RRM), and OZ (RanBP2

Znf) proteins were identified using hmmsearch in HMMER (version 3.2.1)[85] with

corresponding domain and motif profiles. PPR hallmark domain and motif

profiles were obtained from a previous study[34] and searched using hmmsearch

with parameters "--domtblout -noali -E 0.1". All hmmsearch scores for P, P1, P2,

S1, S2, L1, L2, E1, E2, and E+ domains or motifs were greater than 0, with SS

motif greater than 10, and DYW/DYW:KP domain greater than 30. As most PPR

hallmark domain-containing proteins are localized to mitochondria and/or

plastids[19,86], they are defined as PPR proteins. In addition, PPR proteins were

required to contain at least three of P, P1, P2, L1, L2, or SS motifs, unless they

contained a DYW or DYW:KP domain.

The MORF domain profile, constructed using hmmbuild in HMMER (version

3.2.1) with nine MORF domains from *A. thaliana*, was searched using

hmmsearch with parameters "--domtblout -noali -E 1e-10". As MORF domain

was homologous to PA domain, only hits with hmmsearch score greater than

36 were kept. The RRM domain profile was obtained from Pfam (PF00076.26)[87]

and searched using hmmsearch with parameters "--domtblout -noali -E 1e-10".

The hmmsearch score was greater than 0. GR domains, present in some

MORF and ORRM proteins, were defined as regions containing at least eight

glycines within a 20-amino-acid sliding window. The RanBP2 Znf profile was

obtained from Pfam (PF00641.22) and searched using hmmsearch with the

same parameters as for the RRM domain. As MORF, ORRM, and OZ proteins

were localized to mitochondrion and/or plastids, their subcellular localization

was predicted using DeepLoc (version 2.1)[88] with "Accurate" mode, TargetP

(version 2)[89], Plant-mSub[90] (https://bioinfo.usu.edu/Plant-mSubP), and

LOCALIZER[91] (https://localizer.csiro.au). Only localization results supported by

at least two tools were retained. The domain composition was identified using

InterPro (https://www.ebi.ac.uk/interpro)[92].

WGD Inference

To infer WGD, intra-genome alignments were conducted using BLASTP in the

Basic Local Alignment Search Tool (BLAST, version 2.12.0+)[93] with an E-value

cutoff of 1e-5. Based on the alignments, homologous dotplots were generated

using WGDI[94], and synteny blocks were inferred using MCScanX[95]. Ks values

of homologs in synteny blocks were calculated using KaKs_Calulator (version

3.0)[96] and ParaAT (version 2.0)[97] with parameters "-p proc -m mafft -f axt -g -k".

The median Ks value for each synteny block was calculated. Collectively, Ks

peak counts and synteny blocks in the homologous dotplot were used to infer

the number of WGD events.

Identification and classification of duplicated gene pairs

To identify duplicated gene pairs, all protein sequences within a genome were

aligned using BLASTP (BLAST, version 2.12.0+) with an E-value cutoff of 1e-5.

The highest-scoring alignment for each gene was considered a duplicated pair.

Alignments covering more than 60% of the longer gene (query or subject) were

required to have a sequence similarity exceeding 20%. Alternatively, for

alignments covering less than 60% of the longer gene, the alignment was

required to span at least 105 amino acid residues with a sequence similarity

exceeded 40%. Otherwise, the query and subject genes were considered

singletons. Based on genomic loci, duplicated gene pairs were classified into

four categories: tandem pairs (adjacent genes), proximal pairs (within 10 loci

on the same chromosome), dispersed pairs (more than 10 loci apart on the

same chromosome or on different chromosomes), and WGD pairs (located in

synteny blocks).

TE annotation

TEs were identified by a combination of *ab initio* and homology-based methods.

For *ab initio* prediction, a consensus sequence library was built using

RepeatModeler (version 2.0.6)[98] with the parameter "-LTRStruct". An LTR

library was then constructed using LTR_Finder (version

1.07)[99]/LTR_FINDER_parallel (version 1.1)[100], LTRharvest (version 1.6.6)[101]

with parameters "-minlenltr 100 -maxlenltr 7000", and LTR_retriever (version

2.9.0)[102]. Both libraries were used to annotate TEs using RepeatMasker

(version 4.1.8)[103]. For homology-based detection, TEs were identified using

RepeatMasker (version 4.1.8) with Dfam (version 3.9)[104] database. TE

annotations obtained from both methods were then combined.

Retroposition inference

Retroposition, underlying dispersed duplication gene pairs, was characterized

by intron loss, polyA enrichment downstream of the 3'UTR, and flanking TEs.

First, intron loss in one dispersed duplication gene pair was required to meet

one of following criteria: (1) both genes had no intron; (2) a gene had no intron,

while the other had at least one intron; (3) considering intron retention, a gene

contained one intron, while the other had at least one intron; (4) both genes had

more than one intron, differing by more than 10 introns. Second, polyA

enrichment downstream of the 3'UTR of the intronless gene (or polyT

enrichment upstream of the 5'UTR) was defined as at least eight

adenines/thymines within a 10-base window. Genomic annotations were used

to detect polyA/T enrichment in regions extending 1 kb downstream of the

3'UTR or upstream of the 5'UTR. If the 3'UTR or 5'UTR of a gene was not

annotated, the shortest corresponding UTR in the genome was used. If no

3'UTR or 5'UTR was annotated in the genome, they were defined as the 100-

bp regions downstream of the stop codon and upstream of the start codon,

respectively. Third, dispersed duplication gene pairs with TEs located within 20

kb upstream and downstream were defined as TE-flanked.

Identification of DYW and DYW:KP domain homologs and HGT

inference

To explore HGT of plant DYW and DYW:KP domains, 10,407 protein

sequences (hmmsearch score ≥ 30, length ≥ 130 aa) of 68 representative

species were collected (Supplementary Table 8). DYW and DYW:KP domain

homologs were searched in the NCBI Non-redundant Protein Sequence

Database (NR) (acquisition date: 2025.01.19) using BLASTP (BLAST, version

2.12.0+) with an E-value cutoff of 1e-5. Protein sequences retrieved from NR

originated from 40 lineages (Supplementary Table 9). Homologs exhibiting

more than 20% sequence similarity to plant DYW and DYW:KP domains were

further confirmed via hmmsearch in HMMER (version 3.2.1) using a score cutoff

of 30. To avoid contamination, non-plant DYW and DYW:KP domains were

retained only if present in at least two species within the corresponding lineage.

Otherwise, non-plant DYW and DYW:KP domains detected in a single species

were retained only if present in at least one additional genome assembly.

To reconstruct the phylogenetic tree of DYW and DYW:KP domains from plants

and non-plants, domains from six representative plants (*P. pearsonii*, *T.*

lepidozioides, *I. sinensis*, *A. spinulosa*, *G. montanum*, and *A. trichopoda*) and

all non-plant counterparts were aligned using MAFFT (version 7.525), with

poorly aligned regions trimmed by trimAl (version 1.5.rev0)[105] using parameters

"-gt 0.7 -cons 60 -st 0.1". The phylogenetic tree was inferred using IQ-TREE

(version 2.1.2) (Best-fit model: JTT + R10) with parameters "-B 1000 -bnni -m

MFP -T AUTO --safe" and visualized with iTOL (version 7)[106]. Bootstraps were

from 1,000 replicates. Multiple sequence alignment of DYW domains from

plants and non-plants was performed using MEGA (version 11)[107] and

visualized with Jalview (version 2.11.4.0)[108].

Identification of MORF domain homologs

To identify homologs of plant MORF domains, 334 protein sequences

(hmmsearch score ≥ 0, length ≥ 86 aa) from 33 representative species were

collected (Supplementary Table 14). These MORF domain homologs searched

in the NCBI NR using BLASTP (BLAST, version 2.12.0+) with an E-value cutoff

of 1e-5. The protein sequences retrieved from NR originated from 45 lineages

(Supplementary Table 9). Homologs with more than 20% similarity to plant

MORF domains were retained. The MORF domains and their homologs were

aligned using MAFFT (version 7.525), and the phylogenetic tree was

constructed using FastTree[109] (Model: Jones-Taylor-Thorton, CAT

approximation with 20 rate categories) with parameters "-pseudo -gamma" and

visualized by iTOL (version 7).

Phylogenetic analysis of MORF domains

To reconstruct the phylogenetic tree of MORF domains, all MORF domains from

gymnosperms and angiosperms were selected. These domains were aligned

to MORF1~MORF9 domains from *A. thaliana* to assign their types. Five PA

domains from non-seed plants (Dicom.22G070000.1.p, HaB36G042,

Phcar.S3G189600.t1, Phsp.C2G192800.t1, XP_024533311.1), exhibiting the

highest similarity to MORF domains, were used as the outgroup. All MORF

domains, along with the outgroup sequences, were aligned using MAFFT

(version 7.525). The phylogenetic tree was inferred using IQ-TREE (version

2.1.2) (Best-fit model: JTT + R8) with parameters "-B 1000 -bnni -m MFP -T

AUTO" and visualized by iTOL (version 7)[106]. Bootstraps were from 1,000

replicates.

Analysis of PPR and MORF gene expression

To calculate gene expression, reference genome indexes were built using

hisat2-build in HISAT2 (version 2.0.5)[110] and rsem-prepare-reference in RSEM

(version 1.3.1)[111]. Raw RNA sequencing data were quality-filtered using fastp

(version 0.20.1)[112] with parameters "-q 20 -u 40 -l 15 -g -x -r -W 4 -M 20 -w 12".

Strand specificity was inferred using infer_experiment.py in RSeQC (version

2.6.4)[113]. Reads mapping was conducted using STAR (version 2.7.1a)[114]. Gene

expression matrix was generated using rsem-calculate-expression in RSEM

(version 1.3.1) with parameters "--keep-intermediate-files --time --star --

append-names --output-genome-bam --sort-bam-by-coordinate".

Protein structure prediction and alignment

620 Protein structures were predicted using ColabFold
621 (https://colab.research.google.com/github/sokrypton/ColabFold)[115], and
622 different protein structures were aligned using "Matchmaker" in ChimeraX
623 (version 1.9)[116].

Data availability

625 All data used and generated in this study are available at the Open Archive for
626 Miscellaneous Data (OMIX: OMIX012585)[117] in the National Genomics Data
627 Center (NGDC), China National Center for Bioinformation / Beijing Institute of
628 Genomics, Chinese Academy of Sciences[118].

Code availability

630 The computational codes used in this study have been deposited in GitHub
631 (https://github.com/Bioinfo-Ming/PlantEditosomeEvolution).

ACKNOWLEDGEMENTS

We thank Shuhui Song, Lina Ma, Chao Zhang, Tianyi Xu, Tongtong Zhu,
Dechang Yang, Shuangyang Wu, Lin Liu, Zhao Li, Yang Zhang, Xing Zheng,
Yue Qi, Xinyu Zhou, Wenzhuo Cheng, Yuxin Qin, Miaomiao Wang, Shiting
Wang, Lingjie Wang, Zihan Wang, Kehua Ma, Pan Li, Yiran Zhan, and Zheng
Luo for their valuable discussions and advice. This work was supported by
grants from National Natural Science Foundation of China [T2425005 and
32030021] and International Partnership Program of the Chinese Academy of
Sciences [153F11KYSB20160008].

CONTRIBUTOR INFORMATION

M.C. collected and analyzed the data, and wrote the manuscript. Z.Z. conceived, initiated and supervised this study. Q.M.Q. and Z.Z. revised the manuscript.

FUNDING

The authors declare no competing interests.

REFERENCES

Castandet, B. & Araya, A. RNA editing in plant organelles. Why make it easy? *Biochemistry (Moscow)* **76**, 924-931 (2011).

Hao, W. *et al.* RNA editing and its roles in plant organelles. *Frontiers in Genetics* **12**, 757109 (2021).

Takenaka, M., Zehrmann, A., Verbitskiy, D., Haertel, B. & Brennicke, A. RNA editing in plants and its evolution. *Annual Review of Genetics* **47**, 335-352 (2013).

Fujii, S. & Small, I. The evolution of RNA editing and pentatricopeptide repeat genes. *New Phytologist* **191**, 37-47 (2011).

Hu, R. *et al.* Adaptive evolution of the enigmatic Takakia now facing climate change in Tibet. *Cell* **186**, 3558-3576. e3517 (2023).

Small, I. D., Schallenberg‑Rüdinger, M., Takenaka, M., Mireau, H. & Ostersetzer‑Biran, O. Plant organellar RNA editing: what 30 years of research has revealed. *The Plant Journal* **101**, 1040-1056 (2020).

Wang, Y. *et al.* Empty Pericarp21 encodes a novel PPR-DYW protein that is required for mitochondrial RNA editing at multiple sites, complexes I and V biogenesis, and seed development in maize. *PLoS Genetics* **15**, e1008305 (2019).

8   Sun, Y. K., Gutmann, B., Yap, A., Kindgren, P. & Small, I. Editing of
chloroplast rps14 by PPR editing factor EMB2261 is essential for
Arabidopsis development. *Frontiers in Plant Science* **9**, 841 (2018).
9   Li, J. *et al.* The Pentatricopeptide Repeat Protein OsPPR674 Regulates
Rice Growth and Drought Sensitivity by Modulating RNA Editing of the
Mitochondrial Transcript ccmC. *International Journal of Molecular Sciences*
26, 2646 (2025).
10   Shi, X., Germain, A., Hanson, M. R. & Bentolila, S. RNA recognition motif-
containing protein ORRM4 broadly affects mitochondrial RNA editing and
impacts plant development and flowering. *Plant Physiology* **170**, 294-309
(2016).
11   Yang, Y. *et al.* The RNA editing factor SlORRM4 is required for normal fruit
ripening in tomato. *Plant Physiology* **175**, 1690-1702 (2017).
12   Luo, Z. *et al.* Pentatricopeptide repeat gene-mediated mitochondrial RNA
editing impacts on rice drought tolerance. *Frontiers in Plant Science* **13**,
926285 (2022).
13   Yuan, H. & Liu, D. Functional disruption of the pentatricopeptide protein
SLG1 affects mitochondrial RNA editing, plant development, and
responses to abiotic stresses in Arabidopsis. *The Plant Journal* **70**, 432-
444 (2012).
14   Sun, T., Bentolila, S. & Hanson, M. R. The unexpected diversity of plant
organelle RNA editosomes. *Trends in Plant Science* **21**, 962-973 (2016).
15   Wang, T. & Takenaka, M. The molecular basis and evolution of the
organellar RNA editosome by complementary DYW deaminases in seed
plants. *Plant Physiology* **197**, kiaf142 (2025).
16   Knoop, V. C-to-U and U-to-C: RNA editing in plant organelles and beyond.
Journal of Experimental Botany **74**, 2273-2294 (2023).
17   Yan, J., Zhang, Q. & Yin, P. RNA editing machinery in plant organelles.

Science China Life Sciences **61**, 162-169 (2018).

18 Ichinose, M. & Sugita, M. RNA editing and its molecular mechanism in plant

organelles. *Genes* **8**, 5 (2016).

19 Wang, Y. & Tan, B.-C. Pentatricopeptide repeat proteins in plants: Cellular

functions, action mechanisms, and potential applications. *Plant*

Communications **6** (2025).

20 Yan, J. *et al.* MORF9 increases the RNA-binding activity of PLS-type

pentatricopeptide repeat protein in plastid RNA editing. *Nature Plants* **3**, 1-

8 (2017).

21 Wang, Y. *et al.* Maize PPR-E proteins mediate RNA C-to-U editing in

mitochondria by recruiting the trans deaminase PCW1. *The Plant Cell* **35**,

529-551 (2023).

22 Yang, Y.-Z. *et al.* GRP23 plays a core role in E-type editosomes via

interacting with MORFs and atypical PPR-DYWs in Arabidopsis

mitochondria. *Proceedings of the National Academy of Sciences* **119**,

e2210978119 (2022).

23 Shi, X., Bentolila, S. & Hanson, M. R. Organelle RNA recognition motif-

containing (ORRM) proteins are plastid and mitochondrial editing factors in

Arabidopsis. *Plant Signaling & Behavior* **11**, 294-309 (2016).

24 Sun, T. *et al.* An RNA recognition motif-containing protein is required for

plastid RNA editing in Arabidopsis and maize. *Proceedings of the National*

Academy of Sciences **110**, E1169-E1178 (2013).

25 Sun, T. *et al.* A zinc finger motif-containing protein is essential for

chloroplast RNA editing. *PLoS Genetics* **11**, e1005028 (2015).

26 Gipson, A. B., Hanson, M. R. & Bentolila, S. The RanBP2 zinc finger

domains of chloroplast RNA editing factor OZ1 are required for protein–

protein interactions and conversion of C to U. *The Plant Journal* **109**, 215-

226 (2022).

27 Wang, Z. *et al.* OsOZ1, a zinc finger motif-containing protein, is essential

for chloroplast development through RNA editing and intron splicing in rice.

Plant Growth Regulation **105**, 449-461 (2025).

28 Haag, S. *et al.* Crystal structures of the Arabidopsis thaliana organellar RNA

editing factors MORF1 and MORF9. *Nucleic Acids Research* **45**, 4915-

4928 (2017).

29 Li, J., Yuan, J., Jing, Y. & Lin, R. MORF proteins: A small family regulating

organellar RNA editing and beyond. *Journal of Integrative Plant Biology*

(2025).

30 Takenaka, M. *et al.* Multiple organellar RNA editing factor (MORF) family

proteins are required for RNA editing in mitochondria and plastids of plants.

Proceedings of the National Academy of Sciences **109**, 5104-5109 (2012).

31 Sugita, M., Uchiyama, H. & Ichinose, M. RRM and PPR proteins network

as a global regulator for plastid RNA metabolism. *Endocytobiosis Cell Res.*

27, 41-44 (2016).

32 Baudry, K., Monachello, D., Castandet, B., Majeran, W. & Lurin, C.

Dissecting the molecular puzzle of the editosome core in Arabidopsis

organelles. *Plant Science* **344**, 112101 (2024).

33 Shikanai, T. RNA editing in plant organelles: machinery, physiological

function and evolution. *Cellular and Molecular Life Sciences CMLS* **63**, 698-

708 (2006).

34 Gutmann, B. *et al.* The expansion and diversification of pentatricopeptide

repeat RNA-editing factors in plants. *Molecular plant* **13**, 215-230 (2020).

35 Manna, S. An overview of pentatricopeptide repeat proteins and their

applications. *Biochimie* **113**, 93-99 (2015).

36 Manna, S. & Barth, C. Identification of a novel pentatricopeptide repeat

subfamily with a C-terminal domain of bacterial origin acquired via ancient

horizontal gene transfer. *BMC Research Notes* **6**, 525 (2013).

37 Schallenberg-Rüdinger, M., Lenz, H., Polsakiewicz, M., Gott, J. M. & Knoop,

V. A survey of PPR proteins identifies DYW domains like those of land plant

RNA editing factors in diverse eukaryotes. *RNA Biology* **10**, 1549-1556

(2013).

38 Knoop, V. & Rüdinger, M. DYW-type PPR proteins in a heterolobosean

protist: plant RNA editing factors involved in an ancient horizontal gene

transfer? *FEBS Letters* **584**, 4287-4291 (2010).

39 Schaap, P. *et al.* The Physarum polycephalum genome reveals extensive

use of prokaryotic two-component and metazoan-type tyrosine kinase

signaling. *Genome Biology and Evolution* **8**, 109-125 (2016).

40 Manna, S., Brewster, J. & Barth, C. Identification of pentatricopeptide

repeat proteins in the model organism Dictyostelium discoideum.

International Journal of Genomics **2013**, 586498 (2013).

41 Fu, C.-J., Sheikh, S., Miao, W., Andersson, S. G. & Baldauf, S. L. Missing

genes, multiple ORFs, and C-to-U type RNA editing in Acrasis kona

(Heterolobosea, Excavata) mitochondrial DNA. *Genome Biology and*

Evolution **6**, 2240-2257 (2014).

42 Puttick, M. N. *et al.* The interrelationships of land plants and the nature of

the ancestral embryophyte. *Current Biology* **28**, 733-745. e732 (2018).

43 Wu, W. *et al.* Genomic convergence in terrestrial root plants through

tandem duplication in response to soil microbial pressures. *Cell Reports* **43**

(2024).

44 Shen, Y. *et al.* Chromosome-level and haplotype-resolved genome

provides insight into the tetraploid hybrid origin of patchouli. *Nature*

Communications **13**, 3511 (2022).

45 Xu, Z. *et al.* Crocus genome reveals the evolutionary origin of crocin

biosynthesis. *Acta Pharmaceutica Sinica B* **14**, 1878-1891 (2024).

46 Wang, Z. *et al.* Near-complete assembly and comprehensive annotation of

the wheat Chinese Spring genome. *Molecular Plant* **18**, 892-907 (2025).

47 Panchy, N., Lehti-Shiu, M. & Shiu, S.-H. Evolution of gene duplication in

plants. *Plant Physiology* **171**, 2294-2316 (2016).

48 Freeling, M. Bias in plant gene content following different sorts of

duplication: tandem, whole-genome, segmental, or by transposition.

Annual Review of Plant Biology **60**, 433-453 (2009).

49 Wang, X. *et al.* A recent burst of gene duplications in Triticeae. *Plant*

Communications **3** (2022).

50 Feng, X. *et al.* Genomes of multicellular algal sisters to land plants

illuminate signaling network evolution. *Nature Genetics* **56**, 1018-1031

(2024).

51 Zhang, J. *et al.*   (2020).

52 Li, F.-W. *et al.* Anthoceros genomes illuminate the origin of land plants and

the unique biology of hornworts. *Nature Plants* **6**, 259-272 (2020).

53 Schafran, P. *et al.* Pan-phylum genomes of hornworts reveal conserved

autosomes but dynamic accessory and sex chromosomes. *Nature Plants*

11, 49-62 (2025).

54 Li, C. *et al.* Extraordinary preservation of gene collinearity over three

hundred million years revealed in homosporous lycophytes. *Proceedings*

of the National Academy of Sciences **121**, e2312607121 (2024).

55 Cui, J. *et al.* Chromosome-level reference genome of tetraploid Isoetes

sinensis provides insights into evolution and adaption of lycophytes.

GigaScience **12**, giad079 (2023).

56 Zhang, C., Gschwend, A. R., Ouyang, Y. & Long, M. Evolution of gene

structural complexity: an alternative-splicing-based model accounts for

intron-containing retrogenes. *Plant Physiology* **165**, 412-423 (2014).

57 Schrader, L. & Schmitz, J. The impact of transposable elements in adaptive

evolution. *Molecular Ecology* **28**, 1537-1549 (2019).

58 Takenaka, M. *et al.* DYW domain structures imply an unusual regulation

principle in plant organellar RNA editing catalysis. *Nature catalysis* **4**, 510-

522 (2021).

59 Boussardon, C. *et al.* The cytidine deaminase signature HxE(x)nCxxC of

DYW1 binds zinc and is necessary for RNA editing of ndhD‑1. *New*

Phytologist **203**, 1090-1095 (2014).

60 Yang, Y. *et al.* DYW cytidine deaminase domains have a long‑range impact

on RNA recognition by the PPR array of chimeric plant C-to-U RNA editing

factors and strongly affect target selection. *The Plant Journal* **116**, 840-854

(2023).

61 Hayes, M. L., Giang, K., Berhane, B. & Mulligan, R. M. Identification of two

pentatricopeptide repeat genes required for RNA editing and zinc binding

by C-terminal cytidine deaminase-like domains. *Journal of Biological*

Chemistry **288**, 36519-36529 (2013).

62 Bolhuis, P. Sampling kinetic protein folding pathways using all-atom models.

Lecture Notes in Physics, 703, 393-433 (2006).

63 Hespeels, B. *et al.* Back to the roots, desiccation and radiation resistances

are ancestral characters in bdelloid rotifers. *BMC Biology* **21**, 72 (2023).

64 Wang, W., Yang, Y., Örstan, A., He, Z. & Wang, Q.   Species diversity of

bdelloid rotifer (Rotifera, Bdelloidea) in different areas in China, with a

description of two new species. *ZooKeys* **1229**, 1 (2025).

65 Jain, S. C., Shinde, U., Li, Y., Inouye, M. & Berman, H. M. The crystal

structure of an autoprocessed Ser221Cys-subtilisin E-propeptide complex

at 2.0 Å resolution. *Journal of Molecular Biology* **284**, 137-144 (1998).

66 Li, Y., Hu, Z., Jordan, F. & Inouye, M. Functional Analysis of the Propeptide

of Subtilisin E as an Intramolecular Chaperone for Protein Folding:

Refolding And Inhibitory Abilities of Propeptide Mutants (∗). *Journal of*

Biological Chemistry **270**, 25127-25132 (1995).

67 Wang, D. *et al.* Genome-wide analysis of multiple organellar RNA editing

factor family in poplar reveals evolution and roles in drought stress.

International Journal of Molecular Sciences **20**, 1425 (2019).

68 Liu, P.-Z. *et al.* Genome-wide identification and expression analysis of the

MORF gene family in celery reveals their potential role in chloroplast

development. *Journal of Genetic Engineering and Biotechnology* **22**,

100443 (2024).

69 Searing, A. M., Satyanarayan, M. B., O′Donnell, J. P. & Lu, Y. Two

organelle RNA recognition motif proteins affect distinct sets of RNA editing

sites in the Arabidopsis thaliana plastid. *Plant Direct* **4**, e00213 (2020).

70 Shi, X., Hanson, M. R. & Bentolila, S. Two RNA recognition motif-containing

proteins are plant mitochondrial editing factors. *Nucleic Acids Research* **43**,

3814-3825 (2015).

71 Liang, Z. & Schnable, J. C. Functional divergence between subgenomes

and gene pairs after whole genome duplications. *Molecular Plant* **11**, 388-

397 (2018).

72 Rüdinger, M., Fritz-Laylin, L., Polsakiewicz, M. & Knoop, V. Plant-type

mitochondrial RNA editing in the protist Naegleria gruberi. *RNA* **17**, 2058-

2062 (2011).

73 Yang, J., Harding, T., Kamikawa, R., Simpson, A. G. & Roger, A. J.

Mitochondrial genome evolution and a novel RNA editing system in deep-

branching heteroloboseids. *Genome Biology and Evolution* **9**, 1161-1174

(2017).

74 He, P., Huang, S., Xiao, G., Zhang, Y. & Yu, J. Abundant RNA editing sites

of chloroplast protein-coding genes in Ginkgo biloba and an evolutionary

pattern analysis. *BMC Plant Biology* **16**, 257 (2016).

75 Xhemalçe, B., Miller, K. M. & Gromak, N. Epitranscriptome in action: RNA

860 modifications in the DNA damage response. *Molecular Cell* **84**, 3610-3626

861 (2024).

862 76 Enright, A. J., Iliopoulos, I., Kyrpides, N. C. & Ouzounis, C. A. Protein

863 interaction maps for complete genomes based on gene fusion events.

864 *Nature* **402**, 86-90 (1999).

865 77 Marcotte, E. M., Pellegrini, M., Thompson, M. J., Yeates, T. O. & Eisenberg,

866 D. A combined algorithm for genome-wide prediction of protein function.

867 *Nature* **402**, 83-86 (1999).

868 78 Marcotte, E. M. *et al.* Detecting protein function and protein-protein

869 interactions from genome sequences. *Science* **285**, 751-753 (1999).

870 79 Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. & Zdobnov, E. M.

871 BUSCO update: novel and streamlined workflows along with broader and

872 deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and

873 viral genomes. *Molecular Biology and Evolution* **38**, 4647-4654 (2021).

874 80 Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for

875 comparative genomics. *Genome Biology* **20**, 1-14 (2019).

876 81 Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software

877 version 7: improvements in performance and usability. *Molecular Biology*

878 *and Evolution* **30**, 772-780 (2013).

879 82 Criscuolo, A. & Gribaldo, S. BMGE (Block Mapping and Gathering with

880 Entropy): a new software for selection of phylogenetic informative regions

881 from multiple sequence alignments. *BMC Evolutionary Biology* **10**, 1-21

882 (2010).

883 83 Minh, B. Q. *et al.* IQ-TREE 2: new models and efficient methods for

884 phylogenetic inference in the genomic era. *Molecular Biology and Evolution*

885 **37**, 1530-1534 (2020).

886 84 Mirarab, S. *et al.* ASTRAL: genome-scale coalescent-based species tree

887 estimation. *Bioinformatics* **30**, i541-i548 (2014).

85   Eddy, S. Sequence analysis using profile hidden Markov models.
http://hmmer. wustl. edu (2004).

86   Meng, L. *et al.* PPR proteins in plants: Roles, mechanisms, and prospects
for rice research. *Frontiers in Plant Science* **15**, 1416742 (2024).

87   Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids*
Research **49**, D412-D419 (2021).

88   Ødum, M. T. *et al.* DeepLoc 2.1: multi-label membrane protein type
prediction using protein language models. *Nucleic Acids Research* **52**,
W215-W220 (2024).

89   Armenteros, J. J. A. *et al.* Detecting sequence signals in targeting peptides
using deep learning. *Life Science Alliance* **2** (2019).

90   Sahu, S. S., Loaiza, C. D. & Kaundal, R. Plant-mSubP: a computational
framework for the prediction of single-and multi-target protein subcellular
localization using integrated machine-learning approaches. *AoB Plants* **12**,
plz068 (2020).

91   Sperschneider, J. *et al.* LOCALIZER: subcellular localization prediction of
both plant and effector proteins in the plant cell. *Scientific Reports* **7**, 44598
(2017).

92   Blum, M. *et al.* InterPro: the protein sequence classification resource in
2025. *Nucleic Acids Research* **53**, D444-D456 (2025).

93   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC*
Bioinformatics **10**, 1-9 (2009).

94   Sun, P. *et al.* WGDI: a user-friendly toolkit for evolutionary analyses of
whole-genome duplications and ancestral karyotypes. *Molecular Plant* **15**,
1841-1851 (2022).

95   Wang, Y. *et al.* MCScanX: a toolkit for detection and evolutionary analysis
of gene synteny and collinearity. *Nucleic Acids Research* **40**, e49-e49
(2012).

96 Zhang, Z. KaKs_Calculator 3.0: calculating selective pressure on coding
and non-coding sequences. *Genomics, Proteomics and Bioinformatics* **20**,
536-540 (2022).

97 Zhang, Z. *et al.* ParaAT: a parallel tool for constructing multiple protein-
coding DNA alignments. *Biochemical and Biophysical Research*
Communications **419**, 779-781 (2012).

98 Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of
transposable element families. *Proceedings of the National Academy of*
Sciences **117**, 9451-9457 (2020).

99 Xu, Z. & Wang, H. LTR_FINDER: an efficient tool for the prediction of full-
length LTR retrotransposons. *Nucleic Acids Research* **35**, W265-W268
(2007).

100 Ou, S. & Jiang, N. LTR_FINDER_parallel: parallelization of LTR_FINDER
enabling rapid identification of long terminal repeat retrotransposons.
Mobile DNA **10**, 48 (2019).

101 Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible
software for de novo detection of LTR retrotransposons. *BMC*
Bioinformatics **9**, 1-14 (2008).

102 Ou, S. & Jiang, N. LTR_retriever: a highly accurate and sensitive program
for identification of long terminal repeat retrotransposons. *Plant Physiology*
176, 1410-1422 (2018).

103 Chen, N. Using RepeatMasker to identify repetitive elements in genomic
sequences. *Current Protocols in Bioinformatics* **5**, 4.10. 11-14.10. 14 (2004).

104 Storer, J., Hubley, R., Rosen, J., Wheeler, T. J. & Smit, A. F. The Dfam
community resource of transposable element families, sequence models,
and genome annotations. *Mobile DNA* **12**, 2 (2021).

105 Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses.

Bioinformatics **25**, 1972-1973 (2009).

106 Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: recent updates to
the phylogenetic tree display and annotation tool. *Nucleic Acids Research*
52, W78-W82 (2024).

107 Tamura, K., Stecher, G. & Kumar, S. MEGA11: molecular evolutionary
genetics analysis version 11. *Molecular Biology and Evolution* **38**, 3022-
3027 (2021).

108 Procter, J. B. *et al.* Alignment of biological sequences with Jalview. *Methods*
in Molecular Biology **2231**, 203-224 (2021).

109 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2–approximately
maximum-likelihood trees for large alignments. *PloS One* **5**, e9490 (2010).

110 Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based
genome alignment and genotyping with HISAT2 and HISAT-genotype.
Nature Biotechnology **37**, 907-915 (2019).

111 Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-
Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 1-
16 (2011).

112 Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**, i884-i890 (2018).

113 Wang, L., Wang, S. & Li, W. RSeQC: quality control of RNA-seq
experiments. *Bioinformatics* **28**, 2184-2185 (2012).

114 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29, 15-21 (2013).

115 Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nature*
Methods **19**, 679-682 (2022).

116 Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis.
Protein Science **32**, e4792 (2023).

117 Zhang, S. *et al.* The GSA Family in 2025: A Broadened Sharing Platform

for Multi-Omics and Multimodal Data. *Genomics, Proteomics &*

Bioinformatics, qzaf072 (2025).

118 CNCB-NGDC Members and Partners. Database Resources of the National

Genomics Data Center, China National Center for Bioinformation in 2025.

Nucleic Acids Research **53**, D30-D44 (2025).

Figures

Fig. 1

Fig. 1: The evolutionary landscape of PPR, MORF, ORRM, and OZ genes across Archaeplastida. (a) The maximum likelihood phylogenomic tree of 364 Archaeplastida species, including glaucophytes (*n*=1), rhodophytes (*n*=7),

prasinodermophytes (*n*=1), chlorophytes (*n*=33), charophytes (*n*=7), hornworts

(*n*=22), mosses (*n*=13), liverworts (*n*=4), lycophytes (*n*=6), ferns (*n*=5),

gymnosperms (*n*=13), and angiosperms (*n*=262). The best-fit model of each

single-copy orthologous gene tree is automatically selected according to

Bayesian Information Criterion in IQ-TREE. All gene trees (*n*=138) are merged

and the final species tree is inferred by ASTRAL. Bootstraps are from 1,000

replicates. The distribution of **(b)** PPR, **(c)** MORF, **(d)** ORRM, and **(e)** OZ gene

across Archaeplastida lineages. Two lycophytes, *S. moellendorffii* and *I.*

sinensis, have the highest number of PPR genes (*n*=3,861 and 3,256,

respectively). Orange and purple bars indicate gene expansions occurring

simultaneously across all four gene families, and within individual families,

respectively. Filled and open black circles denote species that have undergone

genome diploidization/polyploidization (D/P), and those with unknown genome

expansion, respectively. D/P: *Arachis hypogaea*, *Atropa bell adonna*, *B. napus*,

C. rotundifolia, *C. sativus*, *Helianthus tuberosus*, *Papaver somniferum*, *P.*

cablin, *Pontederia crassipes*, *T. aestivum*, *Zanthoxylum bungeanum*, and

Zingiber officinale. Unknown D/P: *C. pinnatifida*, *Iris pallida*, and *M. fusca*.

Fig. 2

1001

Fig. 2: Mechanisms of PPR gene expansions. (a) Numbers of duplicated

PPR genes (original and copied) derived from WGD, TD, PD, and DD in seven

representative species (*A. agrestis*, *N. orbicularis*, *P. pearsonii*, *P. phymatodes*,

T. lepidozioides, *I. sinensis*, and *H. asiatica*) and the outgroup (*Z.*

circumcarinatum). A red five-pointed star denotes a whole genome duplication

event. WGD: Whole Genome Duplication, TD: Tandem Duplication, PD:

Proximal Duplication, DD: Dispersed Duplication. **(b)** Percentage of intronless

DD PPR gene pairs. Numbers to the left and right of the colon indicate intron

counts in each DD PPR gene pair. Categories "0:0", "0:n", "1:n", and "n:n

(difference > 10)" denote intron loss in DD PPR gene pairs. **(c)** E2 and **(d)** DYW

subgroup PPR gene pairs (DD) flanked by similar LTR/Copia and LTR/Gypsy,

respectively. **(e)** Percentages of intronless DD PPR gene pairs with polyA tails

and flanked by LTRs.

1015

Fig. 3

Lineage ■ Plant ■ Non-plant **DYW type** ■ Plant DYW ■ Plant DYW:KP ■ Non-plant DYW/DYW:KP I–IV Non-plant DYW clusters

Fig. 3: The maximum likelihood phylogenetic tree of DYW and DYW:KP

domains in plants and non-plants. Three predominant clusters (I, II, and III)

of non-plant DYW domains are embedded within plants. Among non-plant

lineages, Metazoa contain the largest number of DYW domains. Best-fit model

(IQ-TREE): JTT + R10. Bootstraps are from 1,000 replicates.

Fig. 4

Fig. 4: HGT of DYW domains from plants to bdelloid rotifers. (a) Sequence

alignments between plant and non-plant DYW domains. Amino acid residues in

rectangles denote PG box, HxE(x)nCxxC, and DYW tripeptide. **(b)** Protein

structure alignment between DYW domain (protein ID: Tle1c03524.1) in the

moss *T. lepidozioides* and its counterpart (protein ID: UJR34389.1) in the

bdelloid rotifer *A. vage*. **(c)** Expression of PPR genes (DYW, E+, PLS, and P

subgroups) in whole tissues of *A. vage*. TPM: Transcripts Per Million. **(d)**

Expression of DYW subgroup PPR genes in *T. lepidozioides* under control

condition and after 2-hour UV irradiation, followed by 0, 6, 12, and 24-hour

recovery. **(e)** A proposed model of horizontal transfer of DYW subgroup PPR

genes from mosses (e.g., *T. lepidozioides*) to bdelloid rotifers (e.g., *A. vage*).

1036

Fig. 5

1038

Fig. 5: Conservation and diversity of MORF domains. (a) The maximum-
likelihood phylogenetic tree of MORF and PA domains. The tree scale bar
represents the number of substitutions per site. Model (FastTree): Jones-
Taylor-Thorton, CAT approximation with 20 rate categories. **(b)** Sequence and
(c) structure alignments between MORF3 domain (protein ID: Pd03G21970A)
from *Pinus densiflora* and its homologous PA domain (protein ID:
XP_018897422.1) from *Bemisia tabaci*. **(d)** The maximum-likelihood
phylogenetic tree of MORF1~MORF9 domains from angiosperms and
gymnosperms. Five non-seed plant PA domains with highest sequence
similarity to MORF domains are used as the outgroup (protein IDs:
Dicom.22G070000.1.p,   HaB36G042,   Phcar.S3G189600.t1,

Phsp.C2G192800.t1, and XP_024533311.1). Best-fit model (IQ-TREE): JTT +

R8. Bootstraps are from 1,000 replicates.

1052

1053 **Fig. 6**

1054

Fig. 6: Reclassification and diversity of PPR accessory proteins. (a)
Reclassification of MORF proteins based on domain composition. Blue and
green bars represent MORF1~MORF9 and other domains, respectively. SP-
MORF: Single-Pure MORF, SM-MORF: Single-Mixed MORF; MP-MORF:
Multiple-Pure MORF; MM-MORF: Multiple-Mixed MORF. **(b)** The phylogenetic
tree of reclassified MORF proteins from angiosperms and gymnosperms. **(c)**
Fusions among PPR, MORF, RRM, and RanBP2 Znf domains. Numbers in
parentheses indicate domain counts. Proteins marked with solid and dashed
arrows are localized to mitochondria/plastids and other subcellular
compartments, respectively.

1065

1066 **Extended Data Fig. 1**

1067

Extended Data Fig. 1: A proposed model for the evolutionary

diversification of plant RNA editing factors. Editing factor composition varies

from hornworts to angiosperms, potentially maintaining efficient RNA editing.

The question mark indicates that OZ protein is rare in hornworts. Deep-rose

PPR: DYW or DYW:KP subgroup PPR protein; red-orange PPR: DYW

subgroup PPR protein. C=>U: C-to-U RNA editing; C<=>U: C-to-U or U-to-C

RNA editing.