An acyltransferase protects *Caenorhabditis elegans* from thiol reductive stress through an autoinhibitory hypoxia response pathway

Ravi[1] and Jogender Singh[1,*]

[1]Department of Biological Sciences, Indian Institute of Science Education and Research, Mohali, Punjab, 140306, India

[*]For correspondence: jogender@iisermohali.ac.in

Keywords: dithiothreitol; HIF-1; redox homeostasis; reductive stress; RHY-1

Abstract

Cellular redox homeostasis depends on a finely tuned balance between oxidizing and reducing conditions, and disturbances in this balance lead to oxidative or reductive stress. While oxidative stress and its pathological outcomes are well studied, the molecular mechanisms underlying cellular responses to reductive stress remain poorly understood. Using *Caenorhabditis elegans* as a model, we investigate thiol reductive stress induced by dithiothreitol (DTT) and uncover a critical protective role for the hypoxia response pathway. We identify RHY-1, a membrane-associated acyltransferase and known negative regulator of the hypoxia-inducible factor HIF-1, as essential for survival under thiol reductive stress. Notably, *rhy-1* is a direct transcriptional target of HIF-1, and overexpression of *rhy-1* fully rescues the sensitivity of *hif-1* loss-of-function mutants to DTT. We demonstrate that RHY-1 functions in an autoinhibitory feedback loop, where elevated RHY-1 levels suppress activation of the hypoxia response pathway even during reductive stress. Finally, we show that RHY-1 physically interacts with CYSL-1, a cysteine synthase-like protein and positive regulator of HIF-1, and likely inhibits its function through this interaction. Together, our findings establish RHY-1 as both a regulatory and effector component of the hypoxia response pathway that mediates cellular protection against thiol reductive stress.

Introduction

Redox homeostasis, the finely tuned balance between oxidizing and reducing environments, is essential for cellular survival (Ravi and Singh, 2025). Reactive oxygen species (ROS), continuously generated as byproducts of metabolism, play crucial roles in cell signaling and homeostasis (Sies et al., 2022). However, excessive ROS accumulation leads to oxidative stress, which contributes to various pathological conditions, including neurodegeneration, cardiovascular disease, and cancer (Forman and Zhang, 2021; Schieber and Chandel, 2014). To counter oxidative insults, cells have evolved complex antioxidant defense systems comprising thioredoxin, glutathione (GSH), glutathione peroxidases, catalase, and superoxide dismutases (Kurutas, 2016). In addition to endogenous systems, dietary antioxidants are frequently consumed under both physiological and pathological conditions to mitigate ROS-mediated damage (Kurutas, 2016). Paradoxically, excessive production or intake of antioxidants can disrupt redox equilibrium, leading to reductive stress (Mercola 2025; Zhang et al. 2025). While oxidative stress responses have been extensively characterized, the mechanisms underlying reductive stress remain poorly understood (Ravi and Singh, 2025).

Reductive stress has been implicated in several disorders, including cardiomyopathy, neurodegeneration, and metabolic dysfunction (Mercola 2025; Ravi and Singh 2025; Zhang et al. 2025). The limited understanding of reductive stress responses arises in part from the lack of robust experimental models capable of reliably inducing such conditions (Pan et al., 2024; Yang et al., 2022). Nevertheless, recent studies have begun to elucidate cellular mechanisms that sense and respond to reductive stress. For example, folliculin-interacting protein 1 (FNIP1) has been identified as a redox-sensitive molecule; its cysteine residues directly sense reductive stress, leading to proteasomal degradation of FNIP1 and subsequent increases in mitochondrial activity and ROS generation (Manford et al., 2021, 2020). In parallel, cells undergo extensive metabolic reprogramming under reductive stress to alleviate toxicity (Xiao and Loscalzo, 2020; Yang et al., 2025).

Thiols, characterized by their sulfhydryl groups, constitute a major line of defense against oxidative stress by undergoing reversible oxidation to disulfides (Ulrich and Jakob, 2019). However, excessive exposure to thiol-containing compounds, such as GSH, N-acetylcysteine, or dithiothreitol (DTT), can trigger reductive stress, adversely affecting development and lifespan in *Caenorhabditis elegans* (Gokul and Singh, 2022; Gusarov et al., 2021; Ravi et al., 2023; Winter et al., 2022). DTT, traditionally thought to act primarily

within the endoplasmic reticulum (ER) due to its oxidizing environment (Braakman et al., 1992; Lodish and Kong, 1993; Tatu et al., 1993), has recently been shown to exert broader cellular effects. In addition to inducing ER stress, DTT can paradoxically generate oxidative stress by promoting futile redox cycling in the ER (Maity et al., 2016). Moreover, DTT perturbs the methionine-homocysteine cycle in *C. elegans* by upregulating the S-adenosylmethionine (SAM)-dependent methyltransferase RIPS-1 (Gokul and Singh, 2022; Winter et al., 2022). This induction occurs via the hypoxia-inducible factor HIF-1-mediated hypoxia response pathway (Ravi et al., 2023). Although thiol reductive stress activates the hypoxia response, the physiological significance of this crosstalk remains unclear. It is not known whether activation of the hypoxia response represents a protective adaptation, nor have the downstream effector(s) mediating this protection been identified.

In this study, we demonstrated that the HIF-1-mediated hypoxia response pathway protects *C. elegans* from DTT-induced thiol stress. We further identified an acyltransferase, RHY-1, a transcriptional target of HIF-1, as a key protective factor against thiol stress. Interestingly, RHY-1 also functions as a negative regulator of the hypoxia pathway by inhibiting CYSL-1, a positive regulator of HIF-1 activation. Our data revealed that RHY-1 operates in an autoinhibitory feedback loop, where its overexpression suppresses hypoxia pathway activation even under thiol reductive stress conditions. Finally, we showed that RHY-1 physically interacts with CYSL-1 *in vivo*, likely repressing its function through direct binding. Collectively, our findings establish RHY-1 as both a critical effector and regulator of the hypoxia response pathway during thiol reductive stress.

Results

The hypoxia response pathway protects *C. elegans* from thiol reductive stress

We previously reported that DTT upregulates the SAM-dependent methyltransferase *rips-1* through the HIF-1-mediated hypoxia response pathway (Gokul and Singh, 2022; Ravi et al., 2023). Elevated *rips-1* expression contributes to DTT toxicity, and *rips-1* loss-of-function mutants display resistance to 10 mM DTT (Gokul and Singh, 2022). Since *hif-1* loss-of-function mutants lack induction of *rips-1* expression (Ravi et al., 2023), they were expected to exhibit similar resistance. However, *hif-1(ia4)* mutants were unable to develop on 10 mM DTT (Ravi et al., 2023). This observation suggested that HIF-1 may play a protective role

under thiol reductive stress. Alternatively, the increased sensitivity of *hif-1* mutants could arise from residual basal *rips-1* expression, which is toxic in the presence of DTT.

To test whether basal *rips-1* expression accounts for the heightened DTT sensitivity of *hif-1(ia4)* mutants compared to *rips-1* mutants, we generated a *hif-1(ia4);rips-1(jsn11)* double mutant. The *rips-1(jsn11)* allele carries an early stop codon, resulting in a complete loss of functional RIPS-1 protein (Ravi et al., 2023). Therefore, *hif-1(ia4);rips-1(jsn11)* worms should lack any functional RIPS-1. As expected, *rips-1(jsn11)* single mutants developed normally on 10 mM DTT; however, *hif-1(ia4);rips-1(jsn11)* worms failed to develop (Fig. 1a and b). This result demonstrated that the increased sensitivity of *hif-1* mutants to DTT cannot be attributed to basal *rips-1* expression.

DTT is known to activate the hypoxia response pathway via the upstream regulator CYSL-1 (Fig. 1c). Similar to *hif-1(ia4)* worms, *cysl-1* loss-of-function mutants fail to induce *rips-1* expression and are sensitive to 10 mM DTT (Ravi et al., 2023). Consistently, *cysl-1(jsn17);rips-1(jsn11)* double mutants also failed to develop on 10 mM DTT (Fig.1a and b). Together, these findings indicated that activation of the HIF-1-mediated hypoxia response pathway by DTT confers protection against thiol reductive stress (Fig. 1c).

RHY-1 protects against thiol reductive stress

To further delineate the role of the hypoxia response pathway in defense against thiol reductive stress, we examined the effects of loss-of-function mutations in *rhy-1*, *egl-9*, and *vhl-1*. These genes are key regulators of the hypoxia response pathway and are required for the proteasomal degradation of the transcription factor HIF-1 (Epstein et al., 2001; Shen et al., 2006). Inactivation of any of these genes leads to stabilization of HIF-1 and constitutive activation of hypoxia-responsive genes (Shen et al., 2006). Because *rhy-1*, *egl-9*, and *vhl-1* mutants exhibit elevated *rips-1* expression (Ravi et al., 2023), they are expected to display increased sensitivity to DTT-induced reductive stress. To eliminate confounding effects arising from *rips-1* activity, we generated double mutants, *rhy-1(jsn13jsn14);rips-1(jsn11)*, *egl-9(sa307);rips-1(jsn11)*, and *vhl-1(jsn16);rips-1(jsn11)*, and assessed their sensitivity to 10 mM DTT. We chose this concentration because higher DTT levels can introduce secondary effects related to ER stress (Gokul and Singh, 2022). Unexpectedly, while *rips-1(jsn11)* worms developed normally on 10 mM DTT, *rhy-1(jsn13jsn14);rips-1(jsn11)* worms failed to develop, whereas *egl-9(sa307);rips-1(jsn11)* and *vhl-1(jsn16);rips-1(jsn11)* worms reached

adulthood (Fig. 2a and b). This finding was surprising, as *rhy-1* mutants were anticipated to exhibit enhanced tolerance to DTT due to increased activation of the hypoxia response pathway.

To rule out nonspecific effects from the *rips-1(jsn11)* background, we next evaluated the DTT sensitivity of single mutants *rhy-1(jsn13jsn14)*, *egl-9(jsn15)*, and *vhl-1(jsn16)* across a range of DTT concentrations. As expected, all three mutants were more sensitive to DTT than wild-type N2 worms, consistent with elevated *rips-1* expression (Fig. 2c-g). However, *rhy-1(jsn13jsn14)* worms exhibited significantly greater sensitivity than either *egl-9(jsn15)* or *vhl-1(jsn16)* worms (Fig. 2c-g), suggesting a unique protective role for RHY-1 against DTT toxicity.

DTT toxicity can be alleviated by vitamin B12 supplementation, which restores SAM levels depleted by increased *rips-1* expression (Gokul and Singh, 2022; Winter et al., 2022). We therefore tested whether vitamin B12 supplementation could rescue the DTT sensitivity of mutants with constitutively active hypoxia signaling. While vitamin B12 supplementation effectively rescued DTT-induced toxicity in N2, *egl-9(jsn15)*, and *vhl-1(jsn16)* worms, it failed to ameliorate the effects in *rhy-1(jsn13jsn14)* mutants (Supplementary Fig. 1a and b). Consistently, vitamin B12 also failed to rescue DTT toxicity in an independent *rhy-1* allele, *rhy-1(ok1402)* (Supplementary Fig. 1c and d). Taken together, these results demonstrated that RHY-1 protects from the toxic effects of DTT.

RHY-1 functions downstream of HIF-1 to protect from thiol toxicity

The *rhy-1* gene encodes an acyltransferase that not only regulates HIF-1 activity but is itself a transcriptional target of HIF-1 (Shen et al., 2006). Consistent with this, *rhy-1* expression increases upon DTT exposure in a HIF-1-dependent manner (Ravi et al., 2023). Given that RHY-1 protects against DTT-induced toxicity, we investigated whether the protective role of HIF-1 under thiol reductive stress is mediated through RHY-1. To test this hypothesis, we carried out *rhy-1* rescue experiments. First, we examined whether overexpression of *rhy-1* under its native promoter (*rhy-1p::rhy-1*) could rescue the DTT sensitivity of *rhy-1(jsn13jsn14)* mutants. Indeed, *rhy-1* overexpression restored normal development of *rhy-1(jsn13jsn14)* worms on 10 mM DTT in the presence of 500 nM vitamin B12 (Supplementary Fig. 2a-c).

We next introduced the *rhy-1p::rhy-1* transgene into *hif-1(ia4)* mutants and assessed their development under thiol stress. *hif-1(ia4)* worms overexpressing *rhy-1* exhibited markedly improved development on 10 mM DTT compared to non-transgenic *hif-1(ia4)* worms (Fig. 3a-c). Similarly, supplementation with 500 nM vitamin B12 further enhanced their development relative to non-transgenic controls (Fig. 3a-c). Together, these results demonstrated that *rhy-1* overexpression fully rescues the DTT sensitivity of *hif-1* mutants, indicating that HIF-1 confers protection against thiol reductive stress primarily through the induction of RHY-1.

RHY-1 functions in an autoinhibitory feedback loop to protect from thiol toxicity and physically interacts with CYSL-1

Previous studies have shown that *rhy-1* participates in a negative feedback loop that represses the targets of the hypoxia response pathway (Shen et al., 2006). However, the functional significance of *rhy-1* within this regulatory loop remained unclear. Our findings suggested that RHY-1 not only acts as a negative regulator of the hypoxia response pathway but also serves as an effector protein that mitigates thiol-induced toxicity. To test whether RHY-1 overexpression can repress hypoxia pathway activation even under conditions that normally induce it, we overexpressed *rhy-1p::rhy-1* in *rips-1p::GFP* reporter worms. Adult worms were exposed to varying concentrations of DTT for 12 hours, followed by measurement of green fluorescent protein (GFP) levels. Strikingly, *rhy-1* overexpression completely suppressed the DTT-induced upregulation of *rips-1* (Fig. 4a and b). Collectively, these data indicated that RHY-1 confers protection against DTT toxicity and that elevated RHY-1 levels block activation of the hypoxia response pathway by DTT.

Genetic analyses have previously shown that RHY-1 acts upstream of CYSL-1 and inhibits its function in regulating the hypoxia response pathway under diverse stimuli, including hydrogen sulfide (H_2S) and DTT (Ma et al., 2012; Ravi et al., 2023). CYSL-1, a cysteine synthase-like protein, negatively regulates EGL-9 activity through direct physical interaction (Ma et al., 2012). Given that RHY-1 is a negative regulator of CYSL-1, we hypothesized that RHY-1 may directly interact with CYSL-1 to inhibit its function. To test this possibility, we employed the bimolecular fluorescence complementation (BiFC) assay, a well-established approach for detecting *in vivo* protein-protein interactions in diverse organisms (Hudry et al., 2011; Saka et al., 2007; Shyu et al., 2008). We generated transgenic

C. elegans expressing RHY-1 fused to the N-terminal fragment of Venus (VN173) and CYSL-1 fused to the C-terminal fragment (VC155), both under the control of the *hsp-16.41* promoter. Following heat shock at 33°C for 2 hours and recovery at 20°C for 10 hours, robust Venus fluorescence was observed in worms co-expressing RHY-1::VN173 and CYSL-1::VC155, but not in control worms expressing only vector constructs (Fig. 4c-e). This fluorescence indicated successful Venus reconstitution, providing direct evidence for a physical interaction between RHY-1 and CYSL-1 *in vivo*.

Together, these results suggest that RHY-1 regulates the hypoxia response pathway through a direct physical interaction with CYSL-1. Overall, our findings support a model in which HIF-1 protects against thiol reductive stress via RHY-1, which in turn modulates both DTT toxicity and hypoxia pathway activity through an autoinhibitory feedback loop (Fig. 4f).

Discussion

Our study identifies RHY-1 as a critical downstream effector of the hypoxia response pathway that protects *C. elegans* from DTT-induced reductive stress. Although RHY-1 was previously characterized as an ER membrane protein that negatively regulates HIF-1 under normoxic conditions (Shen et al., 2006), the physiological significance of its HIF-1-dependent upregulation remained unclear. Here, we demonstrate that RHY-1 is essential for survival under thiol reductive stress and fully rescues the DTT sensitivity of *hif-1* loss-of-function mutants. Thus, RHY-1 functions not only as a regulator of the hypoxia response pathway but also as an effector protein that mitigates thiol stress, an inducer of *rhy-1* expression itself through the hypoxia response pathway. This dual regulatory-effector role positions RHY-1 as a key component of an autoinhibitory feedback loop that modulates hypoxia signaling and thiol stress resistance. In the absence of stress, RHY-1 likely inhibits its own production by interacting with and suppressing CYSL-1, thereby preventing unwarranted activation of the hypoxia response pathway (Fig. 4f).

RHY-1 has been implicated in modulating sensitivity to multiple thiol-containing molecules (Horsman et al., 2019; Warnhoff et al., 2024), although its precise function in thiol metabolism appears complex. Unlike its role in DTT sensitivity, *rhy-1* loss-of-function mutants exhibit resistance to H_2S and cysteine (Horsman et al., 2019; Warnhoff et al., 2024). These effects are mediated through HIF-1, suggesting that HIF-1 regulates distinct effector proteins that confer protection against specific thiol stressors. In the case of DTT, however,

RHY-1 overexpression fully restores the DTT sensitivity of *hif-1* mutants, indicating that RHY-1 is likely the principal HIF-1 effector responsible for protection from DTT toxicity. Interestingly, RHY-1 also protects from H_2S toxicity in an HIF-1-independent but SKN-1-dependent manner (Horsman et al., 2019). Increased activity of SKN-1, the *C. elegans* ortholog of mammalian NRF2, can upregulate *rhy-1* expression and compensate for *hif-1* loss under H_2S exposure (Horsman et al., 2019). Furthermore, RHY-1 contributes to cysteine metabolism, as *rhy-1* mutants reduce sulfite production from cysteine and suppress toxicity in a sulfite-sensitized *suox-1(gk738847)* background (Warnhoff et al., 2024). Collectively, these findings suggest that RHY-1 plays multifaceted roles in both the regulation of hypoxia signaling and the metabolism of thiol-containing molecules. Notably, all three thiol-related stressors—H_2S, cysteine, and DTT—are known activators of the hypoxia response pathway (Budde and Roth, 2010; Ravi et al., 2023; Warnhoff et al., 2024).

A key mechanistic question emerging from our study is how RHY-1 confers protection from thiol toxicity. RHY-1 encodes a multipass ER membrane protein containing an acyltransferase-3 domain, which is typically associated with lipid biosynthesis and metabolic regulation (Shen et al., 2006). One possibility is that RHY-1 indirectly mitigates thiol toxicity by altering cellular metabolism, for instance, through acylation of metabolic intermediates such as lipids or carbohydrates. Another possibility is that RHY-1 directly influences thiol metabolism. While S-acetylation of small-molecule thiols occurs in cells (Di Paola et al., 2022; Filipovic et al., 2018; Locigno et al., 2002), RHY-1 is unlikely to catalyze this reaction directly because it is predicted to possess O-acyltransferase rather than S-acetyltransferase activity. Nonetheless, *rhy-1* mutants display defects in cysteine catabolism and impaired sulfite production (Warnhoff et al., 2024), implying a direct metabolic role for RHY-1 in thiol processing.

In summary, RHY-1 emerges as a multifunctional regulator that integrates hypoxia signaling with thiol metabolism. By functioning in a negative feedback loop downstream of HIF-1, RHY-1 fine-tunes the hypoxia response while simultaneously providing protection from reductive stress. Future studies dissecting the biochemical activities of RHY-1 and its substrate specificity will be crucial for understanding how acyltransferases contribute to cellular adaptation under reductive stress conditions.

Materials and methods

Bacterial strains

Escherichia coli OP50 was used as the food source for *Caenorhabditis elegans* throughout this study. *E. coli* OP50 cultures were grown in Luria-Bertani broth at 37°C with shaking overnight.

C. elegans strains and growth conditions

C. elegans hermaphrodites were maintained at 20°C on nematode growth medium (NGM) plates seeded with *E. coli* OP50 unless otherwise indicated. Bristol N2 was used as the wild-type control unless otherwise indicated. The following strains were used in the study: JSJ11 *rips-1(jsn11)*, JSJ13 *jsnIs1[rips-1p::GFP + myo-2p::mCherry]*, JSJ14 *jsnIs1[rips-1p::GFP + myo-2p::mCherry];rhy-1(jsn13jsn14)*, JSJ15 *jsnIs1[rips-1p::GFP + myo-2p::mCherry];egl-9(jsn15)*, JSJ16 *jsnIs1[rips-1p::GFP + myo-2p::mCherry];vhl-1(jsn16)*, JSJ17 *jsnIs1[rips-1p::GFP + myo-2p::mCherry];cysl-1(jsn17)*, ZG31 *hif-1(ia4)*, JT307 *egl-9(sa307),* and RB1297 *rhy-1(ok1408)*. The following strains were generated by standard genetic crosses: *hif-1(ia4);rips-1(jsn11)*, *egl-9(sa307);rips-1(jsn11)*, *rhy-1(jsn13jsn14);rips-1(jsn11)*, *vhl-1(jsn16);rips-1(jsn11)*, and *rips-1(jsn11);cysl-1(jsn17)*.

Generation of transgenic *C. elegans*

To overexpress *rhy-1*, the genomic region containing the *rhy-1* coding sequence and 995 bp of its upstream promoter was amplified from N2 genomic DNA. The fragment, including the stop codon, was cloned into the pPD95_77 plasmid using SalI and KpnI restriction sites. The primer sequences are listed in Table S1. The resulting construct (*rhy-1p::rhy-1*) was co-injected into N2 worms at 50 ng/µL, along with pCFJ104 (*myo-3p::mCherry*) as a co-injection marker (25 ng/µL). Transgenic worms carrying the extrachromosomal array *jsnEx5[rhy-1p::rhy-1 + myo-3p::mCherry]* were subsequently crossed into *hif-1(ia4)*, *rhy-1(jsn13jsn14)*, and *rips-1p::GFP* backgrounds to generate *hif-1(ia4);jsnEx5[rhy-1p::rhy-1 + myo-3p::mCherry]*, *rhy-1(jsn13jsn14);jsnEx5[rhy-1p::rhy-1 + myo-3p::mCherry]*, and *rips-1p::GFP;jsnEx5[rhy-1p::rhy-1 + myo-3p::mCherry]*.

Supplementation experiments

The following supplements were obtained from HiMedia BioSciences: DTT (#RM525) and vitamin B12 (cyanocobalamin) (#PCT0204). Stock solutions were prepared in water, with DTT stocks freshly prepared prior to each experiment and vitamin B12 stocks stored at 4°C. Stocks were diluted to the desired final concentrations in NGM prior to pouring the plates. NGM plates were seeded with *E. coli* OP50 and incubated at room temperature for at least three days before use.

Development assays

To obtain synchronized *C. elegans* populations, 20-25 gravid adult hermaphrodites were transferred to NGM plates containing the indicated supplements and allowed to lay eggs for 2 hours. The adults were then removed, and the plates were incubated at 20°C for 72 hours. Worms were examined and scored at different developmental stages (L1/L2, L3, and L4/adult). Representative images were captured during assessment. Each condition was tested in at least three independent biological replicates.

Bimolecular fluorescence complementation (BiFC) assay

For BiFC analysis, *rhy-1* and *cysl-1* genomic sequences were cloned downstream of the *hsp-16.41* promoter into plasmids pCE-BiFC-VN173 and pCE-BiFC-VC155 (Addgene), respectively. The *rhy-1* coding sequence was inserted in-frame between the SmaI and KpnI restriction sites, whereas the *cysl-1* coding sequence was inserted in-frame between the NheI and KpnI sites. The primer sequences are listed in Table S1. The BiFC plasmid constructs were microinjected into N2 worms at a final concentration of 10 ng/µL, along with *myo-3p::mCherry* (25 ng/µL) as a co-injection marker. Control worms were injected with empty pCE-BiFC-VN173 and pCE-BiFC-VC155 vectors together with the same marker. For the BiFC assay, transgenic worms were synchronized by egg laying and incubated at 20°C for 82 hours post egg laying. The worms were then subjected to heat shock at 33°C for 2 hours, followed by recovery at 20°C for 10 hours. Fluorescence signals were subsequently visualized using a fluorescence stereomicroscope.

Fluorescence imaging and analysis

The fluorescence reporter strains were picked under a non-fluorescence stereomicroscope to minimize potential bias. Fluorescence imaging was carried out as described previously (Das et al., 2025; Ghosh and Singh, 2024). Briefly, worms were anesthetized in 50 mM sodium azide in M9 buffer and mounted on 2% agarose pads. Fluorescence was visualized using a Nikon SMZ18 stereomicroscope, and images were analyzed using ImageJ software. Quantitative fluorescence intensity data were plotted using GraphPad Prism 8.

Statistical analysis

Statistical analysis was performed using GraphPad Prism 8. Error bars represent mean ± standard deviation (SD). An unpaired, two-tailed, two-sample t-test was used when applicable, and the data were considered statistically significant when $P < 0.05$. In the figures, asterisks (*) denote statistical significance as follows: * for $P < 0.05$, **for $P < 0.01$, *** for $P < 0.001$, as compared with the appropriate controls. All experiments were performed in triplicate.

Data availability

All data generated or analyzed during this study are included in the manuscript and supporting files.

Acknowledgments

We thank the Caenorhabditis Genetics Center (funded by the NIH Office of Research Infrastructure Programs (P40 OD010440)) for providing the strains used in this study.

Funding

This work was supported by the following grants: Ramalingaswami Re-entry Fellowship (Ref. No. BT/RLF/Re-entry/50/2020) and Har-Gobind Khorana-Innovative Young Biotechnologist Fellowship (File No. HRD-17011/2/2023-HRD-DBT) awarded by the Department of Biotechnology, India; Anusandhan National Research Foundation (ANRF)

Core Research Grant (Ref. No. CRG/2023/001136) awarded by DST, India; STARS grant (File No. MoE-STARS/STARS-2/2023-0116) awarded by the Ministry of Education, India; Research Grant (Ref. No. 37/1741/23/EMR-II) awarded by the Council of Scientific & Industrial Research (CSIR), India; and IISER Mohali intramural funds.

Declaration of interests

The authors declare no competing interests.

Reference

Braakman I, Helenius J, Helenius A. 1992. Manipulating disulfide bond formation and protein folding in the endoplasmic reticulum. *EMBO J* **11**:1717–1722. doi:10.1002/j.1460-2075.1992.tb05223.x

Budde MW, Roth MB. 2010. Hydrogen Sulfide Increases Hypoxia-inducible Factor-1 Activity Independently of von Hippel–Lindau Tumor Suppressor-1 in C. elegans. *Mol Biol Cell* **21**:212–217. doi:10.1091/mbc.e09-03-0199

Das P, Ravi, Singh J. 2025. Iron-deplete diet enhances Caenorhabditis elegans lifespan via oxidative stress response pathways. doi:10.1101/2025.02.11.637611

Di Paola R, Modafferi S, Siracusa R, Cordaro M, D'Amico R, Ontario ML, Interdonato L, Salinaro AT, Fusco R, Impellizzeri D, Calabrese V, Cuzzocrea S. 2022. S-Acetyl-Glutathione Attenuates Carbon Tetrachloride-Induced Liver Injury by Modulating Oxidative Imbalance and Inflammation. *Int J Mol Sci* **23**:4429. doi:10.3390/ijms23084429

Epstein ACR, Gleadle JM, McNeill LA, Hewitson KS, O'Rourke J, Mole DR, Mukherji M, Metzen E, Wilson MI, Dhanda A, Tian Y-M, Masson N, Hamilton DL, Jaakkola P, Barstead R, Hodgkin J, Maxwell PH, Pugh CW, Schofield CJ, Ratcliffe PJ. 2001. *C. elegans* EGL-9 and Mammalian Homologs Define a Family of Dioxygenases that Regulate HIF by Prolyl Hydroxylation. *Cell* **107**:43–54. doi:10.1016/S0092-8674(01)00507-4

Filipovic MR, Zivanovic J, Alvarez B, Banerjee R. 2018. Chemical Biology of H2S Signaling through Persulfidation. *Chem Rev* **118**:1253–1337. doi:10.1021/acs.chemrev.7b00205

Forman HJ, Zhang H. 2021. Targeting oxidative stress in disease: promise and limitations of antioxidant therapy. *Nat Rev Drug Discov* **20**:689–709. doi:10.1038/s41573-021-00233-1

Ghosh A, Singh J. 2024. Translation initiation or elongation inhibition triggers contrasting effects on Caenorhabditis elegans survival during pathogen infection. *mBio* **15**:e02485-24. doi:10.1128/mbio.02485-24

Gokul G, Singh J. 2022. Dithiothreitol causes toxicity in C. elegans by modulating the methionine–homocysteine cycle. *eLife* **11**:e76021. doi:10.7554/eLife.76021

Gusarov I, Shamovsky I, Pani B, Gautier L, Eremina S, Katkova-Zhukotskaya O, Mironov A, Makarov AA, Nudler E. 2021. Dietary thiols accelerate aging of C. elegans. *Nat Commun* **12**:4336. doi:10.1038/s41467-021-24634-3

Horsman JW, Heinis FI, Miller DL. 2019. A Novel Mechanism To Prevent H2S Toxicity in Caenorhabditis elegans. *Genetics* **213**:481–490. doi:10.1534/genetics.119.302326

Hudry B, Viala S, Graba Y, Merabet S. 2011. Visualization of protein interactions in living Drosophila embryos by the bimolecular fluorescence complementation assay. *BMC Biol* **9**:5. doi:10.1186/1741-7007-9-5

Kurutas EB. 2016. The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: current state. *Nutr J* **15**:71. doi:10.1186/s12937-016-0186-5

Locigno R, Pincemail J, Henno A, Treusch G, Castronovo V. 2002. S-acetyl-glutathione selectively induces apoptosis in human lymphoma cells through a GSH-independent mechanism. *Int J Oncol* **20**:69–75.

Lodish HF, Kong N. 1993. The secretory pathway is normal in dithiothreitol-treated cells, but disulfide-bonded proteins are reduced and reversibly retained in the endoplasmic reticulum. *J Biol Chem* **268**:20598–20605. doi:10.1016/S0021-9258(20)80767-4

Ma DK, Vozdek R, Bhatla N, Horvitz HR. 2012. CYSL-1 Interacts with the O2-sensing Hydroxylase EGL-9 to Promote H2S-modulated Hypoxia-induced Behavioral Plasticity in C. elegans. *Neuron* **73**:925–940. doi:10.1016/j.neuron.2011.12.037

Maity S, Rajkumar A, Matai L, Bhat A, Ghosh A, Agam G, Kaur S, Bhatt NR, Mukhopadhyay A, Sengupta S, Chakraborty K. 2016. Oxidative Homeostasis Regulates the Response to Reductive Endoplasmic Reticulum Stress through Translation Control. *Cell Rep* **16**:851–865. doi:10.1016/j.celrep.2016.06.025

Manford AG, Mena EL, Shih KY, Gee CL, McMinimy R, Martínez-González B, Sherriff R, Lew B, Zoltek M, Rodríguez-Pérez F, Woldesenbet M, Kuriyan J, Rape M. 2021. Structural basis and regulation of the reductive stress response. *Cell* **184**:5375-5390.e16. doi:10.1016/j.cell.2021.09.002

Manford AG, Rodríguez-Pérez F, Shih KY, Shi Z, Berdan CA, Choe M, Titov DV, Nomura DK, Rape M. 2020. A Cellular Mechanism to Detect and Alleviate Reductive Stress. *Cell* **183**:46-61.e21. doi:10.1016/j.cell.2020.08.034

Mercola J. 2025. Reductive stress and mitochondrial dysfunction: The hidden link in chronic disease. *Free Radic Biol Med* **233**:118–131. doi:10.1016/j.freeradbiomed.2025.03.029

Pan X, Heacock ML, Abdulaziz EN, Violante S, Zuckerman AL, Shrestha N, Yao C, Goodman RP, Cross JR, Cracan V. 2024. A genetically encoded tool to increase cellular NADH/NAD+ ratio in living cells. *Nat Chem Biol* **20**:594–604. doi:10.1038/s41589-023-01460-w

Ravi, Kumar A, Bhattacharyya S, Singh J. 2023. Thiol reductive stress activates the hypoxia response pathway. *EMBO J* **42**:e114093. doi:10.15252/embj.2023114093

Ravi, Singh J. 2025. Redox imbalance and hypoxia-inducible factors: a multifaceted crosstalk. *FEBS J* **292**:3833–3848. doi:10.1111/febs.70013

Saka Y, Hagemann AI, Piepenburg O, Smith JC. 2007. Nuclear accumulation of Smad complexes occurs only after the midblastula transition in Xenopus. *Development* **134**:4209–4218. doi:10.1242/dev.010645

Schieber M, Chandel NS. 2014. ROS function in redox signaling and oxidative stress. *Curr Biol CB* **24**:R453-462. doi:10.1016/j.cub.2014.03.034

Shen C, Shao Z, Powell-Coffman JA. 2006. The *Caenorhabditis elegans rhy-1* Gene Inhibits HIF-1 Hypoxia-Inducible Factor Activity in a Negative Feedback Loop That Does Not Include *vhl-1*. *Genetics* **174**:1205–1214. doi:10.1534/genetics.106.063594

Shyu YJ, Hiatt SM, Duren HM, Ellis RE, Kerppola TK, Hu C-D. 2008. Visualization of protein interactions in living Caenorhabditis elegans using bimolecular fluorescence complementation analysis. *Nat Protoc* **3**:588–596. doi:10.1038/nprot.2008.16

Sies H, Belousov VV, Chandel NS, Davies MJ, Jones DP, Mann GE, Murphy MP, Yamamoto M, Winterbourn C. 2022. Defining roles of specific reactive oxygen species (ROS) in cell biology and physiology. *Nat Rev Mol Cell Biol* **23**:499–515. doi:10.1038/s41580-022-00456-z

Tatu U, Braakman I, Helenius A. 1993. Membrane glycoprotein folding, oligomerization and intracellular transport: effects of dithiothreitol in living cells. *EMBO J* **12**:2151–2157. doi:10.1002/j.1460-2075.1993.tb05863.x

Ulrich K, Jakob U. 2019. The role of thiols in antioxidant systems. *Free Radic Biol Med*, Early Life on Earth and Oxidative Stress **140**:14–27. doi:10.1016/j.freeradbiomed.2019.05.035

Warnhoff K, Bhattacharya S, Snoozy J, Breen PC, Ruvkun G. 2024. Hypoxia-inducible factor induces cysteine dioxygenase and promotes cysteine homeostasis in Caenorhabditis elegans. *eLife* **12**:RP89173. doi:10.7554/eLife.89173

Winter AD, Tjahjono E, Beltrán LJ, Johnstone IL, Bulleid NJ, Page AP. 2022. Dietary-derived vitamin B12 protects Caenorhabditis elegans from thiol-reducing agents. *BMC Biol* **20**:228. doi:10.1186/s12915-022-01415-y

Xiao W, Loscalzo J. 2020. Metabolic Responses to Reductive Stress. *Antioxid Redox Signal* **32**:1330–1347. doi:10.1089/ars.2019.7803

Yang R, Guo Z, Li B. 2025. NADH reductive stress drives metabolic reprogramming. *Trends Cell Biol* **0**. doi:10.1016/j.tcb.2025.07.005

Yang R, Yang C, Ma L, Zhao Y, Guo Z, Niu J, Chu Q, Ma Y, Li B. 2022. Identification of purine biosynthesis as an NADH-sensing pathway to mediate energy stress. *Nat Commun* **13**:7031. doi:10.1038/s41467-022-34850-0

Zhang Shiyi, Wang N, Gao Z, Gao J, Wang X, Xie H, Wang C-Y, Zhang Shu. 2025. Reductive stress: The key pathway in metabolic disorders induced by overnutrition. *J Adv Res*. doi:10.1016/j.jare.2025.01.012

Figures

Figure 1. Hypoxia response pathway protects *C. elegans* from thiol reductive stress

(a) Representative images of N2, *rips-1(jsn11)*, *hif-1(ia4)*, *hif-1(ia4);rips-1(jsn11)*, *cysl-1(jsn17)*, and *cysl-1(jsn17);rips-1(jsn11)* worms on 0 mM and 10 mM DTT on *E. coli* OP50 diet after 72 h of egg laying at 20°C. Scale bar = 200 μm.

(b) Quantification of different developmental stages of N2, *rips-1(jsn11)*, *hif-1(ia4)*, *hif-1(ia4);rips-1(jsn11)*, *cysl-1(jsn17)*, and *cysl-1(jsn17);rips-1(jsn11)* worms on 0 mM and 10 mM DTT on *E. coli* OP50 diet after 72 h of egg laying at 20°C (*n* = 3 biological replicates; worms per condition per replicate > 70).

(c) Schematic representation of the HIF-1-mediated hypoxia response pathway and its role in modulating thiol toxicity.

Figure 2. RHY-1 protects against thiol reductive stress

(a) Representative images of *rips-1(jsn11)*, *rhy-1(jsn13jsn14);rips-1(jsn11)*, *egl-9(sa307);rips-1(jsn11)*, and *vhl-1(jsn16);rips1(jsn11)* worms on 0 mM and 10 mM DTT on *E. coli* OP50 diet after 72 h of egg laying at 20°C. Scale bar = 200 µm.

(b) Quantification of different developmental stages of *rips-1(jsn11)*, *rhy-1(jsn13jsn14);rips-1(jsn11)*, *egl-9(sa307);rips-1(jsn11)*, and *vhl-1(jsn16);rips1(jsn11)* worms on 0 mM and 10 mM DTT on *E. coli* OP50 diet after 72 h of egg laying at 20°C (n = 3 biological replicates; worms per condition per replicate > 75).

(c) Representative images of N2, JSJ14 (*rhy-1(jsn13jsn14)*), JSJ15 (*egl-9(jsn15)*), and JSJ16 (*vhl-1(jsn16)*) worms on 0 mM, 2 mM, 4 mM, and 6 mM DTT on *E. coli* OP50 diet after 72 hours of egg laying. Scale bar = 200 μm.

(d)-(g) Quantification of different developmental stages of N2, JSJ14 (*rhy-1(jsn13jsn14)*), JSJ15 (*egl-9(jsn15)*), and JSJ16 (*vhl-1(jsn16)*) worms on 0 mM (d), 2 mM (e), 4 mM (f), and 6 mM (g) DTT on *E. coli* OP50 diet after 72 hours of egg laying ($n = 3$ biological replicates; worms per condition per replicate > 70).

Figure 3. RHY-1 functions downstream of HIF-1 to protect from thiol toxicity

(a) Representative images of *hif-1(ia4)* and *hif-1(ia4);rhy-1p::rhy-1* worms exposed to 0 mM DTT, 10 mM DTT, or 10 mM DTT supplemented with 500 nM vitamin B12 on *E. coli* OP50 for 72 h at 20°C. Scale bar = 200 μm.

(b)-(c) Quantification of different developmental stages of *hif-1(ia4)* (b) and *hif-1(ia4);rhy-1p::rhy-1* (c) worms exposed to 0 mM DTT, 10 mM DTT, or 10 mM DTT supplemented with 500 nM vitamin B12 on *E. coli* OP50 for 72 h at 20°C (n = 3 biological replicates; worms per condition per replicate > 65).

Figure 4. RHY-1 functions in an autoinhibitory feedback loop to protect from thiol toxicity and physically interacts with CYSL-1

(a) Representative fluorescence images of *rips-1p::GFP* and *rips-1p::GFP;rhy-1p::rhy-1* worms after exposure of synchronized gravid adults to 0 mM, 5 mM, 7.5 mM, and 10 mM DTT for 12 hours. Scale bar = 200 μm.

(b) Quantification of fluorescence levels of *rips-1p::GFP* and *rips-1p::GFP;rhy-1p::rhy-1* worms after exposure of synchronized gravid adults to 0 mM, 5 mM, 7.5 mM, and 10 mM

DTT for 12 hours. ***$P < 0.001$, **$P < 0.01$, and *$P < 0.05$ via the *t*-test. n.s., nonsignificant ($n = 20$-21 worms each).

(c) Representative fluorescence images of worms expressing the indicated BiFC plasmid combinations with or without a 2-hour heat shock. Scale bar $= 200$ μm.

(d) Quantification of fluorescence intensity in worms expressing empty pCE-BiFC-VN173 and pCE-BiFC-VC155 vectors with or without heat shock. ns, nonsignificant via the *t*-test ($n = 35$ worms each).

(e) Quantification of fluorescence intensity in worms expressing RHY-1::VN173 and CYSL-1::VC155 with or without heat shock. ***$P < 0.001$ via the *t*-test ($n = 35$ worms each).

(f) Model for the autoinhibitory feedback loop of the hypoxia response pathway that protects from thiol toxicity.

Supplementary Figures

Supplementary Figure 1. RHY-1 protects from thiol toxicity

(a) Representative images of N2, JSJ14 (*rhy-1(jsn13jsn14)*), JSJ15 (*egl-9(jsn15)*), and JSJ16 (*vhl-1(jsn16)*) worms exposed to 0 mM DTT or 10 mM DTT supplemented with 500 nM vitamin B12 on *E. coli* OP50 for 72 h at 20°C. Scale bar = 200 μm.

(b) Quantification of different developmental stages of N2, JSJ14 (*rhy-1(jsn13jsn14)*), JSJ15 (*egl-9(jsn15)*), and JSJ16 (*vhl-1(jsn16)*) worms exposed to 0 mM DTT or 10 mM DTT supplemented with 500 nM vitamin B12 on *E. coli* OP50 for 72 h at 20°C ($n = 3$ biological replicates; worms per condition per replicate > 80).

(c) Representative images of N2 and *rhy-1(ok1402)* worms exposed to 0 mM DTT, 10 mM DTT, or 10 mM DTT supplemented with 500 nM vitamin B12 on *E. coli* OP50 for 72 h at 20°C. Scale bar = 200 μm.

(d) Quantification of different developmental stages of N2 and *rhy-1(ok1402)* worms exposed to 0 mM DTT, 10 mM DTT, or 10 mM DTT supplemented with 500 nM vitamin B12 on *E. coli* OP50 for 72 h at 20°C ($n = 3$ biological replicates; worms per condition per replicate > 55).

Supplementary Figure 2. RHY-1 protects from thiol toxicity

(a) Representative images of *rhy-1(jsn13jsn14)* and *rhy-1(jsn13jsn14);rhy-1p::rhy-1* worms exposed to 0 mM DTT, 10 mM DTT, or 10 mM DTT supplemented with 500 nM vitamin B12 on *E. coli* OP50 for 72 h at 20°C. Scale bar = 200 μm.

(b)-(c) Quantification of different developmental stages of *rhy-1(jsn13jsn14)* (b) and *rhy-1(jsn13jsn14);rhy-1p::rhy-1* (c) worms exposed to 0 mM DTT, 10 mM DTT, or 10 mM DTT supplemented with 500 nM vitamin B12 on *E. coli* OP50 for 72 h at 20°C (n = 3 biological replicates; worms per condition per replicate > 35).

Table S1: Primers used in the study

Cloning primers	
Primer name	Primer sequence (5'-3')
rhy-1p::rhy-1 _Sal1_F	ACCGTCGACACCGCATTGTCACACCATCT
*rhy-1p::rhy-1*_Kpn1_R	ACCGGTACCTTCCAGGTGAGGATCCGTTC
*cysl-1*_F_Nhe1_VC155	GACGCTAGCATGGCTGATCGCAACTCTAT
*cysl-1*_R_Kpn1_VC155	GCAGGTACCTCCATAATATCATAGTTGTGAAGAC
*rhy-1*_F_Sma1_VN173	GACCCCGGGATGTCATCATCGCCACACACTCAC
*rhy-1*_R_Kpn1_VN173	GCAGGTACCCAGGCCCAGTCGTGTGGCATATT

Genotyping primers	
Primer name	Primer sequence (5'-3')
*cysl-1(jsn17)*_F	GGACTTCCAGGGACAGTTGG
*cysl-1(jsn17)*_R	TCAATGAGCGTGGTCACTCC
*rips-1(jsn11)*_F	TCGCCGTCAGCAATGTAATCT
*rips-1(jsn11)*_R	TTCACTTGCCCCGCTGTTAT
*rhy-1(jsn13jsn14)*_F	AGCCTCAAGCGATCAGTCAA
*rhy-1(jsn13jsn14)*_R	CCCAAAAACCACAAGCGGAG
*vhl-1(jsn16)*_F	AGAAGTATTCTGGCCGGAGC
*vhl-1(jsn16)*_F	CATTGCTGAGGTCTCTGGGG
*rhy-1(ok1402)*_F1	TTGGAGCTCTCATGGGAAAC
*rhy-1(ok1402)*_F2	ATTCCGATCTGTATTCGCTA
*rhy-1(ok1402)*_R1	AGCTTGAAGGAATGGGAGCC
*egl-9(sa307)*_F1	GACAGGTGTTGCGAATGAGC
*egl-9(sa307)*_F2	CGTGGTCTCGCACTAAGCAT
*egl-9(sa307)*_R1	AGTCACTCACTTCGCGGTAA
*hif-1(ia4)*_F1	TTGAGGAATGCCGCATGTTC
*hif-1(ia4)*_F2	TTTGAGCAAAGCACACTCAGC
*hif-1(ia4)*_R1	ACGCCCTTTAGGACTCACTTT