Title

Age Associated Increase in Microglia Inflammation and Phagocytosis in the Adult Neural

Stem Cell Niche.

Authors

Ronald Cutler[1,=], Samuel Harrison[1,=], Natalia Sandoval-Kuhn[1], Stephen Tomlinson[2], Erzsebet

Kokovay[1]

Affiliations

[1] UTH-SA, Department of Cell Systems and Anatomy, San Antonio, TX, USA

[2] Medical University of South Carolina, Department of Pharmacology and Immunology,

Charleston, SC, USA

[=] Authors contributed equally to this work

[*] Correspondence to E.K. (KokovayE@uthscsa.edu)

Summary

Neurogenesis occurs throughout life but is reduced aging. We show that during aging, microglia

in the Subventricular Zone become activated and display an inflammatory phenotype. We show

that young SVZ microglia have reduced phagocytic capacity compared to whole brain microglia

in vitro, suggesting that these niche microglia are specialized not to phagocytize stem cells and

progenitors. However, during aging, SVZ microglia exhibit increased phagocytosis. We show that

during aging, there is a significant increase in the number of microglia that have NSC and

progenitor markers within the lysosomes, strongly suggesting that during aging microglia engulf

NSCs and progenitors. Reducing inflammation improves proliferation but had no significant

reduction on phagocytosis suggesting that inflammation is not a key driver in the age-associated

phagocytosis but is important in reduced proliferation. This indicates that SVZ microglia are key

regulators of neurogenesis in the aged SVZ.

Introduction

Adult neural stem cells (NSCs) reside in two neurogenic niches of the murine brain named the subgranular zone (SGZ) of the dentate gyrus and the subventricular zone (SVZ). These NSCs give rise to neurons, astrocytes, and oligodendrocytes. Neurogenesis, the production of new neurons, occurs throughout life in the SVZ and SGZ niches. Here, we are focused on the NSCs in the SVZ niche, which during normal brain homeostasis, are in a quiescent state (qNSCs) or actively dividing state (aNSCs). aNSCs produce a progenitor cell called transit amplifying cells (TACs), which proliferate multiple times to eventually yield neuroblasts. These neuroblasts migrate from the SVZ through the rostral migratory stream (RMS) and into the olfactory bulb (OB), where they mature and integrate into the neuronal circuitry of the OB (Alvarez-Buylla and García-Verdugo, 2002). During aging, there is a drastic decline in the proliferation of NSCs, increased NSC quiescence, and reduced NSC number (Ahlenius et al., 2009; Enwere et al., 2004; Kalamakis et al., 2019; Maslov et al., 2004; Solano Fonseca et al., 2016).

Microglia, the innate immune cells of the brain, function as monocytes and extend processes for surveillance of the local tissue. Microglia are myeloid progenitors that originate in the yolk sac and infiltrate the CNS during development, which comprise around 10% of the cell population of the adult brain (Ginhoux et al., 2010). Microglia are versatile cells that work to clear cellular debris or dying cells via phagocytosis, produce signaling molecules to maintain brain homeostasis, maintain neuronal synapses by synaptic pruning, and help to repair tissue following injury (Colonna and Butovsky, 2017). It has been observed that microglia that reside in the SVZ niche have distinct properties from other brain regions in morphology, ability to support NSC proliferation and neurogenesis (Goings et al., 2006; Marshall et al., 2014; Ribeiro Xavier et al., 2015; Shigemoto-Mogami et al., 2014; Solano Fonseca et al., 2016). However, we have previously

observed that during aging SVZ microglia become more activated and proinflammatory. In fact,

age-related increase in SVZ microglial activation precedes the decline of NSC proliferation,

suggesting that microglia activation may be a driver of NSC dysfunction (Solano Fonseca et al.,

2016). Here, we found that during aging, SVZ microglia not only display significant upregulation

of pro-inflammatory signatures, but also show increased lysosomal marker and morphology of

activatation. We show that SVZ microglia become more phagocytic during aging i*n vitro* and *in*

vivo, where they engulf aNSCs and TACs and express more inflammation associated genes.

Finally, we performed interventions to reduce inflammation and saw a partial rescue of both

proliferation and TAC number, however this was not accompanied by a reduction in TAC

phagocytosis, suggesting that microglia inflammation contributes to reduced NSC proliferation

but is not a driver of NSC phagocytosis during aging.

Age-related changes in SVZ microglia morphology

We have previously shown that SVZ neurogenesis is reduced by 12 months of age (Solano

et al., 2016). To understand how SVZ microglia change during aging, we performed

immunohistochemistry for the microglia marker Iba1 and the lysosomal marker CD68 in the SVZs

dissected as wholemounts to preserve the 3-D cytoarchitecture of the niche, from 3- and 12-month-

old mice. In young adult mice (3-month-old), we observed ramified cell morphology throughout

the SVZ indicated by Iba1+ signal with relatively low amounts of CD68+ signal within Iba1+ cells

(Fig. 1A). In aged mice (12-month-old), we observed a more activated cell morphology with fewer

processes, larger cell bodies, and higher amounts of Cd68+ signal within Iba1+ microglia in the

SVZ (Fig. 1C). During neuronal development, it has been shown that microglia adopt a phagocytic

phenotype, reminiscent of the phenotype seen during SVZ aging, and was associated with

phagocytosis of precursor cells during development. This led us to hypothesize that activated

microglia in the aging SVZ may also have increased phagocytotic activity.

Figure 1. Age-Associated SVZ Microglia Morphological Changes and Increased Activation.
Representative confocal projection images of SVZ microglia stained with Iba1 (*green*) and
microglia lysosomes stained with CD68 (*red*). (A) Young (3-month) SVZ microglia imaged at 40x
magnification. Scale bar: 50µm. (B) Young SVZ microglia imaged at 100x magnification. Scale
bar: 20µm. (C) Aged (12-month) SVZ microglia imaged at 40x magnification. Scale bar: 50µm.
(D) 12-month SVZ microglia imaged at 100x magnification. Scale bar: 20µm.

Gene expression in the aging SVZ suggest increased phagocytosis of NSCs by microglia

To determine if there are age-related transcriptional changes in SVZ microglia as well as NSCs related to phagocytosis and inflammation, we reanalyzed a published data set that profiled 14,685 single-cell transcriptomes from the SVZ of young (3-month-old) and old (28/29-month-old) male mice. With this, we created pseudobulk samples for each cell type and biological replicate and performed differential gene expression (DGE) analysis to compare old vs. young, followed by gene set enrichment analysis (GSEA) (Supplemental Fig. 1). We first focused on the age-related changes in the microglia, where we found 254 and 54 genes significantly (adjusted p-value < 0.05) up- and down-regulated, respectively (Supplemental Fig. 1C). GSEA of the upregulated genes revealed processes involved in translation and secretion of inflammatory cytokines, and lipid metabolism, which has been previously reported (Yen et al., 2024). GSEA of downregulated genes revealed processes involved in cytoskeleton organization, which has not been reported before and may suggest dysfunction in migration (Supplemental Fig. 1D).

Notably, we did not find enrichments of phagocytosis from GSEA, however, we did find genes that act as receptors for phagocytosis, such as *Axl* and *Tlr2*, to be amongst the genes with the highest increase in gene expression during microglia aging (Supplemental Fig. 1C). Thus, we curated comprehensive gene sets related to several aspects of phagocytosis from the molecular signatures database (Liberzon et al., 2024) and overlapped these with the set of age-related differentially expressed genes (ARDEGs; adjusted p-value < 0.05). Out of 19 genes related to phagocytosis that overlapped with the microglia ARDEG set, 18 were found to be significantly upregulated in old microglia as compared to the young (Fig. 2A). Notably, 8 of these genes (*Cyba, Atp6v0e, Rab7b, Cybb, Vim, B2m, Tlr2,* and *Hvcn1*) were associated with the formation of phagocytic vesicles. We extended our phagocytosis gene set to include genes identified through

screens from previous studies (Haney et al., 2018), (Pluvinage et al., 2019), and found that all 9

genes that overlapped with the microglia ARDEGs were significantly upregulated in old microglia

as compared to the young (Supplementary Figs. 2A,B). However, it is possible that the

upregulation of these genes may be part of a compensatory response to the increase in phagocytic

activity. In the same vein, we found that the expression of the receptor essential for microglial

function, *Csf1r*, was downregulated in aged microglia while the ligand *Csf1* showed the opposite

trend and thus may be a compensatory response to chronic microglia activation (Hu et al., 2024)

(Supplementary Fig. 2C). Finally, we found that the broad upregulation of phagocytosis-related

gene expression was also accompanied by an increase in the expression of genes related to

microglia activation (Supplementary Figs. 2C-D). Overall, these findings suggest that aging causes

complex gene expression reprogramming which shifts the expression profiles of aged microglia to

resemble those in the states of phagocytosis, activation, and development (Izgi et al., 2022). This

is further supported by the phenotypic resemblance of aged SVZ microglia to microglia in the

developmental context in terms of an activated morphology and increased phagocytosis of neural

progenitors (Cunningham et al., 2013).

We next asked if the upregulation of phagocytosis-associated gene expression was specific

to any of the known cell surface receptors pathways required for this process. By using a curated

list of the components of each receptor pathway known to be associated with phagocytosis, we

found gene expression upregulation of the receptors in the Complement, TAM, and Toll-like

pathways. On the other hand, for the FC gamma, Scavenger, P2ry6, and Trem2 receptor pathways,

we found largely consistent upregulation of gene expression for the components of these pathways,

which did not coincide with upregulation of the core receptor (Deczkowska et al., 2018).Of note,

while we did not detect increased expression of the receptor Trem2 in aged microglia, which is

known to promote anti-inflammatory responses and phagocytosis, we did find an expression

pattern in aged microglia consistent with the disease-activated microglia (DAM) stage 2 signature

that has been reported to be Trem2-dependent (Supplementary Figs. 2E). Overall, these results

suggest that during the aging of microglia, there is a complex upregulation of expression for genes

across multiple phagocytosis pathways that corresponds with a unique Trem2-independent DAM

state

We then turned our attention to analyzing the transcriptomes of the neural progenitor cells

where we asked if we could detect any age-related expression changes in genes related to target

cell engulfment, such as apoptosis, "find-me," "eat-me," and "don't eat-me" signals. Compared to

microglia, there were very few ARDEGs, as only 13 genes were found to be significantly (adjusted

p-value < 0.05) upregulated in aged Astrocyte-qNSC as compared to young, while none were

identified in aged aNSC, TACs or Neuroblasts when compared to young (Supplementary Figs. 1E-

J). Because of the lack of significant ARDEGs, instead of overlapping gene sets of interest with

the ARDEGs for each of the neural progenitor cell types, we utilized GSEA, as this method does

not rely on setting significance thresholds and considers changes of sets of genes rather than

individual genes (Methods). We paired this approach with curated gene sets that contained genes

associated with apoptosis, "find-me," "eat-me," and "don't eat-me" signals (Supplementary Table

2). With this, we found a significant (adjusted p-value < 0.05) age-related upregulation of the

expression of the apoptosis gene set in Astrocyte-qNSC cells, which contrasted with an age-related

decrease in aNSC, TACs and Neuroblasts (Fig. 2C; Supplementary Fig. 3A). Although we were

unable to find any significant enrichments when using genes sets for "find-me" or "eat-me"

signals, we did find significant age-related upregulation of the expression for complement system

and opsonization gene sets (known to be associated with "eat-me" signals (Li, 2012)) in Astrocyte-

qNSC and Neuroblasts, but not in aNSC or TACs (Fig. 3C; Supplementary Fig. 3B-C).

Intriguingly, we also found significant age-related upregulation of the expression for the "don't

eat-me" gene set in Astrocyte-qNSC and Neuroblasts but not in aNSC-NPCs (Fig. 2C;

Supplementary Fig. 3D). The "don't eat-me" gene set includes genes that are part of the MHC

Class I complex, as they are known to function as a "don't eat-me signal" against macrophages in

certain contexts, especially when co-expressed with the *B2m* subunit (Khalaji et al., 2023).

Importantly, the receptor of MHC I, *Lilrb4a*, is among the genes whose expression is significantly

upregulated in aged microglia (Supplementary Fig. 1C). Because of this, we also looked for age-

related expression changes using the antigen processing and presentation gene set (MHC pathway).

The expression pattern of this gene set matched the trends of the "don't eat-me" gene set, which is

suggestive of anti-phagocytic function of the MHC Class I complex in aged Astrocyte-qNSC and

Neuroblasts, but not aNSC-NPCs. In summary, while all 3 neural progenitor cell types have

increased expression of complement system genes that are associated with "eat-me" signals and

thus may promote phagocytosis, Astrocyte-qNSC and Neuroblasts appear to counteract this signal

through increased expression of "don't eat-me" signals such as *Cd200, Pecam1*, and the MHC

class 1 complex (*H2-K1, H2-D1, B2m*). This suggest that the age-related increase in phagocytic

activity of microglia may be targeted towards aNSC and TACs.

Figure 2. Re-analysis of aging SVZ scRNA data reveals age-related increase of microglia phagocytosis.

(A) Pseudobulk level analysis of differential gene expression (DGE) changes during aging of young (3 months old) and aged (28–29 months old) microglia in the SVZ. Red points genes that overlapped with the phagocytosis-related gene sets (Supplementary table 2) and are log2 fold change > 1 and adjusted p-value < 0.05. Statistics: n=3 mice/group, DGE results obtained using Wald test followed by Benjamini-Hochberg correction for multiple hypothesis testing. (B) Heatmap of age-related DEGs (adjusted p-value < 0.05) in aging SVZ microglia (obtained from analysis in (A)) that overlapped with phagocytosis-related gene sets (Supplementary table 2) Statistics: n=3 mice/group. (C) Gene set enrichment analysis of age-related gene expression changes in neural progenitor cells using apoptosis, "find-me," "eat-me," and "don't eat-me" signal gene sets (Supplementary table 2). Statistics: n=3 mice/group, GSEA p-values obtained using a permutation test followed by followed by Benjamini-Hochberg correction for multiple hypothesis testing.

Microglia in the SVZ have an age-related increase of hagocytosis of NSCs and TACs

To assess SVZ microglia phagocytic activity, we dissected out the SVZ from the brain of mice, isolated microglia from either the SVZ or whole brain (excluding the SVZ) from 3- and 12-month old mice (young and old), and then expanded them *in vitro* where engulfment of yeast particles conjugated to pHrodo, a pH sensitive dye (Kapellos et al., 2016), was measured via live imaging (Methods). Interestingly, we observed relatively low phagocytic activity in the 3-month-old SVZ microglia compared to 3-month old microglia isolated from the whole brain, suggesting that microglia in the SVZ niche are specialized to be less phagocytic than non-germinal areas in the brain. However, microglia from 12-month-old SVZ showed a significant age-related increase in phagocytosis activity compared to the 3-month-old SVZ microglia, which approached levels of phagocytic activity of microglia isolated from whole brain (Fig. 3A).

Next, we sought to validate the findings of an age-related increase in SVZ microglia phagocytosis activity in our *in vitro* system by examining SVZ microglia phagocytosis *in vivo*. First, we asked if age-associated increases in microglia lysosomal activity in the SVZ was due to increased phagocytosis of apoptotic NSCs and TACs. We performed immunohistochemistry for cleaved caspase 3 on coronal sections from 3-month-old and 12-month-old mice. We did not observe a significant increase in apoptosis in the aged SVZ (Fig.3B) Next, to detect if phagocytosis increased during aging, we performed immunohistochemistry (IHC) on SVZ wholemounts against the microglia marker Iba1, lysosome marker CD68, and the aNSC/TAC marker Mash1 (Kim et al., 2011; Lim and Alvarez-Buylla, 2016). Interestingly, we observed colocalization of the lysosome marker CD68 and Mash1 within microglia in both 3- and 12-month-old SVZs (Fig. 3C

&D). Quantification revealed approximately 20% of SVZ microglia contain Mash1 within their

lysosomes in 3-month-old SVZ, which significantly increased by 12 months of age (Fig. 3C).

Figure 3. Age-related increase in SVZ microglia phagocytosis *in vitro* and *in vivo*. (A) Time-
lapse *in vitro* phagocytosis assay of young (3-month) or aged (12-month) SVZ or whole brain
(WB) microglia labeled with isolectin and incubated with Zymosan-pHrodo (Zymo),
CytochalasinD and Zymosan-pHrodo (CytoD), or isolectin only (Control). Phagocytosis is
quantified as pHrodo+ area indicated by red fluorescence / microglia number. Statistics: n=3
mice/group, microglia > 1,000 per mouse. (B) Quantification of Caspase 3 in coronal brain
sections. Statistics: n=6 mice/group. (C) Quantification of *in vivo* phagocytosis of 3- or 12-month
SVZ microglia measured as Iba1+CD68+Mash1+ microglia / total Iba1+ microglia. *n*=11
mice/group. (D) Representative confocal projection images of young or aged SVZ microglia
stained with Iba1 (*green*), microglia lysosomes stained with CD68 (*red*), and TAC transcription
factor stained with Mash1 (*magenta*). Scale bar: 10µm. P-value legend: ns: no symbol; *: p ≤ 0.05,
**: p ≤ 0.01, ***: p ≤ 0.001, ****: p ≤ 0.0001.

CR2-crry complement 3 intervention increased NSCs proliferation but not phagocytosis

To identify possible mechanisms for the age-related increase in microglia phagocytosis of

aNSC/TAC in the SVZ, we performed qRT-PCR for pro-inflammatory genes shown to be

upregulated in microglia during aging from our analysis of RNAseq databases. We reasoned that

pro-inflammatory genes that are very low or absent in the young SVZ compared to microglia

isolated from outside the SVZ that are increased with age in SVZ microglia would be good

candidates based on our *in vitro* phagocytosis results. One gene identified to be higher in microglia

isolated from outside the germinal region compared to young SVZ microglia, that was highly

upregulated upon aging was Complement 3 (C3) (Fig. 2, Supplementary Fig. 4). We found that

receptors for C3, *ITGAM* (Complement receptor 3), and *ITGAX* (Complement receptor 4) were

upregulated in SVZ microglia during aging (Supplementary Fig 4). To inhibit C3 activity we

utilized CR2-crry, a protein developed for targeted inhibition of C3 cleavage and C3 deposition

(Atkinson et al., 2005; Song et al., 2003). 3- and 12-month-old mice (young and old) mice received

CR2-crry or saline treatment every other day for a total of 3 doses followed by an EdU pulse 2

hours before sacrifice to label proliferating cells (Fig. 4A). SVZ wholemounts were collected and

IHC was performed as described above .

CR2-crry treatment had no significant effect on proliferation (Fig. 4F), SVZ microglia

CD68 area (Fig. 4B), or SVZ microglia phagocytosis of TACs (Fig. 4G) in the SVZs isolated from

3-month-old mice. We observed an expected decline in SVZ proliferation from 3- and 12-month-

old saline treated mice (Fig. 4F). There was a significant increase in SVZ proliferation in 12-

month-old CR2-crry treated groups compared to 12-month-old saline treated groups indicated by

EdU+ cells (Fig. 4F). We observed an increase in CD68 area/Iba1+ cells in 12-month-old saline

treated groups compared to 3-month-old saline treated groups (Fig. 4B), with a non-significant

decrease in 12-month-old CR2-crry treatment CD68 area/Iba1+ cell compared to 12-month-old

controls (Fig. 4B). There was an increase of Iba1+CD68+Mash1+ colocalization within Iba1+ cells

in 12-month-old saline treated controls compared to 3-month old saline treated controls (Fig. 4G)

with an increase of Iba1+CD68+Mash1+ colocalization within Iba1+ cells in 12-month-old CR2-

crry treatment vs 12-month-old controls (Fig. 4G). These findings suggest that C3 is involved in

age-related declines in proliferation but not aNSC and TAC phagocytosis. The increase in

phagocytosis of aNSCs and TACs during aging suggests that increasing the number of aNSCs and

TACs produced via cell division results in more cells being present to undergo phagocytosis.

Figure 4. Minocycline treatment restores some proliferation in the aged SVZ but does not reduce SVZ microglia phagocytosis. (A) Graphic representation of experimental design. (B) Quantification of EdU positive cells in the SVZ of young (3-month) or aged (12-month) mice. (C) Confocal micrographs the SVZ from young vehicle treated, old vehicle treated and minocycline treated mice. (D) Quantification of CD68 area Iba1+ microglia n=5 mice/group. (E) Quantification of percent Iba+ Mash1+ CD68+ cells n=5mice/group. P-value legend: ns: no symbol; *: p ≤ 0.05, **: p ≤ 0.01, ***: p ≤ 0.001, ****: p ≤ 0.0001.

Minocycline anti-inflammatory intervention effects on SVZ microglia and NSCs

We have previously seen that SVZ microglia have high levels of activation with increased

CD68 compared to non-germinal regions during aging, resulting in a pro-inflammatory milieu in

the SVZ (Solano Fonseca et al., 2016). We wanted to determine if broadly reducing the activation

of aged SVZ microglia would result in increased proliferation and/or reduced phagocytosis. We

treated 3- and 12-month old mice (young and old) mice with minocycline, an antibiotic with potent

anti-inflammatory properties (Elewa et al., 2006; Garrido-Mesa et al., 2013; Plane et al., 2010;

Tikka et al., 2001; Yrjänheikki et al., 1998) or saline for 14 days followed by a 2 hour EdU pulse

to label proliferating cells (Fig. 5A) before euthanasia. SVZ wholemounts were collected for

immunohistochemistry as previously described. At 3 months of age, minocycline treatment did not

have a significant effect on microglial CD68 expression (Fig. 5B), proliferation (Fig. 5F), or SVZ

microglia phagocytosis of Mash1 positive cells (Fig 5G). At 12 months of age, minocycline

treatment resulted in a significant reduction in microglia expression of CD68 compared to saline

treated controls (Fig. 5B), indicating that minocycline reduced inflammation in the aged SVZ.

Minocycline treatment resulted in increased proliferation in the aged SVZ (Fig. 5F) although this

did not reach levels measured in young mice. As expected, we saw a significant increase in the

number of CD68 positive lysosomes containing Mash1 in the 12-month-old SVZs compared to

young 3-month-old SVZs (Fig. 5G). Surprisingly, we observed a significant increase in Mash1

containing lysosomes in the SVZs of 12-month-old mice treated with minocycline compared to

saline treated controls (Fig. 5G), indicating that anti-inflammatory treatment alone does not reduce

phagocytosis in the SVZ.

Figure 5. Complement inhibition partially rescues SVZ proliferation in aged mice and further increases SVZ microglia phagocytosis. (A) Graphic representation of CR2-crry or saline administration to young (3-month) or aged (12-month) mice. (B) Quantification of microglia activation measured by CD68 area / number of microglia. (C) Confocal projection images of young control, aged control, aged CR2-crry treated SVZ microglia stained with Iba1 (*green*), microglia lysosomes stained with CD68 (*red*), and TAC transcription factor stained with Mash1 (*magenta*). Arrowheads in aged control and aged CR2-crry treated indicate colocalization of Mash1+ signal within microglia lysosomes (Iba1+CD68+). Scale bar: 10μm. (D) Quantification of proliferating EdU-positive cells in the SVZ of young or aged control or CR2-crry treated mice. (E) Quantification of Mash1 containing microglia measured by Iba1+CD68+Mash1+ / number of Iba1+ cells from young or aged control or CR2-crry treated mice. *n*=4 mice/group. P-value legend: ns: no symbol; *: p ≤ 0.05, **: p ≤ 0.01, ***: p ≤ 0.001, ****: p ≤ 0.0001.

Discussion

The adult SVZ produces new neurons for the distant olfactory bulb throughout life which

contribute to olfactory learning and discrimination (Lledo and Vally 2016). In addition, following

brain injury neural progenitors and neuroblast undergo a proliferative burst and then migrate to

damaged areas where they give rise to a limited number of neurons, astrocytes and

oligodendrocytes and secrete pro-reparative factors (Liang…Carmichael 2019, Yamashita 2006 J

Neuro, Imotola 2004, Robin, 2006). The blockade of this response results in larger legions and

worsens behavioral deficits (Carmichael Ohab 2006). This reparative ability makes the SVZ an

enticing therapeutic target to boost the endogenous repair response. During aging SVZ

neurogenesis becomes drastically reduced with the most-rapid reductions between 9-12 months of

age in the rodent (Luo…Conover 2006 Aging Cell) corresponding to middle-age and persists for

the duration of aging. The reduction is coincidental to periods of increasing risk for brain injury

associated with aging. Thus, understanding drivers of SVZ NSC age-associated dysfunction is

imperative. Our findings indicate that pro-inflammation and upregulation of phagocytotic

signatures within the SVZ niche by microglia plays a role in decreased NSC number and reduced

proliferation reported during aging.

During embryonic neurogenesis, SVZ microglia have ameboid morphology and regulate

the size of the precursor pool by phagocytosis of non-dying neuroblasts (Cunningham et al., 2013;

Morrison et al., 2023) In the adult dentate gyrus, another neurogenic niche, unactivated microglia

phagocytose newborn neurons undergoing apoptosis In contrast, in the adult SVZ, very little

apoptosis occurs (Luo et al., 2006). In contrast, in the olfactory bulb where newborn neurons

generated in the SVZ must integrate into the circuitry widespread phagocytosis has been reported

(Xavier et al., 2015). Using immunohistochemistry, we did not observe an increase in apoptosis in

the aging SVZ suggesting that the observed increased phagocytosis of aNSCs/TACs was not due

to increased phagocytosis.

Microglia are distributed throughout the brain, however there are regional differences that

make them heterogenous within their local microenvironment such as differences in cell density,

morphology, transcriptomes, and receptor expression (Grabert et al., 2016; Tan et al., 2020). SVZ

microglia have distinct characteristics compared to microglia from non-germinal regions

indicating the specialization of microglia within the SVZ niche (Marshall et al., 2014; Ribeiro

Xavier et al., 2015; Shigemoto-Mogami et al., 2014; Solano Fonseca et al., 2016)..Riberio Xavier

et al. reported that SVZ microglia in young adult mice not only have a distinct morphological

phenotype, but also have specialized gene and receptor expression compared to microglia

distributed in non-germinal regions. They also showed the necessity of SVZ microglia for

neuroblast survival and migration from the SVZ; while suggesting that SVZ microglia perform

little amounts of phagocytosis within the young SVZ niche, possibly because of low expression of

the P2Y12 receptor (Ribeiro Xavier et al., 2015). We found that young SVZ microglia *in* vitro

exhibit less phagocytosis then microglia from other brain regions which supports the notion that

they are specialized to be less phagovcytic in the young SVZ microenvironment.

To understand the roles of the observed age-associated increase in SVZ microglia

phagocytosis on aging NSCs of the SVZ, we utilized minocycline and CR2-crry in vivo

interventions to reduce SVZ microglia inflammation and phagocytosis respectively. Due to the

high levels of microglia activation in the SVZ, minocycline treatment was used to investigate if a

decrease in microglia inflammation in the aged SVZ would rescue NSC proliferation and/or result

in a decrease of TAC phagocytosis. Minocycline is often used as a general anti-inflammatory, that

has been shown to offer neuroprotection and reduces the activation of microglia (Ekdahl et al.,

2003; Plane et al., 2010; Tikka et al., 2001; Wu et al., 2002; Yrjänheikki et al., 1998). CR2-crry

treatment, a murine C3 activation and deposition inhibitor (Atkinson et al., 2005; Song et al., 2003)

that crosses the blood brain barrier (Alawieh et al., 2018), was used to selectively inhibit C3 of the

complement pathway. With minocycline treatment we observed a significant decrease in SVZ

microglia activation in the 12-month-old treated mice, indicating that minocycline was reducing

inflammation in the aged SVZ. Importantly, the aged mice that received minocycline had an

increase in SVZ proliferation, indicating that the age-associated decline of SVZ proliferation can

be partially rescued by the administration of minocycline. Similar to the minocycline results, CR2-

crry intervention in aged mice resulted in a trending decline in SVZ activation as well as an

increase in SVZ proliferation. Suggesting that treating aged mice with CR2-crry not only reduced

microglia activation in the SVZ but also partially rescued the age-associated decline of SVZ

proliferation in a complement mediated fashion. Both in vivo interventions had strikingly similar

results. including an increase in SVZ microglia phagocytosis of aNSCs and TACs in the aged

groups treated with either minocycline or CR2-crry. With both interventions we saw a further

increase in aNSCs and TAC phagocytosis. The increase in microglia phagocytosis observed in the

12 month treated mice occurred in the presence of increased SVZ proliferation, which has not been

previously described. Our findings suggest, based on the increase in number of proliferating SVZ

cells, that SVZ microglia inflammation is responsible at least in part for the age-related reduction

in neurogenesis. However, the increases seen in SVZ microglia phagocytosis suggest that more

cells in the aged SVZ further increases phagocytosis. This could be potentially due to the increase

of cells within the SVZ expressing "eat me" signals without upregulation of the "don't eat me"

signals that were associated with qNSCs and neuroblasts, which was detected in our analysis of

the available RNAseq data. An increase in cells that express "eat me" signals will be an important

future direction to explore as a possible mechanistic answer to our observed age-associated

increase in SVZ microglia phagocytosis.

Methods

<u>Animals and Interventions</u>

Male and female C57BL/6 mice were used in this study either bred at the UTHSA vivarium or

acquired from the National Institute of Aging. Young mice (2-4 months of age) and aged mice (12-

14 months of age) were housed in accordance with the University of Texas Health San Antonio

Institutional Animal Care and Use Committee and performed in accordance with institutional and

federal guidelines and allowed free access to water and standard rodent chow ad libitum.

Both male and female C57BL/6 mice (2-4 months or 12-14 months) received CR2-crry (10µg/1g

mouse; Stephen Tomlinson Lab, Medical University of South Carolina) or minocycline (50µg/1g

mouse; Sigma-aldrich, Cat. No. M9511) with respective saline controls via intraperitoneal

injection. Minocycline or saline was administered daily for 2 weeks. CR2-crry or saline was

administered every other day for a total of 3 doses.

<u>Immunohistochemistry</u>

The SVZ was isolated for wholemount (WM) tissue analysis; mice were anesthetized with

isoflurane, a cervical dislocation was performed, followed by brain extraction. Brains were placed

into cold HIB where they were halved and the septum and hippocampus were removed, and the

SVZ was microdissected from the surrounding brain tissue. The SVZ WM was then fixed in fresh

4% paraformaldehyde (PFA) for 30 minutes. SVZ WMs were permeabilized in 2% PBST and

blocked overnight at 4° C in 10% normal donkey serum (NDS). The SVZ WMs were then

incubated at 4°C for 4 days with primary antibody in blocking solution, followed by overnight

incubation at 4°C with secondary antibody made in blocking solution, and a 5-minute incubation

at room temperature (RT) with DAPI (1:1000; Invitrogen, Cat. No. D1306) then mounted on a

slide in ProLong Gold antifade reagent (Invitrogen, Cat. No. P36930). Primary antibodies that

were used include Iba1 (1:200; Wako, Cat. No. 019-19741 and 1:200; Novus, Cat. No. NB100-

1028), CD68 (1:250; BioRad, Cat. No. MCA1957), and Mash1 (1:5000; CosmoBioUSA, Cat. No.

SK-T01-003) Cleaved caspase 3 (1:200 Cell Signaling Cat. No D175 5A1E). Species-specific

secondary antibodies (Jackson ImmunoResearch) were used at a concentration of 1:500 in SVZ

WMs.

<u>Immunohistochemistry Quantification</u>

EdU was quantified in the SVZ by making a "Z Project" 2-dimensional image of the Z-stack in

ImageJ on 10 random fields of view and hand counting the number of EdU+ nuclei. The number

of microglia were quantified by hand counting Iba1+ cells, excluding cells whose cell body

touched the image boundary on Zen Microscopy Software (Zeiss). Total CD68 area was quantified

by making a "Z project" 2-dimensional image of the Z-stack, using the "Threshold" function, and

then the "Analyze particle" function to give total area in µm2 that was then averaged by total

number of microglia in each image. Total number of Iba1+ microglia were quantified similarly to

EdU+ cells described above. Mash1 colocalization within Iba1+ microglia were quantified by hand

counting in Zen Microscopy Software (Zeiss) by examining each Iba1+ microglia's lysosome

(CD68+) to determine if each Iba1+CD68+ microglia contained Mash1 colocalization. The

number of Mash1+CD68+Iba1+ microglia were then averaged to total number of Iba1+ microglia

in each image. There were no microglia that contained Mash1 that were not CD68+. SVZ WMs

were analyzed from 15µm thick Z-stacks (15 Z-stacks per WM). Confocal images were obtained

with a Zeiss LSM 710 confocal microscope at 40x objective.

<u>Single-cell RNA-sequencing Quality control filtering</u>

Quality control of downloaded single-cell RNA-Seq data (see Data Availability) was first

performed, which filtered cells where the number of genes or UMIs detected were greater or less

than 5 median absolute deviations (MADs) or percent mitochondria reads greater or less than 3

MADs, relative to all cells in the dataset. Genes were filtered if detected in less than 10 cells out

of all cells in the dataset. Cell identity labels were those assigned by the original analysis.

<u>Single-cell RNA-sequencing differential expression and gene set enrichment analysis</u>

Pseudobulk samples were created by summing counts of each identified cell type for each

biological replicate. The DESeq2 package (MI et al., 2014) was then used to perform differential

expression analysis to compare old vs. young groups. Following this the ClusterProfiler package

(G et al., 2012) was then used to perform gene set enrichment analysis (GSEA) using the gene

ontology biological process database.

<u>Microglia phagocytosis gene set analysis</u>

Genes associated with known phagocytosis receptors and/or pathways active in microglia were

manually curated by searching the Molecular Signatures Database (msigDB) and the literature (i.e.

Complement, Toll-like receptor, TAM receptor, Trem2 receptor, FC gamma, Pr2y6 receptor,

Scavenger receptor, Mannose receptor, LRP1 receptor, Galectin-3 adaptor protein, Microglia

Activation, Microglia Proliferation, Disease-Associated Microglia (DAM), and Microglia

Development). In addition, gene sets related to screens of phagocytosis regulators was obtained

from the literature (see Data Availability). These gene sets were then overlapped with the genes

that were found to be significantly differently expressed (adjusted p-value < 0.05) in the microglia

old vs. young comparison and plotted as a heatmap using the ComplexHeatmap package.

<u>Neural stem cell gene set analysis</u>

Genes associated with pathways known to be expressed in cells that are targeted for phagocytosis

were manually curated by searching the Molecular Signatures Database (msigDB) and the

literature (i.e. Apoptosis, Opsonization, Antigen Processing and Presentation, "eat-me," "find-

me," and "don't eat me"). Because of the lack of genes which were significantly differentially

expressed in neural stem cells (i.e. Astrocytes/qNSCs, aNSCs/NPCs and Neuroblasts) in the old

vs. young comparison, the ClusterProfiler package (ref) was then used to perform gene set

enrichment analysis (GSEA) using the manually curated gene sets. The Log_2 fold change of genes

within the gene sets found to be significantly enriched in age-related genes expression of neural

stem cells were then plotted as a heatmap using the ComplexHeatmap package.

<u>Microglia Isolation and Cell Culture</u>

465 Microglia were cultured in two ways, either plated following MACS for microglia specific culture

466 or through mixed glial culture. C57BL/6 mice were injected intraperitoneally with pentobarbital

467 (25µL/mouse) then perfused with cold 1x PBS. SVZ WMs or the remainder of the whole brain

468 excluding the SVZ and cerebellum, hereby referred to as whole brain (WB), were isolated

469 following perfusion. The tissues were minced and dissociated in 50U papain in DMEM/F12 at

470 37°C for 45 minutes and manually triturated and spun 3x in DMEM. SVZ and WB homogenate

471 were gently pressed through a 70µm cell strainer to remove debris. For mixed glial culture, SVZ

472 homogenate were then plated in a T25 flask coated with poly-L-ornithine (PLO) in neural growth

473 media (DMEM/F12, 5% fetal bovine serum, 1x N2, 10ng/mL EGF, and 10ng/mL bFGF) until a

474 confluent cell monolayer formed. The culture was then fed microglia proliferation media

475 (DMEM/F12, 10% fetal bovine serum, 1x N2, and 20 ng/mL GM-CSF) until microglia could be

476 seen atop the monolayer. Once microglia were present, they were shaken off from the cell

477 monolayer and the SVZ microglia could be plated for further experimentation.

478 For microglia isolated by MACS, following the cell strainer step, WB homogenate was

479 resuspended in HIB and centrifugated at 400G with no brake in 22% Percoll (Sigma-aldrich, Cat.

480 No. P4937) solution to remove myelin from the samples. The WB homogenate was then

481 resuspended in 0.5% bovine serum albumin and incubated with 10µL CD11b Microbeads (cat#

482 130-093-636) per 107 cells. The homogenate was then placed in an LS column (MiltenyiBiotec,

483 Cat. No. 130-042-401) in a QuadroMACS™ separator (MiltenyiBiotec, Cat. No. 130-090-976) to

484 allow for MACS sorting of CD11b + microglia following manufacturer protocols

485 (MiltenyiBiotec). WB microglia were plated in a 6-well plate coated with PLO/laminin (10µg/mL;

486 Sigma-aldrich, Cat. No. L2020) at 250k cells/well in microglia expansion media (DMEM/F12,

487 100 µg/mL PenStrep, 5 µg/mL n-acetyl-L-cysteine, 5 µg/mL Insulin, 100 µg/mL apo-transferrin,

100 µg/mL sodium selenite, 1 µg/mL heparin sulfate, 10% fetal bovine serum, 1.5 µg/mL

cholesterol, 40 ng/mL M-CSF, 20 ng/mL GM-CSF, and 2 ng/mL TGF-β) adapted from [Bohlen et

al., 2017].

<u>Microglia *in vitro* Phagocytosis Assay</u>

The phagocytosis assay was performed using live cell imaging by Incucyte S3 (Essen BioScience)

for WB or SVZ microglia phagocytosis from 3- or 12-month mice. WB microglia were passaged

from a 6-well plate with Trypsin (Fisher, Cat. No. 25-200-072) following MACS isolation and

SVZ microglia were collected from mixed glial culture and microglia were plated in a 96 well

plate coated with PLO/laminin at 20k cells/well in microglia expansion media for 3 days. After 3

days, SVZ and WB microglia were moved to serum free media (DMEM/F12, 100 µg/mL PenStrep,

5 µg/mL n-acetyl-L-cysteine, 5 µg/mL Insulin, 100 µg/mL apo-transferrin, 100 µg/mL sodium

selenite, 1 µg/mL heparin sulfate, 1.5 µg/mL cholesterol, 40 ng/mL M-CSF, and 2 ng/mL TGF-β)

adapted from [Bohlen et al., 2017] for 1 week. All microglia were then incubated with 5 µg/mL

Isolectin IB4 Alexa Fluor™ 488 Conjugate (Invitrogen, Cat. No. I21411) for 60 minutes at 37°C.

SVZ or WB microglia of both ages were treated with Zymosan S. cerevisiae BioParticles™ (10

mg/mL; Invitrogen, Cat. No. Z2849) conjugated pHrodo™ Red SE (Molecular Probes™, Cat. No.

P36600) following manufacturer protocols (ThermoFisher) and adapted from [Haney et al., 2018;

Pluvinage et al., 2019], both Zymosan-pHrodo and Cytochalasin D (10 mM/µL Sigma-aldrich,

Cat. No. C8273) to inhibit phagocytosis, or isolectin only as a negative control. The microglia

were then incubated in the Incucyte S3 with fluorescent images captured every hour over the

course of 48 hours for live cell imaging of microglia phagocytosis. The images were analyzed in

the IncuCyte software (Essen BioScience) quantifying colocalization of the green fluorescence of

Isolectin IB4 labelled microglia and the red fluorescence of engulfed Zymosan-pHrodo within the

microglia lysosome demonstrating microglia phagocytosis. Total pHrodo area / number of

microglia was calculated for each well and condition yielding amount of in vitro phagocytosis

occurring at each time point.

<u>RNA Extraction and qPCR</u>

RNA was extracted from freshly isolated microglia following magnetic-activated cell sorting

(MACS) isolation, using a TRIzol-chloroform method (Rio et al., 2010) where cells were lysed in

TRIzol (Invitrogen, Cat. No. 15596018), aqueous layer separated with chloroform, RNA was

precipitated with isopropanol, washed in ethanol, dried, then resuspended in RNase-free water.

cDNA was synthesized using High Capacity cDNA Reverse Transcription Kit (Applied

Biosystems, Cat. No. 4368814). Quantitative polymerase chain reaction (qPCR) was performed

using a C1000 Touch Thermal Cycler (BioRad) with iTaq™ Universal SYBR® Green Supermix

(BioRad, Cat. No. 1725121). Reactions were performed in doublet with glyceraldehyde-3-

phosphate dehydrogenase (GAPDH) used as an endogenous control. Negative control reactions

were performed using the same conditions without cDNA. Quantification was analyzed using the

ΔΔCt method as described [Livak and Schmittgen, 2001]. Primers were used for the following

transcripts: C1QA, P2Y12, TREM2, LPAR, C3, CR1, ITGAM, ITGAX, GAPDH.

Primers	Forward	Reverse
C1QA	AAAGGCAATCCAGGCAATATCA	TGGTTCTGGTATGGACTCTCC
C3	CCAGCTCCCCATTAGCTCTG	GCACTTGCCTCTTTAGGAAGTC

TREM2	CTGGAACCGTCACCATCACTC	CGAAACTCGATGACTCCTCGG
LPAR	AGCCATGAACGAACAACAGTG	CATGATGAACACGCAAACAGTG
CR1	ATGGAGGTCTCTTCTCGGAGT	GGCCGAAGGCTACAAGGAG
ITGAM	GGGAGGACAAAAACTGCCTCA	ACAACTAGGATCTTCGCAGCAT
ITGAX	CTGGATAGCCTTTCTTCTGCTG	GCACACTGTGTCCGAACTCA
GAPDH	AGGTCGGTGTGAACGGATTTG	TGTAGACCATGTAGTTGAGGTCA

EdU Labeling and Staining

Mice were injected intraperitoneally (50µg EdU/1g mouse) 2 hours before sacrifice. Their brains

were removed and SVZ was isolated and fixed for 30 minutes in 4% PFA. EdU staining was

performed following manufacturer protocols (Fisher, Click-it EdU Imaging Kit, Cat. No. C10337).

Data Analysis

Student's t-tests and two-way ANOVA were used to analyze the data on GraphPad Prism 8

Software. All data are expressed as mean ± standard error of the mean.

Data Availability

Normalized single-cell RNA-sequencing data and cell type annotations of the SVZ from young (3

months old) and old (28–29 months old) male mice (n = 3) was downloaded from Supplementary

Table 2 of the Dulken et al., 2019 study. Results of screen for phagocytosis regulator genes were

downloaded from Supplementary Table 5 of the Haney et al., 2019 study and Supplementary Table

2 of the Pluvinage et al., 2019 study. Marker genes of major cell types in the SVZ of young (2-4

months old) male and female mice were downloaded from Table S1 of the Zywitza et al., 2018

study.

Code Availability

Code to reproduce bioinformatics analyses are available at the following repository:

https://github.com/cutleraging/nsc-aging

References

Ahlenius, H., Visan, V., Kokaia, M., Lindvall, O., and Kokaia, Z. (2009). Neural stem and

progenitor cells retain their potential for proliferation and differentiation into functional neurons

despite lower number in aged brain. J. Neurosci. *29*, 4408–4419.

https://doi.org/10.1523/JNEUROSCI.6003-08.2009.

Alvarez-Buylla, A., and García-Verdugo, J.M. (2002). Neurogenesis in adult subventricular zone.

J. Neurosci. *22*, 629–634. https://doi.org/10.1523/jneurosci.22-03-00629.2002.

Atkinson, C., Song, H., Lu, B., Qiao, F., Burns, T.A., Holers, V.M., Tsokos, G.C., and Tomlinson,

S. (2005). Targeted complement inhibition by C3d recognition ameliorates tissue injury without

apparent increase in susceptibility to infection. J. Clin. Invest. *115*, 2444–2453.

https://doi.org/10.1172/JCI25208.

Colonna, M., and Butovsky, O. (2017). Microglia function in the central nervous system during

health and neurodegeneration. Annu. Rev. Immunol. *35*, 441–468.

https://doi.org/10.1146/annurev-immunol-051116-052358.

Elewa, H.F., Hilali, R., Hess, D.C., Machado, L.S., and Fagan, S.C. (2006). Minocycline for short-

term neuroprotection. Pharmacotherapy *26*, 515–521. https://doi.org/10.1592/phco.26.4.515.

Enwere, E., Shingo, T., Gregg, C., Fujikawa, H., Ohta, S., and Weiss, S. (2004). Aging results in

reduced epidermal growth factor receptor signaling, diminished olfactory neurogenesis, and

deficits in fine olfactory discrimination. J. Neurosci. *24*, 8354–8365.

https://doi.org/10.1523/JNEUROSCI.2751-04.2004.

Garrido-Mesa, N., Zarzuelo, A., and Gálvez, J. (2013). Minocycline: Far beyond an antibiotic. Br.

J. Pharmacol. *169*, 337–352. https://doi.org/10.1111/bph.12139.

Ginhoux, F., Greter, M., Leboeuf, M., Nandi, S., See, P., Gokhan, S., Mehler, M.F., Conway, S.J.,

Ng, L.G., Stanley, E.R., et al. (2010). Fate mapping analysis reveals that adult microglia derive

from primitive macrophages. Science (80-.). *330*, 841–845.

https://doi.org/10.1126/science.1194637.

Goings, G.E., Kozlowski, D.A., and Szele, F.G. (2006). Differential activation of microglia in

neurogenic versus non-neurogenic regions of the forebrain. Glia *54*, 329–342.

https://doi.org/10.1002/glia.20381.

Kalamakis, G., Brüne, D., Ravichandran, S., Bolz, J., Fan, W., Ziebell, F., Stiehl, T., Catalá-

Martinez, F., Kupke, J., Zhao, S., et al. (2019). Quiescence Modulates Stem Cell Maintenance and

Regenerative Capacity in the Aging Brain. Cell *176*, 1407-1419.e14.

https://doi.org/10.1016/j.cell.2019.01.040.

Kapellos, T.S., Taylor, L., Lee, H., Cowley, S.A., James, W.S., Iqbal, A.J., and Greaves, D.R.

(2016). A novel real time imaging platform to quantify macrophage phagocytosis. Biochem.

Pharmacol. *116*, 107–119. https://doi.org/10.1016/j.bcp.2016.07.011.

Kim, E.J., Ables, J.L., Dickel, L.K., Eisch, A.J., and Johnson, J.E. (2011). Ascl1 (Mash1) Defines

Cells with Long-Term Neurogenic Potential inSubgranular and Subventricular Zones in Adult

Mouse Brain. PLoS One *6*, 18472. https://doi.org/10.1371/JOURNAL.PONE.0018472.

Lim, D.A., and Alvarez-Buylla, A. (2016). The adult ventricular–subventricular zone (V-SVZ)

and olfactory bulb (OB) neurogenesis. Cold Spring Harb. Perspect. Biol. *8*, a018820.

https://doi.org/10.1101/cshperspect.a018820.

Marshall, G.P., Deleyrolle, L.P., Reynolds, B.A., Steindler, D.A., and Laywell, E.D. (2014).

Microglia from neurogenic and non-neurogenic regions display differential proliferative potential

and neuroblast support. Front. Cell. Neurosci. *8*, 180. https://doi.org/10.3389/fncel.2014.00180.

Maslov, A.Y., Barone, T.A., Plunkett, R.J., and Pruitt, S.C. (2004). Neural Stem Cell Detection,

Characterization, and Age-Related Changes in the Subventricular Zone of Mice. J. Neurosci. *24*,

1726–1733. https://doi.org/10.1523/JNEUROSCI.4608-03.2004.

Plane, J.M., Shen, Y., Pleasure, D.E., and Deng, W. (2010). Prospects for minocycline

neuroprotection. Arch. Neurol. *67*, 1442–1448. https://doi.org/10.1001/archneurol.2010.191.

Ribeiro Xavier, A.L., Kress, B.T., Goldman, S.A., De Lacerda Menezes, J.R., and Nedergaard, M.

(2015). A distinct population of microglia supports adult neurogenesis in the subventricular zone.

J. Neurosci. *35*, 11848–11861. https://doi.org/10.1523/JNEUROSCI.1217-15.2015.

Shigemoto-Mogami, Y., Hoshikawa, K., Goldman, J.E., Sekino, Y., and Sato, K. (2014).

Microglia enhance neurogenesis and oligodendrogenesis in the early postnatal subventricular

zone. J. Neurosci. *34*, 2231–2243. https://doi.org/10.1523/JNEUROSCI.1619-13.2014.

Solano Fonseca, R., Mahesula, S., Apple, D.M., Raghunathan, R., Dugan, A., Cardona, A.,

O'Connor, J., and Kokovay, E. (2016). Neurogenic niche microglia undergo positional remodeling

and progressive activation contributing to age-associated reductions in neurogenesis. Stem Cells

Dev. *25*, 542–555. https://doi.org/10.1089/scd.2015.0319.

Song, H., He, C., Knaak, C., Guthridge, J.M., Holers, V.M., and Tomlinson, S. (2003).

Complement receptor 2-mediated targeting of complement inhibitors to sites of complement

activation. J. Clin. Invest. *111*, 1875–1885. https://doi.org/10.1172/JCI17348.

Tikka, T., Fiebich, B.L., Goldsteins, G., Keinänen, R., and Koistinaho, J. (2001). Minocycline, a

tetracycline derivative, is neuroprotective against excitotoxicity by inhibiting activation and

proliferation of microglia. J. Neurosci. *21*, 2580–2588. https://doi.org/10.1523/jneurosci.21-08-

02580.2001.

Yrjänheikki, J., Keinänen, R., Pellikka, M., Hökfelt, T., and Koistinaho, J. (1998). Tetracyclines

inhibit microglial activation and are neuroprotective in global brain ischemia. Proc. Natl. Acad.

Sci. U. S. A. *95*, 15769–15774. https://doi.org/10.1073/pnas.95.26.15769.

Cunningham, C. L., Martínez-Cerdeño, V., & Noctor, S. C. (2013). Microglia Regulate the Number

625   of   Neural   Precursor   Cells   in   the   Developing   Cerebral   Cortex.

626   https://doi.org/10.1523/JNEUROSCI.3441-12.2013

Deczkowska, A., Keren-Shaul, H., Weiner, A., Colonna, M., Schwartz, M., & Amit, I. (2018).

Disease-Associated Microglia: A Universal Immune Sensor of Neurodegeneration. In A.

W. Hadas Keren-Shaul 1 2 5, Marco Colonna 3, Michal Schwartz 4, Ido Amit 1 (Ed.). Cell.

G, Y., LG, W., Y, H., & QY, H. (2012). clusterProfiler: an R package for comparing biological

themes among gene clusters. *Omics : a journal of integrative biology*, *16*(5).

https://doi.org/10.1089/omi.2011.0118

Haney, M. S., Bohlen, C. J., Morgens, D. W., Ousey, J. A., Barkal, A. A., Tsui, C. K., Ego, B. K.,

Levin, R., Kamber, R. A., Collins, H., Tucker, A., Li, A., Vorselen, D., Labitigan, L., Crane,

E., Boyle, E., Jiang, L., Chan, J., Rincón, E., . . . Bassik, M. C. (2018). Identification of

phagocytosis regulators using magnetic genome-wide CRISPR screens. *Nat Genet*, *50*(12),

1716-1727. https://doi.org/10.1038/s41588-018-0254-1

Hu, B., Fujian Provincial Key Laboratory of Neurodegenerative Disease and Aging Research, S.

o. M., Institute of Neuroscience, Xiamen University, Xiamen, China, Duan, S., Fujian

Provincial Key Laboratory of Neurodegenerative Disease and Aging Research, S. o. M.,

Institute of Neuroscience, Xiamen University, Xiamen, China, Wang, Z., Fujian Provincial

Key Laboratory of Neurodegenerative Disease and Aging Research, S. o. M., Institute of

Neuroscience, Xiamen University, Xiamen, China, Li, X., Fujian Provincial Key

Laboratory of Neurodegenerative Disease and Aging Research, S. o. M., Institute of

Neuroscience, Xiamen University, Xiamen, China, Zhou, Y., Fujian Provincial Key

Laboratory of Neurodegenerative Disease and Aging Research, S. o. M., Institute of

Neuroscience, Xiamen University, Xiamen, China, Zhang, X., Fujian Provincial Key

Laboratory of Neurodegenerative Disease and Aging Research, S. o. M., Institute of

Neuroscience, Xiamen University, Xiamen, China, Zhang, Y.-W., Fujian Provincial Key

Laboratory of Neurodegenerative Disease and Aging Research, S. o. M., Institute of Neuroscience, Xiamen University, Xiamen, China, Xu, H., Fujian Provincial Key Laboratory of Neurodegenerative Disease and Aging Research, S. o. M., Institute of Neuroscience, Xiamen University, Xiamen, China, Zheng, H., Fujian Provincial Key Laboratory of Neurodegenerative Disease and Aging Research, S. o. M., Institute of Neuroscience, Xiamen University, Xiamen, China, & Basic Medical Sciences, S. o. M., Xiamen University, Xiamen, China. (2024). Insights Into the Role of CSF1R in the Central Nervous System and Neurological Disorders. *Frontiers in Aging Neuroscience*, *13*, 789834. https://doi.org/10.3389/fnagi.2021.789834

Izgi, H., Han, D., Isildak, U., Huang, S., Kocabiyik, E., Khaitovich, P., Somel, M., & Dönertaş, H. M. (2022). Inter-tissue convergence of gene expression during ageing suggests age-related loss of tissue and cellular identity. https://doi.org/doi:10.7554/eLife.68048

Khalaji, A., Yancheshmeh, F. B., Farham, F., Khorram, A., Sheshbolouki, S., Zokaei, M., Vatankhah, F., & Soleymani-Goloujeh, M. (2023). Don't eat me/eat me signals as a novel strategy in cancer immunotherapy. *Heliyon*, *9*(10), e20507. https://doi.org/10.1016/j.heliyon.2023.e20507

Li, W. (2012). Eat-me signals: keys to molecular phagocyte biology and "appetite" control. *J Cell Physiol*, *227*(4), 1291-1297. https://doi.org/10.1002/jcp.22815

Liberzon, A., Broad Institute of MIT and Harvard, M. S. C., MA 02142, USA, Birger, C., Broad Institute of MIT and Harvard, M. S. C., MA 02142, USA, Thorvaldsdóttir, H., Broad Institute of MIT and Harvard, M. S. C., MA 02142, USA, Ghandi, M., Broad Institute of MIT and Harvard, M. S. C., MA 02142, USA, Mesirov, J. P., Broad Institute of MIT and Harvard, M. S. C., MA 02142, USA, Department of Medicine, U. S. D., La Jolla, CA

673 92093, USA, Moores Cancer Center, U. S. D., La Jolla, CA 92093, USA, Tamayo, P., Broad

674 Institute of MIT and Harvard, M. S. C., MA 02142, USA, Department of Medicine, U. S.

675 D., La Jolla, CA 92093, USA, & Moores Cancer Center, U. S. D., La Jolla, CA 92093,

676 USA. (2024). The Molecular Signatures Database (MSigDB) hallmark gene set collection.

677 *Cell systems*, *1*(6), 417. https://doi.org/10.1016/j.cels.2015.12.004

678 Luo, J., SB, D., JB, L., RQ, N., & JC, C. (2006). The aging neurogenic subventricular zone. *Aging*

679 *cell*, *5*(2). https://doi.org/10.1111/j.1474-9726.2006.00197.x

680 MI, L., W, H., & S, A. (2014). Moderated estimation of fold change and dispersion for RNA-seq

681 data with DESeq2. *Genome biology*, *15*(12). https://doi.org/10.1186/s13059-014-0550-8

682 Pluvinage, J. V., Haney, M. S., Smith, B. A. H., Sun, J., Iram, T., Bonanno, L., Li, L., Lee, D. P.,

683 Morgens, D. W., Yang, A. C., Shuken, S. R., Gate, D., Scott, M., Khatri, P., Luo, J.,

684 Bertozzi, C. R., Bassik, M. C., & Wyss-Coray, T. (2019). CD22 blockade restores

685 homeostatic microglial phagocytosis in ageing brains. *Nature*, *568*(7751), 187-192.

686 https://doi.org/10.1038/s41586-019-1088-4

687 Sierra, A., JM, E., JJ, D., JH, C., G, E., LS, O.-W., SE, T., & M, M.-S. (2010). Microglia shape

688 adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell stem cell*,

689 *7*(4). https://doi.org/10.1016/j.stem.2010.08.014

690 Solano, F., S, M., DM, A., R, R., A, D., A, C., J, O. C., & E, K. (2016). Neurogenic Niche Microglia

691 Undergo Positional Remodeling and Progressive Activation Contributing to Age-

692 Associated Reductions in Neurogenesis. *Stem cells and development*, *25*(7).

693 https://doi.org/10.1089/scd.2015.0319

694 Xavier, A. R., BT, K., SA, G., JR, L. d. M., & M, N. (2015). A Distinct Population of Microglia

695 Supports Adult Neurogenesis in the Subventricular Zone. *The Journal of neuroscience :*

the   official   journal   of   the   Society   for   Neuroscience,   35(34).

https://doi.org/10.1523/JNEUROSCI.1217-15.2015

Yen, J.-H. J., Department of Microbiology and Immunology, I. U. S. o. M., Indianapolis, IN, USA,

Yu, I.-C. I., & Department of Anatomy, C. B., and Physiology, Indiana University School

of Medicine, Fort Wayne, IN, USA. (2024). The role of ApoE-mediated microglial lipid

metabolism in brain aging and disease. *Immunometabolism (Cobham (Surrey, England)*,

5(1). https://doi.org/10.1097/IN9.0000000000000018

Supplementary Figure 1. Quality control, differential expression, and gene set enrichment analysis of major cell types in Dulken *et al*., 2019 aging SVZ scRNA dataset

A) Number of cells captured per sample. O=old (28-29 months old) and Y=young (3 months old), number indicates mouse replicate. B) Distribution of genes identified per cell in each sample. C) SVZ microglia pseudobulk differential gene expression of old vs. young. Red points are log₂ fold change > 1 and adjusted p-value < 0.05. Statistics: n=3 mice/group, DGE results obtained using Wald test followed by Benjamini-Hochberg correction for multiple hypothesis testing. D) SVZ microglia gene set enrichment analysis of gene ontology biological process for old vs. young differentially expressed genes shown in (C). Statistics: GSEA p-values obtained using a permutation test followed by followed by Benjamini-Hochberg correction for multiple hypothesis testing. E-F) Same as in B-C, but for astrocytes-quiescent neural stem cells (qNSCs). G-H) Same as in B-C, but for activate neural stem cells (aNSCs) – neural progenitor cells (NPCs). I-J) Same as in B-C, but for neuroblasts.

Supplementary Figure 2. Extended analysis of aging SVZ microglia gene expression changes

A-B) Heatmap of age-related differentially expressed genes (ARDEGs; adjusted p-value < 0.05) in aging SVZ microglia that overlapped with gene sets derived from screens to identify genes regulating phagocytosis from Haney *et al*., 2018 (A) and Pluvinage *et al*., 2019 (B). Statistics: n=3 mice/group C) Same as in (A), but for ARDEGs that overlapped with the Molecular Signatures Database MICROGLIAL CELL ACTIVATION or MICROGLIAL CELL PROLIFERATION gene sets (Supplementary Table 2). D) SVZ microglia pseudobulk differential gene expression of old vs. young where points highlighted in red are genes that are log$_2$ fold change > 1 and adjusted p-value < 0.05 and that overlapped with the Molecular Signatures Database entries: DESCARTES FETAL CEREBELLUM MICROGLIA, DESCARTES FETAL CEREBRUM MICROGLIA, DESCARTES MAIN FETAL MICROGLIA, FAN EMBRYONIC CTX BIG GROUPS MICROGLIA, FAN EMBRYONIC CTX MICROGLIA 1,FAN EMBRYONIC CTX MICROGLIA 2, FAN EMBRYONIC CTX MICROGLIA 3 gene sets (Supplementary Table 2). Statistics: n=3 mice/group. E) Same as in (A), but for ARDEGs that overlapped with the disease associated microglia (DAM) gene sets reviewed in Deczkowska *et al*., 2018 (Supplementary Table 2).

Supplementary Figure 3. Extended analysis of aging SVZ neural progenitor cell gene expression changes

A) Average Log$_2$ fold changes in gene expression changes for astrocytes-qNSC, aNSCs-NPCs, and neuroblasts for genes that overlapped with the Reactome Apoptosis gene set (Supplementary Table 2). Fold changes are averages calculated form pseudobulk differential expression analysis (Methods). Statistics: n=3 mice/group. B) Same as in (A), but for genes that overlapped with the Molecular Signatures Database HALLMARK COMPLEMENT gene set (Supplementary Table 2). C) Same as in (A), but for genes that overlapped with the Molecular Signatures Database GOBP OPSONIZATION gene set (Supplementary Table 2). D) Same as in (A), but for genes that overlapped with the curated DON'T EAT ME gene set (Supplementary Table 2). E) Same as in (A), but for genes that overlapped with the Molecular Signatures Database GOBP ANTIGEN PROCESSING AND PRESENTATION gene set (Supplementary Table 2).

Supplementary Figure 4. qPCR validation of genes selected from the scRNA-Seq results
(A-B) Quantification of gene expression of microglia from the SVZ (SVZ MG) or whole brain (WB MG) in young (3 months) and aged (12 months) mice. Statistics: n=3 mice/group; σ: not detected. P-value legend: ns: no symbol; *: p ≤ 0.05, **: p ≤ 0.01, ***: p ≤ 0.001, ****: p ≤ 0.0001.