Title

Age Associated Increase in Microglia Inflammation and Phagocytosis in the Adult Neural Stem
Cell Niche

Authors

Ronald Cutler[1, =], Samuel Harrison[1, =], Natalia Sandoval-Kuhn[1], Stephen Tomlinson[2], Erzsebet
Kokovay[1, *]

Affiliations

[1] UTH-SA, Department of Cell Systems and Anatomy, San Antonio, TX, USA 78229

[2] Medical University of South Carolina, Department of Pharmacology and Immunology,
Charleston, SC, USA 29425

[=] Authors contributed equally to this work

[*] Correspondence to E.K. (KokovayE@uthscsa.edu)

Number of pages = 38

Number of figures: Main = 4, Supplementary = 5

Number of tables: Main = 0, Supplementary = 6

Number of words: Abstract = 137, Introduction = 569, Discussion = 1,272

Conflicts of interest: The authors declare no competing financial interests.

Acknowledgements: This work was supported by funding from the NIH under grants
R01NS102448 and T32 AG0201890

Author contributions: E.K. and R.C. designed the research; R.C., S.H., and N.S.K performed
the research; S.T. contributed unpublished reagents/analytic tools; R.C. and S.H. analyzed data;
R.C. and E.K. wrote the paper.

Abstract

Adult SVZ neurogenesis declines with age, but the niche mechanisms remain incompletely defined. Here, we show that SVZ microglia acquire activation-associated morphology with aging in male and female C57BL/6 mice. Reanalysis of published SVZ single-cell RNA-Seq data revealed that aged microglia upregulate inflammatory and lysosome/phagosome-related gene programs, including multiple phagocytosis-associated receptors and effectors. Functionally, young SVZ-derived microglia exhibit reduced basal phagocytic activity compared with non-SVZ microglia *in vitro*, whereas aged SVZ microglia show increased phagocytosis. *In vivo* SVZ wholemounts reveal increased progenitor-derived material within lysosomal compartments of microglia with age, without an accompanying increase in apoptosis. Inhibiting complement C3 activation partially restores SVZ proliferation in aged mice and coincides with a reduction of progenitor-derived material within lysosomal compartments of aged microglia. These data support a model in which microglial inflammatory signaling and engagement with neural progenitors contributes to age-related suppression of SVZ proliferation.

Significance Statement

Adult neural stem cells support plasticity and may aid repair after brain injury, but this capacity declines with age. We show that microglia in the aging subventricular zone acquire inflammatory and phagocytic programs, increase their lysosomal association with progenitor-derived material, and are linked to reduced progenitor proliferation. Complement inhibition partially restores proliferative output in aged mice, connecting inflammatory signaling to this niche defect. These findings identify age-dependent microglia–progenitor interactions as a mechanism that may limit neural stem cell activity in the adult brain and suggest immune pathways as possible targets for preserving or restoring neurogenic potential during aging.

Introduction

Adult neural stem cells (NSCs) reside in two principal neurogenic niches of the rodent brain: the subgranular zone (SGZ) of the dentate gyrus and the subventricular zone (SVZ) lining the lateral ventricles. In the current study, we are focused on the NSCs in the SVZ niche, which exist in quiescent (qNSCs) and activated (aNSCs) states and generate transit amplifying cells (TACs), that expand and then produce migratory neuroblasts (Alvarez-Buylla & García-Verdugo, 2002; Doetsch et al., 1999; Lim & Alvarez-Buylla, 2016). These neuroblasts migrate from the SVZ through the RMS into the OB where they differentiate and integrate into local circuitry (Alvarez-Buylla & García-Verdugo, 2002). With aging, multiple studies report reduced SVZ neurogenesis accompanied by decreased NSC proliferation, increased NSC quiescence, and diminished NSC pool size (Ahlenius et al., 2025; Cutler & Kokovay, 2020; Enwere et al., 2004; Kalamakis et al., 2019; Maslov et al., 2004; Solano et al., 2016). These findings motivate work aimed at identifying niche-derived mechanisms that constrain activation and lineage progression in the aged SVZ.

Microglia are brain-resident macrophage-lineage cells that survey the parenchyma and shape neural homeostasis through debris clearance, phagocytosis, synapse remodeling, and secretion of signaling factors (Deczkowska et al., 2018). Fate-mapping indicates that microglia originate from early embryonic yolk sac–derived myeloid progenitors that seed the central nervous system and largely self-maintain thereafter (Ginhoux et al., 2010). Microglia constitute a substantial fraction of brain cells (commonly estimated at ~5–15% depending on region and quantification) and display pronounced context-dependent states across development, aging, and disease (Deczkowska et al., 2018).

Several studies suggest that microglia within the SVZ are not interchangeable with microglia in

non-neurogenic parenchyma. SVZ-associated microglia exhibit distinct morphological and

antigenic features and have been implicated in regulating neurogenesis and progenitor dynamics

(Goings et al., 2006; Marshall et al., 2014; Shigemoto-Mogami et al., 2014; Solano et al., 2016;

Xavier et al., 2015). Aging is accompanied by microglial state changes in the SVZ, including

increased inflammatory and lysosomal programs that precedes the decline of NSC proliferation,

raising the possibility that altered microglia–NSC interactions contribute to reduced neurogenesis

in the aged SVZ (Solano et al., 2016). Intriguingly, these age-related shifts in microglia towards

an activated state are reminiscent of their phenotype in the developing cerebral cortex and SVZ

where they have been shown to phagocytose neural precursor cells (Cunningham et al., 2013;

Morrison et al., 2023). Furthermore, in a model of cortical ischemic stroke, microglia were

reported to phagocytose dying neuroblasts, constraining the endogenous neurogenic response after

injury (Nath et al., 2024). Despite evidence that niche inflammation can constrain activation and

lineage output, it remains unclear whether aging-associated microglial activation in the SVZ is

coupled to changes in microglial phagocytic behavior toward specific NSC populations, and

whether inflammatory signaling and phagocytic engagement represent linked or separable

mechanisms.

In this study, we test the hypothesis that aging alters SVZ microglial state and microglia–NSC

interactions in ways that contribute to diminished progenitor proliferation. We found that during

aging, SVZ microglia are characterized by increases in a lysosomal marker and enlarged

morphologies, alongside upregulation of pro-inflammatory and phagocytic gene expression

signatures, consistent with a pro-phagocytic phenotype. Aged SVZ microglia show increased

phagocytosis *in vitro* and *in vivo*, where we found evidence for the engulfment of aNSCs/TACs.

Finally, inhibition of complement C3 activation partially restored proliferative measures and

reduced aNSC/TAC engulfment, suggesting that microglial inflammatory signaling and progenitor

phagocytosis contributes to reduced NSC proliferation during aging.

Materials and Methods

Animals and Interventions

Male and female C57BL/6 mice were used in this study either bred at the UTHSA vivarium or acquired from the National Institute of Aging. Young mice (3 months) and aged mice (12 months) were housed in accordance with the University of Texas Health San Antonio Institutional Animal Care and Use Committee and performed in accordance with institutional and federal guidelines and allowed free access to water and standard rodent chow ad libitum.

Both male and female C57BL/6 mice received CR2-crry (10µg/1g mouse; Stephen Tomlinson Lab, Medical University of South Carolina) with saline controls via intraperitoneal injection. CR2-crry or saline was administered every other day for a total of 3 doses.

Immunohistochemistry

The SVZ was isolated for wholemount (WM) tissue analysis; mice were anesthetized with isoflurane, a cervical dislocation was performed, followed by brain extraction. Brains were placed into cold HIB where they were halved and the septum and hippocampus were removed, and the SVZ was microdissected from the surrounding brain tissue. The SVZ WM was then fixed in fresh 4% paraformaldehyde (PFA) for 30 minutes. SVZ WMs were permeabilized in 2% PBST and blocked overnight at 4° C in 10% normal donkey serum (NDS). The SVZ WMs were then incubated at 4°C for 4 days with primary antibody in blocking solution, followed by overnight incubation at 4°C with secondary antibody made in blocking solution, and a 5-minute incubation at room temperature (RT) with DAPI (1:1000; Invitrogen, Cat. No. D1306) then mounted on a

slide in ProLong Gold antifade reagent (Invitrogen, Cat. No. P36930). Primary antibodies that

were used include Iba1 (1:200; Wako, Cat. No. 019-19741 and 1:200; Novus, Cat. No. NB100-

1028), CD68 (1:250; BioRad, Cat. No. MCA1957), and Mash1 (1:5000; CosmoBioUSA, Cat. No.

SK-T01-003) Cleaved caspase 3 (1:200 Cell Signaling Cat. No D175 5A1E). Species-specific

secondary antibodies (Jackson ImmunoResearch) were used at a concentration of 1:500 in SVZ

WMs.

Immunohistochemistry Quantification

EdU was quantified in the SVZ by making a "Z Project" 2-dimensional image of the Z-stack in

ImageJ on 10 random fields of view and hand counting the number of EdU+ nuclei. The number

of microglia were quantified by hand counting Iba1+ cells, excluding cells whose cell body

touched the image boundary on Zen Microscopy Software (Zeiss). Total CD68 area was quantified

by making a "Z project" 2-dimensional image of the Z-stack, using the "Threshold" function, and

then the "Analyze particle" function to give total area in μm2 that was then averaged by total

number of microglia in each image. Total number of Iba1+ microglia were quantified similarly to

EdU+ cells described above. Mash1 colocalization within Iba1+ microglia were quantified by hand

counting in Zen Microscopy Software (Zeiss) by examining each Iba1+ microglia's lysosome

(CD68+) to determine if each Iba1+CD68+ microglia contained Mash1 colocalization. The

number of Mash1+CD68+Iba1+ microglia were then averaged to total number of Iba1+ microglia

in each image. There were no microglia that contained Mash1 that were not CD68+. SVZ WMs

were analyzed from 15μm thick Z-stacks (15 Z-stacks per WM). Confocal images were obtained

with a Zeiss LSM 710 confocal microscope at 40x objective.

Single-cell RNA-sequencing quality control and filtering

Quality control of downloaded single-cell RNA-Seq data (see Data Availability) was first performed, which filtered cells where the number of genes or UMIs detected were greater or less than 5 median absolute deviations (MADs) or percent mitochondria reads greater or less than 3 MADs, relative to all cells in the dataset. Genes were filtered if detected in less than 10 cells out of all cells in the dataset. Cell identity labels were those assigned by the original analysis.

Single-cell RNA-sequencing differential expression and gene set enrichment analysis

Pseudobulk samples were created by summing counts of each identified cell type for each biological replicate. The DESeq2 package (Love et al., 2014) was then used to perform differential expression analysis to compare old vs. young groups. Following this the ClusterProfiler package (Yu et al., 2012) was then used to perform gene set enrichment analysis using the gene ontology biological process database or custom annotations.

Microglia phagocytosis gene set analysis

Genes associated with known phagocytosis receptors and/or pathways active in microglia were manually curated by searching the Molecular Signatures Database (MsigDB) and the literature (i.e. Complement, Toll-like receptor, TAM receptor, Trem2 receptor, FC gamma, Pr2y6 receptor,

Scavenger receptor, Mannose receptor, LRP1 receptor, Galectin-3 adaptor protein, Microglia

Activation, Microglia Proliferation, Disease-Associated Microglia (DAM), and Microglia

Development). In addition, gene sets related to screens of phagocytosis regulators was obtained

from the literature (see Data Availability). These gene sets were then overlapped with the genes

that were found to be significantly differentially expressed (adjusted p-value < 0.05) in the

microglia old vs. young comparison.

Neural stem cell gene set analysis

Genes associated with pathways known to be expressed in cells that are targeted for phagocytosis

were manually curated by searching the Molecular Signatures Database (MsigDB) and the

literature (i.e. Apoptosis, Opsonization, Antigen Processing and Presentation, "eat-me," "find-

me," and "don't eat me"). Because of the lack of genes which were significantly differentially

expressed in neural stem cells (i.e. Astrocytes/qNSCs, aNSCs/NPCs and Neuroblasts) in the old

vs. young comparison, the ClusterProfiler package was then used to perform gene set enrichment

analysis using the manually curated gene sets. The Log$_2$ fold change of genes within the gene sets

found to be significantly enriched in age-related genes expression of neural stem cells.

Microglia Isolation and Cell Culture

SVZ microglia were derived by mixed-glial expansion whereas whole brain microglia were

MACS-isolated. C57BL/6 mice were injected intraperitoneally with pentobarbital (25µL/mouse)

then perfused with cold 1x PBS. SVZ WMs or the remainder of the whole brain excluding the

SVZ and cerebellum, hereby referred to as whole brain (WB), were isolated following perfusion.

The tissues were minced and dissociated in 50U papain in DMEM/F12 at 37°C for 45 minutes and

manually triturated and spun 3x in DMEM. SVZ and WB homogenate were gently pressed through

a 70μm cell strainer to remove debris. For mixed glial culture, SVZ homogenate were then plated

in a T25 flask coated with poly-L-ornithine (PLO) in neural growth media (DMEM/F12, 5% fetal

bovine serum, 1x N2, 10ng/mL EGF, and 10ng/mL bFGF) until a confluent cell monolayer formed.

The culture was then fed microglia proliferation media (DMEM/F12, 10% fetal bovine serum, 1x

N2, and 20 ng/mL GM-CSF) until microglia could be seen atop the monolayer. Once microglia

were present, they were shaken off from the cell monolayer and the SVZ microglia could be plated

for further experimentation.

For microglia isolated by MACS, following the cell strainer step, WB homogenate was

resuspended in HIB and centrifugated at 400G with no brake in 22% Percoll (Sigma-aldrich, Cat.

No. P4937) solution to remove myelin from the samples. The WB homogenate was then

resuspended in 0.5% bovine serum albumin and incubated with 10μL CD11b Microbeads (cat#

130-093-636) per 107 cells. The homogenate was then placed in an LS column (MiltenyiBiotec,

Cat. No. 130-042-401) in a QuadroMACS™ separator (MiltenyiBiotec, Cat. No. 130-090-976) to

allow for MACS sorting of CD11b + microglia following manufacturer protocols

(MiltenyiBiotec). WB microglia were plated in a 6-well plate coated with PLO/laminin (10μg/mL;

Sigma-aldrich, Cat. No. L2020) at 250k cells/well in microglia expansion media (DMEM/F12,

100 μg/mL PenStrep, 5 μg/mL n-acetyl-L-cysteine, 5 μg/mL Insulin, 100 μg/mL apo-transferrin,

100 μg/mL sodium selenite, 1 μg/mL heparin sulfate, 10% fetal bovine serum, 1.5 μg/mL

cholesterol, 40 ng/mL M-CSF, 20 ng/mL GM-CSF, and 2 ng/mL TGF-β) adapted from [Bohlen et

al., 2017].

Microglia *in vitro* Phagocytosis Assay

The phagocytosis assay was performed using live cell imaging by Incucyte S3 (Essen BioScience)

for WB or SVZ microglia phagocytosis from 3- or 12-month mice. WB microglia were passaged

from a 6-well plate with Trypsin (Fisher, Cat. No. 25-200-072) following MACS isolation and

SVZ microglia were collected from mixed glial culture and microglia were plated in a 96 well

plate coated with PLO/laminin at 20k cells/well in microglia expansion media for 3 days. After 3

days, SVZ and WB microglia were moved to serum free media (DMEM/F12, 100 µg/mL PenStrep,

5 µg/mL n-acetyl-L-cysteine, 5 µg/mL Insulin, 100 µg/mL apo-transferrin, 100 µg/mL sodium

selenite, 1 µg/mL heparin sulfate, 1.5 µg/mL cholesterol, 40 ng/mL M-CSF, and 2 ng/mL TGF-β)

adapted from [Bohlen et al., 2017] for 1 week. All microglia were then incubated with 5 µg/mL

Isolectin IB4 Alexa Fluor™ 488 Conjugate (Invitrogen, Cat. No. I21411) for 60 minutes at 37°C.

SVZ or WB microglia of both ages were treated with Zymosan S. cerevisiae BioParticles™ (10

mg/mL; Invitrogen, Cat. No. Z2849) conjugated pHrodo™ Red SE (Molecular Probes™, Cat. No.

P36600) following manufacturer protocols (ThermoFisher) and adapted from [Haney et al., 2018;

Pluvinage et al., 2019], both Zymosan-pHrodo and Cytochalasin D (10 mM/µL Sigma-aldrich,

Cat. No. C8273) to inhibit phagocytosis, or isolectin only as a negative control. The microglia

were then incubated in the Incucyte S3 with fluorescent images captured every hour over the

course of 48 hours for live cell imaging of microglia phagocytosis. The images were analyzed in

the IncuCyte software (Essen BioScience) quantifying colocalization of the green fluorescence of

Isolectin IB4 labelled microglia and the red fluorescence of engulfed Zymosan-pHrodo within the

microglia lysosome demonstrating microglia phagocytosis. Total pHrodo area / number of

microglia was calculated for each well and condition yielding amount of in vitro phagocytosis

occurring at each time point.

RNA Extraction and qPCR

RNA was extracted from freshly isolated microglia following magnetic-activated cell sorting

(MACS) isolation, using a TRIzol-chloroform method (Rio et al., 2010) where cells were lysed in

TRIzol (Invitrogen, Cat. No. 15596018), aqueous layer separated with chloroform, RNA was

precipitated with isopropanol, washed in ethanol, dried, then resuspended in RNase-free water.

cDNA was synthesized using High Capacity cDNA Reverse Transcription Kit (Applied

Biosystems, Cat. No. 4368814). Quantitative polymerase chain reaction (qPCR) was performed

using a C1000 Touch Thermal Cycler (BioRad) with iTaq™ Universal SYBR® Green Supermix

(BioRad, Cat. No. 1725121). Reactions were performed in doublet with glyceraldehyde-3-

phosphate dehydrogenase (GAPDH) used as an endogenous control. Negative control reactions

were performed using the same conditions without cDNA. Quantification was analyzed using the

ΔΔCt method as described [(Livak & Schmittgen, 2001)]. Primers were used for the following

transcripts: C1QA, P2Y12, TREM2, LPAR, C3, CR1, ITGAM, ITGAX, GAPDH.

Primers	Forward	Reverse
C1QA	AAAGGCAATCCAGGCAATATCA	TGGTTCTGGTATGGACTCTCC
C3	CCAGCTCCCCATTAGCTCTG	GCACTTGCCTCTTTAGGAAGTC
TREM2	CTGGAACCGTCACCATCACTC	CGAAACTCGATGACTCCTCGG
LPAR	AGCCATGAACGAACAACAGTG	CATGATGAACACGCAAACAGTG

CR1	ATGGAGGTCTCTTCTCGGAGT	GGCCGAAGGCTACAAGGAG
ITGAM	GGGAGGACAAAAACTGCCTCA	ACAACTAGGATCTTCGCAGCAT
ITGAX	CTGGATAGCCTTTCTTCTGCTG	GCACACTGTGTCCGAACTCA
GAPDH	AGGTCGGTGTGAACGGATTTG	TGTAGACCATGTAGTTGAGGTCA

EdU Labeling and Staining

275 Mice were injected intraperitoneally (50μg EdU/1g mouse) 2 hours before sacrifice. Their brains

276 were removed and SVZ was isolated and fixed for 30 minutes in 4% PFA. EdU staining was

277 performed following manufacturer protocols (Fisher, Click-it EdU Imaging Kit, Cat. No. C10337).

Experimental Design and Statistical Analysis

280 Student's t-tests and two-way ANOVA were used to analyze the data on GraphPad Prism 8

281 Software. All data are expressed as mean ± standard error of the mean.

Data Accessibility

Normalized single-cell RNA-sequencing data and cell type annotations of the SVZ from young (3

months old) and old (28–29 months old) male mice (n = 3) was downloaded from Supplementary

Table 2 of the Dulken et al., 2019 study. Results of screen for phagocytosis regulator genes were

downloaded from Supplementary Table 5 of the Haney et al., 2019 study and Supplementary Table

2 of the Pluvinage et al., 2019 study. Marker genes of major cell types in the SVZ of young (2-4

months old) male and female mice were downloaded from Table S1 of the Zywitza et al., 2018

study.

Code Accessibility

Code to reproduce single-cell RNA-sequencing data analyses is available at the following

repository: https://github.com/cutleraging/nsc-aging

Results

Age-related changes in SVZ microglia morphology

We have previously reported that murine SVZ neurogenesis is reduced by 12 months of age (Solano et al., 2016). To assess how SVZ microglia change during aging, we performed immunohistochemistry on SVZ wholemount preparations (to preserve 3D niche cytoarchitecture) from young (3 month) and aged (12 month) mice, staining for the microglial marker Iba1 and the lysosomal marker CD68. In young mice, Iba1+ microglia exhibited a ramified morphology throughout the SVZ with relatively low CD68 signal within Iba1+ cells (Fig. 1A-B). In aged mice, Iba1+ microglia displayed morphology consistent with activation, including fewer processes and enlarged somata, together with increased CD68 signal within Iba1+ cells (Fig. 1C-D). Because microglia can adopt a lysosome-rich, phagocytic phenotype during neuronal development that is associated with engulfment of neural precursor cells, we hypothesized that age-associated microglial activation in the SVZ is accompanied by increased phagocytic activity toward SVZ progenitors (Cunningham et al., 2013).

Figure 1. Age-Associated SVZ Microglia Morphological Changes and Increased Activation. Representative confocal projection images of SVZ microglia stained with Iba1 (green), lysosomes stained with CD68 (red), and nuclei stained with DAPI (blue). **(A-B)** Young (3 month) SVZ microglia imaged at 40X magnification (A) or 100X magnification (B). **(C-D)** Aged (12 month) SVZ microglia imaged at 40X magnification (C) or 100X magnification (D). Scale bars in (A) and (C) = 20μm, scale bars in (B) and (D) = 50μm

Gene expression in the aging SVZ suggests increased microglial phagocytic programs

To assess age-related transcriptional changes in SVZ microglia and NSC populations relevant to inflammation and phagocytosis, we reanalyzed a published single-cell RNA-Seq dataset comprising of 14,685 cells from the SVZ of young (3 month) and old (28/29 month) male mice (Dulken et al., 2019). We generated pseudobulk profiles for each cell type and biological replicate, performed differential gene expression analysis (old vs. young), followed with gene set enrichment analysis (GSEA) (Supplementary Figs. 1A-B).

In microglia, we identified 254 genes significantly upregulated and 54 genes significantly downregulated with age (adjusted p-value < 0.05; Supplementary Fig. 1C; Supplementary Table 1). GSEA of genes upregulated with aging revealed enrichment of processes related to translation, secretion of inflammatory cytokines, and lipid metabolism, consistent with prior analyses (Dulken et al., 2019; Yen & Yu, 2024). In contrast, downregulated genes were enriched for cytoskeleton organization terms, which has not been previously reported and may suggest dysfunction in motility or morphology-related programs (Supplementary Fig. 1D).

Although canonical "phagocytosis" terms were not significantly enriched by GSEA

(Supplementary Fig. 1D), several phagocytosis-associated receptors, including *Axl* and *Tlr2*, were

among the most strongly upregulated transcripts with age (Supplementary Fig. 1C). We therefore

curated comprehensive gene sets representing multiple components of phagocytosis from MSigDB

(Liberzon et al., 2024) and intersected these with age-related differentially expressed genes

(ARDEGs; adjusted p-value < 0.05; Supplementary Table 2). 19 phagocytosis-related genes

overlapped with the microglia ARDEG set, of which 18 were upregulated in aged microglia (Fig.

2A; Supplementary Table 3). Notably, 8 of these genes (*Cyba, Atp6v0e, Rab7b, Cybb, Vim, B2m,*

Tlr2, and *Hvcn1*) are linked to phagosome formation/maturation and vesicular acidification. We

further expanded our phagocytosis gene list to include genes identified in prior functional screens

(Haney et al., 2018; Pluvinage et al., 2019), and found that all intersecting genes (9/9) were

upregulated in aged microglia (Supplementary Figs. 2A-B). Finally, we found that the broad

upregulation of phagocytosis-related gene expression was also accompanied by an increase in the

expression of genes related to microglia activation (Supplementary Figs. 2C-D). While these

transcriptional shifts are consistent with increased lysosomal/phagocytic programs, we note that

elevated expression of such genes could also reflect compensatory responses to altered phagocytic

demand or efficiency rather than a direct readout of increased engulfment per se. For instance, the

expression of the receptor essential for microglial function, *Csf1r*, was downregulated in aged

microglia while the ligand *Csf1* showed the opposite trend and thus may be a compensatory

response to chronic microglia activation (Supplementary Fig. 2C) (Hu et al., 2024). Together, these

findings suggest that aging is accompanied by transcriptional reprogramming of SVZ microglia

toward states featuring inflammatory signaling and lysosome/phagosome-related programs, with

partial resemblance to developmental and phagocytic microglial states (Cunningham et al., 2013;

Izgi et al., 2022).

We next asked whether age-associated changes were concentrated within particular receptor-

mediated phagocytosis pathways. Using a curated list for the major phagocytosis pathways, we

observed upregulation of the receptors in the Complement, TAM, and Toll-like pathways (Fig. 2B;

Supplementary Table 3). In contrast, for the FC gamma, Scavenger, P2ry6, and Trem2 receptor

pathways, multiple pathway components were consistently upregulated without a corresponding

increase in the core receptor transcript (Deczkowska et al., 2018). Notably, while Trem2 itself was

not increased, aged microglia displayed expression patterns consistent with a disease-associated

microglia (DAM) stage 2 signature that has been reported to be Trem2-dependent in other contexts

(Supplementary Fig. 2E). Overall, these results indicate broad multi-pathway upregulation of

phagocytosis-related gene expression during aging and suggest that aged SVZ microglia may

adopt a unique Trem2-independent DAM-like state.

We then analyzed NSC populations for age-associated expression changes in signals relevant to

target-cell recognition and engulfment, including apoptosis, "find-me," "eat-me," and "don't eat-

me" programs. Compared to microglia, there were very few ARDEGs: 13 genes were significantly

upregulated in aged astrocyte-qNSC as compared to young (adjusted p-value < 0.05), whereas no

genes met this threshold for aged aNSC, TACs or neuroblasts (Supplementary Figs. 1E-J). Given

this limited power for threshold-based overlap analyses, we used GSEA with curated gene sets for

apoptosis and phagocytosis-related signaling (Supplementary Table 2).

Using this approach, we detected age-associated upregulation of an apoptosis gene set in astrocyte-

qNSC cells (adjusted p-value < 0.05), whereas aNSC, TACs and neuroblasts showed the opposite

trend (Fig. 2C; Supplementary Fig. 3A). We did not detect significant enrichment for curated "find-

me" or generic "eat-me" gene sets. However, we observed age-associated upregulation of

complement system and opsonization gene sets (often considered "eat-me"–adjacent signals) in

astrocyte-qNSCs and neuroblasts, but not in aNSCs or TACs (Fig. 2C; Supplementary Fig. 3B,C)

(Li, 2012). Intriguingly, astrocyte-qNSCs and neuroblasts also showed increased expression of a

curated "don't eat-me" gene set (Fig. 2C; Supplementary Fig. 3D). This set includes genes related

to MHC class I antigen presentation such as *H2-K1* and *H2-D1*, which in some contexts can oppose

macrophage phagocytosis, particularly when expressed with *B2m* (Supplementary Fig. 3D)

(Khalaji et al., 2023). Consistent with a potential receptor–ligand axis, *Lilrb4a*, an inhibitory

receptor for MHC class I, was among the age-upregulated microglial ARDEGs (Supplementary

Fig. 1C). An antigen processing/presentation gene set showed a similar pattern to the "don't eat-

me" set, further supporting increased MHC class I–associated programs in aged astrocyte-qNSCs

and neuroblasts, but not aNSC-NPCs (Supplementary Fig. 1E).

In summary, transcriptomic changes in SVZ microglia with age are consistent with increased

inflammatory and lysosome/phagosome-related programs across multiple receptor pathways. In

aged NSC populations, astrocyte-qNSCs and neuroblasts showed increased complement pathway

components ("eat-me" signals) together with increased expression of putative "don't eat-me" anti-

phagocytic signals, while aNSC-NPCs did not show comparable upregulation of those signals.

These patterns are consistent with a model in which age-associated microglial phagocytic

programs may preferentially engage aNSC/TACs.

Figure 2. Gene expression in the aging SVZ suggests increased microglial phagocytic programs

(A) Pseudobulk level analysis of differential gene expression changes during aging of young (3 month) and aged (28–29 month) microglia in the SVZ. Red points indicate genes that overlapped with the phagocytosis-related gene sets (Supplementary table 2) which have absolute log2 fold change > 1 and adjusted p-value < 0.05. Statistics: n=3 mice/group, DGE results obtained using Wald test followed by Benjamini-Hochberg correction for multiple hypothesis testing. **(B)** Heatmap of age-related DEGs (adjusted p-value < 0.05) in aging SVZ microglia (obtained from analysis in (A)) that overlapped with phagocytosis-related gene sets (Supplementary table 2)

Cutler, Harrison, et al.

419 Statistics: n=3 mice/group. **(C)** Gene set enrichment analysis of age-related gene expression

420 changes in neural progenitor cells using apoptosis, "find-me," "eat-me," and "don't eat-me" signal

421 gene sets (Supplementary table 2). Statistics: n=3 mice/group, GSEA p-values obtained using a

422 permutation test followed by Benjamini-Hochberg correction for multiple hypothesis testing.

424 SVZ microglia show an age-associated increase in phagocytic activity toward

425 SVZ progenitors

To assess SVZ microglia phagocytic activity, we microdissected the SVZ and isolated microglia

either from the SVZ or from the remainder of the brain (excluding the SVZ) from young (3 month)

and aged (12 month) mice. Isolated microglia were expanded *in vitro*, and phagocytosis was

quantified by live imaging of engulfment of pHrodo-conjugated yeast particles (pH-sensitive

reporter that fluoresces upon acidification in phagolysosomes) (Kapellos et al., 2016). In young

animals, SVZ-derived microglia displayed lower phagocytic activity than microglia isolated from

non-SVZ brain regions under our assay conditions, consistent with niche-associated specialization

toward reduced basal phagocytosis (Fig. 3A) (Xavier et al., 2015). In aged animals, SVZ-derived

microglia showed a significant increase in phagocytic activity relative to young SVZ microglia,

approaching the levels observed in whole-brain microglia.

We next sought *in vivo* evidence that the age-associated increase in SVZ microglial phenotypes

reflects increased phagocytic engagement with SVZ progenitors. As an initial test, we quantified

apoptosis in the SVZ by immunohistochemistry for cleaved caspase-3 on coronal sections from

young (3 month) and aged (12 month) mice and did not detect a significant age-associated increase

in apoptotic cells in the SVZ (Fig. 3B). Next, we sought to validate the findings of an age-related

Cutler, Harrison, et al.   23

increase in SVZ microglia phagocytosis activity in our *in vitro* system by examining SVZ

microglia phagocytosis *in vivo*. We assessed microglial phagolysosomal content directly by

staining SVZ wholemounts for Iba1 (microglia), CD68 (lysosomes), and Mash1 (aNSC/TAC

specific transcription factor) (Kim et al., 2011; Lim & Alvarez-Buylla, 2016). In both young and

aged SVZ, we observed Mash1 signal within CD68+ compartments inside Iba1+ microglia (Fig.

3D). Quantification indicated that ~20% of SVZ microglia contained Mash1+ material within

CD68+ compartments in young mice, and this fraction increased over 2-fold at 12 months of age

(Fig. 3C). Together, these in vitro and in vivo data support an age-associated increase in SVZ

microglial phagocytic activity and increased microglial phagolysosomal association with

progenitor-derived material in the aged SVZ.

Figure 3. SVZ microglia show an age-associated increase in phagocytic activity toward SVZ progenitors. (A) Time-lapse *in vitro* phagocytosis assay of young (3 months) or aged (12 months) SVZ or whole brain (WB) microglia labeled with isolectin and incubated with Zymosan-pHrodo (Zymo), CytochalasinD and Zymosan-pHrodo (CytoD, negative control), or isolectin only (Control). Phagocytosis is quantified as pHrodo+ area indicated by red fluorescence / microglia number. Statistics: n=3 mice/group, ≥ 1,000 microglia quantified per mouse. (B) Quantification of apoptosis in SVZ whole mounts. Statistics: n=6 mice/group, p-value estimated using two-way ANOVA. (C) Quantification of *in vivo* phagocytosis of 3- or 12-month SVZ microglia measured

as Iba1+CD68+Mash1+ microglia / total Iba1+ microglia (Methods). *n*=11 mice/group, ≥100

microglia quantified per mouse. (D) Representative confocal projection images of young or aged

SVZ microglia stained with Iba1 (green), microglia lysosomes stained with CD68 (red), and TAC

transcription factor stained with Mash1 (magenta). Scale bar: 10µm, p-value estimated using two-

way ANOVA. P-value legend: ns: no symbol; ****: p ≤ 0.0001.

Complement inhibition partially rescues SVZ proliferation in aged mice

To explore possible mechanisms that might contribute to the age-associated increase in SVZ

microglial phagocytic engagement of aNSC/TAC populations in the SVZ, we examined expression

of candidate pro-inflammatory factors in SVZ microglia guided by our transcriptomic reanalysis

(Fig. 2A). We reasoned that pro-inflammatory genes that were low or absent in the young SVZ

microglia relative to non-SVZ microglia that are increased with age in SVZ microglia would be

good candidates based on our *in vitro* phagocytosis results (Fig. 3A). Among the candidates, *C3*

(Complement component 3) showed higher expression in non-SVZ microglia relative to young

SVZ microglia and was strongly upregulated with age in SVZ microglia (Supplementary Fig. 4).

In addition, transcripts encoding *C3* receptors *Itgam* and *Itgax* were increased in aged SVZ

microglia. Based on these observations, we tested whether inhibiting *C3* activation would alter

SVZ proliferation and/or microglial phagocytic readouts *in vivo*.

To inhibit C3 cleavage and deposition, we used CR2-Crry, a targeted complement inhibitor that

crosses the blood brain barrier and localizes to sites of complement activation via CR2 to then

block C3 activation through Crry (Atkinson et al., 2005; Song et al., 2003). Young (3 month) and

aged (12 month) mice received CR2-Crry or saline every other day for three total doses, followed

by a 2-hour EdU pulse prior to sacrifice to label proliferating cells (Fig. 4A). SVZ wholemounts

were collected and processed for immunohistochemistry as described above (Methods).

In young mice, CR2-Crry treatment did not significantly affect SVZ proliferation as indicated by

EdU+ cells, microglial CD68 area, or the fraction of microglia containing Mash1+ material within

CD68+ compartments (Fig. 4C). As expected, saline-treated aged mice showed reduced SVZ

proliferation relative to saline-treated young mice (Fig. 4B). In aged mice, CR2-Crry significantly

increased SVZ proliferation and Mash1+ cells compared with age-matched saline controls as

indicated by EdU+ cells (Fig. 4B; Supplementary Fig. 5). CD68 area per Iba1+ microglia was

increased in aged saline-treated mice relative to young controls, while CR2-Crry produced a non-

significant reduction in this measure in aged mice (Fig. 4D). Similarly, the microglia containing

Mash1+ material within CD68+ compartments increased with age, and CR2-Crry treatment

resulted in a non-significant reduction relative to aged saline controls (Fig. 4E). Together, these

data indicate that complement inhibition via CR2-Crry can partially restore proliferative output in

the aged SVZ.

Figure 4. Complement inhibition partially rescues SVZ proliferation in aged mice (A) Graphic representation of CR2-crry or saline administration to young (3 month) or aged (12 month) mice. **(B)** Quantification of proliferating EdU-positive cells in the SVZ of young or aged control or CR2-crry treated mice. Statistics: *n*=4 mice/group, ≥ 100 image fields quantified per mouse, p-values estimated using two-way ANOVA. **(C)** Confocal projection images of young control, aged control, aged CR2-crry treated SVZ microglia stained with Iba1 (green), microglia lysosomes stained with CD68 (red), and aNSC/TAC transcription factor stained with Mash1 (magenta). Arrowheads in aged control and aged CR2-crry treated indicate colocalization of Mash1+ signal within microglia lysosomes (Iba1+CD68+). Scale bar: 10μm. **(D)** Quantification of microglia activation measured by CD68 area / Iba1+ cells. Statistics: *n*=4 mice/group, ≥ 1,000 microglia quantified per mouse, p-values estimated using two-way ANOVA. **(E)** Quantification of Mash1 containing microglia (Iba1+CD68+Mash1+ / Iba1+ cells / Mash1+ cells) from young or aged control or CR2-crry treated mice. Statistics: Same as in (D). P-value legend: ns: no symbol; *: p ≤ 0.05, **: p ≤ 0.01, ***: p ≤ 0.001, ****: p ≤ 0.0001.

Discussion

Adult SVZ NSCs generate new neurons that migrate to the OB and contribute to olfactory learning and discrimination throughout life (Lledo & Valley, 2016). In addition, following brain injury SVZ neural progenitors and neuroblast undergo a proliferative burst and then migrate to damaged areas where they give rise to a limited number of neurons, astrocytes and oligodendrocytes and can also secrete pro-reparative factors (Liang et al., 2019; Robin et al., 2006; Yamashita et al., 2006). The blockade of this response results in larger lesions and worsens behavioral deficits (Ohab et al., 2006). Together, these observations support the SVZ as a candidate target for therapies aimed at enhancing endogenous repair. However, SVZ neurogenesis declines drastically in aged rodents, with the most-rapid reductions between midlife (9-12 months) and persists for the duration of aging (Luo et al., 2006). This decline coincides with increasing risk of brain injury with age, motivating a need to identify niche mechanisms that constrain SVZ stem/progenitor function. Our data support a model in which aging is accompanied by increased microglial inflammatory and phagocytic programs within the SVZ niche, together contributing to reduced stem/progenitor output during aging.

During embryonic neurogenesis, SVZ-associated microglia display amoeboid morphology and can regulate precursor pool size through phagocytosis of non-apoptotic neuroblasts (Cunningham et al., 2013; Morrison et al., 2023). In the adult dentate gyrus, another neurogenic niche, microglia phagocytose apoptotic newborn neurons as part of routine homeostasis, whereas apoptosis within the adult SVZ is generally reported to be low (Luo et al., 2006). In the OB, where many SVZ-derived neurons fail to integrate, phagocytic clearance is more prominent (Xavier et al., 2015). In our study, we did not detect an age-associated increase in apoptotic signal in the SVZ, arguing against a simple model in which increased microglial lysosomal content in aging is driven

Cutler, Harrison, et al.   30

primarily by increased clearance of apoptotic SVZ progenitors. Instead, our *in vivo* colocalization

and *in vitro* phagocytosis assays are consistent with increased microglial phagocytic programs and

increased microglial association with Mash1+ progenitor-derived material in the aged niche,

although we emphasize that colocalization-based metrics are proxies and do not, on their own,

establish engulfment kinetics or target selectivity.

Reanalysis of published single-cell transcriptomes from young and old SVZ provided an

independent line of evidence consistent with increased inflammatory and phagocytosis-related

programs in aged microglia. In pseudobulk AGDGE and gene-set analyses, aged microglia

upregulated inflammatory signaling and lysosome/phagosome-associated genes, including

multiple receptors and effectors linked to phagocytic function (e.g., *Axl*, *Tlr2*, *Lgal3*, and *C3*), and

showed broader increases across several phagocytosis pathway components (e.g. Complement, FC

gamma, and Toll-like). In contrast, neural stem/progenitor populations exhibited comparatively

few significant age-related DE genes, motivating gene-set–level testing rather than threshold

overlaps. Those analyses suggested that astrocyte-qNSCs and neuroblasts increase

complement/opsonization-related expression (i.e. "eat-me" programs) while also increasing

expression of putative anti-phagocytic programs, including MHC class I/antigen presentation–

associated genes (i.e. "don't eat-me" programs). aNSCs-NPCs did not show the same pattern of

increased "don't eat-me" programs, which is consistent with, but does not prove, a model in which

aNSCs/ TACs may be relatively vulnerable to microglial phagocytic engagement in the aged niche.

Importantly, transcriptomic signatures cannot distinguish increased phagocytic capacity from

compensatory responses to altered substrate availability, nor can they assign directionality without

functional validation. The strength of the scRNA-seq analysis is therefore as mechanistic context

that complements the *in vivo* and *in vitro* assays.

Microglia are distributed throughout the brain, but exhibit region-dependent heterogeneity in

density, morphology, transcriptional state, and receptor expression shaped by local

microenvironments (Grabert et al., 2016; Tan et al., 2020). SVZ microglia are reported to be

phenotypically distinct from non-germinal regions, supporting the idea of niche specialization

(Goings et al., 2006; Marshall et al., 2014; Shigemoto-Mogami et al., 2014; Solano et al., 2016;

Xavier et al., 2015). Riberio Xavier et al. showed that SVZ microglia in young adults have distinct

morphology and receptor profiles, are required for neuroblast survival and migration, and display

comparatively low phagocytic activity in the young SVZ niche, potentially linked to reduced

expression of specific purinergic receptors (Xavier et al., 2015). Our *in vitro* assay is consistent

with this niche-specific baseline state. SVZ-derived microglia from young mice exhibited lower

phagocytic activity than microglia isolated from non-SVZ brain regions. With aging, we observe

a shift toward increased CD68+ lysosomal features and increased phagocytic readouts, consistent

with remodeling of this niche-specialized state.

To probe whether inflammatory signaling contributes causally to SVZ aging phenotypes, we used

CR2-Crry, a targeted inhibitor of complement C3 activation and deposition (Atkinson et al., 2005;

Song et al., 2003) that crosses the blood brain barrier (Alawieh et al., 2018). This intervention

partially rescued SVZ proliferation in aged mice, supporting the conclusion that complement-

linked signaling contributes to the age-associated reduction in proliferative output. Although

insignificant, CR2-Crry treatment reduced microglia inflammation and Mash1+ material within

CD68+ compartments in aged microglia. Future work will be aimed at the direct quantification of

engulfment dynamics, and mechanistic tests of candidate receptor–ligand axes suggested by the

transcriptomic analyses.

Finally, our transcriptomic observations raise a testable hypothesis about target susceptibility.

Aging may increase "eat-me" programs in multiple SVZ populations, while "don't eat-me"

programs may be relatively stronger in astrocyte-qNSCs and neuroblasts than in aNSCs and TACs.

If validated, this could help explain why increased microglial phagocytic programs in aging are

associated with progenitor loss while sparing other SVZ populations. Future work should directly

test whether differences in cell-surface inhibitory cues (e.g., *Cd200*, *Pecam1*, MHC class I–linked

programs) and corresponding microglial inhibitory receptors shape selective engagement of

progenitors in the aged niche, and whether manipulating these interactions can decouple restored

proliferation from progenitor loss, potentially paving the way for future therapeutics.

Several limitations should be considered when interpreting these findings. First, our *in vivo*

phagocytosis readout relies on confocal colocalization of Mash1 signal within CD68+

compartments in Iba1+ microglia in SVZ wholemounts. While this pattern is consistent with

phagolysosomal internalization, additional validation using phagosome markers or ultrastructural

approaches will strengthen localization and interpretation. Second, the *in vitro* phagocytosis assay

uses pHrodo-conjugated yeast/zymosan-like particles, which can engage receptors such as TLR2

and dectin-1. Consequently, the observed age- and region-dependent differences may reflect

pathway-specific responsiveness rather than global phagocytic capacity. Testing alternative cargos

(e.g., apoptotic bodies) and defined opsonization states will be necessary to assess generality.

Third, although the CR2-Crry intervention is designed to inhibit C3 activation and deposition, we

did not directly measure complement activation or target engagement within the SVZ following

treatment. This limits causal inference about the extent and spatial specificity of complement

inhibition in the niche and makes it difficult to distinguish direct complement effects from

downstream or systemic consequences.

Additional caveats apply to generalizability and mechanistic resolution. Our work focuses on

discrete ages (3 and 12 months for most experiments), which captures a midlife transition but does

not resolve the temporal ordering of microglial activation, phagocytic engagement, and progenitor

decline across aging; denser longitudinal sampling would better establish sequence and potential

thresholds. The CR2-Crry intervention may not be microglia-specific and could act on other niche

cells or peripheral immune pathways, complicating attribution of effects to microglia versus the

broader SVZ environment. Moreover, increases in EdU+ counts provide a useful proliferative

readout but do not by themselves resolve whether the rescued proliferation reflects altered cell-

cycle dynamics, survival, lineage progression, or changes in the composition of proliferating

populations; additional phenotyping of EdU+ cells would sharpen interpretation. Finally, the

scRNA-seq reanalysis provides supportive mechanistic context, but transcriptomic signatures

cannot distinguish increased phagocytic "capacity" from compensatory programs or assign

directionality of cell–cell interactions; functional perturbations of candidate receptor–ligand axes

and direct measures of engulfment dynamics will be required to establish causality.

References

Ahlenius, H., Visan, V., Kokaia, M., Lindvall, O., & Kokaia, Z. (2009). Neural stem and progenitor cells retain their potential for proliferation and differentiation into functional neurons despite lower number in aged brain. *The Journal of Neuroscience*, *29*(14), 4408–4419. https://doi.org/10.1523/JNEUROSCI.6003-08.2009

Alawieh, A., Zandian, J., & Feng, W. (2018). Factors affecting post-stroke motor recovery: Implications on neurotherapy after brain injury. *Behavioural Brain Research*, *340*, 94–101. https://doi.org/10.1016/j.bbr.2016.08.029

Alvarez-Buylla, A., & García-Verdugo, J. M. (2002). Neurogenesis in adult subventricular zone. *The Journal of Neuroscience*, *22*(3), 629–634. https://doi.org/10.1523/JNEUROSCI.22-03-00629.2002

Atkinson, C., Song, H., Lu, B., Qiao, F., Burns, T. A., Holers, V. M., Tsokos, G. C., & Tomlinson, S. (2005). Targeted complement inhibition by C3d recognition ameliorates tissue injury without apparent increase in susceptibility to infection. *Journal of Clinical Investigation*, *115*(9), 2444–2453. https://doi.org/10.1172/JCI25208

Cunningham, C. L., Martínez-Cerdeño, V., & Noctor, S. C. (2013). Microglia regulate the number of neural precursor cells in the developing cerebral cortex. *The Journal of Neuroscience*, *33*(10), 4216–4233. https://doi.org/10.1523/JNEUROSCI.3441-12.2013

Cutler, R. R., & Kokovay, E. (2020). Rejuvenating subventricular zone neurogenesis in the aging brain. *Current Opinion in Pharmacology*, *50*, 1–8. https://doi.org/10.1016/j.coph.2019.10.005

Deczkowska, A., Keren-Shaul, H., Weiner, A., Colonna, M., Schwartz, M., & Amit, I. (2018). Disease-associated microglia: A universal immune sensor of neurodegeneration. *Cell*, *173*(5), 1073–1081. https://doi.org/10.1016/j.cell.2018.05.003

Doetsch, F., Caillé, I., Lim, D. A., García-Verdugo, J. M., & Alvarez-Buylla, A. (1999). Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell*, *97*(6), 703–716. https://doi.org/10.1016/s0092-8674(00)80783-7

Dulken, B. W., Buckley, M. T., Navarro Negredo, P., Saligrama, N., Cayrol, R., Leeman, D. S., ... & Brunet, A. (2019). Single-cell analysis reveals T cell infiltration in old neurogenic niches. *Nature*, *571*(7764), 205–210. https://doi.org/10.1038/s41586-019-1362-5

Enwere, E., Shingo, T., Gregg, C., Fujikawa, H., Ohta, S., & Weiss, S. (2004). Aging results in reduced epidermal growth factor receptor signaling, diminished olfactory neurogenesis, and deficits in fine olfactory discrimination. *The Journal of Neuroscience*, *24*(38), 8354–8365. https://doi.org/10.1523/JNEUROSCI.2751-04.2004

Ginhoux, F., Greter, M., Leyane, M., Nandi, S., See, P., Sivasankar, B., ... & Merad, M. (2010). Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science*, *330*(6005), 841–845. https://doi.org/10.1126/science.1194637

Goings, D. E., Kozlowski, K. A., & Szele, F. G. (2006). Differential activation of microglia in neurogenic versus non-neurogenic regions of the forebrain. *Glia*, *54*(4), 329–342. https://doi.org/10.1002/glia.20381

Grabert, K., Michoel, T., Karavolos, M. H., Clohisey, S., Baillie, J. K., Stevens, M. P., ... & McColl, B. W. (2016). Microglial brain region-dependent diversity and selective regional sensitivities to aging. *Nature Neuroscience*, *19*(3), 504–516. https://doi.org/10.1038/nn.4222

Haney, M. S., Bohlen, C. J., Morgens, D. W., Ousey, J. A., Barkal, A. A., Tsui, C. K., ... & Bassik, M. C. (2018). Identification of phagocytosis regulators using magnetic genome-wide CRISPR screens. *Nature Genetics*, *50*(12), 1716–1727. https://doi.org/10.1038/s41588-018-0254-1

Hu, B., Duan, S., Wang, Z., Li, X., Zhou, Y., Zhang, X., ... & Zheng, H. (2021). Insights into the role of CSF1R in the central nervous system and neurological disorders. *Frontiers in Aging Neuroscience*, *13*, 789834. https://doi.org/10.3389/fnagi.2021.789834

Izgi, H., Han, D., Isildak, U., Huang, S., Kocabiyik, E., Khaitovich, P., Somel, M., & Dönertaş, H. M. (2022). Inter-tissue convergence of gene expression during ageing suggests age-related loss of tissue and cellular identity. *eLife*, *11*, e68048. https://doi.org/10.7554/eLife.68048

Kalamakis, G., Brüne, D., Ravichandran, S., Bolz, J., Fan, W., Ziebell, F., ... & Martin-Villalba, A. (2019). Quiescence modulates stem cell maintenance and regenerative capacity in the aging brain. *Cell*, *176*(6), 1407–1419. https://doi.org/10.1016/j.cell.2019.01.040

Kapellos, T. S., Taylor, L., Lee, H., Cowley, S. A., James, W. S., Iqbal, A. J., & Greaves, D. R. (2016). A novel real time imaging platform to quantify macrophage phagocytosis. *Biochemical Pharmacology*, *116*, 107–119. https://doi.org/10.1016/j.bcp.2016.07.011

Khalaji, A., Yancheshmeh, F. B., Farham, F., Khorram, A., Sheshbolouki, S., Zokaei, M., Vatankhah, F., & Soleymani-Goloujeh, M. (2023). Don't eat me/eat me signals as a novel strategy in cancer immunotherapy. *Heliyon*, *9*(10), e20507. https://doi.org/10.1016/j.heliyon.2023.e20507

Kim, E. J., Ables, J. L., Dickel, L. K., Eisch, A. J., & Johnson, J. E. (2011). Ascl1 (Mash1) defines cells with long-term neurogenic potential in subgranular and subventricular zones in adult mouse brain. *PLoS One*, *6*(3), e18472. https://doi.org/10.1371/journal.pone.0018472

Li, W. (2012). Eat-me signals: Keys to molecular phagocyte biology and "appetite" control. *Journal of Cellular Physiology*, *227*(4), 1291–1297. https://doi.org/10.1002/jcp.22815

Liang, H., Zhang, H., Geng, A., Mao, M., Tong, S., Tu, S., ... & Chen, S. T. (2019). Region-
specific and activity-dependent regulation of SVZ neurogenesis and recovery after stroke.
Proceedings of the National Academy of Sciences, *116*(27), 13621–13630.
https://doi.org/10.1073/pnas.1811825116

Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J. P., & Tamayo, P. (2015).
The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Systems*, *1*(6),
417–425. https://doi.org/10.1016/j.cels.2015.12.004

Lim, D. A., & Alvarez-Buylla, A. (2016). The adult ventricular–subventricular zone (V-SVZ)
and olfactory bulb (OB) neurogenesis. *Cold Spring Harbor Perspectives in Biology*, *8*(5),
a018820. https://doi.org/10.1101/cshperspect.a018820

Lledo, P. M., & Valley, M. T. (2016). Adult olfactory bulb neurogenesis. *Cold Spring Harbor*
Perspectives in Biology, *8*(8), a018945. https://doi.org/10.1101/cshperspect.a018945

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.
https://doi.org/10.1186/s13059-014-0550-8

Luo, J., Daniels, S. B., Lennington, J. B., Notti, R. Q., & Conover, J. C. (2006). The aging
neurogenic subventricular zone. *Aging Cell*, *5*(2), 139–152. https://doi.org/10.1111/j.1474-
9726.2006.00197.x

Marshall, G. P., Deleyrolle, L. P., Reynolds, B. A., Steindler, D. A., & Laywell, E. D. (2014).
Microglia from neurogenic and non-neurogenic regions display differential proliferative
potential and neuroblast support. *Frontiers in Cellular Neuroscience*, *8*, 180.
https://doi.org/10.3389/fncel.2014.00180

Maslov, A. Y., Barone, T. A., Plunkett, R. J., & Pruitt, S. C. (2004). Neural stem cell detection,
characterization, and age-related changes in the subventricular zone of mice. *The Journal of*
Neuroscience, *24*(7), 1726–1733. https://doi.org/10.1523/JNEUROSCI.4608-03.2004

Morrison, V., Holloway, M. G., Taylor, J. B., Bassik, A. A., Kim, P. J., Kim, K. A., ... & Carter,
B. D. (2023). Jedi-1/MEGF12-mediated phagocytosis controls the pro-neurogenic properties of
microglia in the ventricular-subventricular zone. *bioRxiv*.
https://doi.org/10.1101/2023.03.03.531012

Nath, S., Martínez Santamaría, J. C., Chu, Y. H., Choi, J. S., Conforti, P., Lin, J. D., ... &
Schachtrup, C. (2024). Interaction between subventricular zone microglia and neural stem cells
impacts the neurogenic response in a mouse model of cortical ischemic stroke. *Nature*
Communications, *15*(1), 9095. https://doi.org/10.1038/s41467-024-53217-1

Ohab, J. J., Fleming, S., Blesch, A., & Carmichael, S. T. (2006). A neurovascular niche for
neurogenesis after stroke. *The Journal of Neuroscience*, *26*(50), 13007–13016.
https://doi.org/10.1523/JNEUROSCI.4323-06.2006

Pluvinage, J. V., Haney, M. S., Smith, B. A. H., Sun, J., Iram, T., Bonanno, L., ... & Wyss-
Coray, T. (2019). CD22 blockade restores homeostatic microglial phagocytosis in ageing brains.
Nature, *568*(7751), 187–192. https://doi.org/10.1038/s41586-019-1088-4

Robin, A. M., Zhang, Z. G., Wang, L., Zhang, R. L., Katakowski, M., Zhang, L., ... & Chopp, M.
(2006). Stromal cell-derived factor 1alpha mediates neural progenitor cell motility after focal
cerebral ischemia. *Journal of Cerebral Blood Flow & Metabolism*, *26*(1), 125–134.
https://doi.org/10.1038/sj.jcbfm.9600172

Shigemoto-Mogami, Y., Hoshikawa, K., Goldman, J. E., Sekino, Y., & Sato, K. (2014).
Microglia enhance neurogenesis and oligodendrogenesis in the early postnatal subventricular
zone. *The Journal of Neuroscience*, *34*(6), 2231–2243.
https://doi.org/10.1523/JNEUROSCI.1619-13.2014

Solano, F. G., Ma, S., Ameloot, D. M., Rivera, R., D'Amours, A., Chemtob, S., ... & Kokovay,
E. (2016). Neurogenic niche microglia undergo positional remodeling and progressive activation
contributing to age-associated reductions in neurogenesis. *Stem Cells and Development*, *25*(7),
542–555. https://doi.org/10.1089/scd.2015.0319

Song, H., He, C., Knaak, C., Guthridge, J. M., Holers, V. M., & Tomlinson, S. (2003).
Complement receptor 2–mediated targeting of complement inhibitors to sites of complement
activation. *Journal of Clinical Investigation*, *111*(12), 1875–1885.
https://doi.org/10.1172/JCI17348

Tan, Y., Zheng, Y., & Tang, L. (2020). Microglial regional heterogeneity and its role in the
brain. *Molecular Psychiatry*, *25*(2), 351–367. https://doi.org/10.1038/s41380-019-0609-8

Xavier, A. L., Kress, B. T., Goldman, S. A., de Menezes, J. R., & Nedergaard, M. (2015). A
distinct population of microglia supports adult neurogenesis in the subventricular zone. *The
Journal of Neuroscience*, *35*(34), 11848–11861. https://doi.org/10.1523/JNEUROSCI.1217-
15.2015

Yamashita, T., Ninomiya, M., Hernández Acosta, P., García-Verdugo, J. M., Sunabori, T.,
Sakaguchi, M., ... & Sawamoto, K. (2006). Subventricular zone-derived neuroblasts migrate and
differentiate into mature neurons in the post-stroke adult striatum. *The Journal of Neuroscience*,
26(24), 6541–6552. https://doi.org/10.1523/JNEUROSCI.0149-06.2006

Yen, J. H., & Yu, I. C. (2024). The role of ApoE-mediated microglial lipid metabolism in brain
aging and disease. *Immunometabolism*, *5*(1), e00018.
https://doi.org/10.1097/IN9.0000000000000018

Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). clusterProfiler: An R package for comparing
biological themes among gene clusters. *OMICS: A Journal of Integrative Biology*, *16*(5), 284–
287. https://doi.org/10.1089/omi.2011.0118